

*Pacific  
Journal of  
Mathematics*

**SELF-SIMILAR SOLUTIONS OF THE  $p$ -LAPLACE HEAT  
EQUATION: THE FAST DIFFUSION CASE**

MARIE FRANÇOISE BIDAUT-VÉRON

Volume 227 No. 2

October 2006



## SELF-SIMILAR SOLUTIONS OF THE $p$ -LAPLACE HEAT EQUATION: THE FAST DIFFUSION CASE

MARIE FRANÇOISE BIDAUT-VÉRON

We study the self-similar solutions of the equation  $u_t - \operatorname{div}(|\nabla u|^{p-2}\nabla u) = 0$  in  $\mathbb{R}^N$ , where  $N \geq 1$ ,  $p \in (1, 2)$ . We provide a complete description of the signed solutions of the form  $u(x, t) = (\pm t)^{-\alpha/\beta} w((\pm t)^{-1/\beta}|x|)$ , regular or singular at  $x = 0$ , with  $\alpha, \beta$  real,  $\beta \neq 0$ , and possibly not defined on all of  $\mathbb{R}^N \times (0, \pm\infty)$ .

### 1. Introduction and main results

In this article we study the existence of self-similar solutions of the degenerate parabolic equation involving the  $p$ -Laplace operator in  $\mathbb{R}^N$ ,  $N \geq 1$ ,

$$(E_u) \quad u_t - \operatorname{div}(|\nabla u|^{p-2}\nabla u) = 0,$$

with  $1 < p < 2$ . In the sequel we set

$$\delta = \frac{p}{2-p},$$

so  $\delta > 1$ . Two critical values  $P_1, P_2$  are involved in the problem

$$P_1 = \frac{2N}{N+1}, \quad P_2 = \frac{2N}{N+2};$$

see [DiBenedetto and Herrero 1990], for example. They are connected with  $\delta$  through the relations

$$p > P_1 \iff \delta > N, \quad p > P_2 \iff \delta > \frac{N}{2}.$$

If  $u(x, t)$  is a solution and  $\alpha, \beta \in \mathbb{R}$ , then  $u_\lambda(x, t) = \lambda^\alpha u(\lambda x, \lambda^\beta t)$  is a solution of  $(E_u)$  if and only if

$$\beta = p - (2-p)\alpha = (2-p)(\delta - \alpha);$$

---

MSC2000: primary 35K65; secondary 34C35.

Keywords: degenerate parabolic equations, self-similar solutions.

thus  $\beta > 0$  if and only if  $\alpha < \delta$ . For given  $\alpha \in \mathbb{R}$  such that  $\alpha \neq \delta$ , the natural way to construct particular solutions is to search for self-similar solutions, radially symmetric in  $x$ , of the form

$$(1-1) \quad u = u(x, t) = (\varepsilon\beta t)^{-\alpha/\beta} w(r), \quad r = (\varepsilon\beta t)^{-1/\beta} |x|,$$

where  $\varepsilon = \pm 1$ . By translation, for any real  $T$ , we obtain solutions defined for any  $t > T$  when  $\varepsilon\beta > 0$ , or  $t < T$  when  $\varepsilon\beta < 0$ . The hypersurfaces  $\{r = \text{constant}\}$  are of focusing type if  $\beta > 0$  and of spreading type if  $\beta < 0$ . We are led to the equation

$$(E_w) \quad (|w'|^{p-2} w')' + \frac{N-1}{r} |w'|^{p-2} w' + \varepsilon(rw' + \alpha w) = 0 \quad \text{in}(0, \infty).$$

If we look for solutions of  $(E_u)$  under the form

$$u = Ae^{-\varepsilon\mu t} w(r), \quad r = Me^{-\varepsilon\mu t/\delta} |x|, \quad \mu > 0,$$

then  $w$  solves  $(E_w)$  provided  $M = \delta/\alpha$  and  $A = (\delta^p/\alpha^{p-1}\mu)^{1/(2-p)}$ , where  $\alpha > 0$  is arbitrary. This is another motivation for studying equation  $(E_w)$  for any real  $\alpha$ .

In the huge literature on self-similar solutions of parabolic equations, many results deal with positive solutions  $u$  defined and smooth on  $\mathbb{R}^N \times (0, \infty)$ . Equation  $(E_w)$  was studied in [Qi and Wang 1999] when  $\alpha > 0$ ,  $\varepsilon = 1$ . In our work we provide an exhaustive description of the self-similar solutions of equation  $(E_u)$ , possibly not defined on all of  $(0, \infty)$ , with constant or changing sign. In particular, for suitable values of  $\alpha$ , we prove the existence of solutions  $w$  oscillating with respect to 0 as  $r$  tends to 0 or  $\infty$ , or constant-sign solutions oscillating with respect to some nonzero constant. Our main tool is the reduction of the problem to an autonomous system with two variables and two parameters,  $p$  and  $\alpha$ . We are led to a dynamical system, which we study by phase-plane techniques. When  $p = \frac{3}{2}$ , this system is nearly quadratic, and many devices from the theory of algebraic dynamical systems can be used. In the general case such structures do not exist; then we use energy functions associated to the system. The behavior of the solutions presents great diversity, according to the possible values of  $p$  and  $\alpha$ .

In the sequel we set

$$\eta = \frac{N-p}{p-1};$$

thus  $\eta > 0$  if  $N \geq 2$ , and  $\eta = -1$  if  $N = 1$ . Observe the relation connecting  $\eta$ ,  $\delta$  and  $N$ :

$$(1-2) \quad \frac{\delta - N}{p - 1} = \delta - \eta = \frac{N - \eta}{2 - p}.$$

**Explicit solutions.** Obviously if  $w$  is a solution of  $(E_w)$ , so is  $-w$ . Many particular solutions are well-known.

**The infinite point source solution  $U_\infty$ .** The simplest positive solutions of equation  $(E_w)$ , which exist for any  $\alpha$  such that  $\varepsilon(\delta - N)(\delta - \alpha) > 0$ , are given by

$$(1-3) \quad w(r) = \ell r^{-\delta},$$

where

$$(1-4) \quad \ell = \left( \varepsilon \delta^{p-1} \frac{\delta - N}{\delta - \alpha} \right)^{1/(2-p)} > 0.$$

They correspond to a unique solution  $u$  of  $(E_u)$  called  $U_\infty$  in [Chasseigne and Vazquez 2002], singular at  $x = 0$ , for any  $t \neq 0$ :

$$U_\infty(x, t) = \left( \frac{Ct}{|x|^p} \right)^{1/(2-p)}, \quad C = (2 - p)\delta^{p-1}(\delta - N).$$

**The case  $\alpha = N$ .** Here the equation  $(E_w)$  has a first integral

$$(1-5) \quad w + \varepsilon r^{-1} |w'|^{p-2} w' = Cr^{-N}.$$

All the solutions corresponding to  $C = 0$  are given by

$$(1-6) \quad \begin{aligned} w &= w_{K,\varepsilon}(r) = \pm(\varepsilon\delta^{-1}r^{p'} + K)^{-\delta/p'}, \\ u &= \pm u_{K,\varepsilon}(x, t) = (\varepsilon\beta_N t)^{-N/\beta_N} (\varepsilon\delta^{-1}(\varepsilon\beta_N t)^{-p'/\beta_N} |x|^{p'} + K)^{-(p-1)/(2-p)}, \end{aligned}$$

$K \in \mathbb{R}$ ,

with  $\beta = \beta_N = (N + 1)(p - P_1)$ . For  $p > P_1$ ,  $\varepsilon = 1$ ,  $K > 0$ , the solutions are named after Barenblatt [1952]. For given  $c > 0$ , the function  $u_{K,1}$ , defined on  $\mathbb{R}^N \times (0, \infty)$ , is the unique solution of equation  $(E_u)$  with initial data  $u(0) = c\delta_0$ ,  $\delta_0$  being the Dirac mass at 0 and  $K$  begin determined by  $\int_{\mathbb{R}^N} u_K(x, t) dt = c$ ; see for example [Zhao 1993]. Moreover the functions  $u_{K,1}$ , with  $K > 0$ , are the only nonnegative solutions defined on  $\mathbb{R}^N \times (0, \infty)$ , such that  $u(x, 0) = 0$  for any  $x \neq 0$ ; see [Kamin and Vázquez 1992]. In the case  $K = 0$ , we find again the function  $U_\infty$ , and  $U_\infty$  is the limit of the functions  $u_{K,1}$  as  $K \rightarrow 0$ , or equivalently  $c \rightarrow \infty$ .

**The case  $\alpha = \eta$ .** We exhibit a family of solutions of  $(E_w)$ :

$$(1-7) \quad w(r) = Cr^{-\eta}, \quad u(t, x) = C|x|^{-\eta} = C|x|^{(p-N)/(p-1)}, \quad C \neq 0,$$

Solutions  $u$ , independent of  $t$ , are the fundamental  $p$ -harmonic solutions of the equation when  $p > P_1$ .

**The case  $\alpha = -p'$ .** Equation  $(E_w)$  admits solutions of the form

$$(1-8) \quad \begin{aligned} w(r) &= \pm K(N(Kp')^{p-2} + \varepsilon r^{p'}), \\ u(x, t) &= \pm K(N(Kp')^{p-2}t + \varepsilon|x|^{p'}), \quad K > 0, \end{aligned}$$

and the functions  $u$  are solutions of the form  $\psi(t) + \Phi(|x|)$  with  $\Phi$  nonconstant. They have constant sign when  $\varepsilon = 1$ , and a changing sign when  $\varepsilon = -1$ .

**The case  $\alpha = 0$ .** Here equation  $(E_w)$  can be explicitly solved: either  $w' \equiv 0$  (hence  $w \equiv a \in \mathbb{R}$ , and  $u$  is a constant solution of  $(E_u)$ ), or there exists  $K \in \mathbb{R}$  such that

$$(1-9) \quad |w'| = r^{(1-N)/(p-1)} \times \begin{cases} \left(K + \frac{\varepsilon}{\delta - N} r^{N-\eta}\right)^{-1/(2-p)} & \text{if } \delta \neq N, \\ \left(\frac{2-p}{p-1}(K + \varepsilon \ln r)\right)^{-1/(2-p)} & \text{if } \delta = N, \end{cases}$$

which gives  $w$  by integration, up to a constant, and then  $u(x, t) = w(|x|/(\varepsilon pt)^{1/p})$ .

**The case  $N = 1$  and  $\alpha = (p - 1)/(2 - p) > 0$ .** Here again we obtain explicit solutions:

$$w(r) = \pm(\varepsilon K(r - (K\alpha)^{p-1}))^{-\alpha}, \quad u(x, t) = \pm(\varepsilon K(|x| - \varepsilon(K\alpha)^{p-1}t))^{-\alpha}, \quad K > 0.$$

All the functions  $w$  above are defined on intervals of the form  $(R, 0)$ ,  $R \geq 0$  if  $\varepsilon = 1$ , and  $(0, S)$ ,  $S \leq \infty$  if  $\varepsilon = -1$ .

**Note.** When  $\alpha = \delta$ , equation  $(E_u)$  is invariant under the transformation  $u_\lambda(x, t) = \lambda^\alpha u(\lambda x, t)$ ; searching solutions of the form  $u(x, t) = |x|^{-\delta} \psi(t)$ , we find again the function  $U_\infty$ .

**Different kinds of singularities.** Consider equation  $(E_w)$ . It is easy to get local existence and uniqueness near any point  $r_1 > 0$ ; thus any solution  $w$  is defined on a maximal interval  $(R_w, S_w)$ , with  $0 \leq R_w < S_w \leq \infty$ ; and in fact  $S_w = \infty$  when  $\varepsilon = 1$ , and  $R_w = 0$  when  $\varepsilon = -1$  (see Theorem 2.2). Returning to solution the  $u$  of  $(E_u)$  associated to  $w$  by (1-1), it is defined on a subset of  $\mathbb{R}^N \setminus \{0\} \times (0, \pm\infty)$ :

$$D_w = \{(x, t) : x \in \mathbb{R}^N, \varepsilon\beta t > 0, (\varepsilon\beta t)^{1/\beta} R_w < |x| < (\varepsilon\beta t)^{1/\beta} S_w\}.$$

When  $w$  is defined on  $(0, \infty)$ , then  $u$  is defined on  $\mathbb{R}^N \setminus \{0\} \times (0, \pm\infty)$ .

**Regular solutions.** Among the solutions of  $(E_w)$  defined near 0, we also show the existence and uniqueness of solutions  $w = w(\cdot, a) \in C^2([0, S_w))$  such that, for some  $a \in \mathbb{R}$ ,

$$(1-10) \quad w(0) = a, \quad w'(0) = 0.$$

These are called *regular solutions*. Obviously, they are defined on  $[0, \infty)$  when  $\varepsilon = 1$ . If  $w$  is regular, then  $D_w = \mathbb{R}^N \times (0, \pm\infty)$ , and  $u(\cdot, t) \in C^1(\mathbb{R}^N)$  for  $t \neq 0$ ; we will say that  $u$  is *regular*. This does not imply the regularity up to  $t = 0$ : indeed  $u$  presents a singularity at time  $t = 0$  if and only if  $0 < \alpha < \delta$ . In the sequel we shall not mention the trivial solution  $w \equiv 0$ , corresponding to  $a = 0$ .

**Singular solutions.** If  $R_w = 0$  and  $w$  is not regular,  $u$  presents a singularity at  $x = 0$  for  $t \neq 0$ , called a *standing singularity*. Following [Vazquez and Véron 1996; Chasseigne and Vazquez 2002], for such a solution, we say that  $x = 0$  is a *weak singularity* if  $x \mapsto w(|x|) \in L^1_{\text{loc}}(\mathbb{R}^N)$ , or equivalently if  $u(\cdot, t) \in L^1_{\text{loc}}(\mathbb{R}^N)$  for  $t \neq 0$ ; and a *strong singularity* if not. If  $u$  has a strong/weak singularity, and  $\lim_{t \rightarrow 0} u(t, x) = 0$  for any  $x \neq 0$ , we call  $u$  a *strong/weak razor blade*. If  $u(\cdot, t) \in L^1(\mathbb{R}^N)$  for  $t \neq 0$ , then  $u$  is called *integrable*.

**Solutions with a reduced domain.** If  $R_w > 0$  or  $S_w < \infty$ , we say that  $u$  and  $w$  have a *reduced domain*. Then  $D_w$  has a lateral boundary of the form  $\Sigma_w = \{|x| = C(\varepsilon\beta t)^{1/\beta}\}$ , of parabolic type if  $\beta > 0$  and of hyperbolic type if  $\beta < 0$ , and  $u$  has an explosion near  $\Sigma_w$ . In Proposition 2.15 we calculate the blow-up rate, which is of the order of  $d(x, t)^{-(p-1)/(2-p)}$ , where  $d(x, t)$  is the distance to  $\Sigma_w$ .

**Main results.** We give a summary of our main results, expressed in terms of the function  $u$ , avoiding for simplicity particular cases (such as  $N = 1$ , or  $\alpha = \delta$ , or  $p = P_1$ ) and solutions with a reduced domain (although there exist many such). All cases omitted here and detailed statements in terms of  $w$  can be found inside each section. An important critical value of  $\alpha$  is given by

$$(1-11) \quad \alpha^* = \delta + \frac{\delta(N - \delta)}{(p - 1)(2\delta - N)};$$

it appears when  $\varepsilon = 1$ ,  $p > P_2$ , and then  $\alpha^* > 0$ , or  $\varepsilon = -1$ ,  $p < P_2$ , and then  $\alpha^* < 0$ .

**Note.** To return from  $w$  to  $u$ , consider any solution  $w$  of  $(E_w)$  defined on  $(0, \infty)$ , such that for some  $\lambda \geq 0$  and  $\mu \in \mathbb{R}$ ,  $\lim_{r \rightarrow 0} r^\lambda w = c \neq 0$  and  $\lim_{r \rightarrow 0} r^\mu w = c' \neq 0$ . Then:

- (i) For fixed  $t$ ,  $u$  has a singularity in  $|x|^{-\lambda}$  near  $x = 0$ , and a behavior in  $|x|^{-\mu}$  for large  $|x|$ . Thus  $x = 0$  is a *weak singularity* if and only if  $\lambda < N$ , and  $u$  is integrable if and only if  $\lambda < N < \mu$ .
- (ii) For fixed  $x \neq 0$ , the behavior of  $u$  near  $t = 0$ , depends on the sign of  $\beta$ :

$$\begin{aligned} \lim_{t \rightarrow 0} |x|^\mu |t|^{(\alpha-\mu)/\beta} u(x, t) &= C \neq 0 \quad \text{if } \alpha < \delta, \\ \lim_{t \rightarrow 0} |x|^\lambda |t|^{(\alpha-\lambda)/\beta} u(x, t) &= C \neq 0 \quad \text{if } \delta < \alpha. \end{aligned}$$

**Solutions defined for  $t > 0$ .** Here we look for solutions  $u$  of  $(E_u)$  on  $\mathbb{R}^N \setminus \{0\} \times (0, \infty)$  of the form (1-1). That means  $\varepsilon\beta > 0$ , or equivalently  $\varepsilon = 1$  and  $\alpha < \delta$  (see Section 3) or  $\varepsilon = -1$ ,  $\delta < \alpha$  (see Section 4). We begin with the case  $\varepsilon = 1$ , and examine the dependence on the sign of  $p - P_1$ . For proofs, see Theorems 3.2, 3.4 and 3.5.

**Theorem 1.1.** Assume  $\varepsilon = 1$ ,  $-\infty < \alpha < \delta$ ,  $p > P_1$ , and  $N \geq 2$ . Then  $U_\infty$  is a solution on  $\mathbb{R}^N \setminus \{0\} \times (0, \infty)$  and a strong razor blade. There exist also positive solutions having a strong singularity in  $|x|^{-\delta}$  and satisfying  $\lim_{t \rightarrow 0} |x|^\alpha u = L > 0$  (for  $x \neq 0$ ). For  $\alpha \leq N$ , any function  $u(\cdot, t)$  has at most one zero at time  $t$ .

- (1) For  $\alpha < N$ , all regular solutions on  $\mathbb{R}^N \times (0, \infty)$  have constant sign, are not integrable, and they are solutions of  $(E_u)$  with initial data  $L|x|^{-\alpha} \in L^1_{\text{loc}}(\mathbb{R}^N)$ . There exist **positive integrable razor blades** having a singularity in  $|x|^{-\eta}$ . There exist also positive solutions having a weak regularity in  $|x|^{-\eta}$  and satisfying  $\lim_{t \rightarrow 0} |x|^\alpha u = L$ ; in particular if  $\alpha = \eta$ , then  $u \equiv C|x|^{-\eta}$ . There exist solutions with one zero and a weak or a strong singularity.
- (2) For  $\alpha = N$ , all regular (Barenblatt) solutions have constant sign and are integrable. There exist solutions with one zero and a weak singularity.
- (3) For  $N < \alpha$ , all regular solutions have at least one zero. If  $\alpha < \alpha^*$ , any solution has a **finite number of zeros**. If  $N < \alpha^*$ , there exists  $\check{\alpha} \in (\alpha^*, \delta)$  such that if  $\check{\alpha} < \alpha$ , regular solutions are **oscillating around 0** for large  $|x|$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ ; and there exists precisely a solution  $u$  such that  $r^\delta w$  is **periodic in  $\ln r$** .

**Theorem 1.2.** Assume  $\varepsilon = 1$ ,  $-\infty < \alpha < \delta$ , and  $p < P_1$ . Then all regular solutions on  $\mathbb{R}^N \times (0, \infty)$  have constant sign, are not integrable, and are solutions of  $(E_u)$  with initial data  $L|x|^{-\alpha} \in L^1_{\text{loc}}(\mathbb{R}^N)$ . There is no other solution on  $\mathbb{R}^N \setminus \{0\} \times (0, \infty)$ .

If  $\alpha > 0$ , all the solutions  $w$  tend to 0 at  $\infty$ , whereas if  $\alpha < 0$ , some of the solutions are unbounded near  $\infty$ .

Next we come to the case  $\varepsilon = -1$ , which is treated in Theorems 4.1 and 4.2.

**Theorem 1.3.** Assume  $\varepsilon = -1$ ,  $\delta < \alpha$ ,  $p > P_1$ , and  $N \geq 2$ . There is no regular solution on  $\mathbb{R}^N \times (0, \infty)$ . Besides the function  $U_\infty$ , which is a strong razor blade, there exist **positive integrable razor blades** having a singularity in  $|x|^{-\eta}$ , and positive solutions having a strong singularity in  $|x|^{-\alpha}$  and satisfying  $\lim_{t \rightarrow 0} |x|^\alpha u = L$ .

**Theorem 1.4.** Assume  $\varepsilon = -1$ ,  $\delta < \alpha$ ,  $p < P_1$  ( $N \geq 2$ ). There is no regular solution on  $\mathbb{R}^N \times (0, \infty)$ . There exists a positive solution on  $\mathbb{R}^N \setminus \{0\} \times (0, \infty)$  with a singularity in  $|x|^{-\alpha}$  (strong if and only if  $N \leq \alpha$ ), and  $\lim_{t \rightarrow 0} |x|^\alpha u = L$ .

**Note.** Weak singularities can occur even if  $p > P_1$ . For example, the solutions  $u(t, x) = C|x|^{-\eta} = C|x|^{(p-N)/(p-1)}$  ( $N \geq 2$ ) given in (1-7) have a weak singularity. There even exist positive solutions  $u$  with a standing singularity, and integrable; see Theorems 1.1 and 1.3. This is not contradictory with the regularizing effect  $L^1_{\text{loc}}(\mathbb{R}^N) \rightarrow L^\infty_{\text{loc}}(\mathbb{R}^N)$ , which concerns solutions in  $(0, \infty) \times \mathbb{R}^N$ . The functions constructed above are solutions in  $(0, \infty) \times \mathbb{R}^N \setminus \{0\}$ , and the singularity  $x = 0$  is not removable.

**Solutions defined for  $t < 0$ .** Next we consider the solutions defined for  $t < 0$ , and more generally for  $t < T$ . They correspond to  $\varepsilon = 1$ ,  $\delta < \alpha$  (Section 5), or  $\varepsilon = -1$ ,  $\alpha < \delta$  (Section 6). A main question in that case is the *extinction problem*: do there exist regular solutions  $u$  vanishing identically on  $\mathbb{R}$  at time  $T$ ? Do there exist singular razor blades, vanishing on  $\mathbb{R}^N \setminus \{0\}$  at time  $T$ ? Are they integrable?

One of our most significant results is the existence of two critical values  $\alpha_{\text{crit}} > 0$  (when  $P_2 < p < P_1$ ) and  $\alpha^{\text{crit}} < 0$  (when  $1 < p < P_2$ ), for which the *regular solutions*  $u_{\alpha_{\text{crit}}}$  are *positive, integrable*, and vanish identically at time 0. Another new phenomena is the existence of positive solutions such that  $C_1 U_\infty \leq u \leq C_2 U_\infty$  for some  $C_1, C_2 > 0$ , with a periodicity property, see Theorems 1.6 and 1.8.

First assume  $\varepsilon = 1$ . From Theorems 5.1 when  $p > P_1$  and 5.4, 5.6, 5.7 when  $p < P_1$ , we deduce:

**Theorem 1.5.** *Assume  $\varepsilon = 1$ ,  $\delta < \alpha$ ,  $p > P_1$ , with  $N \geq 2$ . Any solution  $u$  on  $\mathbb{R}^N \setminus \{0\} \times (0, -\infty)$ , in particular the regular ones, is **oscillating around 0** for fixed  $t < 0$  and large  $|x|$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ . There exists a solution such that  $r^\delta w$  is **periodic** in  $\ln r$ . There exist weak integrable razor blades, with a singularity in  $|x|^{-\eta}$ .*

**Theorem 1.6.** *Assume  $\varepsilon = 1$ ,  $\delta < \alpha$ ,  $p < P_1$ . Then  $U_\infty$  is a solution on  $\mathbb{R}^N \setminus \{0\} \times (0, -\infty)$ , and a weak razor blade.*

- (1) *If  $p < P_2$ , all regular solutions on  $\mathbb{R}^N \times (0, -\infty)$  have **constant sign**, are not integrable, and vanish identically at  $t = 0$ , with  $\|u(\cdot, t)\|_{L^\infty(\mathbb{R}^N)} \leq C|t|^{\alpha/|\beta|}$ . All the solutions have a finite number of zeros.*
- (2) *For  $\alpha < \eta$ , regular solutions have **constant sign**, with the same behavior (given by (1–6) if  $\alpha = N$ ). There exists a positive solution  $u$ , which is not integrable, with a singularity in  $|x|^{-\alpha}$  (a strong one if and only if  $\alpha \geq N$ ), and  $\lim_{t \rightarrow 0} |x|^\alpha u = L$ . If  $\alpha = \eta$ , then  $u(t, x) = C|x|^{-\eta}$  is a solution with a strong singularity.*
- (3) *If  $p > P_2$ , there exists a **critical value**  $\alpha_{\text{crit}}$  such that  $\eta < \alpha_{\text{crit}} < \alpha^*$  and the **regular solutions**  $u_{\alpha_{\text{crit}}}$  have **constant sign**, are **integrable**, and vanish identically at  $t = 0$ , with  $\|u(\cdot, t)\|_{L^\infty(\mathbb{R}^N)} \leq C|t|^{\alpha/|\beta|}$ .*
- (4) *If  $\alpha \in (\alpha_{\text{crit}}, \alpha^*)$ , there exist **positive** solutions  $u$  such that  $r^\delta w$  is **periodic** in  $\ln r$ ; thus*

$$C_1 U_\infty \leq u \leq C_2 U_\infty \quad \text{for some } C_1, C_2 > 0.$$

*There exist positive solutions  $u$ , with the same bounds, such that  $r^\delta w$  is asymptotically periodic near 0. There exist **positive integrable** solutions  $u$  such that  $r^\delta w$  is asymptotically periodic near 0.*

- (5) If  $\alpha_{\text{crit}} < \alpha$ , all regular solutions are oscillating around 0 for fixed  $t < 0$  and large  $|x|$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ . There exist solutions **oscillating around 0**, such that  $r^\delta w$  is **periodic**. If  $\alpha^* < \alpha$ , there exist **positive integrable razor blades**, with a singularity in  $|x|^{-\delta}$ .

Finally suppose  $\varepsilon = -1$ . From Theorems 6.1, 6.2 when  $p > P_1$  and 6.4, 6.6, 6.8, 6.9 when  $p < P_1$ , we obtain:

**Theorem 1.7.** Assume  $\varepsilon = -1$ ,  $\alpha < \delta$  and  $p > P_1$ , with  $N \geq 2$ . If  $\alpha > 0$ , there exist **positive** solutions  $u$  with a weak singularity in  $|x|^{-\eta}$ , integrable if and only if  $\alpha > N$ , and  $\lim_{t \rightarrow 0} |x|^\alpha u = L$ . If  $\alpha < 0$ , any solution has at least a zero. If  $-p' < \alpha$ , there is no regular solution on  $\mathbb{R}^N \times (0, -\infty)$ . If  $\alpha = -p'$ , all regular solutions, given by (1–8), have one zero.

**Theorem 1.8.** Assume  $\varepsilon = -1$ ,  $\alpha < \delta$  and  $p < P_1$ . Then  $U_\infty$  is a solution on  $\mathbb{R}^N \setminus \{0\} \times (0, -\infty)$ , and a weak razor blade.

- (1) If  $p > P_2$ , all the solutions have a finite number of zeros. There exist **positive integrable razor blades**, with a singularity in  $|x|^{-\delta}$ .
- (2) If  $-p' < \alpha$ , there is no regular solution on  $\mathbb{R}^N \times (0, -\infty)$ . There exist positive integrable razor blades as above. If  $\alpha > 0$ , there exist **positive** solutions  $u$  with a weak singularity in  $|x|^{-\delta}$ , integrable if and only if  $\alpha > N$ , and  $\lim_{t \rightarrow 0} |x|^\alpha u = L$ . If  $-p' < \alpha < 0$ , there exist solutions with one zero and the same behavior. If  $\alpha = -p'$ , all regular solutions, given by (1–8), have one zero.
- (3) If  $p < P_2$ , there exists a critical value  $\alpha^{\text{crit}}$  such that  $\alpha^* < \alpha^{\text{crit}} < -p'$  for which the **regular solutions**  $u_{\alpha^{\text{crit}}}$  have **constant sign**, are **integrable**, and vanish identically at  $t = 0$ , with  $\|u(\cdot, t)\|_{L^\infty(\mathbb{R}^N)} \leq C|t|^{\alpha/|\beta|}$ .
- (4) If  $p < P_2$  and  $\alpha \in (\alpha^*, \alpha^{\text{crit}})$ , there exist **positive** solutions  $u$  such that  $r^\delta w$  is **periodic** in  $\ln r$ , and thus

$$C_1 U_\infty \leq u \leq C_2 U_\infty \quad \text{for some } C_1, C_2 > 0.$$

There exist positive solutions with a weak singularity in  $|x|^{-\delta}$ , with the same bounds, such that  $r^\delta w$  is asymptotically periodic near  $\infty$ . The regular solutions have **constant sign**, are not integrable, vanish identically at  $t = 0$ , and  $r^\delta w$  is asymptotically periodic near  $\infty$ .

- (5) If  $p < P_2$  and  $\alpha < \alpha^{\text{crit}}$ , there exist solutions **oscillating around 0**, such that  $r^\delta w$  is **periodic**. There exist solutions **oscillating around 0**, integrable, such that  $r^\delta w$  is asymptotically periodic. If  $\alpha \leq \alpha^*$ , all regular solutions have **constant sign**, are not integrable, and vanish identically at  $t = 0$ .

**Note.** If  $p < P_1$ , recall that the Harnack inequality does not hold, as can be shown by the regular positive solutions constructed in Theorem 1.6, in particular those given by (1–6) when  $\alpha = N$ . The two kinds of regular, integrable, solutions constructed for the critical values  $\alpha_{\text{crit}} > 0$  and  $\alpha^{\text{crit}} < 0$  are of different types: the first, constructed for  $p > P_2$ , disappears in a spreading way, the second, for  $p < P_2$ , disappears in a focusing way.

The case  $p > 2$  will be treated in a second article [Bidaut-Véron 2006b], where we complete the results of [Gil and Vázquez 1997].

## 2. General properties

**Different formulations of the problem.** In the remainder of the article we can assume that  $\alpha \neq 0$ , since the solutions are given explicitly by (1–9) when  $\alpha = 0$ . Defining

$$(2-1) \quad \begin{aligned} J_N(r) &= r^N(w + \varepsilon r^{-1}|w'|^{p-2}w'), \\ J_\alpha(r) &= r^{\alpha-N}J_N(r), \end{aligned}$$

$(E_w)$  can be written in an equivalent way under the form

$$(2-2) \quad \begin{aligned} J'_N(r) &= r^{N-1}(N - \alpha)w, \\ J'_\alpha(r) &= -\varepsilon(N - \alpha)r^{\alpha-2}|w'|^{p-2}w'. \end{aligned}$$

If  $\alpha = N$ , then  $J_N$  is constant, so we find again (1–5).

We shall often use the following logarithmic substitution; for given  $d \in \mathbb{R}$ , setting

$$(2-3) \quad w(r) = r^{-d}y_d(\tau), \quad Y_d = -r^{(d+1)(p-1)}|w'|^{p-2}w', \quad \tau = \ln r,$$

we obtain the equivalent system

$$(2-4) \quad \begin{aligned} y'_d &= dy_d - |Y_d|^{(2-p)/(p-1)}Y_d, \\ Y'_d &= (p-1)(d-\eta)Y_d + \varepsilon e^{(p+(p-2)d)\tau}(\alpha y_d - |Y_d|^{(2-p)/(p-1)}Y_d). \end{aligned}$$

And  $y_d, Y_d$  satisfy the equations

$$(2-5) \quad \begin{aligned} y''_d + (\eta - 2d)y'_d - d(\eta - d)y_d \\ + \frac{\varepsilon}{p-1}e^{((p-2)d+p)\tau}|dy_d - y'_d|^{2-p}(y'_d + (\alpha - d)y_d) = 0, \end{aligned}$$

$$(2-6) \quad \begin{aligned} Y''_d + (p-1)(\eta - 2d - p')Y'_d + \varepsilon e^{((p-2)d+p)\tau}|Y_d|^{\frac{2-p}{p-1}}\left(\frac{Y'_d}{p-1} + (\alpha - d)Y_d\right) \\ - (p-1)^2(\eta - d)(p' + d)Y_d = 0. \end{aligned}$$

**Reduction to an autonomous system.** The substitution (2-3) with  $d = \delta$  is the most helpful: setting

$$(2-7) \quad y = y_d, \quad w(r) = r^{-\delta} y(\tau), \quad Y = -r^{(\delta+1)(p-1)} |w'|^{p-2} w', \quad \tau = \ln r,$$

we are led to the *autonomous* system that plays a key role in the sequel:

$$(S) \quad \begin{aligned} y' &= \delta y - |Y|^{(2-p)/(p-1)} Y, \\ Y' &= (\delta - N)Y + \varepsilon(\alpha y - |Y|^{(2-p)/(p-1)} Y). \end{aligned}$$

Since  $N - \delta p = \eta - 2\delta$  and  $N - \delta = (p - 1)(\eta - \delta)$ , Equation (2-5) takes the form

$$(E_y) \quad (p - 1)y'' + (N - \delta p)y' + \delta(\delta - N)y + \varepsilon|\delta y - y'|^{2-p}(y' + (\alpha - \delta)y) = 0,$$

while Equation (2-6) becomes

$$(E_Y) \quad Y'' + (N - 2\delta)Y' + \frac{\varepsilon}{p - 1} |Y|^{(2-p)/(p-1)} Y' + \varepsilon(\alpha - \delta) |Y|^{(2-p)/(p-1)} Y + \delta(\delta - N)Y = 0.$$

When  $w$  has constant sign, we define two functions associated to  $(y, Y)$ :

$$(2-8) \quad \begin{aligned} \zeta(\tau) &= \frac{|Y|^{(2-p)/(p-1)} Y}{y}(\tau) = -\frac{r w'(r)}{w(r)}, \\ \sigma(\tau) &= \frac{Y}{y}(\tau) = -\frac{|w'(r)|^{p-2} w'(r)}{r w(r)}. \end{aligned}$$

They play an essential role in the asymptotic behavior:  $\zeta$  describes the behavior of  $w'/w$  and  $\sigma$  is the slope in the phase plane  $(y, Y)$ . They satisfy the equations

$$(2-9) \quad \zeta' = \zeta(\zeta - \eta) + \frac{\varepsilon}{p - 1} |\zeta y|^{2-p} (\alpha - \zeta) = \zeta \left( \zeta - \eta + \frac{\varepsilon(\alpha - \zeta)}{(p - 1)\sigma} \right),$$

$$(2-10) \quad \sigma' = \varepsilon(\alpha - N) + (|\sigma y|^{2-p} \sigma - N)(\sigma - \varepsilon) = \varepsilon(\alpha - N) + (\zeta - N)(\sigma - \varepsilon).$$

**Note.** Since (S) is autonomous, for any solution  $w$  of  $(E_w)$  of the problem, all the functions  $w_\xi(r) = \xi^\delta w(\xi r)$ ,  $\xi > 0$ , are also solutions. From uniqueness, all regular solutions are completely described from one of them:  $w(r, a) = a w(a^{1/\delta} r, 1)$ ; thus they present the same behavior at infinity.

System (S) will be studied by using phase plane techniques, which was not done in [Qi and Wang 1999], and gives our main results. The set of trajectories of system (S) in the phase plane  $(y, Y)$  is symmetric with respect to  $(0, 0)$ . We define

$$(2-11) \quad \mathcal{M} = \{(y, Y) \in \mathbb{R}^2 : |Y|^{(2-p)/(p-1)} Y = \delta y\},$$

which is the set of the extremal points of  $y$ . We denote the four quadrants by

$$\mathfrak{Q}_1 = (0, \infty) \times (0, \infty), \quad \mathfrak{Q}_2 = (-\infty, 0) \times (0, \infty), \quad \mathfrak{Q}_3 = -\mathfrak{Q}_1, \quad \mathfrak{Q}_4 = -\mathfrak{Q}_2.$$

**Remarks 2.1.** (i) The field at any point  $(\xi, 0)$  with  $\xi > 0$  satisfies  $y' = -\xi^{1/(p-1)} < 0$ , and so points toward  $\mathfrak{Q}_2$ . The field at any point  $(\varphi, 0)$  with  $\varphi > 0$  satisfies  $Y' = \varepsilon\alpha\varphi$ , and so points toward  $\mathfrak{Q}_1$  if  $\varepsilon\alpha > 0$  and toward  $\mathfrak{Q}_4$  if  $\varepsilon\alpha < 0$ .

(ii) The pair  $(y, Y)$  defined by (2-7) is related to  $J_N$  by the identity

$$(2-12) \quad J_N(r) = r^{N-\delta}(y(\tau) - \varepsilon Y(\tau)), \quad \tau = \ln r,$$

and the formulae (2-2) can be recovered from the relations

$$(2-13) \quad \begin{aligned} (y - \varepsilon Y)' &= (\delta - \alpha)y + \varepsilon(N - \delta)Y = (\delta - \alpha)(y - \varepsilon Y) + \varepsilon(N - \alpha)Y \\ &= (\delta - N)(y - \varepsilon Y) + (N - \alpha)y. \end{aligned}$$

(iii) In the sequel the monotonicity of the functions  $y_d, Y_d$ , in particular  $y, Y, \zeta$  and  $\sigma$  plays an important role. At any extremal point  $\tau$ , these functions satisfy

$$(2-14) \quad y_d''(\tau) = y_d(\tau) \left( d(\eta - d) - \frac{\varepsilon(\alpha - d)}{p-1} e^{((p-2)d+p)\tau} |dy_d(\tau)|^{2-p} \right),$$

$$(2-15) \quad \begin{aligned} Y_d''(\tau) &= Y_d(\tau) \left( (p-1)^2(\eta - d)(p' + d) \right. \\ &\quad \left. - \varepsilon(\alpha - d) e^{((p-2)d+p)\tau} |Y_d(\tau)|^{(2-p)/(p-1)} \right), \end{aligned}$$

$$(2-16) \quad \begin{aligned} (p-1)y''(\tau) &= \delta^{2-p}y(\tau) \left( \delta^{p-1}(N - \delta) - \varepsilon(\alpha - \delta) |y(\tau)|^{2-p} \right) \\ &= -|Y(\tau)|^{(2-p)/(p-1)} Y'(\tau), \end{aligned}$$

$$(2-17) \quad Y''(\tau) = Y(\tau) \left( \delta(N - \delta) - \varepsilon(\alpha - \delta) |Y(\tau)|^{(2-p)/(p-1)} \right) = \varepsilon\alpha y'(\tau),$$

$$(2-18) \quad \begin{aligned} (p-1)\zeta''(\tau) &= \varepsilon(2-p) \left( (\alpha - \zeta) |\zeta|^{2-p} |y|^{-p} y y' \right) (\tau) \\ &= \varepsilon(2-p) \left( (\alpha - \zeta) (\delta - \zeta) |\zeta y|^{2-p} \right) (\tau), \end{aligned}$$

$$(2-19) \quad \begin{aligned} (p-1)\sigma''(\tau) &= (2-p) \left( (\sigma - \varepsilon) |\sigma|^{(2-p)/(p-1)} Y |y|^{(4-3p)/(p-1)} y' \right) (\tau) \\ &= \zeta'(\tau) (\sigma(\tau) - \varepsilon). \end{aligned}$$

**Energy functions for the system (S).** There is a classical energy function associated to equation ( $E_w$ ):

$$(2-20) \quad E(r) = \frac{1}{p'} |w'|^p + \varepsilon \frac{\alpha}{2} w^2,$$

which is nonincreasing when  $\varepsilon = 1$ , since  $E'(r) = -(N - 1)r^{-1}|w'|^p - \varepsilon r w'^2$ . This is not sufficient in our study: we need energy functions adapted to  $y$  and  $Y$ .

Using the ideas of [Bidaut-Véron 1989], we construct two of them by using the Anderson and Leighton formula [1968].

We find a first function  $W$  given by

$$(2-21) \quad W(\tau) = \mathfrak{W}(y(\tau), Y(\tau)), \quad \text{where}$$

$$\mathfrak{W}(y, Y) = \varepsilon \left( \frac{(2\delta - N)\delta^{p-1}}{p} |y|^p + \frac{|Y|^{p'}}{p'} - \delta y Y \right) + \frac{\alpha - \delta}{2} y^2.$$

It satisfies

$$\begin{aligned} W'(\tau) &= \varepsilon(2\delta - N)(\delta y - |Y|^{(2-p)/(p-1)}Y)(|\delta y|^{p-2}\delta y - Y) - (\delta y - |Y|^{(2-p)/(p-1)}Y)^2 \\ &= (\delta y - |Y|^{(2-p)/(p-1)}Y)(|\delta y|^{p-2}\delta y - Y) \\ &\quad \times \left( \varepsilon(2\delta - N) - \frac{\delta y - |Y|^{(2-p)/(p-1)}Y}{|\delta y|^{p-2}\delta y - Y} \right). \end{aligned}$$

When  $\varepsilon(2\delta - N) \leq 0$ , then  $W$  is nonincreasing. When  $\varepsilon(2\delta - N) > 0$ , we consider the curve

$$\mathcal{L} = \{(y, Y) \in \mathbb{R}^2 : H(y, Y) = \varepsilon(2\delta - N)\},$$

where

$$H(y, Y) := \frac{\delta y - |Y|^{(2-p)/(p-1)}Y}{|\delta y|^{p-2}\delta y - Y}$$

and by convention this quotient takes the value  $|\delta y|^{2-p}/(p-1)$  if  $|\delta y|^{p-2}\delta y = Y$ .  $\mathcal{L}$  is a closed curve surrounding  $(0, 0)$ , symmetric with respect to  $(0, 0)$ . Let  $\mathcal{S}_{\mathcal{L}}$  be the domain with boundary  $\mathcal{L}$  and containing  $(0, 0)$ :

$$(2-22) \quad \mathcal{S}_{\mathcal{L}} = \{(y, Y) \in \mathbb{R}^2 : H(y, Y) \leq \varepsilon(2\delta - N)\}.$$

Then  $W'(\tau) \geq 0$  if  $(y(\tau), Y(\tau)) \in \mathcal{S}_{\mathcal{L}}$  and  $W'(\tau) \leq 0$  if  $(y(\tau), Y(\tau)) \notin \mathcal{S}_{\mathcal{L}}$ . Observe that  $\mathcal{S}_{\mathcal{L}}$  is bounded: indeed, for any  $(y, Y) \in \mathbb{R}^2$ ,

$$(2-23) \quad H(y, Y) \geq \frac{1}{2}((\delta y)^{2-p} + |Y|^{(2-p)/(p-1)}).$$

Also  $\mathcal{S}_{\mathcal{L}}$  is connected; more precisely, for any  $(y, Y) \in \mathcal{S}_{\mathcal{L}}$  and any  $\theta \in [0, 1]$ , we have  $(\theta y, \theta^{p-1}Y) \in \mathcal{S}_{\mathcal{L}}$ .

A second function, denoted by  $V$ , is also given by Anderson formula, or by multiplication by  $Y'$  in  $(E_Y)$ : let

$$(2-24) \quad V(\tau) = \mathfrak{V}(Y(\tau), Y'(\tau)), \quad \text{where}$$

$$\mathfrak{V}(Y, Z) = \frac{\varepsilon}{2}(\delta(\delta - N)Y^2 + Y'^2) + \frac{\alpha - \delta}{p'}|Y|^{p'};$$

then

$$V'(\tau) = \left( \varepsilon(2\delta - N) - \frac{1}{p-1}|Y|^{(2-p)/(p-1)} \right) Y'^2.$$

When  $\varepsilon(2\delta - N)$  is not positive,  $V$  is nonincreasing. When it is positive, we have  $V'(\tau) \geq 0$  whenever  $|Y(\tau)| \leq D$ , where

$$(2-25) \quad D = (\varepsilon(2\delta - N)(p - 1))^{(p-1)/(2-p)}.$$

The function  $W$  gives more information on the system, because  $\mathcal{G}_{\mathcal{L}}$  is bounded, whereas the set of zeros of  $V'$  is unbounded.

**Stationary points of system (S).** If  $\alpha = \delta = N$ , system (S) has infinitely many stationary points, given by  $\pm(k, (\delta k)^{p-1})$ ,  $k \geq 0$ . Otherwise, if  $\varepsilon(\delta - N)(\delta - \alpha) \leq 0$ , the system has a unique stationary point  $(0, 0)$ . If  $\varepsilon(\delta - N)(\delta - \alpha) > 0$ , it admits the three stationary points

$$(2-26) \quad (0, 0), \quad M_\ell = (\ell, (\delta\ell)^{p-1}) \in \mathcal{Q}_1, \quad M'_\ell = -M_\ell \in \mathcal{Q}_3,$$

where  $\ell$  is defined in (1-4). In that case, we find again that  $w \equiv \ell r^{-\delta}$  is a particular solution of equation ( $E_w$ ).

**Local behavior at (0, 0).** The linearized problem at  $(0, 0)$  is given by

$$y' = \delta y, \quad Y' = (\delta - N)Y + \varepsilon\alpha y,$$

and has eigenvalues  $\mu_1 = \delta - N$  and  $\mu_2 = \delta$ . Thus  $(0, 0)$  is a saddle point when  $\delta < N$  and a source when  $N < \delta$ . One can choose a basis of eigenvectors  $v_1 = (0, -1)$  and  $v_2 = (N, \varepsilon\alpha)$ .

**Local behavior at  $M_\ell$ .** Setting

$$(2-27) \quad y = \ell + \bar{y}, \quad Y = (\delta\ell)^{p-1} + \bar{Y},$$

system (S) is equivalent in  $\mathcal{Q}_1$  to

$$(2-28) \quad \bar{y}' = \delta\bar{y} - \varepsilon v(\alpha)\bar{Y} - \Psi(\bar{Y}), \quad \bar{Y}' = \varepsilon\alpha\bar{y} + (\delta - N - v(\alpha))\bar{Y} - \varepsilon\Psi(\bar{Y}),$$

where

$$(2-29) \quad v(\alpha) = \frac{\delta(N - \delta)}{(p - 1)(\alpha - \delta)},$$

$$\Psi(\vartheta) = ((\delta\ell)^{p-1} + \vartheta)^{1/(p-1)} - \delta\ell - \frac{(\delta\ell)^{2-p}}{p-1}\vartheta,$$

with  $\vartheta > -(\delta\ell)^{p-1}$ . The linearized problem is given by

$$\bar{y}' = \delta\bar{y} - \varepsilon v(\alpha)\bar{Y}, \quad \bar{Y}' = \varepsilon\alpha\bar{y} + (\delta - N - v(\alpha))\bar{Y}.$$

Its eigenvalues  $\lambda_1 \leq \lambda_2$  are the solutions of the equation

$$(2-30) \quad \lambda^2 - (2\delta - N - v(\alpha))\lambda + p'(N - \delta) = 0.$$

The discriminant  $\Delta$  of this equation is

$$(2-31) \quad \Delta = (2\delta - N - v(\alpha))^2 - 4p'(N - \delta) = (N + v(\alpha))^2 - 4v(\alpha)\alpha.$$

The critical value  $\alpha^*$  of  $\alpha$ , given in (1-11), arises when  $\varepsilon(\delta - N/2) > 0$ :

$$\alpha = \alpha^* \iff \lambda_1 + \lambda_2 = 0.$$

When  $\delta < N$  and  $\varepsilon = 1$ , then  $\delta < \alpha$  and  $M_\ell$  is a sink when  $\delta \leq N/2$  or  $\delta > N/2$  and  $\alpha < \alpha^*$ , and a source when  $\delta > N/2$  and  $\alpha > \alpha^*$ . When  $\delta < N$ , and  $\varepsilon = -1$ , then  $\alpha < \delta$  and  $M_\ell$  is a source when  $\delta \geq N/2$  or  $\delta < N/2$  and  $\alpha > \alpha^*$ , and a sink when  $\delta < N/2$  and  $\alpha < \alpha^*$ . When  $N < \delta$ , then  $M_\ell$  is always a saddle point, but, as we will see later, the value  $\alpha^*$  also plays a role.

More specifically, the sign of  $\alpha^*$  and its position with respect to  $N$  or  $\eta$  play a role. By computation,

$$(2-32) \quad \begin{aligned} \alpha^* &= \frac{p'(\delta^2 - 3\delta + 2N)}{2(2\delta - N)} = \eta + \frac{(\delta - N)^2}{(p - 1)(2\delta - N)} \\ &= N + \frac{(\delta - N)(\delta^2 - (N + 3)\delta + N)}{(2\delta - N)(\delta - 1)}. \end{aligned}$$

Thus, if  $\varepsilon = 1$ , then  $\alpha^* > \eta > 0$  if  $N \geq 2$ ; if  $N = 1$ ,  $\alpha^* > 0$  if  $p > \frac{4}{3}$ . If  $\varepsilon = -1$ , then  $\alpha^* < -p' < 0$ .

Otherwise, when  $\Delta > 0$  a basis of eigenvectors  $u_1 = (-\varepsilon v(\alpha), \lambda_1 - \delta)$ ,  $u_2 = (\varepsilon v(\alpha), \delta - \lambda_2)$  can be chosen. If  $\Delta \geq 0$ , then  $\delta$  is exterior to the roots if  $\varepsilon\alpha > 0$ , and  $\lambda_1 < \delta < \lambda_2$  if  $\varepsilon\alpha < 0$ .

### Existence of solutions of equation $(E_w)$ .

**Theorem 2.2.** (i) *Take  $r_1 > 0$  ( $r_1 \geq 0$  if  $N = 1$ ) and let  $a, a'$  be reals. There exists a unique solution  $w$  of equation  $(E_w)$  in a neighborhood  $\mathcal{V}$  of  $r_1$ , such that  $w \in C^2(\mathcal{V})$  and  $w(r_1) = a$ ,  $w'(r_1) = a'$ . It has a unique extension to a maximal interval of the form*

$$\begin{aligned} (R_w, \infty) \quad \text{with } 0 \leq R_w \quad &\text{if } \varepsilon = 1, \\ (0, S_w) \quad \text{with } S_w \leq \infty \quad &\text{if } \varepsilon = -1. \end{aligned}$$

*If  $0 < R_w$  or  $S_w < \infty$ , as the case may be,  $w$  is monotone near  $R_w$  or  $S_w$  with an infinite limit.*

(ii) *For any  $a \in \mathbb{R}$ , there exists a unique regular solution  $w$  of  $(E_w)$  satisfying (1-10), and*

$$(2-33) \quad \lim_{r \rightarrow 0} |w'|^{p-2} w' / r w = -\varepsilon\alpha / N.$$

(iii) If  $N \geq 2$ , any solution defined near 0 and bounded is regular. If  $N = 1$ , it satisfies  $\lim_{r \rightarrow 0} w' = b \in \mathbb{R}$ , and  $\lim_{r \rightarrow 0} w = a \in \mathbb{R}$ .

*Proof.* (i) Local existence and uniqueness near  $r_1 > 0$  follow directly from Cauchy's theorem applied to equation  $(E_w)$  or to system  $(S)$ , since the map  $\xi \mapsto f_p(\xi) = |\xi|^{(2-p)/(p-1)}\xi$  is of class  $C^1$ . If  $N = 1$ , we can take  $r_1 = 0$ , obtain a local solution in a neighborhood of 0 in  $\mathbb{R}$  and reduce it to  $[0, \infty)$ .

Any local solution around  $r_1$  has a unique extension to a maximal interval  $(R_w, S_w)$ . Suppose that  $0 < R_w$  (or  $S_w < \infty$ ) and that  $w$  is oscillating around 0 near  $R_w$  (or  $S_w$ ). Making the substitution (2-3), with  $d \neq 0$ , if  $\tau$  is a maximal point of  $|y_d|$ , we see that (2-14) holds. If we take  $d$  such that  $\varepsilon(d - \alpha) > 0$ , the sequence  $(y_d(\tau))$  stays bounded, since the exponential has a positive limit; for that reason  $y_d$  stays bounded,  $w$  is bounded near  $R_w$  (or  $S_w$ ) and then also  $J'_N$ ,  $J_N$  and  $w'$ , which is contradictory. Thus  $w$  keeps a constant sign, for example  $w > 0$ , near  $R_w$  (or  $S_w$ ). At each extremal point  $r$  such that  $w(r) > 0$ , we find  $(|w'|^{p-2}w')'(r) = -\varepsilon\alpha w(r)$ ; thus  $r$  is unique since  $\alpha \neq 0$ . Thus  $w$  is strictly monotone near  $R_w$  (or  $S_w$ ), and  $w$  and  $|w'|$  tend to  $\infty$ .

First suppose  $\varepsilon = 1$ . We show that  $S_w = \infty$ . This is easy when  $\alpha > 0$ : since  $E$  is nondecreasing,  $w$  and  $w'$  are bounded for  $r > r_1$ . Assume  $\alpha < 0$  and  $S_w < \infty$ . Then for example  $w$  is positive near  $S_w$ , nondecreasing, and  $\lim_{r \rightarrow S_w} w = \infty$ . Then  $J_\alpha$  is nonincreasing and nonnegative near  $S_w$ ; hence again  $w$  and  $w'$  are bounded, which is contradictory.

Next suppose  $\varepsilon = -1$ . If  $R_w > 0$ , for example,  $w$  is positive and nonincreasing and  $\lim_{r \rightarrow R_w} w = \infty$ . Then either  $\alpha < N$  and  $J_N$  is nonnegative and nondecreasing near  $R_w$ , and thus bounded, or  $\alpha \geq N$  and  $J_\alpha$  is nonnegative and nondecreasing near  $R_w$ , and still bounded. In either case we reach a contradiction, then  $R_w = 0$ .

(ii) By symmetry we can suppose  $a \geq 0$ . Let  $\rho > 0$ . By (2-1) and (2-2), any regular solution  $w$  on  $[0, \rho]$  satisfies

$$(2-34) \quad \begin{aligned} w(r) &= a - \varepsilon \int_0^r f_p(sT(w)) ds, \\ T(w)(r) &= w(r) + (\alpha - N) \int_0^1 \theta^{N-1} w(r\theta) d\theta. \end{aligned}$$

Conversely, any function  $w \in C^0([0, \rho])$  that solves (2-34) satisfies  $w \in C^1((0, \rho])$  and  $|w'|^{p-2}w'(r) = rT(w)$ ; hence  $|w'|^{p-2}w' \in C^1((0, \rho])$  and  $w$  satisfies  $(E_w)$  in  $(0, \rho]$ . And  $\lim_{r \rightarrow 0} rT(w) = 0$ , thus  $w \in C^1([0, \rho])$  and  $|w'|^{p-2}w' \in C^1([0, \rho])$ . Then  $w$  satisfies  $(E_w)$  in  $[0, \rho]$  and  $w'(0) = 0$ . From  $(E_w)$ , we have

$$\lim_{r \rightarrow 0} |w'|^{p-2}w'/rw = -\varepsilon\alpha/N,$$

and therefore  $w - a = O(r^{p'})$  near 0. We look for  $w$  of the form  $a + r^{p'}\zeta(r)$ , with

$$\zeta \in \mathcal{B}_{\rho, M} = \{\zeta \in C^0([0, \rho]) : \|\zeta\|_{C^0([0, \rho])} = \max_{r \in [0, \rho]} |\zeta(r)| \leq M\}.$$

We are led to the problem  $\zeta = \Theta(\zeta)$ , where

$$\begin{aligned} \Theta(\zeta)(r) &= -\varepsilon \int_0^1 \theta^{1/(p-1)} f_p(T(a + (r\theta)^{p'}\zeta(r\theta))) d\theta \\ &= -\varepsilon \int_0^1 \theta^{1/(p-1)} f_p\left(\frac{\alpha a}{N} + T((r\theta)^{p'}\zeta(r\theta))\right) d\theta. \end{aligned}$$

Taking for example  $M = (|\alpha|a)^{1/(p-1)}$ , it follows that  $\Theta$  is a strict contraction from  $\mathcal{B}_{\rho, M}$  into itself for  $\rho$  small enough, hence existence and uniqueness hold in  $[0, \rho]$ .

(iii) If  $w$  is defined in  $(0, \rho)$  and bounded, then  $J'_N$  is integrable. Set

$$l = \lim_{r \rightarrow 0} J_N(r).$$

Then  $|w'|^{p-2}w' = \varepsilon l r^{1-N}(1 + o(1))$ . If  $N \geq 2$ , this implies  $l = 0$ ; thus from above,  $w$  is regular. If  $N = 1$ , then  $\lim_{r \rightarrow 0} w' = b \in \mathbb{R}$ , and  $\lim_{r \rightarrow 0} w = a \in \mathbb{R}$ . □

**Definition.** Suppose  $p > 1$ . Let  $\mathcal{T}_r$  be the trajectory in the plane  $(y, Y)$  (see (2-7)) starting from  $(0, 0)$  at  $-\infty$ , with slope  $\varepsilon\alpha/N$  and  $y > 0$  near time  $-\infty$ . Its opposite  $-\mathcal{T}_r$  is also a trajectory with the same properties (except that  $y < 0$ ). Both are called *regular trajectories*. In this situation we say that  $y$  is regular. Observe that  $\mathcal{T}_r$  starts in  $\mathcal{Q}_1$  if  $\varepsilon\alpha > 0$ , and in  $\mathcal{Q}_4$  if  $\varepsilon\alpha < 0$ .

**Remark 2.3.** Let  $w$  be any solution of  $(E_w)$  such that  $w > 0$  on some interval  $I$ .

- (i) The function  $w$  has at most one extremal point on  $I$ , since  $(|w'|^{p-2}w')' = -\varepsilon\alpha w$ , and this point is a maximum if  $\varepsilon\alpha > 0$  and a minimum if  $\varepsilon\alpha < 0$ .
- (ii) From (2-33), if  $w$  is regular and  $w > 0$  in  $(0, r_1)$ ,  $r_1 \leq \infty$ , then  $w' < 0$  in  $(0, r_1)$  when  $\varepsilon\alpha > 0$ ; thus  $\mathcal{T}_r$  is in  $\mathcal{Q}_1$ . And  $w' > 0$  in  $(0, r_1)$  when  $\varepsilon\alpha < 0$ ; hence  $\mathcal{T}_r$  is in  $\mathcal{Q}_3$  in  $(-\infty, \ln r_1)$ .

**Remark 2.4.** In the case  $\delta \neq N$ , we can give a shorter proof of Theorem 2.2(ii). Indeed,  $(0, 0)$  is either a source or a saddle point. Thus there exists precisely one trajectory starting from  $(0, 0)$  at  $-\infty$ , with  $y > 0$ , with slope  $\varepsilon\alpha/N$ . The corresponding solutions are regular: the slope  $\sigma$  defined in (2-8) satisfies  $\lim_{\tau \rightarrow -\infty} \sigma = \varepsilon\alpha/N$ . Thus  $\lim_{r \rightarrow 0} |w'|^{p-2}w'/rw = -\varepsilon\alpha/N$ , implying that  $w^{(2-p)/(p-1)}$  has a limit  $a > 0$ . Since  $\lim_{r \rightarrow 0} w' = 0$ , this function  $w$  satisfies (1-10), and any  $a$  is obtained by scaling.

**Notation.** For any point  $P_0 = (y_0, Y_0) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ , the unique trajectory in the phase plane  $(y, Y)$  going through  $P_0$  is denoted by  $\mathcal{T}_{[P_0]}$ . Notice that  $\mathcal{T}_{[-P_0]} = -\mathcal{T}_{[P_0]}$ , from the symmetry of system  $(S)$ .

**First sign properties.**

**Proposition 2.5.** *Let  $w \not\equiv 0$  be any solution of  $(E_w)$ .*

- (i) *If  $\varepsilon = 1$  and  $\alpha \leq \max(N, \eta)$ , then  $w$  has at most one zero, and no zero if  $w$  is regular.*
- (ii) *If  $\varepsilon = 1$  and  $N < \min(\delta, \alpha)$  and  $w$  is regular, then  $w$  has at least one zero.*
- (iii) *If  $\varepsilon = -1$  and  $\alpha \geq \min(0, \eta)$ , then  $w$  has at most one zero. If  $\alpha > 0$  and  $w$  is regular, then it has no zero.*
- (iv) *If  $\varepsilon = -1$  and  $-p' \leq \alpha < \min(0, \eta)$ , then  $w'$  has at most one zero; consequently  $w$  has at most two zeros, and at most one if  $w$  is regular.*

*Proof.* (i) Let  $\varepsilon = 1$ . Take two consecutive zeros  $\rho_0 < \rho_1$  of  $w$ , with  $w > 0$  on  $(\rho_0, \rho_1)$ , so  $w'(\rho_1) < 0 < w'(\rho_0)$ . If  $\alpha \leq N$ , we find, using the function  $J_N$  of (2-1),

$$J_N(\rho_1) - J_N(\rho_0) = -\rho_1^{N-1} |w'(\rho_1)|^{p-2} - \rho_0^{N-1} w'(\rho_0)^{p-1} = (N - \alpha) \int_{\rho_0}^{\rho_1} s^{N-1} w \, ds,$$

which is contradictory; thus  $w$  has at most one zero. If  $w$  is regular with  $w(0) > 0$  and  $\rho_1$  is a first zero, then

$$J_N(\rho_1) = -\rho_1^{N-1} |w'(\rho_1)|^{p-1} = (N - \alpha) \int_0^{\rho_1} s^{N-1} w \, ds \geq 0,$$

again a contradiction. Next suppose  $0 < \alpha \leq \eta$  and use the substitution (2-3), with  $d = \alpha$ . Then  $y_\alpha$  has at most one zero: indeed, if  $y_\alpha$  has a maximal point  $\tau$  where it is positive, and is not constant, then from (2-14),

$$(2-35) \quad y''_\alpha(\tau) = \alpha(\eta - \alpha)y_d(\tau);$$

hence  $y''_\alpha(\tau) < 0$ , which is impossible. In the same way the regular solution satisfies  $\lim_{\tau \rightarrow -\infty} y_\alpha = 0$  since  $\alpha > 0$ , and  $y_\alpha$  has no maximal point; thus  $y_\alpha$  is positive and increasing.

(ii) Let  $\varepsilon = 1$  and  $w > 0$  on  $[0, \infty)$ . If  $N < \alpha$ , then  $J_N(r) = (N - \alpha) \int_0^r s^{N-1} w \, ds < 0$ . The function  $r \mapsto \delta r^{p'} - w^{(p-2)/(p-1)}$  is nonincreasing; hence  $w = O(r^{-\delta})$  at  $\infty$ , so  $y$  is bounded at  $\infty$ . For any  $r \geq 1$ , one gets  $J_N(r) \leq J_N(1) < 0$ , hence  $y(\tau) + |J_N(1)|e^{(\delta-N)\tau} \leq Y(\tau)$  for any  $\tau \geq 0$ , from (2-12). Then  $\lim_{\tau \rightarrow \infty} Y = \infty$ , implying by (S) that  $\lim_{\tau \rightarrow \infty} y' = -\infty$ , which is impossible.

(iii) Let  $\varepsilon = -1$  and  $\alpha \geq \min(\eta, 0)$ . We use again the substitution (2-3) for some  $d \neq 0$ . If  $y_d$  is not constant and has a maximal point where it is positive, then (2-14) holds. Taking  $d \in (0, \min(\alpha, \eta))$  if  $N \geq 2$  and  $\alpha > 0$  and  $d = -1$  if  $N = 1$  and  $\eta = -1 \leq \alpha$ , we reach a contradiction. Now suppose  $w$  is regular and  $\alpha > 0$ . Then  $w' > 0$  near 0, from Theorem 2.2, and as long as  $w$  stays positive, any extremal point  $r$  is a strict minimum; thus in fact  $w' > 0$  on  $[0, S_w)$ .

(iv) Let  $\varepsilon = -1$  and  $-p' \leq \alpha < \min(0, \eta)$ . Suppose that  $w'$  has two consecutive zeros  $\rho_1 < \rho_2$ , and use (2-3) again with  $d = \alpha$ . If the function  $Y_\alpha$  is not constant and has a maximal point  $\tau$  where it is positive, we get from (2-15)

$$(2-36) \quad Y''_\alpha(\tau) = (p - 1)^2(\eta - \alpha)(p' + \alpha)Y_\alpha(\tau);$$

thus  $Y''_\alpha(\tau) < 0$ , and  $Y_\alpha$  has at most one zero. Next consider regular solutions: they satisfy  $Y_\alpha = e^{(\alpha(p-1)+p)\tau}(|\alpha|a/N)(1 + o(1))$  near  $-\infty$ , by Theorem 2.2 and (2-3). Thus  $\lim_{\tau \rightarrow -\infty} Y_\alpha = 0$ ; as above  $Y_\alpha$  cannot have any extremal point, so  $Y_\alpha$  is positive and increasing. Then  $w' < 0$  from (2-3), and  $w$  has at most one zero.  $\square$

**Remark 2.6.** From (2-35) and (2-36) we see that if  $0 < \alpha \leq \eta$  then  $y_\alpha$  has only minimal points on any set where it is positive, and the same conclusion holds for  $Y_\alpha$  when  $-p' \leq \alpha \leq \min(\eta, 0)$ .

**Proposition 2.7.** *Let  $y$  be any solution of  $(E_y)$ , linked with  $w$  by (2-7), and having constant sign in a semi-interval around the point  $\ln R_w$  or  $\ln S_w$ .*

- (i) *If  $y$  is not strictly monotone near that same point, then  $R_w = 0$  or  $S_w = \infty$ . If  $y$  is not constant, then either  $\varepsilon = 1$  and  $\delta < N < \alpha$  or  $\varepsilon = -1$  and  $\alpha < \delta < N$ . In any case,  $y$  oscillates around  $\ell$ .*
- (ii) *If  $y$  is strictly monotone near  $\ln R_w$  or  $\ln S_w$ , then also  $Y, \zeta, \sigma$  are monotone near the same point.*

*Proof.* Let  $s = R_w$  or  $S_w$ , and suppose that  $y$  has constant sign near  $s$ . Then so does  $Y$ , by Remark 2.3.

(i) At each point  $\tau$  where  $y'(\tau) = 0$ , we have  $y''(\tau) \neq 0$ , and (2-16) holds with  $y > 0$ . Suppose that  $y$  is not strictly monotone near  $s$ . There exists a strictly monotone sequence  $(\tau_n)$  converging to  $s$  and such that  $y'(\tau_n) = 0$ ,  $y''(\tau_{2n}) > 0$ ,  $y''(\tau_{2n+1}) < 0$ . Then either  $\varepsilon = 1$  and  $\delta < \min(\alpha, N)$ , or  $\varepsilon = -1$  and  $\alpha < \delta < N$ ; and  $y(\tau_{2n}) < \ell < y(\tau_{2n+1})$ . This cannot happen if  $s$  is finite, because  $y$  tends to  $\infty$ . It is also impossible when  $\varepsilon = 1$  and  $\alpha \leq N$ ; indeed, there exist at least two points  $\theta_1 < \theta_2$  such that  $y(\theta_1) = y(\theta_2) = \ell$  and  $y \geq \ell$  on  $(\theta_1, \theta_2)$ , with  $y'(\theta_1) > 0 > y'(\theta_2)$ . Then from (S),  $Y(\theta_1) < (\delta\ell)^{p-1} < Y(\theta_2)$ . And from (2-13),  $(e^{(N-\delta)\tau}(y - Y))' = (N - \alpha)e^{(N-\delta)\tau}y$ ; and the constant  $(\ell, (\delta\ell)^{p-1})$  is also a solution of (S), hence

$$(e^{(N-\delta)\tau}(y - \ell - Y + (\delta\ell)^{p-1}))' = (N - \alpha)e^{(N-\delta)\tau}(y - \ell) \geq 0$$

on  $(\theta_1, \theta_2)$ . A contradiction follows by integration on this interval.

(ii) Suppose  $y$  strictly monotone near  $s$ . At any extremal point  $\tau$  of  $Y$ , we find  $Y''(\tau) = \varepsilon\alpha y'(\tau)$  from (2-17); hence  $y'(\tau) \neq 0$ , and  $Y''(\tau)$  has constant sign; thus  $\tau$  is unique, and  $Y$  is strictly monotone near  $s$ .

Next consider the function  $\zeta$  satisfying (2–9). If there exists  $\tau_0$  such that  $\zeta(\tau_0) = \alpha$ , then  $\zeta'(\tau_0) = \alpha(\alpha - \eta)$ . If  $\alpha \neq \eta$ , then  $\tau_0$  is unique, so  $\alpha - \zeta$  has a constant sign near  $s$ . Then also  $\zeta''(\tau)$  has constant sign at any extremal point  $\tau$  of  $\zeta$ , from (2–18). Then  $\zeta$  is strictly monotone near  $s$ . If  $\alpha = \eta$ , then  $\zeta \equiv \alpha$ .

Finally consider  $\sigma$ , which satisfies (2–10). At each point  $\tau$  such that  $\sigma'(\tau) = 0$ , (2–19) holds and  $Y$  has a constant sign. If there exists  $\tau_0$  such that  $\sigma(\tau_0) = \varepsilon$ , then  $\sigma'(\tau_0) = \varepsilon(\alpha - N)$ . If  $\alpha \neq N$ , then  $\tau_0$  is unique, and  $\sigma - \varepsilon$  has constant sign near  $s$ . Thus  $\sigma''(\tau)$  has constant sign at any extremal point  $\tau$  of  $\sigma$ , by (2–19), since  $Y$  has constant sign near  $s$ . If  $\alpha = N$ , then  $\sigma \equiv \varepsilon$ .  $\square$

**Behavior of  $w$  near 0 or  $\infty$ .** Here we suppose  $w$  is defined near 0 or  $\infty$ , which means the function  $y$  of (2–7) is defined near  $\pm\infty$ . We study the behavior of  $y$  and then return to  $w$ . First we suppose  $y$  monotone, so we can assume  $y > 0$  near  $\pm\infty$ . We do not look for a priori estimates, which could be obtained by successive approximations as in [Bidaut-Véron 2006a]. Our method is based on monotonicity and L'Hospital's rule, which is much more rapid and efficient.

**Proposition 2.8.** *Let  $(y, Y)$  be any solution of (S) such that  $y$  is strictly monotone and  $y > 0$  near  $s = \pm\infty$ . Then  $\zeta$  has a finite limit  $\lambda$  near  $s$ , which is equal to 0,  $\alpha$ ,  $\eta$ ,  $\delta$ . More precisely, we are in one of the following cases:*

- (i)  $(y, Y)$  converges to a stationary point different from  $(0, 0)$ . Then  $\lambda = \delta$ , and  $\varepsilon(\delta - N)(\delta - \alpha) > 0$  or  $\alpha = \delta = N$ .
- (ii)  $(y, Y)$  converges to  $(0, 0)$ . Then
  - either  $\lambda = 0$ ,  $s = -\infty$ , and  $y$  is regular, or  $N = 1$ ;
  - or  $\lambda = \eta$ ; then either  $(s = \infty, \delta < N)$  or  $(s = \infty, \delta = N, \varepsilon(\alpha - N) < 0)$  or  $(s = -\infty, N < \delta)$  or  $(s = -\infty, \delta = N, \varepsilon(\alpha - N) > 0)$ .
- (iii)  $\lim_{\tau \rightarrow s} y = \infty$  and  $\lambda = \alpha$ . Then either  $(s = \infty, \alpha < \delta)$  or  $(s = \infty, \alpha = \delta, \varepsilon(\delta - N) < 0)$  or  $(s = -\infty, \delta < \alpha)$  or  $(s = -\infty, \alpha = \delta, \varepsilon(\delta - N) > 0)$ .

*Proof.* From Proposition 2.7, the functions  $y, Y, \sigma, \zeta$  are monotone; hence  $\zeta$  has a limit  $\lambda \in [-\infty, \infty]$  and  $\sigma$  has a limit  $\mu \in [-\infty, \infty]$ , and  $(y, Y)$  converges to a stationary point, or  $\lim y = \infty$ . Then  $\lim |Y| = \infty$ , since  $\alpha \neq 0$  from system (S). To apply the L'Hospital's rule, we consider the two quotients

$$(2-37) \quad \frac{Y'}{y'} = \frac{(\delta - N)\sigma + \varepsilon(\alpha - \zeta)}{\delta - \zeta}$$

and

$$(2-38) \quad \begin{aligned} \frac{(|Y|^{(2-p)/(p-1)}Y)'}{y'} &= \frac{\zeta(\delta - N + \varepsilon(\alpha - \zeta))/\sigma}{(p - 1)(\delta - \zeta)} \\ &= \frac{\zeta(\delta - N) + \varepsilon(\alpha - \zeta)|\zeta y|^{2-p}}{(p - 1)(\delta - \zeta)}. \end{aligned}$$

(i) *First case:*  $\varepsilon(\delta - N)(\delta - \alpha) > 0$  and  $(y, Y)$  converges to the point  $M_\ell$  defined by (2–26). Then obviously  $\lambda = \delta$ ; or  $\alpha = \delta = N$  and  $\lim_{\tau \rightarrow s} y = k > 0$ ; then  $\lim_{\tau \rightarrow s} Y = (\delta k)^{p-1}$ , so  $\lambda = \delta$ .

(ii) *Second case:*  $(y, Y)$  converges to  $(0, 0)$ . Then  $\lambda$  is finite; indeed, if  $\lambda = \pm\infty$ , the quotient (2–38) converges to  $(N - \delta)/(p - 1)$ , because  $|\zeta y| = |Y|^{1/(p-1)} = o(1)$ ; thus  $\zeta = |Y|^{(2-p)/(p-1)} Y/y$  has the same limit, from L'Hospital's rule, which is contradictory.

We next consider  $N$  in relation to  $\delta$ . If  $N < \delta$ , then  $(0, 0)$  is a source, thus  $s = -\infty$ . Using the eigenvectors, either  $\mu = \varepsilon\alpha/N$ , then  $|\zeta|^{p-1} = |\mu|y^{2-p}(1 + o(1))$ , thus  $\lambda = 0$  and  $w$  is regular, from Remark 2.4. Or  $\mu = \pm\infty$ ; then  $\lambda = \lambda(\delta - N)/(p - 1)(\delta - \zeta)$  from (2–38), thus  $\lambda = 0$  or  $\lambda = \eta$ . If  $\lambda = 0$ , then  $\zeta'/\zeta$  converges to  $-\eta$  from (2–9), and  $s = -\infty$ , thus necessarily  $\eta < 0$ , which means  $N = 1$ .

If  $\delta < N$  (so  $N \geq 2$ ), then  $(0, 0)$  is a saddle point. Thus either  $s = -\infty$  and  $\mu = \varepsilon\alpha/N$ ,  $\lambda = 0$  and  $w$  is regular. Or  $s = \infty$ ,  $\mu = \pm\infty$ , and as above,  $\lambda = 0$  or  $\lambda = \eta$ . Now if  $\lambda = 0$  the quotient (2–37) converges to  $\mp\infty$ , which is contradictory. Thus  $\lambda = \eta$ .

If  $\delta = N$  (so  $N \geq 2$ ), either  $\lambda = 0$ , so  $y' > 0$ ,  $s = -\infty$ , and  $\mu = \varepsilon\alpha/N$  by (2–38); or else  $\lambda > 0$ , in which case  $\lambda = N = \eta$  from (2–38). Moreover if  $s = \infty$ , then  $\varepsilon(\alpha - N) < 0$ ; if  $s = -\infty$ , then  $\varepsilon(\alpha - N) > 0$ . Indeed  $(\varepsilon y - Y)' = \varepsilon(N - \alpha)y$  and  $y - \varepsilon Y$  converges to 0; thus if  $s = \infty$  and  $\varepsilon(N - \alpha) \geq 0$ , or  $s = -\infty$  and  $\varepsilon(N - \alpha) \leq 0$ , then  $\mu \leq \varepsilon$ , but  $\mu = \infty$ , we reach again a contradiction.

(iii) *Third case:*  $y$  tends to  $\infty$ . If  $s = \infty$ , then  $y' > 0$ , thus  $\zeta < \delta$ ; if  $s = -\infty$ , then  $\zeta > \delta$ . If  $\lambda = \pm\infty$ , then the quotient (2–38) converges to  $\varepsilon\infty$ ; thus  $\lambda = \varepsilon\infty$  and  $s = -\varepsilon\infty$ . In any case,  $\zeta' < 0$ , so  $|\mu| \leq 1/(p - 1)$  by (2–9), and  $\mu = \varepsilon$  by (2–37); thus  $Y' = -\varepsilon|Y|^{(2-p)/(p-1)} Y(1 + o(1))$ , and we reach a contradiction by integration. Thus  $\lambda$  is finite; moreover  $\lambda \neq 0$  for otherwise we would have  $\mu = 0$ , seeing that  $\sigma = |\zeta y|^{p-2}\zeta$ ; but  $\mu = \alpha/\delta$  by (2–37).

If  $\alpha \neq \delta$ , then  $\lambda = \alpha$  or  $\delta$ , by (2–38). In turn  $\sigma = |\lambda y|^{p-2}\lambda(1 + o(1))$ , thus  $\mu = 0$ . From (2–37), necessarily  $\lambda = \alpha$ . And if  $s = \infty$ , then  $y' > 0$ , thus  $\zeta < \delta$ , thus  $\alpha < \delta$ . If  $s = -\infty$ , then similarly  $\alpha > \delta$ .

If  $\alpha = \delta$ , then  $\lambda = \alpha = \delta \neq N$ , and  $\varepsilon(\delta - N)(\delta - \zeta) < 0$  from (2–38); thus if  $s = \infty$ , then  $\varepsilon(\delta - N) < 0$  since  $\zeta < \delta$ ; if  $s = -\infty$ , then  $\varepsilon(\delta - N) > 0$ .  $\square$

Next we improve Proposition 2.8 by giving a precise behavior of  $w$  in any case:

**Proposition 2.9.** *We keep the assumptions of Proposition 2.8.*

(i) *If  $\lambda = \alpha \neq \delta$ , then  $\lim r^\alpha w = L > 0$  (near 0, or  $\infty$ ).*

(ii) *If  $\lambda = \eta > 0$ ,  $\eta \neq N$ , then  $\lim r^\eta w = c > 0$ .*

(iii) If  $\lambda = \alpha = \delta \neq N$ , then

$$(2-39) \quad \lim r^\delta (\ln r)^{-1/(2-p)} w = \kappa := ((2-p)\delta^{p-1}|N-\delta|)^{1/(2-p)}.$$

(iv) If  $\lambda = \eta = N = \delta \neq \alpha$ , then

$$(2-40) \quad \lim r^N (\ln r)^{(N+1)/2} w = \rho := \frac{1}{N} \left( \frac{N(N-1)}{2|\alpha-N|} \right)^{(N+1)/2}.$$

(v) If  $N = 1$ ,  $\lambda = \eta = -1$  or  $\lambda = 0$  (near 0) then

$$(2-41) \quad \lim_{r \rightarrow 0} w = a \in \mathbb{R}, \quad \lim_{r \rightarrow 0} w' = b;$$

and  $b \neq 0$ ; moreover,  $a = 0$  (hence  $b > 0$ ) if and only if  $\lambda = -1$ .

*Proof.* (i) Let  $\lambda = \alpha \neq \delta$ . From (2-8) we have  $rw'(r) = -\alpha w(r)(1 + O(1))$ . Next we apply Proposition 2.8, and are led to two cases:

If  $s = \infty$  and  $\alpha < \delta$ , then for any  $\gamma > 0$  we have  $w = O(r^{-\alpha+\gamma})$  and  $1/w = O(r^{\alpha+\gamma})$  near  $\infty$  and  $w' = O(r^{-\alpha-1+\gamma})$ . Then  $J'_\alpha(r) = O(r^{\alpha(2-p)-p-1+\gamma})$ , so  $J'_\alpha$  is integrable, hence  $J_\alpha$  has a limit  $L$ , and  $\lim r^\alpha w = L$ , seeing that  $J_\alpha(r) = r^\alpha w(1 + o(1))$ . If  $L = 0$ , then  $r^\alpha w = O(r^{\alpha(2-p)-p+\gamma})$ , which contradicts the estimate of  $1/w = O(r^{\alpha+\gamma})$  for  $\gamma$  small enough. Thus  $L > 0$ .

Otherwise, we have  $s = -\infty$  and  $\delta < \alpha$ ; hence  $\lim_{\tau \rightarrow s} y = \infty$ ,  $w = O(r^{-\alpha-\gamma})$ ,  $1/w = O(r^{\alpha-\nu})$ ,  $w' = O(r^{-\alpha-1-\gamma})$  near 0, and  $J'_\alpha(r) = O(r^{\alpha(2-p)-p-1-\gamma})$ . Thus  $J'_\alpha$  is still integrable; hence  $\lim r^\alpha w = L \geq 0$ . If  $L = 0$ , then  $r^\alpha w = O(r^{\alpha(2-p)-p-\gamma})$ , which contradicts the estimate of  $1/w$ . Therefore we again obtain  $L > 0$ .

(ii) Let  $\lambda = \eta > 0$ ,  $\eta \neq N$ . From Proposition 2.8, either  $s = \infty$ ,  $\delta < N$  or  $s = -\infty$ ,  $N < \delta$ . As above we get  $w = O(r^{-\eta\pm\gamma})$  and  $1/w = O(r^{\eta\pm\gamma})$  near  $\infty$  or 0. Here we make the substitution (2-3) with  $d = \eta$ . We find  $y_\eta = O(e^{\pm\gamma\tau})$ ,  $1/y_\eta = O(e^{\pm\gamma\tau})$ ,  $y'_\eta = O(e^{\pm\gamma\tau})$ , thus  $Y_\eta = O(e^{\pm\gamma\tau})$ , and from (2-4),  $Y'_\eta = O(e^{\pm\gamma\tau})$ . Substituting in (2-4), we deduce  $Y'_\eta = O(e^{(2-p)((\delta-\eta)\pm\gamma)\tau})$ . When  $s = \infty$ , then  $\delta < \eta$ , when  $s = -\infty$ , then  $\delta > \eta$  from (1-2). In any case,  $Y'_\eta$  is integrable, hence  $Y_\eta$  has a limit  $k$ , and  $Y_\eta - k = O(e^{(2-p)((\delta-\eta)\pm\gamma)\tau})$ . Now  $(e^{-\eta\tau} y_\eta)' = -e^{-\eta\tau} Y_\eta^{1/(p-1)}$ , thus  $y_\eta$  has a limit  $c = k^{1/(p-1)}/\eta$ ; in other words,  $\lim r^\eta w = c$ . If  $c = 0$ , then  $Y_\eta = O(e^{(2-p)((\delta-\eta)\pm\gamma)\tau})$ ,  $y_\eta = O(e^{((2-p)((\delta-\eta)\pm\gamma)/(p-1))\tau})$ , which contradicts  $1/y_\eta = O(e^{\gamma\tau})$  for  $\gamma$  small.

(iii) Now suppose  $\lambda = \alpha = \delta \neq N$ . Then either  $s = \infty$  and  $\varepsilon(\delta - N) < 0$  or  $s = -\infty$  and  $\varepsilon(\delta - N) > 0$ ; moreover,  $\lim_{\tau \rightarrow s} y = \infty$ . Then  $Y = (\delta y)^{p-1}(1 + o(1))$ , and  $\mu = 0$ ; hence  $y - \varepsilon Y = y(1 + o(1))$ , and from (2-13),

$$(y - \varepsilon Y)' = \varepsilon(N - \delta)Y = \varepsilon(N - \delta)\delta^{p-1}(y - \varepsilon Y)^{p-1}(1 + o(1)).$$

Then  $y = (|N - \delta|\delta^{p-1}(2-p)|\tau|)^{1/(2-p)}(1 + o(1))$ , which is equivalent to (2-39) by (2-7).

(iv) Let  $\lambda = \eta = N = \delta \neq \alpha$ . Then either  $s = \infty$  and  $\varepsilon(\alpha - N) < 0$  or  $s = -\infty$  and  $\varepsilon(\alpha - N) > 0$ ; moreover,  $\lim_{\tau \rightarrow s} y = 0$ . Then  $Y = (Ny)^{p-1}(1 + o(1))$  and  $\mu = \infty$ , so  $Y - \varepsilon y = Y(1 + o(1))$ , and from (2-13) we have

$$(Y - \varepsilon y)' = \varepsilon(\alpha - N)y = \varepsilon(\alpha - N)N^{-1}(Y - \varepsilon y)^{1/(p-1)}(1 + o(1)).$$

Hence  $y = c|\tau|^{-(N+1)/2}(1 + o(1))$  with  $c = N^{-1}(N(N-1)/2|\alpha - N|)^{(N+1)/2}$ , and (2-40) follows from (2-7).

(v) Let  $\lambda = 0$ . Then also  $rw' = o(w)$ ; thus by integration we get  $w + |w'| = O(r^{-k})$  for any  $k > 0$ . Then  $J_1'$  is integrable, so  $J_1$  has a limit at 0, and  $\lim_{r \rightarrow 0} rw = 0$ . Therefore  $\lim_{r \rightarrow 0} w' = b \in \mathbb{R}$  and  $\lim_{r \rightarrow 0} w = a \geq 0$ . Then  $b \neq 0$ , since regular solutions satisfy (2-33), and  $a \neq 0$ , since  $a = 0$  would imply  $w = -br(1 + o(1))$ ,  $\zeta = -1$ . If  $\lambda = \eta = -1$ , then from (2-8),  $w$  is nondecreasing, so it has a limit  $a \geq 0$  at 0, leading to  $w' = -a\lambda r^{-1}(1 + o(1))$ , and by integration  $a = 0$ . And  $((w')^{p-1})' = \varepsilon(1 - \alpha)w(1 + o(1))$ , so  $w'$  has a limit  $b \neq 0$ .  $\square$

Next we consider the cases where  $y$  is not monotone and possibly changes sign.

**Proposition 2.10.** *Assume  $\varepsilon = 1$ .*

- (i) *Suppose that  $N \leq \delta < \alpha$ , or  $N < \delta \leq \alpha$ . Then any solution  $y$  has a infinite number of zeros near  $\infty$ .*
- (ii) *Suppose that  $y$  has a infinite number of zeros near  $\pm\infty$ . Then either*

$$N < \alpha < \delta \text{ and } |y| < \ell \text{ and } |Y| < (\delta\ell)^{p-1} \text{ near } \pm\infty,$$

*or  $N < \delta = \alpha$ , or  $\max(\delta, N, \eta) < \alpha$ . If moreover  $\delta < N < \alpha$ , then  $|y|$  exceeds  $\ell$  at its extremal points and  $|Y|$  exceeds  $(\delta\ell)^{p-1}$  at its extremal points.*

*Proof.* (i) Suppose the conclusion does not hold. Then for example  $y > 0$  for large  $\tau$ ; and  $y$  is monotone, from Proposition 2.7(i). Applying Proposition 2.8 with  $s = \infty$ , we reach a contradiction.

(ii) Suppose that  $y$  is oscillating around 0 near  $\pm\infty$ . Then from (2-16), at the extremal points,

$$(2-42) \quad |y(\tau)|^{2-p}(\delta - \alpha) < (\delta - N)\delta^{p-1},$$

and the inequality is strict, because in case of equality,  $y$  is constant by uniqueness. Similarly  $Y$  is oscillating around 0, and at the extremal points one finds, from (2-17),

$$(2-43) \quad |Y(\tau)|^{(2-p)/(p-1)}(\delta - \alpha) < (\delta - N)\delta.$$

Then  $\max(N, \eta) < \alpha$ , thanks to Proposition 2.5; and the conclusions follow from (2-42) and (2-43).  $\square$

We can complete these results according to the sign of  $\delta - N/2$ :

**Proposition 2.11.** *Suppose that  $\varepsilon(\delta - N/2) \leq 0$ . Then any solution  $y$  has a finite number of zeros near  $\ln R_w$  or  $\ln S_w$ . If  $y$  is defined near  $\pm\infty$  and nonmonotone, then  $(y, Y)$  converges to  $\pm M_\ell$ . There is no cycle or homoclinic orbit in  $\mathbb{R}^2$ .*

*Proof.* (i) Suppose that  $y$  has an infinity of zeros. Then  $R_w = 0$  or  $S_w = \infty$ , and there exists a strictly monotone sequence  $(r_n)$  of consecutive zeros of  $w$ , converging to 0 or  $\infty$ . Since  $\varepsilon(\delta - N/2) \leq 0$ , the energy function  $V$  defined in (2–24) is nonincreasing. We claim that  $V$  is bounded. This is not easy to prove; we define for the purpose a function

$$U(r) = r^N \left( \frac{1}{2} w^2 + \varepsilon r^{-1} |w'|^{p-2} w' w \right) = e^{(N-2\delta)\tau} y \left( \frac{1}{2} y - \varepsilon Y \right).$$

Then

$$U'(r) = r^{N-1} \left( \left( \frac{1}{2} N - \alpha \right) w^2 + \varepsilon |w'|^p \right) = e^{(N-1-2\delta)\tau} \left( \left( \frac{1}{2} N - \alpha \right) y^2 + \varepsilon |Y|^{p'} \right).$$

If  $\varepsilon = 1$ , then  $\delta \leq N/2 < N < \alpha$ . If  $\varepsilon = -1$ , then  $\alpha < 0$ , by Proposition 2.10. Then  $U(r_n) = 0$  and  $\varepsilon U'(r_n) > 0$ . Therefore there exists another sequence  $(s_n)$  such that  $s_n \in (r_n, r_{n+1})$ ,  $U(s_n) = 0$ , and  $\varepsilon U'(s_n) \leq 0$ . At the point  $\tau_n = e^{s_n}$  we find  $2^{1-p'} y^{2p'} = 2|Y|^{p'} \leq \varepsilon(2\alpha - N)y^2$ , so  $(y(\tau_n), Y(\tau_n))$  is bounded. Hence  $(V(\tau_n))$  is bounded, so  $V$  is bounded near  $\pm\infty$ . Therefore  $V$  has a finite limit  $\chi$ , and  $Y$  and  $Y'$  are bounded because  $\varepsilon(\alpha - \delta) > 0$ ; in turn,  $(y, Y)$  is bounded. Otherwise  $(0, 0)$  and  $\pm M_\ell$  are not in the limit set at  $\pm\infty$ , since  $(0, 0)$  is a saddle point, and  $\pm M_\ell$  is a source or a sink. Then the trajectory has a limit cycle, and there exists a periodic solution  $(y, Y)$ . The corresponding function  $V$  is periodic and monotone, hence constant; then  $V' \equiv 0$  implies that  $Y$  is constant and hence also  $y$ , by (S). But this is a contradiction.

(ii) Suppose that  $y$  is positive near  $\pm\infty$ , and nonmonotone. If  $\varepsilon = 1$ , then  $\delta \leq N/2 < N < \alpha$ ; if  $\varepsilon = -1$ , then  $\alpha < \delta < N$ , by Proposition 2.7, and  $y$  oscillates around  $\ell$ . There exists a sequence of minimal points  $(\tau_n)$ , where  $y(\tau_n) < \ell$ , and  $|Y(\tau_n)| = \delta y(\tau_n)$ ; thus again  $(y(\tau_n), Y(\tau_n))$  is bounded, and as above  $(y, Y)$  is bounded. The trajectory has no limit cycle, and hence converges to  $M_\ell$ . Finally, if there is an homoclinic orbit, then  $\mathcal{T}_r$  is homoclinic. Then  $\lim_{\tau \rightarrow -\infty} V = \lim_{\tau \rightarrow \infty} V = 0$ ; hence  $V \equiv 0$ , and as above  $(y, Y)$  is constant, so  $(y, Y) \equiv (0, 0)$ , again a contradiction.  $\square$

**Proposition 2.12.** *If  $y$  is not monotone near  $\varepsilon\infty$  (positive or changing sign), then  $y$  and  $Y$  are bounded.*

*Proof.* From Proposition 2.11, it follows that  $\varepsilon(\delta - N/2) > 0$ . When  $\varepsilon = 1$ , and  $y$  is changing sign and  $N < \alpha < \delta$ , then  $|y|$  is bounded by  $\ell$  from above. Apart from this case, if  $y$  is changing sign, then  $\varepsilon(\alpha - \delta) > 0$ , from Proposition 2.11. If  $y$  stays positive, either  $\varepsilon = 1$ ,  $\delta < \min(\alpha, N)$ , or  $\varepsilon = -1$ ,  $\alpha < \delta < N$ , by Proposition 2.7.

In any case  $\varepsilon(\alpha - \delta) > 0$ . Here we use the energy function  $W$  defined by (2–21). We can write  $\mathcal{W}(y, Y)$  in the form

$$(2-44) \quad \mathcal{W}(y, Y) = \varepsilon(F(y, Y) + G(y)),$$

with

$$(2-45) \quad F(y, Y) = \frac{|Y|^{p'}}{p'} - \delta y Y + \frac{|\delta y|^p}{p}, \quad G(y) = \frac{(\delta - N)\delta^{p-1}}{p} |y|^p + \frac{\varepsilon(\alpha - \delta)}{2} y^2.$$

Observe that  $F(y, Y) \geq 0$ , so  $\varepsilon \mathcal{W}(y, Y) \geq G(y) > 0$  for large  $|y|$ . Then  $W'(\tau) \leq 0$  whenever  $(y(\tau), Y(\tau)) \notin \mathcal{S}_{\mathcal{G}}$ , where  $\mathcal{S}_{\mathcal{G}}$  is given in (2–22). Let  $\tau_0$  be arbitrary in the interval of definition of  $y$ . Since  $\mathcal{S}_{\mathcal{G}}$  is bounded, there exists  $k > 0$  large enough that  $\varepsilon W(\tau) \leq k$  for any  $\tau$  such that  $\varepsilon(\tau - \tau_0) \geq 0$  and  $(y(\tau), Y(\tau)) \in \mathcal{S}_{\mathcal{G}}$ , and we can choose  $k > W(\tau_0)$ . Then  $\varepsilon W(\tau) \leq k$  for  $\varepsilon(\tau - \tau_0) \geq 0$ ; hence  $y$  and  $Y$  are bounded near  $\varepsilon\infty$ .  $\square$

**Further sign properties.** We can improve Proposition 2.5 using Propositions 2.8 and 2.9:

**Proposition 2.13.** *Assume  $\varepsilon = 1$ ,  $-\infty < \alpha \leq \delta$  and  $\alpha < N$ . Then all regular solutions have constant sign,  $y$  is strictly monotone and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . Any solution has at most one zero, and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ .*

*Proof.* Regular solutions have constant sign by Proposition 2.5. Moreover  $J_N$  is increasing from 0; thus it is positive for  $r > 0$ , which means  $Y < y$ . And  $y$  is monotone near  $\infty$ , by Proposition 2.7. From Proposition 2.8, we have three possibilities: either  $\alpha < N < \delta$  and  $\lim_{\tau \rightarrow \infty} \zeta = \delta$ , in which case  $\lim_{\tau \rightarrow \infty} Y/y = (\delta - \alpha)/(\delta - N) > 1$ , which is impossible; or  $\delta \leq N$  and  $\lim_{\tau \rightarrow \infty} \zeta = \eta \geq N$ , in which case  $\lim_{\tau \rightarrow \infty} Y/y = \infty$ , which is also contradictory, or finally  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . Moreover  $y$  is increasing on  $\mathbb{R}$  from 0 to  $\infty$ ; indeed, if  $y$  has a local maximum for some  $\tau$ , we get  $\alpha < N < \delta$  and  $y(\tau) \leq \ell$  from (2–16), and moreover  $\ell < \delta^{(p-1)/(2-p)}$ ; but  $\delta y(\tau) = Y(\tau)^{1/(p-1)} < y(\tau)^{1/(p-1)}$ , which is contradictory.

For the second statement, we see from Proposition 2.5 that any solution  $w \neq 0$  has at most one zero. If  $w(r_1) = 0$  and, say,  $w > 0$  on  $(r_1, \infty)$ , we get  $w'(r_1) > 0$ ; thus  $J_N(r) \geq J_N(r_1) > 0$  for  $r \geq r_1$ , and we conclude as above.  $\square$

**Proposition 2.14.** *Assume  $\varepsilon = -1$ .*

- (i) *If  $\alpha < 0$  and  $N \leq \delta$ , all regular solutions have at least one zero.*
- (ii) *If  $0 < \alpha$ , all regular solutions have constant sign and satisfy  $S_w < \infty$ .*
- (iii) *If  $-p' < \alpha < \min(0, \eta)$ , all regular solutions have precisely one zero and  $S_w < \infty$ .*

*Proof.* (i) Let  $\alpha < 0$  and  $N \leq \delta$ . Since  $\varepsilon\alpha > 0$ , the trajectory  $\mathcal{T}_r$  starts in  $\mathfrak{D}_1$ . Suppose that  $y$  stays positive. Then  $\mathcal{T}_r$  stays in  $\mathfrak{D}_1$ , from Remark 2.3. If  $N \leq \delta$ , then  $y$  is monotone, since it can only have minimal points, from (2–16); and  $(0, 0)$  is the only stationary point. Then  $\lim_{\tau \rightarrow \infty} y = \infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha < 0$  from Proposition 2.8; thus  $(y, Y)$  is in  $\mathfrak{D}_4$  for large  $\tau$ , which is impossible.

(ii) Let  $0 < \alpha$ . Then  $\varepsilon\alpha < 0$ , so that  $\mathcal{T}_r$  starts in  $\mathfrak{D}_4$ . Moreover  $y > 0$  on  $\mathbb{R}$ , by Proposition 2.5. And  $\mathcal{T}_r$  stays in  $\mathfrak{D}_4$ , by Remark 2.1(i) on page 211. Thus  $y' = \delta y + |Y|^{1/(p-1)} > 0$ . If  $S_w = \infty$ , we see from Proposition 2.8 that  $\lim_{\tau \rightarrow \infty} \zeta = \alpha > 0$ ; hence  $(y, Y)$  ends up in  $\mathfrak{D}_1$ , which is false. Then  $S_w < \infty$ .

(iii) Let  $-p' < \alpha < \min(0, \eta)$ . Then  $\mathcal{T}_r$  starts in  $\mathfrak{D}_1$ . By Proposition 2.5,  $Y_\alpha$  stays positive,  $\mathcal{T}_r$  stays in  $\mathfrak{D}_1 \cup \mathfrak{D}_2$ , and  $Y_\alpha$  is increasing:

$$Y'_\alpha = -(p-1)(\eta-\alpha)Y_\alpha + e^{(p-(2-p)\alpha)\tau}(Y_\alpha^{1/(p-1)} - \alpha y_\alpha) > 0.$$

Suppose that  $S_w = \infty$ . Then  $\lim_{\tau \rightarrow \infty} Y_\alpha(\tau) \geq C > 0$ , so  $r^{\alpha+1}w'(r) \leq -C^{1/(p-1)}$  for large  $r$ , and, by integration,  $r^\alpha w(r) \leq -C^{1/(p-1)}/2$ . In particular, we obtain from (2–3) that  $\lim_{\tau \rightarrow \infty} y = -\infty$ . From Propositions 2.7, 2.8, and 2.9, it follows that  $\lim_{r \rightarrow \infty} r^\alpha w = L < 0$ ; thus  $\lim_{\tau \rightarrow \infty} Y_\alpha(\tau) = (\alpha L)^{p-1}$ . And there exists a unique  $\tau_0$  such that  $y_\alpha(\tau_0) = 0$ , by Remark 2.1(i). But

$$\begin{aligned} (2-46) \quad & Y''_\alpha(\tau) - (p-1)^2(\eta-\alpha)(\alpha+p')Y_\alpha \\ &= \frac{Y'_\alpha}{Y_\alpha} \left( \frac{1}{p-1} e^{(p-(2-p)\alpha)\tau} Y_\alpha^{1/(p-1)} - (p-1)(\eta-2\alpha-p')Y_\alpha \right) \\ &\geq \frac{Y'_\alpha}{Y_\alpha} \left( \frac{\alpha}{p-1} e^{(p-(2-p)\alpha)\tau} y_\alpha + (\eta-\alpha)(2-p) + (p-1)(\alpha+p')Y_\alpha \right). \end{aligned}$$

Thus  $Y''_\alpha(\tau) > 0$  for any  $\tau \geq \tau_0$ , an impossibility. Then  $S_w < \infty$ ,  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ , and  $y$  has a zero. □

**Behavior of  $w$  near  $R_w > 0$  or  $S_w < \infty$ .**

**Proposition 2.15.** *Let  $w$  be any solution of  $(E_w)$  with a reduced domain (so either  $\varepsilon = 1$  and  $R_w > 0$  or  $\varepsilon = -1$  and  $S_w < \infty$ ). Let  $s = R_w$  or  $S_w$ . Then*

$$(2-47) \quad \lim_{r \rightarrow s} |r-s|^{(p-1)/(2-p)} s^{1/(2-p)} w = \pm \left( \frac{p-1}{2-p} \right)^{\frac{p-1}{2-p}} \quad \text{and} \quad \lim_{\tau \rightarrow \ln s} \sigma = \varepsilon.$$

*Proof.* From Proposition 2.5, we can suppose that  $\varepsilon w$  is decreasing near  $s$  and  $\lim_{r \rightarrow s} w = \infty$ ; thus  $y > 0$ ,  $\varepsilon Y > 0$  near  $\ln s$ , and  $\lim_{\tau \rightarrow \ln s} y = \infty$ . Also,  $\sigma$  is monotone near  $\ln s$ , by Proposition 2.7; thus it has a limit  $\mu$  such that  $\varepsilon\mu \in [0, \infty]$ . Suppose that  $\mu = 0$ . Then  $Y = o(y) = o(y - \varepsilon Y)$ ; from (2–13) we get

$$(y - \varepsilon Y)' = (\delta - \alpha)(y - \varepsilon Y) + \varepsilon(N - \alpha)Y = (\delta - \alpha + o(1))(y - \varepsilon Y),$$

so  $y$  cannot blow up in finite time. In the same way, if  $\mu = \infty$ , then  $y = o(\varepsilon Y) = o(\varepsilon Y - y)$ , and

$$(y - \varepsilon Y)' = (\delta - N)(y - \varepsilon Y) + (N - \alpha)y = (\delta - N + o(1))(y - \varepsilon Y),$$

again leading to a contradiction; thus  $\varepsilon\mu \in (0, \infty)$ . Therefore  $\lim_{\tau \rightarrow \ln R_w} \zeta = \varepsilon\infty$  and  $\mu = \varepsilon$  from (2-37); then  $w'w^{-1/(p-1)} + (\varepsilon + o(1))r^{1/(p-1)} = 0$ , and (2-47) holds.  $\square$

**More information on stationary points.**

**The Hopf bifurcation point.** When  $\varepsilon(\delta - N/2) > 0$ , a Hopf bifurcation appears at the critical value  $\alpha = \alpha^*$  given by (1-11). Then some cycles do appear near  $\alpha^*$ , by the Poincaré–Andronov–Hopf theorem; see [Hale and Koçak 1991, p. 344]. We get more precise results by using the Lyapunov test for a weak sink or source; it requires an expansion up to the order 3 near  $M_\ell$ , in a suitable basis of eigenvectors, where the linearized problem has a rotation matrix.

**Theorem 2.16.** *Let  $\varepsilon(\delta - N/2) > 0$ .*

- (i) *Suppose  $\varepsilon = 1$ . If  $\alpha = \alpha^*$ , then  $M_\ell$  is a weak source. If  $\alpha < \alpha^*$  with  $\alpha^* - \alpha$  small enough, there exists a unique limit cycle in  $\mathcal{D}_1$  attracting at  $-\infty$ .*
- (ii) *Suppose  $\varepsilon = -1$ . If  $\alpha = \alpha^*$ , then  $M_\ell$  is a weak sink. If  $\alpha > \alpha^*$  with  $\alpha - \alpha^*$  small enough, there exists a unique limit cycle in  $\mathcal{D}_1$ , attracting at  $\infty$ .*

*Proof.* The eigenvalues are given by  $\lambda_1 = -ib$ ,  $\lambda_2 = ib$ , with  $b = \sqrt{p'(N - \delta)}$ . From (2-29) we get

$$v(\alpha^*) = 2\delta - N = \frac{\delta(N - \delta)}{(p - 1)(\alpha^* - \delta)} = \frac{\varepsilon(\delta\ell)^{2-p}}{(p - 1)}.$$

First we make the substitution (2-27) as above, which leads to (2-28). The function  $\Psi$  defined in (2-29) has an expansion near  $t = 0$  of the form

$$\Psi(\vartheta) = B_2\vartheta^2 + B_3\vartheta^3 + \dots,$$

where

$$B_2 = \frac{(2 - p)(\delta\ell)^{3-2p}}{2(p - 1)^2}, \quad B_3 = \frac{(2 - p)(3 - 2p)(\delta\ell)^{4-3p}}{6(p - 1)^6} = \frac{2(3 - 2p)B_2^2}{3(2 - p)v(\alpha^*)}.$$

Next we make the substitution

$$\tau = -\theta/b, \quad \bar{y}(\tau) = \varepsilon v(\alpha)x_1(\theta), \quad \bar{Y}(\tau) = \delta x_1(\theta) + bx_2(\theta),$$

and obtain

$$x_1'(\theta) = x_2 + \frac{\varepsilon}{bv(\alpha)}\Psi(\delta x_1 + bx_2), \quad x_2'(\theta) = -x_1 - \frac{\varepsilon(N - \delta)}{b^2v(\alpha)}\Psi(\delta x_1 + bx_2).$$

We write the expansion of order 3 in the form

$$\begin{aligned} x'_1 &= x_2 + \varepsilon(a_{2,0}x_1^2 + a_{1,1}x_1x_2 + a_{0,2}x_2^2 + a_{3,0}x_1^3 + a_{2,1}x_1^2x_2 + a_{1,2}x_1x_2^2 + a_{0,3}x_2^3 + \dots), \\ x'_1 &= -x_1 + \varepsilon(b_{2,0}x_1^2 + b_{1,1}x_1x_2 + b_{0,2}x_2^2 + b_{3,0}x_1^3 + b_{2,1}x_1^2x_2 + b_{1,2}x_1x_2^2 + b_{0,3}x_2^3 + \dots), \end{aligned}$$

and we compute the Lyapunov coefficient

$$\begin{aligned} L_C &= \varepsilon(3a_{3,0} + a_{1,2} + b_{2,1} + 3b_{0,3}) \\ &\quad - a_{2,0}a_{1,1} + b_{1,1}b_{0,2} - 2a_{0,2}b_{0,2} - a_{0,2}a_{1,1} + 2a_{2,0}b_{2,0} + b_{1,1}b_{2,0}. \end{aligned}$$

After simplification, we obtain

$$\begin{aligned} \frac{(2-p)b\nu(\alpha)^2}{2B_2^2(\delta^2 + b^2)} L_C &= (N - 2\delta)(1 - \varepsilon(3 - 2p)) \\ &= \begin{cases} 2(N - 2\delta)(p - 1) < 0 & \text{if } \varepsilon = 1, \\ 2(N - 2\delta)(2 - p) > 0 & \text{if } \varepsilon = -1, \end{cases} \end{aligned}$$

The nature of  $M_\ell$  follows from [Hubbard and West 1995, p. 292], taking into account that  $\theta$  has opposite sign from  $\tau$ . If  $\varepsilon = 1$ ,  $M_\ell$  is a weak source, and there exists a small limit cycle attracting at  $-\infty$  for all  $\alpha$  near  $\alpha^*$  such that  $M_\ell$  is a sink; this means that  $\alpha < \alpha^*$ . If  $\varepsilon = -1$ ,  $M_\ell$  is a weak sink and there exists a small limit cycle attracting at  $\infty$  for all  $\alpha$  near  $\alpha^*$  such that  $M_\ell$  is a source; this means  $\alpha^* < \alpha$ . □

**Node points or spiral points.** When the system ( $S$ ) has three stationary points, and  $M_\ell$  is a source or a sink (so  $\delta < N$ ), it is interesting to know if  $M_\ell$  is a node point. When  $\alpha^*$  exists, it is a spiral point, by (2–30).

If  $\varepsilon = 1$ , we see from (2–31) that  $M_\ell$  is a node point when  $\delta \leq N/2 - \sqrt{p'(N - \delta)}$  or  $\delta > N/2 - \sqrt{p'(N - \delta)}$  and  $\alpha \leq \alpha_1$ , or  $\delta > N/2 + \sqrt{p'(N - \delta)}$  and  $\alpha_2 \leq \alpha$ , where

$$\begin{aligned} (2-48) \quad \alpha_1 &= \delta + \frac{\delta(N - \delta)}{(p - 1)(2\delta - N + 2\sqrt{p'(N - \delta)})}, \\ \alpha_2 &= \delta + \frac{\delta(N - \delta)}{(p - 1)(2\delta - N - 2\sqrt{p'(N - \delta)})}. \end{aligned}$$

If  $\varepsilon = -1$ ,  $M_\ell$  is a node when  $\delta \geq N/2 + \sqrt{p'(N - \delta)}$ , or  $\delta < N/2 + \sqrt{p'(N - \delta)}$  and  $\alpha_2 \leq \alpha$ , or  $\delta < N/2 - \sqrt{p'(N - \delta)}$  and  $\alpha \leq \alpha_1$ . In any case  $\alpha_1 < \alpha_2$ .

**Remarks 2.17.** (i) Let  $\varepsilon = 1$ . One can verify that  $N \leq \alpha_1$  and that  $N = \alpha_1$  if and only if  $N = \delta/(p - 1) = p'/(2 - p)$ . Also  $\alpha_1 < \eta$  if and only if  $\delta^2 + (7 - N)\delta + N > 0$ , which is true for  $N \leq 14$ , but not always.

(ii) Let  $\varepsilon = -1$ . It is easy to see that  $\alpha_2 \leq 0$  and that  $\alpha_2 = 0$  if and only if  $N(2 - p) = \delta$ , or equivalently  $N = p/(2 - p)^2$ . Also  $\alpha_2 > -p'$  if and only if  $\delta^2 + 7\delta - 8N < 0$ , which is true for  $\delta < N/2 < 9$ , but not always.

**Nonexistence of cycles.** If the system  $(S)$  admits a cycle  $\mathbb{C}$  in  $\mathbb{R}^2$ , then  $\mathbb{C}$  surrounds at least one stationary point. If it surrounds  $(0, 0)$ , the corresponding solutions  $y$  are not of constant sign. If it only surrounds  $M_\ell$ , then it stays in  $\mathfrak{Q}_1$ , so  $y$  stays positive. Indeed  $\alpha \neq 0$  from (1-9), and  $\mathbb{C}$  cannot intersect  $\{(\varphi, 0), \varphi > 0\}$  at two points, and similarly  $\{(0, \xi), \xi > 0\}$ , by Remark 2.1(i) on page 211.

For suitable values of  $\alpha, \delta, N$ , we can show that cycles cannot exist, by using Bendixson's criterion or the Poincaré map. Writing  $(S)$  under the form

$$(2-49) \quad y' = f_1(y, Y), \quad Y' = f_2(y, Y),$$

we obtain

$$(2-50) \quad \frac{\partial f_1}{\partial y}(y, Y) + \frac{\partial f_2}{\partial Y}(y, Y) = 2\delta - N - \varepsilon|Y|^{(2-p)/(p-1)}.$$

For example, as a direct consequence of Bendixson's criterion, if  $\varepsilon(\delta - N/2) < 0$ , we find again the nonexistence of any cycle in  $\mathbb{R}^2$ , which was obtained in Proposition 2.11. Now we consider cycles in  $\mathfrak{Q}_1$ .

First we extend to system  $(S)$  a general property of quadratic systems, proved in [Chicone and Tian 1982], stating that there cannot exist a closed orbit surrounding a node point. Note that the restriction of our system to  $\mathfrak{Q}_1$  is quadratic if  $p = \frac{3}{2}$ .

**Theorem 2.18.** *Let  $\delta < N$  and  $\varepsilon(\delta - \alpha) < 0$ . When  $M_\ell$  is a node point, there is no cycle or homoclinic orbit in  $\mathfrak{Q}_1$ .*

*Proof.* We use the linearization (2-27), (2-28), (2-29). Consider the line  $L$  with equation  $A\bar{y} + \bar{Y} = 0$ , where  $A$  is a real parameter. The points of  $L$  are in  $\mathfrak{Q}_1$  whenever  $-(\delta\ell)^{p-1} < \bar{Y}$  and  $-\ell < \bar{y}$ . As in [Chicone and Tian 1982], we study the orientation of the vector field along  $L$ : we find

$$A\bar{y}' + \bar{Y}' = (\varepsilon v(\alpha)A^2 + (N + v(\alpha))A + \varepsilon\alpha)\bar{y} - (A + \varepsilon)\Psi(\bar{Y}).$$

Using (2-31), apart from the case  $\varepsilon = 1, \alpha = N = \alpha_1$ , we can find an  $A$  such that  $\varepsilon v(\alpha)A^2 + (N + v(\alpha))A + \varepsilon\alpha = 0$ , and  $A + \varepsilon \neq 0$ . Moreover  $\Psi(\bar{Y}) \geq 0$  on  $L \cap \mathfrak{Q}_1$ ; indeed,  $(p - 1)\Psi'(t) = ((\delta\ell)^{p-1} + t)^{(2-p)/(p-1)} - t^{(2-p)/(p-1)}$ , so  $\Psi$  has a minimum on  $(-\delta\ell)^{p-1}, \infty)$  at 0, and hence is nonnegative on this interval. Then the orientation of the vector field does not change along  $L \cap \mathfrak{Q}_1$ ; in particular no cycle can exist in  $\mathfrak{Q}_1$ ; and similarly no homoclinic trajectory can exist. In the case  $\varepsilon = 1, \alpha = N = \alpha_1, Y \equiv y \in [0, \ell)$  defines the trajectory  $\mathcal{T}_r$ , corresponding to the solutions given by (1-6) with  $K > 0$ , and again no cycle can exist in  $\mathfrak{Q}_1$ : it would intersect  $\mathcal{T}_r$ . □

Next we prove the nonexistence of cycles on one side of the Hopf bifurcation point:

**Theorem 2.19.** *Assume  $\delta < N$  and  $\varepsilon(\delta - \alpha) < 0 < \varepsilon(\delta - N/2)$ . If  $\varepsilon(\alpha - \alpha^*) \geq 0$ , there exists no cycle or homoclinic orbit in  $\mathcal{Q}_1$ .*

*Proof.*  $M_\ell$  is a source or weak source if  $\varepsilon = 1$ , and a sink or weak sink if  $\varepsilon = -1$ . Suppose there exists a cycle in  $\mathcal{Q}_1$ . Then any trajectory starting from  $M_\ell$  at  $-\varepsilon\infty$  has a limit cycle in  $\mathcal{Q}_1$ , which is attracting at  $\varepsilon\infty$ . Such a cycle is not unstable (if  $\varepsilon = 1$ ) or not stable (if  $\varepsilon = -1$ ); in other words the Floquet integral on the period  $[0, \mathcal{P}]$  is nonpositive if  $\varepsilon = 1$  and nonnegative if  $\varepsilon = -1$ . From (2-50) we then get

$$(2-51) \quad \varepsilon \int_0^{\mathcal{P}} \left( \frac{\partial f_1}{\partial y}(y, Y) + \frac{\partial f_2}{\partial Y}(y, Y) \right) d\tau = \int_0^{\mathcal{P}} \left( |2\delta - N| - \frac{1}{p-1} Y^{(2-p)/(p-1)} \right) d\tau \leq 0.$$

Now, from (2-28),

$$\begin{aligned} 0 &= \delta \int_0^{\mathcal{P}} \bar{y} d\tau - v(\alpha) \int_0^{\mathcal{P}} \bar{Y} d\tau - \int_0^{\mathcal{P}} \Psi(\bar{Y}) d\tau, \\ 0 &= \alpha \int_0^{\mathcal{P}} \bar{y} d\tau + (\delta - N - v(\alpha)) \int_0^{\mathcal{P}} \bar{Y} d\tau - \int_0^{\mathcal{P}} \Psi(\bar{Y}) d\tau. \end{aligned}$$

Moreover, since  $\Psi$  is nonnegative,

$$\int_0^{\mathcal{P}} \Psi(\bar{Y}) d\tau = -p' \int_0^{\mathcal{P}} \bar{y} d\tau = -\frac{p'(N - \delta)}{\alpha - \delta} \int_0^{\mathcal{P}} \bar{Y} d\tau > 0;$$

and since  $y' = \delta y - Y^{1/(p-1)}$ ,

$$\int_0^{\mathcal{P}} Y^{1/(p-1)} dt = \delta \int_0^{\mathcal{P}} y dt < \delta \ell \mathcal{P}.$$

From this, (2-51), and Jensen's inequality, it follows that

$$\begin{aligned} (p-1)|2\delta - N| &\leq \int_0^{\mathcal{P}} Y^{(2-p)/(p-1)} d\tau \\ &\leq \mathcal{P}^{p-1} \left( \int_0^{\mathcal{P}} Y^{1/(p-1)} d\tau \right)^{2-p} < (\delta \ell)^{2-p} = \frac{\varepsilon \delta (N - \delta)}{\alpha - \delta}. \end{aligned}$$

Hence  $\varepsilon(\alpha - \alpha^*) < 0$ , a contradiction. Next, suppose that there is an homoclinic orbit. From [Hubbard and West 1995, Theorem 9.3, p. 303] we see that the saddle connection is repelling if  $\varepsilon = 1$  and attracting if  $\varepsilon = -1$ , because the sum of the eigenvalues  $\mu_1, \mu_2$  of the linearized problem at  $(0, 0)$  is  $2\delta - N$ . That means that the solutions just inside it spiral toward the loop near  $-\varepsilon\infty$ . Because  $M_\ell$  is a

source or weak source or sink or weak sink, such solutions have a limit cycle that is attracting at  $\varepsilon\infty$ . As before, we reach a contradiction.  $\square$

Finally we get the nonexistence of cycles in nonobvious cases, where we have shown that any solution has at most one or two zeros.

**Theorem 2.20.** *Assume  $\delta < N$  and  $\varepsilon(\delta - \alpha) < 0 < \varepsilon(\delta - N/2)$ . If  $\varepsilon = 1$  and  $\alpha \leq \eta$ , or  $\varepsilon = -1$  and  $-p' \leq \alpha < 0$ , there exists no cycle and no homoclinic orbit in  $\mathfrak{Q}_1$ .*

*Proof.* (i) Suppose there exists a cycle. There are two possibilities:

Suppose  $\varepsilon = 1$  and  $\alpha \leq \eta$ .  $M_\ell$  is a sink since  $\alpha < \alpha^*$ , so any trajectory converging to  $M_\ell$  at  $\infty$  has a limit cycle  $\mathbb{O}$  in  $\mathfrak{Q}_1$ , attracting at  $-\infty$ . Let  $(y, Y)$  describe the orbit  $\mathbb{O}$ , of period  $\mathcal{P}$ . Then  $\mathbb{O}$  is not stable, so the Floquet integral is nonnegative, and from (2-51),

$$\int_0^{\mathcal{P}} \left( 2\delta - N - \frac{1}{p-1} Y^{(2-p)/(p-1)} \right) d\tau \geq 0.$$

Otherwise  $y$  is bounded from above and below; thus the function  $y_\alpha$ , defined by (2-3) with  $d = \alpha$ , satisfies  $\lim_{\tau \rightarrow -\infty} y_\alpha = 0$  and  $\lim_{\tau \rightarrow \infty} y_\alpha = \infty$ ; moreover  $y_\alpha$  has only minimal points, from (2-35), since  $\alpha \leq \eta$ ; thus  $y'_\alpha > 0$  on  $\mathbb{R}$ . From (2-5) and (2-4) with  $d = \alpha$ ,

$$\frac{y''_\alpha}{y'_\alpha} + \eta - 2\alpha + \frac{1}{p-1} Y^{(2-p)/(p-1)} = \frac{\alpha(\eta - \alpha)y_\alpha}{y'_\alpha} = \frac{\alpha(\eta - \alpha)y_\alpha}{\alpha y_\alpha - Y_\alpha^{1/(p-1)}} > \eta - \alpha.$$

Upon integration over  $[0, \mathcal{P}]$ , this implies  $\eta - 2\alpha + 2\delta - N > \eta - \alpha$ , which is impossible, since  $\delta - N + \delta - \alpha < 0$ .

Alternatively, suppose  $\varepsilon = -1$  and  $-p' \leq \alpha < 0$ .  $M_\ell$  is a source since  $\alpha^* < \alpha$ , and any trajectory converging to it at  $-\infty$  has a limit cycle  $\mathbb{O}'$  attracting at  $\infty$ . Let  $(y, Y)$  describe the orbit  $\mathbb{O}'$ , of period  $\mathcal{P}$ . Then  $\mathbb{O}'$  is not unstable, so the Floquet integral is nonpositive, hence

$$\int_0^{\mathcal{P}} \left( 2\delta - N + \frac{1}{p-1} Y^{(2-p)/(p-1)} \right) d\tau \leq 0.$$

Moreover  $Y$  is bounded from above and below; thus  $Y_\alpha$ , defined by (2-3) with  $d = \alpha$ , satisfies  $\lim_{\tau \rightarrow -\infty} Y_\alpha = \infty$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = 0$ . And  $Y_\alpha$  has only minimal points, by (2-36), since  $-p' \leq \alpha < 0$ ; thus  $Y'_\alpha < 0$  on  $\mathbb{R}$ . From (2-6) and (2-4) we get

$$\begin{aligned} \frac{Y''_\alpha}{Y'_\alpha} + (p-1)(\eta - 2\alpha - p') - \frac{1}{p-1} Y^{(2-p)/(p-1)} &= \frac{(p-1)^2(\eta - \alpha)(p' + \alpha)Y_\alpha}{Y'_\alpha} \\ &< -(p-1)(p' + \alpha). \end{aligned}$$

Upon integration over  $[0, \mathcal{P}]$ , this implies  $(p-1)(\eta - 2\alpha - p') + 2\delta - N < -(p-1) \times (p' + \alpha)$ , which means  $p\delta + (p-1)|\alpha| < 0$ ; but this is false.

(ii) Suppose there exists an homoclinic orbit. Since  $\delta < N$ , the origin is a saddle point, so  $\mathcal{T}_r$  is the only trajectory starting from  $(0, 0)$  in  $\mathcal{Q}_1$ , and there exists a unique trajectory  $\mathcal{T}_s$  converging to  $(0, 0)$ , lying in  $\mathcal{Q}_1$  for large  $\tau$ , having infinite slope at  $(0, 0)$ , and satisfying  $\lim_{r \rightarrow 0} r^\eta w = c > 0$ .

If  $\varepsilon = 1$ , then  $\mathcal{T}_r$  satisfies  $\lim_{\tau \rightarrow -\infty} e^{-\alpha\tau} y_\alpha = a > 0$ , so  $\lim_{\tau \rightarrow -\infty} y_\alpha = 0$ ; also  $y_\alpha$  has only minimal points, so it is increasing and positive; and  $\mathcal{T}_s$  satisfies  $\lim_{\tau \rightarrow \infty} e^{(\eta-\alpha)\tau} y_\alpha = c > 0$ . If  $\alpha < \eta$ , then  $\lim_{\tau \rightarrow \infty} y_\alpha = 0$ , thus  $\mathcal{T}_r \neq \mathcal{T}_s$ . If  $\alpha = \eta$ ,  $\mathcal{T}_s$  is given explicitly by (1-7), that means  $y_\alpha$  is constant, thus again  $\mathcal{T}_r \neq \mathcal{T}_s$ .

If  $\varepsilon = -1$ , then  $\mathcal{T}_s$  satisfies  $\lim_{\tau \rightarrow -\infty} e^{(\eta-\alpha)(p-1)\tau} Y_\alpha > 0$ , because  $\lim_{\tau \rightarrow -\infty} \zeta = \eta$ ; so  $\lim_{\tau \rightarrow \infty} Y_\alpha = 0$ . Moreover  $Y_\alpha$  has only minimal points, and hence is increasing and positive; otherwise  $\mathcal{T}_r$  satisfies  $\lim_{\tau \rightarrow -\infty} e^{-(\alpha(p-1)+p)\tau} Y_\alpha = -a\alpha/N > 0$ , by (2-33). If  $\alpha > -p'$ , we get  $\lim_{\tau \rightarrow \infty} Y_\alpha = 0$ , which implies  $\mathcal{T}_r \neq \mathcal{T}_s$ . If  $\alpha = -p'$ , then  $\mathcal{T}_r$  is given explicitly by (1-8); in other words  $Y_\alpha$  is constant, and again  $\mathcal{T}_r \neq \mathcal{T}_s$ . □

**Boundedness of cycles.** When there do exist cycles, except for a few cases, we cannot prove their uniqueness, but we can show:

**Theorem 2.21.** *When nonempty, the set  $\mathcal{C}$  of cycles of system (S) is bounded in  $\mathbb{R}^2$ .*

*Proof.* Suppose there exists a cycle  $\mathcal{C}$  in  $\mathbb{R}^2$ . By Propositions 2.5, 2.7, 2.10, 2.11 and Theorem 2.20, this can happen only in four cases:  $\varepsilon = 1, N < \alpha < \delta$ ;  $\varepsilon = 1, N < \delta = \alpha$ ;  $\varepsilon = 1, \max(\delta, N, \eta) < \alpha, N/2 < \delta$ ;  $\varepsilon = -1, \delta < N/2, \alpha < -p'$ . In the first case,  $\mathcal{C}$  is bounded and lies in  $(-\ell, \ell) \times (-(\delta\ell)^{p-1}, (\delta\ell)^{p-1})$ , by Proposition 2.10. In the other cases we use the energy function  $W$ . Let  $(y, Y)$  describe the trajectory  $\mathcal{C}$ . Then  $W$  is periodic, and its maximum and minimum points are precisely the points of the curve  $\mathcal{L}$ . Indeed if  $W'(\tau_1) = 0$  and the point  $(y(\tau_1), Y(\tau_1))$  is not on  $\mathcal{L}$ , it lies on the curve  $\mathcal{M}$  defined in (2-11); hence  $y'(\tau_1) = 0$  and  $y''(\tau_1) \neq 0$ , since  $\mathcal{C}$  is not just a stationary point. Therefore  $(\delta y - |Y|^{(2-p)/(p-1)} Y)(|\delta y|)^{p-2} \delta y - Y > 0$  near  $\tau_1$ ; then  $W'$  has constant sign, and  $\tau_1$  is not a maximum or a minimum. In this way we obtain estimates for  $W$  independently of the trajectory:

$$\max_{\tau \in \mathbb{R}} |W(\tau)| = M = \max_{(y, Y) \in \mathcal{L}} |W(y, Y)|.$$

At the maximal points  $\tau$  of  $y$ , one has  $|Y(\tau)|^{(2-p)/(p-1)} Y(\tau) = \delta y(\tau)$ , so

$$W(\tau) = \frac{\varepsilon(\delta - N)\delta^{p-1}}{p} |y(\tau)|^p + \frac{\alpha - \delta}{2} y^2(\tau).$$

By the Hölder inequality,  $y$  is bounded by a constant independent of the trajectory, and

$$\frac{|Y|^{p'}}{p'} \leq \delta y Y + \frac{|2\delta - N|\delta^{p-1}}{p} |y|^p + \frac{|\alpha - \delta|}{2} y^2 + M.$$

Thus  $Y$  is also uniformly bounded, and  $\mathcal{C}$  is bounded. □

### 3. The case $\varepsilon = 1$ , $\alpha < \delta$ or $\alpha = \delta < N$

**Lemma 3.1.** *Assume  $\varepsilon = 1$  and  $-\infty < \max(\alpha, N) < \delta(\alpha \neq 0)$ . In the phase plane  $(y, Y)$ , there exist*

- (i) *a trajectory  $\mathcal{T}_1$  converging to  $M_\ell$  at  $\infty$ , such that  $y$  is increasing as long as it is positive;*
- (ii) *a trajectory  $\mathcal{T}_2$  in  $\mathcal{Q}_1 \cup \mathcal{Q}_4$  converging to  $M_\ell$  at  $-\infty$ , and unbounded at  $\infty$ , with  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ ;*
- (iii) *a trajectory  $\mathcal{T}_3$  converging to  $M_\ell$  at  $-\infty$ , such that  $y$  has at least one zero;*
- (iv) *a trajectory  $\mathcal{T}_4$  in  $\mathcal{Q}_1$ , converging to  $M_\ell$  at  $\infty$ , with  $\lim_{\tau \rightarrow \ln R_w} Y/y = 1$ ;*
- (v) *trajectories  $\mathcal{T}_5$  in  $\mathcal{Q}_1 \cup \mathcal{Q}_4$  unbounded at  $\pm\infty$ , with*

$$\lim_{\tau \rightarrow \infty} \zeta = \alpha \quad \text{and} \quad \lim_{\tau \rightarrow \ln R_w} Y/y = 1.$$

*Proof.* Here the system (S) has three stationary points, defined by (2–26). The point  $(0, 0)$  is a source, and the point  $M_\ell$  is a saddle point. The eigenvalues satisfy  $\lambda_1 < 0 < \lambda_2 < \delta$ . The eigenvectors  $u_1 = (-v(\alpha), \lambda_1 - \delta)$  and  $u_2 = (v(\alpha), \delta - \lambda_2)$  form a positively oriented basis, and  $u_1$  points toward  $\mathcal{Q}_3$ , while  $u_2$  points toward  $\mathcal{Q}_1$ . There exist four particular trajectories converging to  $M_\ell$  at  $\pm\infty$ , namely:

- $\mathcal{T}_1$  converging to  $M_\ell$  at  $\infty$ , with tangent vector  $u_1$ ; then  $y < \ell$  and  $Y < (\delta\ell)^{p-1}$  and  $y' > 0$  near  $\infty$ ; as above,  $y$  cannot have a local minimum, so  $y' > 0$  whenever  $y > 0$ .
- $\mathcal{T}_2$  converging to  $M_\ell$  at  $-\infty$ , with tangent vector  $u_2$ ; then  $y' > 0$  near  $-\infty$ . If  $y$  has a local maximum at some  $\tau$ , then  $y''(\tau) \leq 0$ , so that  $y(\tau) \leq \ell$  from (2–16), which is impossible. Then  $y$  is increasing on  $\mathbb{R}$  and  $\lim_{\tau \rightarrow \infty} y = \infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$  from Proposition 2.8. In particular  $\mathcal{T}_2$  stays in  $\mathcal{Q}_1$  if  $\alpha > 0$ , and enters  $\mathcal{Q}_4$  if  $\alpha < 0$ .
- $\mathcal{T}_3$  converging to  $M_\ell$  at  $-\infty$ , with tangent vector  $-u_2$ ; then  $y' < 0$  near  $-\infty$ . If  $y$  has a local minimum at some  $\tau$ , then  $y(\tau) \geq \ell$ , which is still impossible. Thus  $y$  is decreasing at long as the trajectory stays in  $\mathcal{Q}_1$ . It cannot stay in it, because it cannot converge to  $(0, 0)$ . It cannot enter  $\mathcal{Q}_4$  by Remark 2.1(i) on page 211. Then it enters  $\mathcal{Q}_2$  and  $y$  has at least one zero.
- $\mathcal{T}_4$  converging to  $M_\ell$  at  $\infty$ , with tangent vector  $-u_1$ ; then  $y' < 0$  near  $\infty$ . As above,  $y$  cannot have a local maximum, it is decreasing and  $\lim_{\tau \rightarrow \ln R_w} y = \infty$ . From Proposition 2.8,  $y$  cannot be defined near  $-\infty$ , hence  $R_w > 0$  and  $\lim_{\tau \rightarrow \ln R_w} Y/y = 1$ .

For any trajectory  $\mathcal{T}$  in the domain delimited by  $\mathcal{T}_2$  and  $\mathcal{T}_4$ , the function  $y$  is positive, and  $\mathcal{T}$  cannot converge to  $M_\ell$  at  $\infty$ , and  $y$  is monotone for large  $\tau$  from

Proposition 2.7, because  $\alpha < \delta$ ; thus  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$  from Proposition 2.8, and  $y$  is not defined near  $-\infty$ , and  $\mathcal{T}$  is of type (5).  $\square$


We now study the various global behaviors, according to the values of  $\alpha$ . The results are expressed in terms of  $w$ .

$$\alpha \leq N < \delta$$

**Theorem 3.2.** *Assume the  $\varepsilon = 1$  and  $-\infty < \alpha \leq N < \delta$ , with  $\alpha \neq 0$ . All regular solutions  $w$  of  $(E_w)$  have constant sign, and  $\lim_{r \rightarrow \infty} r^\alpha |w| = L > 0$  if  $\alpha < N$ ,  $\lim_{r \rightarrow \infty} r^\delta |w| = \ell$  if  $\alpha = N$ . And  $w(r) = \ell r^{-\delta}$  is also a solution. There exist solutions satisfying any one of these characterizations:*

- (1) (only if  $\alpha < N$ )  $w$  is positive,  $\lim_{r \rightarrow 0} r^\eta w = c > 0$ , if  $N \geq 2$  (and (2–41) holds with  $a > 0 > b$  if  $N = 1$ ), and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;
- (2)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\delta w = \ell$ ,  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$ ;
- (3)  $w$  has precisely one zero,  $\lim_{r \rightarrow 0} r^\delta w = \ell$ ,  $\lim_{r \rightarrow \infty} r^\alpha w(r) = L < 0$ ;
- (4)  $w$  is positive,  $R_w > 0$ ,  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;
- (5)  $w$  is positive,  $R_w > 0$ ,  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$ ;
- (6)  $w$  has one zero,  $R_w > 0$ , and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ ;
- (7) (only if  $\alpha < N$ )  $w$  is positive,  $\lim_{r \rightarrow 0} r^\eta w = c > 0$  if  $N \geq 2$  (and (2–41) holds with  $a > 0 > b$  if  $N = 1$ ), and  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$ ;
- (8)  $w$  has one zero, with  $\lim_{r \rightarrow 0} r^\eta w = c > 0$  if  $N \geq 2$  (and (2–41) holds with  $a > 0 > b$  if  $N = 1$ ), and  $\lim_{r \rightarrow \infty} r^\alpha w = -L < 0$ ;
- (9)  $N = 1$ ,  $w > 0$  and (2–41) holds with  $a \geq 0$ ,  $b > 0$  and  $\lim_{r \rightarrow \infty} r^\alpha w = L$ .

Up to symmetry, all the solutions of  $(E_w)$  are as above.



**Figure 1.** Theorem 3.2:  $N = 2 < \delta = 3$ .

*Proof.* (i) We first assume that  $\alpha \neq N$ , and refer to Figure 1, left. The trajectory  $\mathcal{T}_r$  starts in  $\mathcal{D}_1$  for  $\alpha > 0$ , in  $\mathcal{D}_4$  for  $\alpha < 0$ , and  $y$  stays positive. Then  $\lim_{\tau \rightarrow \infty} y = \infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ , and  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$ , by Propositions 2.10 and 2.13, since  $\alpha < N$ . Moreover  $y$  is increasing: indeed if it has a local maximum, at this point  $y \leq \ell$ , and then  $y$  has no local minimum, since at such a point  $y \geq \ell$ , so that  $y$  cannot tend to  $\infty$ . Then  $\mathcal{T}_r$  stays in  $\mathcal{D}_1$ , and  $Y$  is increasing from 0 to  $\infty$ . Indeed each extremal point  $\tau$  of  $Y$  is a local minimum, from (2–17). If  $\alpha < 0$ , in the same way, then  $Y$  is decreasing from 0 to  $-\infty$ , and  $\mathcal{T}_r$  stays in  $\mathcal{D}_4$ .

First we follow the trajectory  $\mathcal{T}_1$ : it does not intersect  $\mathcal{T}_r$ , and cannot enter  $\mathcal{D}_2$  by Remark 2.1(i). Thus  $y$  stays positive and increasing. It cannot enter  $\mathcal{D}_4$ , seeing that it does not meet  $\mathcal{T}_r$  if  $\alpha > 0$ , or (by the same remark) if  $\alpha < 0$ . Thus  $\mathcal{T}_1$  stays in  $\mathcal{D}_1$ , and  $(y, Y)$  converges necessarily to  $(0, 0)$ . If  $N \geq 2$ , then  $\lim_{\tau \rightarrow -\infty} \zeta = \eta$ ,  $\lim_{r \rightarrow 0} r^\eta w = c > 0$  from Proposition 2.8 and 2.9. If  $N = 1$ , since  $\mathcal{T}_1$  stays in  $\mathcal{D}_1$ , then necessarily  $\lim_{\tau \rightarrow -\infty} \zeta = 0$ , thus (2–41) holds with  $a > 0 > b$ .

Next we follow  $\mathcal{T}_3$ : here  $y$  has a zero, which is unique by Proposition 2.5, since  $\alpha < N$ . Then  $y < 0$ , and  $\lim_{\tau \rightarrow \infty} y = -\infty$ ,  $\lim_{r \rightarrow \infty} r^\alpha w = -L < 0$  by Propositions 2.8 and 2.9.  $\mathcal{T}_3$  stays in  $\mathcal{D}_2$  if  $\alpha < 0$ , or goes from  $\mathcal{D}_2$  into  $\mathcal{D}_3$  if  $\alpha > 0$ .

Trajectories  $\mathcal{T}_2, \mathcal{T}_4, \mathcal{T}_5$  of Lemma 3.1 yield solutions  $w$  of type (2), (4), (5).

For any trajectories  $\mathcal{T}_6$  in the domain delimited by  $\mathcal{T}_3, \mathcal{T}_4$ ,  $y$  has one zero, and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ ; and  $w$  is of type (6).

The solutions of type (7) correspond to the trajectories  $\mathcal{T}$  in the domain delimited by  $\mathcal{T}_r, \mathcal{T}_1, \mathcal{T}_2$ . Indeed  $\lim_{\tau \rightarrow \infty} y = \infty$ , and  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$ . And  $\lim_{\tau \rightarrow -\infty} y = 0$ . If  $N \geq 2$ , then  $\lim_{\tau \rightarrow -\infty} \zeta = \eta$ ,  $\lim_{r \rightarrow 0} r^\eta w = c > 0$ , from Proposition 2.8 and 2.9. If  $N = 1$ ,  $\mathcal{T}$  cannot meet  $\mathcal{T}_r$ , thus necessarily  $\lim_{\tau \rightarrow -\infty} \zeta = 0$ , and (2–41) holds with  $a > 0 > b$ .

Up to a change of  $w$  into  $-w$ , the solutions of type (8) and (9) correspond to the trajectories in the domain delimited by  $-\mathcal{T}_r, \mathcal{T}_1, \mathcal{T}_3$ . Indeed they satisfy  $\lim_{\tau \rightarrow \infty} y = -\infty$ , and  $\lim_{r \rightarrow \infty} r^\alpha w = L < 0$ ; and  $\lim_{\tau \rightarrow -\infty} y = 0$ . If  $N \geq 2$ , then  $\lim_{r \rightarrow 0} r^\eta w = c > 0$  and  $w$  has a zero. If  $N = 1$ , either (2–41) holds with  $a = 0 > b$  and  $w$  stays negative, or  $a < 0, b < 0$  and  $w$  has a zero. Such solutions exist from Theorem 2.2. By symmetry, all the solutions are described.

(ii) Now assume  $\alpha = N$  (Figure 1, right). Then  $M_\ell$  belongs to the line  $y = Y$ , and

$$u_1 = (-\delta/(p - 1), -\delta/(p - 1))$$

has the same direction. Moreover  $J_N$  is constant, which means  $y - Y = Ce^{(\delta-N)\tau}$ , with  $C \in \mathbb{R}$ . The solutions corresponding to  $C = 0$  satisfy  $y \equiv Y$ , thus  $\mathcal{T}_1 = \mathcal{T}_r = \{(\xi, \xi) : \xi \in [0, \ell)\}$ , corresponding to the regular Barenblatt solutions. And  $\mathcal{T}_4 = \{(\xi, \xi) : \xi > \ell\}$  yields the solutions defined by (1–6) for  $K < 0$ . All other solutions exist as before, apart from type (7). □

**Note.** The trajectory  $\mathcal{T}_1$  is the only one joining the stationary points  $(0, 0)$  and  $M_\ell$ . Hence, for  $\alpha < N$ , solutions  $w$  of type (1) are unique, up to the scaling mentioned in the note on page 210. Solutions of types (2), (4), and (5) are also unique.

$$N < \alpha < \delta$$

Here we prove that some periodic trajectories can exist, according to the value of  $\alpha$  with respect to  $\alpha^*$ . By (2–32),  $N < \alpha^*$  whenever  $\delta^2 - (N + 3)\delta + N > 0$ , and in particular for any  $p \leq \frac{3}{2}$ . Our main tool is the Poincaré–Bendixson theorem, using the level curves of the energy function  $\mathcal{W}$ :

**Lemma 3.3.** *Assume  $\varepsilon = 1$  and  $N < \alpha < \delta$ . Consider, for  $k \in \mathbb{R}$ , the level curves*

$$\mathcal{C}_k = \{(y, Y) \in \mathbb{R}^2 : \mathcal{W}(y, Y) = k\}$$

*of the function  $\mathcal{W}$  defined in (2–21). They are symmetric with respect to  $(0, 0)$ . Let*

$$k_\ell = \mathcal{W}(\ell, (\delta\ell)^{p-1}) = \frac{1}{2}(\delta - N)\delta^{p-2}\ell^p.$$

*If  $k > k_\ell$ , then  $\mathcal{C}_k$  has two unbounded connected components. If  $0 < k < k_\ell$ ,  $\mathcal{C}_k$  has three connected components, of which one is bounded. If  $k = k_\ell$ ,  $\mathcal{C}_{k_\ell}$  is connected with a double point at  $M_\ell$ . If  $k = 0$ , one of the three connected components of  $\mathcal{C}_0$  is  $\{(0, 0)\}$ . If  $k < 0$ ,  $\mathcal{C}_k$  has two unbounded connected components.*

*Proof.* The energy  $k_\ell$  of the statement is positive. Also  $(y, Y) \in C_k$  if and only if  $F(y) = k - G(y)$ , where  $F, G$  are defined in (2–45). By symmetry we can reduce the study of  $C_k$  to the set  $y > 0$ . Let  $\varphi(s) = |s|^{p'/p'} - s + 1/p$  for any  $s \in \mathbb{R}$ , and set  $\theta = Y/(\delta y)^{p-1}$ . Then (2–44) reduces to

$$\varphi(\theta) = (k - G(y))/(\delta y)^p.$$

The function  $\varphi$  is decreasing on  $(-\infty, 1)$  from  $\infty$  to 0, and increasing on  $(1, \infty)$  from 0 to  $\infty$ . Let  $\psi_1$  be the inverse of the restriction of  $\varphi$  to  $(-\infty, 1]$  and  $\psi_2$  the inverse of the restriction of  $\varphi$  to  $[1, \infty)$ , both defined on  $[0, \infty)$ . For any  $y > 0$ ,

$$y \in \mathcal{C}_k \iff Y = \Phi_1(y) < (\delta y)^{p-1} \text{ or } Y = \Phi_2(y) \geq (\delta y)^{p-1},$$

where

$$(3-1) \quad \Phi_i(y) = (\delta y)^{p-1} \psi_i\left(\frac{k - G(y)}{(\delta y)^p}\right) \text{ for } i = 1, 2,$$

$\Phi_1$  lies below  $\mathcal{M}$  whereas  $\Phi_2$  lies above  $\mathcal{M}$ , and  $\Phi_1, \Phi_2 \in C^1((0, \infty))$ . The function  $G$  has a maximal point at  $y = \ell$ , and  $G(\ell) = k_\ell$ . Using symmetry we see that either  $k > k_\ell$  and  $y$  ranges over  $\mathbb{R}$ , in which case  $\mathcal{C}_k$  has two unbounded connected components; or  $0 < k < k_\ell$  and  $\mathcal{C}_k$  has three connected components, one of which,  $\mathcal{C}_k^b$ , is bounded; or  $k = k_\ell$  and  $\mathcal{C}_{k_\ell}$  is connected with a double point at  $M_\ell$ ; or yet  $k = 0$  and one of the three connected components of  $\mathcal{C}_0$  is  $\{(0, 0)\}$ ; or  $k < 0$  and  $\mathcal{C}_k$

has two unbounded connected components. The unbounded components satisfy  $\lim_{|y| \rightarrow \infty} Y/y^{2/p'} = \pm(p'(\delta\alpha)/2)^{1/p'}$ , by (3-1). The zeros of  $\Phi'_i$  are contained in

$$\mathcal{N} = \{(y, Y) \in \mathbb{R}^2 : y > 0, \delta Y = -(\delta - \alpha)y + (2\delta - N)(\delta y)^{p-1}\},$$

and  $\mathcal{N}$  lies above  $\mathcal{M}$  as long as  $y < \ell$ .

We now describe  $\mathcal{C}_k^b$  when  $0 < k \leq k_\ell$ . The function  $\Phi_1$  is increasing on a segment  $[0, \bar{y}]$  such that  $\bar{y} < \ell$ , and  $\Phi_1(0) = -(kp')^{1/p'}$  and  $(\bar{y}, \Phi_1(\bar{y})) \in \mathcal{M}$ , with an infinite slope at this point;  $\Phi_2$  is increasing on some interval  $[0, \tilde{y}]$  such that  $(\tilde{y}, \Phi_2(\tilde{y})) \in \mathcal{N}$  and then decreasing on  $(\tilde{y}, \bar{y}]$ , and  $\Phi_2(0) = (kp')^{1/p'}$  and  $\Phi_2(\bar{y}) = \Phi_1(\bar{y})$ . By symmetry with respect to  $(0, 0)$ , the curve  $\mathcal{C}_k^b$  is completely described.

Next consider  $\mathcal{C}_{k_\ell}$  for  $y > 0$ : the function  $\Phi_2$  is increasing on  $[0, \infty)$  from  $(p'k_\ell)^{1/p'}$  to  $\infty$ , and  $\Phi_2(\ell) = (\delta\ell)^{p-1}$ ; the function  $\Phi_1$  is increasing on some interval  $[0, \hat{y}]$  such that  $(\hat{y}, \Phi_1(\hat{y})) \in \mathcal{N}$ , so  $\hat{y} > \ell$ ; and  $(\hat{y}, \Phi_1(\hat{y}))$  is below  $\mathcal{M}$ , and  $\Phi_1(\ell) = (\delta\ell)^{p-1}$ , and  $\Phi_1$  is decreasing on  $(\hat{y}, \infty)$ , with  $\lim_{y \rightarrow \infty} \Phi_1 = -\infty$ . Setting  $\mathcal{C}_{k_\ell,1} = \{(y, \Phi_1(y)) : y > \ell\}$  and  $\mathcal{C}_{k_\ell,2} = \{(y, \Phi_2(y)) : y > \ell\}$ , one has  $\mathcal{C}_{k_\ell} = \mathcal{C}_{k_\ell}^b \cup \pm\mathcal{C}_{k_\ell,1} \cup \mathcal{C}_{k_\ell,2}$ . □

**Theorem 3.4.** *Assume  $\varepsilon = 1$  and  $N < \alpha < \delta$ . Then  $w(r) = lr^{-\delta}$  is a solution of  $(E_w)$ .*

- (i) *If  $\alpha \leq \alpha^*$ , any solution of  $(E_w)$  has at most a finite number of zeros.*
- (ii) *There exist  $\check{\alpha}$  such that  $\max(N, \alpha^*) < \check{\alpha} < \delta$ , such that if  $\alpha > \check{\alpha}$ , in the phase plane  $(y, Y)$ , there exists a cycle surrounding  $(0, 0)$ .*
- (iii) *Let  $\alpha$  be such that there exists no such cycle. Then all regular solutions have a finite positive number of zeros and  $\lim_{r \rightarrow \infty} r^\alpha w = L_r \neq 0$  or  $\lim_{r \rightarrow \infty} r^\delta w = \pm \ell$ . There exist solutions of types (2)–(6) of Theorem 3.2, and solutions such that*
  - (1') *(only if  $L_r \neq 0$ )  $\lim_{r \rightarrow 0} r^\delta w = \ell$ , and  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  (or (2-41) holds if  $N = 1$ );*
  - (7')  *$\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  (or (2-41) holds if  $N = 1$ ) and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ .*
- (iv) *For any  $\alpha$  such that there exists such a cycle, there exist solutions  $w$  which oscillate near 0 and  $\infty$ , and  $r^\delta w$  is periodic in  $\ln r$ . All regular solutions  $w$  oscillate near  $\infty$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ . There exist solutions of types (2), (4), (5), and solutions*
  - (1'') *with precisely one zero,  $R_w > 0$ , and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;*
  - (3'') *such that  $\lim_{r \rightarrow 0} r^\delta w = \ell$ , and oscillating near  $\infty$ ;*
  - (9) *such that  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  (or (2-41) holds if  $N = 1$ ) and oscillating near  $\infty$ ;*
  - (10) *with precisely one zero,  $R_w > 0$ , and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ ;*
  - (11) *with  $R_w > 0$  and oscillating near  $\infty$ .*

*Proof.* There always exist solutions of type (2), (4), and (5), by Lemma 3.1.

(i) Assume  $\alpha \leq \alpha^*$  (see Figure 2, left). Consider any trajectory  $\mathcal{T}$ . Suppose  $y$  has infinitely many zeros near  $\pm\infty$ . From Proposition 2.10,  $\mathcal{T}$  is contained in the set

$$\mathcal{D} = \{(y, Y) \in \mathbb{R}^2 : |y| < \ell, |Y| < (\delta\ell)^{p-1}\}$$

near  $\pm\infty$ . Then  $\mathcal{T}$  is bounded near  $\pm\infty$ , hence the limit set at  $\pm\infty$  is contained in  $\mathcal{D}$ . But  $M_\ell \notin \mathcal{D}$ , and  $(0, 0)$  is a source and a node point, so it cannot be in the limit set  $\Gamma$  at  $\infty$ . From the Poincaré–Bendixson theorem,  $\Gamma$  is a closed orbit, so that there exists a cycle. Moreover, from (2–25), (2–49) and (2–50),

$$\frac{\partial f_1}{\partial y}(y, Y) + \frac{\partial f_2}{\partial Y}(y, Y) = \frac{1}{p-1}(D^{(2-p)/(p-1)} - |Y|^{(2-p)/(p-1)});$$

thus, by Bendixson’s criterion, the set  $\{|Y| < D\}$  contains no cycle. Now note that


$$(3-2) \quad \alpha \leq \alpha^* \iff (\delta\ell)^{p-1} \leq D.$$

Then there is no cycle in  $\mathcal{D}$ , and we reach a contradiction.

(ii) Now assume  $\alpha > \max(N, \alpha^*)$ . The curve  $\mathcal{L}$  intersects  $\mathcal{M}$  at  $(\delta^{-1}D^{1/(p-1)}, D)$ . Then

$$\mathcal{S}_{\mathcal{L}} \cap \mathcal{M} = \{(\delta^{-1}(\theta D)^{1/(p-1)}, \theta D) : \theta \in [0, 1]\};$$

and  $D < (\delta\ell)^{p-1}$  by (3–2), so  $\mathcal{S}_{\mathcal{L}}$  does not contain  $M_\ell$ . We can find  $k_1 > 0$  small enough that  $\mathcal{C}_{k_1}^b$  is interior to  $\mathcal{S}_{\mathcal{L}}$ . Next we search for  $k \in (0, k_\ell)$  such that  $\mathcal{L}$  is in the domain delimited by  $C_k^b$ . By symmetry we only consider the points of  $\mathcal{L}$  such that  $y \geq 0$ . In any case for any point of  $\mathcal{L}$  we have  $|\delta y|^p + |Y|^{p'} \leq M = (2(2\delta - N))^\delta$ , by (2–23) and by convexity. By a straightforward computation this implies that



**Figure 2.** Theorem 3.4:  $\varepsilon = 1, N = 2 < \alpha < \delta = 3$ .

${}^{\circ}W(y, Y) \leq KM$ , where  $K = \max(2/p', (3\delta - N)/\delta p)$ . Let  $\check{\alpha} = \check{\alpha}(\delta, N)$  be given by  $KM = k_{\ell}$ . This means that

$$\delta - \check{\alpha} = \left( \frac{\delta - N}{2K\delta^{2-p}} \right)^{1/\delta} \frac{\delta^{p-1}(\delta - N)}{2(2\delta - N)}.$$

If  $\alpha > \check{\alpha}$ , there exists  $k_2 < k_{\ell}$  such that  $\mathcal{L}$  is contained in the set

$$\{(y, Y) \in \mathbb{R}^2 : {}^{\circ}W(y, Y) < k_2\},$$

which has three connected components; because  $\mathcal{S}_{\mathcal{L}}$  is connected, it is contained in the interior to  $\mathcal{C}_{k_2}^b$ . Then the domain delimited by  $\mathcal{C}_{k_1}^b$  and  $\mathcal{C}_{k_2}^b$  is bounded and forward invariant. It does not contain any stationary point, and so it contains a cycle, by the Poincaré–Bendixson theorem (see Figure 2, right).

(iii) Let  $\alpha$  be such that there exists no cycle. Since  $N < \alpha$ , all regular solutions  $y$  have at least one zero. They have a finite number of zeros. For if not,  $(y, Y)$  is bounded near  $\infty$ , so it has a limit cycle. Then either  $\lim_{\tau \rightarrow \infty} y = \pm\infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha > 0$ , so that the trajectory  $\mathcal{T}_r$  ends up in  $\mathcal{Q}_1 \cup \mathcal{Q}_3$  and  $\lim r^{\alpha} w = L_r \neq 0$ , or else  $\lim_{\tau \rightarrow \infty} y = \pm\ell$  and  $\lim_{r \rightarrow \infty} r^{\delta} w = \pm\ell$ .

$\mathcal{T}_3$  cannot meet  $\mathcal{T}_r$  or  $-\mathcal{T}_r$ , thus  $y$  has a unique zero, and  $\lim_{\tau \rightarrow \infty} y = -\infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . The same happens for the trajectories  $\mathcal{T}_6$  in the domain delimited by  $\mathcal{T}_3, \mathcal{T}_4$ . Thus there exist solutions of types (3) and (6).

Suppose  $L_r \neq 0$  and consider  $\mathcal{T}_1$ : the trajectories  $\mathcal{T}_r, -\mathcal{T}_r, \mathcal{T}_1$  have a last intersection point at time  $\tau_0$  with the half-axis  $\{y = 0, Y < 0\}$  at some points  $P_r, P'_r, P_1$ , and  $P_1 \in [P_r, P'_r]$ . The domain delimited by  $\mathcal{T}_r, -\mathcal{T}_r$  and  $[P_r, P'_r]$  is bounded and backward invariant, by Remark 2.1(i) on page 211. Then  $\mathcal{T}_1$  stays in it for  $\tau < \tau_0$ , it has a finite number of zeros, and converges to  $(0, 0)$  near  $-\infty$ ; thus  $w$  is of type (1'). If  $N \geq 2$ , then  $\lim_{\tau \rightarrow \infty} \zeta = \eta$ , so that  $y$  has at least one zero.

Since  $(0, 0)$  is a source, there exist other solutions converging to  $(0, 0)$  near  $-\infty$ , they have a finite number of zeros, and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ , and  $w$  is of type (7').

(iv) Let  $\alpha$  such that there exists a cycle, thus  $\mathcal{T}_r$  has a limit cycle  $\mathcal{C}$ .

Consider again  $\mathcal{T}_1$ . Since  $M_{\ell} \notin \mathcal{S}_{\mathcal{L}}$ , the function  $W$  is decreasing near  $\infty$ , so that  $W(\tau) > k_{\ell}$ ; thus  $\mathcal{T}_1$  is exterior to  $\mathcal{C}_{k_{\ell}}^b$  for large  $\tau$ , in the domain exterior to  $\mathcal{C}_{k_{\ell}}^b$  delimited by  $\mathcal{C}_{k_{\ell},1}$  and  $-\mathcal{C}_{k_{\ell},2}$ ; and it cannot cut  $\mathcal{C}_{k_{\ell}}$ . Moreover  $y$  is decreasing as long as  $y > 0$ , then  $\mathcal{T}_1$  enters  $\mathcal{Q}_4$  as  $\tau$  decreases. It cannot stay in it, because it would converge to  $(0, 0)$ , which is impossible. Then  $y$  has at least one zero, and  $\mathcal{T}_1$  enters  $\mathcal{Q}_3$ . It stays in it, since it cannot cross  $-\mathcal{C}_{k_{\ell},2}$ . Thus  $y$  has a unique zero, and  $\lim_{\tau \rightarrow -\infty} y = -\infty$ , and  $R_w > 0$  from Proposition 2.8, because  $\mathcal{T}_1$  cannot converge to  $(0, 0)$  at  $-\infty$ , and  $w$  is of type (1'').

Next consider  $\mathcal{T}_3$ . Here  $W$  is decreasing near  $-\infty$ , hence  $W(\tau) < k_{\ell}$ ; thus  $\mathcal{T}_3$  is in the interior of  $\mathcal{C}_{k_{\ell}}^b$  near  $-\infty$ . Now the domain delimited by  $\mathcal{C}_{k_1}^b$  and  $\mathcal{C}_{k_{\ell}}^b$  is

forward invariant, thus  $\mathcal{T}_3$  stays in it; then it is bounded, and has a limit cycle at  $\infty$ , and  $w$  is of type (3'').

The solutions of type (9) correspond to trajectories  $\mathcal{T}$  in the domain delimited by  $\mathcal{O}$ , and distinct from  $\mathcal{T}_r$ . Indeed  $\mathcal{T}$  is bounded, in particular the limit-set at  $-\infty$  is  $(0, 0)$ , or a closed orbit. But  $\mathcal{T}$  cannot intersect  $\mathcal{T}_r$ . Then  $\mathcal{T}$  converges to  $(0, 0)$  near  $-\infty$ .

The solutions of type (10) correspond to a trajectory  $\mathcal{T}$  in the domain delimited by  $\mathcal{T}_1 \cup \mathcal{T}_2$  (or its opposite): indeed  $y$  has constant sign near  $\infty$  and near  $\ln R_w$ , and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ , and  $R_w > 0$ , from Proposition 2.8. Then  $\mathcal{T}$  starts in  $\mathcal{Q}_3$ , and ends up in  $\mathcal{Q}_1$ ; and  $y$  has at most one zero, because at such a point  $y' = -|Y|^{1/(p-1)}Y > 0$ , thus it has precisely one zero.

Solutions of type (11) correspond to a trajectory  $\mathcal{T}$  in the domain delimited by  $\mathcal{T}_1, \mathcal{T}_4, -\mathcal{T}_1, -\mathcal{T}_4$ . Then  $y$  cannot have constant sign near  $\infty$ : indeed this implies  $\lim \zeta = \alpha > 0$ ; this is impossible since the line  $Y = y$  is an asymptotic direction for  $\mathcal{T}_1, \mathcal{T}_4$ . Thus  $\mathcal{T}$  is bounded near  $\infty$ , and it has a limit cycle at  $\infty$ . Near  $-\infty$ ,  $y$  a constant sign, because  $\mathcal{T}$  cannot meet  $\mathcal{T}_3$ ; and  $R_w > 0$  from Proposition 2.8, and  $\mathcal{T}$  has the same asymptotic direction  $Y = y$  as  $\mathcal{T}_1, \mathcal{T}_4$ . □

**Note.** From numerical studies, we conjecture that  $\check{\alpha}$  is unique, and the number of zeros of  $w$  increases with  $\alpha$  in the range  $(N, \check{\alpha})$ ; and moreover there exists  $\alpha_1 = N < \alpha_2 < \dots < \alpha_n < \alpha_{n+1} < \dots$ , such that regular solutions have  $n$  zeros for any  $\alpha \in (\alpha_n, \alpha_{n+1})$ , with  $\lim_{r \rightarrow \infty} r^\alpha w = L_r \neq 0$ , and  $n + 1$  zeros for  $\alpha = \alpha_{n+1}$ , with  $\lim_{r \rightarrow \infty} r^\delta w = \pm \ell$ .

$$\alpha \leq \delta \leq N, \alpha \neq N$$


Here  $(0, 0)$  is the only stationary point, and  $N \geq 2$ .

**Theorem 3.5.** *Assume  $\varepsilon = 1$  and  $-\infty < \alpha \leq \delta \leq N, \alpha \neq 0, N$ . Then all regular solutions of  $(E_w)$  have constant sign, and the positive ones satisfy  $\lim_{r \rightarrow \infty} r^\alpha w(r) = L > 0$  if  $\alpha \neq \delta$ , or (2-39) holds if  $\alpha = \delta$ . All the other solutions have a reduced domain ( $R_w > 0$ ). Among them, there exist solutions satisfying any one of these characterizations:*

- (1)  $w$  is positive,  $\lim_{r \rightarrow \infty} r^\eta w = c \neq 0$  if  $\delta < N$ , or  $\lim_{r \rightarrow \infty} r^N (\ln r)^{(N+1)/2} w = \varrho$  defined in (2-40) if  $\delta = N$ ;
- (2)  $w$  is positive,  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$  if  $\alpha \neq \delta$ , or (2-39) holds if  $\alpha = \delta$ ;
- (3)  $w$  has one zero, and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$  if  $\alpha \neq \delta$ , or (2-39) holds if  $\alpha = \delta$ .

*Up to symmetry, all the solutions are as above.*

*Proof.* Any solution has at most one zero, by Proposition 2.5. The trajectory  $\mathcal{T}_r$  starts in  $\mathcal{Q}_4$  if  $\alpha < 0$  (Figure 3, left) and in  $\mathcal{Q}_1$  if  $\alpha > 0$  (Figure 3, right). Moreover  $y$



**Figure 3.** Theorem 3.5:  $\varepsilon = 1$ ,  $\alpha < \delta = 3 < N = 4$ .

stays positive, and  $\lim_{\tau \rightarrow \infty} y = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ , by Proposition 2.13. Then  $\lim_{r \rightarrow \infty} r^\alpha w(r) = L > 0$  if  $\alpha < \delta$ , or (2–39) holds if  $\alpha = \delta$ , from Proposition 2.9. Moreover  $y$  is increasing: indeed it has no local maximum from (2–16). Thus  $\mathcal{T}_r$  does not meet  $\mathcal{M}$ , and so stays below  $\mathcal{M}$ . If  $\alpha > 0$ , then  $\mathcal{T}_r$  stays in  $\mathcal{Q}_1$ , and  $Y$  is increasing from 0 to  $\infty$ ; indeed each extremal point  $\tau$  of  $Y$  is a local minimum, by (2–17). Likewise, if  $\alpha < 0$  the function  $Y$  is decreasing from 0 to  $-\infty$ , and  $\mathcal{T}_r$  stays in  $\mathcal{Q}_4$ . The only solutions  $y$  defined on  $(0, \infty)$  are the regular ones, by Proposition 2.8.

For any point  $P = (\varphi, (\delta\varphi)^{p-1}) \in \mathbb{R}^2$  with  $\varphi > 0$ , in other words on the curve  $\mathcal{M}$ , the trajectory  $\mathcal{T}_{[P]}$  intersects  $\mathcal{M}$  transversally: the vector field is  $(0, -(N-\alpha)\varphi)$ . Moreover the solution going through this point at time  $\tau_0$  satisfies  $y''(\tau_0) > 0$  from  $(E_y)$ , then  $\tau_0$  is a point of local minimum. From (2–16),  $\tau_0$  is unique, so that it is a minimum. Then  $y > 0$ ,  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ ,  $\lim_{\tau \rightarrow \ln R_w} Y/y = 1$ , and  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_1$  if  $\alpha > 0$ , or goes from  $\mathcal{Q}_1$  into  $\mathcal{Q}_3$  if  $\alpha < 0$ . The corresponding  $w$  is of type (2).

For any point  $P = (0, \xi)$ ,  $\xi > 0$ , the trajectory  $\mathcal{T}_{[P]}$  goes through  $P$  from  $\mathcal{Q}_1$  into  $\mathcal{Q}_2$ , by Remark 2.1(i). Then  $y$  has only one zero, and as above, it is decreasing on  $\mathbb{R}$  and  $\lim_{\tau \rightarrow \infty} y = -\infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ ,  $\lim_{\tau \rightarrow \ln R_w} Y/y = 1$ . Thus  $\mathcal{T}_{[P]}$  starts in  $\mathcal{Q}_1$ , then stays in  $\mathcal{Q}_2$  if  $\alpha < 0$ , and enters  $\mathcal{Q}_3$  and stays in it if  $\alpha > 0$ . The corresponding  $w$  is of type (3).

It remains to prove the existence of a solution of type (1). If  $\delta < N$ , then  $(0, 0)$  is a saddle point. There exists a trajectory  $\mathcal{T}_1$  converging to  $(0, 0)$  at  $\infty$ , with  $y > 0$ , and  $\lim_{\tau \rightarrow \infty} \zeta = \eta > 0$ , thus in  $\mathcal{Q}_1$  near  $\infty$ , with  $y' < 0$ . As above,  $y$  has no local maximum, it is increasing, so that  $y > 0$ . If  $\delta = N$ , we consider the sets

$$\begin{aligned} \mathcal{A} &= \{P \in (0, \infty) \times \mathbb{R} : \mathcal{T}_{[P]} \cap \mathcal{M} \neq \emptyset\}, \\ \mathcal{B} &= \{P \in (0, \infty) \times \mathbb{R} : \mathcal{T}_{[P]} \cap \{(0, \xi) : \xi > 0\} \neq \emptyset\}. \end{aligned}$$

They are nonempty, and open, because the intersections are transverse. Since  $\mathcal{T}_r$  is below  $\mathcal{M}$ , the sets  $\mathcal{A}$  and  $\mathcal{B}$  are contained in the domain  $\mathcal{R}$  of  $\mathcal{Q}_1 \cup \mathcal{Q}_2$  above  $\mathcal{T}_r$ , and  $\mathcal{A} \cup \mathcal{B} \neq \mathcal{R}$ . As a result there exists at least a trajectory  $\mathcal{T}_1$  above  $\mathcal{T}_r$ , which does not intersects  $\mathcal{M}$  and the set  $\{(0, \xi) : \xi > 0\}$ . The corresponding  $y$  is monotone. Suppose that  $y$  is increasing, then  $\lim_{\tau \rightarrow -\infty} y = 0$ ; it is impossible since  $\mathcal{T}_1 \neq \mathcal{T}_r$ . Then  $y$  is decreasing, and  $\lim_{\tau \rightarrow \infty} y = 0$ . In any case  $w$  is of type (1), by Propositions 2.8 and 2.9. All the solutions are described, because any solution has at most one zero, and at most one extremum point. And  $\mathcal{T}_1$  is unique when  $\delta < N$ . □

#### 4. The case $\varepsilon = -1, \delta < \alpha$


$$N < \delta < \alpha$$

**Theorem 4.1.** *Assume  $\varepsilon = -1$  and  $N < \delta < \alpha$ . Then all regular solutions of  $(E_w)$  have constant sign and satisfy  $S_w < \infty$ . And  $w \equiv \ell r^{-\delta}$  is a solution. There exist solutions satisfying any one of these characterizations:*

- (1)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  if  $N \geq 2$  (and  $\lim_{r \rightarrow 0} w = a > 0, \lim_{r \rightarrow 0} w' = b(a) < 0$  if  $N = 1$ ) and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;
- (2)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $S_w < \infty$ ;
- (3)  $w$  has one zero,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $S_w < \infty$ ;
- (4)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$  and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;
- (5)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$  and  $S_w < \infty$ ;
- (6)  $w$  has one zero,  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$  and  $S_w < \infty$ ;
- (7)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  if  $N \geq 2$  (and  $\lim_{r \rightarrow 0} w = a > 0$  for any  $a > 0$  and  $\lim_{r \rightarrow 0} w' = b < 0, b \neq b(a)$  if  $N = 1$ ), and  $S_w < \infty$ ;
- (8)  $w$  has one zero and the same behavior;
- (9) (only if  $N = 1$ )  $w$  is positive,  $\lim_{r \rightarrow 0} w = a > 0$ , and  $\lim_{r \rightarrow 0} w' = b > 0$ , and  $S_w < \infty$ .

*Up to symmetry, all solutions are as above.*

*Proof.* Here we still have three stationary points,  $(0, 0)$  is a source and  $M_\ell$  a saddle point (see Figure 4). By Propositions 2.5 and 2.14, all regular solutions have constant sign and satisfy  $S_w < \infty$ . Also,  $\mathcal{T}_r$  stays in  $\mathcal{Q}_4$  by Remark 2.3, and  $\lim_{\tau \rightarrow \ln S_w} Y/y = -\infty$  by Proposition 2.15. Since  $\alpha > 0$ , any solution  $y$  has at most one zero, by Proposition 2.5, and  $y$  is monotone near  $\ln S_w$  (finite or not) and near  $-\infty$ , by Proposition 2.7. In the linearization near  $M_\ell$  the eigenvectors  $u_1 = (\nu(\alpha), \lambda_1 - \delta)$  and  $u_2 = (-\nu(\alpha), \delta - \lambda_2)$  form a positively oriented basis, where



**Figure 4.** Theorem 4.1:  $\varepsilon = -1$ ,  $N < \delta = 3 < \alpha$ .

now  $v(\alpha) < 0$  and  $\lambda_1 < \delta < \lambda_2$ ; thus  $u_1$  points toward  $\mathcal{Q}_3$  and  $u_2$  points toward  $\mathcal{Q}_4$ . There exist four particular trajectories converging to  $M_\ell$  near  $\pm\infty$ , namely:

- $\mathcal{T}_1$  converging to  $M_\ell$  at  $\infty$ , with tangent vector  $u_1$ . Here  $y$  is increasing near  $\infty$ , and as long as  $y > 0$ ; indeed, if there exists a minimal point  $\tau$ ,  $(E_y)$  shows that  $y(\tau) > \ell$ . And  $\mathcal{T}_1$  stays in  $\mathcal{Q}_1$  on  $\mathbb{R}$ , by Remark 2.1(i) on page 211. Therefore  $\mathcal{T}_1$  converges to  $(0, 0)$  at  $-\infty$ , and  $w$  is of type (1), where  $b(a)$  is a function of  $a$ , by the note on page 210.
- $\mathcal{T}_2$  converging to  $M_\ell$  at  $-\infty$ , with tangent vector  $u_2$ . Here again  $y' > 0$  as long as  $y > 0$ . Also  $Y' < 0$  near  $-\infty$ , and  $Y$  is decreasing as long as  $Y > 0$ : if there exists a minimal point of  $Y$  in  $\mathcal{Q}_1$ ,  $(E_Y)$  shows that  $Y(\tau) > (\delta\ell)^{p-1}$ . But  $(y, Y)$  cannot stay in  $\mathcal{Q}_1$ , as this would imply  $\lim_{\tau \rightarrow \infty} y = \infty$ , which is impossible by Proposition 2.8. Thus  $\mathcal{T}_2$  enters  $\mathcal{Q}_4$  at some point  $(\xi_2, 0)$  with  $\xi_2 > 0$  and stays in it since  $y' > 0$ . Thus  $S_w < \infty$  and  $\lim_{\tau \rightarrow \infty} Y/y = -1$ , and  $w$  is of type (2).
- $\mathcal{T}_3$  converging to  $M_\ell$  at  $-\infty$ , with tangent vector  $-u_2$ . Here again  $y' < 0$  as long as  $y > 0$ . And  $Y' > 0$  as long as  $Y > 0$ ; thus  $Y' > 0$  on  $\mathbb{R}$ . Then again  $(y, Y)$  cannot stay in  $\mathcal{Q}_1$ , so  $y$  has a unique zero, and  $\mathcal{T}_3$  enters  $\mathcal{Q}_2$  at some point  $(0, \xi_3)$  with  $\xi_3 > 0$  and stays in it. Hence  $S_w < \infty$  and  $\lim_{\tau \rightarrow \infty} Y/y = -1$ , and  $w$  is of type (3).
- $\mathcal{T}_4$  converging to  $M_\ell$  at  $\infty$ , with tangent vector  $-u_1$ . In the same way,  $y$  is decreasing near  $\infty$ , and  $y$  is everywhere decreasing: if there exists a maximal point  $\tau$ , then  $y(\tau) < \ell$  by  $(E_y)$ . Then  $Y$  stays positive, thus  $\mathcal{T}_4$  stays in  $\mathcal{Q}_1$ . By Proposition 2.8,  $\lim_{\tau \rightarrow -\infty} y = \infty$  and  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha$ , so  $w$  is of type (4).

Next we describe all the other trajectories  $\mathcal{T}_{[P]}$  with one point  $P$  in the domain  $\mathcal{R}$  above  $\mathcal{T}_r \cup (-\mathcal{T}_r)$ .

If  $P = (\varphi, 0)$  with  $\varphi > \xi_2$ , then  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_4$  after  $P$ , because it cannot meet  $\mathcal{T}_2$ ; before  $P$  it stays in  $\mathcal{Q}_1$ , by Remark 2.1(i). Thus again  $S_w = \infty$ , and  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha > 0$ , and  $y$  has a unique minimal point, and  $w$  is of type (5). For any  $P$  is in the domain delimited by  $\mathcal{T}_2, \mathcal{T}_4$ , the trajectory  $\mathcal{T}_{[P]}$  is of the same type.

If  $P = (0, \xi)$  with  $\xi > \xi_3$ , then  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_2$  after  $P$ , in  $\mathcal{Q}_1$  before  $P$ , since it cannot meet  $\mathcal{T}_2, \mathcal{T}_4$ . Then  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha > 0$ , and  $S_w = \infty$ , and  $w$  is of type (6). If  $P$  is in the domain delimited by  $\mathcal{T}_3, \mathcal{T}_4$ , then  $\mathcal{T}_{[P]}$  is of the same type.

If  $P = (\varphi, 0)$  with  $\varphi \in (0, \xi_2)$ , then  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_4$  after  $P$ , in  $\mathcal{Q}_1$  before  $P$ ; it cannot meet  $\mathcal{T}_r$ , thus  $S_w < \infty$ ; and  $\mathcal{T}_{[P]}$  converges to  $(0, 0)$  in  $\mathcal{Q}_1$  at  $-\infty$ ; thus  $w$  is of type (7), by Theorem 2.2. If  $P$  is in the domain delimited by  $\mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_r$ , then  $\mathcal{T}_{[P]}$  is of the same type.

If  $P = (0, \xi)$  for some  $\xi \in (0, \xi_3)$ , then  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_2$  after  $P$ , in  $\mathcal{Q}_1$  before  $P$ ; and  $\mathcal{T}$  cannot meet  $-\mathcal{T}_r$ , so that  $S_w < \infty$ . Then  $\mathcal{T}_{[P]}$  converges to  $(0, 0)$  in  $\mathcal{Q}_1$  at  $-\infty$ , and  $w$  is of type (8).

If  $P$  lies in the domain delimited by  $\mathcal{T}_1, \mathcal{T}_3$  and  $-\mathcal{T}_r$ , either  $y$  has one zero, and  $\mathcal{T}_{[P]}$  is of the same type; or  $y < 0$  on  $\mathbb{R}$ , and  $y' = \delta y - Y^{1/(p-1)} < 0$ . Hence  $S_w < \infty$  and  $\mathcal{T}_{[P]}$  converges to  $(0, 0)$  in  $\mathcal{Q}_2$  at  $-\infty$ . It implies  $N = 1$  (see Figure 4, right), and  $-w$  is of type (9), by Propositions 2.8 and 2.9; and such a solution does exist, by Theorem 2.2. Up to symmetry, all the solutions have been obtained. Here again, up to a scaling, the solutions  $w$  of types (1)–(4) are unique.  $\square$


$$\delta \leq \min(\alpha, N) \text{ (apart from } \alpha = \delta = N)$$

**Theorem 4.2.** *Suppose  $\varepsilon = -1$  and  $\delta \leq \min(\alpha, N)$  (apart from  $\alpha = \delta = N$ ). Then all regular solutions of  $(E_w)$  have constant sign and a reduced domain ( $S_w < \infty$ ). There exist solutions satisfying any one of these characterizations:*

- (1)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$  and  $\lim_{r \rightarrow \infty} r^\eta w = c \neq 0$  if  $\delta < N$ , or (2–40) holds if  $\delta = N < \alpha$ ;
- (2)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$  if  $\delta < \alpha$ , or (2–39) holds if  $\alpha = \delta < N$ , and  $S_w < \infty$ ;
- (3)  $w$  has one zero and the same behavior.

*Up to symmetry, all solutions are as above.*

*Proof.* Here  $(0, 0)$  is the only one stationary point, and  $N \geq 2$  (Figure 5). By Propositions 2.5 and 2.14, all regular solutions have constant sign, and  $S_w < \infty$ . Moreover  $w' > 0$  near 0, by Theorem 2.2; and  $w$  can only have minimal points, by Remark 2.3, so  $w' > 0$  on  $(0, S_w)$ . In other words,  $\mathcal{T}_r$  stays in  $\mathcal{Q}_4$ , and  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ . By Propositions 2.5 and 2.7, any solution  $y$  has at most one zero and is monotone at the extremities. By Proposition 2.8, apart from  $\mathcal{T}_r$ , any



**Figure 5.** Theorem 4.2:  $\varepsilon = -1$ ,  $\delta = 3 < N = 4$ .

trajectory  $\mathcal{T}$  satisfies  $\lim_{\tau \rightarrow -\infty} |y| = \infty$ , and so  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha > 0$ ; hence  $\mathcal{T}$  starts from  $\mathcal{Q}_1$  or  $\mathcal{Q}_3$  at  $-\infty$ .

For any  $P = (\varphi, 0)$  with  $\varphi > 0$ , the trajectory  $\mathcal{T}_{[P]}$  goes from  $\mathcal{Q}_1$  into  $\mathcal{Q}_4$  at  $P$ , by Remark 2.1(i) on page 211; it stays in  $\mathcal{Q}_4$  after  $P$ , since it cannot meet  $\mathcal{T}_r$ ; and it stays in  $\mathcal{Q}_1$  before  $P$ : it cannot start from  $\mathcal{Q}_3$ , because it does not meet  $-\mathcal{T}_r$ . Thus  $y$  remains positive and  $w$  is of type (2).

For any  $P = (0, \xi)$  with  $\xi > 0$ ,  $\mathcal{T}_{[P]}$  goes from  $\mathcal{Q}_1$  into  $\mathcal{Q}_2$  by the same remark; thus  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_2$  after  $P$ , since it cannot meet  $-\mathcal{T}_r$ , and in  $\mathcal{Q}_1$  before  $P$ , and  $w$  is of type (3).

It remains to prove the existence of solutions of type (1). If  $\delta < N$ , the origin is a saddle point, so there exists a trajectory  $\mathcal{T}_1$  converging to  $(0, 0)$  at  $\infty$ ; and  $\lim_{\tau \rightarrow \infty} \zeta = \eta > 0$ , by Proposition 2.8. Thus  $\mathcal{T}_1$  lies in  $\mathcal{Q}_1$  for large  $\tau$ , and stays there, because  $\mathcal{Q}_1$  is backward invariant. The conclusion follows. If  $\delta = N$ , we consider the sets


$$\begin{aligned} \mathcal{A} &= \{P \in \mathcal{Q}_1 : \mathcal{T}_{[P]} \cap \{(\varphi, 0) : \varphi > 0\} \neq \emptyset\}, \\ \mathcal{B} &= \{P \in \mathcal{Q}_1 : \mathcal{T}_{[P]} \cap \{(0, \xi) : \xi > 0\} \neq \emptyset\}. \end{aligned}$$

They are nonempty and open, since the vector field is transverse at  $(\varphi, 0)$  and  $(0, \xi)$ ; thus  $\mathcal{A} \cup \mathcal{B} \neq \emptyset$ . Hence there exists a trajectory  $\mathcal{T}_1$  staying in  $\mathcal{Q}_1$ ; therefore  $S_w = \infty$  and  $\mathcal{T}_1$  converges to  $(0, 0)$  at  $\infty$ , and  $w$  is of type (1), by Proposition 2.9. All solutions have been described, up to symmetry.  $\square$

### 5. The case $\varepsilon = 1$ , $\delta \leq \alpha$

$$N \leq \delta \leq \alpha$$

**Theorem 5.1.** Assume  $\varepsilon = 1$ ,  $N \leq \delta \leq \alpha$  and  $\alpha \neq N$ .



**Figure 6.** Theorem 5.1:  $\varepsilon = 1, < \delta = 3 < \alpha = 3.5$ .


- (i) *There exists a cycle surrounding  $(0, 0)$ , and thus also solutions  $w$  of  $(E_w)$  with changing sign and such that  $r^\delta w$  is periodic in  $\ln r$ . All other solutions  $w$ , in particular the regular ones, are oscillating near  $\infty$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ . There exist solutions  $w$  such that  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  if  $2 \leq N < \delta$  and (2–40) holds if  $N = \delta$ , or (2–41) holds if  $N = 1$ .*
- (ii) *There exist solutions such that  $R_w > 0$ , or  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$  if  $\alpha \neq \delta$ , or (2–39) holds if  $\alpha = \delta$ .*

*Proof.* (i) Here  $(0, 0)$  is the only stationary point. From Proposition 2.8, any trajectory is bounded and  $y$  is oscillating around 0 near  $\infty$ .

First assume  $N < \delta < \alpha$  (Figure 6, left). Then  $(0, 0)$  is a source and all trajectories have a limit cycle at  $\infty$  or are periodic. In particular there exists at least one cycle, with orbit  $\mathbb{O}_p$ . The trajectory  $\mathcal{T}_r$  has a limit cycle  $\mathbb{O} \subseteq \mathbb{O}_p$ . There exist also trajectories  $\mathcal{T}_s$  starting from  $(0, 0)$  with an infinite slope, such that  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$  if  $N \geq 2$  or (2–41) if  $N = 1$ , and all the  $\mathcal{T}_s$  have the same limit cycle  $\mathbb{O}$ .

Next assume  $N = \delta < \alpha$  (Figure 6, right). Then  $\mathcal{T}_r$  cannot converge to  $(0, 0)$ , since it would intersect itself. Thus again the limit set at  $\infty$  is a closed orbit  $\mathbb{O}$ . No trajectory can converge to  $(0, 0)$  at  $\infty$ , as it would spiral around this point and hence intersect  $\mathcal{T}_r$ . Consider any trajectory  $\mathcal{T} \neq \mathcal{T}_r$  in the connected component of  $\mathbb{O}$  containing  $(0, 0)$ .  $\mathcal{T}$  is bounded, so its limit set at  $-\infty$  is  $(0, 0)$  or a closed orbit. The second case is impossible, since  $\mathcal{T}$  does not meet  $\mathcal{T}_r$ . Thus  $\mathcal{T}$  is of the form  $\mathcal{T}_s$ , and the corresponding  $w$  satisfies (2–40).

(ii) By Theorem 2.21, all cycles are contained in a ball  $B$  of  $\mathbb{R}^2$ . Take any point  $P_0$  exterior to  $B$ . Then  $\mathcal{T}_{[P_0]}$  has a limit cycle at  $\infty$  contained in  $B$ . If it has a limit cycle at  $-\infty$ , it is contained in  $B$ , so  $\mathcal{T}_{[P_0]}$  is contained in  $B$ , which is impossible. Thus  $y$  has constant sign near  $\ln R_w$ . By Proposition 2.8, either  $R_w > 0$  or  $y$  is defined near  $-\infty$ . □



**Figure 7.** Theorem 5.2:  $\varepsilon = 1, \alpha = \delta = N = 3$ .

**Theorem 5.2.** Assume  $\varepsilon = 1$  and  $\alpha = \delta = N$ . All regular solutions of  $(E_w)$  have constant sign, and are given by (1–6). For any  $k \in \mathbb{R}$ ,  $w(r) = kr^{-N}$  is a solution. There exist solutions satisfying any one of these characterizations:

- (1)  $w$  is positive,  $\lim_{r \rightarrow 0} r^N w = c_1 > 0$ ,  $\lim_{r \rightarrow \infty} r^N w = c_2 > 0$  ( $c_2 \neq c_1$ );
- (2)  $w$  has one zero,  $\lim_{r \rightarrow 0} r^N w = c_1 > 0$  and  $\lim_{r \rightarrow \infty} r^N w = c_2 < 0$ ;
- (3)  $w$  is positive,  $R_w > 0$ , and  $\lim_{r \rightarrow 0} r^N w = c \neq 0$ ;
- (4)  $w$  has one zero and the same behavior.

Up to symmetry, all solutions are as above.

*Proof.* Since  $\alpha = N$ , equation  $(E_w)$  admits the first integral (1–5), so  $J_N \equiv C$  for  $C \in \mathbb{R}$ . We gave in (1–6) the regular (Barenblatt) solutions for the case  $C = 0$ . Since  $\delta = N$ , (1–5) is equivalent to the equation  $Y \equiv y - C$ , by (2–12) (refer to Figure 7). For any  $k \in \mathbb{R}$ ,  $(y, Y) \equiv (k, |Nk|^{p-2}Nk)$  is a solution of the system  $(S)$  located on the curve  $\mathcal{M}$ , so that  $w(r) = kr^{-N}$  is a solution. Any solution has at most one zero, by Proposition 2.5. By Propositions 2.8 and 2.10, any trajectory converges to a point  $(k, |Nk|^{p-2}Nk)$  of  $\mathcal{M}$  at  $\infty$ . Let  $\bar{C} < 0$  be such that the line  $Y = y - \bar{C}$  is tangent to  $\mathcal{M}$ . For any  $C \in (\bar{C}, 0)$ , the line  $Y = y - C$  cuts  $\mathcal{M}$  at three points  $k_1 < 0 < k_2 < k_3$ . And  $y' > 0$  if the trajectory is below  $\mathcal{M}$  and  $y' < 0$  if it is above  $\mathcal{M}$ . We find two solutions  $y$  defined on  $\mathbb{R}$ : one is positive and  $\lim_{\tau \rightarrow -\infty} y = k_2$ ,  $\lim_{\tau \rightarrow -\infty} y = k_3$ , and the other has one zero. All other solutions satisfy  $R_w > 0$ ,  $\lim_{\tau \rightarrow \ln R_w} Y/y = 1$ ; some of them are positive, the others have one zero.  $\square$

$$\delta < \min(\alpha, N)$$

Here the system has three stationary points:  $(0, 0)$  is a saddle point, while  $M_\ell, M'_\ell$  are sinks if  $\delta \leq N/2$ , or  $N/2 < \delta$  and  $\alpha < \alpha^*$ , and sources when  $N/2 < \delta$  and  $\alpha > \alpha^*$ , and node points whenever  $\alpha \leq \alpha_1$  or  $\alpha_2 \leq \alpha$ , where  $\alpha_1, \alpha_2$  are defined in (2–48) (recall that  $\alpha_1$  can be greater or less than  $\eta$ ). This case is one

of the most delicate, since two types of periodic trajectories can appear, either surrounding  $(0, 0)$ , corresponding to changing sign solutions, or located in  $\mathcal{Q}_1$  or  $\mathcal{Q}_3$ , corresponding to solutions of constant sign. Notice that  $\delta < N$  implies  $\delta < N < \eta$  by (1–2), and  $N/2 < \delta$  implies  $\eta < \alpha^*$  by (2–32).

**Remark 5.3.** (i)  $\mathcal{T}_r$  starts in  $\mathcal{Q}_1$ . Since  $(0, 0)$  is a saddle point, Propositions 2.8 and 2.9 show there is a unique trajectory  $\mathcal{T}_s$  converging to  $(0, 0)$ , residing in  $\mathcal{Q}_1$  for large  $\tau$ , having an infinite slope at  $(0, 0)$ , and satisfying  $\lim_{r \rightarrow 0} r^\eta w = c > 0$ . Moreover if  $\mathcal{T}_r$  does not stay in  $\mathcal{Q}_1$ , then  $\mathcal{T}_s$  stays in it, and is bounded and contained in the domain delimited by  $\mathcal{Q}_1 \cap \mathcal{T}_r$ , by Remark 2.1(i). Thus if  $\mathcal{T}_r$  is homoclinic, it stays in  $\mathcal{Q}_1$ .

(ii) Any trajectory  $\mathcal{T}$  is bounded near  $\infty$ , by Propositions 2.8 and 2.12. From the strong form of the Poincaré–Bendixson theorem [Hubbard and West 1995, p. 239], any trajectory  $\mathcal{T}$  bounded at  $\pm\infty$  either converges to  $(0, 0)$  or  $\pm M_\ell$ , or its limit set  $\Gamma_\pm$  at  $\pm\infty$  is a cycle, or it is homoclinic hence  $\mathcal{T} = \mathcal{T}_r$  and  $\Gamma_\pm = \overline{\mathcal{T}_r}$  (indeed, for any  $P \in \Gamma_\pm$ ,  $\mathcal{T}_{[P]}$  converges at  $\infty$  and  $-\infty$  to  $(0, 0)$  or  $\pm M_\ell$ ; if one of them is  $\pm M_\ell$ , then  $\pm M_\ell \in \overline{\mathcal{T}_{[P]}} \subset \Gamma_\pm$ , and  $M_\ell$  is a source or a sink, so  $\mathcal{T}$  converges to  $\pm M_\ell$ ; otherwise  $\mathcal{T}$  is homoclinic and  $\mathcal{T}_{[P]} = \mathcal{T}_r$ ).

(iii) If there exists a limit cycle around  $(0, 0)$ , then by (2–42) this cycle also surrounds the points  $\pm M_\ell$ .

We begin with the case  $\alpha \leq \eta$ , where there exists no cycle in  $\mathcal{Q}_1$  and no homoclinic orbit, by Theorem 2.20.

**Theorem 5.4.** *Assume that  $\varepsilon = 1$  and  $\delta < \min(\alpha, N)$ , and  $\alpha \leq \eta$ . Then all regular solutions of  $(E_w)$  have constant sign, and  $\lim_{r \rightarrow \infty} r^\delta |w(r)| = \ell$ . And  $w(r) = \ell r^{-\delta}$  is a solution.*


*If  $\alpha < \eta$ , there exist solutions satisfying any one of these characterizations:*

- (1)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = L$  and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;
- (2)  $w$  is positive,  $R_w > 0$  and  $\lim_{r \rightarrow \infty} r^\eta w = c > 0$ ;
- (3)  $w$  is positive,  $R_w > 0$  and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;
- (4)  $w$  has one zero,  $R_w > 0$  and  $\lim_{r \rightarrow \infty} r^\delta w = \ell$ ;

*If  $\alpha = \eta$ , then  $w = Cr^{-\eta}$  is a solution and there exist solutions of type (4), but not of type (2) or (3).*

*Proof.* By Proposition 2.5 and Remark 2.3,  $\mathcal{T}_r$  stays in  $\mathcal{Q}_1$  and converges to  $M_\ell$  at  $\infty$ ; indeed there is no cycle in  $\mathcal{Q}_1$ , by Propositions 2.8, 2.12 and 2.20.

(i) Assume  $\alpha < \eta$  (Figure 8, left). Consider any trajectory in  $\mathcal{Q}_1$ . Then  $Y_\alpha > 0$ . If there exists  $\tau$  such that  $Y'_\alpha(\tau) = 0$ , at this point  $Y''_\alpha(\tau) \geq 0$  by (2–36), and  $\tau$



**Figure 8.** Theorem 5.4:  $\varepsilon = 1, \delta = 3 < N = 4 < \eta = 5$ .

is a local minimum.  $\mathcal{T}_r$  satisfies  $\lim_{\tau \rightarrow -\infty} Y_\alpha = 0$ , and so  $Y'_\alpha > 0$  on  $\mathbb{R}$ . This is equivalent to  $\alpha y > Y^{1/(p-1)} + (p-1)(\eta - \alpha)Y$ . Therefore  $\mathcal{T}_r$  stays strictly below the curve

$$\mathcal{M}_\alpha = \{(y, Y) \in \mathcal{D}_1 : \alpha y = Y^{1/(p-1)} + (p-1)(\eta - \alpha)Y\}.$$

First consider  $\mathcal{T}_s$ . Since  $\alpha < \eta$ , this trajectory satisfies  $\lim_{\tau \rightarrow \infty} Y_\alpha = 0$ . Then  $Y'_\alpha < 0$  on  $(\ln R_w, \infty)$ , so  $\mathcal{T}_s$  stays strictly above  $\mathcal{M}_\alpha$ . Hence it stays above  $\mathcal{M}$ : indeed, if it meets  $\mathcal{M}$  at a first point  $(y_1, (\delta y_1)^{p-1})$ , the function  $y$  has a maximum at this point. Thus by (2–16), we have  $\ell < y_1$  and

$$(\alpha - \delta)y_1^{2-p} = \delta^{p-1}(p-1)(\eta - \alpha) < \delta^{p-1}(p-1)(\eta - \delta),$$

contradicting (1–2) and (1–4). This shows that  $y' < 0$ . Suppose that  $y$  is defined on  $\mathbb{R}$ ; then  $\lim_{\tau \rightarrow -\infty} y = \infty$  and  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha$ . If  $\zeta' > 0$  on  $\mathbb{R}$ , then  $\zeta(\mathbb{R}) = (\alpha, \eta)$ , which contradicts (2–9). Then  $\zeta$  has at least one extremal point  $\tau$ , and  $\zeta(\tau)$  is exterior to  $(\alpha, \eta)$ , by (2–9); if it is a minimum,  $\zeta(\tau) > \alpha$  by (2–18), since  $y' < 0$ , and if it is a maximum,  $\zeta(\tau) < \alpha$ . Thus we reach again a contradiction. Therefore  $R_w > 0$  and  $\lim_{\tau \rightarrow \ln R_w} Y/y = 1$ , and the corresponding  $w$  is of type (2).

For any  $P = (\varphi, 0)$ ,  $\varphi > 0$ , the trajectory  $\mathcal{T}_{[P]}$  stays in  $\mathcal{D}_1$  after  $P$ . The solution  $(y, Y)$  originating at  $P$  at time 0 satisfies  $Y_\alpha(0) = 0$ ; hence  $Y'_\alpha(\tau) > 0$  for any  $\tau \geq 0$ . Thus  $\mathcal{T}_{[P]}$  stays below  $\mathcal{M}_\alpha$ . Moreover it enters  $\mathcal{D}_4$  as  $\tau$  decreases. But  $y' > 0$  in  $\mathcal{D}_4$ , by (S); thus  $\mathcal{T}_{[P]}$  does not stay in  $\mathcal{D}_4$ , by Proposition 2.8; it goes into  $\mathcal{D}_3$  and must stay there because it cannot meet  $-\mathcal{T}_s$ . This shows that  $R_w > 0$  and  $y$  has precisely one zero, and  $w$  is of type (4).

Next consider any trajectory  $\mathcal{T}_{[P_1]}$  going through some point  $P_1 = (y_1, Y_1)$  in  $\mathcal{D}_1$ , lying below  $\mathcal{T}_s$  and such that  $\alpha y_1 < Y_1^{1/(p-1)}$ . Such a trajectory exists because  $y = Y$  is an asymptotic direction of  $\mathcal{T}_s$ . Let  $(y, Y)$  be the solution issuing from  $P_1$  at time 0. Suppose  $y$  is defined on  $\mathbb{R}$ ; then  $\lim_{\tau \rightarrow -\infty} y = \infty$  and  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha$ .

Also,  $\zeta(0) > \alpha$ . Then  $\zeta > \delta$  on  $(-\infty, 0)$ : otherwise there would exist  $\tau < 0$  such that  $\zeta(\tau) = \alpha$  and  $\zeta'(\tau) \geq 0$ , contradicting (2–9). Thus  $y' < 0$  on  $(-\infty, \tau_1)$ . Either  $\zeta' > 0$  on  $(-\infty, 0)$ , in which case  $\zeta > \eta > 0$  by (2–9), which is impossible; or  $\zeta$  has at least an extremal point  $\tau$ . If it is a minimum, then  $\zeta(\tau) > \alpha$  from (2–18); if it is a maximum, then  $\zeta(\tau) < \alpha$ ; and again we reach a contradiction. Therefore  $R_w > 0$ , and the trajectory stays in  $\mathfrak{D}_1$  and converges to  $M_\ell$ , because there is no cycle in  $\mathfrak{D}_1$ , by Theorem 2.20. Hence  $w$  is of type (3).

Let  $\mathfrak{O}$  be the domain of  $\mathfrak{D}_1$  bounded above by  $\mathcal{T}_s$ . It is forward invariant. Any trajectory going through any point of  $\mathfrak{O}$  converges to  $M_\ell$  at  $\infty$ . Either it meets the axis  $Y = 0$  at some point  $(\xi, 0)$  with  $\xi > 0$ , or it stays in  $\mathfrak{O}$ , satisfies  $R_w > 0$  and  $\lim_{\tau \rightarrow \ln R_w} T/y = 1$ , and meets  $M_\alpha$ , since  $M_\ell$  lies strictly below  $M_\alpha$ . Let

$$\begin{aligned} \mathcal{A} &= \{P \in \mathfrak{O} : \mathcal{T}_{[P]} \cap \{(\varphi, 0) : \varphi > 0\} \neq \emptyset\}, \\ \mathcal{B} &= \{P \in \mathfrak{O} : \mathcal{T}_{[P]} \cap M_\alpha \neq \emptyset\}. \end{aligned}$$

These sets are nonempty and open: indeed, one can check that the intersection with  $M_\alpha$  is transverse, because  $\alpha \neq \eta$ . Thus  $\mathcal{A} \cup \mathcal{B} \neq \mathfrak{O}$ , so there exists a trajectory  $\mathcal{T}_1$  with  $w$  of type (1).

(ii) Assume  $\alpha = \eta$  (Figure 8, right). There is no positive solution with  $R_w > 0$ , thus no solution of type (2) or (3). Indeed all the trajectories stay below  $\mathcal{T}_s$ , and  $\mathcal{T}_s$  is defined by the equation  $\zeta \equiv \eta$ , meaning that  $w \equiv Cr^{-\eta}$ , or equivalently  $Y_\eta \equiv C$ ; thus  $Y'_\eta \equiv 0$  and  $\mathcal{T}_s = M_\eta$ . Consider any trajectory  $\mathcal{T}_{[P]}$  going through some point  $P = (\varphi, 0)$  with  $\varphi > 0$ , and the solution  $(y, Y)$  issuing from  $P$  at time 0. Then  $Y_\eta(0) = 0$  and  $Y_\eta < 0$ , so  $Y'_\eta = \eta y - |Y|^{(2-p)/(p-1)} Y > 0$  on  $(-\infty, 0)$ , seeing that  $\mathcal{T}_{[P]}$  does not meet  $-\mathcal{T}_s$ . Suppose  $R_w = 0$ . Then  $\mathcal{T}_{[P]}$  starts from  $\mathfrak{D}_3$ , with  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha = \eta$ . Then  $\lim_{\tau \rightarrow -\infty} y_\eta = L < 0$ ; thus  $\lim_{\tau \rightarrow -\infty} Y_\eta = -(\alpha|L|)^{(2-p)/(p-1)}$ . A straightforward computation gives

$$Y''_\eta = Y'_\eta \left( N - \frac{1}{p-1} |Y|^{(2-p)/(p-1)} \right).$$

This leads to  $Y''_\eta < 0$  near  $-\infty$ , which is impossible; thus  $R_w < \infty$  and  $w$  is of type (4). □

**Remarks.** (i) For  $\alpha \leq \eta$ , both trajectories  $\mathcal{T}_r$  and  $\mathcal{T}_s$  stay in  $\mathfrak{D}_1$ .

(ii) When  $\alpha \leq N$ , one can verify that the regular positive solution  $y$  is increasing and  $y \leq \ell$  on  $\mathbb{R}$ , so  $r^\delta w(r) \leq \ell$  for any  $r \geq 0$ .

(iii) When  $\alpha = N$ , we have  $\mathcal{T}_r = \{(\xi, \xi) : \xi \in [0, \ell)\}$ , and the corresponding solutions  $w$  are given by (1–6) with  $K > 0$ . And  $\mathcal{T}_3 = \{(\xi, \xi) : \xi > \ell\}$  is a trajectory corresponding to particular solutions  $w$  of type (3), given by (1–6) with  $K < 0$ .

Next we come to the most interesting case, where  $\eta < \alpha$ .

**Lemma 5.5.** *Assume  $\varepsilon = 1$ ,  $\delta < \min(\alpha, N)$  and  $\eta < \alpha$ . If  $N/2 < \delta$  and  $\alpha < \alpha^*$  and  $\mathcal{T}_s$  stays in  $\mathcal{Q}_1$ ,  $\mathcal{T}_s$  has a limit cycle at  $-\infty$  in  $\mathcal{Q}_1$  or is homoclinic. If  $\delta \leq N/2$ , then  $\mathcal{T}_s$  does not stay in  $\mathcal{Q}_1$ .*

*Proof.* In any case  $M_\ell$  is a sink, so  $\mathcal{T}_s$  cannot converge to  $M_\ell$  at  $-\infty$ . Suppose  $\mathcal{T}_s$  has no limit cycle in  $\mathcal{Q}_1$ , and is not homoclinic and stays in  $\mathcal{Q}_1$ . (This happens when  $\delta \leq N/2$ , by Proposition 2.11.) Then either  $\lim_{\tau \rightarrow -\infty} y = \infty$  and  $\lim_{r \rightarrow 0} r^\alpha w = \Lambda \neq 0$ , or  $R_w > 0$ . In either case, for any  $d \in (\eta, \alpha)$ , the function  $y_d(\tau) = r^d w = r^{d-\delta} y$  satisfies  $\lim_{\tau \rightarrow \ln R_w} y_d = \infty = \lim_{\tau \rightarrow \infty} y_d$ . Then it has a minimum point, contradicting (2-5).  $\square$

**Theorem 5.6.** *Assume  $\varepsilon = 1$  and  $N/2 < \delta < \min(\alpha, N)$ . Then  $w(r) = \ell r^{-\delta}$  is still a solution.*

(i) *There exists a (maximal) critical value  $\alpha_{\text{crit}}$  of  $\alpha$ , such that*


$$\max(\eta, \alpha_1) < \alpha_{\text{crit}} < \alpha^*,$$

*and the regular trajectory is homoclinic: all regular solutions of  $(E_w)$  have constant sign and satisfy  $\lim_{r \rightarrow \infty} r^\eta w = c \neq 0$ .*

- (ii) *For any  $\alpha \in (\alpha_{\text{crit}}, \alpha^*)$ , there does exist a cycle in  $\mathcal{Q}_1$ , in other words there exist positive solutions  $w$  such that  $r^\delta w$  is periodic in  $\ln r$ . There exist positive solutions such that  $r^\delta w$  is asymptotically periodic in  $\ln r$  near 0 and  $\lim_{r \rightarrow \infty} r^\delta w = \delta$ . There exist positive solutions such that  $r^\delta w$  is asymptotically periodic in  $\ln r$  near 0 and  $\lim_{r \rightarrow \infty} r^\eta w = c \neq 0$ .*
- (iii) *For any  $\alpha \geq \alpha^*$  there does not exist such a cycle, but there exist positive solutions such that  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $\lim_{r \rightarrow \infty} r^\eta w = c > 0$ .*
- (iv) *For any  $\alpha > \alpha_{\text{crit}}$ , there exists also a cycle, surrounding  $(0, 0)$  and  $\pm M_\ell$ , thus  $r^\delta w$  is changing sign and periodic in  $\ln r$ . All regular solutions change signs and are oscillating at  $\infty$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ . There exist solutions such that  $R_w > 0$ , or  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$ , and oscillating at  $\infty$ , and  $r^\delta w$  is asymptotically periodic in  $\ln r$ .*

*Proof.* (i) For any  $\alpha \in (\alpha_1, \alpha_2)$  such that  $\eta \leq \alpha$ , we have by Remark 5.3 three possibilities for the regular trajectory  $\mathcal{T}_r$ :

- $\mathcal{T}_r$  converges to  $M_\ell$  and spirals around it, or else it has a limit cycle in  $\mathcal{Q}_1$  around  $M_\ell$ . Then  $\mathcal{T}_r$  meets the set  $\mathcal{E} = \{(\ell, Y) : Y > (\delta\ell)^{p-1}\}$  at a first point  $(\ell, Y_r(\alpha))$ . Note that  $\ell$  and  $\mathcal{E}$  depend continuously on  $\alpha$ . Then  $\mathcal{T}_s$  meets  $\mathcal{E}$  at some last point  $(\ell, Y_s(\alpha))$  such that  $Y_s(\alpha) - Y_r(\alpha) > 0$ . See Figure 9, top left.



**Figure 9.** Theorem 5.6:  $\varepsilon = -1$ ,  $N/2 < \delta = 3 < N = 4$ .

- $\mathcal{T}_r$  does not stay in  $\mathcal{Q}_1$ ; then  $\mathcal{T}_s$  is bounded at  $-\infty$ , and so converges to  $M_\ell$  at  $-\infty$  and spirals around this point, or it has a limit cycle around  $M_\ell$ . Then  $\mathcal{T}_s$  meets  $\mathcal{E}$  at a last point  $(\ell, Y_s(\alpha))$  and  $\mathcal{T}_r$  meets  $\mathcal{E}$  at a first point  $(\ell, Y_r(\alpha))$  such that  $Y_s(\alpha) - Y_r(\alpha) < 0$ . See Figure 9, bottom row.
- $\mathcal{T}_r$  is homoclinic, which is equivalent to  $Y_s(\alpha) - Y_r(\alpha) = 0$ . See Figure 9, top right.

Now the function  $\alpha \mapsto g(\alpha) = Y_s(\alpha) - Y_r(\alpha)$  is continuous. If  $\alpha_1 < \eta$ , then  $g(\eta)$  is defined and  $g(\eta) > 0$ , by Theorem 5.4. If  $\eta \leq \alpha_1$ , we observe that for  $\alpha = \alpha_1$ , the trajectory  $\mathcal{T}_s$  leaves  $\mathcal{Q}_1$ , by Theorem 2.18, because  $\alpha_1$  is a sink, and does so transversally by Remark 2.1(i). The same holds for  $\alpha = \alpha_1 + \gamma$  for  $\gamma$  small enough, by continuity, so  $\mathcal{T}_r$  stays in  $\mathcal{Q}_1$  and  $g(\alpha_1 + \gamma) > 0$ . If  $\alpha \geq \alpha^*$  (Figure 9, bottom right), then  $M_\ell$  is a source or a weak source, by Theorem 2.16; thus  $\mathcal{T}_r$  cannot converge to  $M_\ell$ . By Theorem 2.19, there exists no cycle in  $\mathcal{Q}_1$  and no homoclinic orbit. By Remark 5.3(i),  $\mathcal{T}_r$  cannot stay in  $\mathcal{Q}_1$ , so  $g(\alpha) < 0$  for  $\alpha^* \leq \alpha < \alpha_2$ . As a

consequence, there exists at least one  $\alpha_{\text{crit}} \in (\max(\eta, \alpha_1), \alpha^*)$  such that  $g(\alpha_{\text{crit}}) = 0$ . If it is not unique, we can choose the largest one.

(ii) Suppose  $\alpha < \alpha^*$ . The existence and uniqueness of the desired cycle  $\mathbb{C}$  in  $\mathcal{D}_1$  follows by Theorem 2.16 when  $\alpha$  is close to  $\alpha^*$  (Figure 9, bottom left). In fact, existence holds for any  $\alpha \in (\alpha_{\text{crit}}, \alpha^*)$ ; indeed  $g(\alpha) < 0$  on this interval, and  $\mathcal{T}_s$  cannot converge to  $M_\ell$  at  $-\infty$ , so it has a limit cycle around  $M_\ell$  at  $-\infty$ . Since  $M_\ell$  is a sink, there exist also trajectories converging to  $M_\ell$  at  $\infty$ , with a limit cycle at  $-\infty$  contained in  $\mathbb{C}$ . Now  $\mathcal{T}_r$  does not stay in  $\mathcal{D}_1$  and is bounded at  $\infty$ , so it has a limit cycle at  $\infty$  containing the three stationary points.

(iii) Suppose  $\alpha \geq \alpha^*$ . Then  $\mathcal{T}_s$  stays in  $\mathcal{D}_1$ , is bounded on  $\mathbb{R}$ , and converges at  $-\infty$  to  $M_\ell$ . At the same time,  $\mathcal{T}_r$  does not stay in  $\mathcal{D}_1$  for the same reason as above; thus it has a limit cycle at  $\infty$ , containing the three stationary points (see Figure 9, bottom right).

(iv) For any  $\alpha > \alpha_{\text{crit}}$ , apart from  $\mathcal{T}_s$  and the cycles, all the trajectories have a limit cycle at  $\infty$  containing the three stationary points. By Theorem 2.21, all the cycles are contained in a ball  $B$  of  $\mathbb{R}^2$ . Take any point  $P$  exterior to  $B$ . By Remark 5.3(ii),  $\mathcal{T}_{[P]}$  has a limit cycle at  $\infty$  contained in  $B$  and cannot have a limit cycle at  $-\infty$ . Thus  $y$  has constant sign near  $\ln R_w$ . By Proposition 2.8, either  $R_w > 0$  or  $y$  is defined near  $-\infty$  and  $\lim_{\tau \rightarrow -\infty} \zeta = L$ ,  $\lim_{r \rightarrow 0} r^\alpha w = L$ . □

**Note.** It is an open question whether  $\alpha_{\text{crit}}$  is unique. It can be shown that if there exist two critical values  $\alpha_{\text{crit}}^1 > \alpha_{\text{crit}}^2$ , the first orbit is contained in the second.

When  $\delta \leq N/2$ , or equivalently  $p \leq P_2$ , there are no cycles in  $\mathbb{R}^2$  and we get:

**Theorem 5.7.** *Assume  $\varepsilon = 1$ ,  $\delta \leq N/2$ , and  $\delta < \alpha$ . All regular solutions of  $(E_w)$  have constant sign, and  $\lim_{r \rightarrow \infty} r^\delta |w| = \ell$ . All solutions have a finite number of zeros. The function  $w(r) = \ell r^{-\delta}$  is a solution. If  $\alpha \leq \eta$ , Theorem 5.4 applies. If  $\eta < \alpha$ , all other solutions have at least one zero. There exist solutions satisfying  $\lim_{r \rightarrow \infty} r^n w = c \neq 0$  and having  $m$  zeros, for some  $m > 0$ . All other solutions satisfy  $\lim_{r \rightarrow \infty} r^\delta w = \pm \ell$ , and have  $m$  or  $m + 1$  zeros. There exist solutions with  $m + 1$  zeros.*

*Proof.* (i) By Proposition 2.11, all solutions have a finite number of zeros. Since  $\delta \leq N/2$ , the function  $W$  defined in (2–21) is nonincreasing. The regular solutions  $(y, Y)$  satisfy  $\lim_{\tau \rightarrow -\infty} W(\tau) = 0$ , so  $W(\tau) \leq 0$  on  $\mathbb{R}$ . If  $y(\tau_0) = 0$  for some real  $\tau_0$ , then  $W(\tau_0) = |Y(\tau_0)|^{p'} > 0$ , and we reach a contradiction. From Propositions 2.8 and 2.11 we obtain  $\lim_{\tau \rightarrow \infty} y = \pm \ell$ , so  $\lim_{r \rightarrow \infty} r^\delta w = \pm \ell$ .

(ii) Assume  $\eta < \alpha$ . By Lemma 5.5,  $\mathcal{T}_s$  does not stay in  $\mathcal{D}_1$ . By Propositions 2.8 and 2.15,  $\mathcal{T}_s$  cannot stay in  $\mathcal{D}_4$ , so it intersects the line  $y = 0$  at points  $(0, \xi_1), \dots, (0, \xi_m)$ . By Remark 5.3, any trajectory other than  $\mathcal{T}_s$  converges to  $\pm M_\ell$ . Given

any  $P = (0, \xi)$ , with  $\xi > |\xi_i|$  for  $1 \leq i \leq m$ , the trajectory  $\mathcal{T}_{[P]}$  cannot intersect  $\mathcal{T}_s$  or  $-\mathcal{T}_s$ , so  $y$  has  $m + 1$  zeros. Any other solution has  $m$  or  $m + 1$  zeros, because the trajectory does not meet  $\mathcal{T}_r$  or  $-\mathcal{T}_r$  or  $\mathcal{T}_{[P]}$ . Finally,  $R_w > 0$  or  $\lim_{r \rightarrow 0} r^\alpha w = L \neq 0$ .  $\square$

**Note.** Theorems 5.4, 5.6 and 5.7 recover, in particular, the results in [Qi and Wang 1999, Theorem 2].

### 6. The case $\varepsilon = -1, \alpha \leq \delta$

$$\max(\alpha, N) \leq \delta$$

Here  $(0, 0)$  is the only stationary point, and it is a source when  $\delta \neq N$ . We first suppose  $0 < \alpha$ .


**Theorem 6.1.** *Suppose  $\varepsilon = -1, \max(\alpha, N) \leq \delta$  and  $0 < \alpha$ .*

- (i) *Suppose  $\alpha \neq N$  or  $\alpha \neq \delta$ . Then all regular solutions of  $(E_w)$  have constant sign and a reduced domain  $(S_w < \infty)$ . There exist solutions satisfying any one of these characterizations:*
  - (1)  *$w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = c \neq 0$  if  $N \geq 2$  ( $\lim_{r \rightarrow 0} w = a > 0, \lim_{r \rightarrow 0} w' = b < 0$  if  $N = 1$ ), and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$  if  $\alpha \neq \delta$ , or (2–39) holds if  $\alpha = \delta$ ;*
  - (2)  *$w$  is positive,  $\lim_{r \rightarrow 0} r^\alpha w = c \neq 0$  if  $N \geq 2$  ( $\lim_{r \rightarrow 0} w = a > 0, \lim_{r \rightarrow 0} w' = b \neq 0$ , or  $a = 0 < b$  if  $N = 1$ ), and  $S_w < \infty$ ;*
  - (3)  *$w$  has one zero,  $\lim_{r \rightarrow 0} r^\alpha w = c \neq 0$  if  $N \geq 2$  ( $\lim_{r \rightarrow 0} w = a > 0, \lim_{r \rightarrow 0} w' = b < 0$  if  $N = 1$ ), and  $S_w < \infty$ .*
- (ii) *Suppose  $\alpha = \delta = N$ . Then the regular solutions, given by (1–6), have constant sign, with  $S_w < \infty$ . For any  $k \in \mathbb{R}, w(r) = kr^{-N}$  is a solution. Moreover there exist positive solutions such that  $\lim_{r \rightarrow 0} r^N w = c > 0$  and  $S_w < \infty$ , and solutions with one zero, such that  $\lim_{r \rightarrow 0} r^N w = c > 0$  and  $S_w < \infty$ .*

*Up to symmetry, all solutions are as above.*

*Proof.* (i) Here  $\alpha \neq N$  or  $\alpha \neq \delta$  (Figure 10, left). Since  $\alpha > 0$ , Propositions 2.5, 2.7 and 2.14 imply that  $y > 0$  and  $S_w < \infty$  for  $\mathcal{T}_r$ ; and any solution  $y$  has at most one zero, and  $y, Y$  are monotone near  $-\infty$  and near  $\ln S_w$ . By Proposition 2.8, any trajectory  $\mathcal{T}$  converges to  $(0, 0)$  at  $-\infty$ ; and apart from  $\mathcal{T}_r$ , such a trajectory is tangent to the axis  $y = 0$ . Now suppose  $y > 0$  near  $-\infty$ . If  $N \geq 2$ , then  $\mathcal{T}$  starts in  $\mathcal{Q}_1$ , since  $\lim_{\tau \rightarrow -\infty} \zeta = \eta > 0$ ; if  $N = 1$ , then  $\lim_{r \rightarrow 0} w = a \geq 0$  and  $\lim_{r \rightarrow 0} w' = b$ , and  $\mathcal{T}$  starts in  $\mathcal{Q}_1$  if  $b < 0$  and in  $\mathcal{Q}_4$  if  $b > 0$  (in particular when  $a = 0$ ).

For any  $P = (\varphi, 0)$  with  $\varphi > 0$ , the trajectory  $\mathcal{T}_{[P]}$  satisfies  $y > 0$  on  $\mathbb{R}$ , and by Remark 2.1(i), it stays in  $\mathcal{Q}_4$  after  $P$ , because it cannot meet  $\mathcal{T}_r$  (hence  $S_w < \infty$ ); also it stays in  $\mathcal{Q}_1$  before  $P$ , so  $w$  is of type (2). In the same way for any  $P = (0, \xi)$



**Figure 10.** Theorem 6.1:  $\varepsilon = -1$ .

with  $\xi > 0$ , the trajectory  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_2$  after  $P$ , since it cannot meet  $-\mathcal{T}_r$  (hence  $S_w < \infty$ ), and it stays in  $\mathcal{Q}_1$  before  $P$ , so  $w$  is of type (3).

Next consider the sets

$$\mathcal{A} = \{P \in \mathcal{Q}_1 : \mathcal{T}_{[P]} \cap \{(\varphi, 0) : \varphi > 0\} \neq \emptyset\},$$

$$\mathcal{B} = \{P \in \mathcal{Q}_1 : \mathcal{T}_{[P]} \cap \{(0, \xi) : \xi > 0\} \neq \emptyset\}.$$

From the previous discussion we know they are nonempty and open, so  $\mathcal{A} \cup \mathcal{B} \neq \mathcal{Q}_1$ . There exists a trajectory  $\mathcal{T}_1$  starting at  $(0, 0)$  and staying in  $\mathcal{Q}_1$ . By Proposition 2.8, necessarily  $\lim_{\tau \rightarrow \infty} y = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha > 0$ , so  $w$  is of type (1) by Proposition 2.9.

Finally we describe all other trajectories  $\mathcal{T}_{[P]}$  with one point  $P$  in the domain  $\mathcal{R}$  above  $\mathcal{T}_r \cup (-\mathcal{T}_r)$ . If  $P$  is in the domain delimited by  $\mathcal{T}_r, \mathcal{T}_1$ , then  $w$  is still of the type (2). If  $P$  is in the domain delimited by  $-\mathcal{T}_r, \mathcal{T}_1$ , then either  $y$  has a zero and  $w$  is of type (3), or  $N = 1, y < 0$  and  $-w$  is of type (2). Up to a symmetry, all the solutions have been obtained.

(ii) Here  $\alpha = \delta = N$  (Figure 10, right). Since  $\alpha = N$  equation (1–5) holds, and the regular solutions relative to  $C = 0$  are given by (1–6). Since  $\delta = N$ , (1–5) is equivalent to  $y + Y \equiv C$ , from (2–12). For any  $k \in \mathbb{R}$ ,  $(y, Y) \equiv P_k = (k, |Nk|^{p-2}Nk)$  is a solution of system  $(S)$ , located on the curve  $\mathcal{M}$ , thus  $w(r) = kr^{-N}$  is a solution of  $(E_w)$ . Any solution has at most one zero, by Proposition 2.5. From Propositions 2.8, and 2.10, any other trajectory converges to a point  $P_k \in \mathcal{M}$  at  $\infty$ , and  $S_w < \infty$ . There exists trajectories such that  $y$  has constant sign, and other ones such that  $y$  has one zero. All solutions have been obtained.  $\square$

Next we suppose  $\alpha < 0$ , and distinguish the cases  $N \geq 2$  and  $N = 1$ .

**Theorem 6.2.** *Suppose  $\varepsilon = -1$  and  $\alpha < 0 < 2 \leq N \leq \delta$ . Then any solution of  $(E_w)$  has a finite number of zeros. Regular solutions have at least one zero, and*

precisely one if  $-p' \leq \alpha$ . Any solution has at least one zero, and any nonregular solution satisfies  $\lim_{r \rightarrow 0} r^\eta w = c \neq 0$ .


If  $-p' < \alpha$ , all regular solutions have a reduced domain ( $S_w < \infty$ ), and they fall into the following types, all of which occur:

- (1) solutions with two zeros and  $S_w < \infty$ ;
- (2) solutions with one zero and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ ;
- (3) solutions with one zero and  $S_w < \infty$ .

If  $\alpha = -p'$ , all regular solutions satisfy  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ . The other solutions are of type (1).

*Proof.* By Proposition 2.8, any trajectory converges necessarily to  $(0, 0)$  at  $-\infty$ , and apart from  $\mathcal{T}_r$ , it is tangent to the axis  $y = 0$ . Any solution  $y$  has a finite number of zeros, and  $y$  is monotone near  $-\infty$ , and near  $S_w$  (finite or not), by Propositions 2.7 and 2.11, since  $\delta > N/2$ . Either  $S_w < \infty$ , so  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ , or  $S_w = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha < 0$ . In any case  $(y, Y)$  is in  $\mathcal{Q}_2$  or  $\mathcal{Q}_4$  for large  $\tau$ . By Proposition 2.14,  $\mathcal{T}_r$  has at least one zero, and starts in  $\mathcal{Q}_1$ . Since  $N \geq 2$ , any trajectory  $\mathcal{T} \neq \pm \mathcal{T}_r$  satisfies  $\lim_{\tau \rightarrow -\infty} \zeta = \eta > 0$ . Thus it starts in  $\mathcal{Q}_1$  (or  $\mathcal{Q}_3$ ), and has at least one zero. Any trajectory  $\mathcal{T}$  starting in  $\mathcal{Q}_1$  enters  $\mathcal{Q}_2$ , by Remark 2.1(i). And  $y' = \delta y - Y^{1/(p-1)}$ , so  $y$  decreases as long as  $\mathcal{T}$  stays in  $\mathcal{Q}_2$ . Then either  $\mathcal{T}$  enters  $\mathcal{Q}_3$ , hence also  $\mathcal{Q}_4$ , and  $y$  has at least two zeros; or it stays in  $\mathcal{Q}_2$ , and either  $S_w < \infty$  and  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ , or  $S_w = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ .

(i) Suppose  $-p' < \alpha$  (Figure 11, left). Then  $\mathcal{T}_r$  has precisely one zero, by Proposition 2.14, thus it stays in  $\mathcal{Q}_2$ , and  $S_w < \infty$ ,  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ . Any other solution has at most two zeros, because the trajectory does not meet  $\pm \mathcal{T}_r$ . Recall that the function  $Y_\alpha$  defined by (2–3) with  $d = \alpha$  has only minimal points on the



**Figure 11.** Theorem 6.2:  $\varepsilon = -1$ ,  $N = 2 < \delta = 3$ .

sets where it is positive, by Remark 2.6. By Proposition 2.14,  $\mathcal{T}_r$  satisfies

$$Y'_\alpha = -(p-1)(\eta-\alpha)Y_\alpha + e^{(p-(2-p)\alpha)\tau}(Y_\alpha^{1/(p-1)} - \alpha y_\alpha) > 0,$$

which is equivalent to

$$(6-1) \quad Y^{1/(p-1)} - (p-1)(\eta-\alpha)Y > \alpha y.$$

$\mathcal{T}_r$  stays strictly to the right of the curve

$$(6-2) \quad \mathcal{N}_\alpha = \{(y, Y) \in \mathbb{R} \times (0, \infty) : \alpha y = Y^{1/(p-1)} - (p-1)(\eta-\alpha)Y\},$$

which intersects the axis  $y = 0$  at the points  $(0, 0)$  and  $(0, (p-1)(\eta-\alpha))$ .

For  $\bar{P} = (\varphi, 0)$  with  $\varphi < 0$ , the trajectory  $\mathcal{T}_{[\bar{P}]}$  enters  $\mathcal{Q}_3$  after  $\bar{P}$ , by Remark 2.1(i); the solution passing through  $\bar{P}$  at  $\tau = 0$  satisfies  $Y_\alpha(0) = 0$  (so  $Y_\alpha$  stays positive for  $\tau < 0$ ) and  $Y'_\alpha(\tau) < 0$ , since  $Y_\alpha$  has no maximal point. Thus  $\mathcal{T}_{[\bar{P}]}$  stays in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$  before  $P$ , to the left of  $\mathcal{N}_\alpha$ , and starts and ends in  $\mathcal{Q}_1$  and ends up in  $\mathcal{Q}_4$ . Hence  $y$  has two zeros. If  $S_w = \infty$  then  $\lim_{\tau \rightarrow \infty} |y| = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha < 0$ ; this is impossible, because  $\mathcal{T}_{[\bar{P}]}$  does not meet  $-\mathcal{T}_r$ . Thus  $S_w < \infty$ , and  $w$  is of type (1).

Next consider  $\mathcal{T}_{[P]}$ , for  $P = (\varphi, \xi) \in \mathcal{N}_\alpha$ , with  $\varphi \leq 0$ . The solution going through  $P$  at  $\tau = 0$  satisfies  $Y'_\alpha(0) = 0$ ,  $Y_\alpha(0) > 0$ , and  $0$  is a minimal point; hence  $Y''_\alpha(0) > 0$ . Indeed, if  $Y''_\alpha(0) = 0$ , then  $Y_\alpha$  is constant on  $\mathbb{R}$  by uniqueness; by (2-6), in turn, we have  $Y_\alpha \equiv 0$  (since  $\alpha \neq -p'$ ); but this is false. Therefore  $Y'_\alpha(\tau) > 0$  for  $\tau > 0$  and  $Y'_\alpha(\tau) < 0$  for  $\tau < 0$ . Thus  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$ , to the right of  $\mathcal{N}_\alpha$  after  $P$ , with  $y < 0$  by Remark 2.1(i); it stays to the left of  $\mathcal{N}_\alpha$  before  $P$ , and converges to  $(0, 0)$  at  $-\infty$  in  $\mathcal{Q}_1$ . Suppose that  $S_w = \infty$ . Then  $\lim_{\tau \rightarrow \infty} |y| = \infty$ ,  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ , and  $\lim_{\tau \rightarrow \infty} y_\alpha = L < 0$  by Proposition 2.9; thus  $\lim_{\tau \rightarrow \infty} Y_\alpha = (\alpha L)^{p-1}$ . As in Proposition 2.14, one finds that  $Y''_\alpha(\tau) > 0$  for any  $\tau > 0$ , which is impossible. Thus  $\mathcal{T}_{[P]}$  satisfies  $S_w < \infty$ , showing that  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ . The corresponding  $w$  is of type (3).

Finally, let  $\mathcal{R}$  be the domain of  $\mathcal{Q}_1 \cup \mathcal{Q}_2$  delimited by  $\mathcal{T}_r$  and containing  $\mathcal{N}_\alpha$ , and define the sets

$$(6-3) \quad \begin{aligned} \mathcal{A} &= \{P \in \mathcal{R} : \mathcal{T}_{[P]} \cap \{(\varphi, 0) : \varphi < 0\} \neq \emptyset\}, \\ \mathcal{B} &= \{P \in \mathcal{R} : \mathcal{T}_{[P]} \cap \mathcal{N}_\alpha \neq \emptyset\}, \end{aligned}$$

corresponding to trajectories of type (1) or (3). These sets are nonempty and open, because here again the intersection with  $\mathcal{N}_\alpha$  is transverse (recall that  $\alpha \neq -p'$ ). Thus  $\mathcal{A} \cup \mathcal{B} \neq \mathcal{R}$ . There exists a trajectory in  $\mathcal{R}$  disjoint from  $\mathcal{N}_\alpha$ , starting at  $(0, 0)$  in  $\mathcal{Q}_1$  and ending up in  $\mathcal{Q}_2$ . It cannot satisfy  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ , so  $S_w = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . Hence  $w$  is of type (2).

(ii) Suppose  $\alpha = -p'$  (Figure 11, right). The regular solutions are given by (1–8), they have one zero, but  $S_w = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . They satisfy  $Y_{-p'} \equiv C$ , thus  $Y'_{-p'} \equiv 0$ , thus  $\mathcal{T}_r = \mathcal{M}_{-p'}$ . Consider  $\mathcal{T}_{[\bar{P}]}$ ; the solution passing through  $\bar{P}$  at  $\tau = 0$  satisfies and  $Y_{-p'}(0) = 0$ , thus  $Y_{-p'}$  stays negative for  $\tau > 0$  and  $Y'_{-p'} < 0$ . Suppose that  $S_w = \infty$ , then  $\lim_{\tau \rightarrow \infty} y_\alpha = L > 0$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = -(|\alpha|L)^{p-1}$ . But as in (2-46),  $Y''_\alpha(\tau) < 0$  for any  $\tau > 0$ , which leads to a contradiction. Thus  $S_w < \infty$ , and  $w$  is of type (1). Finally suppose that there exists a trajectory  $\mathcal{T} \neq \mathcal{T}_r$  staying in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$ . Then  $Y_\alpha > 0$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = 0$ , and it cannot meet  $\mathcal{T}_r$ , thus  $S_w = \infty$ , and  $\lim_{\tau \rightarrow -\infty} Y_\alpha = \infty$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = C > 0$ . As in Proposition 2.14, it is impossible. Thus there does not exist solution of type (2) or (3).  $\square$

**Theorem 6.3.** *Suppose  $\varepsilon = -1$  and  $\alpha < 0 < N = 1 < \delta$ . Then any solution of  $(E_w)$  has still a finite number of zeros. Regular solutions have at least one zero, and precisely one if  $-p' \leq \alpha$ .*


*If  $-1 < \alpha < 0$ , all regular solutions have a reduced domain ( $S_w < \infty$ ). Moreover:*

- (1) *the solutions with  $\lim_{r \rightarrow 0} w = a > 0$  and  $\lim_{r \rightarrow 0} w' = b < 0$  have one zero and  $S_w < \infty$ ;*
- (2) *the solutions with  $\lim_{r \rightarrow 0} w = 0$  and  $\lim_{r \rightarrow 0} w' = b > 0$  are positive and  $S_w < \infty$ ;*
- (3) *there exist solutions with one zero and  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b > 0$  and  $S_w < \infty$ ;*
- (4) *there exist positive solutions with  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b > 0$  and  $S_w < \infty$ ;*
- (5) *for any  $a > 0$  there exists  $b > 0$  such that  $w$  is positive and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ .*

*If  $\alpha = -1$ , for any  $b > 0$ ,  $w \equiv br$  is a solution. The other solutions such that  $\lim_{r \rightarrow 0} w \neq 0$  have one zero, and satisfy  $S_w < \infty$ .*

*If  $-p' < \alpha < -1$ , then*

- (6) *there exist solutions with one zero, with  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b < 0$ , and  $S_w < \infty$ ;*
- (7) *the solutions with  $\lim_{r \rightarrow 0} w = 0$  and  $\lim_{r \rightarrow 0} w' = b > 0$  have one zero and  $S_w < \infty$ ;*
- (8) *there exist solutions with one zero, with  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b > 0$  and  $S_w < \infty$ ;*
- (9) *there exist solutions with  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b < 0$ , with two zeros and  $S_w < \infty$ ;*



**Figure 12.** Theorem 6.3:  $\varepsilon = -1$ ,  $N = 1 < \delta = 3$ .

- (10) for any  $a > 0$  there exists  $b > 0$  and a solution with  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b < 0$ , with one zero and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ .

*Proof.* The case  $N = 1$  is still the more complex one, since some trajectories start in  $\mathcal{Q}_2$  (or  $\mathcal{Q}_4$ ), corresponding to the solutions such that  $\lim_{r \rightarrow 0} w = a$  and  $\lim_{r \rightarrow 0} w' = b$ , with  $b \neq 0$ ,  $ab \geq 0$ . Any solution has still a finite number of zeros, by Proposition 2.11.

(i) Suppose  $-1 < \alpha < 0$  (Figure 12, left). By Proposition 2.5, any solution has at most one zero, so regular solutions have precisely one zero. Thus  $\mathcal{T}_r$  meets the axis  $y = 0$  at some point  $(0, \xi_r)$ .

Consider the trajectory  $\mathcal{T}_s$  such that  $\lim_{r \rightarrow 0} w = 0$  and  $\lim_{r \rightarrow 0} w' = b < 0$  (which means  $\lim_{\tau \rightarrow -\infty} \zeta = \eta = -1$ ), starting from  $(0, 0)$  in  $\mathcal{Q}_2$ , so  $w < 0$  near 0. For any  $d \in (-1, \alpha)$ , the function  $y_d$  satisfies  $y_d(\tau) = be^{(d+1)\tau}(1 + o(1))$  near  $-\infty$ , so  $\lim_{\tau \rightarrow -\infty} y_d = 0$ . Then  $y_d$  has no zeros, because  $|y_d|$  has no maximal point, by (2-14); thus  $\mathcal{T}_s$  stays in  $\mathcal{Q}_2$ . If  $\mathcal{T}_s$  satisfies  $S_w = \infty$ , then  $\lim_{\tau \rightarrow \infty} y_\alpha = L < 0$ , so  $\lim_{\tau \rightarrow \infty} y_d = 0$ , which is impossible; thus  $w$  is of type (2). The domain is reduced since  $\mathcal{T}_r$  cannot meet  $\mathcal{T}_s$ .

For  $\bar{P} = (\varphi, 0)$  with  $\varphi < 0$ , the trajectory  $\mathcal{T}_{[\bar{P}]}$  does not meet  $\mathcal{T}_s$ , thus converges to  $(0, 0)$  at  $-\infty$  in  $\mathcal{Q}_2$ ; then  $\lim_{r \rightarrow 0} (-w) = a > 0$  and  $\lim_{r \rightarrow 0} (-w)' = b > 0$ , and  $\mathcal{T}_{[\bar{P}]}$  ends up in  $\mathcal{Q}_4$ ; thus  $y$  has one zero and  $-w$  is of type (3).

For  $P = (0, \xi)$ , with  $\xi \in (0, \xi_r)$ ,  $\mathcal{T}_{[P]}$  has one zero and converges to  $(0, 0)$  at  $-\infty$  in  $\mathcal{Q}_1$ ; hence  $\lim_{r \rightarrow 0} w = a > 0$  and  $\lim_{r \rightarrow 0} w' = b < 0$ . The domain is reduced since  $\mathcal{T}_{[P]}$  and  $\mathcal{T}_s$  do not meet. Thus  $w$  is of type (1). Conversely, any solution such that  $\lim_{r \rightarrow 0} w = a > 0$  and  $\lim_{r \rightarrow 0} w' = b < 0$  has one zero and satisfies  $S_w < \infty$ .

Next consider a trajectory  $\mathcal{T}$  such that  $\lim_{r \rightarrow 0} (-w) = a > 0$  and  $\lim_{r \rightarrow 0} (-w)' = b > 0$ ; that is,  $\mathcal{T}$  starts in  $\mathcal{Q}_2$  below  $\mathcal{T}_s$ . Then  $\zeta(\tau) = -(b/a)e^\tau(1 + o(1))$  near  $-\infty$ ,

so  $\lim_{\tau \rightarrow -\infty} \zeta = 0$ . If  $\zeta$  has an extremal point  $\theta$ , we have

$$(p - 1)\zeta''(\theta) = (2 - p)(\zeta - \alpha)(\delta - \zeta)|\zeta y|^{2-p},$$

by (2-18); thus  $\theta$  is a minimal point if  $\zeta(\theta) > \alpha$ , and maximal if  $\zeta(\theta) < \alpha$ . (Equality is impossible since it would require  $\zeta \equiv \alpha$ .) Thus either  $\zeta$  has a first zero  $\tau_1$  and  $\alpha < \zeta(\tau) < 0$  for  $\tau < \tau_1$ , and  $\mathcal{T}$  is one of the  $\mathcal{T}_{[\bar{p}]}$ ; or  $\zeta$  remains negative, in which case if  $S_w = \infty$ , then  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ , so  $\zeta$  is necessarily decreasing, and  $\alpha < \zeta(\tau) < 0$  for any  $\tau$ . In both cases,  $\mathcal{T}$  stays below the curve

$$\mathcal{M}' = \{(y, Y) \in \mathbb{R} \times (0, \infty) : \alpha y = Y^{1/(p-1)}\},$$

as long as it is in  $\mathcal{Q}_2$ . Hence, for any  $P \in \mathcal{Q}_2$  such that  $P$  is on or above  $\mathcal{M}'$ , the trajectory  $\mathcal{T}_{[P]}$  satisfies  $S_w < \infty$ ; in particular on finds again  $\mathcal{T}_s$ . For any  $P$  between  $\mathcal{M}'$  and  $\mathcal{T}_s$ , the solution has constant sign,  $\mathcal{T}_{[P]}$  converges to  $(0, 0)$  at  $-\infty$  and  $\lim_{r \rightarrow 0}(-w) = a > 0$  and  $\lim_{r \rightarrow 0}(-w') = b > 0$ , and  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ , so  $\mathcal{T}_{[P]}$  meets  $\mathcal{M}_\alpha$ . Thus  $-w$  is of type (4).

Finally, let  $\mathcal{R}_1$  be the domain of  $\mathcal{Q}_2$  delimited by  $\mathcal{T}_s$  and the axis  $Y = 0$ , and set

$$\begin{aligned} \mathcal{A}_1 &= \{P \in \mathcal{R}_1 : \mathcal{T}_{[P]} \cap \{(\varphi, 0) : \varphi < 0\} \neq \emptyset\}, \\ \mathcal{B}_1 &= \{P \in \mathcal{R}_1 : \mathcal{T}_{[P]} \cap \mathcal{N}_\alpha \neq \emptyset\}. \end{aligned}$$

These sets are open, since the intersection is transverse (recall that  $\alpha \neq -1$ ). They are also nonempty, so  $\mathcal{A}_1 \cup \mathcal{B}_1 \neq \mathcal{R}_1$ , and there exists a trajectory such that  $y$  is defined on  $\mathbb{R}$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . By scaling, we can find for any  $a > 0$  at least one  $b$  such that the corresponding  $w$  has constant sign and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ ; thus  $|w|$  is of type (5).

(ii) Suppose  $\alpha = -1$ . Then  $\mathcal{T}_s$  is given explicitly by  $w \equiv br$ , so  $Y \equiv -y^{p-1}$ , or equivalently  $Y_{-1} \equiv b$ ; hence  $\mathcal{T}_s = \mathcal{N}_{-1}$ . For any other solution, one finds  $Y''_{-1} = Y'_{-1}(1 + e^{2\tau}|Y_{-1}|^{(2-p)/(p-1)})$ , so  $Y_{-1}$  is strictly monotone, by uniqueness, and  $Y''_{-1}$  has the sign of  $Y'_{-1}$ . Any trajectory such that  $\lim_{r \rightarrow 0} w = a > 0$  and  $\lim_{r \rightarrow 0} w' = b < 0$ , starting in  $\mathcal{Q}_1$ , satisfies  $Y'_{-1} > 0$ , and  $Y_{-1}$  is convex. Thus  $Y_{-1}$  cannot have a finite limit,  $S_w < \infty$ , and the trajectory ends up in  $\mathcal{Q}_2$ , so  $y$  has a zero. Any trajectory such that  $\lim_{r \rightarrow 0}(-w) = a > 0$  and  $\lim_{r \rightarrow 0}(-w)' = b > 0$ , starting in  $\mathcal{Q}_2$ , satisfies  $Y'_{-1} < 0$ , so  $Y_{-1}$  has a zero and the trajectory ends up in  $\mathcal{Q}_4$ . Hence, apart from  $\mathcal{T}_s$ , all trajectories satisfy  $S_w < \infty$ , and  $y$  has one zero.

(iii) Suppose  $-p' < \alpha < -1$  (Figure 12, right). Then  $\mathcal{T}_r$  starts in  $\mathcal{Q}_1$ ,  $y$  has one zero from Proposition 2.14, and  $\mathcal{T}_r$  ends up in  $\mathcal{Q}_2$ , with  $S_w < \infty$ . Any solution has at most two zeros.

Consider  $\mathcal{T}_s$ : we claim that it cannot stay in  $\mathcal{Q}_2$ . Suppose that it stays in it, thus  $y < 0 < Y$ . Then  $\zeta < 0$ , and  $\lim_{\tau \rightarrow -\infty} \zeta = \eta = -1$ , and  $\zeta$  is monotone near  $-\infty$ ; if  $\zeta' \leq 0$ , then  $\zeta \leq -1$  near  $-\infty$ , and we reach a contradiction from (2-9). Then

$\zeta' \geq 0$  near  $-\infty$ ; but any extremal point of  $\zeta$  is a minimal point by (2–18). Hence  $\zeta$  remains increasing, is defined on  $\mathbb{R}$  and has a limit  $\lambda \in [-1, 0]$ ; but  $\lambda = \alpha$ , by Proposition 2.8, again leading to a contradiction. Therefore  $\mathcal{T}_s$  enters  $\mathcal{D}_3$  at some point  $(\varphi_s, 0)$  with  $\varphi_s < 0$ , then enters  $\mathcal{D}_4$ , and  $y$  has precisely one zero; and  $w$  is of type (7).

Any solution such that  $\lim_{r \rightarrow 0}(-w) = a > 0$  and  $\lim_{r \rightarrow 0}(-w)' = b > 0$  also has one zero, since its trajectory stays under  $\mathcal{T}_s$  in  $\mathcal{D}_2$ ; thus  $w$  is of type (8).

As in the case  $N \geq 2$ , for any  $P = (\varphi, \xi) \in \mathcal{N}_\alpha$  with  $\varphi \leq 0$ ,  $\mathcal{T}_{[P]}$  stays in  $\mathcal{D}_1 \cup \mathcal{D}_2$  and  $S_w < \infty$ . In particular for  $P_0 = (0, \xi_0)$ , where  $\xi_0 = ((p-1)(-1-\alpha))^{(p-1)/(2-p)}$ , the trajectory  $\mathcal{T}_{[P_0]}$  starts from  $\mathcal{D}_1$ , so  $\lim_{r \rightarrow 0} w = a > 0$ ,  $\lim_{r \rightarrow 0} w' = b_0(a) > 0$ ; also  $w$  has one zero, and  $S_w < \infty$ . Thus  $w$  is of type (6).

The sets  $\mathcal{A}, \mathcal{B}$  defined as in (6–3) are still open in this case, and  $\mathcal{B}$  contains  $\mathcal{T}_{[P_0]}$ . Also,  $\mathcal{A}$  contains  $\mathcal{T}_s$ ; hence  $\mathcal{A}$  contains any  $\mathcal{T}_{[P]}$ , where  $P = (\varphi, 0)$  with  $\varphi < \varphi_s$ . Such a trajectory satisfies  $\lim_{r \rightarrow 0} w = a > 0$  and  $\lim_{r \rightarrow 0} w' = b < 0$ , and  $w$  is of type (9). Moreover  $\mathcal{A} \cup \mathcal{B} \neq \mathcal{R}$ ; thus for any  $a > 0$  there exists  $b < 0$  such that the corresponding  $w$  has one zero and  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ , so  $w$  is of type (10). □

$$\alpha < \delta < N$$

As in the case  $\varepsilon = 1, \delta < \min(\alpha, N)$  of page 246, here two kinds of periodic trajectories can appear, and the study is delicate. Here also  $N \geq 2$ , and we still have three stationary points, and  $(0, 0)$  is a saddle point.  $M_\ell$  is a source if  $N/2 \leq \delta$  or  $\delta < N/2$  and  $\alpha^* < \alpha$ , and a sink if  $\delta < N/2$  and  $\alpha < \alpha^*$ ; notice that  $\alpha^* < -p' < 0$ , by (2–32). Also  $M_\ell$  is a node whenever  $\alpha \leq \alpha_1$  or  $\alpha_2 \leq \alpha$ , where  $\alpha_1, \alpha_2$  are defined in (2–48), and  $\alpha_2$  can be greater or less than  $-p'$ . We begin with the simplest case.

**Theorem 6.4.** *Assume  $\varepsilon = -1$  and  $0 < \alpha < \delta < N$ . All regular solutions have constant sign and a reduced domain ( $S_w < \infty$ ). The function  $w \equiv \ell r^{-\delta}$  is a solution. There exist solutions satisfying any one of these characterizations:*

- (1)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $S_w < \infty$ ;
- (2)  $w$  has one zero,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $S_w < \infty$ ;
- (3)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $\lim_{r \rightarrow \infty} r^\eta w = c > 0$ ;
- (4)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $\lim_{r \rightarrow \infty} r^\alpha w = L > 0$ .

*Up to symmetry, all solutions are as above.*

*Proof.* Since  $\alpha > 0$ , regular solutions have constant sign and satisfy  $S_w < \infty$ , by Propositions 2.5 and 2.14. Here  $\mathcal{T}_r$  starts in  $\mathcal{D}_4$  and stays in it, by Remark 2.3 (Figure 13). Any solution has at most one zero by Proposition 2.5. The point  $M_\ell$  is a source, and a node point, by Remark 2.17, and  $0 < \lambda_1 < \delta < \lambda_2$ . The eigenvectors  $u_1 = (\nu(\alpha), \lambda_1 - \delta)$  and  $u_2 = (-\nu(\alpha), \delta - \lambda_2)$  form a positively oriented basis, where

now  $v(\alpha) < 0$ ; thus  $u_1$  points toward  $\mathcal{Q}_3$  and  $u_2$  toward  $\mathcal{Q}_4$ . There are two particular trajectories  $\mathcal{T}_1, \mathcal{T}_2$  starting from  $M_\ell$  at  $-\infty$ , with respective tangent vectors  $u_2$  and  $-u_2$ . All other trajectories  $\mathcal{T}$  approaching  $M_\ell$  at  $-\infty$  do so along  $u_1$ ; and  $y$  is monotone at the extremities, by Proposition 2.7, since  $\mathcal{T}$  cannot meet  $\mathcal{T}_1, \mathcal{T}_2$ .

First consider  $\mathcal{T}_1$ . The function  $y$  is nondecreasing near  $-\infty$  and remains so as long as  $\mathcal{T}_1$  stays in  $\mathcal{Q}_1$ ; indeed,  $Y$  is nonincreasing near  $-\infty$ , so  $Y(\tau) < (\delta\ell)^{p-1}$ . If  $y$  has a maximal point  $\tau$ , then  $y(\tau) > \ell$  by (2-16), and  $Y^{1/(p-1)} = \delta y$ ; hence  $Y(\tau) > (\delta\ell)^{p-1}$ , so  $Y$  has a minimal point  $\tau_1$  in  $\mathcal{Q}_1$ ; therefore  $Y(\tau_1) < (\delta\ell)^{p-1}$  by  $(E_Y)$ ; and  $Y'(\tau_1) = 0$ , so  $\alpha\ell < \alpha y(\tau_1) < (N - \delta)\alpha Y(\tau_1)/(\delta - \alpha)$ , a contradiction. If  $\mathcal{T}_1$  stays in  $\mathcal{Q}_1$ , then  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha > 0$  by Proposition 2.8, which is also contradictory. Thus  $\mathcal{T}_1$  enters  $\mathcal{Q}_4$  at some point  $(\varphi_1, 0)$  and stays in it;  $S_w < \infty$  because  $\mathcal{T}_1$  and  $\mathcal{T}_r$  don't meet, so  $w$  is of type (1).


Next consider  $\mathcal{T}_2$ . Near  $-\infty$ , the function  $Y$  is nondecreasing, and  $y$  is nonincreasing;  $y$  is monotone as long as  $y > 0$ : if there existed a minimal point  $\tau$ , we would have  $y(\tau) > \ell$  by (2-16). Also  $Y$  is nondecreasing as long as  $Y > 0$ : if  $Y$  has a maximal point  $\tau$ , then  $Y(\tau) > (\delta\ell)^{p-1}$  by  $(E_Y)$ ; and

$$\alpha\ell > \alpha y(\tau) > (N - \delta)\alpha Y(\tau)/(\delta - \alpha),$$

which is again impossible. Thus  $\mathcal{T}_2$  cannot stay in  $\mathcal{Q}_1$ ; it enters  $\mathcal{Q}_2$  at some point  $(0, \xi_2)$  and stays in  $\mathcal{Q}_2$ , since it does not meet  $-\mathcal{T}_r$ . Hence  $S_w < \infty$ , and  $w$  is of type (2).

There exists also a unique trajectory  $\mathcal{T}_3$  converging to  $(0, 0)$  at  $\infty$ , ending up in  $\mathcal{Q}_1$ , since  $(0, 0)$  is a saddle point. It stays in the domain of  $\mathcal{Q}_1$  delimited by  $\mathcal{T}_1, \mathcal{T}_2$ , because  $\mathcal{Q}_1$  is backward invariant. Thus  $\mathcal{T}_3$  converges to  $M_\ell$  at  $-\infty$ , tangentially to  $u_1$ . And  $y$  is increasing on  $\mathbb{R}$ : indeed  $y' < 0$  near  $\pm\infty$ , and  $y$  cannot have two extremal points. Then  $w$  is of type (3).

For any point  $P = (\varphi, 0)$  with  $\varphi > \varphi_1$ , the trajectory  $\mathcal{T}_{[P]}$  goes from  $\mathcal{Q}_1$  into  $\mathcal{Q}_4$ , by Remark 2.1(i). It does not meet  $\mathcal{T}_r$  or  $\mathcal{T}_1$ ; hence it stays in  $\mathcal{Q}_4$  after  $P$ , and



**Figure 13.** Theorem 6.4:  $\varepsilon = -1, 0 < \alpha = 2 < \delta = 3 < N = 4$ .

$S_w < \infty$ . Before  $P$ , it stays in  $\mathfrak{D}_1$  because it does not meet  $\mathcal{T}_1$  or  $\mathcal{T}_2$ , by the same remark. By Proposition 2.8, either  $\lim_{\tau \rightarrow -\infty} \zeta = \alpha < \delta$ , so  $y' = y(\delta - \zeta) > 0$  near  $-\infty$ , and  $\lim_{\tau \rightarrow -\infty} y = \infty$ , which is impossible; or (necessarily)  $\mathcal{T}_{[P]}$  converges to  $M_\ell$ , tangentially to  $u_1$ , and  $\mathcal{T}_{[P]}$  is of type (2). Similarly, for any  $P' = (0, \xi)$  with  $\xi > \xi_2$ , the trajectory  $\mathcal{T}_{[P']}$  goes from  $\mathfrak{D}_1$  into  $\mathfrak{D}_2$ ; it remains there after  $P$  (so  $S_w < \infty$ ) and remains in  $\mathfrak{D}_1$  before  $P$ , converging to  $M_\ell$  at  $-\infty$ , tangentially to  $-u_1$ . Thus  $w$  is still of type (2).

The sets

$$\mathcal{A} = \{P \in \mathfrak{D}_1 : \mathcal{T}_{[P]} \cap \{(\varphi, 0) : \varphi > 0\} \neq \emptyset\},$$

$$\mathcal{B} = \{P \in \mathfrak{D}_1 : \mathcal{T}_{[P]} \cap \{(0, \xi) : \xi > 0\} \neq \emptyset\},$$

are open and nonempty, so  $\mathcal{A} \cup \mathcal{B} \neq \mathfrak{D}_1$ . There is at least one trajectory  $\mathcal{T}_4$  in  $\mathfrak{D}_1$  converging to  $M_\ell$  at  $-\infty$  and such that  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ ; thus  $w$  is of type (4).

For any point  $P$  in the bounded domain  $\mathcal{R}'$  of  $\mathfrak{D}_1$  delimited by  $\mathcal{T}_2$  and  $\mathcal{T}_3$ , the trajectory  $\mathcal{T}_{[P]}$  is confined to  $\mathcal{R}'$  before  $P$ , and  $y$  has no maximal point; thus  $y$  is monotone, and  $\mathcal{T}$  converges to  $M_\ell$  at  $-\infty$ . It cannot stay in  $\mathfrak{D}_1$  since it cannot converge to  $(0, 0)$ . Thus it goes from  $\mathfrak{D}_1$  into  $\mathfrak{D}_2$  and stays there, because it does not meet  $-\mathcal{T}_r$ . Thus  $S_w < \infty$ , and  $w$  is again of type (2).

For any  $P$  in the domain of  $\mathfrak{D}_1$  delimited by  $\mathcal{T}_1$  and  $\mathcal{T}_3$ , the trajectory  $\mathcal{T}_{[P]}$  converges to  $M_\ell$  at  $-\infty$ , tangentially to  $u_1$ ; it enters  $\mathfrak{D}_4$  and stays there. Thus  $S_w < \infty$  and  $w$  is of type (1). No trajectory can stay in  $\mathfrak{D}_4(\mathfrak{D}_2)$  except  $\mathcal{T}_r(-\mathcal{T}_r)$ ; thus all the solutions have been described, up to a symmetry.  $\square$

Now we come to the case  $\alpha < 0$ , and discuss according to the sign of  $\alpha - p'$ . This situation is different from the case  $\varepsilon = 1$ ,  $\delta < \min(\alpha, N)$  discussed on page 246, by Remark (i) on page 249 and part (i) of the next remark.

**Remark 6.5.** Assume  $\varepsilon = -1$  and  $\alpha < 0$ .

(i) The regular trajectory  $\mathcal{T}_r$  starts in  $\mathfrak{D}_1$ . There exists a unique trajectory  $\mathcal{T}_s$  converging to  $(0, 0)$ , lying in  $\mathfrak{D}_1$  for large  $\tau$ , having an infinite slope at  $(0, 0)$ , and satisfying  $\lim_{r \rightarrow 0} r^\eta w = c > 0$ . If  $\mathcal{T}_s$  does not stay in  $\mathfrak{D}_1$ , then  $\mathcal{T}_r$  does stay in it, and it is bounded and contained in the domain delimited by  $\mathfrak{D}_1 \cap \mathcal{T}_s$ , by Remark 2.1(i). If  $\mathcal{T}_r$  is homoclinic, it stays in  $\mathfrak{D}_1$ .

Conversely, if  $\mathcal{T}_s$  stays in  $\mathfrak{D}_1$  and is not homoclinic,  $\mathcal{T}_r$  does not stay in  $\mathfrak{D}_1$ , for the following reason.  $\mathcal{T}_s$  either converges to  $M_\ell$  at  $-\infty$  or has a limit cycle around it; if  $\mathcal{T}_r$  stays in  $\mathfrak{D}_1$ , either the corresponding  $y$  is increasing, so  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ ; or  $\lim_{\tau \rightarrow \infty} \zeta = \alpha < 0$ , by Propositions 2.15 and 2.8, so  $\mathcal{T}_r$  enters  $\mathfrak{D}_4$  and we reach a contradiction; or  $y$  oscillates around  $\ell$  near  $\infty$ , by Proposition 2.7, so it meets  $\mathcal{T}_s$ , which is impossible.

(ii) Any trajectory  $\mathcal{T}$  is bounded near  $-\infty$  from Propositions 2.8 and 2.10. Any trajectory  $\mathcal{T}$  bounded at  $\pm\infty$  converges to  $(0, 0)$  or  $\pm M_\ell$ , or its limit set  $\Gamma_\pm$  at  $\pm\infty$  is a limit cycle; or  $\mathcal{T}_r$  is homoclinic and  $\Gamma_\pm = \overline{\mathcal{T}_r}$ .

(iii) If there exists a limit cycle around  $(0, 0)$ , it also surrounds  $\pm M_\ell$ , by (2–42) and (2–43).

Next we study the case  $-p' \leq \alpha$ , where there is no cycle and no homoclinic orbit in  $\mathfrak{Q}_1$ , by Theorem 2.20.

**Theorem 6.6.** (i) Assume  $\varepsilon = -1$  and  $-p' < \alpha < 0 < \delta < N$ . Then all regular solutions have precisely one zero, and  $S_w < \infty$ . The function  $w \equiv \ell r^{-\delta}$  is a solution. There exist solutions satisfying any one of these characterizations:

- (1)  $w$  is positive,  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $\lim_{r \rightarrow \infty} r^\eta w = c > 0$ ;
- (2)  $w$  has one zero,  $\lim_{r \rightarrow 0} r^\delta w = \ell$ , and  $\lim_{r \rightarrow \alpha} r^\alpha w = L < 0$ ;
- (3)  $w$  has one zero,  $\lim_{r \rightarrow 0} r^\delta w = \ell$ , and  $S_w < \infty$ ;
- (4)  $w$  has two zeros,  $\lim_{r \rightarrow 0} r^\delta w = \ell$ , and  $S_w < \infty$ .


(ii) Assume  $\alpha = -p'$ . Then the regular solutions, given by (1–8), have one zero, and  $\lim_{r \rightarrow \alpha} r^\alpha w = L < 0$ . There exist solutions of type (1) and (4).

Up to symmetry, all solutions are as above.

*Proof.* (i) Assume  $-p' < \alpha < 0$  (Figure 14, left). By Proposition 2.5, any solution  $y$  has at most two zeros, and  $Y$  has at most one zero.

First consider  $\mathcal{T}_s$ . The function  $Y_\alpha$  defined by (2–3) with  $d = \alpha$  satisfies  $Y_\alpha = O(e^{(\alpha-\eta)\tau})$  near  $\infty$ , thus  $\lim_{\tau \rightarrow \infty} Y_\alpha = 0$ . Then from Remark 2.6,  $Y_\alpha$  is decreasing, thus  $Y_\alpha > 0$ , and  $\mathcal{T}_s$  stays in  $\mathfrak{Q}_1 \cup \mathfrak{Q}_2$ . In fact it stays in  $\mathfrak{Q}_1$ , by Remark 2.1(i). From Propositions 2.8, 2.7, 2.11, and Theorem 2.20,  $\mathcal{T}_s$  converges to  $M_\ell$  at  $-\infty$ . Indeed if  $\lim y = \infty$ , then  $\lim_{\tau \rightarrow \infty} \zeta = \alpha < 0$ ; if  $S_w < \infty$ , then  $\lim Y/y = -1$ ; which contradicts  $Y > 0$ . Then  $w$  is of type (1).

The trajectory  $\mathcal{T}_r$  stays in  $\mathfrak{Q}_1 \cup \mathfrak{Q}_2$ , and  $y$  has precisely one zero, and  $S_w < \infty$ , so  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ . We claim that  $\mathcal{T}_r$  cannot stay in  $\mathfrak{Q}_1$ . Indeed, it cannot



**Figure 14.** Theorem 6.6:  $\varepsilon = -1$ ,  $\delta = 3 < N/2 < N = 9$ .

converge to  $M_\ell$ , which is a source, or oscillate around  $\mathcal{Q}_1$ , because it does not meet  $\mathcal{T}_s$ , or tend to  $\infty$ , or satisfy  $S_w < \infty$  with  $Y > 0$ . Thus  $y$  has precisely one zero,  $\mathcal{T}_r$  enters  $\mathcal{Q}_2$  and stays in it. Moreover the corresponding  $Y_\alpha$  satisfies  $Y'_\alpha > 0$ , or equivalently (6–1). Consider again the curve  $\mathcal{N}_\alpha$  defined in (6–2). Here  $\mathcal{T}_r$  stays strictly to the right of  $\mathcal{N}_\alpha$ , and  $\mathcal{T}_s$  to the left of  $\mathcal{N}_\alpha$ .

For any  $\bar{P} = (\varphi, 0)$  with  $\varphi < 0$ , the trajectory  $\mathcal{T}_{[\bar{P}]}$  enters  $\mathcal{Q}_3$  after  $\bar{P}$ , by Remark 2.1(i). The solution going through  $\bar{P}$  at  $\tau = 0$  satisfies  $Y_\alpha(0) = 0$ ; thus  $Y_\alpha$  stays positive as before, and  $Y'_\alpha < 0$ , since  $Y_\alpha$  has no maximal point, by Remark 2.6. Thus  $\mathcal{T}_{[\bar{P}]}$  stays in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$  before  $\bar{P}$ , to the left of  $\mathcal{N}_\alpha$ . It cannot stay in  $\mathcal{Q}_2$ , by Propositions 2.7 and 2.8. As  $\tau$  decreases, it enters  $\mathcal{Q}_1$ , and converges to  $M_\ell$ , by Theorem 2.20. If  $S_w = \infty$ , then  $\lim |y| = \infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = \alpha < 0$ ; this is impossible, since  $\mathcal{T}_{[\bar{P}]}$  does not meet  $-\mathcal{T}_r$ . Thus  $S_w < \infty$ ,  $\lim Y/y = -1$ ,  $\mathcal{T}_{[\bar{P}]}$  goes from  $\mathcal{Q}_3$  into  $\mathcal{Q}_4$  and stays in it, and  $w$  is of type (4). The solution  $y$  has precisely two zeros.

Next consider  $\mathcal{T}_{[P]}$  for any  $P = (\varphi, \xi) \in \mathcal{N}_\alpha$  with  $\varphi < 0$ . The solution passing through  $P$  at  $\tau = 0$  satisfies  $Y'_\alpha(0) = 0$  and  $Y_\alpha(0) > 0$ , and 0 is a minimal point. Therefore  $Y''_\alpha(0) > 0$ ; indeed, if  $Y''_\alpha(\tau) = 0$ , we conclude from uniqueness that  $Y_\alpha$  is constant on  $\mathbb{R}$ ; then (2–6) yields  $Y_\alpha \equiv 0$ , since  $\alpha \neq -p'$ . But this cannot be. Therefore  $Y'_\alpha(\tau) > 0$  for  $\tau > 0$ ,  $Y'_\alpha(\tau) < 0$  for  $\tau < 0$ , and  $\mathcal{T}_{[P]}$  stays in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$ , to the right of  $\mathcal{N}_\alpha$  after  $P$ , with  $y < 0$  by Remark 2.1(i), and to the left of  $\mathcal{N}_\alpha$  before  $P$ . As above it cannot stay in  $\mathcal{Q}_2$  near  $-\infty$ , and converges to  $M_\ell$ . Suppose that it satisfies  $S_w = \infty$ . Then  $\lim |y| = \infty$ ,  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ , and  $\lim_{\tau \rightarrow \infty} y_\alpha = L < 0$  by Proposition 2.9; hence  $\lim_{\tau \rightarrow \infty} Y_\alpha = (\alpha L)^{p-1}$ . As in Proposition 2.5(iii), we find  $Y''_\alpha(\tau) > 0$  for any  $\tau > 0$ , which is impossible. Then  $S_w < \infty$ , so  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$  and  $w$  is of type (3).

Finally consider the domain  $\mathcal{R}$  of  $\mathcal{Q}_1 \cup \mathcal{Q}_2$  delimited by  $\mathcal{T}_r$  and  $\mathcal{T}_s$  and containing  $\mathcal{N}_\alpha$ . Form the sets

$$\begin{aligned} \mathcal{A} &= \{P \in \mathcal{R} : \mathcal{T}_{[P]} \cap \mathcal{N}_\alpha \neq \emptyset\}, \\ \mathcal{B} &= \{P \in \mathcal{R} : \mathcal{T}_{[P]} \cap \{(\xi, 0) : \xi > 0\} \neq \emptyset\}, \end{aligned}$$

corresponding to trajectories of type (3) or (4). They are nonempty and open, since here again the intersection with  $\mathcal{N}_\alpha$  is transverse ( $\alpha \neq -p'$ ). Thus  $\mathcal{A} \cup \mathcal{B}$  is distinct from  $\mathcal{R}$ : there exists a trajectory in  $\mathcal{R}$  that does not meet  $\mathcal{N}_\alpha$ ; it converges to  $M_\ell$  at  $-\infty$  or oscillates around it, and it is located below  $\mathcal{N}_\alpha$  in  $\mathcal{Q}_2$ . It cannot satisfy  $\lim_{\tau \rightarrow \ln S_w} Y/y = -1$ , so  $S_w = \infty$  and we have  $\lim_{\tau \rightarrow \infty} \zeta = \alpha$ . Hence  $w$  is of type (2).

(ii) Assume  $\alpha = -p'$  (Figure 14, right). Then regular solutions have a different behavior: they are given explicitly by (1–8). They satisfy  $Y_{-p'} \equiv C$ , thus  $Y'_{-p'} \equiv 0$ ,

thus  $\mathcal{T}_r = \mathcal{M}_{-p'}$ . Here  $y$  has a zero, and  $S_w = \infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = -p'$ . As above  $\mathcal{T}_s$  stays in  $\mathcal{Q}_1$  and  $w$  is of type (1).

Next consider again  $\mathcal{T}_{[\bar{P}]}$ . The solution going through  $\bar{P}$  at  $\tau = 0$  satisfies  $Y_{-p'}(0) = 0$ , thus  $Y_{-p'}$  stays negative for  $\tau > 0$  and  $Y'_{-p'} < 0$ . Suppose that  $S_w = \infty$ , and  $\lim_{\tau \rightarrow \infty} \zeta = -p'$ , then  $\lim_{\tau \rightarrow \infty} y_\alpha = L > 0$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = -(|\alpha|L)^{p-1}$ . But as in (2-46),  $Y''_\alpha(\tau) < 0$  for any  $\tau > 0$ , which leads to a contradiction. Then  $S_w < \infty$  and  $w$  is of type (4).

Finally suppose that there exists a trajectory  $\mathcal{T} \neq \mathcal{T}_r$  staying in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$ . Then it converges to  $M_\ell$ , thus  $Y_\alpha > 0$ ,  $S_w = \infty$ , and  $\lim_{\tau \rightarrow -\infty} Y_\alpha = \infty$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = C > 0$ . If  $\mathcal{T}$  has a minimal point, then it has an inflection point where  $Y'_\alpha > 0$ , which as above is impossible. Then  $Y'_\alpha < 0$ ; (2-6) yields

$$(p - 1)Y''_{-p'} = Y'_{-p'}(e^{p'\tau} Y_{-p'}^{(2-p)/(p-1)} - N(p - 1)) = Y'_{-p'}(Y - N(p - 1)),$$

and  $\lim_{\tau \rightarrow \infty} Y = \infty$ , so  $Y''_{-p'} < 0$  for large  $\tau$ , which is impossible. Thus there exist no solutions of type (2) or (3). □

We now come to the most difficult case:  $\alpha < -p'$ .

**Lemma 6.7.** *Assume  $\varepsilon = -1$  and  $\alpha < -p'$ . If  $\delta < N/2$  and  $\alpha^* < \alpha$ , either  $\mathcal{T}_r$  has a limit cycle in  $\mathcal{Q}_1$ , or is homoclinic, or all regular solutions have at least two zeros. If  $N/2 \leq \delta < N$ , they have at least two zeros.*

*Proof.* In any case  $M_\ell$  is a source. Suppose that  $\mathcal{T}_r$  has no limit cycle in  $\mathcal{Q}_1$ , or is not homoclinic (in particular it happens when  $N/2 \leq \delta < N$ , by Proposition 2.11), and stays in  $\mathcal{Q}_1 \cup \mathcal{Q}_2$ , thus  $Y$  stays positive. Then from Propositions 2.8, 2.9 and 2.15, either  $\lim_{\tau \rightarrow -\infty} y = \infty$ ,  $\lim_{\tau \rightarrow \infty} y_\alpha = L \neq 0$ ,  $\lim_{\tau \rightarrow \infty} Y_\alpha = (\alpha L)^{p-1}$ , or  $S_w < \infty$ . In any case, for any  $d \in (\alpha, -p')$ , the function  $Y_d = e^{(d-\alpha)\tau} Y_\alpha$  satisfies  $\lim_{\tau \rightarrow \ln S_w} Y_d = \infty = \lim_{\tau \rightarrow \infty} Y_d$ . Then it has a minimum point, and this contradicts (2-15). Thus  $\mathcal{T}_r$  enters  $\mathcal{Q}_3$ . If it stays in it, it has a limit cycle; then  $-\mathcal{T}_r$  has a limit cycle in  $\mathcal{Q}_1$ . But  $-\mathcal{T}_r$  does not meet  $\mathcal{T}_r$ , and  $M_\ell$  is in the domain of  $\mathcal{Q}_1$  delimited by  $\mathcal{T}_r$ , since  $\mathcal{T}_r$  meets  $\mathcal{M}$  to the right of  $M_\ell$ , by (2-16); this is impossible. Then  $\mathcal{T}_r$  enters  $\mathcal{Q}_4$ , and  $y$  has at least two zeros. □


**Theorem 6.8.** *Assume  $\varepsilon = -1$  and  $\delta < N/2$ ,  $\alpha < -p'$ . Then  $w(r) = \ell r^{-\delta}$  is still a solution.*

(i) *There exists a (minimal) critical value  $\alpha^{\text{crit}}$  of  $\alpha$ , such that*

$$\alpha^* < \alpha^{\text{crit}} < \min(-p', \alpha_2) < 0,$$

*and  $\mathcal{T}_r$  is homoclinic: all regular solutions have constant sign and satisfy*

$$\lim_{r \rightarrow \infty} r^\eta w = c \neq 0.$$



**Figure 15.** Theorem 6.8:  $\varepsilon = -1$ ,  $\delta = 3 < N/2 < N = 9$ .

- (ii) For any  $\alpha \in (\alpha^*, \alpha^{\text{crit}})$  there does exist a cycle in  $\mathcal{D}_1$ ; equivalently there exist solutions such that  $r^\delta w$  is periodic in  $\ln r$ . All regular solutions have constant sign and  $r^\delta w$  is asymptotically periodic in  $\ln r$ . There exist positive solutions such that  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $r^\delta w$  is asymptotically periodic in  $\ln r$ .
- (iii) For any  $\alpha \leq \alpha^*$ , there does not exist such a cycle, regular solutions have constant sign, and  $\lim_{r \rightarrow \infty} r^\delta |w| = \ell$ .
- (iv) For any  $\alpha < \alpha^{\text{crit}}$ , there exists also a cycle surrounding  $(0, 0)$  and  $\pm M_\ell$ , thus  $w$  is changing sign and  $r^\delta w$  is periodic in  $\ln r$ . There exist solutions oscillating near 0, and  $r^\delta w$  is asymptotically periodic in  $\ln r$ , and  $\lim_{r \rightarrow \infty} r^\eta w = c \neq 0$ . There exist solutions oscillating near 0, and  $r^\delta w$  is asymptotically periodic in  $\ln r$ , and  $S_w < \infty$  or  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ .

*Proof.* (i) For any  $\alpha \in (\alpha_1, \alpha_2)$ , such that  $\alpha \leq -p'$  we have three possibilities, by Remark 6.5:

- $\mathcal{T}_s$  converges to  $M_\ell$  at  $-\infty$ , spiraling around this point, since  $\alpha$  is a spiral point, or it has a limit cycle around  $M_\ell$ . Then  $\mathcal{T}_s$  meets the set  $\mathcal{E} = \{(\ell, Y) :$

$Y > (\delta\ell)^{p-1}$  at a first point  $(\ell, Y_s(\alpha))$ ; and  $\mathcal{T}_r$  meets  $\mathcal{E}$  at a last point  $(\ell, Y_r(\alpha))$  such that  $Y_r(\alpha) - Y_s(\alpha) > 0$ . Moreover  $\mathcal{T}_r$  enters  $\mathcal{Q}_2$ , by Proposition 2.8. See Figure 15, top left.

- $\mathcal{T}_s$  enters  $\mathcal{Q}_4$ ; hence  $\mathcal{T}_r$  converges to  $M_\ell$  at  $\infty$  and spirals around this point, or it has a limit cycle around  $M_\ell$ . Then  $\mathcal{T}_s$  meets  $\mathcal{E}$  at a last point  $(\ell, Y_s(\alpha))$ ,  $\mathcal{T}_r$  meets  $\mathcal{E}$  at a first point  $(\ell, Y_r(\alpha))$  such that  $Y_r(\alpha) - Y_s(\alpha) < 0$ . See Figure 15, bottom row.
- $\mathcal{T}_r$  is homoclinic, or equivalently  $Y_r(\alpha) - Y_s(\alpha) = 0$ . See Figure 15, top right.

Now the function  $\alpha \mapsto h(\alpha) = Y_r(\alpha) - Y_s(\alpha)$  is continuous. If  $-p' < \alpha_2$ , then  $h(-p')$  is defined and  $h(-p') > 0$ , by Theorem 6.6. If  $\alpha_2 \leq -p'$ , we observe that for  $\alpha = \alpha_2$ , by Theorem 2.18,  $\mathcal{T}_r$  must leave  $\mathcal{Q}_1$  (because  $\alpha_2$  is a source) and does so transversally; thus the same holds for  $\alpha = \alpha_2 - \gamma$  if  $\gamma > 0$  is small enough. Therefore  $\mathcal{T}_s$  stays in  $\mathcal{Q}_1$  by Remark 6.5, so  $h(\alpha_2 - \gamma) > 0$ . If  $\alpha \leq \alpha^*$ , then  $M_\ell$  is a sink or a weak sink, by Theorem 2.16; therefore  $\mathcal{T}_s$  cannot converge to  $M_\ell$  at  $-\infty$ . By Theorem 2.19, there are no cycles in  $\mathcal{Q}_1$  and no homoclinic orbits. By Remark 6.5,  $\mathcal{T}_s$  cannot stay in  $\mathcal{Q}_1$ ; hence  $\mathcal{T}_r$  stays in  $\mathcal{Q}_1$  and is bounded and converges at  $\infty$  to  $M_\ell$ . Thus  $h(\alpha) < 0$  for  $\alpha_1 < \alpha \leq \alpha^*$ , so there exists at least an  $\alpha^{\text{crit}} \in (\alpha^*, \min(-p', \alpha_2))$  such that  $h(\alpha^{\text{crit}}) = 0$ . If it is not unique, we choose the smallest one.

(ii) Let  $\alpha > \alpha^*$ . The existence and uniqueness of such a cycle in  $\mathcal{Q}_1$  follows from Theorem 2.16 if  $\alpha - \alpha^*$  is small enough (Figure 15, lower left). For any  $\alpha \in (\alpha^*, \alpha^{\text{crit}})$ , we still have existence: indeed,  $h(\alpha) < 0$  on this interval, so  $\mathcal{T}_r$  stays in  $\mathcal{Q}_1$ , and  $\mathcal{T}_r$  cannot converge to  $M_\ell$  at  $\infty$ , hence it has a limit cycle around  $M_\ell$  at  $\infty$ . Since  $M_\ell$  is a source, there also exist trajectories converging to  $M_\ell$  at  $-\infty$ , with a limit cycle at  $\infty$ . And  $\mathcal{T}_s$  does not stay in  $\mathcal{Q}_1$ , and it is bounded at  $-\infty$ . Thus it has a limit cycle at  $-\infty$  surrounding  $(0, 0)$  and  $\pm M_\ell$ .

(iii) Let  $\alpha \leq \alpha^*$  (Figure 15, lower right). Then  $\mathcal{T}_r$  stays in  $\mathcal{Q}_1$ , is bounded on  $\mathbb{R}$ , and converges to  $M_\ell$  at  $\infty$ , while  $\mathcal{T}_s$  does not stay in  $\mathcal{Q}_1$  as above. Thus  $\mathcal{T}_s$  has a limit cycle at  $-\infty$ , containing the three stationary points.

(iv) For any  $\alpha < \alpha^{\text{crit}}$  apart from  $\mathcal{T}_r$  and the cycles, all trajectories have a limit cycle at  $-\infty$  containing the three stationary points. By Theorem 2.21, all the cycles are contained in a ball  $B$  of  $\mathbb{R}^2$ . Take any point  $P$  exterior to  $B$ . By Remark 6.5,  $\mathcal{T}_{[P]}$  has a limit cycle at  $-\infty$  contained in  $B$  and cannot have a limit cycle at  $\infty$ . Therefore  $y$  has constant sign near  $\ln S_w$ . By Proposition 2.8, either  $S_w < \infty$  or  $y$  is defined near  $\infty$  and  $\lim_{\tau \rightarrow \infty} \zeta = L$ ,  $\lim_{r \rightarrow \infty} r^\alpha w = L$ .  $\square$

Finally we consider the case  $N/2 \leq \delta$ , where no cycle can exist.

**Theorem 6.9.** *Assume  $\varepsilon = -1$  and  $\alpha < 0 < N/2 \leq \delta < N$ . Then all solutions of  $(E_w)$  have a finite number of zeros, and  $w(r) = \ell r^{-\delta}$  is a solution. If  $-p' \leq \alpha$ , Theorem 6.6 applies. If  $\alpha < -p'$ , there exist positive solutions such that  $\lim_{r \rightarrow 0} r^\delta w = \ell$  and  $\lim_{r \rightarrow \infty} r^\eta w = c > 0$ . All regular solutions have the same number  $m \geq 2$  of zeros. All other solutions satisfy  $\lim_{r \rightarrow -\infty} r^\delta w = \pm \ell$ , and have  $m$  or  $m + 1$  zeros; there exist solutions with  $m + 1$  zeros.*

*Proof.* By Proposition 2.11, all solutions have a finite number of zeros, and any solution is monotone near 0 and  $\ln S_w$ , or converges to  $\pm M_\ell$ . By Remark 6.5, apart from  $\mathcal{T}_r$ , all trajectories converge to  $\pm M_\ell$  at  $-\infty$ . The functions  $V$  and  $W$  are nonincreasing. The trajectory  $\mathcal{T}_s$  satisfies  $\lim_{\tau \rightarrow \infty} V = \lim_{\tau \rightarrow \infty} W = 0$ , so  $V \geq 0$ ,  $W \geq 0$ . If  $y$  has a zero at some point  $\tau$ , then  $W(\tau) = -|Y(\tau)|^{p'}/p'$ , which is impossible. If  $Y$  has a zero at some point  $\theta$ , then  $V(\theta) = -Y'(\theta)^2/2$ , also a contradiction. Thus  $\mathcal{T}_s$  stays in  $\mathcal{D}_1$ . By Remark 6.5 and Proposition 2.11,  $\mathcal{T}_r$  does not stay in  $\mathcal{D}_1$ , but enters  $\mathcal{D}_2$ . By Lemma 6.7,  $\mathcal{T}_r$  enters  $\mathcal{D}_4$ , and  $y$  has at least two zeros. Let  $m$  be the number of its zeros. Then  $\mathcal{T}_r$  cuts the axis  $y = 0$  at points  $(0, \xi_1), \dots, (0, \xi_m)$ . Consider any trajectory  $\mathcal{T}_{[P]}$  with  $P = (0, \xi)$ , where  $\xi > |\xi_i|$  for  $1 \leq i \leq m$ . It cannot intersect  $\mathcal{T}_r$  or  $-\mathcal{T}_r$ , so  $y$  has  $m + 1$  zeros. Any trajectory has  $m$  or  $m + 1$  zeros, because it does not meet  $\mathcal{T}_r$  or  $-\mathcal{T}_r$  or  $\mathcal{T}_{[P]}$ . And  $S_w < \infty$  or  $\lim_{r \rightarrow \infty} r^\alpha w = L \neq 0$ .  $\square$

### Acknowledgment

We thank Hector Giacomini (University of Tours) for interesting discussions on the existence of cycles in dynamical systems.

### References

- [Anderson and Leighton 1968] L. R. Anderson and W. Leighton, “Liapunov functions for autonomous systems of second order”, *J. Math. Anal. Appl.* **23** (1968), 645–664. MR 37 #5488 Zbl 0186.41602
- [Barenblatt 1952] G. I. Barenblatt, “On self-similar motions of a compressible fluid in a porous medium”, *Akad. Nauk SSSR. Prikl. Mat. Meh.* **16** (1952), 679–698. MR 14,699h Zbl 0047.19204
- [Bidaut-Véron 1989] M.-F. Bidaut-Véron, “Local and global behavior of solutions of quasilinear equations of Emden–Fowler type”, *Arch. Rational Mech. Anal.* **107**:4 (1989), 293–324. MR 90f:35066 Zbl 0696.35022
- [Bidaut-Véron 2006a] M. F. Bidaut-Véron, “The  $p$ -Laplace heat equation with a source term: self-similar solutions revisited”, *Adv. Nonlinear Stud.* **6**:1 (2006), 69–108. MR MR2196892
- [Bidaut-Véron 2006b] M. F. Bidaut-Véron, “Self-similar solutions of the  $p$ -Laplace heat equation: the case  $p > 2$ ”, 2006. In preparation.
- [Chasseigne and Vazquez 2002] E. Chasseigne and J. L. Vazquez, “Theory of extended solutions for fast-diffusion equations in optimal classes of data. Radiation from singularities”, *Arch. Ration. Mech. Anal.* **164**:2 (2002), 133–187. MR 2003i:35148 Zbl 1018.35048

- [Chicone and Tian 1982] C. Chicone and J. H. Tian, “On general properties of quadratic systems”, *Amer. Math. Monthly* **89**:3 (1982), 167–178. MR 84f:34044 Zbl 0466.34011
- [DiBenedetto and Herrero 1990] E. DiBenedetto and M. A. Herrero, “Nonnegative solutions of the evolution  $p$ -Laplacian equation. Initial traces and Cauchy problem when  $1 < p < 2$ ”, *Arch. Rational Mech. Anal.* **111**:3 (1990), 225–290. MR 92g:35088
- [Gil and Vázquez 1997] O. Gil and J. L. Vázquez, “Focusing solutions for the  $p$ -Laplacian evolution equation”, *Adv. Differential Equations* **2**:2 (1997), 183–202. MR 97g:35069 Zbl 1023.35514
- [Hale and Koçak 1991] J. K. Hale and H. Koçak, *Dynamics and bifurcations*, Texts in Applied Mathematics **3**, Springer, New York, 1991. MR 93e:58047 Zbl 0745.58002
- [Hubbard and West 1995] J. H. Hubbard and B. H. West, *Differential equations: a dynamical systems approach*, v. 2: *Higher dimensional systems*, Texts in Applied Mathematics **18**, Springer, New York, 1995. MR 96e:34001 Zbl 0824.34001
- [Kamin and Vázquez 1992] S. Kamin and J. L. Vázquez, “Singular solutions of some nonlinear parabolic equations”, *J. Anal. Math.* **59** (1992), 51–74. MR 94e:35079 Zbl 0802.35066
- [Qi and Wang 1999] Y.-W. Qi and M. Wang, “The global existence and finite time extinction of a quasilinear parabolic equation”, *Adv. Differential Equations* **4**:5 (1999), 731–753. MR 2000d:35106 Zbl 0959.35104
- [Vazquez and Véron 1996] J. L. Vazquez and L. Véron, “Different kinds of singular solutions of nonlinear parabolic equations”, pp. 240–249 in *Nonlinear problems in applied mathematics: in honor of Ivar Stakgold on his 70th birthday*, Soc. Ind. Appl. Math., Philadelphia, 1996. Zbl 0886.35078
- [Zhao 1993] J. N. Zhao, “The asymptotic behaviour of solutions of a quasilinear degenerate parabolic equation”, *J. Differential Equations* **102**:1 (1993), 33–52. MR 94c:35040 Zbl 0816.35070

Received February 4, 2005. Revised January 2, 2006.

MARIE FRANÇOISE BIDAUT-VÉRON  
LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE  
CNRS UMR 6083  
FACULTÉ DES SCIENCES ET TECHNIQUES  
PARC GRANDMONT  
37200 TOURS  
FRANCE  
veronmf@univ-tours.fr

