

Pacific Journal of Mathematics

**IRREDUCIBLE REPRESENTATIONS FOR THE ABELIAN
EXTENSION OF THE LIE ALGEBRA OF DIFFEOMORPHISMS
OF TORI IN DIMENSIONS GREATER THAN 1**

CUIPO JIANG AND QIFEN JIANG

Volume 231 No. 1

May 2007

IRREDUCIBLE REPRESENTATIONS FOR THE ABELIAN EXTENSION OF THE LIE ALGEBRA OF DIFFEOMORPHISMS OF TORI IN DIMENSIONS GREATER THAN 1

CUIPO JIANG AND QIFEN JIANG

We classify the irreducible weight modules of the abelian extension of the Lie algebra of diffeomorphisms of tori of dimension greater than 1, with finite-dimensional weight spaces.

1. Introduction

Let $W_{\nu+1}$ be the Lie algebra of diffeomorphisms of the $(\nu+1)$ -dimensional torus. If $\nu = 0$, the universal central extension of the complex Lie algebra W_1 is the Virasoro algebra, which, together with its representations, plays a very important role in many areas of mathematics and physics [Belavin et al. 1984; Dotsenko and Fateev 1984; Di Francesco et al. 1997]. The representation theory of the Virasoro algebra has been studied extensively; see, for example, [Kac 1982; Kaplansky and Santharoubane 1985; Chari and Pressley 1988; Mathieu 1992].

If $\nu \geq 1$, however, the Lie algebra $W_{\nu+1}$ has no nontrivial central extension [Ramos et al. 1990]. But $W_{\nu+1}$ has abelian extensions whose abelian ideals are the central parts of the corresponding toroidal Lie algebras; see [Berman and Billig 1999], for example. There is a close connection between irreducible integrable modules of the toroidal Lie algebra and irreducible modules of the abelian extension \mathcal{L} ; see [Berman and Billig 1999; Eswara Rao and Moody 1994; Jiang and Meng 2003], for instance. In fact, the classification of integrable modules of toroidal Lie algebras and their subalgebras depends heavily on the classification of irreducible representations of \mathcal{L} and its subalgebras. See [Billig 2003] for the constructions of the abelian extensions for the group of diffeomorphisms of a torus.

In this paper we study the irreducible weight modules of \mathcal{L} , for $\nu \geq 1$. If V is an irreducible weight module of \mathcal{L} some of whose central charges c_0, \dots, c_ν are nonzero, one can assume that c_0, \dots, c_N are \mathbb{Z} -linearly independent and $c_{N+1} = \dots = c_\nu = 0$, where $N \geq 0$. We prove that if $N \geq 1$, then V must have weight

MSC2000: primary 17B67, 17B65; secondary 17B68.

Keywords: irreducible representation, abelian extension, central charge.

Work supported in part by NSF of China, grants No. 10271076 and No. 10571119.

spaces which are infinite-dimensional. So if all the weight spaces of V are finite-dimensional, N vanishes. We classify the irreducible modules of \mathcal{L} with finite-dimensional weight spaces and some nonzero central charges. We prove that such a module V is isomorphic to a highest weight module. The highest weight space T is isomorphic to an irreducible $(\mathcal{A}_v + W_v)$ -module all of whose weight spaces have the same dimension, where \mathcal{A}_v is the ring of Laurent polynomials in v commuting variables, regarded as a commutative Lie algebra. An important step is to characterize the \mathcal{A}_v -module structure of T . It turns out that the action of \mathcal{A}_v on T is essentially multiplication by polynomials in \mathcal{A}_v . Therefore T can be identified with Larsson's construction [1992] by a result in [Eswara Rao 2004]. That is, T is a tensor product of gl_v -module with \mathcal{A}_v .

When all the central charges of V are zero, we prove that the abelian part acts on V as zero if V is a uniformly bounded \mathcal{L} -module. So the result in this case is not complete.

Throughout the paper, \mathbb{C} , \mathbb{Z}_+ and \mathbb{Z}_- denote the sets of complex numbers, positive integers and negative integers.

2. Basic concepts and results

Let $\mathcal{A}_{v+1} = \mathbb{C}[t_0^{\pm 1}, t_1^{\pm 1}, \dots, t_v^{\pm 1}]$ ($v \geq 1$) be the ring of Laurent polynomials in commuting variables t_0, t_1, \dots, t_v . For $\underline{n} = (n_1, n_2, \dots, n_v) \in \mathbb{Z}^v$, $n_0 \in \mathbb{Z}$, we denote $t_0^{n_0} t_1^{n_1} \dots t_v^{n_v}$ by $t_0^{n_0} t^{\underline{n}}$. Let $\tilde{\mathcal{K}}$ be the free \mathcal{A}_{v+1} -module with basis $\{k_0, k_1, \dots, k_v\}$ and let $d\tilde{\mathcal{K}}$ be the subspace spanned by all elements of the form

$$\sum_{i=0}^v r_i t_0^{r_0} t^{\underline{r}} k_i, \quad \text{for } (r_0, \underline{r}) = (r_0, r_1, \dots, r_v) \in \mathbb{Z}^{v+1}.$$

Set $\mathcal{K} = \tilde{\mathcal{K}}/d\tilde{\mathcal{K}}$ and denote the image of $t_0^{r_0} t^{\underline{r}} k_i$ still by itself. Then \mathcal{K} is spanned by the elements $\{t_0^{r_0} t^{\underline{r}} k_p \mid p = 0, 1, \dots, v, r_0 \in \mathbb{Z}, \underline{r} \in \mathbb{Z}^v\}$ with relations

$$(2-1) \quad \sum_{p=0}^v r_p t_0^{r_0} t^{\underline{r}} k_p = 0.$$

Let \mathcal{D} be the Lie algebra of derivations on \mathcal{A}_{v+1} . Then

$$\mathcal{D} = \left\{ \sum_{p=0}^v f_p(t_0, t_1, \dots, t_v) d_p \mid f_p(t_0, t_1, \dots, t_v) \in \mathcal{A}_{v+1} \right\},$$

where $d_p = t_p \partial / \partial t_p$, $p = 0, 1, \dots, v$. From [Berman and Billig 1999] we know that the algebra \mathcal{D} admits two nontrivial 2-cocycles with values in \mathcal{K} :

$$\tau_1(t_0^{m_0} t^{\underline{m}} d_a, t_0^{n_0} t^{\underline{n}} d_b) = -n_a m_b \sum_{p=0}^v m_p t_0^{m_0+n_0} t^{\underline{m}+\underline{n}} k_p,$$

$$\tau_2(t_0^{m_0} t^m d_a, t_0^{n_0} t^n d_b) = m_a n_b \sum_{p=0}^v m_p t_0^{m_0+n_0} t^{m+n} k_p.$$

Let $\tau = \mu_1 \tau_1 + \mu_2 \tau_2$ be an arbitrary linear combination of τ_1 and τ_2 . Then the corresponding abelian extension of \mathcal{D} is

$$\mathcal{L} = \mathcal{D} \oplus \mathcal{H},$$

with the Lie bracket

$$(2-2) \quad [t_0^{m_0} t^m d_a, t_0^{n_0} t^n k_b] = n_a t_0^{m_0+n_0} t^{m+n} k_b + \delta_{ab} \sum_{p=0}^v m_p t_0^{m_0+n_0} t^{m+n} k_p,$$

$$[t_0^{m_0} t^m d_a, t_0^{n_0} t^n d_b] = n_a t_0^{m_0+n_0} t^{m+n} d_b - m_b t_0^{m_0+n_0} t^{m+n} d_a + \tau(t_0^{m_0} t^m d_a, t_0^{n_0} t^n d_b).$$

The sum

$$\mathfrak{h} = \left(\bigoplus_{i=0}^v \mathbb{C} k_i \right) \oplus \left(\bigoplus_{i=0}^v \mathbb{C} d_i \right)$$

is an abelian Lie subalgebra of \mathcal{L} . An \mathcal{L} -module V is called a weight module if

$$V = \bigoplus_{\lambda \in \mathfrak{h}^*} V_\lambda,$$

where $V_\lambda = \{v \in V \mid h \cdot v = \lambda(h)v \text{ for all } h \in \mathfrak{h}\}$. Denote by $P(V)$ the set of all weights. Throughout the paper, we assume that V is an irreducible weight module of \mathcal{L} with finite-dimensional weight spaces. Since V is irreducible, we have

$$k_i|_V = c_i,$$

where the constants c_i , for $i = 0, 1, \dots, v$, are called the central charges of V .

Lemma 2.1. *Let $A = (a_{ij})$ ($0 \leq i, j \leq v$) be a $(v+1) \times (v+1)$ matrix such that $\det A = 1$ and $a_{ij} \in \mathbb{Z}$. There exists an automorphism σ of \mathcal{L} such that*

$$\sigma(t^{\bar{m}} k_j) = \sum_{p=0}^v a_{pj} t^{\bar{m} A^T} k_p, \quad \sigma(t^{\bar{m}} d_j) = \sum_{p=0}^v b_{jp} t^{\bar{m} A^T} d_p, \quad 0 \leq j \leq v,$$

where $t^{\bar{m}} = t_0^{m_0} t^m$, $B = (b_{ij}) = A^{-1}$.

3. The structure of V with nonzero central charges

In this section, we discuss the weight module V which has nonzero central charges. It follows from Lemma 2.1 that we can assume that c_0, c_1, \dots, c_N are \mathbb{Z} -linearly independent, i.e., if $\sum_{i=0}^N a_i c_i = 0$, $a_i \in \mathbb{Z}$, then all a_i ($i = 0, \dots, N$) must be zero,

and $c_{N+1} = c_{N+2} = \cdots = c_\nu = 0$, where $N \geq 0$. For $\bar{m} = (m_0, \underline{m})$, denote $t_0^{m_0} t^{\underline{m}}$ by $t^{\bar{m}}$ as in Lemma 2.1. It is easy to see that V has the decomposition

$$V = \bigoplus_{\bar{m} \in \mathbb{Z}^{\nu+1}} V_{\bar{m}},$$

where $V_{\bar{m}} = \{v \in V \mid d_i(v) = (\gamma_0(d_i) + m_i)v, i = 0, 1, \dots, \nu\}$, with $\gamma_0 \in P(V)$ a fixed weight, and $\bar{m} = (m_0, m_1, \dots, m_\nu) \in \mathbb{Z}^{\nu+1}$. If V has finite-dimensional weight spaces, the $V_{\bar{m}}$ are finite-dimensional, for $\bar{m} \in \mathbb{Z}^{\nu+1}$.

In Lemmas 3.1–3.6 we assume that V has finite-dimensional weight spaces.

Lemma 3.1. *For $p \in \{0, 1, \dots, \nu\}$ and $0 \neq t^{\bar{m}} k_p \in \mathcal{L}$, if there is a nonzero element v in V such that $t^{\bar{m}} k_p v = 0$, then $t^{\bar{m}} k_p$ is locally nilpotent on V .*

Lemma 3.2. *Let $t_0^{m_0} t^{\underline{m}} k_p \in \mathcal{L}$ be such that $\bar{m} = (m_0, \underline{m}) \neq \bar{0}$, and there exists $0 \leq a \leq N$ such that $m_a \neq 0$ if $N < p \leq \nu$. If $t_0^{m_0} t^{\underline{m}} k_p$ is locally nilpotent on V , then $\dim V_{\bar{n}} > \dim V_{\bar{n}+\bar{m}}$ for all $\bar{n} \in \mathbb{Z}^{\nu+1}$.*

Proof. Case 1: $p \in \{0, 1, \dots, N\}$. We first prove that $\dim V_{\bar{n}} \geq \dim V_{\bar{n}+\bar{m}}$ for all $\bar{n} \in \mathbb{Z}^{\nu+1}$. Suppose $\dim V_{\bar{n}} = m$, $\dim V_{\bar{n}+\bar{m}} = n$. Let $\{w_1, w_2, \dots, w_n\}$ be a basis of $V_{\bar{n}+\bar{m}}$ and $\{w'_1, w'_2, \dots, w'_m\}$ a basis of $V_{\bar{n}}$. We can assume that $m_a \neq 0$ for some $0 \leq a \leq \nu$ distinct from p , where $\bar{m} = (m_0, \underline{m}) = (m_0, m_1, \dots, m_\nu)$. Since $t^{\bar{m}} k_p$ is locally nilpotent on V and $V_{\bar{n}+\bar{m}}$ is finite-dimensional, there exists $k > 0$ such that $(t^{\bar{m}} k_p)^k V_{\bar{n}+\bar{m}} = 0$. Therefore

$$(t^{-\bar{m}} d_a)^k (t^{\bar{m}} k_p)^k (w_1, w_2, \dots, w_n) = 0.$$

On the other hand, by induction on k , we can deduce that

$$(t^{-\bar{m}} d_a)^k (t^{\bar{m}} k_p)^k = \sum_{i=0}^k \frac{k! k!}{i! (k-i)! (k-i)!} m_a^i c_p^i (t^{\bar{m}} k_p)^{k-i} (t^{-\bar{m}} d_a)^{k-i}.$$

Therefore

$$\begin{aligned} t^{\bar{m}} k_p \left(\sum_{i=0}^{k-1} \frac{k! k!}{i! (k-i)! (k-i)!} m_a^i c_p^i (t^{\bar{m}} k_p)^{k-1-i} (t^{-\bar{m}} d_a)^{k-1-i} \right) t^{-\bar{m}} d_a (w_1, w_2, \dots, w_n) \\ = -k! m_a^k c_p^k (w_1, w_2, \dots, w_n). \end{aligned}$$

Assume that

$$\begin{aligned} \left(\sum_{i=0}^{k-1} \frac{k! k!}{i! (k-i)! (k-i)!} m_a^i c_p^i (t^{\bar{m}} k_p)^{k-1-i} (t^{-\bar{m}} d_a)^{k-1-i} \right) t^{-\bar{m}} d_a (w_1, w_2, \dots, w_n) \\ = (w'_1, w'_2, \dots, w'_m) C, \end{aligned}$$

with $C \in \mathbb{C}^{m \times n}$, and that

$$(3-1) \quad t^{\bar{m}} k_p (w'_1, w'_2, \dots, w'_m) = (w_1, w_2, \dots, w_n) B,$$

with $B \in \mathbb{C}^{n \times m}$. Then

$$BC = -k! m_a^k c_p^k I.$$

This implies that $m \geq n$. So $\dim V_{\bar{n}} \geq \dim V_{\bar{n}+\bar{m}}$ for all $\bar{n} \in \mathbb{Z}^{v+1}$. Also, by (3-1) and the fact that $r(B) = n$, we know that $m > n$ if and only if there exists $v \in V_{\bar{n}}$ such that $t^{\bar{m}} k_p \cdot v = 0$. Since $t^{\bar{m}} k_p$ is locally nilpotent on V , there exist an integer $s \geq 0$ and $w \in V_{\bar{n}+s\bar{m}}$ such that

$$(t^{\bar{m}} k_p) \cdot w = 0.$$

Therefore $(t^{-\bar{m}} k_p) t^{\bar{m}} k_p \cdot w = t^{\bar{m}} k_p (t^{-\bar{m}} k_p \cdot w) = 0$. If $t^{-\bar{m}} k_p \cdot w = 0$, by the proof above, $\dim V_{\bar{n}+s\bar{m}-\bar{m}} < \dim V_{\bar{n}+s\bar{m}}$, contradicting the fact that $\dim V_{\bar{n}+s\bar{m}-\bar{m}} \geq \dim V_{\bar{n}+s\bar{m}}$. Therefore $(t^{-\bar{m}} k_p)^r \cdot w \neq 0$ for all $r \in \mathbb{N}$. Since

$$(t^{-\bar{m}} k_p)^s t^{\bar{m}} k_p \cdot w = t^{\bar{m}} k_p (t^{-\bar{m}} k_p)^s \cdot w = 0$$

and $(t^{-\bar{m}} k_p)^s \cdot w \in V_{\bar{n}}$, it follows that there is a nonzero element v in $V_{\bar{n}}$ such that $t^{\bar{m}} k_p \cdot v = 0$. Thus $n < m$.

Case 2: $N < p \leq v$. The proof is similar to that of case 1, but we have to consider $t^{-\bar{m}} d_p$ and $t^{\bar{m}} k_p$ instead and use the \mathbb{Z} -linear independence of c_1, \dots, c_N . \square

Lemma 3.3. *Let $0 \neq t^{\bar{m}} k_p \in \mathcal{L}$ and $0 \neq t^{\bar{n}} k_p \in \mathcal{L}$ be such that $(m_0, \dots, m_N) \neq 0$, $(n_0, \dots, n_N) \neq 0$ if $N < p \leq v$, where $\bar{m} = (m_0, m_1, \dots, m_v)$.*

- (1) *If $t^{\bar{m}} k_p$ is locally nilpotent on V , $t^{\bar{m}} k_q$ is locally nilpotent for $q = 0, 1, \dots, v$.*
- (2) *If both $0 \neq t^{\bar{m}} k_p$ and $0 \neq t^{\bar{n}} k_p$ are locally nilpotent on V , then $t^{\bar{m}+\bar{n}} k_p$ is locally nilpotent.*
- (3) *If $0 \neq t^{\bar{m}+\bar{n}} k_p$ is locally nilpotent on V and $(m_0 + n_0, \dots, m_N + n_N) \neq 0$ if $N < p \leq v$, then $t^{\bar{m}} k_p$ or $t^{\bar{n}} k_p$ is locally nilpotent.*

Lemma 3.4. *For $0 \leq p \leq v$, let $0 \neq t^{\bar{m}} k_p \in \mathcal{L}$ be such that $(m_0, \dots, m_N) \neq 0$, where $\bar{m} = (m_0, m_1, \dots, m_v)$. Then $t^{\bar{m}} k_p$ or $t^{-\bar{m}} k_p$ is locally nilpotent on V .*

Proof. The proof occupies the next few pages. We first deal with the case $0 \leq p \leq N$. Without losing generality, we can take $p = 0$.

Suppose the lemma is false. By Lemma 3.2, for any $\bar{r} \in \mathbb{Z}^{v+1}$ we have

$$\dim V_{\bar{r}+\bar{m}} = \dim V_{\bar{r}} = \dim V_{\bar{r}-\bar{m}}, \quad t^{\bar{m}} k_0 V_{\bar{r}} = V_{\bar{r}+\bar{m}}, \quad t^{-\bar{m}} k_0 V_{\bar{r}} = V_{\bar{r}-\bar{m}}.$$

Fix $\bar{r} = (r_0, \underline{r}) \in \mathbb{Z}^{v+1}$ such that $V_{\bar{r}} \neq 0$. Let $\{v_1, \dots, v_n\}$ be a basis of $V_{\bar{r}}$ and set

$$v_i(k\bar{m}) = \frac{1}{c_0} t^{k\bar{m}} k_0 \cdot v_i, \quad i = 1, 2, \dots, n,$$

where $k \in \mathbb{Z} \setminus \{0\}$. Then $\{v_1(k\bar{m}), v_2(k\bar{m}), \dots, v_n(k\bar{m})\}$ is a basis of $V_{\bar{r}+k\bar{m}}$. Let $B_{-\bar{m}, \bar{m}}^{(0)}, B_{\bar{m}, -\bar{m}}^{(0)} \in \mathbb{C}^{n \times n}$ be such that

$$\begin{aligned} \frac{1}{c_0} t^{\bar{m}} k_0(v_1(-\bar{m}), v_2(-\bar{m}), \dots, v_n(-\bar{m})) &= (v_1, v_2, \dots, v_n) B_{\bar{m}, -\bar{m}}^{(0)}, \\ \frac{1}{c_0} t^{-\bar{m}} k_0(v_1(\bar{m}), v_2(\bar{m}), \dots, v_n(\bar{m})) &= (v_1, v_2, \dots, v_n) B_{-\bar{m}, \bar{m}}^{(0)}. \end{aligned}$$

Since $t^{\bar{m}} k_0$ and $t^{-\bar{m}} k_0$ are commutative, it is easy to deduce that

$$B_{\bar{m}, -\bar{m}}^{(0)} = B_{-\bar{m}, \bar{m}}^{(0)}.$$

By Lemma 3.1, $B_{\bar{m}, -\bar{m}}^{(0)}$ is an $n \times n$ invertible matrix.

Claim. $B_{\bar{m}, -\bar{m}}^{(0)}$ does not have distinct eigenvalues.

Proof. Set $c = 1/c_0$. To prove the claim, we need to consider $ct^{\bar{m}} k_0 ct^{-\bar{m}} k_0 - \lambda \text{id}$, where $\lambda \in \mathbb{C}^*$. As in the proof of Lemma 3.1, we can deduce that if there is a nonzero element v in V such that $(ct^{\bar{m}} k_0 ct^{-\bar{m}} k_0 - \lambda \text{id})v = 0$, then $ct^{\bar{m}} k_0 ct^{-\bar{m}} k_0 - \lambda \text{id}$ is locally nilpotent on V . On the other hand, we have

$$(ct^{\bar{m}} k_0 ct^{-\bar{m}} k_0 - \lambda \text{id})^l(v_1, v_2, \dots, v_n) = (v_1, v_2, \dots, v_n)(B_{\bar{m}, -\bar{m}}^{(0)} - \lambda \text{id})^l.$$

Therefore the claim holds. □

For $p \in \{1, 2, \dots, v\}$, let $C_{\bar{m}, \bar{0}}^p, C_{\bar{m}, -\bar{m}}^p \in \mathbb{C}^{n \times n}$ be such that

$$\begin{aligned} t^{\bar{m}} k_p(v_1, v_2, \dots, v_n) &= (v_1(\bar{m}), \dots, v_n(\bar{m})) C_{\bar{m}, \bar{0}}^{(p)}, \\ t^{\bar{m}} k_p(v_1(-\bar{m}), \dots, v_n(-\bar{m})) &= (v_1, v_2, \dots, v_n) C_{\bar{m}, -\bar{m}}^{(p)}. \end{aligned}$$

Since

$$\frac{1}{c_0} t^{-\bar{m}} k_0 t^{\bar{m}} k_p(v_1, v_2, \dots, v_n) = t^{\bar{m}} k_p \frac{1}{c_0} t^{-\bar{m}} k_0(v_1, v_2, \dots, v_n),$$

we have

$$(3-2) \quad C_{\bar{m}, -\bar{m}}^{(p)} = B_{-\bar{m}, \bar{m}}^{(0)} C_{\bar{m}, \bar{0}}^{(p)}.$$

Furthermore, by the fact that

$$\frac{1}{c_0} t^{\bar{m}} k_0 \frac{1}{c_0} t^{-\bar{m}} k_0 t^{\bar{m}} k_p(v_1, v_2, \dots, v_n) = t^{\bar{m}} k_p \frac{1}{c_0} t^{\bar{m}} k_0 \frac{1}{c_0} t^{-\bar{m}} k_0(v_1, v_2, \dots, v_n)$$

and

$$t^{\bar{m}} k_q \frac{1}{c_0} t^{-\bar{m}} k_0 t^{\bar{m}} k_p = t^{\bar{m}} k_p \frac{1}{c_0} t^{-\bar{m}} k_0 t^{\bar{m}} k_q,$$

we deduce that

$$(3-3) \quad B_{-\bar{m}, \bar{m}}^{(0)} C_{\bar{m}, \bar{0}}^{(p)} = C_{\bar{m}, \bar{0}}^{(p)} B_{-\bar{m}, \bar{m}}^{(0)}, \quad C_{\bar{m}, \bar{0}}^{(p)} C_{\bar{m}, \bar{0}}^{(q)} = C_{\bar{m}, \bar{0}}^{(q)} C_{\bar{m}, \bar{0}}^{(p)}, \quad 1 \leq p, q \leq \nu.$$

Hence there exists $D \in \mathbb{C}^{n \times n}$ such that $\{D^{-1} B_{-\bar{m}, \bar{m}}^{(0)} D, D^{-1} C_{\bar{m}, \bar{0}}^{(p)} D \mid 1 \leq p \leq \nu\}$ are all upper triangular matrices. If we set

$$(w_1, w_2, \dots, w_n) = (v_1, v_2, \dots, v_n) D$$

and

$$w_i(k\bar{m}) = \frac{1}{c_0} t^{k\bar{m}} k_0 w_i, \quad 1 \leq i \leq n, k \in \mathbb{Z} \setminus \{0\},$$

then

$$\begin{aligned} \frac{1}{c_0} t^{k\bar{m}} k_0 (w_1(-\bar{m}), w_2(-\bar{m}), \dots, w_n(-\bar{m})) &= (w_1, \dots, w_n) D^{-1} B_{-\bar{m}, \bar{m}}^{(0)} D, \\ t^{\bar{m}} k_p (w_1, w_2, \dots, w_n) &= (w_1(\bar{m}), \dots, w_n(\bar{m})) D^{-1} C_{\bar{m}, \bar{0}}^{(p)} D. \end{aligned}$$

So we can assume that $B_{-\bar{m}, \bar{m}}^{(0)}$, $C_{\bar{m}, \bar{0}}^{(p)}$, and $C_{\bar{m}, -\bar{m}}^{(p)}$, for $1 \leq p \leq \nu$ are all invertible upper triangular matrices. Furthermore, because

$$\left(t^{\bar{m}} k_p \frac{1}{c_0} t^{-\bar{m}} k_0 - \lambda \text{id} \right)^l (v_1, v_2, \dots, v_n) = (v_1, v_2, \dots, v_n) (C_{\bar{m}, -\bar{m}}^{(p)} - \lambda \text{id})^l,$$

the argument used in the proof of the claim shows that $C_{\bar{m}, -\bar{m}}^{(p)}$ also does not have distinct eigenvalues. For $1 \leq p \leq N$, set

$$B_{\bar{m}, -\bar{m}}^{(p)} = \frac{1}{c_p} C_{\bar{m}, -\bar{m}}^{(p)}$$

and for $0 \leq p \leq N$ denote by λ_p the eigenvalue of $B_{\bar{m}, -\bar{m}}^{(p)}$.

Let $A_{k\bar{m}, \bar{0}}^{(a)}$ and $A_{k_1\bar{m}, k_2\bar{m}}^{(a)}$, for $0 \leq a \leq \nu$ and $k, k_1, k_2 \in \mathbb{Z} \setminus \{0\}$, be such that

$$\begin{aligned} t^{k\bar{m}} d_a(v_1, v_2, \dots, v_n) &= (v_1(k\bar{m}), v_2(k\bar{m}), \dots, v_n(k\bar{m})) A_{k\bar{m}, \bar{0}}^{(a)}, \\ t^{k_1\bar{m}} d_a(v_1(k_2\bar{m}), v_2(k_2\bar{m}), \dots, v_n(k_2\bar{m})) \\ &= (v_1(k_1\bar{m} + k_2\bar{m}), \dots, v_n(k_1\bar{m} + k_2\bar{m})) A_{k_1\bar{m}, k_2\bar{m}}^{(a)}. \end{aligned}$$

Case 1: $\nu > 1$. Since $t^{\bar{m}} k_0 = t_0^{m_0} t^{\bar{m}} k_0 \neq 0$, it follows that there exists $1 \leq a \leq \nu$ such that $m_a \neq 0$, where $\underline{m} = (m_1, m_2, \dots, m_\nu)$. Let $b \in \{1, \dots, \nu\}$ be such that $a \neq b$. Consider

$$(3-4) \quad [t^{-\bar{m}} d_a, \frac{1}{c_0} t^{\bar{m}} k_0] = m_a \frac{1}{c_0} k_0, \quad [t^{-\bar{m}} d_a, t^{\bar{m}} k_b] = m_a k_b.$$

Case 1.1: There exists $b \in \{0, 1, \dots, \nu\}$ such that $b \neq 0, a$ and $c_b = 0$. Then

$$A_{-\bar{m}, \bar{m}}^{(a)} = B_{\bar{m}, -\bar{m}}^{(0)} A_{-\bar{m}, \bar{0}}^{(a)} + m_a I, \quad A_{-\bar{m}, \bar{m}}^{(a)} C_{\bar{m}, \bar{0}}^{(b)} = C_{\bar{m}, -\bar{m}}^{(b)} A_{-\bar{m}, \bar{0}}^{(a)}.$$

By (3-2) and (3-3),

$$A_{-\bar{m},\bar{0}}^{(a)} + m_a B_{\bar{m},-\bar{m}}^{(0)}{}^{-1} = C_{\bar{m},\bar{0}}^{(b)} A_{-\bar{m},\bar{0}}^{(a)} C_{\bar{m},\bar{0}}^{(b)}{}^{-1}.$$

But the sum on the left-hand side cannot be similar to $A_{-\bar{m},\bar{0}}^{(a)}$, since $m_a \neq 0$ and $B_{\bar{m},-\bar{m}}^{(0)}{}^{-1}$ is an invertible upper triangular matrix and does not have different eigenvalues. Thus this case is excluded.

Case 1.2: $c_b \neq 0$ for all $b \in \{0, 1, \dots, \nu\}$, $b \neq 0, a$. By (3-4) and (3-2), we have

$$\begin{aligned} B_{\bar{m},-\bar{m}}^{(0)} A_{-\bar{m},\bar{0}}^{(a)} B_{\bar{m},-\bar{m}}^{(0)}{}^{-1} + m_a B_{\bar{m},-\bar{m}}^{(0)}{}^{-1} - m_a B_{\bar{m},-\bar{m}}^{(b)}{}^{-1} \\ = B_{\bar{m},-\bar{m}}^{(0)} C_{\bar{m},\bar{0}}^{(b)} A_{-\bar{m},\bar{0}}^{(a)} C_{\bar{m},\bar{0}}^{(b)}{}^{-1} B_{\bar{m},-\bar{m}}^{(0)}{}^{-1}. \end{aligned}$$

(I) There exists $b \neq 0$ and a such that $\lambda_0 \neq \lambda_b$. Then $m_a B_{\bar{m},-\bar{m}}^{(0)}{}^{-1} - m_a B_{\bar{m},-\bar{m}}^{(b)}{}^{-1}$ is an invertible upper triangular matrix and does not have different eigenvalues. As in case 1.1, we deduce a contradiction.

(II) $\lambda_0 = \lambda_b$ for all $b \in \{1, \dots, \nu\}$ distinct from a .

(II.1) Suppose first that $c_a = 0$ (in this case $N = \nu - 1$, $a = \nu$) or $c_a \neq 0$ and $\lambda_a = \lambda_0$ (in this case $N = \nu$). Since $\sum_{p=0}^{\nu} m_p t^{\bar{m}} k_p = 0$, we have

$$\sum_{p=0}^{\nu} m_p t^{\bar{m}} k_p \frac{1}{c_0} t^{-\bar{m}} k_0 = 0.$$

So $\sum_{p=0}^{\nu} m_p C_{\bar{m},-\bar{m}}^{(p)} = 0$, and therefore

$$\sum_{p=0}^{\nu} m_p c_p = 0,$$

which contradicts the assumption that c_0, \dots, c_N are \mathbb{Z} -linearly independent.

(II.2) Now suppose $c_a \neq 0$, $\lambda_a \neq \lambda_0$ and there exists $b \neq 0$ and a such that $m_b \neq 0$. We deduce a contradiction as in case 1.2(I) by interchanging a by b .

(II.3) Suppose $c_a \neq 0$, $\lambda_a \neq \lambda_0$ and $m_b = 0$ for all $b \in \{1, \dots, \nu\}$ distinct from a . Then $m_0 c_0 \lambda_0 + m_a c_a \lambda_a = 0$. The proof of this case is the same as in case 2.2 below.

Case 2.: $\nu = 1$. In this case $a = 1$.

Case 2.1: $c_a = 0$. Since $[t^{-\bar{m}} d_0, t^{\bar{m}} k_0] = [t^{-\bar{m}} k_0, t^{\bar{m}} d_0] = 0$, we have

$$A_{-\bar{m},\bar{m}}^{(0)} = B_{\bar{m},-\bar{m}}^{(0)} A_{-\bar{m},\bar{0}}^{(0)}, \quad A_{\bar{m},-\bar{m}}^{(0)} = B_{-\bar{m},\bar{m}}^{(0)} A_{\bar{m},\bar{0}}^{(0)}.$$

Therefore

$$[t^{-\bar{m}} d_0, t^{\bar{m}} d_0](v_1, v_2, \dots, v_n) = (v_1, v_2, \dots, v_n) B_{-\bar{m},\bar{m}}^{(0)} [A_{-\bar{m},\bar{0}}^{(0)}, A_{\bar{m},\bar{0}}^{(0)}].$$

At the same time, we have

$$[t^{-\bar{m}}d_0, t^{\bar{m}}d_0] = 2m_0d_0 + m_0^2(-\mu_1 + \mu_2)(m_0k_0 + m_1k_1),$$

where $\tau = \mu_1\tau_1 + \mu_2\tau_2$ as above. So

$$(3-5) \quad B_{-\bar{m}, \bar{m}}^{(0)}[A_{-\bar{m}, \bar{0}}^{(0)}, A_{\bar{m}, \bar{0}}^{(0)}] = (2m_0(\gamma_0(d_0) + r_0) + m_0^2(-\mu_1 + \mu_2)(m_0c_0 + m_1c_1))I,$$

where γ_0 is the weight fixed above. Since γ_0 is arbitrary, we can choose it such that

$$2m_0(\gamma_0(d_0) + r_0) + m_0^2(-\mu_1 + \mu_2)(m_0c_0 + m_1c_1) \neq 0.$$

But $B_{-\bar{m}, \bar{m}}^{(0)}$ is an invertible triangular matrix and does not have different eigenvalues, in contradiction with (3-5).

Case 2.2: $c_a \neq 0$. Since

$$\begin{aligned} [t^{-\bar{m}}d_0, t^{\bar{m}}k_0] &= -m_1k_1, [t^{-\bar{m}}d_1, t^{\bar{m}}k_0] = m_1k_0 \text{ and} \\ [t^{\bar{m}}d_0, t^{-\bar{m}}k_0] &= m_1k_1, [t^{\bar{m}}d_1, t^{-\bar{m}}k_0] = -m_1k_0, \end{aligned}$$

we have

$$[k_0t^{-\bar{m}}d_0 + k_1t^{-\bar{m}}d_1, t^{\bar{m}}k_0] = [k_0t^{\bar{m}}d_0 + k_1t^{\bar{m}}d_1, t^{-\bar{m}}k_0] = 0.$$

Therefore

$$\begin{aligned} k_0A_{-\bar{m}, \bar{m}}^{(0)} + k_1A_{-\bar{m}, \bar{m}}^{(1)} &= B_{\bar{m}, -\bar{m}}^{(0)}(k_0A_{-\bar{m}, \bar{0}}^{(0)} + k_1A_{-\bar{m}, \bar{0}}^{(1)}), \\ k_0A_{\bar{m}, -\bar{m}}^{(0)} + k_1A_{\bar{m}, -\bar{m}}^{(1)} &= B_{-\bar{m}, \bar{m}}^{(0)}(k_0A_{\bar{m}, \bar{0}}^{(0)} + k_1A_{\bar{m}, \bar{0}}^{(1)}), \end{aligned}$$

and

$$\begin{aligned} [k_0t^{-\bar{m}}d_0 + k_1t^{-\bar{m}}d_1, k_0t^{\bar{m}}d_0 + k_1t^{\bar{m}}d_1](v_1, \dots, v_n) \\ = (v_1, \dots, v_n)B_{\bar{m}, -\bar{m}}^{(0)}[k_0A_{-\bar{m}, \bar{0}}^{(0)} + k_1A_{-\bar{m}, \bar{0}}^{(1)}, k_0A_{\bar{m}, \bar{0}}^{(0)} + k_1A_{\bar{m}, \bar{0}}^{(1)}]. \end{aligned}$$

At the same time, we have

$$\begin{aligned} [k_0t^{-\bar{m}}d_0 + k_1t^{-\bar{m}}d_1, k_0t^{\bar{m}}d_0 + k_1t^{\bar{m}}d_1] \\ = 2(m_0c_0 + m_1c_1)(c_0d_0 + c_1d_1) - (m_0c_0 + m_1c_1)^3(\mu_1 - \mu_2)\text{id}. \end{aligned}$$

Since c_0 and c_1 are \mathbb{Z} -linearly independent, we know that $m_0c_0 + m_1c_1 \neq 0$. As in case 2.1, we deduce a contradiction.

This concludes the first part of the proof. We next turn to the second major case, $N < p \leq v$.

If $N \geq 1$ or $N = 0$, we have $(m_1, \dots, m_v) \neq 0$, and the lemma follows from the first part and Lemma 3.3. Otherwise, let $t^{\bar{m}}k_p = t_0^{m_0}k_p$. Set $\mathcal{L}_0 = \bigoplus_{m_0 \in \mathbb{Z}} \mathbb{C}t_0^{m_0}d_0 \oplus \mathbb{C}k_0$ and $W = U(\mathcal{L}_0)v$, where $v \in V_{\bar{s}}$ is a homogeneous element. Since $c_0 \neq 0$, the sets $\{\dim W_{(n_0, 0) + \bar{s}} \mid n_0 \in \mathbb{Z}\}$ are not uniformly bounded. But if neither $t_0^{m_0}k_p$

nor $t_0^{-m_0}k_p$ is locally nilpotent, then t_0k_p and $t_0^{-1}k_p$ are not locally nilpotent. So by Lemmas 3.2 and 3.1, $\dim V_{(n_0,0)+\bar{s}} = \dim V_{\bar{s}}$ for all $n_0 \in \mathbb{Z}$, which is impossible since $\dim V_{(n_0,0)+\bar{s}} \geq \dim W_{(n_0,0)+\bar{s}}$. This proves Lemma 3.4 \square

For $0 \leq p \leq N$, consider the direct sum

$$\bigoplus_{m_p \in \mathbb{Z}} \mathbb{C}t_p^{m_p}d_p \oplus \mathbb{C}k_p,$$

which is a Virasoro Lie subalgebra of \mathcal{L} . Since $c_p \neq 0$, it follows from [Mathieu 1992] that there is a nonzero $v_p \in V_{\bar{r}}$ for some $\bar{r} \in \mathbb{Z}^{v+1}$ such that

$$(3-6) \quad t_p^{m_p}d_pv_p = 0 \quad \text{for all } m_p \in \mathbb{Z}_+$$

or

$$(3-7) \quad t_p^{m_p}d_pv_p = 0 \quad \text{for all } m_p \in \mathbb{Z}_-.$$

Lemma 3.5. *If $v_p \in V_{\bar{r}}$ satisfies (3-6), the sets*

$$\{t_p^{m_p}k_q \mid m_p \in \mathbb{Z}_+, q = 0, 1, 2, \dots, v, q \neq p\}$$

are all locally nilpotent on V . Likewise for (3-7), with \mathbb{Z}_+ replaced by \mathbb{Z}_- .

Proof. We only prove the first statement. Suppose it is false; then by Lemma 3.3 t_pk_q is not locally nilpotent on V for some $q \in \{0, 1, \dots, v\}$, $q \neq p$. By Lemma 3.4, $t_p^{-1}k_q$ is locally nilpotent. Therefore there exists $k \in \mathbb{Z}_+$ such that

$$(t_p^{-1}k_q)^{k-1}v_p \neq 0, \quad (t_p^{-1}k_q)^kv_p = 0.$$

So

$$\begin{aligned} t_p^2d_p(t_p^{-1}k_q)^kv_p &= -kt_pk_q(t_p^{-1}k_q)^{k-1}v_p + (t_p^{-1}k_q)^kt_p^2d_pv_p \\ &= -kt_pk_q(t_p^{-1}k_q)^{k-1}v_p = 0. \end{aligned}$$

This implies that t_pk_q is locally nilpotent, a contradiction. \square

Lemma 3.6. *If $v_p \in V_{\bar{r}}$ satisfies (3-6), the sets*

$$\{t^{\bar{m}}k_p \mid \bar{m} = (m_0, \dots, m_v) \in \mathbb{Z}^{v+1}, m_p \in \mathbb{Z}_+\}$$

are all locally nilpotent on V . Likewise for (3-7), with \mathbb{Z}_+ replaced by \mathbb{Z}_- .

Proof. Again we only prove the first statement. Without loss of generality, we assume that $p = 0$. Let \mathcal{H}' be the subspace of \mathcal{H} spanned by elements of \mathcal{H} which are locally nilpotent on V . If $t^{\underline{m}}k_0$, for any $\underline{m} \in \mathbb{Z}^v \setminus \{0\}$, is not locally nilpotent on V , the lemma holds thanks to Lemmas 3.3 and 3.5. Suppose $\mathcal{H}' \cap \{t^{\underline{m}}k_0 \mid \underline{m} \in \mathbb{Z}^v\} \neq \{0\}$. By Lemmas 3.2, 3.3 and 3.5, if $t^{\underline{m}}k_0 \in \mathcal{H}'$, then $t^{-\underline{m}}k_0 \notin \mathcal{H}'$, and $t_0^{m_0}t^{\underline{m}}k_0 \in \mathcal{H}'$ for all $m_0 > 0$.

Case 1: Suppose $t_0^{m_0}t^{-\underline{m}}k_0 \in \mathcal{H}'$ for any $t^{\underline{m}}k_0 \in \mathcal{H}'$. Then the lemma is proved.

Case 2: Suppose there exists $0 \neq t^{\underline{m}}k_0 \in \mathcal{H}'$ such that $t_0 t^{-\underline{m}}k_0 \notin \mathcal{H}'$. Since $\underline{m} = (m_1, \dots, m_v) \neq 0$, we can assume that $m_a \neq 0$ for some $a \in \{1, 2, \dots, v\}$. Let $V_{\bar{r}_0}$ be such that

$$\dim V_{\bar{r}_0} = \min\{\dim V_{\bar{s}} \mid V_{\bar{s}} \neq 0, \bar{s} \in \mathbb{Z}^{v+1}\}.$$

Case 2.1: Assume $t_0^i t^{-\underline{m}}k_0 \notin \mathcal{H}'$ for any $i > 0$. Let $l \in \mathbb{Z}_+$ and consider

$$(3-8) \quad \sum_{i=0}^l a_i t_0^{-i} t^{-\underline{m}}k_0 t_0^i t^{-\underline{m}}k_0 v = 0,$$

where $v \in V_{\bar{r}_0} \setminus \{0\}$. By Lemma 3.4, $\{t_0^i t^{\underline{m}}k_0, t_0^{-i} t^{\underline{m}}k_0 \mid i \in \mathbb{Z}_+\} \subseteq \mathcal{H}'$. So by Lemma 3.2, we have

$$t_0^i t^{\underline{m}}k_0 V_{\bar{r}_0} = t_0^{-i} t^{\underline{m}}k_0 V_{\bar{r}_0} = t_0^i t^{\underline{m}}d_p V_{\bar{r}_0} = t_0^{-i} t^{\underline{m}}d_p V_{\bar{r}_0} = 0, \quad i \in \mathbb{Z}_+, 0 \leq p \leq v.$$

Let $j \in \{0, 1, \dots, l\}$. From (3-8) we have

$$t_0^{-j} t^{\underline{m}}d_a t_0^j t^{\underline{m}}d_a \left(\sum_{i=0}^l a_i t_0^{-i} t^{-\underline{m}}k_0 t_0^i t^{-\underline{m}}k_0 \right) v = 0.$$

Therefore

$$\sum_{i=0}^l a_i (-m_a) t_0^{j-i} k_0 (-m_a) t_0^{i-j} k_0 v = a_j m_a^2 c_0^2 v = 0.$$

So $a_j = 0$, $j = 0, 1, \dots, l$. This means $\{t_0^{-i} t^{-\underline{m}}k_0 t_0^i t^{-\underline{m}}k_0 v \mid 0 \leq i \leq l\}$ are linearly independent. Since l can be any positive integer, it follows that $V_{\bar{r}_0 - (0, 2\underline{m})}$ is infinite-dimensional, a contradiction.

Case 2.2: Assume there exists $l \in \mathbb{Z}_+$ such that

$$t_0^{l-1} t^{-\underline{m}}k_0 \notin \mathcal{H}', \quad t_0^l t^{-\underline{m}}k_0 \in \mathcal{H}'.$$

(I) Assume that $t_0^l t^{-i\underline{m}}k_0 \in \mathcal{H}'$ for any $i \in \mathbb{Z}_+$. Let $s > 0$ and consider

$$\sum_{i=1}^s a_i t_0^{-l} t^{i\underline{m}}k_0 t^{-i\underline{m}}k_0 v = 0.$$

Similar to the proof above, we can deduce that $V_{\bar{r}_0 - (l, 0)}$ is infinite-dimensional, in contradiction with the assumption that V has finite-dimensional weight spaces.

(II) Assume there exists $s_1 \in \mathbb{Z}_+$ such that

$$t_0^l t^{-\underline{m}}k_0 \in \mathcal{H}', \quad t_0^l t^{-2\underline{m}}k_0 \in \mathcal{H}', \quad \dots, \quad t_0^l t^{-s_1 \underline{m}}k_0 \in \mathcal{H}', \quad t_0^l t^{-(s_1+1)\underline{m}}k_0 \notin \mathcal{H}'.$$

Then there exist $s_2, s_3, \dots, s_k, \dots$ such that $s_i \geq s_1$ for $i = 2, 3, \dots, k, \dots$ and

$$t_0^{il} t^{(-s_1-s_2-\dots-s_{i-1}-1)\underline{m}}k_0 \in \mathcal{H}', \quad t_0^{il} t^{(-s_1-s_2-\dots-s_{i-1}-2)\underline{m}}k_0 \in \mathcal{H}', \dots,$$

$$t_0^{il} t^{(-s_1-s_2-\dots-s_{i-1}-s_i)\underline{m}} k_0 \in \mathcal{H}', \quad t_0^{il} t^{(-s_1-s_2-\dots-s_{i-1}-s_i-1)\underline{m}} k_0 \notin \mathcal{H}'.$$

Assume that

$$\begin{aligned} & \left(\sum_{i=1}^{s_1} a_i t_0^{-l} t^{i\underline{m}} k_0 t^{-i\underline{m}} k_0 + \sum_{i=1}^{s_2} a_{s_1+i} t_0^{-2l} t^{(s_1+i)\underline{m}} k_0 t_0^l t^{-(s_1+i)\underline{m}} k_0 \right. \\ & \quad + \sum_{i=1}^{s_3} a_{s_1+s_2+i} t_0^{-3l} t^{(s_1+s_2+i)\underline{m}} k_0 t_0^{2l} t^{-(s_1+s_2+i)\underline{m}} k_0 + \dots \\ & \quad \left. + \sum_{i=1}^{s_k} a_{s_1+\dots+s_{k-1}+i} t_0^{-kl} t^{(s_1+\dots+s_{k-1}+i)\underline{m}} k_0 t_0^{(k-1)l} t^{-(s_1+\dots+s_{k-1}+i)\underline{m}} k_0 \right) v = 0. \end{aligned}$$

Let

$$\begin{aligned} & t^{j\underline{m}} d_a t_0^l t^{-j\underline{m}} d_a, & 1 \leq j \leq s_1, \\ & t_0^{-l} t^{(s_1+j)\underline{m}} d_a t_0^{2l} t^{-(s_1+j)\underline{m}} d_a, & 1 \leq j \leq s_2, \\ & \dots, \\ & t_0^{-(k-1)l} t^{(s_1+s_2+\dots+s_{k-1}+j)\underline{m}} d_a t_0^{kl} t^{-(s_1+s_2+\dots+s_{k-1}+j)\underline{m}} d_a, & 1 \leq j \leq s_k \end{aligned}$$

act on the two sides of the above equation respectively. By Lemma 3.4, we deduce that $a_i = 0$, for $i = 1, 2, \dots, s_1$, and that

$$a_{s_1+\dots+s_{j-1}+i} = 0 \quad \text{for } i = 1, 2, \dots, s_j, \quad 2 \leq j \leq k.$$

Since k can be any positive integer, it follows that $V_{\vec{r}_0-(l, \underline{0})}$ is infinite-dimensional, which contradicts our assumption. The lemma is proved. \square

Lemmas 3.1 through 3.6 immediately yield the following result.

Theorem 3.7. *Let V be an irreducible weight module of \mathcal{L} such that c_0, \dots, c_N are \mathbb{Z} -linearly independent and $N \geq 1$. Then V has weight spaces that are infinite-dimensional.*

Let

$$\begin{aligned} \mathcal{L}_+ &= \sum_{p=0}^v t_0 \mathbb{C}[t_0, t_1^{\pm 1}, \dots, t_v^{\pm 1}] k_p \oplus \sum_{p=0}^v t_0 \mathbb{C}[t_0, t_1^{\pm 1}, \dots, t_v^{\pm 1}] d_p, \\ \mathcal{L}_- &= \sum_{p=0}^v t_0^{-1} \mathbb{C}[t_0^{-1}, t_1^{\pm 1}, \dots, t_v^{\pm 1}] k_p \oplus \sum_{p=0}^v t_0^{-1} \mathbb{C}[t_0^{-1}, t_1^{\pm 1}, \dots, t_v^{\pm 1}] d_p, \\ \mathcal{L}_0 &= \sum_{p=0}^v \mathbb{C}[t_1^{\pm 1}, \dots, t_v^{\pm 1}] k_p \oplus \sum_{p=0}^v \mathbb{C}[t_1^{\pm 1}, \dots, t_v^{\pm 1}] d_p. \end{aligned}$$

Then

$$\mathcal{L} = \mathcal{L}_+ \oplus \mathcal{L}_0 \oplus \mathcal{L}_-.$$

Definition 3.8. Let W be a weight module of \mathcal{L} . If there is a nonzero vector $v_0 \in W$ such that

$$\mathcal{L}_+ v_0 = 0, \quad W = U(\mathcal{L})v_0,$$

then W is called a highest weight module of \mathcal{L} . If there is a nonzero vector $v_0 \in W$ such that

$$\mathcal{L}_- v_0 = 0, \quad W = U(\mathcal{L})v_0,$$

then W is called a lowest weight module of \mathcal{L} .

From Lemmas 3.2 and 3.6, we obtain:

Theorem 3.9. *Let V be an irreducible weight module of \mathcal{L} with finite-dimensional weight spaces and with central charges $c_0 \neq 0, c_1 = c_2 = \dots = c_v = 0$. Then V is a highest or lowest weight module of \mathcal{L} .*

In the remainder of this section we assume that V is an irreducible weight module of \mathcal{L} with finite-dimensional weight spaces and with central charges $c_0 \neq 0, c_1 = \dots = c_v = 0$.

Set

$$T = \begin{cases} \{v \in V \mid \mathcal{L}_+ v = 0\} & \text{if } V \text{ is a highest weight module of } \mathcal{L}, \\ \{v \in V \mid \mathcal{L}_- v = 0\} & \text{if } V \text{ is a lowest weight module of } \mathcal{L}. \end{cases}$$

Then T is a \mathcal{L}_0 -module and

$$V = U(\mathcal{L}_-)T \quad \text{or} \quad V = U(\mathcal{L}_+)T.$$

Since V is an irreducible \mathcal{L} -module, T is an irreducible \mathcal{L}_0 -module. T has the decomposition

$$T = \bigoplus_{\underline{m} \in \mathbb{Z}^v} T_{\underline{m}},$$

where $\underline{m} = (m_1, m_2, \dots, m_v)$, $T_{\underline{m}} = \{v \in T \mid d_i v = (m_i + \mu(d_i))v, 1 \leq i \leq v\}$ and μ is a fixed weight of T . As in the proof in [Jiang and Meng 2003; Eswara Rao and Jiang 2005], we can deduce:

Theorem 3.10. (1) *For all $\underline{m}, \underline{n} \in \mathbb{Z}^v$, $p = 1, 2, \dots, v$, we have*

$$\dim T_{\underline{m}} = \dim T_{\underline{n}}, \quad t^{\underline{m}} k_p \cdot T = 0,$$

$$t^{\underline{m}} k_0(v_1(\underline{n}), \dots, v_m(\underline{n})) = c_0(v_1(\underline{m} + \underline{n}), v_2(\underline{m} + \underline{n}), \dots, v_n(\underline{m} + \underline{n})),$$

$$t^{\underline{m}} d_0(v_1(\underline{n}), v_2(\underline{n}), \dots, v_n(\underline{n})) = \mu(d_0)(v_1(\underline{m} + \underline{n}), v_2(\underline{m} + \underline{n}), \dots, v_n(\underline{m} + \underline{n})),$$

where $\{v_1(\underline{0}), \dots, v_m(\underline{0})\}$ is a basis of $T_{\underline{0}}$ and $v_i(\underline{m}) = \frac{1}{c_0} t^{\underline{m}} k_0 v_i(\underline{0})$, for $i = 1, 2, \dots, m$.

(2) As an $(\mathcal{A}_v \oplus \mathcal{D}_v)$ -module, T is isomorphic to

$$F^\alpha(\psi, b) = V(\psi, b) \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_v^{\pm 1}]$$

for some $\alpha = (\alpha_1, \dots, \alpha_v)$, ψ , and b , where $\mathcal{A}_v = \mathbb{C}[t_1^{\pm 1}, \dots, t_v^{\pm 1}]$, \mathcal{D}_v is the derivation algebra of \mathcal{A}_v , and $V(\psi, b)$ is an m -dimensional, irreducible $\mathfrak{gl}_v(\mathbb{C})$ -module satisfying $\psi(I) = b \text{id}_{V(\psi, b)}$ and

$$t^r d_p(w \otimes t^m) = (m_p + \alpha_p)w \otimes t^{r+m} + \sum_{i=1}^v r_i \psi(E_{ip})w \otimes t^{r+m}$$

for $w \in V(\psi, b)$.

Let

$$M = \text{Ind}_{\mathcal{L}_+ + \mathcal{L}_0}^{\mathcal{L}} T \quad \text{or} \quad M = \text{Ind}_{\mathcal{L}_- + \mathcal{L}_0}^{\mathcal{L}} T.$$

Theorem 3.11. *Among the submodules of M intersecting T trivially, there is a maximal one, which we denote by M^{rad} . Moreover $V \cong M/M^{\text{rad}}$.*

4. The structure of V with $c_0 = \dots = c_v = 0$

Assume that V is an irreducible weight module of \mathcal{L} with finite-dimensional weight spaces and $c_0 = \dots = c_v = 0$.

Lemma 4.1. *For any $t^{\bar{r}} k_p \in \mathcal{K}$, $t^{\bar{r}} k_p$ or $t^{-\bar{r}} k_p$ is locally nilpotent on V .*

Lemma 4.2. *If V is uniformly bounded, $t^{\bar{r}} k_p$ is locally nilpotent on V for any $t^{\bar{r}} k_p \in \mathcal{K}$.*

Proof. For $t^{\bar{r}} k_p \in \mathcal{K}$, by Lemma 4.1, $t^{\bar{r}} k_p$ or $t^{-\bar{r}} k_p$ is nilpotent on $V_{\bar{m}}$ for all $\bar{m} \in \mathbb{Z}^{v+1}$. Since V is uniformly bounded, i.e., $\max\{\dim V_{\bar{m}} \mid \bar{m} \in \mathbb{Z}^{v+1}\} < \infty$, there exists $N \in \mathbb{Z}_+$ such that

$$(t^{\bar{r}} k_p t^{-\bar{r}} k_p)^N V = 0, (t^{\bar{r}} k_p t^{-\bar{r}} k_p)^{N-1} V \neq 0$$

If the lemma is false, we can assume that $t^{-\bar{r}} k_p$ is not locally nilpotent on V . Therefore for any $0 \neq v \in V$, we have $t^{-\bar{r}} k_p v \neq 0$. So

$$(t^{\bar{r}} k_p)^N V = 0.$$

Let $t^{-2\bar{r}} d_q \in \mathcal{K}$ be such that $p \neq q$ and $r_q \neq 0$. By the fact that $[t^{-2\bar{r}} d_q, t^{\bar{r}} k_p] = r_q t^{-\bar{r}} k_p$, we deduce that $t^{-\bar{r}} k_p (t^{\bar{r}} k_p)^{N-1} V = 0$, a contradiction. \square

Lemma 4.3. *If there exists $0 \neq v \in V$ such that $t^{\bar{m}} k_p v = 0$ for all $\bar{m} \in \mathbb{Z}^{v+1}$ and $0 \leq p \leq v$. Then $\mathcal{K}(V) = 0$.*

Proof. This follows from (2-2), since \mathcal{K} is commutative and V is an irreducible \mathcal{L} -module. \square

Theorem 4.4. *If V is uniformly bounded, $t^{\bar{r}} k_p V$ vanishes for any $t^{\bar{r}} k_p \in \mathcal{K}$.*

Proof. Let $0 \neq t_i k_p \in \mathcal{H}$. If $t_i k_p V = 0$, it is easy to prove that $\mathcal{H}(V) = 0$. If $t_i k_p V \neq 0$. Since V is uniformly bounded, by Lemma 4.2, there exists $l \in \mathbb{Z}_+$ such that

$$(4-1) \quad (t_i k_p t_i^{-1} k_p)^l V = 0, \quad (t_1 k_p t_1^{-1} k_p)^{l-1} V \neq 0.$$

If there exists $s \in \mathbb{Z}_+$ such that $(t_i^{-1} k_p)^s V = 0$, $(t_i^{-1} k_p)^{s-1} V \neq 0$. By the fact that $[t^{\bar{m}} d_i, t_i^{-1} k_p] = -t_i^{-1} t^{\bar{m}} k_p$ and $[t^{\bar{m}} d_p, t_i^{-1} k_p] = t_i^{-1} t^{\bar{m}} k_i$, we have

$$t^{\bar{r}} k_p (t_i^{-1} k_p)^{s-1} V = t^{\bar{r}} k_i (t_i^{-1} k_p)^{s-1} V = 0 \quad \text{for all } \bar{r} \in \mathbb{Z}^{\nu+1}.$$

If $(t_i^{-1} k_p)^s V \neq 0$ for all $s \in \mathbb{Z}_+$. Then by (4-1) there is $r \geq 0$ such that $(t_i k_p)^{l-i} (t_i^{-1} k_p)^{l+i} V = 0$ for all $0 \leq i \leq r$, and $(t_i k_p)^{l-r-1} (t_i^{-1} k_p)^{l+r+1} V \neq 0$. So for any $\bar{m} \in \mathbb{Z}^{\nu+1}$, we have

$$t^{-\bar{m}} d_i (t_i k_p)^{l-r} (t_i^{-1} k_p)^{l+r+1} V = 0, \quad t^{-\bar{m}} d_p (t_i k_p)^{l-r} (t_i^{-1} k_p)^{l+r+1} V = 0.$$

Therefore

$$\begin{aligned} t^{\bar{r}} k_p (t_i k_p)^{l-r-1} (t_i^{-1} k_p)^{l+r+1} V &= 0, \\ t^{\bar{r}} k_i (t_i k_p)^{l-r-1} (t_i^{-1} k_p)^{l+r+1} V &= 0, \end{aligned}$$

for all $\bar{r} \in \mathbb{Z}^{\nu+1}$.

Case 1: $\nu \in 2\mathbb{Z}_+ + 1$. By the preceding discussion, there exist nonnegative integers l_i and r_i , for $i = 0, 2, 4, \dots, \nu - 1$, such that

$$(t_\nu k_{\nu-1})^{l_{\nu-1}} (t_\nu^{-1} k_{\nu-1})^{r_{\nu-1}} (t_{\nu-2} k_{\nu-3})^{l_{\nu-3}} (t_{\nu-2}^{-1} k_{\nu-3})^{r_{\nu-3}} \dots (t_1 k_0)^{l_0} (t_1^{-1} k_0)^{r_0} V \neq 0$$

and

$$t^{\bar{m}} k_p (t_\nu k_{\nu-1})^{l_{\nu-1}} (t_\nu^{-1} k_{\nu-1})^{r_{\nu-1}} (t_{\nu-2} k_{\nu-3})^{l_{\nu-3}} (t_{\nu-2}^{-1} k_{\nu-3})^{r_{\nu-3}} \dots (t_1 k_0)^{l_0} (t_1^{-1} k_0)^{r_0} V$$

vanishes for all $0 \leq p \leq \nu$ and $\bar{m} \in \mathbb{Z}^{\nu+1}$. By Lemma 4.3, the conclusion of the theorem holds.

Case 2: $\nu \in 2\mathbb{Z}$. Then there exist nonnegative integers l_i and r_i , for $i = 0, 2, 4, \dots, \nu - 2$, such that

$$W = (t_{\nu-1} k_{\nu-2})^{l_{\nu-2}} (t_{\nu-1}^{-1} k_{\nu-2})^{r_{\nu-2}} (t_{\nu-3} k_{\nu-4})^{l_{\nu-4}} (t_{\nu-3}^{-1} k_{\nu-4})^{r_{\nu-4}} \dots (t_1 k_0)^{l_0} (t_1^{-1} k_0)^{r_0} V$$

is nonzero and

$$(4-2) \quad t^{\bar{m}} k_p W = 0$$

for all $0 \leq p \leq \nu - 1$ and $\bar{m} \in \mathbb{Z}^{\nu+1}$. By (2-1), we know that

$$(4-3) \quad t^{\bar{m}} k_\nu W = 0,$$

for $\bar{m} \in \mathbb{Z}^{v+1}$ such that $m_v \neq 0$. If there exists $t^{\bar{r}_0}k_v$ satisfying $t^{\bar{r}_0}k_v W \neq 0$, let

$$\begin{aligned}\mathcal{L}_v &= \text{span} \{t^{\underline{m}}d_i, t^{\bar{m}}d_v, t^{\underline{m}}k_v \mid t^{\underline{m}} = t_0^{m_0}t_1^{m_1} \cdots t_{v-1}^{m_{v-1}}, 0 \leq i \leq v-1, \\ \underline{m} &= (m_0, \dots, m_{v-1}) \in \mathbb{Z}^v, \bar{m} \in \mathbb{Z}^{v+1}\}, \\ W' &= U(\mathcal{L}_v)W.\end{aligned}$$

Then $W' \neq 0$ and

$$t^{\bar{m}}k_p W' = 0, \quad t^{\bar{n}}k_v W' = 0,$$

for all $0 \leq p \leq v-1$, $\bar{m} \in \mathbb{Z}^{v+1}$, and $\bar{n} \in \mathbb{Z}^{v+1}$ such that $n_v \neq 0$. If there exists $0 \neq t^{\underline{m}}k_v$ such that $t^{\underline{m}}k_v W' \neq 0$, we have

$$(t^{-\underline{m}}k_v)^l (t^{\underline{m}}k_v)^l W' = 0 \quad \text{and} \quad (t^{-\underline{m}}k_v)^{l-1} (t^{\underline{m}}k_v)^{l-1} W' \neq 0$$

for some $l \in \mathbb{Z}_+$. As in the preceding proof, we can deduce that there exists a nonzero $v \in W'$ such that

$$t^{\underline{n}}k_v v = 0$$

for all $\underline{n} \in \mathbb{Z}^v$. Therefore

$$t^{\bar{m}}k_p v = 0$$

for all $\bar{m} \in \mathbb{Z}^{v+1}$ and $0 \leq p \leq v$. We have proved that $\mathcal{K}(V) = 0$. \square

References

- [Belavin et al. 1984] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, “Infinite conformal symmetry in two-dimensional quantum field theory”, *Nuclear Phys. B* **241**:2 (1984), 333–380. MR 86m:81097 Zbl 0661.17013
- [Berman and Billig 1999] S. Berman and Y. Billig, “Irreducible representations for toroidal Lie algebras”, *J. Algebra* **221**:1 (1999), 188–231. MR 2000k:17004 Zbl 0942.17016
- [Billig 2003] Y. Billig, “Abelian extensions of the group of diffeomorphisms of a torus”, *Lett. Math. Phys.* **64**:2 (2003), 155–169. MR 2004h:22012 Zbl 1079.58004
- [Chari and Pressley 1988] V. Chari and A. Pressley, “Unitary representations of the Virasoro algebra and a conjecture of Kac”, *Compositio Math.* **67**:3 (1988), 315–342. MR 89h:17025 Zbl 0661.17022
- [Di Francesco et al. 1997] P. Di Francesco, P. Mathieu, and D. Sénéchal, *Conformal field theory*, Springer, New York, 1997. MR 97g:81062 Zbl 0869.53052
- [Dotsenko and Fateev 1984] V. S. Dotsenko and V. A. Fateev, “Conformal algebra and multipoint correlation functions in 2D statistical models”, *Nuclear Phys. B* **240**:3 (1984), 312–348. MR 85i:82061
- [Eswara Rao 2004] S. Eswara Rao, “Partial classification of modules for Lie algebra of diffeomorphisms of d -dimensional torus”, *J. Math. Phys.* **45**:8 (2004), 3322–3333. MR 2005d:17028 Zbl 1071.17020
- [Eswara Rao and Jiang 2005] S. Eswara Rao and C. Jiang, “Classification of irreducible integrable representations for the full toroidal Lie algebras”, *J. Pure Appl. Algebra* **200**:1-2 (2005), 71–85. MR 2006b:17038 Zbl 1070.17009

- [Eswara Rao and Moody 1994] S. Eswara Rao and R. V. Moody, “Vertex representations for n -toroidal Lie algebras and a generalization of the Virasoro algebra”, *Comm. Math. Phys.* **159**:2 (1994), 239–264. MR 94m:17028 Zbl 0808.17018
- [Jiang and Meng 2003] C. Jiang and D. Meng, “Integrable representations for generalized Virasoro-toroidal Lie algebras”, *J. Algebra* **270**:1 (2003), 307–334. MR 2005b:17053 Zbl 1037.17029
- [Kac 1982] V. G. Kac, “Some problems on infinite-dimensional Lie algebras and their representations”, pp. 117–126 in *Lie algebras and related topics* (New Brunswick, NJ, 1981), edited by D. J. Winter, Lecture Notes in Math. **933**, Springer, Berlin, 1982. MR 84e:17010 Zbl 0493.17011
- [Kaplansky and Santharoubane 1985] I. Kaplansky and L. J. Santharoubane, “Harish-Chandra modules over the Virasoro algebra”, pp. 217–231 in *Infinite dimensional groups with applications* (Berkeley, 1984), edited by V. Kac, Math. Sci. Res. Inst. Publ. **4**, Springer, New York, 1985. MR 87d:17013 Zbl 0589.17013
- [Larsson 1992] T. A. Larsson, “Conformal fields: a class of representations of $\text{Vect}(N)$ ”, *Internat. J. Modern Phys. A* **7**:26 (1992), 6493–6508. MR 93h:17053 Zbl 0972.17502
- [Mathieu 1992] O. Mathieu, “Classification of Harish-Chandra modules over the Virasoro Lie algebra”, *Invent. Math.* **107**:2 (1992), 225–234. MR 93d:17034 Zbl 0779.17025
- [Ramos et al. 1990] E. Ramos, C.-H. Sah, and R. E. Shrock, “Algebras of diffeomorphisms of the N -torus”, *J. Math. Phys.* **31**:8 (1990), 1805–1816. MR 91j:17043 Zbl 0733.17014

Received December 3, 2005.

CUIPO JIANG
DEPARTMENT OF MATHEMATICS
SHANGHAI JIAOTONG UNIVERSITY
SHANGHAI 200030
CHINA
cpjiang@sjtu.edu.cn

QIFEN JIANG
DEPARTMENT OF MATHEMATICS
SHANGHAI JIAOTONG UNIVERSITY
SHANGHAI 200030
CHINA
qfjiang@sjtu.edu.cn

