

*Pacific  
Journal of  
Mathematics*

**AN ELEMENTARY, EXPLICIT, PROOF OF THE EXISTENCE  
OF QUOT SCHEMES OF POINTS**

TROND STØLEN GUSTAVSEN, DAN LAKSOV AND ROY MIKAEL SKJELNES



## AN ELEMENTARY, EXPLICIT, PROOF OF THE EXISTENCE OF QUOT SCHEMES OF POINTS

TROND STØLEN GUSTAVSEN, DAN LAKSOV AND ROY MIKAEL SKJELNES

**We give an easy and elementary construction of quotient schemes of modules of relatively finite rank under very general conditions. The construction provides a natural, explicit description of an affine covering of the quotient schemes, that is useful in many situation.**

### Introduction

Quotient schemes were introduced by A. Grothendieck [1966], and belong among the fundamental tools in algebraic geometry. They generalize simultaneously the Hilbert schemes and the Grassmann schemes. In most cases where these schemes appear it suffices to know that they exist. However, there are cases when it is crucial to have an explicit description of the schemes. We give here, for any morphism of schemes  $\text{Proj}(\mathcal{R}) \rightarrow S$ , where  $S$  is arbitrary and  $\mathcal{R}$  is a quasicohherent graded  $\mathbb{O}_S$ -algebra, an elementary construction of quotient schemes parametrizing equivalence classes of surjections from a quasicohherent  $\mathbb{O}_{\text{Proj}(\mathcal{R})}$ -module to coherent modules that are of relatively finite rank over  $S$ . The construction provides a natural and explicit description of an affine covering of the quotient schemes. In a previous article [Gustavsen et al. 2007] we indicated the usefulness of such a description in the case of Hilbert schemes of points, and further evidence of this is given by M. Huibregtse [2002; 2006]. Our proof of the existence of the quotient schemes is a simplification and clarification of the constructions of these works.

The main new idea is the description of a local version of the quot functor. More precisely, let  $A$  be a ring,  $B$  an  $A$ -algebra and  $E$  and  $F$  modules over  $A$  with  $E$  free of finite rank, and fix an  $A$ -module homomorphism  $s : E \rightarrow B \otimes_A F$ . We parametrize  $B$ -module structures on  $E$ , together with  $B$ -module homomorphisms  $u : B \otimes_A F \rightarrow E$  such that  $us = \text{id}_E$ .

When  $S$  is locally noetherian and  $\mathcal{R}$  is locally finitely generated by elements of degree one, our existence result for the quotient schemes of modules of relatively finite rank follows from the more general results of Grothendieck [1966]. Apparently the first detailed existence proof of Grothendieck's result was provided by

---

*MSC2000:* 14D20, 14A15.

*Keywords:* Hilbert schemes, quotient schemes.

A. Altman and S. Kleiman [1980], valid under quite general conditions, where  $S$  is not assumed to be locally noetherian. A much used method for obtaining these results was given by D. Mumford [1966] (see also [Sernesi 2006]). In contrast to these approaches our method relies on simple algebraic constructions and avoids embeddings into high dimensional grassmannians via Castelnuovo–Mumford regularity. As a consequence our local description of the quotient schemes is, in most cases appearing in applications, in terms of explicit natural equations in the affine space of commuting matrices of smallest possible size, that is, the size is equal to the finite rank of the modules.

### 1. The local quot functor

**1.1.** Let  $A$  be a commutative ring with unit, and let  $A \rightarrow B$  be an  $A$ -algebra. Moreover, let  $E$  and  $F$  be  $A$ -modules where  $E$  is free of finite rank, and let

$$s : E \rightarrow B \otimes_A F$$

be an  $A$ -module homomorphism. We want to describe all  $B$ -module structures on  $E$ , together with  $B$ -module homomorphisms  $u : B \otimes_A F \rightarrow E$  such that  $us = \text{id}_E$ . Recall that two surjections are considered *equivalent* if their kernels coincide, and that a  $B$ -module structure on an  $A$ -module  $E$  corresponds to an  $A$ -algebra homomorphism  $B \rightarrow \text{End}_A(E)$ .

More precisely, we want to describe, for every  $A$ -algebra  $A \rightarrow A'$ , the set consisting of an  $A' \otimes_A B$ -module structure on  $A' \otimes_A E$  together with an  $A' \otimes_A B$ -module homomorphism  $A' \otimes_A B \otimes_A F \xrightarrow{u} A' \otimes_A E$  such that the composite  $A'$ -module homomorphism

$$A' \otimes_A E \xrightarrow{\text{id}_{A'} \otimes_A s} A' \otimes_A B \otimes_A F \xrightarrow{u} A' \otimes_A E$$

is the identity. This clearly defines a functor from  $A$ -algebras to sets.

The main objective of Sections 1, 2 and 3 is to show that this functor is representable by an  $A$ -algebra  $Q^s$ , and to give a simple explicit description of this algebra.

We first find a slightly different description of this functor.

**Notation 1.2.** We shall need *evaluation* and *trace* maps. We recall that when  $G$  and  $K$  are  $A$ -modules with  $K$  free of finite rank, we obtain for every submodule  $D$  of  $\text{Hom}_A(G, K)$  the *evaluation* homomorphisms

$$\text{ev} : D \otimes_A G \rightarrow K \quad \text{or} \quad \text{ev} : G \otimes_A D \rightarrow K$$

that maps  $x \otimes_A \varphi$ , or  $\varphi \otimes_A x$ , to  $\varphi(x)$ . Moreover, we have the *trace* homomorphism

$$\text{tr} : A \rightarrow K \otimes_A K^\vee$$

obtained from the dual homomorphism of  $\text{ev} : K^\vee \otimes_A K \rightarrow A$ .

**Proposition 1.3.** *There is a natural bijection between*

- (1) *the set of  $B$ -module structures on  $E$  provided with a surjective homomorphism  $u : B \otimes_A F \rightarrow E$  of  $B$ -modules, and*
- (2) *the set of  $A$ -algebra homomorphisms  $\varphi : B \rightarrow \text{End}_A(E)$  provided with an  $A$ -module homomorphism  $v : F \rightarrow E$  such that the composite map*

$$B \otimes_A F \xrightarrow{\varphi \otimes_A v} \text{End}_A(E) \otimes_A E \xrightarrow{\text{ev}} E$$

*is surjective.*

*The bijection maps a pair  $(v, \varphi)$  in (2) to  $\text{ev}(\varphi \otimes_A v)$  in (1), where  $E$  has the  $B$ -module structure given by  $\varphi$ .*

*Proof.* We first note that  $B$ -module structures on  $E$  correspond to  $A$ -algebra homomorphisms  $\varphi : B \rightarrow \text{End}(E)$ .

Given a  $B$ -module structure on  $E$ , a surjection  $u : B \otimes_A F \rightarrow E$  of  $B$ -modules determines a  $B$ -module structure on  $E$  uniquely. Moreover an  $A$ -module homomorphism  $u : B \otimes_A F \rightarrow E$  determines an  $A$ -module homomorphism  $v : F \rightarrow E$  by *restriction of scalars*, and conversely,  $u$  is determined by  $v$  by *extension of scalars*.

Finally, an  $A$ -module homomorphism  $v : F \rightarrow E$  and a  $B$ -module structure  $\varphi : B \rightarrow \text{End}_A(E)$  on  $E$  makes the composite map

$$B \otimes_A F \xrightarrow{\varphi \otimes_A v} \text{End}_A(E) \otimes_A E \xrightarrow{\text{ev}} E$$

into a  $B$ -module homomorphism, since  $\text{ev}$  is an  $\text{End}_A(E)$ -module homomorphism and  $\varphi : B \rightarrow \text{End}_A(E)$  is a ring homomorphism. □

**Corollary 1.4.** *The bijection of the proposition induces a bijection between*

- (1)  *$B$ -module structures on  $E$  provided with a homomorphism of  $B$ -modules  $u : B \otimes_A F \rightarrow E$  such that*

$$E \xrightarrow{s} B \otimes_A F \xrightarrow{u} E$$

*is the identity, and*

- (2)  *$A$ -algebra homomorphisms  $\varphi : B \rightarrow \text{End}_A(E)$  provided with an  $A$ -module homomorphism  $v : F \rightarrow E$  such that*

$$E \xrightarrow{s} B \otimes_A F \xrightarrow{\varphi \otimes_A v} \text{End}_A(E) \otimes_A E \xrightarrow{\text{ev}} E$$

*is the identity.*

*Proof.* This follows from the proposition since the surjectivity of  $u$  and  $\text{ev}(\varphi \otimes_A v)$  is automatic. □

## 2. The local Hilbert scheme

The material of this section will basically give a construction of Hilbert schemes of points, as in [Gustavsen et al. 2007], but the presentation here is different from that of that paper. We use that the  $A$ -algebra  $\text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee)$  parametrizes module homomorphisms  $u : G \rightarrow \text{End}_A(E)$ . Then we explicitly construct a residue algebra  $H$  of  $\text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee)$  that parametrizes those  $u$  such that the elements of the image commute. Then  $H$  also parametrizes  $A$ -algebra homomorphisms  $\text{Sym}_A(G) \rightarrow \text{End}_A(E)$ .

**Notation 2.1.** Let  $G$  be an  $A$ -module and  $K$  a free  $A$ -module of finite rank. We denote by

$$u_K : \text{Sym}_A(G \otimes_A K^\vee) \otimes_A G \rightarrow \text{Sym}_A(G \otimes_A K^\vee) \otimes_A K$$

the  $\text{Sym}_A(G \otimes_A K^\vee)$ -module homomorphism defined by  $u_K(1 \otimes_A x) = x \otimes_A \text{tr}(1_A)$  for all  $x \in G$ , where  $G \otimes_A K^\vee$  is considered as a submodule of  $\text{Sym}_A(G \otimes_A K^\vee)$ .

**Lemma 2.2.** *For every  $A$ -algebra homomorphism  $A \rightarrow A'$  there is a natural bijection between*

- (1)  $A'$ -module homomorphisms  $A' \otimes_A G \rightarrow A' \otimes_A K$ , and
- (2)  $A$ -algebra homomorphisms  $\text{Sym}_A(G \otimes_A K^\vee) \rightarrow A'$ .

The bijection maps  $\varphi : \text{Sym}_A(G \otimes_A K^\vee) \rightarrow A'$  to the homomorphism  $u$  defined by  $u(1_{A'} \otimes_A x) = (\varphi \otimes_A \text{id}_K)u_K(1 \otimes_A x)$  for all  $x \in G$ .

*Proof.* The set (2) is mapped to the set (1) via three isomorphisms:

- (1)  $\text{Hom}_{A\text{-alg}}(\text{Sym}_A(G \otimes_A K^\vee), A') \rightarrow \text{Hom}_A(G \otimes_A K^\vee, A')$  that follows from the definition of the symmetric algebra.
- (2)  $\text{Hom}_A(G \otimes_A K^\vee, A') \rightarrow \text{Hom}_A(G, A' \otimes_A K)$ , that is a canonical standard isomorphism, when  $K$  is free, that maps  $u : G \otimes_A K^\vee \rightarrow A'$  to the composite homomorphism  $G \xrightarrow{\text{id}_G \otimes_A \text{tr}} G \otimes_A K^\vee \otimes_A K \xrightarrow{u \otimes_A \text{id}_K} A' \otimes_A K$ .
- (3)  $\text{Hom}_A(G, A' \otimes_A K) \rightarrow \text{Hom}_{A'}(A' \otimes_A G, A' \otimes_A K)$ , that is the standard isomorphism obtained by *extension of scalars*.  $\square$

**Notation 2.3.** Let  $u : G \rightarrow K$  be a homomorphism of  $A$ -modules with  $K$  free of finite rank. We denote by  $\mathfrak{J}_Z(u)$  the image of the composite homomorphism

$$G \otimes_A K^\vee \xrightarrow{u \otimes_A \text{id}_{K^\vee}} K \otimes_A K^\vee \xrightarrow{\text{ev}} A.$$

**Lemma 2.4.** *For every  $A$ -algebra  $A \rightarrow A'$  the  $A'$ -module homomorphism*

$$A' \otimes_A G \xrightarrow{\text{id}_{A'} \otimes_A u} A' \otimes_A K$$

is zero if and only if the homomorphism  $A \rightarrow A'$  factors via the residue homomorphism  $A \rightarrow A/\mathfrak{I}_Z(u)$ .

*Proof.* This is an immediate consequence of the definition of  $\mathfrak{I}_Z(u)$ .  $\square$

**Notation 2.5.** Let

$$v : \text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee) \otimes_A G \otimes_A G \rightarrow \text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee) \otimes_A \text{End}_A(E)$$

be the  $\text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee)$ -module homomorphism defined by

$$v(1 \otimes_A x \otimes_A y) = u_{\text{End}_A(E)}(1 \otimes_A x)u_{\text{End}_A(E)}(1 \otimes_A y) - u_{\text{End}_A(E)}(1 \otimes_A y)u_{\text{End}_A(E)}(1 \otimes_A x),$$

where  $u_{\text{End}_A(E)}$  is defined in paragraph 2.1. We denote by  $H$  the residue algebra of  $\text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee)$  modulo the ideal  $\mathfrak{I}_Z(v)$  and let

$$\rho_H : \text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee) \rightarrow H$$

be the residue homomorphism. From the  $\text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee)$ -module homomorphism  $u_{\text{End}_A(E)}$  and the  $A$ -algebra homomorphism  $\rho_H$  we obtain an  $H$ -module homomorphism  $w : H \otimes_A G \rightarrow H \otimes_A \text{End}_A(E)$  defined by  $w(1 \otimes_A x) = (\rho_H \otimes_A 1_{\text{End}_A(E)})u_{\text{End}_A(E)}(1 \otimes_A x)$  for all  $x \in G$ . It follows from the definition of  $\mathfrak{I}_Z(w)$  and  $H$  that the elements of the image of  $w$  commute. Consequently  $w$  gives a unique  $H$ -algebra homomorphism

$$\mu_H : H \otimes_A \text{Sym}_A(G) \rightarrow H \otimes_A \text{End}_A(E)$$

such that  $\mu_H(1_H \otimes_A x) = w(1_H \otimes_A x)$  for all  $x \in G$ .

**Lemma 2.6.** *Let  $A \rightarrow A'$  be an  $A$ -algebra. We have a bijection between*

- (1)  $A'$ -algebra homomorphisms  $A' \otimes_A \text{Sym}_A(G) \rightarrow A' \otimes_A \text{End}_A(E)$ , and
- (2)  $A$ -algebra homomorphisms  $H \rightarrow A'$ .

*The bijection maps  $\varphi : H \rightarrow A'$  to the homomorphism  $u$  defined by  $u(1_{A'} \otimes_A x) = (\varphi \otimes_A \text{id}_{\text{End}_A(E)})\mu_H(1_H \otimes_A x)$  for all  $x \in G$ .*

*Proof.* It follows from Lemma 2.2 that there is a bijection between  $A'$ -module homomorphisms  $u : A' \otimes_A G \rightarrow A' \otimes_A \text{End}_A(E)$  and  $A$ -algebra homomorphisms  $\varphi : \text{Sym}_A(G \otimes_A \text{End}_A(E)^\vee) \rightarrow A'$ . By the definition of  $\mathfrak{I}_Z(w)$  and Lemma 2.4 the homomorphism  $\varphi$  factors via a homomorphism  $\chi : H \rightarrow A'$  if and only if the elements  $u(1_{A'} \otimes_A x)$  commute for all  $x \in X$ . However, the maps  $u : A' \otimes_A G \rightarrow A' \otimes_A \text{End}_A(E)$  whose images consist of commuting elements correspond to  $A'$ -algebra homomorphisms  $\psi : A' \otimes_A \text{Sym}_A(G) \rightarrow A' \otimes_A \text{End}_A(E)$  for which

$$\psi(1_{A'} \otimes_A x) = u(1_{A'} \otimes_A x)$$

for all  $x \in G$ .  $\square$

**Notation 2.7.** Let  $G$  be an  $A$ -module and  $\mathfrak{J}$  an ideal in  $\text{Sym}_A(G)$ . Denote by  $\iota : \mathfrak{J} \rightarrow \text{Sym}_A(G)$  the inclusion map, let  $B = \text{Sym}_A(G)/\mathfrak{J}$ , and let  $\rho_B : \text{Sym}_A(G) \rightarrow B$  be the residue homomorphism. We denote by  $v$  the composite  $H$ -module homomorphism

$$(2-1) \quad H \otimes_A \mathfrak{J} \xrightarrow{\text{id}_H \otimes_A \iota} H \otimes_A \text{Sym}_A(G) \xrightarrow{\mu_H} H \otimes_A \text{End}(E).$$

Moreover we let  $H_B$  be the residue algebra of  $H$  modulo the ideal  $\mathfrak{J}_Z(v)$  and we denote the residue homomorphism by

$$\rho_{H_B} : H \rightarrow H_B.$$

We tensor the modules of (2-1) by  $H_B$  over  $H$  and obtain a homomorphism of  $H_B$ -modules

$$H_B \otimes_A \mathfrak{J} \xrightarrow{\text{id}_{H_B} \otimes_A \iota} H_B \otimes_A \text{Sym}_A(G) \xrightarrow{\text{id}_{H_B} \otimes_H \mu_H} H_B \otimes_A \text{End}_A(E)$$

such that the composite homomorphism is zero by the definition of  $\mathfrak{J}_Z(v)$ . Consequently we obtain a homomorphism of  $H_B$ -algebras

$$\mu_{H_B} : H_B \otimes_A B \rightarrow H_B \otimes_A \text{End}_A(E)$$

such that  $\mu_{H_B}(1 \otimes_A \rho_B(f)) = (\rho_{H_B} \otimes_A \text{id}_{\text{End}_A(E)})\mu_H(1 \otimes_A f)$  for all  $f \in \text{Sym}_A(G)$ .

The algebra  $H_B$  parametrizes all  $B$ -module structures on  $E$ , and  $\mu_{H_B}$  is the *universal homomorphism*.

**Proposition 2.8.** *Let  $A \rightarrow A'$  be an  $A$ -algebra. We have a bijection between*

- (1)  $A'$ -algebra homomorphisms  $A' \otimes_A B \rightarrow A' \otimes_A \text{End}_A(E)$ , and
- (2)  $A$ -algebra homomorphisms  $H_B \rightarrow A'$ .

The bijection maps  $\varphi : H_B \rightarrow A'$  to the homomorphism  $u$  defined by  $u(1_{A'} \otimes_A f) = (\varphi \otimes_A \text{id}_{\text{End}_A(E)})\mu_{H_B}(1_{H_B} \otimes_A f)$  for all  $f \in B$ .

*Proof.* It follows from Lemma 2.4 that an  $A$ -algebra homomorphism  $H \rightarrow A'$  factors via  $\rho_{H_B} : H \rightarrow H_B$  if and only if the composite homomorphism

$$A' \otimes_A \mathfrak{J} \xrightarrow{\text{id}_{A'} \otimes_A \iota} A' \otimes_A \text{Sym}_A(G) \xrightarrow{\text{id}_{A'} \otimes_H \mu_H} A' \otimes_A \text{End}_A(E)$$

is zero. This last condition holds if and only if  $\text{id}_{A'} \otimes_H \mu_H$  factors via an  $A'$ -algebra homomorphism  $A' \otimes_A B \rightarrow A' \otimes_A \text{End}_A(E)$ . □

### 3. The local quot scheme

It follows from Lemma 2.2 that the  $A$ -algebra  $\mathrm{Sym}_A(F \otimes_A E^\vee)$  parametrizes homomorphisms  $F \rightarrow E$ . We use this, together with the properties of the  $A$ -algebra  $H_B$ , to construct a residue algebra  $Q^s$  of  $\mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B$  that parametrizes the *local quot functor* of Section 1.1.

**Notation 3.1.** We defined in 2.1 an  $A$ -module homomorphism

$$u_E : \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A F \rightarrow \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A E$$

such that  $u_E(1 \otimes_A y) = y \otimes_A \mathrm{tr}(1)$  for all  $y \in F$ .

Let  $A \rightarrow B$  be an  $A$ -algebra and fix a presentation  $0 \rightarrow \mathcal{J} \rightarrow \mathrm{Sym}_A(G) \rightarrow B \rightarrow 0$  of  $B$  as in 2.7. Moreover, fix an  $A$ -module homomorphism

$$s : E \rightarrow B \otimes_A F$$

and let

$$v : \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \otimes_A E \rightarrow \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \otimes_A E$$

be the composition of the  $\mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B$ -module homomorphisms

$$\begin{aligned} \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \otimes_A E &\xrightarrow{1 \otimes_A 1 \otimes_A s} \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \otimes_A B \otimes_A F \\ &\xrightarrow{\sim} \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A F \otimes_A H_B \otimes_A B \\ &\xrightarrow{u_E \otimes_A \mu_{H_B}} \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A E \otimes_A H_B \otimes_A \mathrm{End}_A(E) \\ &\xrightarrow{\sim} \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \otimes_A \mathrm{End}_A(E) \otimes_A E \\ &\xrightarrow{\mathrm{id} \otimes_A \mathrm{id} \otimes_A \mathrm{ev}} \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \otimes_A E, \end{aligned}$$

where the isomorphisms without names are the appropriate permutations of the factors in the tensor products. Let  $Q^s$  be the residue algebra of  $\mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B$  modulo the ideal  $\mathcal{J}_Z(v - \mathrm{id})$  and let

$$\rho_{Q^s} : \mathrm{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \rightarrow Q^s$$

be the residue homomorphism.

Denote by  $\rho_1 : \mathrm{Sym}_A(F \otimes_A E^\vee) \rightarrow Q^s$  and  $\rho_2 : H_B \rightarrow Q^s$  the  $A$ -algebra homomorphisms that determine  $\rho_{Q^s}$ , that is,  $\rho_1(f) = \rho_{Q^s}(f \otimes_A 1)$  and  $\rho_2(g) = \rho_{Q^s}(1 \otimes g)$  for all  $f \in \mathrm{Sym}_A(F \otimes_A E^\vee)$  and  $g \in H_B$ . We obtain a universal  $Q^s$ -algebra homomorphism

$$\mu_{Q^s} : Q^s \otimes_A B \rightarrow Q^s \otimes_A \mathrm{End}_A(E)$$

defined by  $\mu_{Q^s}(1_{Q^s} \otimes_A f) = (\rho_2 \otimes_A \mathrm{id}_{\mathrm{End}_A(E)}) \mu_{H_B}(1 \otimes_A f)$  for all  $f \in B$  and a  $Q^s$ -module homomorphism  $u'_{Q^s} : Q^s \otimes_A F \rightarrow Q^s \otimes_A E$  defined by  $u'_{Q^s}(1_{Q^s} \otimes_A y) =$

$(\rho_1 \otimes_A \text{id}_E)u_E(1 \otimes_A y)$  for all  $y \in F$ . When we give  $Q^s \otimes_A E$  the  $Q^s \otimes_A B$ -module structure given by  $\mu_{Q^s}$  we denote by

$$u_{Q^s} : Q^s \otimes_A B \otimes_A F \rightarrow Q^s \otimes_A E$$

the  $Q^s \otimes_A B$ -module homomorphism obtained from  $u'_{Q^s}$  by extension of scalars. Identifying  $Q^s \otimes_A B \otimes_A F$  with  $(Q^s \otimes_A B) \otimes_{Q^s} (Q^s \otimes_A F)$  and  $Q^s \otimes_A \text{End}_A(E) \otimes_A E$  with  $(Q^s \otimes_A \text{End}_A(E)) \otimes_{Q^s} (Q^s \otimes_A E)$  we obtain a composite homomorphism

$$Q^s \otimes_A E \xrightarrow{\text{id}_{Q^s} \otimes_A s} Q^s \otimes_A B \otimes_A F \xrightarrow{\mu_{Q^s} \otimes_A u'_{Q^s}} Q^s \otimes_A \text{End}_A(E) \otimes_A E \xrightarrow{\text{id}_{Q^s} \otimes_A \text{ev}} Q^s \otimes_A E$$

that is the identity by the definition of  $\mathfrak{J}_Z(v - \text{id})$  and  $Q^s$ .

**Theorem 3.2.** *The  $A$ -algebra  $Q^s$  represents the local quot functor defined in Section 1.1. The universal homomorphisms are  $\mu_{Q^s} : Q^s \otimes_A B \rightarrow Q^s \otimes_A \text{End}_A(E)$  and  $u_{Q^s} : Q^s \otimes_A B \otimes_A F \rightarrow Q^s \otimes_A E$ .*

*More precisely, let  $A \rightarrow A'$  be an  $A$ -algebra. We have bijections between the following three sets:*

- (1)  *$A' \otimes_A B$ -module structures on  $A' \otimes_A E$  and  $A' \otimes_A B$ -linear homomorphisms  $u : A' \otimes_A B \otimes_A F \rightarrow A' \otimes_A E$  for the  $A' \otimes_A B$ -module structure such that*

$$A' \otimes_A E \xrightarrow{1 \otimes_A s} A' \otimes_A B \otimes_A F \xrightarrow{u} A' \otimes_A E$$

*is the identity.*

- (2)  *$A'$ -module homomorphisms  $v : A' \otimes_A F \rightarrow A' \otimes_A E$  and  $A'$ -algebra homomorphisms  $\varphi : A' \otimes_A B \rightarrow A' \otimes_A \text{End}(E)$  such that the composite homomorphism*

$$(3-1) \quad A' \otimes_A E \xrightarrow{\text{id}_{A'} \otimes_A s} A' \otimes_A B \otimes_A F \xrightarrow{\varphi \otimes_A v} A' \otimes_A \text{End}_A(E) \otimes_A E \xrightarrow{\text{id}_{A'} \otimes_A \text{ev}} A' \otimes_A E$$

*is the identity.*

- (3)  *$A$ -algebra homomorphisms  $Q^s \rightarrow A'$ .*

*The bijection from the set (2) to the set (1) is described in Proposition 1.3 and the bijection from the set (3) to the set (2) is defined as follows:*

*Let  $\varphi : Q^s \rightarrow A'$  be an  $A$ -algebra homomorphism. The homomorphism  $\varphi \rho_{Q^s} : \text{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \rightarrow A'$  is determined by  $A$ -algebra homomorphisms  $\rho_1 : \text{Sym}_A(F \otimes_A E^\vee) \rightarrow A'$  and  $\rho_2 : H_B \rightarrow A'$ . The homomorphism  $\rho_2$  defines an  $A'$ -algebra homomorphism  $\psi : A' \otimes_A B \rightarrow A' \otimes_A \text{End}_A(E)$  by Proposition 2.8, and  $\rho_1$  defines, by Lemma 2.2 with  $E = K$  and  $G = F$ , an  $A'$ -module homomorphism  $v : A' \otimes_A F \rightarrow A' \otimes_A E$ . We extend  $v$  to an  $A' \otimes_A B$ -module homomorphism  $A' \otimes_A B \otimes_A F \rightarrow A' \otimes_A E$ , when  $A' \otimes_A E$  has the  $A' \otimes_A B$ -module structure given by  $\psi$ .*

*Proof.* The bijection between (1) and (2) is given in Corollary 1.4 when we use the canonical isomorphism of  $A'$ -module  $A' \otimes_A \text{End}_A(E) \rightarrow \text{End}_{A'}(A' \otimes_A E)$ .

From the description of the map from the set in (3) to the set in (2) given in the Theorem it follows that the map is a bijection, since an  $A$ -algebra homomorphism  $\text{Sym}_A(F \otimes_A E^\vee) \otimes_A H_B \rightarrow A'$  factors via  $\rho_{Q^s}$  if and only if the composite homomorphism (3-1) is the identity.

That the map from the set (3) to the set (1) is functorial follows from the explicit description of the maps in the Theorem. □

### 4. The quot functor

**Definition 4.1.** Let  $f : X \rightarrow S$  be a morphism of schemes. For every morphism  $g : S' \rightarrow S$  we write

$$\begin{array}{ccc}
 X' = X \times_S S' & \xrightarrow{g'} & X \\
 f' \downarrow & & \downarrow f \\
 S' & \xrightarrow{g} & S.
 \end{array}$$

Let  $\mathcal{M}$  be a quasicohherent  $\mathcal{O}_X$ -module that is flat over  $S$  and such that  $\text{Supp } \mathcal{M}$  is a scheme that is finite over  $S$ . Here  $\text{Supp } \mathcal{M}$  is the subscheme of  $X$  defined by the annihilator of  $\mathcal{M}$ . We say that  $\mathcal{M}$  is of *relative rank  $n$*  over  $S$  if  $f_*\mathcal{M}$  is a locally free  $\mathcal{O}_S$ -module of rank  $n$ . The latter condition is equivalent to the condition that  $\dim_{\kappa(s)}(f_*\mathcal{M} \otimes_{\mathcal{O}_X} \kappa(s)) = n$  for all points  $s \in S$  (see [Laksov et al. 2000]).

Let  $\mathcal{F}$  be a quasicohherent  $\mathcal{O}_X$ -module. We denote by  $\text{Quot}_{\mathcal{F}/X/S}^n$  the functor from  $S$ -schemes to sets that to a morphism  $g : S' \rightarrow S$  associates the set  $\text{Quot}_{\mathcal{F}/X/S}^n(S')$  of classes of surjections  $g'^*\mathcal{F} \rightarrow \mathcal{M}$  of  $\mathcal{O}_{X'}$ -modules, where  $\mathcal{M}$  is a coherent  $\mathcal{O}_{X'}$ -module which is flat over  $S'$  with support that is a finite scheme over  $S'$  with *relative rank  $n$*  (see [Grothendieck 1966] or [Deligne 1973]).

Let  $\mathcal{E}$  be a locally free  $\mathcal{O}_S$ -module of rank  $n$  and fix an  $\mathcal{O}_S$ -module homomorphism  $s : \mathcal{E} \rightarrow f_*\mathcal{F}$ . We denote by  $\text{Quot}_{\mathcal{F}/X/S}^s$  the subfunctor of  $\text{Quot}_{\mathcal{F}/X/S}^n$  that to an  $S$ -scheme  $S'$  associates the set  $\text{Quot}_{\mathcal{F}/X/S}^s(S')$  of all equivalence classes of surjections  $g'^*\mathcal{F} \rightarrow \mathcal{M}$  such that the composite homomorphism

$$g^*\mathcal{E} \xrightarrow{g^*s} g^*f_*\mathcal{F} \rightarrow f'_*g'^*\mathcal{F} \rightarrow f'_*\mathcal{M}$$

is surjective, that is, an isomorphism.

When convenient we write  $\text{Quot}_{\mathcal{F}}^n$  and  $\text{Quot}_{\mathcal{F}}^s$  for the functors  $\text{Quot}_{\mathcal{F}/X/S}^n$ , respectively  $\text{Quot}_{\mathcal{F}/X/S}^s$ , and indicate in the text that the functors are taken relative to the homomorphism  $f : X \rightarrow S$ .

**Lemma 4.2.** *Let  $A \rightarrow B$  be an  $A$ -algebra and let  $M$  be a  $B$ -module that is free of rank  $n$  as an  $A$ -module. Then  $B/\text{Ann}_B(M)$  is integral over  $A$ .*

*Proof.* Since  $M$  is finitely generated as an  $A$ -module it is finitely generated as a  $B$ -module, say by  $x_1, \dots, x_n$ . We obtain a  $B$ -module homomorphism  $B \rightarrow \prod_{i=1}^n M$  that maps  $b$  to  $(bx_1, \dots, bx_n)$ . This gives an injection  $B/\text{Ann}_B(M) \rightarrow \prod_{i=1}^n M$  of  $B$ -modules. In particular,  $\prod_{i=1}^n M$  is a faithful  $B/\text{Ann}_B(M)$ -module. Moreover, the composite homomorphism  $A \rightarrow B/\text{Ann}_B(M) \rightarrow \prod_{i=1}^n M$  is injective since  $M$  is free as an  $A$ -module. For every element  $f \in B/\text{Ann}_B(M)$ , the module  $A[f]$  is contained in the finitely generated  $A$ -module  $\prod_{i=1}^n M$  that is faithful over  $A[f]$ . Hence  $f$  is integral over  $A$ ; see [Lang 1993, Chapter VII, §1 INT3].  $\square$

**Proposition 4.3.** *Let  $f : X \rightarrow S$  be a morphism of affine schemes. Assume that  $\mathcal{E}$  is a free  $\mathbb{O}_S$ -module and that  $\mathcal{F} = f^*\mathcal{F}_0$  where  $\mathcal{F}_0$  is an  $\mathbb{O}_S$ -module. Then the functor  $\text{Quot}_{\mathcal{F}/X/S}^s$  is representable by  $Q^s$  for  $S = \text{Spec}(A)$ ,  $X = \text{Spec}(B)$ ,  $E = \Gamma(S, \mathcal{E})$ ,  $F = \Gamma(S, \mathcal{F}_0)$  and  $s : E \rightarrow B \otimes_A F$  corresponds to the homomorphism  $s$ .*

*Proof.* In the correspondence between affine schemes over  $S$  and  $A$ -algebras we see that, in order to represent  $\text{Quot}_{\mathcal{F}/X/S}^s$ , we must represent the functor that to  $A'$  associates the  $A' \otimes_A B$ -module homomorphisms  $u : A' \otimes_A B \otimes_A F \rightarrow M$  where  $M$  is a free  $A'$ -module of rank  $n$  such that

$$A' \otimes_A E \xrightarrow{\text{id}_{A'} \otimes_A s} A' \otimes_A B \otimes_A F \xrightarrow{u} M$$

is surjective. This functor is representable by Theorem 3.2, taken into account that the condition  $\text{Supp } M = A' \otimes_A B/\text{Ann}_{A' \otimes_A B}(M)$  is finite over  $A'$  is automatically fulfilled by Lemma 4.2.  $\square$

Like several of the reductions of this section, the next result is well known.

**Lemma 4.4.** *Let  $f : X \rightarrow S$  be a homomorphism of affine schemes. Assume that  $\mathcal{E}$  is a free  $\mathbb{O}_S$ -module. Then  $\text{Quot}_{\mathcal{F}/X/S}^s$  is representable.*

*More precisely, there is a free  $\mathbb{O}_S$ -module  $\mathcal{F}_0$ , a surjection  $u : f^*\mathcal{F}_0 \rightarrow \mathcal{F}$  of  $\mathbb{O}_X$ -modules, and a homomorphism of  $\mathbb{O}_S$ -modules  $s_0 : \mathcal{E} \rightarrow f_*f^*\mathcal{F}_0$  such that  $s = (f_*u)s_0$ . These homomorphisms make  $\text{Quot}_{\mathcal{F}/X/S}^s$  into a closed subfunctor of  $\text{Quot}_{f^*\mathcal{F}_0/X/S}^{s_0}$ .*

*Proof.* Let  $F = \Gamma(X, \mathcal{F})$  and  $E = \Gamma(S, \mathcal{E})$ . Choose a surjection of  $A$ -modules  $F_0 \rightarrow F$  with  $F_0$  free. We then obtain, by extension of scalars, a surjection of  $B$ -modules  $B \otimes_A F_0 \rightarrow F$ , and consequently a surjection  $u : f^*\mathcal{F}_0 \rightarrow \mathcal{F}$  of  $\mathbb{O}_X$ -modules with  $\mathcal{F}_0 = \widetilde{F_0}$ . We lift  $s : E \rightarrow F$  to an  $A$ -module homomorphism  $E \rightarrow B \otimes_A F_0$  via the surjection  $B \otimes_A F_0 \rightarrow F$ . The corresponding lifting  $s_0 : \mathcal{E} \rightarrow f_*f^*\mathcal{F}_0$  has the property that  $s = (f_*u)s_0$ .

Clearly, given an arbitrary  $A$ -algebra  $A \rightarrow A'$ , we have a map  $\text{Quot}_{\mathcal{F}/X/S}^n(A') \rightarrow \text{Quot}_{f^*\mathcal{F}_0/X/S}^n(A')$  that takes  $\mathcal{F} \xrightarrow{v} M$  to the composite homomorphism

$$f^*\mathcal{F}_0 \xrightarrow{u} \mathcal{F} \xrightarrow{v} M,$$

and this map defines a closed immersion of functors  $\text{Quot}_{\mathcal{F}/X/S}^n \rightarrow \text{Quot}_{f^*\mathcal{F}_0/X/S}^n$ . Moreover the closed immersion maps  $\text{Quot}_{\mathcal{F}/X/S}^s$  into  $\text{Quot}_{f^*\mathcal{F}_0/X/S}^{s_0}$  because

$$\mathcal{E} \xrightarrow{s} f_*\mathcal{F} \xrightarrow{v} \mathcal{M}$$

is surjective if and only if  $\mathcal{E} \xrightarrow{s_0} f_*f^*\mathcal{F}_0 \xrightarrow{f_*u} f_*\mathcal{F} \xrightarrow{v} \mathcal{M}$  is surjective.

For the representability of  $\text{Quot}_{f^*\mathcal{F}_0/X/S}^{s_0}$ , use Theorem 3.2 and Proposition 4.3. □

**Proposition 4.5.** *Let  $f : X \rightarrow S$  be a homomorphism of affine schemes. Then  $\text{Quot}_{\mathcal{F}/X/S}^n$  is representable.*

*More precisely, the functor  $\text{Quot}_{\mathcal{F}/X/S}^n$  is covered by open affine subfunctors  $\text{Quot}_{\mathcal{F}/X/S}^s$  for all  $\mathbb{O}_S$ -module homomorphisms  $s : \mathcal{E} \rightarrow f_*\mathcal{F}$ , where  $\mathcal{E}$  is a free  $\mathbb{O}_S$ -module of rank  $n$ .*

*Proof.* It is clear that the subfunctors  $\text{Quot}_{\mathcal{F}/X/S}^s$  of  $\text{Quot}_{\mathcal{F}/X/S}^n$  are open. We have to show that  $\text{Quot}_{\mathcal{F}/X/S}^n$  is covered by these functors. Let  $S = \text{Spec}(A)$ ,  $X = \text{Spec}(B)$ ,  $F = \Gamma(\text{Spec}(B), \mathcal{F})$  and  $E = \Gamma(\text{Spec}(A), \mathcal{E})$ . For every  $A$ -algebra  $A \rightarrow A'$  we let the  $A'$ -module homomorphism  $u : A' \otimes_A F \rightarrow M$  correspond to an element  $g'^*\mathcal{F} \rightarrow \mathcal{M}$  in  $\text{Quot}_{\mathcal{F}/X/S}^n(\text{Spec}(A'))$ . For every maximal ideal  $\mathfrak{p}$  in  $A'$  we can, since  $M$  is a finitely generated  $A'$ -module, find an element  $a' \in A' \setminus \mathfrak{p}$  and an  $A$ -module homomorphism  $s_M : E \rightarrow F$  such that the composite  $A'_{a'}$ -module homomorphism  $A'_{a'} \otimes_A E \xrightarrow{s_M} A'_{a'} \otimes_A F \rightarrow M_{a'}$  is surjective. This shows that the image  $g'^*\mathcal{F}|_{f'^{-1}\text{Spec}(A'_{a'})} \rightarrow \mathcal{M}|_{f'^{-1}\text{Spec}(A'_{a'})}$  of the element  $g'^*\mathcal{F} \rightarrow \mathcal{M}$  by the map  $\text{Quot}_{\mathcal{F}/X/S}^n(\text{Spec}(A')) \rightarrow \text{Quot}_{\mathcal{F}/X/S}^n(\text{Spec}(A'_{a'}))$  lies in  $\text{Quot}_{\mathcal{F}/X/S}^s(\text{Spec}(A'_{a'}))$  so the set  $\text{Quot}_{\mathcal{F}/X/S}^s(\text{Spec}(A'_{a'}))$  covers  $g'^*\mathcal{F}|_{f'^{-1}\text{Spec}(A'_{a'})} \rightarrow \mathcal{M}|_{f'^{-1}\text{Spec}(A'_{a'})}$  considered as an element in  $\text{Quot}_{\mathcal{F}/X/S}^n(\text{Spec}(A'_{a'}))$ .

The representability of  $\text{Quot}_{\mathcal{F}/X/S}^s$  follows from Lemma 4.4 and it follows that the Zariski sheaf  $\text{Quot}_{\mathcal{F}/X/S}^n$  is representable. □

**Lemma 4.6.** *Let  $R$  be a graded  $A$ -algebra. For every prime ideal  $\mathfrak{p}$  of  $A$  write  $\kappa(\mathfrak{p}) = A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$ .*

*Let  $Z$  be a closed subscheme of  $\text{Proj}(\kappa(\mathfrak{p}) \otimes_A R)$  that is finite over  $\text{Spec}(\kappa(\mathfrak{p}))$ . Then there is an element  $a \in A$  not in  $\mathfrak{p}$  and an element  $f \in R_a$  such that  $Z$  is contained in the open subscheme  $\text{Spec}(\kappa(\mathfrak{p}) \otimes_{A_a} (R_a)_{(f)}) = \text{Spec}(\kappa(\mathfrak{p})) \times_{\text{Spec}(A_a)} \text{Spec}((R_a)_{(f)})$  of  $\text{Proj}(\kappa(\mathfrak{p}) \otimes_{A_a} R_a) = \text{Spec}(\kappa(\mathfrak{p})) \times_{\text{Spec}(A_a)} \text{Proj}(R_a)$ .*

*Proof.* Since  $Z$  is finite over  $\text{Spec}(\kappa(\mathfrak{p}))$  the fiber of the induced morphism  $Z \rightarrow \text{Spec}(\kappa(\mathfrak{p}))$  consists of a finite number of points, corresponding to homogeneous prime ideals  $\mathfrak{q}_1, \dots, \mathfrak{q}_k$  in  $\kappa(\mathfrak{p}) \otimes_A R$  that do not contain the irrelevant ideal. Their union consequently does not contain the irrelevant ideal. Hence we can find a homogeneous element  $g \in \kappa(\mathfrak{p}) \otimes_A R$  of positive degree that is not contained in

any of the ideals  $\mathfrak{q}_1, \dots, \mathfrak{q}_k$ . Thus  $Z$  is contained in the open subscheme

$$\mathrm{Spec}((\kappa(\mathfrak{p}) \otimes_A R)_{(g)})$$

of  $\mathrm{Proj}(\kappa(\mathfrak{p}) \otimes_A R)$ .

Clearly we can find an element  $a \in A$  not in  $\mathfrak{p}$  and an element  $f \in R_a$  such that  $1_{\kappa(\mathfrak{p}) \otimes_{A_a} f}$  is the image of  $g$  by the natural isomorphism  $\kappa(\mathfrak{p}) \otimes_A R \rightarrow \kappa(\mathfrak{p}) \otimes_{A_a} R_a$ . However, then  $\mathrm{Spec}((\kappa(\mathfrak{p}) \otimes_A R)_{(g)}) = \mathrm{Spec}(\kappa(\mathfrak{p}) \otimes_{A_a} (R_a)_{(f)})$ , and we have proved the Lemma.  $\square$

**Theorem 4.7.** *Let  $\mathcal{R}$  be a quasicoherent sheaf that is a graded  $\mathbb{O}_S$ -algebra, let  $X = \mathrm{Proj}(\mathcal{R})$ , and let  $\mathcal{F}$  be a quasicoherent  $\mathbb{O}_X$ -module. Then  $\mathrm{Quot}_{\mathcal{F}/X/S}^n$  is representable.*

*More precisely, for every open affine subscheme  $\mathrm{Spec}(A)$  of  $S$  we write  $R = \Gamma(\mathrm{Spec}(A), \mathcal{R})$ . Then  $\mathrm{Quot}_{\mathcal{F}/X/S}^n$  is covered by open subfunctors naturally isomorphic to  $\mathrm{Quot}_{\mathcal{F}|\mathrm{Spec}(R_{(r)})}^n$  relative to  $\mathrm{Spec}(R_{(r)}) \rightarrow \mathrm{Spec}(A)$  for all  $\mathrm{Spec}(A)$  in an open covering of  $S$  and all homogeneous elements  $r$  of positive degree in  $R$ .*

*Proof.* For every affine open subset  $U$  of  $S$  we consider  $\mathrm{Quot}_{\mathcal{F}|f^{-1}(U)}^n$  relative to  $f^{-1}(U) \rightarrow U$  as a subfunctor of  $\mathrm{Quot}_{\mathcal{F}/\mathrm{Proj}(\mathcal{R})/S}$  by letting

$$\mathrm{Quot}_{\mathcal{F}|f^{-1}(U)}^n(S') = \emptyset$$

when  $g : S' \rightarrow S$  does not factor via  $U$ . It is clear that the subfunctors  $\mathrm{Quot}_{\mathcal{F}|f^{-1}(U)}^n$  are open and that they cover  $\mathrm{Quot}_{\mathcal{F}/\mathrm{Proj}(\mathcal{R})/S}^n$ . Consequently we can assume that  $S = \mathrm{Spec}(A)$  is affine.

For every  $a \in A$  and every  $r \in R_a$  we can consider the functor  $\mathrm{Quot}_{a,r} = \mathrm{Quot}_{\mathcal{F}|\mathrm{Spec}((R_a)_{(r)})}^n$  relative to  $\mathrm{Spec}(R_a)_{(r)} \rightarrow \mathrm{Spec}(A_a)$  as a subfunctor of the functor  $\mathrm{Quot}_a = \mathrm{Quot}_{\mathcal{F}|\mathrm{Proj}(R_a)}^n$  relative to  $\mathrm{Proj}(R_a) \rightarrow \mathrm{Spec}(A_a)$ . This is because, if  $g : S' \rightarrow \mathrm{Spec}(A_a)$  is a morphism and  $g'^* \mathcal{F}|f^{-1}(\mathrm{Spec}(R_a)_{(r)}) \rightarrow \mathcal{M}$  represents an element in  $\mathrm{Quot}_{a,r}(S')$ , then  $\mathrm{Supp} \mathcal{M} \subseteq f^{-1}(\mathrm{Spec}((R_a)_{(r)}))$  and  $\mathrm{Supp} \mathcal{M}$  is finite over  $\mathrm{Spec}(A_a)$ , and  $\mathrm{Spec}((R_a)_{(r)}) \rightarrow \mathrm{Spec}(A_a)$  is separated so  $\mathrm{Supp} \mathcal{M}$  is closed in  $\mathrm{Proj}(R_a) \times_{\mathrm{Spec}(A_a)} S'$ . Thus we can extend  $\mathcal{M}$  uniquely by zero to an  $\mathbb{O}_{\mathrm{Proj}(R_a) \times_{\mathrm{Spec}(A_a)} S'}$ -module  $\mathcal{N}$  and the surjection  $g'^* \mathcal{F}|f'^{-1}(\mathrm{Spec}((R_a)_{(r)})) \rightarrow \mathcal{M}$  can be extended to a surjection  $g'^* \mathcal{F} \rightarrow \mathcal{N}$  representing an element in  $\mathrm{Quot}_a(\mathrm{Spec}(A_a))$ . It is clear that the subfunctors  $\mathrm{Quot}_{a,r}$  are open. It remains to show that they cover  $\mathrm{Quot}_{\mathcal{F}/\mathrm{Proj}(R)/\mathrm{Spec}(A)}$ . For this it suffices to show that for every prime ideal  $\mathfrak{p} \subset A$  and every surjection  $g'^* \mathcal{F} \rightarrow \mathcal{M}$ , with  $g' : \mathrm{Spec}(\kappa(\mathfrak{p})) \times_S \mathrm{Proj}(R_a) \rightarrow S' = \mathrm{Spec}(\kappa(\mathfrak{p}))$ , where  $\mathcal{M}$  has finite support over  $\mathrm{Spec}(\kappa(\mathfrak{p}))$  of relative rank  $n$ , there is an  $a \in A \setminus \mathfrak{p}$  and a homogeneous element  $r$  in  $R_a$  such that the support of  $\mathcal{M}$  is contained in the open subscheme  $\mathrm{Spec}(\kappa(\mathfrak{p}) \otimes_A (R_a)_{(r)}) = \mathrm{Spec}(\kappa(\mathfrak{p})) \times_{\mathrm{Spec}(A_a)} \mathrm{Spec}((R_a)_{(r)})$  of  $\mathrm{Proj}(\kappa(\mathfrak{p}) \otimes_A R_a) = \mathrm{Spec}(\kappa(\mathfrak{p})) \times_{\mathrm{Spec}(A_a)} \mathrm{Proj}(R_a)$ . However this is the assertion of Lemma 4.6.  $\square$

### 5. Applications

**5.1. Special cases.** The two best known special cases of Theorem 4.7 are the Hilbert schemes and the Grassmann schemes. As in the Theorem we let  $X = \text{Proj}(\mathcal{R})$  with  $\mathcal{R}$  quasicoherent on the  $S$ -scheme  $X$ , and  $\mathcal{F}$  is a quasicoherent  $\mathcal{O}_X$ -module.

When  $\mathcal{F} = \mathcal{O}_X$  we obtain the existence and construction of the Hilbert scheme  $\text{Hilb}_{X/S}^n$  of  $n$  points in  $X$ , as in [Gustavsen et al. 2007].

When  $X = S$  the Theorem gives the existence and the standard description of the Grassmann scheme  $\text{Grass}_{\mathcal{F}/S}^n$  of  $n$ -quotients of the quasicoherent  $\mathcal{O}_X$ -module  $\mathcal{F}$ .

**5.2. Example: Quot schemes over the affine line.** Let  $A[T]$  be the polynomial ring over  $A$  in the variable  $T$ . Moreover, let  $F$  be a free module of rank  $r$ , and let  $S = \text{Spec}(A)$ ,  $X = \text{Spec}(A[T])$  and  $\mathcal{F} = A[T] \otimes_A F$ . We shall show how our construction gives an open covering of  $\text{Quot}_{\mathcal{F}/X/S}^n$  by affine spaces of relative dimension  $rn$  over  $\text{Spec}(A)$ , with an intersection that contains an affine space of relative dimension  $n$ . In particular, since  $\text{Quot}_{\mathcal{F}/X/S}^n$  can be covered by mutually intersecting open sets that are affine spaces over  $\text{Spec}(A)$  of relative dimension  $rn$ , it is irreducible of relative dimension  $rn$  if and only if  $\text{Spec}(A)$  is irreducible.

Let  $E$  be a vector space with basis  $e_1, \dots, e_n$ , and let  $f_1, \dots, f_r$  be a basis for  $F$ . From Proposition 4.5 we have that  $\text{Quot}_{\mathcal{F}/X/S}^n$  can be covered by open affine subsets  $\text{Quot}_{\mathcal{F}/X/S}^s$ , where the maps  $s : E \rightarrow A[T] \otimes_A F$  are linear.

Let  $X_{ij}$  for  $i, j = 1, \dots, n$  and  $Y_{ij}$  for  $i = 1, \dots, n, j = 1, \dots, r$  be independent variables over  $A$ , and let  $(X_{ij})$ , be the  $n \times n$ -matrix with coordinates  $X_{ij}$ , and  $(Y_{ij})$  the  $n$ -vector with coordinates  $Y_{ij}$  for fixed  $j$ . We denote by  $C$  the ring of polynomials over  $A$  in all these variables, and define maps

$$\varphi : C \otimes_A A[T] \rightarrow C \otimes_A \text{End}_A(E) \quad \text{and} \quad v : C \otimes_A F \rightarrow C \otimes_A E$$

of  $C$ -algebras, respectively of  $C$ -modules, by  $\varphi(T) = (X_{ij})$  with respect to the basis  $e_1, \dots, e_n$ , respectively by  $v(1 \otimes f_j) = (Y_{ij})$ , with respect to the bases  $e_1, \dots, e_n$  and  $f_1, \dots, f_r$ .

In Sections 2 and 3 we proved that  $\text{Quot}_{\mathcal{F}/X/S}^s$  is given as the residue algebra of  $C$  modulo the ideal generated by the relations we obtain requiring that the composite homomorphism

(5-1)

$$C \otimes_A E \xrightarrow{\text{id}_C \otimes_A s} C \otimes_A A[T] \otimes_A F \xrightarrow{\varphi \otimes_A v} C \otimes_A \text{End}_A(E) \otimes_A E \xrightarrow{\text{id}_C \otimes_A \text{ev}} C \otimes_A E$$

be the identity. It follows from the Cayley–Hamilton Theorem used on  $\varphi(T) = (X_{ij})$  that  $\psi := (\text{id}_C \otimes_A \text{ev})(\varphi \otimes_A v)$  is surjective if and only if its restriction to the  $C$ -submodule of  $C \otimes_A A[T] \otimes_A F$  generated by the elements  $1 \otimes T^i \otimes f_j$ , for

$i = 0, \dots, n - 1$  and  $j = 1, \dots, r$ , is surjective. We will describe a collection of maps  $s : E \rightarrow A[T] \otimes_A F$  so that  $\text{Quot}_{\mathcal{F}/X/S}^s$  cover  $\text{Quot}_{\mathcal{F}/X/S}^n$ . To do this we may clearly assume that  $s$  maps the elements of  $e_1, \dots, e_n$  to the elements of the form  $T^i \otimes f_j$  for  $i = 0, \dots, n - 1$  and  $j = 1, \dots, r$ .

Let  $s$  be given, and let  $p_j$  be the number of elements of the form  $T^i \otimes f_j$ , for fixed  $j$  that are in the image by  $s$  of the elements  $e_1, \dots, e_n$ . In particular  $p_1 + \dots + p_r = n$ . Since  $\psi$  is a  $C[T]$ -module homomorphism for the  $C[T]$ -module structure on  $C \otimes_A E$  given by  $\varphi$  we conclude that, if the restriction of  $\psi$  to the  $C$ -module generated by  $s(e_1), \dots, s(e_n)$  is surjective, then  $\psi$  restricted to the  $C$ -module generated by  $1 \otimes f_j, \dots, T^{p_j-1} \otimes f_j$  for  $j = 1, \dots, r$  is surjective. Consequently, we can cover  $\text{Quot}_{\mathcal{F}/X/S}^n$  by the sets  $\text{Quot}_{\mathcal{F}/X/S}^s$  where  $s$  is given by

$$(5-2) \quad s(e_{q_{j-1}+i+1}) = T^i \otimes f_j \quad \text{for } j = 1, \dots, r \quad \text{and } i = 0, \dots, p_j - 1$$

with  $q_j = p_1 + \dots + p_j$  and  $q_0 = 0$ .

The image of  $1 \otimes e_{q_{j-1}+i+1}$  by (5-1) is  $\varphi(T)^i v(1 \otimes f_j)$ . For  $i = 0$  we obtain, in particular, that the condition that (5-1) is the identity is  $(Y_{ij}) = e_{q_{j-1}+1}$  for those  $j = 1, \dots, r$  that satisfy  $p_j \geq 1$ . Hence the image of  $1 \otimes e_{q_{j-1}+i+1}$  by (5-1) is  $\varphi(T)^i e_{q_{j-1}+1}$  for these  $j$ . By induction on  $i$  we easily verify that the condition that (5-1) is the identity is that column number  $i$  in  $(X_{ij})$  is equal to  $e_{q_{j-1}+i+1}$  for  $i = 1, \dots, p_j - 1$ .

Consequently, the condition for Equation (5-1) to be the identity is that

$$v(1 \otimes f_j) = 1 \otimes e_{q_{j-1}+1}$$

for those  $j = 1, \dots, r$  that satisfy  $p_j \geq 1$ , and that  $\varphi(T)$  has column number  $i$  equal to  $e_{i+1}$  when  $i$  is different from  $q_1, \dots, q_r$ , and with no conditions on columns number  $q_1, \dots, q_r$ . Consequently  $\text{Quot}_{\mathcal{F}/X/S}^s$  is affine space of relative dimension  $nr$  when  $s$  is determined by Equation (5-2). Note that, independently of the choice of  $s$  satisfying the conditions (5-2),  $\text{Quot}_{\mathcal{F}/X/S}^s$  contains the affine space of relative dimension  $n$  consisting of the homomorphisms  $\varphi$  such that  $\varphi(T)$  is the companion matrix of the polynomial  $T^n - X_{nn}T^{n-1} - X_{n-1n}T^{n-2} - \dots - X_{1n}$ .

### References

[Altman and Kleiman 1980] A. B. Altman and S. L. Kleiman, “Compactifying the Picard scheme”, *Adv. in Math.* **35**:1 (1980), 50–112. MR 81f:14025a Zbl 0427.14015

[Deligne 1973] P. Deligne, “Cohomologie à supports propres”, pp. 250–480 (exposé XVII) in *Théorie des topos et cohomologie étale des schémas* (SGA4), vol. 3, edited by M. Artin et al., Lecture Notes in Mathematics **305**, Springer, Berlin, 1973. MR 50 #7132 Zbl 0255.14011

[Grothendieck 1966] A. Grothendieck, “Techniques de construction et théorèmes d’existence en géométrie algébrique, IV: Les schémas de Hilbert”, pp. 249–276 (exposé 221) in *Séminaire Bourbaki 1960/1961*, Benjamin, Paris, 1966. Also in vol. 6 of the reprint by Soc. Math. de France, Paris, 1995.

- [Gustavsen et al. 2007] T. S. Gustavsen, D. Laksov, and R. M. Skjelnes, “An elementary, explicit, proof of the existence of Hilbert schemes of points”, *J. Pure Appl. Algebra* **210**:3 (2007), 705–720.
- [Huibregtse 2002] M. E. Huibregtse, “A description of certain affine open subschemes that form an open covering of  $\text{Hilb}_{\mathbb{A}_k^n}^n$ ”, *Pacific J. Math.* **204** (2002), 97–143. MR 2003h:14008 Zbl 1068.14007
- [Huibregtse 2006] M. E. Huibregtse, “An elementary construction of the multigraded Hilbert scheme of points”, *Pacific J. Math.* **223**:2 (2006), 269–315. MR MR2221028 Zbl 05129581
- [Laksov et al. 2000] D. Laksov, Y. Pitteroud, and R. M. Skjelnes, “Notes on flatness and the Quot functor on rings”, *Comm. Algebra* **28**:12 (2000), 5613–5627. MR 2001i:13009 Zbl 0990.13005
- [Lang 1993] S. Lang, *Algebra*, 3rd ed., Addison-Wesley, Reading, MA, 1993. MR 33 #5416 Zbl 0848.13001
- [Mumford 1966] D. Mumford, *Lectures on curves on an algebraic surface*, Annals of Mathematics Studies **59**, Princeton Univ. Press, Princeton, NJ, 1966. MR 35 #187 Zbl 0187.42701
- [Sernesi 2006] E. Sernesi, *Deformations of algebraic schemes*, Grundlehren der Math. Wiss **334**, Springer, Berlin, 2006. MR MR2247603 Zbl 1102.14001

Received January 28, 2007. Revised May 16, 2007.

TROND STØLEN GUSTAVSEN  
 DEPARTMENT OF ECONOMICS  
 BI NORWEGIAN SCHOOL OF MANAGEMENT  
 N-0442 OSLO  
 NORWAY  
 stolen@math.uio.no

DAN LAKSOV  
 DEPARTMENT OF MATHEMATICS  
 KTH  
 S-100 44 STOCKHOLM  
 SWEDEN  
 laksov@math.kth.se

ROY MIKAEL SKJELNES  
 DEPARTMENT OF MATHEMATICS  
 KTH  
 S-100 44 STOCKHOLM  
 SWEDEN  
 skjelnes@math.kth.se



# PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982)

F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Paulo Ney de Souza, Production Manager

Silvio Levy, Senior Production Editor

Alexandru Scorpan, Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.  
PEKING UNIVERSITY  
STANFORD UNIVERSITY

UNIVERSIDAD DE LOS ANDES  
UNIV. OF ARIZONA  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, IRVINE  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF HAWAII  
UNIV. OF MONTANA  
UNIV. OF NEVADA, RENO  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no liability for its contents or policies.

See inside back cover or [www.pjmath.org](http://www.pjmath.org) for submission instructions.

Regular subscription rate for 2007: \$425.00 a year (10 issues). Special rate: \$212.50 a year to individual members of supporting institutions. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 351 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt für Mathematik, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840 is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and at additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2014 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 231 No. 2 June 2007

---

The Euclidean rank of Hilbert geometries	257
OLIVER BLETZ-SIEBERT and THOMAS FOERTSCH	
A volumish theorem for the Jones polynomial of alternating knots	279
OLIVER T. DASBACH and XIAO-SONG LIN	
On the local Nirenberg problem for the $Q$ -curvatures	293
PHILIPPE DELANOË and FRÉDÉRIC ROBERT	
Knot colouring polynomials	305
MICHAEL EISERMANN	
Some new simple modular Lie superalgebras	337
ALBERTO ELDUQUE	
Subfactors from braided $C^*$ tensor categories	361
JULIANA ERLIJMAN and HANS WENZL	
An elementary, explicit, proof of the existence of Quot schemes of points	401
TROND STØLEN GUSTAVSEN, DAN LAKSOV and ROY MIKAEL SKJELNES	
Symplectic energy and Lagrangian intersection under Legendrian deformations	417
HAI-LONG HER	
Harmonic nets in metric spaces	437
JÜRGEN JOST and LEONARD TODJIHOUNDE	
The quantitative Hopf theorem and filling volume estimates from below	445
LUOFEI LIU	
On the variation of a series on Teichmüller space	461
GREG MCSHANE	
On the geometric and the algebraic rank of graph manifolds	481
JENNIFER SCHULTENS and RICHARD WEIDMAN	



0030-8730(200706)231:2;1-F