

*Pacific  
Journal of  
Mathematics*

**VANISHING SECTIONAL CURVATURE ON THE BOUNDARY  
AND A CONJECTURE OF SCHROEDER AND STRAKE**

FENGBO HANG AND XIAODONG WANG

Volume 232 No. 2

October 2007



## VANISHING SECTIONAL CURVATURE ON THE BOUNDARY AND A CONJECTURE OF SCHROEDER AND STRAKE

FENGBO HANG AND XIAODONG WANG

**We prove some rigidity results for compact manifolds with boundary. For a compact Riemannian manifold with nonnegative Ricci curvature and simply connected mean convex boundary, we show that if the sectional curvature vanishes on the boundary, the metric must be flat.**

Schroeder and Strake [1989, Theorem 1] proved a rigidity theorem:

**Theorem.** *Let  $(M, g)$  be a compact Riemannian manifold with convex boundary and nonnegative Ricci curvature. Assume that the sectional curvature is identically zero in some neighborhood  $U$  of  $\partial M$  and that one of these conditions holds:*

- $\partial M$  is simply connected;
- $\dim \partial M$  is even and  $\partial M$  is strictly convex at some point  $p \in \partial M$ .

*Then  $M$  is flat.*

As they remarked, the condition that the metric is flat in a whole neighborhood of  $\partial M$  is very strong. They conjectured that it suffices to only assume that the sectional curvature vanishes on  $\partial M$  and proved this in the special case of a convex metric ball. Xia [1997; 2002] studied the problem and confirmed the conjecture under various additional conditions, such as the boundary has constant mean curvature or constant scalar curvature, or the second fundamental form satisfies some pinching condition. We refer to Xia's papers for the precise statements. Here we present some results related to the conjecture.

**Theorem 1.** *Let  $M$  be a smooth compact connected Riemannian manifold with boundary and nonnegative Ricci curvature. If every component of  $\partial M$  is simply connected and has nonnegative mean curvature and the sectional curvature of  $M$  vanishes on  $\partial M$ , then  $M$  is flat and  $\partial M$  has only one component.*

---

*MSC2000:* primary 53C24; secondary 53C21.

*Keywords:* rigidity, nonnegative Ricci curvature, mean convex boundary, Reilly's formula.

F. Hang is partially supported by NSF grant DMS-0501050 and a Sloan Research Fellowship. X. Wang is partially supported by NSF grant DMS-0505645.

Therefore when  $\partial M$  is simply connected, the conjecture of Schroeder and Strake is true. Moreover, one only needs  $\partial M$  to be mean convex instead of convex. That  $\partial M$  has only one component follows from theorems in [Ichida 1981; Kasue 1983]. Below we present a different argument for it based on Reilly’s formula [1977].

To continue, we need some notation. We will often write  $\langle \cdot, \cdot \rangle$  for the metric on  $M$  and denote its connection by  $D$ . For convenience we write  $\Sigma = \partial M$  and denote the Levi-Civita connection, curvature tensor, and so on of the induced metric on  $\Sigma$  by adding the subscript  $\Sigma$  to the standard notations. Let  $\nu$  be the unit outer normal vector. The shape operator is given by  $A(X) = D_X \nu$  and the second fundamental form is given by  $h(X, Y) = \langle A(X), Y \rangle = \langle D_X \nu, Y \rangle$ , where  $X, Y \in T\Sigma$ . The mean curvature  $H = \text{tr } A$ . Recall Reilly’s formula [1977, formula (14)] for a smooth function  $u$  on  $M$

$$\begin{aligned} \frac{1}{2} \int_M \left( (\Delta u)^2 - |D^2 u|^2 \right) d\mu &= \frac{1}{2} \int_M Rc(\nabla u, \nabla u) d\mu + \int_\Sigma \Delta_\Sigma u \cdot \frac{\partial u}{\partial \nu} dS \\ &\quad + \frac{1}{2} \int_\Sigma H \left( \frac{\partial u}{\partial \nu} \right)^2 dS + \frac{1}{2} \int_\Sigma \langle A(\nabla_\Sigma u), \nabla_\Sigma u \rangle dS. \end{aligned}$$

A special case of theorems in [Ichida 1981; Kasue 1983] claims that if  $M^n$  is a compact connected Riemannian manifold with mean convex boundary  $\Sigma$  and nonnegative Ricci curvature, then  $\Sigma$  has at most two components; moreover if  $\Sigma$  has two components, then  $M$  is isometric to  $\Gamma \times [0, a]$  for some connected compact Riemannian manifold  $\Gamma$  with nonnegative Ricci curvature and  $a > 0$ . For Theorem 1, it is clear that  $M$  cannot have the product metric; hence  $\Sigma$  has one component. Interestingly, one may give an argument for the above special case based on Reilly’s formula. Indeed, assume  $\Sigma$  is not connected, and fix a component  $\Sigma_0$  of  $\Sigma$ ; then we may solve the Dirichlet problem

$$\Delta u = 0 \text{ on } M, \quad u|_{\Sigma_0} = 0, \quad u|_{\Sigma \setminus \Sigma_0} = 1.$$

Applying Reilly’s formula to  $u$ , we get

$$-\int_M |D^2 u|^2 d\mu = \int_M Rc(\nabla u, \nabla u) d\mu + \int_\Sigma H \left( \frac{\partial u}{\partial \nu} \right)^2 dS.$$

Hence  $D^2 u = 0$ . This implies  $|\nabla u| \equiv c > 0$ . Since  $\nabla u = -c\nu$  on  $\Sigma_0$  and  $\nabla u = c\nu$  on  $\Sigma \setminus \Sigma_0$ , we see  $D_X \nu = 0$  for  $X \in T\Sigma$ , that is,  $\Sigma$  is totally geodesic. If we look at the flow generated by  $\nabla u/c$ , then it sends  $\Sigma_0$  to  $\Sigma \setminus \Sigma_0$  at time  $1/c$  and hence  $\Sigma$  has exactly two components. Note that the flow lines are just geodesics. If we fix a coordinate on  $\Sigma_0$ , namely  $\theta^1, \dots, \theta^{n-1}$  and let  $r = u/c$ , then we have  $g = dr \otimes dr + g_{ij}(r, \theta) d\theta^i \otimes d\theta^j$ . Using  $D^2 r = 0$ , we see  $\partial_r g_{ij}(r, \theta) = 0$ . Hence  $M$  is isometric to  $\Sigma_0 \times [0, 1/c]$ .

Under the assumption of Theorem 1 that the sectional curvature of  $M$  vanishes on  $\Sigma$ , it follows from the Gauss and Codazzi equations that

$$R_{\Sigma}(X, Y, Z, W) = h(X, Z)h(Y, W) - h(X, W)h(Y, Z),$$

$$(D_{\Sigma})_X h(Y, Z) = (D_{\Sigma})_Y h(X, Z),$$

where  $X, Y, Z$ , and  $W$  belong to  $T\Sigma$ . By the fundamental theorem for hypersurfaces [Spivak 1999, part (2) of Theorem 21 on p. 63] and that  $\Sigma$  is simply connected, we may find a smooth isometric immersion  $\phi : \Sigma \rightarrow \mathbb{R}^n$  such that the second fundamental form of the immersion has  $h_{\phi} = h$ . If  $\Sigma$  is convex, then  $\phi$  is an embedding by a Hadamard-type theorem of Sacksteder [1960]. With this immersion  $\phi$  in hand, Theorem 1 follows from:

**Proposition 2.** *Let  $M^n$  be a smooth compact connected Riemannian manifold with connected boundary  $\Sigma = \partial M$  and  $Rc \geq 0$ . If  $\phi : \Sigma \rightarrow \mathbb{R}^l$  is an isometric immersion with  $|H_{\phi}| \leq H$  on  $\Sigma$ , where  $H_{\phi}$  is the mean curvature vector of the immersion  $\phi$ , then  $M$  is flat. If  $\phi$  is also an imbedding,  $M$  is isometric to a domain in  $\mathbb{R}^n$ .*

This generalizes [Ros 1988, Theorem 2], a congruence theorem for hypersurface in Euclidean space. Following the Ros’s argument, we will show by Reilly’s formula that the harmonic extension of the map  $\phi$  is in fact an isometric immersion.

*Proof.* We can find a smooth function  $F : M \rightarrow \mathbb{R}^l$  such that  $\Delta F = 0$  in  $M$  and  $F|_{\Sigma} = \phi$ . Applying Reilly’s formula to each component of  $F$  and summing yields

$$-\frac{1}{2} \sum_{\alpha} \int_M |D^2 F^{\alpha}|^2 d\mu$$

$$= \frac{1}{2} \int_M \sum_{\alpha} Rc(\nabla F^{\alpha}, \nabla F^{\alpha}) d\mu + \int_{\Sigma} \sum_{\alpha} \Delta_{\Sigma} \phi^{\alpha} \cdot \frac{\partial F^{\alpha}}{\partial \nu} dS$$

$$+ \frac{1}{2} \int_{\Sigma} \sum_{\alpha} H \left( \frac{\partial F^{\alpha}}{\partial \nu} \right)^2 dS + \frac{1}{2} \int_{\Sigma} \sum_{\alpha} \langle A(\nabla_{\Sigma} \phi^{\alpha}), \nabla_{\Sigma} \phi^{\alpha} \rangle dS.$$

Note that

$$\sum_{\alpha} \langle A(\nabla_{\Sigma} \phi^{\alpha}), \nabla_{\Sigma} \phi^{\alpha} \rangle = \langle Ae_i, e_j \rangle e_i \phi^{\alpha} \cdot e_j \phi^{\alpha} = \langle Ae_i, e_j \rangle \phi_* e_i \cdot \phi_* e_j$$

$$= \langle Ae_i, e_j \rangle \delta_{ij} = \text{tr } A = H,$$

where  $e_1, \dots, e_{n-1}$  is a local orthonormal frame on  $\Sigma$ ; hence

$$0 = \frac{1}{2} \sum_{\alpha} \int_M |D^2 F^{\alpha}|^2 d\mu + \frac{1}{2} \int_M \sum_{\alpha} Rc(\nabla F^{\alpha}, \nabla F^{\alpha}) d\mu$$

$$+ \int_{\Sigma} H_{\phi} \cdot F_* \nu dS + \frac{1}{2} \int_{\Sigma} H |F_* \nu|^2 dS + \frac{1}{2} \int_{\Sigma} H dS$$

$$\geq \frac{1}{2} \sum_{\alpha} \int_M |D^2 F^{\alpha}|^2 d\mu + \frac{1}{2} \int_{\Sigma} H (|F_* \nu|^2 - 2|F_* \nu| + 1) dS.$$

Hence  $D^2 F^{\alpha} = 0$  for all  $\alpha$ . It follows that  $F^* g_{\mathbb{R}^l}$  is parallel on  $M$ . We may find some  $p \in \Sigma$  such that  $|H_{\phi}| > 0$  at  $p$ ; hence  $H(p) > 0$ . From the above argument, this implies  $|F_* \nu| = 1$  at  $p$  and  $F_* \nu$  is perpendicular to  $\phi_* \Sigma_p$ ; hence  $F^* g_{\mathbb{R}^l} = g_M$  at  $p$ . It follows that  $F^* g_{\mathbb{R}^l} = g_M$  on  $M$ , that is,  $F$  is an isometric immersion and  $M$  is flat. Now assume  $\phi$  is an imbedding. Let  $\bar{D}$  be the connection on  $\mathbb{R}^l$ , then  $\bar{D}_X F_* Y - F_* D_X Y = XYF - (D_X Y) F = 0$ . It follows that  $F : M \rightarrow \mathbb{R}^l$  is a totally geodesic submanifold, and hence the image lies in a  $n$  dimensional affine subspace. Without loss of generality, we may assume  $l = n$  and  $\Sigma$  is a compact hypersurface in  $\mathbb{R}^n$ . Then there exists a bounded open domain  $\Omega$  such that  $\partial\Omega = \Sigma$ . Since  $F$  is an immersion, we see  $F(M) \setminus \bar{\Omega}$  is both open and closed in  $\mathbb{R}^n \setminus \bar{\Omega}$ ; hence it must be empty. Based on this, we show  $F : M \rightarrow \bar{\Omega}$  is a covering map, and hence it must be a diffeomorphism.  $\square$

If we assume that  $\partial M$  is convex, then it is clear from the above that  $M$  is isometric to a convex domain in  $\mathbb{R}^n$ . In fact, in this case one may replace the nonnegativity of the Ricci curvature by the much weaker nonnegativity of the scalar curvature, at least when  $M$  is spin.

**Theorem 3.** *Let  $M$  be a smooth compact connected Riemannian manifold with boundary and nonnegative scalar curvature. If  $M$  is spin, each component of  $\partial M$  is convex and simply connected, and the sectional curvature of  $M$  vanishes on  $\partial M$ , then  $M$  is isometric to a convex domain in  $\mathbb{R}^n$ .*

*Proof.* For every component  $\Gamma$  of  $\partial M$ , we have an isometric embedding  $\phi : \Gamma \rightarrow \mathbb{R}^n$  that has  $h$  as the second fundamental form. Let  $\Omega$  be the convex domain enclosed by  $\phi(\Gamma)$ . We glue  $M$  and  $\mathbb{R}^n \setminus \Omega$  along  $\Gamma$  via the diffeomorphism  $\phi$  for all the  $\Gamma$ 's and obtain a complete Riemannian manifold  $N$  that has nonnegative scalar curvature and is flat outside a compact set. Note that the metric is  $C^1$  along the gluing hypersurface. Since  $M$  is spin, we conclude by the generalized positive mass theorem [Shi and Tam 2002, Theorem 3.1] that  $N$  is isometric to  $\mathbb{R}^n$ . It follows that  $M$  is isometric to a convex domain in  $\mathbb{R}^n$ .  $\square$

## References

- [Ichida 1981] R. Ichida, "Riemannian manifolds with compact boundary", *Yokohama Math. J.* **29**:2 (1981), 169–177. MR 83i:53062 Zbl 0493.53033
- [Kasue 1983] A. Kasue, "Ricci curvature, geodesics and some geometric properties of Riemannian manifolds with boundary", *J. Math. Soc. Japan* **35**:1 (1983), 117–131. MR 84g:53068 Zbl 0494.53039
- [Reilly 1977] R. C. Reilly, "Applications of the Hessian operator in a Riemannian manifold", *Indiana Univ. Math. J.* **26**:3 (1977), 459–472. MR 57 #13799 Zbl 0391.53019

- [Ros 1988] A. Ros, “Compact hypersurfaces with constant scalar curvature and a congruence theorem”, *J. Differential Geom.* **27**:2 (1988), 215–223. With an appendix by Nicholas J. Korevaar. MR 89b:53096 Zbl 0638.53051
- [Sacksteder 1960] R. Sacksteder, “On hypersurfaces with no negative sectional curvatures”, *Amer. J. Math.* **82** (1960), 609–630. MR 22 #7087 Zbl 0194.22701
- [Schroeder and Strake 1989] V. Schroeder and M. Strake, “Rigidity of convex domains in manifolds with nonnegative Ricci and sectional curvature”, *Comment. Math. Helv.* **64**:2 (1989), 173–186. MR 90h:53042 Zbl 0692.53020
- [Shi and Tam 2002] Y. Shi and L.-F. Tam, “Positive mass theorem and the boundary behaviors of compact manifolds with nonnegative scalar curvature”, *J. Differential Geom.* **62**:1 (2002), 79–125. MR 2005b:53046 Zbl 1071.53018
- [Spivak 1999] M. Spivak, *A comprehensive introduction to differential geometry*, vol. 4, 3 ed., Publish or Perish, Houston, Tex., 1999.
- [Xia 1997] C. Xia, “Rigidity of compact manifolds with boundary and nonnegative Ricci curvature”, *Proc. Amer. Math. Soc.* **125**:6 (1997), 1801–1806. MR 97i:53043 Zbl 0882.53028
- [Xia 2002] C. Xia, “On a conjecture by Schroeder and Strake”, *Q. J. Math.* **53**:1 (2002), 119–124. MR 2003a:53042 Zbl 1016.53033

Received September 4, 2006. Revised February 2, 2007.

FENGBO HANG  
DEPARTMENT OF MATHEMATICS  
PRINCETON UNIVERSITY  
FINE HALL, WASHINGTON ROAD  
PRINCETON, NJ 08544  
UNITED STATES  
fhang@math.princeton.edu  
<http://www.math.princeton.edu/~fhang>

XIAODONG WANG  
DEPARTMENT OF MATHEMATICS  
MICHIGAN STATE UNIVERSITY  
EAST LANSING, MI 48824  
UNITED STATES  
xwang@math.msu.edu  
<http://www.math.msu.edu/~xwang/>

