

*Pacific  
Journal of  
Mathematics*

UNITARY REPRESENTATIONS OF THE EXTENDED AFFINE  
LIE ALGEBRA  $\widehat{\mathfrak{gl}}_3(\mathbb{C}_q)$

ZITING ZENG

Volume 233 No. 2

December 2007



## UNITARY REPRESENTATIONS OF THE EXTENDED AFFINE LIE ALGEBRA $\widehat{\mathfrak{gl}}_3(\mathbb{C}_q)$

ZITING ZENG

**We present modules for the extended affine Lie algebra  $\widehat{\mathfrak{gl}}_3(\mathbb{C}_q)$  by using the idea of free fields. We give a necessary and sufficient condition for the modules to be unitary.**

### 1. Introduction

Extended affine Lie algebras are a higher dimensional generalization of affine Kac–Moody Lie algebras introduced by Høegh-Krohn and Torr sani [1990] and systematically studied in [Allison et al. 1997; Berman et al. 1996]. It turns out that any extended affine Lie algebra of type  $A$  can be coordinatized by a quantum torus (or a nonassociative torus for some small rank cases). Representations for extended affine Lie algebras coordinatized by quantum tori and Lie algebras related to quantum tori have been studied in [Jakobsen and Kac 1989; Berman and Szmigielski 1999; Gao 2000a; 2000b; 2002; Eswara Rao 2004; 2003; Eswara Rao and Batra 2002; Gao and Zeng 2006; Eswara Rao and Zhao 2004; Lin and Tan 2004; 2006; Golenishcheva-Kutuzova and Kac 1998; Varagnolo and Vasserot 1998; Miki 2004; Zhang and Zhao 1996; Billig and Zhao 2004; Su and Zhu 2005; Lau 2005; Baranovsky et al. 2000] and elsewhere.

Wakimoto’s free fields construction [1986] provides a remarkable way to realize affine Kac–Moody Lie algebras; see also [Feigin and Frenkel 1990; Etingof et al. 1998]. In [Gao and Zeng 2006], we used Wakimoto’s idea to construct a class of representations for  $\widehat{\mathfrak{gl}}_2(\mathbb{C}_q)$  and found the necessary and sufficient condition for the representations to be unitary. Here, we will continue to construct representations for  $\widehat{\mathfrak{gl}}_3(\mathbb{C}_q)$ . As witnessed in [Feigin and Frenkel 1990], the realization for  $\widehat{\mathfrak{gl}}_3(\mathbb{C}_q)$  is much more subtle and complicated than the one for  $\widehat{\mathfrak{gl}}_2(\mathbb{C}_q)$ . This construction for  $\widehat{\mathfrak{gl}}_3(\mathbb{C}_q)$  might shed light on the general construction for  $\widehat{\mathfrak{gl}}_n(\mathbb{C}_q)$  with  $n \geq 4$ .

We then construct a hermitian form and determine when the form is positive definite (so the representations are unitary). Unlike [Gao and Zeng 2006], in which we defined the form on the monomial basis for the module (this idea goes back to [Wakimoto 1985]), we define the form directly on the basis consisting of certain

---

*MSC2000:* primary 06B15; secondary 81R50.

*Keywords:* extended affine Lie algebra, hermitian form, unitary representation.

iterated module actions on a “highest weight vector” 1. This facilitates verifying that the defined form is hermitian.

Let  $q$  be a nonzero complex number. A quantum 2-torus associated to  $q$  [Manin 1991] is the unital associative  $\mathbb{C}$ -algebra  $\mathbb{C}_q[s^{\pm 1}, t^{\pm 1}]$  (or, simply  $\mathbb{C}_q$ ) with generators  $s^{\pm 1}, t^{\pm 1}$  and relations

$$ss^{-1} = s^{-1}s = tt^{-1} = t^{-1}t = 1 \quad \text{and} \quad ts = qst.$$

Define a  $\mathbb{C}$ -linear function  $\kappa : \mathbb{C}_q \rightarrow \mathbb{C}$  by

$$\kappa(s^m t^n) = \delta_{(m,n), (0,0)}.$$

Let  $d_s, d_t$  be the degree operators on  $\mathbb{C}_q$  defined for  $m, n \in \mathbb{Z}$  by

$$d_s(s^m t^n) = ms^m t^n, \quad d_t(s^m t^n) = ns^m t^n.$$

Let  $\mathfrak{gl}_3(\mathbb{C}_q)$  be the Lie algebra of 3 by 3 matrices with entries in  $\mathbb{C}_q$ . We form a natural central extension of  $\mathfrak{gl}_3(\mathbb{C}_q)$  as

$$\widehat{\mathfrak{gl}_3(\mathbb{C}_q)} = \mathfrak{gl}_3(\mathbb{C}_q) \oplus \mathbb{C}c_s \oplus \mathbb{C}c_t$$

with Lie bracket

$$\begin{aligned} (1-1) \quad [E_{ij}(s^{m_1} t^{n_1}), E_{kl}(s^{m_2} t^{n_2})] \\ = \delta_{jk} q^{n_1 m_2} E_{il}(s^{m_1+m_2} t^{n_1+n_2}) - \delta_{il} q^{n_2 m_1} E_{kj}(s^{m_1+m_2} t^{n_1+n_2}) \\ + m_1 q^{n_1 m_2} \delta_{jk} \delta_{il} \delta_{m_1+m_2, 0} \delta_{n_1+n_2, 0} c_s + n_1 q^{n_1 m_2} \delta_{jk} \delta_{il} \delta_{m_1+m_2, 0} \delta_{n_1+n_2, 0} c_t \end{aligned}$$

for  $m_1, m_2, n_1, n_2 \in \mathbb{Z}$  and  $1 \leq i, j, k, l \leq 3$ , where  $E_{ij}(s^m t^n)$  is the matrix whose only nonzero entry is  $s^m t^n$  at the  $(i, j)$  position, and  $c_s$  and  $c_t$  are central elements of  $\widehat{\mathfrak{gl}_3(\mathbb{C}_q)}$ .

The derivations  $d_s$  and  $d_t$  can be extended to derivations on  $\mathfrak{gl}_3(\mathbb{C}_q)$ . Now we can define the semidirect product of the Lie algebra  $\widehat{\mathfrak{gl}_3(\mathbb{C}_q)}$  and those derivations:

$$\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)} = \widehat{\mathfrak{gl}_3(\mathbb{C}_q)} \oplus \mathbb{C}d_s \oplus \mathbb{C}d_t.$$

The Lie algebra  $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$  is an extended affine Lie algebra of type  $A_2$  with nullity 2. See [Allison et al. 1997] and [Berman et al. 1996] for definitions.

### 2. Module for $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$

In this section, we use Wakimoto’s idea [1985] to construct  $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$ -modules as was done in [Gao and Zeng 2006].

Let  $\mathbb{K}_1 = \{(3m + 1, 3n + 1) \mid m, n \in \mathbb{Z}\}$  and  $\mathbb{K}_{-1} = \{(3m - 1, 3n - 1) \mid m, n \in \mathbb{Z}\}$  so that  $\mathbb{K}_{-1} = -\mathbb{K}_1$ . If  $A = (3m + 1, 3n + 1) \in \mathbb{K}_1$ , we always write  $A_1 = m$  and

$A_2 = n$ , and similarly, if  $\mathbf{B} = (3m - 1, 3n - 1) \in \mathbb{K}_{-1}$ , then  $\mathbf{B}_1 = m$  and  $\mathbf{B}_2 = n$ . Let

$$V = \mathbb{C}[x_A, x_B : A \in \mathbb{K}_1, \mathbf{B} \in \mathbb{K}_{-1}]$$

be the (commutative) polynomial ring of infinitely many variables. The operators  $x_{(m,n)}$  and  $\partial/\partial x_{(m,n)}$  act on  $V$  by the usual multiplication and differentiation.

Form a family of  $2 \times 2$  lower triangular matrices

$$X_{m,n} = \begin{pmatrix} a_{(m,n)} & 0 \\ c_{(m,n)} & d_{(m,n)} \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$$

for  $(m, n) \in \mathbb{K}_1 \cup \mathbb{K}_{-1}$  (so that  $a_{(m,n)}d_{(m,n)} = 1$ ). Set

$$P_{(m,n)} = a_{(m,n)} \frac{\partial}{\partial x_{(m,n)}} \quad \text{and} \quad Q_{(m,n)} = c_{(m,n)} \frac{\partial}{\partial x_{(m,n)}} + d_{(m,n)} x_{(m,n)}$$

for  $(m, n) \in \mathbb{K}_1 \cup \mathbb{K}_{-1}$ . Then for  $A, A' \in \mathbb{K}_1$  and  $\mathbf{B}, \mathbf{B}' \in \mathbb{K}_{-1}$ ,

$$[P_A, P_{A'}] = [Q_A, Q_{A'}] = [P_{\mathbf{B}}, P_{\mathbf{B}'}] = [Q_{\mathbf{B}}, Q_{\mathbf{B}'}] = 0$$

$$[P_A, P_{\mathbf{B}}] = [P_A, Q_{\mathbf{B}}] = [Q_A, Q_{\mathbf{B}}] = [P_{\mathbf{B}}, Q_A] = 0$$

$$[P_A, Q_{A'}] = \delta_{A,A'}$$

$$[P_{\mathbf{B}}, Q_{\mathbf{B}'}] = \delta_{\mathbf{B},\mathbf{B}'}$$

Fixing a complex number  $\mu$ , we define operators on  $V$ . For the rest of this paper, sums involving  $A$  and  $A'$  (or any other decorated  $A$ ) will range over  $\mathbb{K}_1$ . Similarly, the  $\mathbf{B}$ 's range over  $\mathbb{K}_{-1}$ . That is, we write  $\sum_{A \in \mathbb{K}_1}$  simply as  $\sum_A$ , and so on.

$$\begin{aligned} e_{21}^{(\mu)}(m_1, n_1) &= -q^{-m_1 n_1} \mu P_{-(3m_1-1, 3n_1-1)} \\ &\quad - \sum_{A, A'} q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1, 3n_1-1)} P_A P_{A'} \\ &\quad - \sum_{A, \mathbf{B}} q^{n_1 A_1 + \mathbf{B}_2 m_1 + \mathbf{B}_2 A_1} Q_{A+\mathbf{B}+(3m_1-1, 3n_1-1)} P_A P_{\mathbf{B}}, \end{aligned}$$

$$e_{12}^{(\mu)}(m_1, n_1) = Q_{(3m_1+1, 3n_1+1)},$$

$$e_{11}^{(\mu)}(m_1, n_1) = \frac{1}{2} \mu \delta_{(m_1, n_1), (0,0)} + \sum_A q^{A_1 n_1} Q_{A+(3m_1, 3n_1)} P_A,$$

$$\begin{aligned} e_{22}^{(\mu)}(m_1, n_1) &= -\frac{1}{2} \mu \delta_{(m_1, n_1), (0,0)} \\ &\quad - \sum_A q^{A_2 m_1} Q_{A+(3m_1, 3n_1)} P_A - \sum_{\mathbf{B}} q^{\mathbf{B}_2 m_1} Q_{\mathbf{B}+(3m_1, 3n_1)} P_{\mathbf{B}}, \end{aligned}$$

$$\begin{aligned} e_{23}^{(\mu)}(m_1, n_1) &= -q^{-m_1 n_1} \mu P_{-(3m_1+1, 3n_1+1)} \\ &\quad - \sum_{A, \mathbf{B}} q^{n_1 \mathbf{B}_1 + A_2 m_1 + A_2 \mathbf{B}_1} Q_{A+\mathbf{B}+(3m_1+1, 3n_1+1)} P_A P_{\mathbf{B}} \\ &\quad - \sum_{\mathbf{B}, \mathbf{B}'} q^{n_1 \mathbf{B}'_1 + \mathbf{B}_2 m_1 + \mathbf{B}_2 \mathbf{B}'_1} Q_{\mathbf{B}+\mathbf{B}'+(3m_1+1, 3n_1+1)} P_{\mathbf{B}} P_{\mathbf{B}'}, \end{aligned}$$

$$\begin{aligned}
 e_{32}^{(\mu)}(m_1, n_1) &= Q_{(3m_1-1, 3n_1-1)}, \\
 e_{31}^{(\mu)}(m_1, n_1) &= \sum_A q^{A_1 n_1} Q_{A+(3m_1-2, 3n_1-2)} P_A, \\
 e_{13}^{(\mu)}(m_1, n_1) &= \sum_B q^{B_1 n_1} Q_{B+(3m_1+2, 3n_1+2)} P_B, \\
 e_{33}^{(\mu)}(m_1, n_1) &= \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} + \sum_B q^{B_1 n_1} Q_{B+(3m_1, 3n_1)} P_B, \\
 D_1^{(\mu)} &= \sum_A A_1 Q_A P_A + \sum_B B_1 Q_B P_B, \\
 D_2^{(\mu)} &= \sum_A A_2 Q_A P_A + \sum_B B_2 Q_B P_B.
 \end{aligned}$$

Although the operators are infinite sums, they are well defined as operators on  $V$ . Now we have the following result:

**Theorem 2.1.** *The linear map  $\pi : \widetilde{\mathfrak{gl}}_3(\mathbb{C}_q) \rightarrow \text{End}V$  given by*

$$\begin{aligned}
 \pi(E_{ij}(s^{m_1} t^{n_1})) &= e^{(\mu)}_{ij}(m_1, n_1), & \pi(d_s) &= D_1^{(\mu)}, & \pi(c_s) &= 0, \\
 & & \pi(d_t) &= D_2^{(\mu)}, & \pi(c_t) &= 0,
 \end{aligned}$$

for  $m_1, n_1 \in \mathbb{Z}$  and  $1 \leq i, j \leq 3$  is a Lie algebra homomorphism.

*Proof.* The proof is straightforward. However, we will provide a few details. It suffices to check the Lie bracket (1-1). We will do this systematically so that we won't miss any cases.

First, we have

$$\begin{aligned}
 & [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{11}(m_2, n_2)] \\
 &= \left[ \sum_A q^{A_1 n_1} Q_{(3m_1, 3n_1)+A} P_A, \sum_{A'} q^{A'_1 n_1} Q_{(3m_2, 3n_2)+A'} P_{A'} \right] \\
 &= \sum_{A'} q^{(m_2+A'_1)n_1+A'_1 n_2} Q_{(3m_1, 3n_1)+(3m_2+3n_2)+A'} P_{A'} + \frac{1}{2} q^{m_2 n_1} \mu \delta_{(m_1+m_2, n_1+n_2), (0, 0)} \\
 &\quad - \sum_A q^{A_1 n_1+(m_1+A_1)n_2} q_{(3m_1, 3n_1)+(3m_2, 3n_2)+A} P_A - \frac{1}{2} q^{m_1 n_2} \mu \delta_{(m_1+m_2, n_1+n_2), (0, 0)} \\
 &= q^{m_2 n_1} e^{(\mu)}_{11}(m_1 + m_2, n_1 + n_2) - q^{m_1 n_2} e^{(\mu)}_{11}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

The next two brackets are easy.

$$\begin{aligned}
 [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{12}(m_2, n_2)] &= q^{m_2 n_1} e^{(\mu)}_{12}(m_1 + m_2, n_1 + n_2), \\
 [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{13}(m_2, n_2)] &= q^{m_2 n_1} e^{(\mu)}_{13}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

Next,

$$\begin{aligned}
& [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{21}(m_2, n_2)] \\
&= \sum_A (-\mu) q^{A_1 n_1 - m_2 n_2} [Q_{A+(3m_1, 3n_1)} P_A, P_{-(3m_2-1, 3n_2-1)}] \\
&\quad - \sum_{A, \bar{A}, \bar{A}'} q^{A_1 n_1 + \bar{A}'_1 n_2 + \bar{A}_2 m_2 + \bar{A}_2 \bar{A}'_1} [Q_{A+(3m_1, 3n_1)} P_A, Q_{\bar{A}+\bar{A}'+(3m_2-1, 3n_2-1)} P_{\bar{A}} P_{\bar{A}'}] \\
&\quad - \sum_{A, \bar{A}, B} q^{A_1 n_1 + n_2 \bar{A}_1 + B_2 m_2 + B_2 \bar{A}_1} [Q_{A+(3m_1, 3n_1)} P_A, Q_{\bar{A}+B+(3m_2-1, 3n_2-1)} P_{\bar{A}} P_B] \\
&= \mu q^{-(m_1+m_2)(n_1+n_2)+m_1 n_2} P_{-(3(m_1+m_2)-1, 3(n_1+n_2)-1)} \\
&\quad - \sum_{\bar{A}, \bar{A}'} q^{(\bar{A}_1+\bar{A}'_1+m_2)n_1+n_2 \bar{A}'_1+\bar{A}_2 m_2+\bar{A}_2 \bar{A}'_1} Q_{(3m_1, 3n_1)+\bar{A}+\bar{A}'+(3m_2-1, 3n_2-1)} P_{\bar{A}} P_{\bar{A}'} \\
&\quad + \sum_{A, \bar{A}} q^{A_1 n_1 + n_2 (m_1+A_1) + \bar{A}_2 m_2 + \bar{A}_2 (m_1+A_1)} Q_{\bar{A}+(3m_1, 3n_1)+A+(3m_2-1, 3n_2-1)} P_{\bar{A}} P_A \\
&\quad + \sum_{A, \bar{A}'} q^{A_1 n_1 + n_2 A'_1 + (n_1+A_2)m_2 + (n_1+A_2)\bar{A}'_1} Q_{A+(3m_1, 3n_1)+\bar{A}'+(3m_2-1, 3n_2-1)} P_{\bar{A}'} P_A \\
&\quad + \sum_{A, B} q^{A_1 n_1 + n_2 (m_1+A_1) + bdb_2 m_2 + B_2 (m_1+A_1)} Q_{A+(3m_1, 3n_1)+B+(3m_2-1, 3n_2-1)} P_B P_A,
\end{aligned}$$

(using that the second and fourth term cancel each other)

$$\begin{aligned}
&= -q^{m_1 n_2} (-\mu q^{-(m_1+m_2)(n_1+n_2)} P_{-(3(m_1+m_2)-1, 3(n_1+n_2)-1)} \\
&\quad - \sum_{A, \bar{A}} q^{A(n_1+n_2)+\bar{A}_2(m_1+m_2)+A_1 \bar{A}_2} Q_{(3(m_1+m_2)-1, 3(n_1+n_2)-1)+A+\bar{A}} P_{\bar{A}} P_A \\
&\quad - \sum_{A, B} q^{A_1(n_1+n_2)+B_2(m_1+m_2)+B_2+A_1} Q_{A+B+(3(m_1+m_2)-1, 3(n_1+n_2)-1)} P_B P_A) \\
&= -q^{m_1 n_2} e^{(\mu)}_{21}(m_1 + m_2, n_1 + n_2).
\end{aligned}$$

We easily verify the next seven brackets.

$$\begin{aligned}
& [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{22}(m_2, n_2)] = 0, \\
& [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{23}(m_2, n_2)] = 0, \\
& [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{31}(m_2, n_2)] = -q^{m_1 n_2} e^{(\mu)}_{31}(m_1 + m_2, n_1 + n_2), \\
& [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] = 0, \\
& [e^{(\mu)}_{11}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] = 0, \\
& [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{12}(m_2, n_2)] = 0, \\
& [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{13}(m_2, n_2)] = 0.
\end{aligned}$$

Next we have

$$\begin{aligned}
 & [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{21}(m_2, n_2)] \\
 &= \mu q^{-m_2 n_2} \delta_{(m_1, n_1), (-m_2, n_2)} \\
 &\quad + \sum_{A'} q^{n_2 A'_1 + n_1 m_2 + n_1 A'_1} Q_{(3m_1+1, 3n_1+1)+A'+(3m_2-1, 3n_2-1)} P_{A'} \\
 &\quad + \sum_A q^{n_2 m_1 + A_2 m_2 + A_2 m_1} Q_{A+(3m_1+1, 3n_1+1)+(3m_2-1, 3n_2-1)} P_A \\
 &\quad + \sum_B q^{n_2 m_1 + B_2 m_2 + B_2 m_1} Q_{(3m_1+1, 3n_1+1)+B+(3m_2-1, 3n_2-1)} P_B \\
 &= q^{n_1 m_2} \left( \sum_{A'} q^{(n_1+n_2)A'_1} Q_{A'+(3m_1+3m_2, 3n_1+3n_2)} P_{A'} + \frac{1}{2} \delta_{(m_1, n_1), (-m_2, n_2)} \right) \\
 &\quad - q^{n_2 m_1} \left( - \sum_A q^{A_2(m_2+m_1)} Q_{A+(3m_1+3m_2, 3n_1+3n_2)} P_A \right. \\
 &\quad \quad \left. - \sum_B q^{B_2(m_2+m_1)} Q_{B+(3m_1+3m_2, 3n_1+3n_2)} P_B - \frac{1}{2} \delta_{(m_1, n_1), (-m_2, n_2)} \right) \\
 &= q^{n_1 m_2} e^{(\mu)}_{11}(m_1 + m_2, n_1 + n_2) - q^{n_2 m_1} e^{(\mu)}_{22}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

The following six brackets can be checked easily.

$$\begin{aligned}
 [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{22}(m_2, n_2)] &= q^{n_1 m_2} e^{(\mu)}_{12}(m_1 + m_2, n_1 + n_2), \\
 [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{23}(m_2, n_2)] &= q^{n_1 m_2} e^{(\mu)}_{13}(m_1 + m_2, n_1 + n_2), \\
 [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{31}(m_2, n_2)] &= -q^{m_1 n_2} e^{(\mu)}_{32}(m_1 + m_2, n_1 + n_2), \\
 [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] &= 0, \\
 [e^{(\mu)}_{12}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] &= 0, \\
 [e^{(\mu)}_{13}(m_1, n_1), e^{(\mu)}_{13}(m_2, n_2)] &= 0.
 \end{aligned}$$

Next,

$$\begin{aligned}
 & [e^{(\mu)}_{13}(m_1, n_1), e^{(\mu)}_{21}(m_2, n_2)] \\
 &= \left[ \sum_B q^{B_1 n_1} Q_{B+(3m_1+2, 3n_1+2)} P_B, -q^{-m_2 n_2} \mu P_{-(3m_2-1, 3n_2-1)} \right. \\
 &\quad \left. - \sum_{A, A'} q^{n_2 A'_1 + A_2 m_2 + A_2 A'_1} Q_{A+A'+(3m_2-1, 3n_2-1)} P_A P_{A'} \right. \\
 &\quad \quad \left. - \sum_{A, B} q^{n_2 A_1 + B_2 m_2 + B_2 A_1} Q_{A+B+(3m_2-1, 3n_2-1)} P_A P_B \right] \\
 &= - \sum_{A, B} q^{(A_1+B_1+m_2)n_1+n_2+A_1+B_2 m_2+B_2 A_1} Q_{(3m_1+2, 3n_1+2)+A+B+(3m_2-1, 3n_2-1)} P_A P_B \\
 &\quad + \sum_{A, B} q^{n_2(m_1+B_1)+A_2 m_2+A_2(m_1+B_1)+B_1 n_1} Q_{A+(3m_1+2, 3n_1+2)+B+(3m_2-1, 3n_2-1)} P_A P_B \\
 &\quad \quad + q^{-m_2 n_2 + (-m_1 - m_2)n_1} \mu P_{(-3m_1-3m_2-1, -3n_1-3n_2-1)}
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{A', B} q^{n_2 A'_1 + (n_1 + B_2)m_2 + (n_1 + B_2)A'_1 + B_1 n_1} Q_{A' + (3m_1 + 2, 3n_1 + 2) + B + (3m_2 - 1, 3n_2 - 1)} P_{A'} P_B \\
 & + \sum_{B, B'} q^{n_2(B'_1 + m_1) + B_2 m_2 + B_2(B'_1 + m_1) + B'_1 n_1} Q_{B' + (3m_1 + 2, 3n_1 + 2) + B + (3m_2 - 1, 3n_2 - 1)} P_{B'} P_B
 \end{aligned}$$

(using that the first and fourth terms cancel each other)

$$\begin{aligned}
 & = q^{n_2 m_1} \left( q^{-(m_1 + m_2)(n_1 + n_2)} \mu P_{-(3m_1 + 3m_2 - 1, 3n_1 + 3n_2 - 1)} \right. \\
 & \quad + \sum_{A, A'} q^{(n_1 + n_2)A'_1 + A_2(m_1 + m_2) + A_2 A'_1} Q_{A + A' + (3m_1 + 3m_2 - 1, 3n_1 + 3n_2 - 1)} P_A P_{A'} \\
 & \quad \left. - \sum_{A, B} q^{(n_1 + n_2)A_1 + B_2(m_1 + m_2) + B_2 A_1} Q_{A + B + (3m_1 + 3m_2 - 1, 3n_1 + 3n_2 - 1)} P_A P_B \right) \\
 & = -q^{n_2 m_1} e^{(\mu)}_{23}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

The following two brackets are easy.

$$\begin{aligned}
 [e^{(\mu)}_{13}(m_1, n_1), e^{(\mu)}_{22}(m_2, n_2)] & = 0, \\
 [e^{(\mu)}_{13}(m_1, n_1), e^{(\mu)}_{23}(m_2, n_2)] & = 0.
 \end{aligned}$$

Next,

$$\begin{aligned}
 & - [e^{(\mu)}_{13}(m_2, n_2), e^{(\mu)}_{31}(m_1, n_1)] = [e^{(\mu)}_{31}(m_1, n_1), e^{(\mu)}_{13}(m_2, n_2)] \\
 & = \sum_{A, B} q^{n_1 A_1 + n_2 B_1} [Q_{(3m_1 - 2, 3n_1 - 2) + A} P_A, Q_{(3m_2 + 2, 3n_2 + 2) + B} P_B] \\
 & = \sum_B q^{n_2 B_1 + n_1(m_2 + B_1)} Q_{(3m_1 + 3m_2, 3n_1 + 3n_2) + B} P_B \\
 & \quad - \sum_A q^{n_1 A_1 + n_2(m_1 + A_1)} Q_{(3m_1 + 3m_2, 3n_1 + 3n_2) + A} P_A \\
 & = q^{n_1 m_2} \left( \sum_B q^{n_2 B_1 + n_1(m_2 + B_1)} Q_{(3m_1 + 3m_2, 3n_1 + 3n_2) + B} P_B + \frac{1}{2} \mu \delta_{(m_1 + m_2, n_1 + n_2), (0, 0)} \right) \\
 & \quad - q^{n_2 m_1} \left( \sum_A q^{n_1 A_1 + n_2(m_1 + A_1)} Q_{(3m_1 + 3m_2, 3n_1 + 3n_2) + A} P_A + \frac{1}{2} \mu \delta_{(m_1 + m_2, n_1 + n_2), (0, 0)} \right) \\
 & = q^{n_1 m_2} e^{(\mu)}_{33}(m_1 + m_2, n_1 + n_2) - q^{n_2 m_1} e^{(\mu)}_{11}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

The next two brackets are easy.

$$\begin{aligned}
 [e^{(\mu)}_{13}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] & = q^{n_1 m_2} e^{(\mu)}_{12}(m_1 + m_2, n_1 + n_2), \\
 [e^{(\mu)}_{13}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] & = q^{n_1 m_2} e^{(\mu)}_{13}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

$$\begin{aligned}
 & [e^{(\mu)}_{21}(m_1, n_1), e^{(\mu)}_{21}(m_2, n_2)] \\
 & = \mu q^{-m_1 n_1} \sum_A q^{(-m_1 - m_2 - A_1)n_2 + A_2 m_2 + A_2(-m_1 - m_2 - A_1)} P_A P_{(-3m_1 - 3m_2, -3n_1 - 3n_2) - A} \\
 & \quad - \mu q^{-m_2 n_2} \sum_{A'} q^{n_1 A'_1 + (-n_1 - n_2 - A'_2)m_1 + (-n_1 - n_2 - A'_2)A'_1} P_{A'} P_{(-3m_1 - 3m_2, -3n_1 - 3n_2) - A'}
 \end{aligned}$$

$$\begin{aligned}
 &+ \left[ \sum_{A,A'} q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1,3n_1-1)} P_A P_{A'}, \right. \\
 &\qquad \qquad \qquad \left. \sum_{A,A'} q^{n_2 A'_1 + A_2 m_2 + A_2 A'_1} Q_{A+A'+(3m_2-1,3n_2-1)} P_A P_{A'} \right] \\
 &+ \left[ \sum_{A,A'} q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1,3n_1-1)} P_A P_{A'}, \right. \\
 &\qquad \qquad \qquad \left. \sum_{A,B} q^{n_2 A_1 + B_2 m_2 + B_2 A_1} Q_{A+B+(3m_2-1,3n_2-1)} P_A P_B \right] \\
 &+ \left[ \sum_{A,B} q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1,3n_1-1)} P_A P_B, \right. \\
 &\qquad \qquad \qquad \left. \sum_{A,A'} q^{n_2 A'_1 + A_2 m_2 + A_2 A'_1} Q_{A+A'+(3m_2-1,3n_2-1)} P_A P_{A'} \right] \\
 &+ \left[ \sum_{A,B} q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1,3n_1-1)} P_A P_B, \right. \\
 &\qquad \qquad \qquad \left. \sum_{A,B} q^{n_2 A_1 + B_2 m_2 + B_2 A_1} Q_{A+B+(3m_2-1,3n_2-1)} P_A P_B \right]
 \end{aligned}$$

(the first and second terms cancel each other)

$$\begin{aligned}
 &= \sum_{A, \bar{A}, \bar{A}'} q^{n_1(\bar{A}_1 + \bar{A}'_1 + m_2) + A_2 m_1 + A_2(\bar{A}_1 + \bar{A}'_1 + m_2) + n_2 \bar{A}'_1 + \bar{A}_2 m_2 + \bar{A}_2 \bar{A}'_1} \\
 &\qquad \qquad \qquad \times Q_{A+\bar{A}+\bar{A}'+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{\bar{A}} P_{\bar{A}'} \\
 &+ \sum_{A', \bar{A}, \bar{A}'} q^{n_1 A'_1 + (\bar{A}_2 + \bar{A}'_2 + n_2) m_1 + (\bar{A}_2 + \bar{A}'_2 + n_2) A'_1 + n_2 \bar{A}'_1 + \bar{A}_2 m_2 + \bar{A}_2 \bar{A}'_1} \\
 &\qquad \qquad \qquad \times Q_{A'+\bar{A}+\bar{A}'+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_{A'} P_{\bar{A}} P_{\bar{A}'} \\
 &- \sum_{A, A', \bar{A}} q^{n_2(A_1 + A'_1 + m_1) + \bar{A}_2 m_2 + \bar{A}_2(A_1 + A'_1 + m_1) + n_1 A'_1 + A_2 m_1 + A_2 A'_1} \\
 &\qquad \qquad \qquad \times Q_{A+A'+\bar{A}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{A'} P_{\bar{A}} \\
 &- \sum_{A, A', \bar{A}'} q^{n_2 \bar{A}'_1 + (A_2 + A'_2 + n_1) m_2 + (A_2 + A'_2 + n_1) \bar{A}'_1 + n_1 A'_1 + A_2 m_1 + A_2 A'_1} \\
 &\qquad \qquad \qquad \times Q_{A+A'+\bar{A}'+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{A'} P_{\bar{A}'} \\
 &- \sum_{A, A', B} q^{n_2(A_1 + A'_1 + m_1) + B_2 m_2 + B_2(A_1 + A'_1 + m_1) + n_1 A'_1 + A_2 m_1 + A_2 A'_1} \\
 &\qquad \qquad \qquad \times Q_{A+A'+B+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{A'} P_B \\
 &+ \sum_{A, A', B} q^{n_1(A_1 + A'_1 + m_2) + B_2 m_1 + B_2(A_1 + A'_1 + m_2) + n_2 A'_1 + A_2 m_2 + A_2 A'_1} \\
 &\qquad \qquad \qquad \times Q_{A+A'+B+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{A'} P_B \\
 &+ \sum_{A, \bar{A}, B} q^{n_1 A_1 + (\bar{A}_2 + B_2 + n_2) m_1 + (\bar{A}_2 + B_2 + n_2) A_1 + n_2 \bar{A}_1 + B_2 m_2 + B_2 \bar{A}_1} \\
 &\qquad \qquad \qquad \times Q_{A+\bar{A}+B+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{\bar{A}} P_B \\
 &- \sum_{A, \bar{A}, \bar{B}} q^{n_2 A_1 + (\bar{A}_2 + \bar{B}_2 + n_1) m_2 + (\bar{A}_2 + \bar{B}_2 + n_1) A_1 + n_1 \bar{A}_1 + \bar{B}_2 m_1 + \bar{B}_2 \bar{A}_1} \\
 &\qquad \qquad \qquad \times Q_{A+\bar{A}+\bar{B}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)} P_A P_{\bar{A}} P_{\bar{B}} \\
 &= 0,
 \end{aligned}$$

where we used that the first term cancels the fourth, the second term cancels the third, the fifth term cancels the seventh, and the sixth term cancels the eighth. Next,

$$\begin{aligned}
 & - [e^{(\mu)}_{21}(m_2, n_2), e^{(\mu)}_{22}(m_1, n_1)] = [e^{(\mu)}_{22}(m_1, n_1), e^{(\mu)}_{21}(m_2, n_2)] \\
 & = \mu \sum_A q^{A_2 m_1 - m_2 n_2} [Q_{(3m_1, 3n_1) + A} P_A, P_{-(3m_2 - 1, 3n_2 - 1)}] \\
 & \quad + \sum_{A, A', \bar{A}} q^{A_2 m_1 + n_2 A'_1 + \bar{A}_2 m_2 + \bar{A}_2 A'_1} [Q_{(3m_1, 3n_1) + A} P_A, Q_{\bar{A} + A' + (3m_2 - 1, 3n_2 - 1)} P_{A'} P_{\bar{A}}] \\
 & \quad + \sum_{A, \bar{A}, B} q^{A_2 m_1 + n_2 \bar{A}_1 + B_2 m_2 + B_2 \bar{A}_1} [Q_{(3m_1, 3n_1) + A} P_A, Q_{\bar{A} + B + (3m_2 - 1, 3n_2 - 1)} P_A P_B] \\
 & \quad + \sum_{A, B, \bar{B}} q^{B_2 m_1 + n_2 A_1 + \bar{B}_2 m_2 + \bar{B}_2 A_1} [Q_{(3m_1, 3n_1) + B} P_B, Q_{A + \bar{B} + (3m_2 - 1, 3n_2 - 1)} P_A P_{\bar{B}}] \\
 & = -\mu q^{-m_2 n_2 + (-n_1 - n_2)m_1} P_{-3m_1 - 3m_2 + 1, -3n_1 - 3n_2 + 1} \\
 & \quad + \sum_{A, A'} q^{(A_2 + A'_2 + n_2)m_1 + n_2 A'_1 + A_2 m_2 + A_2 A'_1} Q_{(3m_1, 3n_1) + A + A' + (3m_2 - 1, 3n_2 - 1)} P_A P_{A'} \\
 & \quad - \sum_{A, \bar{A}} q^{n_2(m_1 + \bar{A}_1) + A_2 m_2 + A_2(m_1 + \bar{A}_1) + \bar{A}_2 m_1} Q_{A + \bar{A} + (3m_1, 3n_1) + (3m_2 - 1, 3n_2 - 1)} P_A P_{\bar{A}} \\
 & \quad - \sum_{A', \bar{A}} q^{n_2 A'_1 + (\bar{A}_2 + n_1)m_2 + (\bar{A}_2 + n_1)A'_1 + \bar{A}_2 m_1} Q_{A' + \bar{A} + (3m_1, 3n_1) + (3m_2 - 1, 3n_2 - 1)} P_{A'} P_{\bar{A}} \\
 & \quad - \sum_{A, B} q^{n_2(A_1 + m_1) + B_2 m_2 + B_2(A_1 + m_1) + A_2 m_1} Q_{A + B + (3m_1, 3n_1) + (3m_2 - 1, 3n_2 - 1)} P_A P_B \\
 & \quad + \sum_{A, B} q^{(A_2 + B_2 + n_2)m_1 + n_2 A_1 + B_2 m_2 + B_2 A_1} Q_{A + B + (3m_1, 3n_1) + (3m_2 - 1, 3n_2 - 1)} P_A P_B \\
 & \quad - \sum_{A, B} q^{n_2 A_1 + (n_1 + B_2)m_2 + (n_1 + B_2)A_1 + B_2 m_1} Q_{A + B + (3m_1, 3n_1) + (3m_2 - 1, 3n_2 - 1)} P_A P_B
 \end{aligned}$$

(using that the second and third terms cancel, as do the fifth and sixth)

$$\begin{aligned}
 & = q^{n_1 m_2} (-\mu q^{-(n_1 + n_2)(m_1 + m_2)} P_{-(3(m_1 + m_2) - 1, 3(n_1 + n_2) - 1)} \\
 & \quad - \sum_{\bar{A}, A'} q^{(n_1 + n_2)A'_1 + (m_1 + m_2)\bar{A}_2 + \bar{A}_2 A'_1} Q_{(3m_1 + 3m_2 - 1, 3n_1 + 3n_2 - 1) + A' + \bar{A}} P_{A'} P_{\bar{A}} \\
 & \quad - \sum_{A, B} q^{(n_1 + n_2)A_1 + (m_1 + m_2)B_2 + B_2 A_1} Q_{(3m_1 + 3m_2 - 1, 3n_1 + 3n_2 - 1) + A + B} P_A P_B \\
 & = q^{n_1 m_2} e^{(\mu)}_{21}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

Further,

$$\begin{aligned}
 & [e^{(\mu)}_{21}(m_1, n_1), e^{(\mu)}_{23}(m_2, n_2)] \\
 & = \mu q^{-m_1 n_1} \sum_A q^{n_2(-m_1 - m_2 - A_1) + A_2 m_2 + A_2(-m_1 - m_2 - A_1)} P_A P_{(-3m_1 - 3m_2, -3n_1 - 3n_2) - A} \\
 & \quad + \left[ \sum_{A, A'} q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A + A' + (3m_1 - 1, 3n_1 - 1)} P_A P_{A'}, \right. \\
 & \quad \left. \sum_{A, B} q^{n_2 B_1 + A_2 m_2 + A_2 B_1} Q_{(3m_2 + 1, 3n_2 + 1) + A + B} P_A P_B \right]
 \end{aligned}$$

$$\begin{aligned}
 & -\mu q^{-m_2 n_2} \sum_A q^{n_1 A_1 + (-n_1 - n_2 - A_2)m_1 + (-n_1 - n_2 - A_2)A_1} P_A P_{(-3m_1 - 3m_2, -3n_1 - 3n_2) - A} \\
 & + \left[ \sum_{A, B} q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B, \right. \\
 & \qquad \qquad \qquad \left. \sum_{A, B} q^{n_2 B_1 + A_2 m_2 + A_2 B_1} Q_{(3m_2+1, 3n_2+1)} P_A P_B \right] \\
 & + \left[ \sum_{A, B} q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B, \right. \\
 & \qquad \qquad \qquad \left. \sum_{B, B'} q^{n_2 B'_1 + B_2 m_2 + B_2 B'_1} Q_{(3m_2+1, 3n_2+1)+B+B'} P_B P_{B'} \right]
 \end{aligned}$$

(the first and third term cancel)

$$\begin{aligned}
 & = \sum_{A, \bar{A}, B} q^{n_1(m_2 + \bar{A}_1 + B_1) + A_2 m_1 + A_2(m_1 + \bar{A}_1 + B_1) + n_2 B_1 + \bar{A}_2 m_2 + \bar{A}_2 B_1} \\
 & \qquad \qquad \qquad \times Q_{A+\bar{A}+B+(3m_1+3m_2, 3n_1+3n_2)} P_A P_{\bar{A}} P_B \\
 & + \sum_{A', \bar{A}, B} q^{n_1 A'_1 + (n_2 + \bar{A}_2 + B_2)m_1 + (n_2 + \bar{A}_2 + B_2)\bar{A}_1 + n_2 B_1 + \bar{A}_2 m_2 + \bar{A}_2 B_1} \\
 & \qquad \qquad \qquad \times Q_{A'+\bar{A}+B+(3m_1+3m_2, 3n_1+3n_2)} P_{A'} P_{\bar{A}} P_B \\
 & - \sum_{A, A', B} q^{n_2 B_1 + (A_2 + A'_2 + n_1)m_2 + (A_2 + A'_2 + n_1)B_1 + n_1 A'_1 + A_2 m_1 + A_2 A'_1} \\
 & \qquad \qquad \qquad \times Q_{A'+A+B+(3m_1+3m_2, 3n_1+3n_2)} P_{A'} P_A P_B \\
 & + \sum_{A, B, B'} q^{n_1(m_2 + A_1 + B'_1) + B_2 m_1 + B_2(m_2 + A_1 + B'_1) + n_2 B'_1 + A_2 m_2 + A_2 B'_1} \\
 & \qquad \qquad \qquad \times Q_{A+B+B'+(3m_1+3m_2, 3n_1+3n_2)} P_A P_B P_{B'} \\
 & - \sum_{A, \bar{A}, B} q^{n_2(A_1 + B_1 + m_1) + \bar{A}_2 m_2 + \bar{A}_2(A_1 + B_1 + m_1) + n_1 A_1 + B_2 m_1 + B_2 A_1} \\
 & \qquad \qquad \qquad \times Q_{A+\bar{A}+B+(3m_1+3m_2, 3n_1+3n_2)} P_A P_{\bar{A}} P_B \\
 & + \sum_{A, B, B'} q^{n_1 A_1 + (n_2 + B_2 + B'_2)m_1 + (n_2 + B_2 + B'_2)A_1 + n_2 B'_1 + B_2 m_2 + B_2 B'_1} \\
 & \qquad \qquad \qquad \times Q_{A+B+B'+(3m_1+3m_2, 3n_1+3n_2)} P_A P_B P_{B'} \\
 & - \sum_{A, B, \bar{B}} q^{n_2(A_1 + \bar{B}_1 + m_1) + B_2 m_2 + B_2(A_1 + \bar{B}_1 + m_1) + n_1 A_1 + \bar{B}_2 m_1 + \bar{B}_2 A_1} \\
 & \qquad \qquad \qquad \times Q_{A+B+\bar{B}+(3m_1+3m_2, 3n_1+3n_2)} P_A P_B P_{\bar{B}} \\
 & - \sum_{A, B', \bar{B}} q^{n_2 B'_1 + (A_2 + \bar{B}_2 + n_1)m_2 + (A_2 + \bar{B}_2 + n_1)B'_1 + n_1 A_1 + \bar{B}_2 m_1 + \bar{B}_2 A_1} \\
 & \qquad \qquad \qquad \times Q_{A+B'+\bar{B}+(3m_1+3m_2, 3n_1+3n_2)} P_A P_{B'} P_{\bar{B}} \\
 & = 0,
 \end{aligned}$$

where the first term and the third, the second and the fifth, the fourth and the eighth, and the sixth and the seventh cancel.

The following three brackets are easy.

$$\begin{aligned}
 & [e^{(\mu)}_{21}(m_1, n_1), e^{(\mu)}_{31}(m_2, n_2)] = 0, \\
 & [e^{(\mu)}_{21}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] = -q^{n_2 m_1} e^{(\mu)}_{31}(m_1 + m_2, n_1 + n_2), \\
 & [e^{(\mu)}_{21}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] = 0.
 \end{aligned}$$

Next,

$$\begin{aligned}
 & [e^{(\mu)}{}_{22}(m_1, n_1), e^{(\mu)}{}_{22}(m_2, n_2)] \\
 &= \sum_{A, A'} q^{A_2 m_1 + A'_2 m_2} [Q_{(3m_1, 3n_1) + A} P_A, Q_{(3m_2, 3n_2) + A'} P_{A'}] \\
 &\quad + \sum_{B, B'} q^{B_2 m_1 + B'_2 m_2} [Q_{(3m_1, 3n_1) + B} P_B, Q_{(3m_2, 3n_2) + B'} P_{B'}] \\
 &= \sum_{A'} q^{(n_2 + A'_2) m_1 + A'_2 m_2} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + A'} P_{A'} \\
 &\quad - \sum_A q^{A_2 m_1 + (n_1 + A_2) m_2} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + A} P_A \\
 &\quad + \sum_{B'} q^{(n_2 + B'_2) m_1 + B'_2 m_2} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + B'} P_{B'} \\
 &\quad - \sum_B q^{B_2 m_1 + (n_1 + B_2) m_2} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + B} P_B \\
 &= q^{n_1 m_2} \left( - \sum_A q^{A_2(m_1 + m_2)} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + A} P_A \right. \\
 &\quad \left. - \sum_B q^{B_2(m_1 + m_2)} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + B} P_B - \frac{1}{2} \mu \delta_{(m_1 + m_2, n_1 + n_2), (0, 0)} \right) \\
 &\quad - q^{n_2 m_1} \left( - \sum_A q^{A_2(m_1 + m_2)} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + A} P_A \right. \\
 &\quad \left. - \sum_B q^{B_2(m_1 + m_2)} Q_{(3m_1, 3n_1) + (3m_2, 3n_2) + B} P_B - \frac{1}{2} \mu \delta_{(m_1 + m_2, n_1 + n_2), (0, 0)} \right) \\
 &= q^{n_1 m_2} e^{(\mu)}{}_{22}(m_1 + m_2, n_1 + n_2) - q^{n_2 m_1} e^{(\mu)}{}_{22}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

$$\begin{aligned}
 & [e^{(\mu)}{}_{22}(m_1, n_1), e^{(\mu)}{}_{23}(m_2, n_2)] \\
 &= \left[ - \sum_A q^{A_2 m_1} Q_{(3m_1, 3n_1) + A} P_A - \sum_B q^{B_2 m_1} Q_{(3m_1, 3n_1) + B} P_B, \right. \\
 &\quad - q^{-m_2 n_2} \mu P_{-(3m_2 + 1, 3n_2 + 1)} - \sum_{A, B} q^{n_2 B_1 + A_2 m_2 + A_2 B_1} Q_{A + B + (3m_2 + 1, 3n_2 + 1)} P_A P_B \\
 &\quad \left. - \sum_{B, B'} q^{n_2 B'_1 + B_2 m_2 + B_2 B'_1} Q_{B' + B + (3m_2 + 1, 3n_2 + 1)} P_B P_{B'} \right] \\
 &= \sum_{A, B} q^{(n_2 + A_2 + B_2) m_1 + n_2 B_1 + A_2 m_2 + A_2 B_1} Q_{(3m_1, 3n_1) + A + B + (3m_2 + 1, 3n_2 + 1)} P_A P_B \\
 &\quad - \sum_{A, B} q^{n_2 B_1 + (n_1 + A_2) m_2 + (A_2 + n_1) B_1 + A_2 m_1} Q_{(3m_1, 3n_1) + A + B + (3m_2 + 1, 3n_2 + 1)} P_A P_B \\
 &\quad \quad - q^{-m_2 n_2 + (-n_2 - n_1) m_1} \mu P_{-(3m_2 + 1, 3n_2 + 1) - (3m_1, 3n_1)} \\
 &\quad - \sum_{A, B} q^{n_2(m_1 + B_1) + A_2 m_2 + A_2(m_1 + B_1) + B_2 m_1} Q_{(3m_1, 3n_1) + A + B + (3m_2 + 1, 3n_2 + 1)} P_A P_B \\
 &\quad + \sum_{B, B'} q^{(n_2 + B_2 + B'_2) m_1 + n_2 B'_1 + B_2 m_2 + B_2 B'_1} Q_{(3m_1, 3n_1) + B + B' + (3m_2 + 1, 3n_2 + 1)} P_B P_{B'}
 \end{aligned}$$

$$\begin{aligned}
 & - \sum_{\mathbf{B}, \mathbf{B}'} q^{n_2(\mathbf{B}'_1+m_1)+\mathbf{B}_2m_2+\mathbf{B}_2(\mathbf{B}'_1+m_1)+\mathbf{B}'_2m_1} Q_{(3m_1, 3n_1)+\mathbf{B}+\mathbf{B}'+(3m_2+1, 3n_2+1)} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & - \sum_{\mathbf{B}, \mathbf{B}'} q^{n_2\mathbf{B}'_1+(n_1+\mathbf{B}_2)m_2+(\mathbf{B}_2+n_1)\mathbf{B}'_1+\mathbf{B}_2m_1} Q_{(3m_1, 3n_1)+\mathbf{B}+\mathbf{B}'+(3m_2+1, 3n_2+1)} P_{\mathbf{B}} P_{\mathbf{B}'}
 \end{aligned}$$

(the first term cancels the fourth, while the fifth cancels the sixth)

$$\begin{aligned}
 & = q^{m_2n_1} (-q^{-(m_1+m_2)(n_1+n_2)}) \mu P_{-(3m_1+3m_2+1, 3n_1+3n_2+1)} \\
 & - \sum_{\mathbf{A}, \mathbf{B}} q^{A_2(m_1+m_2)+(n_1+n_2)\mathbf{B}_1+A_2\mathbf{B}_1} Q_{\mathbf{A}+\mathbf{B}+(3m_1+3m_2+1, 3n_1+3n_2+1)} P_{\mathbf{A}} P_{\mathbf{B}} \\
 & - \sum_{\mathbf{B}, \mathbf{B}'} q^{\mathbf{B}'_1(n_1+n_2)+\mathbf{B}_2(m_1+m_2)\mathbf{B}_2\mathbf{B}'_1} Q_{\mathbf{B}+\mathbf{B}'+(3m_1+3m_2+1, 3n_1+3n_2+1)} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & = q^{m_2n_1} e^{(\mu)}_{23}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

The next three brackets are easy.

$$\begin{aligned}
 & [e^{(\mu)}_{22}(m_1, n_1), e^{(\mu)}_{31}(m_2, n_2)] = 0, \\
 & [e^{(\mu)}_{22}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] = -q^{n_2m_1} e^{(\mu)}_{32}(m_1 + m_2, n_1 + n_2), \\
 & [e^{(\mu)}_{22}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] = 0.
 \end{aligned}$$

Next,

$$\begin{aligned}
 & [e^{(\mu)}_{23}(m_1, n_1), e^{(\mu)}_{23}(m_2, n_2)] \\
 & = \left[ -q^{-m_1n_1} \mu P_{-(3m_1+1, 3n_1+1)} - \sum_{\mathbf{A}, \mathbf{B}} q^{n_1\mathbf{B}_1+A_2m_1+A_2\mathbf{B}_1} Q_{\mathbf{A}+\mathbf{B}+(3m_1+1, 3n_1+1)} P_{\mathbf{A}} P_{\mathbf{B}} \right. \\
 & \quad - \sum_{\mathbf{B}, \mathbf{B}'} q^{n_1\mathbf{B}'_1+\mathbf{B}_2m_1+\mathbf{B}_2\mathbf{B}'_1} Q_{\mathbf{B}+\mathbf{B}'+(3m_1+1, 3n_1+1)} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & \quad - q^{-m_2n_2} \mu P_{-(3m_2+1, 3n_2+1)} - \sum_{\mathbf{A}, \mathbf{B}} q^{n_2\mathbf{B}_1+A_2m_2+A_2\mathbf{B}_1} Q_{\mathbf{A}+\mathbf{B}+(3m_2+1, 3n_2+1)} P_{\mathbf{A}} P_{\mathbf{B}} \\
 & \quad \left. - \sum_{\mathbf{B}, \mathbf{B}'} q^{n_2\mathbf{B}'_1+\mathbf{B}_2m_2+\mathbf{B}_2\mathbf{B}'_1} Q_{\mathbf{B}+\mathbf{B}'+(3m_2+1, 3n_2+1)} P_{\mathbf{B}} P_{\mathbf{B}'} \right] \\
 & = \sum_{\mathbf{B}} \mu q^{-m_1n_1+n_2(-m_1-m_2-\mathbf{B}_1)+\mathbf{B}_2m_2+\mathbf{B}_2(-m_1-m_2-\mathbf{B}_1)} \\
 & \quad \times P_{\mathbf{B}} P_{-(3m_1+1, 3n_1+1)-(3m_2+1, 3n_2+1)-\mathbf{B}} \\
 & - \sum_{\mathbf{B}} \mu q^{-m_2n_2+n_1\mathbf{B}_1+(-n_2-n_1-\mathbf{B}_2)m_1+(-n_2-n_1-\mathbf{B}_2)\mathbf{B}_1} \\
 & \quad \times P_{\mathbf{B}} P_{-(3m_1+1, 3n_1+1)-(3m_2+1, 3n_2+1)-\mathbf{B}} \\
 & + \sum_{\mathbf{B}, \bar{\mathbf{B}}, \mathbf{B}'} q^{n_1(\mathbf{B}'_1+\bar{\mathbf{B}}_1+m_2)+\mathbf{B}_2m_1+\mathbf{B}_2(\mathbf{B}'_1+\bar{\mathbf{B}}_1+m_2)+n_2\mathbf{B}'_1+\bar{\mathbf{B}}_2m_2+\bar{\mathbf{B}}_2\mathbf{B}'_1} \\
 & \quad \times Q_{\mathbf{B}+\mathbf{B}'+\bar{\mathbf{B}}+(3m_1+1, 3n_1+1)+(3m_2+1, 3n_2+1)} P_{\mathbf{B}} P_{\mathbf{B}'} P_{\bar{\mathbf{B}}} \\
 & + \sum_{\mathbf{B}, \bar{\mathbf{B}}, \mathbf{B}'} q^{n_1\mathbf{B}'_1+(\mathbf{B}_2+\bar{\mathbf{B}}_2+n_2)m_1+\mathbf{B}'_1(\mathbf{B}_2+\bar{\mathbf{B}}_2+n_2)+n_2\bar{\mathbf{B}}_1+\mathbf{B}_2m_2+\mathbf{B}_2\bar{\mathbf{B}}_1} \\
 & \quad \times Q_{\mathbf{B}+\mathbf{B}'+\bar{\mathbf{B}}+(3m_1+1, 3n_1+1)+(3m_2+1, 3n_2+1)} P_{\mathbf{B}} P_{\mathbf{B}'} P_{\bar{\mathbf{B}}}
 \end{aligned}$$

$$\begin{aligned}
 & - \sum_{\mathbf{B}, \bar{\mathbf{B}}, \mathbf{B}'} q^{n_2(\mathbf{B}'_1 + \bar{\mathbf{B}}_1 + m_1) + \mathbf{B}_2 m_2 + \mathbf{B}_2(\mathbf{B}'_1 + \bar{\mathbf{B}}_1 + m_1) + n_1 \mathbf{B}'_1 + \bar{\mathbf{B}}_2 m_1 + \bar{\mathbf{B}}_2 \mathbf{B}'_1} \\
 & \quad \times Q_{\mathbf{B} + \mathbf{B}' + \bar{\mathbf{B}} + (3m_1 + 1, 3n_1 + 1) + (3m_2 + 1, 3n_2 + 1)} P_{\mathbf{B}} P_{\mathbf{B}'} P_{\bar{\mathbf{B}}} \\
 & - \sum_{\mathbf{B}, \bar{\mathbf{B}}, \mathbf{B}'} q^{n_2 \mathbf{B}'_1 + (\mathbf{B}_2 + \bar{\mathbf{B}}_2 + n_1) m_2 + (\mathbf{B}_2 + \bar{\mathbf{B}}_2 + n_1) \mathbf{B}'_1 + n_1 \bar{\mathbf{B}}_1 + \mathbf{B}_2 m_1 + \mathbf{B}_2 \bar{\mathbf{B}}_1} \\
 & \quad \times Q_{\mathbf{B} + \mathbf{B}' + \bar{\mathbf{B}} + (3m_1 + 1, 3n_1 + 1) + (3m_2 + 1, 3n_2 + 1)} P_{\mathbf{B}} P_{\mathbf{B}'} P_{\bar{\mathbf{B}}} \\
 & + \sum_{\mathbf{A}, \mathbf{B}, \mathbf{B}'} q^{n_1(\mathbf{B}_1 + \mathbf{B}'_1 + m_2) + \mathbf{A}_2 m_1 + \mathbf{A}_2(\mathbf{B}_1 + \mathbf{B}'_1 + m_2) + n_2 \mathbf{B}'_1 + \mathbf{B}_2 m_2 + \mathbf{B}_2 \mathbf{B}'_1} \\
 & \quad \times Q_{\mathbf{A} + \mathbf{B} + \mathbf{B}' + (3m_1 + 1, 3n_1 + 1) + (3m_2 + 1, 3n_2 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & + \sum_{\mathbf{A}, \mathbf{B}, \mathbf{B}'} q^{n_1 \mathbf{B}_1 + (\mathbf{A}_2 + \mathbf{B}'_2 + n_2) m_1 + (\mathbf{A}_2 + \mathbf{B}'_2 + n_2) \mathbf{B}_1 + n_2 \mathbf{B}'_1 + \mathbf{A}_2 m_2 + \mathbf{A}_2 \mathbf{B}'_1} \\
 & \quad \times Q_{\mathbf{A} + \mathbf{B} + \mathbf{B}' + (3m_1 + 1, 3n_1 + 1) + (3m_2 + 1, 3n_2 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & - \sum_{\mathbf{A}, \mathbf{B}, \mathbf{B}'} q^{n_2(\mathbf{B}_1 + \mathbf{B}'_1 + m_1) + \mathbf{A}_2 m_2 + \mathbf{A}_2(\mathbf{B}_1 + \mathbf{B}'_1 + m_1) + n_1 \mathbf{B}'_1 + \mathbf{B}_2 m_1 + \mathbf{B}_2 \mathbf{B}'_1} \\
 & \quad \times Q_{\mathbf{A} + \mathbf{B} + \mathbf{B}' + (3m_1 + 1, 3n_1 + 1) + (3m_2 + 1, 3n_2 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & - \sum_{\mathbf{A}, \mathbf{B}, \mathbf{B}'} q^{n_2 \mathbf{B}_1 + (\mathbf{A}_2 + \mathbf{B}'_2 + n_1) m_2 + (\mathbf{A}_2 + \mathbf{B}'_2 + n_1) \mathbf{B}_1 + n_1 \mathbf{B}'_1 + \mathbf{A}_2 m_1 + \mathbf{A}_2 \mathbf{B}'_1} \\
 & \quad \times Q_{\mathbf{A} + \mathbf{B} + \mathbf{B}' + (3m_1 + 1, 3n_1 + 1) + (3m_2 + 1, 3n_2 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} P_{\mathbf{B}'} \\
 & = 0,
 \end{aligned}$$

where the first two terms cancel, as do the third and the sixth, the fourth and the fifth, and the last two.

$$\begin{aligned}
 & [e^{(\mu)}_{23}(m_1, n_1), e^{(\mu)}_{31}(m_2, n_2)] \\
 & = \left[ -q^{-m_1 n_1} \mu P_{-(3m_1 + 1, 3n_1 + 1)} - \sum_{\mathbf{A}, \mathbf{B}} q^{n_1 \mathbf{B}_1 + \mathbf{A}_2 m_1 + \mathbf{A}_2 \mathbf{B}_1} Q_{\mathbf{A} + \mathbf{B} + (3m_1 + 1, 3n_1 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} \right. \\
 & \quad \left. - \sum_{\mathbf{B}, \mathbf{B}'} q^{n_1 \mathbf{B}'_1 + \mathbf{B}_2 m_1 + \mathbf{B}_2 \mathbf{B}'_1} Q_{\mathbf{B} + \mathbf{B}' + (3m_1 + 1, 3n_1 + 1)} P_{\mathbf{B}} P_{\mathbf{B}'} \right. \\
 & \quad \left. \sum_{\mathbf{A}} q^{A_1 n_2} Q_{(3m_2 - 2, 3n_2 - 2) + \mathbf{A}} P_{\mathbf{A}} \right] \\
 & = -q^{-m_1 n_1 + (-m_1 - m_2) n_2} \mu P_{-(3m_1 + 1, 3n_1 + 1) - (3m_2 - 2, 3n_2 - 2)} \\
 & - \sum_{\mathbf{A}, \mathbf{A}'} q^{n_1(m_2 + \mathbf{A}'_1) + \mathbf{A}_2 m_1 + \mathbf{A}_2(m_2 + \mathbf{A}'_1) + \mathbf{A}'_1 n_2} Q_{\mathbf{A} + \mathbf{A}' + (3m_2 - 2, 3n_2 - 2) + (3m_1 + 1, 3n_1 + 1)} P_{\mathbf{A}} P_{\mathbf{A}'} \\
 & + \sum_{\mathbf{A}, \mathbf{B}} q^{(\mathbf{A}_1 + \mathbf{B}_1 + m_1) n_2 + \mathbf{A}_2 m_1 + \mathbf{A}_2 \mathbf{B}_1 + n_1 \mathbf{B}_1} Q_{\mathbf{A} + \mathbf{B} + (3m_2 - 2, 3n_2 - 2) + (3m_1 + 1, 3n_1 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} \\
 & - \sum_{\mathbf{A}, \mathbf{B}} q^{n_1(m_2 + \mathbf{A}_1) + \mathbf{B}_2 m_1 + \mathbf{B}_2(m_2 + \mathbf{A}_1) + \mathbf{A}_1 n_2} Q_{\mathbf{A} + \mathbf{B} + (3m_2 - 2, 3n_2 - 2) + (3m_1 + 1, 3n_1 + 1)} P_{\mathbf{A}} P_{\mathbf{B}} \\
 & - \sum_{\mathbf{A}, \mathbf{B}} q^{n_1 \mathbf{B}_1 + (\mathbf{A}_2 + n_2) m_1 + (\mathbf{A}_2 + n_2) \mathbf{B}_1 + \mathbf{A}_1 n_2} Q_{\mathbf{A} + \mathbf{B} + (3m_2 - 2, 3n_2 - 2) + (3m_1 + 1, 3n_1 + 1)} P_{\mathbf{A}} P_{\mathbf{B}}
 \end{aligned}$$

(the third term and the fifth cancel)

$$\begin{aligned}
 & = q^{n_1 m_2} \left( -q^{-(m_1 + m_2)(n_1 + n_2)} \mu P_{-(3m_1 + 3m_2 - 1, 3n_1 + 3n_2 - 1)} \right. \\
 & \quad \left. - \sum_{\mathbf{A}, \mathbf{A}'} q^{(n_1 + n_2) \mathbf{A}'_1 + \mathbf{A}_2(m_1 + m_2) + \mathbf{A}_2 \mathbf{A}'_1} Q_{\mathbf{A} + \mathbf{A}' + (3m_2 + 3m_1 - 1, 3n_2 + 3n_1 - 1)} P_{\mathbf{A}} P_{\mathbf{A}'} \right)
 \end{aligned}$$

$$\begin{aligned}
 & - \sum_{A, B} q^{(n_1+n_2)A_1+B_2(m_1+m_2)+B_2A_1} Q_{A+B+(3m_2+3m_1-1, 3n_2+3n_1-1)} P_A P_B \Big) \\
 & = q^{n_1 m_2} e^{(\mu)}_{21}(m_1 + m_2, n_1 + n_2). \\
 [e^{(\mu)}_{23}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] \\
 & = \left[ -\mu q^{-m_1 n_1} P_{-(3m_1+1, 3n_1+1)} - \sum_{A, B} q^{n_1 B_1+A_2 m_1+A_2 B_1} Q_{(3m_1+1, 3n_1+1)+A+B} P_A P_B \right. \\
 & \quad \left. - \sum_{B, B'} q^{n_1 B'_1+B_2 m_1+B_2 B'_1} Q_{(3m_1+1, 3n_1+1)+B+B'} P_B P_{B'}, Q_{(3m_2-1, 3n_2-1)} \right] \\
 & = -\mu q^{-m_1 n_1} \delta_{(-m_1, -n_1), (m_2, n_2)} \\
 & \quad - \sum_A q^{n_1 m_2+A_2 m_1+A_2 m_2} Q_{(3m_1+1, 3n_1+1)+A+(3m_2-1, 3n_2-1)} P_A \\
 & \quad - \sum_B q^{n_1 m_2+B_2 m_1+B_2 m_2} Q_{(3m_1+1, 3n_1+1)+B+(3m_2-1, 3n_2-1)} P_B \\
 & \quad - \sum_{B'} q^{n_1 B'_1+n_2 m_1+n_2 B'_1} Q_{(3m_1+1, 3n_1+1)+B'+(3m_2-1, 3n_2-1)} P_{B'} \\
 & = q^{n_1 m_2} \left( - \sum_A q^{A_2(m_1+m_2)} Q_{(3m_1+3m_2, 3n_1+3n_2)+A} P_A \right. \\
 & \quad \left. - \sum_B q^{B_2(m_1+m_2)} Q_{(3m_1+3m_2, 3n_1+3n_2)+B} P_B - \frac{1}{2} \mu \delta_{(m_1+m_2, n_1+n_2), (0,0)} \right) \\
 & \quad - q^{n_2 m_1} \left( \sum_B q^{B_1(n_1+n_2)} Q_{(3m_1+3m_2, 3n_1+3n_2)+B} P_B + \frac{1}{2} \mu \delta_{(m_1+m_2, n_1+n_2), (0,0)} \right) \\
 & = q^{n_1 m_2} e^{(\mu)}_{22}(m_1 + m_2, n_1 + n_2) - q^{n_2 m_1} e^{(\mu)}_{33}(m_1 + m_2, n_1 + n_2). \\
 [e^{(\mu)}_{23}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] \\
 & = \left[ -\mu q^{-m_1 n_1} P_{-(3m_1+1, 3n_1+1)} - \sum_{A, B} q^{n_1 B_1+A_2 m_1+A_2 B_1} Q_{(3m_1+1, 3n_1+1)+A+B} P_A P_B \right. \\
 & \quad \left. - \sum_{B, B'} q^{n_1 B'_1+B_2 m_1+B_2 B'_1} Q_{(3m_1+1, 3n_1+1)+B+B'} P_B P_{B'}, \right. \\
 & \quad \left. \sum_B q^{B_1 n_2} Q_{(3m_2, 3n_2)+B} P_B \right] \\
 & = -q^{-m_1 n_1 + (-m_1 - m - 2)n_2} \mu P_{-(3m_1+1, 3n_1+1)-(3m_2, 3n_2)} \\
 & \quad - \sum_{A, B} q^{n_1(m_2+B_1)+A_2 m_1+B_2(m_2+B_1)+B_1 n_2} Q_{A+B+(3m_1+1, 3n_1+1)+(3m_2, 3n_2)} P_A P_B \\
 & \quad - \sum_{B, B'} q^{n_1(m_2+B'_1)+B_2 m_1+B_2(m_2+B'_1)+B'_1 n_2} Q_{B+(3m_2, 3n_2)+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \\
 & \quad - \sum_{B, B'} q^{n_1 B'_1+(n_2+B_2)m_1+(B_2+n_2)B'_1+B_1 n_2} Q_{B+(3m_2, 3n_2)+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \\
 & \quad + \sum_{B, B'} q^{(B_1+B'_1+m_1)n_2+n_1 B'_1+B_2 m_1+B_2 B'_1} Q_{B+(3m_2, 3n_2)+B'+(3m_1+1, 3n_1+1)} P_B P_{B'}
 \end{aligned}$$

(the last two terms cancel)

$$\begin{aligned}
 &= q^{n_1 m_2} \left( -q^{-(m_1+m_2)(n_1+n_2)} \mu P_{-(3m_1+3m_2+1, 3n_1+3n_2+1)} \right. \\
 &\quad - \sum_{A, B} q^{(n_1+n_2)\mathbf{B}_1 + A_2(m_1+m_2) + A_2 \mathbf{B}_1} Q_{A+\mathbf{B}+(3m_1+3m_2+1, 3n_1+3n_2+1)} P_A P_B \\
 &\quad \left. - \sum_{B, B'} q^{(n_1+n_2)\mathbf{B}'_1 + \mathbf{B}_2(m_1+m_2) + \mathbf{B}_2 \mathbf{B}'_1} Q_{\mathbf{B}+\mathbf{B}'+(3m_1+3m_2+1, 3n_1+3n_2+1)} P_B P_{B'} \right) \\
 &= q^{n_1 m_2} e^{(\mu)}_{23}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

The next five brackets are easy.

$$\begin{aligned}
 [e^{(\mu)}_{31}(m_1, n_1), e^{(\mu)}_{31}(m_2, n_2)] &= 0, \\
 [e^{(\mu)}_{31}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] &= 0, \\
 [e^{(\mu)}_{31}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] &= -q^{m_1 n_2} e^{(\mu)}_{31}(m_1 + m_2, n_1 + n_2), \\
 [e^{(\mu)}_{32}(m_1, n_1), e^{(\mu)}_{32}(m_2, n_2)] &= 0, \\
 [e^{(\mu)}_{32}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] &= -q^{m_1 n_2} e^{(\mu)}_{32}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

$$\begin{aligned}
 &[e^{(\mu)}_{33}(m_1, n_1), e^{(\mu)}_{33}(m_2, n_2)] \\
 &= \left[ \sum_B q^{\mathbf{B}_1 n_1} Q_{(3m_1, 3n_1)+\mathbf{B}} P_B, \sum_B q^{\mathbf{B}_1 n_2} Q_{(3m_2, 3n_2)+\mathbf{B}} P_B \right] \\
 &= \sum_B q^{(\mathbf{B}_1+m_2)n_1 + \mathbf{B}_1 n_2} Q_{(3m_1, 3n_1)+(3m_2, 3n_2)+\mathbf{B}} P_B \\
 &\quad - \sum_B q^{(\mathbf{B}_1+m_1)n_2 + \mathbf{B}_1 n_1} Q_{(3m_1, 3n_1)+(3m_2, 3n_2)+\mathbf{B}} P_B \\
 &= q^{n_1 m_2} \sum_B q^{\mathbf{B}_1(n_1+n_2)} Q_{(3m_1, 3n_1)+(3m_2, 3n_2)+\mathbf{B}} P_B \\
 &\quad - q^{m_1 n_2} \sum_B q^{\mathbf{B}_1(n_1+n_2)} Q_{(3m_1, 3n_1)+(3m_2, 3n_2)+\mathbf{B}} P_B \\
 &= q^{n_1 m_2} e^{(\mu)}_{33}(m_1 + m_2, n_1 + n_2) - q^{n_2 m_1} e^{(\mu)}_{33}(m_1 + m_2, n_1 + n_2).
 \end{aligned}$$

Next we check the brackets involving  $D_1^{(\mu)}$  and  $D_2^{(\mu)}$ .

$$\begin{aligned}
 &[D_1^{(\mu)}, e^{(\mu)}_{11}(m_1, n_1)] \\
 &= \left[ \sum_A A_1 Q_A P_A + \sum_B B_1 Q_B P_B, \sum_A q^{A_1 n_1} Q_{(3m_1, 3n_1)+A} P_A + \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} \right] \\
 &= \sum_A (m_1 + A_1) q^{A_1 n_1} Q_{(3m_1, 3n_1)+A} P_A - \sum_A A_1 q^{A_1 n_1} Q_{(3m_1, 3n_1)+A} P_A \\
 &= m_1 \left( \sum_A q^{A_1 n_1} Q_{(3m_1, 3n_1)+A} P_A + \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} \right) = m_1 e^{(\mu)}_{11}(m_1, n_1).
 \end{aligned}$$

The next two brackets are easy.

$$\begin{aligned}
 [D_1^{(\mu)}, e^{(\mu)}_{12}(m_1, n_1)] &= m_1 e^{(\mu)}_{12}(m_1, n_1), \\
 [D_1^{(\mu)}, e^{(\mu)}_{13}(m_1, n_1)] &= m_1 e^{(\mu)}_{13}(m_1, n_1).
 \end{aligned}$$

$$\begin{aligned}
 & [D_1^{(\mu)}, e^{(\mu)}_{21}(m_1, n_1)] \\
 &= \left[ \sum_A A_1 Q_A P_A + \sum_B B_1 Q_B P_B, \right. \\
 & \quad - q^{-m_1 n_1} \mu P_{-(3m_1-1, 3n_1-1)} - \sum_{A, A'} q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1, 3n_1-1)} P_A P_{A'} \\
 & \quad \quad \quad \left. - \sum_{A, B} q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B \right] \\
 &= q^{-m_1 n_1} \mu (-m_1) P_{-(3m_1-1, 3n_1-1)} \\
 & \quad - \sum_{A, A'} (A_1 + A'_1 + m_1) q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1, 3n_1-1)} P_A P_{A'} \\
 & \quad + \sum_{A, A'} A'_1 q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1, 3n_1-1)} P_A P_{A'} \\
 & \quad + \sum_{A, A'} A_1 q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1, 3n_1-1)} P_A P_{A'} \\
 & \quad - \sum_{A, B} (A_1 + B_1 + m_1) q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B \\
 & \quad + \sum_{A, B} B_1 q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B \\
 & \quad + \sum_{A, B} A_1 q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B \\
 &= m_1 \left( -q^{-m_1 n_1} \mu P_{-(3m_1-1, 3n_1-1)} \right. \\
 & \quad \quad \quad - \sum_{A, A'} q^{n_1 A'_1 + A_2 m_1 + A_2 A'_1} Q_{A+A'+(3m_1-1, 3n_1-1)} P_A P_{A'} \\
 & \quad \quad \quad \left. - \sum_{A, B} q^{n_1 A_1 + B_2 m_1 + B_2 A_1} Q_{A+B+(3m_1-1, 3n_1-1)} P_A P_B \right) \\
 &= m_1 e^{(\mu)}_{21}(m_1, n_1).
 \end{aligned}$$

$$\begin{aligned}
 & [D_1^{(\mu)}, e^{(\mu)}_{22}(m_1, n_1)] \\
 &= \left[ \sum_A A_1 Q_A P_A + \sum_B B_1 Q_B P_B, - \sum_A q^{A_2 m_1} Q_{(3m_1, 3n_1)+A} P_A \right. \\
 & \quad \quad \quad \left. - \sum_B q^{B_2 m_1} Q_{(3m_1, 3n_1)+B} P_B - \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} \right] \\
 &= - \sum_A (A_1 + m_1) q^{A_2 m_1} Q_{(3m_1, 3n_1)+A} P_A + \sum_A A_1 q^{A_2 m_1} Q_{(3m_1, 3n_1)+A} P_A \\
 & \quad - \sum_B (m_1 + B_1) q^{B_2 m_1} Q_{(3m_1, 3n_1)+B} P_B + \sum_B B_1 q^{B_2 m_1} Q_{(3m_1, 3n_1)+B} P_B
 \end{aligned}$$

$$\begin{aligned}
 &= m_1 \left( - \sum_A q^{A_2 m_1} Q_{(3m_1, 3n_1)+A} P_A - \sum_B q^{B_2 m_1} Q_{(3m_1, 3n_1)+B} P_B - \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} \right) \\
 &= m_1 e^{(\mu)}_{22}(m_1, n_1), \\
 & [D_1^{(\mu)}, e^{(\mu)}_{23}(m_1, n_1)] \\
 &= \left[ \sum_A A_1 Q_A P_A + \sum_B B_1 Q_B P_B, \right. \\
 & \quad - q^{-m_1 n_1} \mu P_{-(3m_1+1, 3n_1+1)} - \sum_{A, B} q^{n_1 B_1 + A_2 m_1 + A_2 B_1} Q_{A+B+(3m_1+1, 3n_1+1)} P_A P_B \\
 & \quad \left. - \sum_{B, B'} q^{n_1 B'_1 + B_2 m_1 + B_2 B'_1} Q_{B+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \right] \\
 &= q^{-m_1 n_1} \mu (-m_1) P_{-(3m_1+1, 3n_1+1)} \\
 & \quad - \sum_{A, B} (A_1 + B_1 + m_1) q^{n_1 B_1 + A_2 m_1 + A_2 B_1} Q_{A+B+(3m_1+1, 3n_1+1)} P_A P_B \\
 & \quad + \sum_{A, B} A_1 q^{n_1 B_1 + A_2 m_1 + A_2 B_1} Q_{A+B+(3m_1+1, 3n_1+1)} P_A P_B \\
 & \quad + \sum_{A, B} B_1 q^{n_1 B_1 + A_2 m_1 + A_2 B_1} Q_{A+B+(3m_1+1, 3n_1+1)} P_A P_B \\
 & \quad - \sum_{B, B'} (B_1 + B'_1 + m_1) q^{n_1 B'_1 + B_2 m_1 + B_2 B'_1} Q_{B+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \\
 & \quad + \sum_{B, B'} B_1 q^{n_1 B'_1 + B_2 m_1 + B_2 B'_1} Q_{B+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \\
 & \quad + \sum_{B, B'} B'_1 q^{n_1 B'_1 + B_2 m_1 + B_2 B'_1} Q_{B+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \\
 &= m_1 \left( -q^{-m_1 n_1} \mu P_{-(3m_1+1, 3n_1+1)} \right. \\
 & \quad - \sum_{A, B} q^{n_1 B_1 + A_2 m_1 + A_2 B_1} Q_{A+B+(3m_1+1, 3n_1+1)} P_A P_B \\
 & \quad \left. - \sum_{B, B'} q^{n_1 B'_1 + B_2 m_1 + B_2 B'_1} Q_{B+B'+(3m_1+1, 3n_1+1)} P_B P_{B'} \right) \\
 &= m_1 e^{(\mu)}_{23}(m_1, n_1).
 \end{aligned}$$

The following two brackets are easy.

$$[D_1^{(\mu)}, e^{(\mu)}_{31}(m_1, n_1)] = m_1 e^{(\mu)}_{31}(m_1, n_1),$$

$$[D_1^{(\mu)}, e^{(\mu)}_{32}(m_1, n_1)] = m_1 e^{(\mu)}_{32}(m_1, n_1).$$

$$\begin{aligned}
 & [D_1^{(\mu)}, e^{(\mu)}_{33}(m_1, n_1)] \\
 &= \left[ \sum_A A_1 Q_A P_A + \sum_B B_1 Q_B P_B, \sum_B q^{B_1 n_1} Q_{(3m_1, 3n_1)+B} P_B + \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_B (m_1 + \mathbf{B}_1) q^{\mathbf{B}_1 n_1} Q_{(3m_1, 3n_1) + \mathbf{B}} P_B - \sum_B \mathbf{B}_1 q^{\mathbf{B}_1 n_1} Q_{(3m_1, 3n_1) + \mathbf{B}} P_B \\
 &= m_1 \left( \sum_B q^{\mathbf{B}_1 n_1} Q_{(3m_1, 3n_1) + \mathbf{B}} P_B + \frac{1}{2} \mu \delta_{(m_1, n_1), (0, 0)} \right) \\
 &= m_1 e^{(\mu)}_{33}(m_1, n_1).
 \end{aligned}$$

Similarly, we can get

$$[D_2^{(\mu)}, e^{(\mu)}_{ij}(m_1, n_1)] = n_1 e^{(\mu)}_{ij}(m_1, n_1)$$

for  $1 \leq i, j \leq 3$ . Finally,

$$[D_1^{(\mu)}, D_2^{(\mu)}] = \left[ \sum_A \mathbf{A}_1 Q_A P_A + \sum_B \mathbf{B}_1 Q_B P_B, \sum_A \mathbf{A}_2 Q_A P_A + \sum_B \mathbf{B}_2 Q_B P_B \right] = 0.$$

Hence  $\pi : \widetilde{\mathfrak{gl}}_3(\mathbb{C}_q) \rightarrow \text{End}(V)$  is a Lie algebra homomorphism. □

### 3. Hermitian form for $\widetilde{\mathfrak{gl}}_3(\mathbb{C}_q)$ -module

From now on we need to assume that  $|q| = 1$ .

Define a  $\mathbb{R}$ -linear map  $\omega : \widetilde{\mathfrak{gl}}_3(\mathbb{C}_q) \mapsto \widetilde{\mathfrak{gl}}_3(\mathbb{C}_q)$  by

$$\begin{aligned}
 \omega(\lambda x) &= \bar{\lambda} \omega(x) \quad \text{for all } \lambda \in \mathbb{C} \text{ and } x \in \widetilde{\mathfrak{gl}}_3(\mathbb{C}_q), \\
 \omega(E_{ij}(a)) &= (-1)^{i+j} E_{ji}(\bar{a}) \quad \text{for } a \in \mathbb{C}_q, \\
 \omega(d_s) &= d_s, \quad \omega(c_s) = c_s, \\
 \omega(d_t) &= d_t, \quad \omega(c_t) = c_t.
 \end{aligned}$$

Here, the  $\mathbb{R}$ -linear function  $\bar{\phantom{x}} : \mathbb{C}_q \rightarrow \mathbb{C}_q$  is defined as  $\overline{\lambda s^m t^n} = \bar{\lambda} t^{-n} s^{-m} = \bar{\lambda} q^{mn} s^{-m} t^{-n}$ , where  $\bar{\lambda}$  is the complex conjugate for any  $\lambda \in \mathbb{C}$  and  $m, n \in \mathbb{Z}$ .

From [Gao and Zeng 2006, Lemma 3.4], we have

**Lemma 3.1.**  $\omega$  is an antilinear antiinvolution of  $\widetilde{\mathfrak{gl}}_3(\mathbb{C}_q)$ .

We write  $\pi(E_{ij}(r)) \cdot v$  more simply as  $E_{ij}(r) \cdot v$ , for any  $v \in V, r \in \mathbb{C}_q$ .

In [Gao and Zeng 2006], we defined a hermitian form on the basis consisting of monomials and then used another basis consisting of iterated module actions on the “highest weight vector” 1 to determine the condition for the form being positive definite. Here we will use the second basis directly to define the hermitian form which is much simpler.

**Lemma 3.2.** Consider, for  $k, l \in \mathbb{Z}_+ \cup \{0\}$ , the “vectors”

$$E_{12}(\alpha_1) E_{12}(\alpha_2) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot 1,$$

where

$$\alpha_i = s^{m_i} t^{n_i} \quad \text{for } i = 1, \dots, k,$$

$$\beta_j = s^{u_j} t^{v_j} \quad \text{for } j = 1, \dots, l,$$

and  $m_i, n_i, u_j, v_j \in \mathbb{Z}$ . We say each is in level  $(k, l)$  in  $W$ , and together they form a basis for  $V$ .

*Proof.* Define

$$f_{\mathbf{A}, \mathbf{B}} = \prod_{(m, n) \in \mathbb{Z}^2} x_{(3m+1, 3n+1)}^{\mathbf{A}(m, n)} \cdot \prod_{(m', n') \in \mathbb{Z}^2} x_{(3m'-1, 3n'-1)}^{\mathbf{B}(m', n')},$$

for  $\mathbf{A}(m, n), \mathbf{B}(m', n') \in \mathbb{Z}_+ \cup \{0\}$ , where only finitely many  $\mathbf{A}(m, n), \mathbf{B}(m', n')$  are nonzero. The  $f_{\mathbf{A}, \mathbf{B}}$  form a basis for  $V$ .

Let

$$g_{\mathbf{A}} = \prod_{(m, n) \in \mathbb{Z}^2} x_{(3m+1, 3n+1)}^{\mathbf{A}(m, n)} \quad \text{and} \quad h_{\mathbf{B}} = \prod_{(m', n') \in \mathbb{Z}^2} x_{(3m'-1, 3n'-1)}^{\mathbf{B}(m', n')}.$$

In a way similar to [Gao and Zeng 2006, Lemma 4.2], we can write  $g_{\mathbf{A}}$  as a linear combination of  $E_{12}(\alpha_1) \cdots E_{12}(\alpha_k) \cdot 1$  for  $k \leq \sum_{(m, n)} \mathbf{A}(m, n)$ , and  $h_{\mathbf{B}}$  can be written as a linear combination of  $E_{32}(\beta_1) \cdots E_{32}(\beta_l) \cdot 1$  for  $l \leq \sum_{(m', n')} \mathbf{B}(m', n')$ .

Since  $E_{12}(\alpha) E_{32}(\beta) \cdot u = E_{32}(\beta) E_{12}(\alpha) \cdot u$  for any  $u \in V$ , we can write  $f_{(\mathbf{A}, \mathbf{B})}$  as a linear combination of  $E_{12}(\alpha_1) \cdots E_{12}(\alpha_k) E_{32}(\beta_1), \dots, E_{32}(\beta_l) \cdot 1$ . Hence the collection of  $E_{12}(\alpha_1) \cdots E_{12}(\alpha_k) E_{32}(\beta_1), \dots, E_{32}(\beta_l) \cdot 1$  form a basis for  $V$ .  $\square$

Denote this basis in  $V$  by

$$\mathcal{B} = \{E_{12}(\alpha_1) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) \cdots E_{32}(\beta_l) \cdot 1 \mid \text{for all } k, l \in \mathbb{N}, \alpha_i, \beta_j \in \mathbb{C}_q\}.$$

**Lemma 3.3.** For any  $v \in V$ ,

- $\text{lev}(v) = \text{lev}(E_{ii}(a) \cdot v)$  for  $i = 1, 2, 3$ ;
- $\text{lev}(E_{12}(a)(v)) = \text{lev}(v) + (1, 0)$ ;
- $\text{lev}(E_{32}(a) \cdot v) = \text{lev}(v) + (0, 1)$ ;
- $\text{lev}(E_{21}(a) \cdot v) = \text{lev}(v) - (1, 0)$  or  $E_{21}(a) \cdot v = 0$  if  $\text{lev}(v) - (1, 0) \notin \mathbb{Z}_+^2$ ;
- $\text{lev}(E_{23}(a) \cdot v) = \text{lev}(v) - (0, 1)$  or  $E_{23}(a) \cdot v = 0$  if  $\text{lev}(v) - (0, 1) \notin \mathbb{Z}_+^2$ ,

for any nonzero  $a \in \mathbb{C}_q$ .

*Proof.* We only check those  $v$  in the basis  $\mathcal{B}$ .

$$\begin{aligned} & E_{22}(a) E_{12}(\alpha_1) E_{12}(\alpha_2) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot 1 \\ &= E_{12}(\alpha_1) E_{22}(a) E_{12}(\alpha_2) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot 1 \\ &\quad - E_{12}(\alpha_1 a) E_{12}(\alpha_2) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot 1 \\ &= E_{12}(\alpha_1) E_{12}(\alpha_2) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot \left(\frac{1}{2}\mu\right) \kappa(a) \cdot 1 \end{aligned}$$

$$\begin{aligned}
& - \sum_{i=1}^k E_{12}(\alpha_1) \cdots E_{12}(\alpha_i a) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot 1 \\
& + \sum_{i=1}^l E_{12}(\alpha_1) E_{12}(\alpha_2) \cdots E_{12}(\alpha_k) E_{32}(\beta_1) \cdots E_{32}(\beta_i a) \cdots E_{32}(\beta_l) \cdot 1,
\end{aligned}$$

so  $\text{lev}(v) = \text{lev}(E_{22}(a) \cdot v)$ . It is similar for  $E_{11}(a)$ ,  $E_{33}(a)$ .

Further,  $\text{lev}(E_{12}(a)(v)) = \text{lev}(v) + (1, 0)$  and  $\text{lev}(E_{32}(a) \cdot v) = \text{lev}(v) + (0, 1)$  are the definition of level.

For  $E_{21}(a) \cdot v$ , we prove by induction on the level of  $v$ :  $E_{21}(a) \cdot v = 0$  if  $\text{lev}(v) = (0, n)$ ,  $n \in \mathbb{Z}_+ \cup \{0\}$ . If  $n = 0$ , it is obvious that  $E_{21}(a) \cdot 1 = 0$ . Suppose it is true for  $n$ , then

$$\begin{aligned}
& E_{21}(a) E_{32}(\beta_1) E_{32}(\beta_2) \cdots E_{32}(\beta_{n+1}) \cdot 1 \\
& = E_{32}(\beta_1) E_{21}(a) E_{32}(\beta_2) \cdots E_{32}(\beta_{n+1}) \cdot 1 - E_{31}(\beta_1 a) E_{32}(\beta_2) \cdots E_{32}(\beta_{n+1}) \cdot 1 \\
& = - E_{32}(\beta_2) \cdots E_{32}(\beta_{n+1}) E_{31}(\beta_1 a) \cdot 1 \\
& = 0
\end{aligned}$$

by induction.

Supposing  $\text{lev}(E_{21}(a) \cdot v) = \text{lev}(v) - (1, 0)$  or  $E_{21}(a) \cdot v = 0$  is true for  $\text{lev}(v) = (m - 1, n)$ , then for  $v = E_{12}(b)v'$  with  $\text{lev}(v') = (m - 1, n)$  and  $0 \neq b \in \mathbb{C}_q$ , we have

$$E_{21}(a) \cdot E_{12}(b) \cdot v' = E_{12}(b) E_{21}(a) \cdot v' + E_{22}(ab) \cdot v' - E_{11}(ba) \cdot v'.$$

Since  $\text{lev}(E_{21}(a) \cdot v') = (m - 2, n)$ , we have  $\text{lev}(E_{21}(a) \cdot E_{12}(b) \cdot v') = (m - 1, n)$  or  $E_{21}(a) \cdot E_{12}(b) \cdot v' = 0$ . It is similar for  $E_{23}(a)$ .  $\square$

We easily define a contravariant (with respect to  $\pi, \omega$ ) hermitian form on  $V$  by using the basis  $\mathcal{B}$ .

Assuming that  $\mu$  is a real number, define the conjugate bilinear form on the elements in  $\mathcal{B}$  by induction on the level:

$$(1, 1) = 1, (1, f) = 0 \quad \text{if} \quad \text{lev}(f) \neq (0, 0)$$

Suppose for any  $v \in \mathcal{B}$ ,  $(u, v)$  is defined for any  $u$  such that  $\text{lev}(u) = (k', l')$  with  $k' + l' = r - 1$  if  $\text{lev}(u) = (k, l)$ , with  $k + l = r$ . Then there exists a  $u'$  such that  $\text{lev}(u') = (k - 1, l)$  or  $\text{lev}(u') = (k, l - 1)$  and some  $a \in \mathbb{C}_q$  such that  $u = E_{12}(a) \cdot u'$  or  $u = E_{32}(a) \cdot u'$ .

**Theorem 3.4.** *The conjugate bilinear form defined through*

$$(E_{12}(a) \cdot u', v) = (u', \omega(E_{12}(a)) \cdot v) \quad \text{and} \quad (E_{32}(a) \cdot u', v) = (u', \omega(E_{32}(a)) \cdot v)$$

*is a hermitian form on  $V$ .*

*Proof.* We must check that  $(E_{ij}(a) \cdot u, v) = (u, \omega(E_{ij}(a)) \cdot v)$  for  $1 \leq i, j \leq 3$ , and  $a \in \mathbb{C}_q$ . We must also check  $(D_i \cdot u, v) = (u, \omega(D_i) \cdot v)$  for  $i = 1, 2$ .

By definition,

$$(E_{12}(a)u, v) = (u, \omega(E_{12}(a))v) \quad \text{and} \quad (E_{32}(a)u, v) = (u, \omega(E_{32}(a))v),$$

and so

$$\begin{aligned} (E_{13}(a)u, v) &= ([E_{12}(1), E_{23}(a)]u, v) \\ &= (E_{12}(1)E_{23}(a)u, v) - (E_{23}(a)E_{12}(1)u, v) \\ &= (u, \omega(E_{23}(a))\omega(E_{12}(1))v) - (u, \omega(E_{12}(1))\omega(E_{23}(a))v) \\ &= (u, -\omega([E_{23}(a), E_{12}(1)])v) = (u, \omega(E_{13}(a))v). \end{aligned}$$

We use induction on  $\text{lev}(u)$  to prove  $(E_{11}(a) \cdot u, v) = (u, \omega(E_{11}(a)) \cdot v)$ . For any  $v \in \mathfrak{B}$ ,

$$(E_{11}(a)1, v) = \frac{1}{2}\mu\kappa(a)(1, v) = \frac{1}{2}\mu\kappa(a)\delta_{1,v}.$$

Since  $\text{lev}(E_{11}(a) \cdot v) = \text{lev}(v)$  for any  $v \in \mathfrak{B}$ ,

$$(1, \omega(E_{11}(a)) \cdot v) = (1, E_{11}(\bar{a}) \cdot v) = \frac{1}{2}\mu\kappa(\bar{a})\delta_{1,v}.$$

Hence

$$(E_{11}(a) \cdot 1, v) = (1, \omega(E_{11}(a)) \cdot v).$$

Suppose  $(E_{11}(a) \cdot u, v) = (u, \omega(E_{11}(a)) \cdot v)$  holds true for any  $\text{lev}(u) = (l, k)$  with  $l + k = r - 1$ . Then for  $\text{lev}(u) = (l, k)$  with  $l + k = r$ , we have  $u = E_{32}(b) \cdot u'$ , with  $\text{lev}(u') = (l, k - 1)$ , and

$$\begin{aligned} (E_{11}(a)E_{32}(b) \cdot u', v) &= (E_{32}(b)E_{11}(a) \cdot u', v) = (E_{11}(a) \cdot u', \omega(E_{32}(b)) \cdot v) \\ &= (u', \omega(E_{11}(a))\omega(E_{32}(b)) \cdot v) \\ &= (u', \omega(E_{32}(b))\omega(E_{11}(a)) \cdot v) \\ &= (E_{32}(b) \cdot u', \omega(E_{11}(a)) \cdot v) = (u, \omega(E_{11}(a)) \cdot v), \end{aligned}$$

or  $u = E_{12}(b) \cdot u'$ , with  $\text{lev}(u') = (l - 1, k)$ , and

$$\begin{aligned} (E_{11}(a)E_{12}(b) \cdot u', v) &= (E_{12}(b)E_{11}(a) \cdot u', v) + ([E_{11}(a), E_{12}(b)] \cdot u', v) \\ &= (E_{11}(a) \cdot u', \omega(E_{12}(b)) \cdot v) + (u', \omega([E_{11}(a), E_{12}(b)]) \cdot v) \\ &= (u', \omega(E_{11}(a))\omega(E_{12}(b)) \cdot v) - (u', [\omega(E_{11}(a)), \omega(E_{12}(b))] \cdot v) \\ &= (u', \omega(E_{12}(b))\omega(E_{11}(a)) \cdot v) \\ &= (E_{12}(b)u', \omega(E_{11}(a)) \cdot v) = (u, \omega(E_{11}(a)) \cdot v). \end{aligned}$$

Thus  $(E_{11}(a) \cdot u, v) = (u, \omega(E_{11}(a)) \cdot v)$ ; and

$$\begin{aligned}
 (E_{22}(a) \cdot u, v) &= ([E_{21}(a), E_{12}(1)] \cdot u, v) + (E_{11}(a)u, v) \\
 &= (E_{21}(a)E_{12}(1) \cdot u, v) - (E_{12}(1)E_{21}(a) \cdot u, v) + (E_{11}(a)u, v) \\
 &= (u, \omega(E_{12}(1))\omega(E_{21}(a)) \cdot v) \\
 &\quad - (u, \omega(E_{21}(a))\omega(E_{12}(1)) \cdot v) + (u, \omega(E_{11}(a)) \cdot v) \\
 &= (u, \omega([E_{21}(a), E_{12}(1)] \cdot v)) + (u, \omega(E_{11}(a)) \cdot v) \\
 &= (u, \omega(E_{22}(a)) \cdot v).
 \end{aligned}$$

It is similar for  $(E_{33}(a) \cdot u, v) = (u, \omega(E_{33}(a)) \cdot v)$ .

For  $D_1, D_2$ , we also proceed by induction on the level of  $u$ . It is obvious that  $(D_1 \cdot 1, v) = 0$  for any  $v \in \mathfrak{B}$ , and so  $(D_1 \cdot 1, 1) = (1, D_1 \cdot 1) = 0$ . Suppose  $(1, D_1 \cdot v) = 0$  is true for those  $\text{lev}(v) = (k', l')$  with  $k' + l' = r > 0$ . Then

$$\begin{aligned}
 (1, D_1 E_{12}(s^m t^n) \cdot v) &= (1, E_{12} D_1 \cdot v) + (1, m \cdot v) = 0, \\
 (1, D_1 E_{32}(s^m t^n) \cdot v) &= (1, E_{32} D_1 \cdot v) + (1, m \cdot v) = 0.
 \end{aligned}$$

Thus  $(D_1 \cdot 1, v) = (1, D_1 \cdot v)$ .

Suppose for any  $v \in \mathfrak{B}$  that  $(D_1 \cdot u, v) = (u, D_1 \cdot v)$  is true for all  $\text{lev}(u) = (k', l')$  such that  $k' + l' = r$ , then

$$\begin{aligned}
 (D_1 \cdot E_{12}(s^m t^n) \cdot u, v) &= (E_{12}(s^m t^n) D_1 \cdot u, v) + (m \cdot u, v) \\
 &= (D_1 \cdot u, \omega(E_{12}(s^m t^n)) \cdot v) + (u, m \cdot v) \\
 &= (u, D_1 \omega(E_{12}(s^m t^n)) \cdot v) + (u, m \cdot v) \\
 &= (u, \omega(E_{12}(s^m t^n)) D_1 \cdot v) \\
 &= (E_{12}(s^m t^n) u, D_1 \cdot v).
 \end{aligned}$$

It is similar for  $(D_1 \cdot E_{32}(s^m t^n) \cdot u, v) = (E_{32}(s^m t^n) \cdot u, D_1 \cdot v)$ .

Hence  $(D_1 \cdot u, v) = (u, D_1 \cdot v)$ , and  $(D_2 \cdot u, v) = (u, D_2 \cdot v)$ . Note that  $\omega(D_i) = D_i$  for  $i = 1, 2$ .  $\square$

#### 4. Conditions for unitarity

In this section we will determine when the hermitian form given last section is positive definite.

Let  $i \in \mathbb{N}$ ,  $\gamma = (\gamma_1, \dots, \gamma_s)$  be the  $s$ -partition of  $i$ . We denote by  $\text{Par}_s(i)$  be the set of all  $s$ -partition of  $i$ .

Let  $\gamma \in \text{Par}_s(N)$ . We say  $\pi i'_1 \times \pi'_2 \in S_N \times S_N$  is equivalent to  $\pi_1 \times \pi_2 \in S_N \times S_N$ , where  $S_N$  is the permutation group of  $N$  letters, if, for all  $z_1, \dots, z_N \in \mathbb{C}_q$  and

$w_1, \dots, w_N \in \mathbb{C}_q$ ,

$$\kappa(z_{\pi'_1(1)} w_{\pi'_2(1)}, \dots, z_{\pi'_1(\gamma_1)} w_{\pi'_2(\gamma_1)}) \cdots \kappa(z_{\pi'_1(\gamma_1+\dots+\gamma_{s-1}+1)} w_{\pi'_2(\gamma_1+\dots+\gamma_{s-1}+1)}, \dots, z_{\pi'_1(N)} w_{\pi'_2(N)})$$

can be obtained from the analogous expression for  $\pi_1 \times \pi_2$  by only rotating the variables. For example,  $\kappa(z_1 w_1 z_2 w_2 z_3 w_3) = \kappa(z_3 w_3 z_1 w_1 z_2 w_2)$ .

The following lemma is due to Jakobsen and Kac [1989].

**Lemma 4.1.** *Let  $z_1, z_2, \dots, z_N, w_1, w_2, \dots, w_N \in \mathbb{C}_q[s^{\pm 1}, t^{\pm 1}]$*

$$(4-2) \quad \begin{pmatrix} 0 & z_1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & z_2 \\ 0 & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & z_N \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ w_1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ w_2 & 0 \end{pmatrix} \cdots \begin{pmatrix} 0 & 0 \\ w_N & 0 \end{pmatrix} \cdot 1 \\ = \sum_{s=1}^N \sum_{\gamma \in \text{Par}_s(N)} \sum_{[\pi_1 \times \pi_2] \in (S_N \times S_N)(\gamma)} (-1)^{\gamma_1-1} (-\mu) \kappa(z_{\pi_1(1)} w_{\pi_2(1)} \cdots z_{\pi_1(\gamma_1)} w_{\pi_2(\gamma_1)}) \\ \cdot (-1)^{\gamma_2-1} (-\mu) \kappa(z_{\pi_1(\gamma_1+1)} w_{\pi_2(\gamma_1+1)} \cdots z_{\pi_1(\gamma_2)} w_{\pi_2(\gamma_2)}) \cdot \\ \cdots (-1)^{\gamma_s-1} (-\mu) \kappa(z_{\pi_1(\gamma_1+\dots+\gamma_{s-1}+1)} w_{\pi_2(\gamma_1+\dots+\gamma_{s-1}+1)} \cdots z_{\pi_1(N)} w_{\pi_2(N)}) \cdot 1$$

**Lemma 4.2.** *Let  $a_i, c_i, b_j, d_j \in \mathbb{C}_q$  for  $i = 1, \dots, m$  and  $j = 1, \dots, n$ . Let  $R = (a_i c_j)_{m \times m}$  and  $U = (b_i d_j)_{n \times n}$ , and set*

$$\Lambda = \begin{pmatrix} R & 0 \\ 0 & U \end{pmatrix}_{(m+n) \times (m+n)} = (\lambda_{i,j})_{(m+n) \times (m+n)}$$

Then

$$(4-3) \quad E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_n) E_{12}(c_1) \cdots E_{12}(c_m) E_{32}(d_1) \cdots E_{32}(d_n) \cdot 1 \\ = \sum_{s=1}^{m+n} \sum_{\gamma \in \text{Par}_s(m+n)} \sum_{[\pi_1 \times \pi_2] \in (S_{m+n} \times S_{m+n})(\gamma)} (-1)^{\gamma_1-1} (-\mu) \kappa(\lambda_{\pi_1(1), \pi_2(1)} \cdots \lambda_{\pi_1(\gamma_1), \pi_2(\gamma_1)}) \\ \cdot (-1)^{\gamma_2-1} (-\mu) \kappa(\lambda_{\pi_1(\gamma_1+1), \pi_2(\gamma_1+1)} \cdots \lambda_{\pi_1(\gamma_2), \pi_2(\gamma_2)}) \cdot \\ \cdots (-1)^{\gamma_s-1} (-\mu) \kappa(\lambda_{\pi_1(\gamma_1+\dots+\gamma_{s-1}+1), \pi_2(\gamma_1+\dots+\gamma_{s-1}+1)} \cdots \lambda_{\pi_1(N), \pi_2(N)}) \cdot 1.$$

**Remark 4.3.** It is easy to see that  $\lambda_{i,j}$  in every summand should be from different rows and different columns of  $\Lambda$ . And if the summand of (4-3) contains some  $\lambda_{i,j} = 0$ , then this summand is 0. Hence (4-3) is in fact the sum of those  $\lambda_{i,j}$  from  $R$  and  $U$ .

*Proof.* We proceed by induction on  $n$ . For  $n = 0$ , (4-3) is just (4-2). Next assume (4-3) is true up to  $n - 1$ ,

$$E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_n) E_{12}(c_1) \cdots E_{12}(c_m) E_{32}(d_1) \cdots E_{32}(d_n) \cdot 1$$

$$\begin{aligned}
&= E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) \\
&\quad \cdot (E_{12}(c_1) E_{23}(b_n) - E_{13}(c_1 b_n)) E_{12}(c_2) \cdots E_{12}(c_m) E_{32}(d_1) \cdots E_{32}(d_n) \cdot 1 \\
&= E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) E_{12}(c_1) E_{23}(b_n) E_{12}(c_2) \\
&\quad \cdots E_{12}(c_m) E_{32}(d_1) \cdots E_{32}(d_n) \cdot 1 \\
&\quad - E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) E_{12}(c_1) \cdots E_{12}(c_m) (E_{12}(c_1 b_n d_1) \\
&\quad + E_{32}(d_1) E_{13}(c_1 b_n)) E_{32}(d_2) \cdots E_{32}(d_n) \cdot 1 \\
&= E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) E_{12}(c_1) E_{23}(b_n) E_{12}(c_2) \\
&\quad \cdots E_{12}(c_m) E_{32}(d_1) \cdots E_{32}(d_n) \cdot 1 \\
&\quad + \sum_{i=1}^n E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) \\
&\quad \quad \cdot E_{12}(-c_1 b_n d_i) E_{12}(c_2) \cdots E_{12}(c_m) E_{32}(d_1) \cdots \widehat{E_{32}(d_i)} \cdots E_{32}(d_n) \cdot 1 \\
&= E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) E_{12}(c_1) E_{12}(c_2) \\
&\quad \cdots E_{12}(c_m) E_{23}(b_n) E_{32}(d_1) \cdots E_{32}(d_n) \cdot 1 \\
&\quad + \sum_{i=1}^n \sum_{j=1}^m E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) \\
&\quad \quad \cdot E_{12}(c_1) \cdots E_{12}(-c_j b_n d_i) \cdots E_{12}(c_m) E_{32}(d_1) \cdots \widehat{E_{32}(d_i)} \cdots E_{32}(d_n) \cdot 1 \\
&= \sum_{i=1}^n \sum_{j>i} E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) E_{12}(c_1) E_{12}(c_2) \cdots E_{12}(c_m) \\
&\quad \quad \cdot E_{32}(d_1) \cdots \widehat{E_{32}(d_i)} \cdots E_{32}(d_{j-1}) E_{32}(-d_i b_n d_j - d_j b_n d_i) \cdots E_{32}(d_n) \cdot 1 \\
&\quad + \sum_{i=1}^n E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) \\
&\quad \quad \cdot E_{12}(c_1) E_{12}(c_2) \cdots E_{12}(c_m) E_{32}(d_1) \cdots \widehat{E_{32}(d_i)} \cdots E_{32}(d_n) (-\mu) \kappa(b_n d_i) \cdot 1 \\
&\quad + \sum_{i=1}^n \sum_{j=1}^m E_{21}(a_1) \cdots E_{21}(a_m) E_{23}(b_1) \cdots E_{23}(b_{n-1}) \\
&\quad \quad \cdot E_{12}(c_1) \cdots E_{12}(-c_j b_n d_i) \cdots E_{12}(c_m) E_{32}(d_1) \cdots \widehat{E_{32}(d_i)} \cdots E_{32}(d_n) \cdot 1.
\end{aligned}$$

Because (4-3) is true for  $n - 1$ , we expand to see it is also true for  $n$ .  $\square$

**Lemma 4.4.** *The levels are orthogonal with respect to the the hermitian form; that is, the form vanishes when applied to two vectors from different levels.*

*Proof.* Only need to prove those elements in the basis  $\mathcal{B}$ . Let

$$u = E_{12}(a_1) \cdots E_{12}(a_m) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1,$$

$$v = E_{12}(c_1) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1,$$

and suppose  $(m, n) \neq (k, l)$ .

First we prove  $(u, v) = 0$  with  $m = 0$ . If  $k = 0$ , then, supposing  $n > l$ , we have

$$\begin{aligned}(u, v) &= (E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1, E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1) \\ &= ((-1)^l E_{23}(\bar{d}_l) \cdots E_{23}(\bar{d}_1) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1, 1).\end{aligned}$$

Then, by Lemma 3.3,  $\text{lev}(E_{23}(\bar{d}_l) \cdots E_{23}(\bar{d}_1) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1) = (0, n - l)$ , or, if  $E_{23}(\bar{d}_l) \cdots E_{23}(\bar{d}_1) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1 = 0$ , then  $(u, v) = 0$ .

For  $k > 0$ ,

$$\begin{aligned}(u, v) &= (E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1, E_{12}(c_1) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1) \\ &= (-E_{21}(\bar{c}_1) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1, \\ &\quad E_{12}(c_2) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1).\end{aligned}$$

Then from Lemma 3.3, we have  $-E_{21}(\bar{c}_1) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1 = 0$ , and  $(u, v) = 0$ .

Without loss of generality, we can assume that  $m \leq k$ ; then

$$\begin{aligned}(u, v) &= (E_{12}(a_1) \cdots E_{12}(a_m) E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1, \\ &\quad E_{12}(c_1) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1) \\ &= (E_{32}(b_1) \cdots E_{32}(b_n) \cdot 1, \\ &\quad (-1)^m E_{21}(\bar{a}_m) \cdots E_{21}(\bar{a}_1) \cdot E_{12}(c_1) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1).\end{aligned}$$

From Lemma 3.3,

$$\text{lev}(E_{21}(\bar{a}_m) \cdots E_{21}(\bar{a}_1) \cdot E_{12}(c_1) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1) = (k - m, n)$$

or

$$E_{21}(\bar{a}_m) \cdots E_{21}(\bar{a}_1) \cdot E_{12}(c_1) \cdots E_{12}(c_k) E_{32}(d_1) \cdots E_{32}(d_l) \cdot 1 = 0.$$

Then, going back to the case  $m = 0$ , we get  $(u, v) = 0$ .  $\square$

Similarly to [Gao and Zeng 2006, Proposition 4.11] and together with Lemma 4.2, we have

**Proposition 4.5.** *The hermitian form on the element  $h$  in level  $(m, n)$  is a polynomial in  $\mu$ , with leading term  $c(-1)^{m+n}(-\mu)^{m+n} = c\mu^{m+n}$  for some constant  $c > 0$ .*

**Theorem 4.6.**  *$(\pi, V)$  can be made unitary if and only if  $\mu > 0$ .*

*Proof.* From [Gao and Zeng 2006, Theorem 4.12], the hermitian form in level  $(0, n)$  and  $(m, 0)$  is positive definite if and only if  $\mu > 0$ .

Define

$$T_{a,b}(s^{m_1} t^{n_1} s^{m_2} t^{n_2} \cdots s^{m_k} t^{n_k}) = s^{m_1+a} t^{n_1+b} s^{m_2+a} t^{n_2+b} \cdots s^{m_k+a} t^{n_k+b}$$

for  $a, b \in \mathbb{Z}$ . Extend this operator to the linear operator  $\widetilde{T}_{a,b}$  on  $V$  by

$$\begin{aligned} \widetilde{T}_{a,b}(E_{12}(\alpha_1)E_{12}(\alpha_2) \cdots E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2) \cdots E_{32}(\beta_l) \cdot 1) \\ = E_{12}(T_{a,b}\alpha_1)E_{12}(T_{a,b}\alpha_2) \cdots E_{12}(T_{a,b}\alpha_k) \\ \cdot E_{32}(T_{a,b}\beta_1)E_{32}(T_{a,b}\beta_2) \cdots E_{32}(T_{a,b}\beta_l) \cdot 1. \end{aligned}$$

Following Lemma 4.2,  $\widetilde{T}_{a,b}$  preserves the hermitian form on  $V$ . Denote

$$L_{l,r}(M, N) = \text{span}\{E_{12}(s^{m_1}t^{n_1}) \cdots E_{12}(s^{m_l}t^{n_l})E_{32}(s^{j_1}t^{k_1}) \cdots E_{32}(s^{j_r}t^{k_r}) \cdot 1$$

for  $|m_i, n_i| \geq 0$  and  $i = 1, \dots, l$ , with  $j_i, k_i \geq 0$ ,

$$\left\{ \sum_{i=1}^l m_i + \sum_{i=1}^r j_i \leq M, \quad \text{and} \quad \sum_{i=1}^r n_i + \sum_{i=1}^r k_i \leq N \right\}.$$

Since the hermitian form on two different levels is 0, we will prove the unitarity by induction on the level.

For any  $\mu > 0$ , the form is definite in level  $(0, n)$ ; see [Gao and Zeng 2006, Theorem 4.12]. Suppose it is definite in level  $(r, n)$  for those  $r < m$  and it is not definite in level  $(m, n)$ .

From Proposition 4.5, we know that the hermitian form restricted to this level should be positive definite for  $\mu$  big enough. Assuming it is not positive definite for some  $\mu > 0$ , there exist  $M, N$  such that the form restricted to  $L_{m,n}(M, N)$  is not positive definite. From Proposition 4.5, the form on  $L_{l,r}(M, N)$  varies smoothly with  $\mu$ . Then we can find a  $\mu_0$  for which the form is not positive definite and, for all  $\mu > \mu_0$ , it is positive definite. We write  $(\cdot, \cdot)_\mu$  for the hermitian form at  $\mu$ .

Thus the radical of the form is nontrivial at  $\mu_0$ , that is, there exists a nonzero  $\tilde{h} \in L_{m,n}(M, N)$  such that, for any  $h \in L_{m,n}(M, N)$ , we have

$$(\tilde{h}, h)_{\mu_0} = 0.$$

Therefore for any arbitrary element  $h_{m-1,n}$  in  $L_{m-1,n}(M, N)$  and any  $c \in \mathbb{C}$ , we have

$$(E_{21}(c) \cdot \tilde{h}, h_{m-1,n})_{\mu_0} = 0.$$

Since the form is positive definite in level  $(m-1, n)$ , we have  $E_{21}(c) \cdot \tilde{h} = 0$  for any  $c \in \mathbb{C}$ . Replacing  $\tilde{h}$  by  $\widetilde{T_{-a,-b}}(\tilde{h})$  if necessary, we can write

$$\tilde{h} = \sum_{i=1}^m a_i (E_{12}(1))^i x_i,$$

where  $x_i = \sum E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m)E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1$  (here it is a finite sum),  $a_i, \beta_j$  is the of form  $s^l t^k$ , and  $l, k$  cannot both be 0.

Let  $i_0$  be the smallest index such that  $a_{i_0} \neq 0$ ; then  $i_0 \geq 1$ .

Since

$$\begin{aligned}
 & E_{21}(c)(E_{12}(1))^i E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &= (E_{12}(1))^i E_{21}(c) E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &\quad + i(E_{12}(1))^{i-1} (E_{22}(c) - E_{11}(c)) E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &\quad + (-2c) \frac{i(i-1)}{2} (E_{12}(1))^{i-1} E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &= (E_{12}(1))^i E_{21}(c) E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &\quad + i(E_{12}(1))^{i-1} ((-2c)(m-i) - n) E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &\quad + i(E_{12}(1))^{i-1} E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) (E_{22}(c) - E_{11}(c)) \cdot 1 \\
 &\quad + (-2c) \frac{i(i-1)}{2} (E_{12}(1))^{i-1} E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &= (E_{12}(1))^i E_{21}(c) E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1 \\
 &\quad + [ic(-\mu_0) + i((-2c)(m-i) - n) + (-2c) \frac{i(i-1)}{2}]. \\
 &\quad (E_{12}(1))^{i-1} E_{12}(\alpha_{i+1}) \cdots E_{12}(\alpha_m) E_{32}(\beta_1) \cdots E_{32}(\beta_n) \cdot 1,
 \end{aligned}$$

we have

$$E_{21}(c)\tilde{h} = \gamma a_{i_0} (E_{12}(1))^{i_0-1} x_{i_0} + R,$$

where  $R$  contains those terms with powers of  $E_{12}(1)$  greater than  $i_0 - 1$  and

$$\begin{aligned}
 \gamma &= i_0 c(-\mu_0) + i_0((-2c)(m-i_0) - n) + (-2c) \frac{i_0(i_0-1)}{2} \\
 &= ci_0(-\mu_0 - (m-i_0) - (m-1)).
 \end{aligned}$$

Since  $m \geq i_0 \geq 1$ ,  $\mu_0 \geq 0$ , and  $\gamma \neq 0$ , this contradicts  $E_{21}(c)\tilde{h} = 0$ . Thus, for any  $\mu > 0$ , the hermitian form is positive definite.  $\square$

### Acknowledgments

I am grateful to my supervisors Professors Nantel Bergeron and Yun Gao for their encouragement and support during the preparation of this paper. I am especially grateful to Professor Yun Gao for drawing my attention to this subject.

### References

- [Allison et al. 1997] B. N. Allison, S. Azam, S. Berman, Y. Gao, and A. Pianzola, "Extended affine Lie algebras and their root systems", *Mem. Amer. Math. Soc.* **126**:603 (1997), x+122. MR 97i:17015 Zbl 0879.17012
- [Baranovsky et al. 2000] V. Baranovsky, S. Evens, and G. V., "Representations of quantum tori and double-affine Hecke algebras", preprint, 2000. math.RT/0005024
- [Berman and Szmigielski 1999] S. Berman and J. Szmigielski, "Principal realization for the extended affine Lie algebra of type  $\mathfrak{sl}_2$  with coordinates in a simple quantum torus with two generators", pp. 39–67 in *Recent developments in quantum affine algebras and related topics* (Raleigh,

- NC, 1998), edited by N. Jing and K. C. Misra, *Contemp. Math.* **248**, Amer. Math. Soc., Providence, RI, 1999. MR 2001h:17038 Zbl 1031.17014
- [Berman et al. 1996] S. Berman, Y. Gao, and Y. S. Krylyuk, “Quantum tori and the structure of elliptic quasi-simple Lie algebras”, *J. Funct. Anal.* **135**:2 (1996), 339–389. MR 97b:17007 Zbl 0847.17009
- [Billig and Zhao 2004] Y. Billig and K. Zhao, “Vertex operator representations of quantum tori at roots of unity”, *Comm. Contemp. Math.* **6**:1 (2004), 195–220. MR 2005b:17054 Zbl pre02078551
- [Eswara Rao 2003] S. Eswara Rao, “Unitary modules for EALAs co-ordinatized by a quantum torus”, *Comm. Algebra* **31**:5 (2003), 2245–2256. MR 2004e:17024 Zbl 1106.17029
- [Eswara Rao 2004] S. Eswara Rao, “A class of integrable modules for the core of EALA coordinatized by quantum tori”, *J. Algebra* **275**:1 (2004), 59–74. MR 2005a:17024 Zbl pre02105787
- [Eswara Rao and Batra 2002] S. Eswara Rao and P. Batra, “A new class of representations of EALA coordinated by quantum tori in two variables”, *Canad. Math. Bull.* **45**:4 (2002), 672–685. MR 2003k:17031 Zbl 1034.17015
- [Eswara Rao and Zhao 2004] S. Eswara Rao and K. Zhao, “Highest weight irreducible representations of rank 2 quantum tori”, *Math. Res. Lett.* **11**:5-6 (2004), 615–628. MR 2005j:17006 Zbl 1094.17011
- [Etingof et al. 1998] P. I. Etingof, I. B. Frenkel, and A. A. Kirillov, Jr., *Lectures on representation theory and Knizhnik–Zamolodchikov equations*, Mathematical Surveys and Monographs **58**, American Mathematical Society, Providence, RI, 1998. MR 2001b:32028 Zbl 0903.17006
- [Feĭgin and Frenkel 1990] B. L. Feĭgin and E. V. Frenkel, “Affine Kac–Moody algebras and semi-infinite flag manifolds”, *Comm. Math. Phys.* **128** (1990), 161–189. MR 92f:17026 Zbl 0722.17019
- [Gao 2000a] Y. Gao, “Representations of extended affine Lie algebras coordinatized by certain quantum tori”, *Compositio Math.* **123**:1 (2000), 1–25. MR 2002c:17036 Zbl 0958.17008
- [Gao 2000b] Y. Gao, “Vertex operators arising from the homogeneous realization for  $\widehat{\mathfrak{gl}}_N$ ”, *Comm. Math. Phys.* **211**:3 (2000), 745–777. MR 2001k:17044 Zbl 0982.17012
- [Gao 2002] Y. Gao, “Fermionic and bosonic representations of the extended affine Lie algebra  $\mathfrak{gl}_N(\mathbb{C}_q)$ ”, *Canad. Math. Bull.* **45**:4 (2002), 623–633. MR 2003h:17030 Zbl 1040.17017
- [Gao and Zeng 2006] Y. Gao and Z. Zeng, “Hermitian representations of the extended affine Lie algebra  $\widehat{\mathfrak{gl}}_2(\mathbb{C}_q)$ ”, *Adv. Math.* **207**:1 (2006), 244–265. MR 2007g:17023 Zbl 1106.17030
- [Golenishcheva-Kutuzova and Kac 1998] M. I. Golenishcheva-Kutuzova and V. G. Kac, “ $\Gamma$ -conformal algebras”, *J. Math. Phys.* **39**:4 (1998), 2290–2305. MR 99d:81048 Zbl 1031.81527
- [Høegh-Krohn and Torrésani 1990] R. Høegh-Krohn and B. Torrésani, “Classification and construction of quasisimple Lie algebras”, *J. Funct. Anal.* **89**:1 (1990), 106–136. MR 91a:17008 Zbl 0792.17019
- [Jakobsen and Kac 1989] H. P. Jakobsen and V. Kac, “A new class of unitarizable highest weight representations of infinite-dimensional Lie algebras. II”, *J. Funct. Anal.* **82**:1 (1989), 69–90. MR 89m:17032 Zbl 0688.17007
- [Lau 2005] M. Lau, “Bosonic and fermionic representations of Lie algebra central extensions”, *Adv. Math.* **194**:2 (2005), 225–245. MR 2005m:17008 Zbl pre02183827
- [Lin and Tan 2004] W. Lin and S. Tan, “Representations of the Lie algebra of derivations for quantum torus”, *J. Algebra* **275**:1 (2004), 250–274. MR 2005a:17014 Zbl 1106.17020
- [Lin and Tan 2006] W. Lin and S. Tan, “Harish–Chandra modules for the  $q$ -analog Virasoro-like algebra”, *J. Algebra* **297**:1 (2006), 254–272. MR 2007a:17040 Zbl pre05028847

- [Manin 1991] Y. I. Manin, *Topics in noncommutative geometry*, M. B. Porter Lectures, Princeton University Press, Princeton, NJ, 1991. MR 92k:58024 Zbl 0724.17007
- [Miki 2004] K. Miki, “Integrable irreducible highest weight modules for  $\mathfrak{sl}_2(\mathbb{C}_p[x^{\pm 1}, y^{\pm 1}])$ ”, *Osaka J. Math.* **41**:2 (2004), 295–326. MR 2005d:17024 Zbl 02111459
- [Su and Zhu 2005] Y. Su and L. Zhu, “Representations of the derivation algebra of the localization of the quantum plane at  $q = -1$ ”, *Comm. Algebra* **33**:12 (2005), 4375–4382. MR 2006m:17007 Zbl pre02248208
- [Varagnolo and Vasserot 1998] M. Varagnolo and E. Vasserot, “Double-loop algebras and the Fock space”, *Invent. Math.* **133**:1 (1998), 133–159. MR 99g:17035 Zbl 0904.17014
- [Wakimoto 1985] M. Wakimoto, “Extended affine Lie algebras and a certain series of Hermitian representations”, manuscript, 1985.
- [Wakimoto 1986] M. Wakimoto, “Fock representations of the affine Lie algebra  $A_1^{(1)}$ ”, *Comm. Math. Phys.* **104**:4 (1986), 605–609. MR 87m:17011 Zbl 0587.17009
- [Zhang and Zhao 1996] H. Zhang and K. Zhao, “Representations of the Virasoro-like algebra and its  $q$ -analog”, *Comm. Algebra* **24**:14 (1996), 4361–4372. MR 97m:17031 Zbl 0891.17016

Received October 2, 2006. Revised November 12, 2006.

ZITING ZENG  
SCHOOL OF MATHEMATICAL SCIENCES  
BEIJING NORMAL UNIVERSITY  
100875 BEIJING  
CHINA  
zengzt@bnu.edu.cn

