

*Pacific
Journal of
Mathematics*

**CLASSIFICATION OF IRREDUCIBLE INTEGRABLE MODULES
FOR TWISTED TOROIDAL LIE ALGEBRAS
WITH FINITE-DIMENSIONAL WEIGHT SPACES**

S. ESWARA RAO AND PUNITA BATRA

Volume 237 No. 1

September 2008

CLASSIFICATION OF IRREDUCIBLE INTEGRABLE MODULES FOR TWISTED TOROIDAL LIE ALGEBRAS WITH FINITE-DIMENSIONAL WEIGHT SPACES

S. ESWARA RAO AND PUNITA BATRA

This paper classifies irreducible, integrable modules for “twisted toroidal Lie algebras” with finite-dimensional weight spaces. We prove that these modules turn out to be modules of appropriate direct sums of finitely many copies of affine Kac–Moody Lie algebras.

Introduction

Let \mathfrak{g} be a finite-dimensional simple Lie algebra over the complex field \mathbb{C} . Let $A_n = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ be a Laurent polynomial ring in n commuting variables t_1, t_2, \dots, t_n . For $n > 1$, we consider the multiloop Lie algebra $\mathfrak{g} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. The universal central extension of $\mathfrak{g} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ is called a toroidal Lie algebra. If $n = 1$, then it is well known that the universal central extension of a loop algebra $\mathfrak{g} \otimes \mathbb{C}[t^{\pm 1}]$ is one-dimensional. In this case, the universal central extension is an affine Kac–Moody Lie algebra. On the other hand, in the toroidal case, that is, when $n > 1$, the universal central extension is infinite-dimensional. Toroidal Lie algebras are described in [Rao and Moody 1994]. Representations of toroidal Lie algebras have been studied in [Berman and Billig 1999] and [Rao 2004].

In Section 1, we define the twisted toroidal Lie algebra “ $\tau(\mu)$ ”, where μ is a Dynkin diagram automorphism of \mathfrak{g} of order m . In the one variable case, $\tau(\mu)$ is just the twisted affine Kac–Moody Lie algebra [Kac 1990]. Rao [1993] gave some representations of the twisted affine Kac–Moody Lie algebra. Batra [2004] classified finite-dimensional irreducible representations of the twisted multiloop Lie algebras.

In this paper, we twist several variables at the same time, because twisting one variable restricts the class of twisted toroidal Lie algebras to one for each variable. By twisting several variables, we are covering a larger class of twisted toroidal Lie algebras. Now $GL(n, \mathbb{Z})$, the group of $n \times n$ matrices with entries in \mathbb{Z} , acts on \mathbb{Z}^n naturally, and an element of $GL(n, \mathbb{Z})$ defines an automorphism of the nontwisted toroidal Lie algebra “ τ ”. This automorphism is change of coordinates as described

MSC2000: 17B65.

Keywords: twisted toroidal Lie algebras, integrable modules.

in Section 1. We consider $\tau(\mu)$ up to a suitable change of coordinates, and in this way, we can reduce the number of twisted variables to a minimum of two variables; see Remark 1.1.

Among all the modules of a toroidal Lie algebra, the so-called integrable modules (where the real root spaces act locally nilpotently) are very important since they are the ones that lift to representations of the corresponding group. Rao [2004] classified irreducible integrable modules for nontwisted toroidal Lie algebras with finite-dimensional weight spaces. Here we classify the irreducible integrable modules for twisted toroidal Lie algebras having finite-dimensional weight spaces, where the center acts nontrivially. Some of the ideas used in [Rao 2004] go through here, but this is a more complex case and needs new ideas.

In Section 1, we describe the root system for $\tau(\mu)$ and the Weyl group of $\tau(\mu)$, and we define integrable $\tau(\mu)$ -modules. In Section 2, we define central operators on an irreducible module for $\tau(\mu)$. In this paper we classify those irreducible integrable $\tau(\mu)$ -modules V with finite-dimensional weight spaces in which some zero degree generators of the center of $\tau(\mu)$ act nontrivially. Then by change of coordinates, we assume that one of the zero degree generators of the center of $\tau(\mu)$, namely “ K_n ”, acts nontrivially, while the others act trivially. Assuming that K_n acts as a positive integer, we prove in Proposition 2.6 the existence of a highest weight vector in V .

In Section 3, we give the notion of a non-graded $\tau(\mu)$ -module $V(\psi)$, define the graded $\tau(\mu)$ -module $V(\bar{\psi})$, and, in Proposition 3.3, establish a one-to-one correspondence between non-graded and graded modules. So our problem reduces to classifying irreducible integrable non-graded $\tau(\mu)$ -modules. In Section 4, we prove that any such non-graded module is actually a module for $(\mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]) \cap \tau(\mu)$, where $\mathfrak{g}_{\text{aff}} = \mathfrak{g} \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}K_n$ is an affine Lie algebra. In Proposition 4.1, we give the conditions under which the Lie algebra homomorphism $\phi(\mu)$ from $(\mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]) \cap \tau(\mu)$ to $\bigoplus \mathfrak{g}_{\text{aff}}$ (the direct sum of finitely many copies of $\mathfrak{g}_{\text{aff}}$) becomes surjective. In Proposition 5.3, we prove that non-graded modules are zero on kernel of $\phi(\mu)$, and hence the non-graded modules actually become modules for $\bigoplus \mathfrak{g}_{\text{aff}}$. It is well known that the irreducible integrable modules of $\bigoplus \mathfrak{g}_{\text{aff}}$ are tensor products of irreducible highest weight modules [Rao 2005]. The main result of this paper is Theorem 3.4. Proved in Section 5, it says that any irreducible integrable module for $\tau(\mu)$ with finite-dimensional weight spaces, on which the center acts nontrivially, is an irreducible component of $V(\psi) \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$.

We consider cases $m = 2$ and $m = 3$ separately. The modules for $\tau(\mu)$, where μ is a Dynkin diagram automorphism of order 3, are worked out in Section 6.

1. Preliminaries

Let \mathfrak{g} be a finite-dimensional simple Lie algebra of rank d over the complex field \mathbb{C} . Let \mathfrak{h} be a Cartan subalgebra of \mathfrak{g} . Fix a positive integer $n \geq 2$. Let $A = A_n = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ be a Laurent polynomial ring in n commuting variables t_1, t_2, \dots, t_n . For $\underline{m} = (m_1, \dots, m_n) \in \mathbb{Z}^n$, let $t^{\underline{m}} = t_1^{m_1} \cdots t_n^{m_n} \in A$. Then $\mathfrak{g}_A = \mathfrak{g} \otimes A$ can be made into a Lie algebra by defining

$$[X \otimes t^{\underline{r}}, Y \otimes t^{\underline{s}}] = [X, Y] \otimes t^{\underline{r} + \underline{s}}$$

for $X, Y \in \mathfrak{g}$ and $\underline{r}, \underline{s} \in \mathbb{Z}^n$. Suppose Ω_A is a complex vector space with basis $\{t^{\underline{m}} K_i \mid 1 \leq i \leq n, \underline{m} \in \mathbb{Z}^n\}$, and suppose d_A is the subspace of Ω_A spanned by $\{\sum m_i t^{\underline{m}} K_i \mid \underline{m} \in \mathbb{Z}^n\}$. Let (\cdot, \cdot) be a nondegenerate, symmetric, \mathfrak{g} -invariant bilinear form on \mathfrak{g} . Let $\tau_0 = \mathfrak{g} \otimes A \oplus \Omega_A/d_A$ be a Lie algebra defined by

- $[X \otimes t^{\underline{r}}, Y \otimes t^{\underline{s}}] = [X, Y] \otimes t^{\underline{r} + \underline{s}} + (X, Y) \sum r_i t^{\underline{r} + \underline{s}} K_i$ and
- Ω_A/d_A is central in τ_0 .

The Lie algebra τ_0 is the universal central extension of $\mathfrak{g} \otimes A$ [Morita and Yoshii 1985] and is naturally \mathbb{Z}^n -graded. To reflect this fact, we consider a bigger Lie algebra. Let $\tau = \tau_0 \oplus D$, where D is the complex linear span of derivations d_1, \dots, d_n . We now define a Lie algebra structure on τ by

$$\begin{aligned} [d_i, X \otimes t^{\underline{r}}] &= r_i X \otimes t^{\underline{r}} && \text{for } \underline{r} \in \mathbb{Z}^n \text{ and } 1 \leq i \leq n, \\ [d_i, t^{\underline{r}} K_j] &= r_i t^{\underline{r}} K_j && \text{for } \underline{r} \in \mathbb{Z}^n \text{ and } 1 \leq i, j \leq n, \\ [d_i, d_j] &= 0 && \text{for } 1 \leq i, j \leq n. \end{aligned}$$

We call τ the nontwisted toroidal Lie algebra. We will now define the twisted toroidal Lie algebra. Let μ be a Dynkin diagram automorphism of \mathfrak{g} of order m and ϵ a primitive m -th root of unity. Let $S = \{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$ for $k > 0$. Define an automorphism of τ , again denoted by μ , such that

$$\begin{aligned} \mu(X \otimes t^{\underline{r}}) &= (\epsilon)^{r_{i_1} + \dots + r_{i_k}} \mu(X) \otimes t^{\underline{r}}, \\ \mu(t^{\underline{r}} K_i) &= (\epsilon)^{r_{i_1} + \dots + r_{i_k}} t^{\underline{r}} K_i, \\ \mu(d_i) &= d_i. \end{aligned}$$

It can be easily checked that μ is an automorphism of τ of order m . The subalgebra of μ -fixed points of τ is called the twisted toroidal Lie algebra and is denoted by $\tau(\mu)$. Since μ leaves \mathfrak{h} invariant, we let $\mathfrak{h}(\mu)$ be the μ -fixed points of \mathfrak{h} under μ . Letting $m = 2$, we have the decomposition $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ of \mathfrak{g} , where $\mathfrak{g}_0 = \{X \in \mathfrak{g} \mid \mu(X) = X\}$ and $\mathfrak{g}_1 = \{X \in \mathfrak{g} \mid \mu(X) = \epsilon X\}$.

Suppose $(X, Y) \neq 0$ implies $X, Y \in \mathfrak{g}_0$ or $X, Y \in \mathfrak{g}_1$. Then the center of the twisted toroidal Lie algebra, which we denote by $(\Omega_A/d_A)^0$, will be

$$(\Omega_A/d_A)^0 = \text{span}\{t^m K_i \mid m_{i_1} + \dots + m_{i_k} \cong 0 \pmod{2}\}.$$

Also μ leaves $\mathfrak{g} \otimes A$ invariant. Let $(\mathfrak{g} \otimes A)_\mu$ be the μ -fixed points of $\mathfrak{g} \otimes A$. Then

$$\tau(\mu) = (\mathfrak{g} \otimes A)_\mu \oplus (\Omega_A/d_A)^0 \oplus D.$$

It is known from [Berman and Krylyuk 1995] that $(\mathfrak{g} \otimes A)_\mu \oplus (\Omega_A/d_A)^0$ is the universal central extension of $(\mathfrak{g} \otimes A)_\mu$. Let $\tau_0(\mu) = (\mathfrak{g} \otimes A)_\mu \oplus (\Omega_A/d_A)^0$.

Change of coordinates. We recall that $\text{GL}(n, \mathbb{Z})$, the group of $n \times n$ matrices with entries in \mathbb{Z} and determinant ± 1 , acts naturally on \mathbb{Z}^n . Denote the action as $A\underline{m}$ for $A \in \text{GL}(n, \mathbb{Z})$ and $\underline{m} \in \mathbb{Z}^n$. We define an automorphism on τ by

$$\begin{aligned} A.X \otimes t^m &= X \otimes t^{Am}, \\ A.d(t^r)t^s &= d(t^{Ar})t^{As}, \quad \text{where } d(t^r)t^s = \sum r_i t^{r+s} K_i. \end{aligned}$$

Let $(d_1^1, \dots, d_n^1) = (A^T)^{-1}(d_1, \dots, d_n)$. Then define $A(d_i) = d_i^1$ for $1 \leq i \leq n$. It is easy to check that A is an automorphism on τ . This automorphism is nothing but the change of coordinates. For example, set $s_i = t^{Ae_i}$, where $\{e_1, \dots, e_n\}$ is the standard basis of \mathbb{Z}^n . Then clearly

$$\mathbb{C}[s_1^{\pm 1}, \dots, s_n^{\pm 1}] = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}].$$

Remark 1.1. We will consider $\tau(\mu)$ up to a suitable change of coordinates. In this process, the set S will be changed. In general, the set S can be reduced to a minimum of two element set. Consider an example: Let μ be a Dynkin diagram automorphism of \mathfrak{g} of order 2. Let $S = \{i_1, i_2, \dots, i_k\} \subset \{1, 2, \dots, n\}$. We assume that k is odd. Put

$$s_1 = t_{i_1} t_{i_2} \cdots t_{i_k}, \quad s_2 = t_{i_1}, \quad s_3 = t_{i_1} t_{i_2}, \quad \dots, \quad s_k = t_{i_{k-2}} t_{i_{k-1}}.$$

For $i \in \{1, 2, \dots, n\} - S$, we put $s_i = t_i$. Then it is easy to see that

$$\mathbb{C}[s_1^{\pm 1}, \dots, s_k^{\pm 1}] = \mathbb{C}[t_{i_1}^{\pm 1}, \dots, t_{i_k}^{\pm 1}] \quad \text{and} \quad \mathbb{C}[s_1^{\pm 1}, \dots, s_n^{\pm 1}] = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}].$$

Now the set S changes with new variables s_1, s_2, \dots, s_n , and $S = \{1, 2\}$.

We note that $\tau(\mu)$ depends on the set S . But in this paper, we use a change of coordinates to reduce S to $\{1, n\}$. So we are not including S in the notation of $\tau(\mu)$.

Assume

$$\tilde{\mathfrak{h}} = \mathfrak{h} \oplus \sum_{i=1}^n \mathbb{C}K_i \oplus \sum_{i=1}^n \mathbb{C}d_i,$$

which is a Cartan subalgebra of τ . Consider μ -fixed points $\tilde{\mathfrak{h}}(\mu)$ of $\tilde{\mathfrak{h}}$ under μ . Then $\tilde{\mathfrak{h}}(\mu) = \mathfrak{h}(\mu) \oplus \sum_{i=1}^n \mathbb{C}K_i \oplus \sum_{i=1}^n \mathbb{C}d_i$ is a Cartan subalgebra of $\tau(\mu)$. It is well known that \mathfrak{g}_0 , the fixed point subalgebra of \mathfrak{g} under μ , is a simple Lie algebra [Kac 1990], and $\mathfrak{h}(\mu)$ is a Cartan subalgebra of \mathfrak{g}_0 .

Let $\alpha_1, \alpha_2, \dots, \alpha_l \in \mathfrak{h}(\mu)^*$ be the simple roots of \mathfrak{g}_0 . For $1 \leq i \leq l$, we extend the α_i to $\tilde{\mathfrak{h}}(\mu)^*$ by defining $\alpha_i(K_j) = \alpha_i(d_j) = 0$ for $1 \leq j \leq n$. Let $\alpha_1^\vee, \dots, \alpha_l^\vee \in \mathfrak{h}(\mu)$ be the coroots of \mathfrak{g}_0 . For $1 \leq i \leq n$, define $\delta_i \in \tilde{\mathfrak{h}}(\mu)^*$ and $w_i \in \tilde{\mathfrak{h}}(\mu)^*$ by

$$\begin{aligned} \delta_i(\alpha_j^\vee) &= 0, & w_i(\alpha_j^\vee) &= 0 & \text{for } 1 \leq j \leq l, \\ \delta_i(K_j) &= 0, & w_i(K_j) &= \delta_{ij} & \text{for } 1 \leq j \leq n, \\ \delta_i(d_j) &= \delta_{ij}, & w_i(d_j) &= 0 & \text{for } 1 \leq j \leq n, \end{aligned}$$

Then it is easy to see that $\alpha_1, \alpha_2, \dots, \alpha_l, \delta_1, \dots, \delta_n, w_1, \dots, w_n$ is a \mathbb{C} -basis of $\tilde{\mathfrak{h}}(\mu)^*$.

Recall that the nondegenerate, symmetric, \mathfrak{g} -invariant bilinear form (\cdot, \cdot) remains nondegenerate on \mathfrak{g}_0 as well as on $\mathfrak{h}(\mu)$. The form (\cdot, \cdot) induces a nondegenerate form $\langle \cdot, \cdot \rangle$ on $\tilde{\mathfrak{h}}(\mu)^*$ defined as

$$\begin{aligned} \langle \alpha_i, \alpha_j \rangle &= (\alpha_i, \alpha_j) & \text{for } 1 \leq i, j \leq l, \\ \langle \alpha_i, \delta_j \rangle &= 0 = \langle \alpha_i, w_j \rangle & \text{for } 1 \leq i \leq l \text{ and } 1 \leq j \leq n, \\ \langle \delta_i, \delta_j \rangle &= 0 = \langle w_i, w_j \rangle, \\ \langle \delta_i, w_j \rangle &= \delta_{ij} & \text{for } 1 \leq i, j \leq n. \end{aligned}$$

For $\underline{m} = (m_1, \dots, m_n) \in \mathbb{Z}^n$, define $\delta_{\underline{m}} = \sum m_i \delta_i$ and note that $\langle \delta_{\underline{m}}, \delta_{\underline{m}} \rangle = 0$.

Root system for $\tau(\mu)$. Let $m = 2$. Then

$$\tau(\mu) = \mathfrak{g}_0 \otimes \sum_{m_1 + \dots + m_{i_k} \cong 0 \pmod{2}} \mathbb{C}t^m \oplus \mathfrak{g}_1 \otimes \sum_{m_1 + \dots + m_{i_k} \cong 1 \pmod{2}} \mathbb{C}t^m \oplus (\Omega_A/d_A)^0 \oplus D.$$

Let Δ_0 be the roots of \mathfrak{g}_0 and Δ_0^s be the short roots of \mathfrak{g}_0 . Notice that the nonzero weights of the \mathfrak{g}_0 -module \mathfrak{g}_1 are all short roots of \mathfrak{g}_0 except in $A_{2l}^{(2)}$.

Define

$$\begin{aligned} \Delta_{\text{re}} &= \{ \alpha + \delta_{\underline{m}} \mid \alpha \in \Delta_0, m_{i_1} + \dots + m_{i_k} \cong 0 \pmod{2} \} \\ &\quad \cup \{ \alpha + \delta_{\underline{m}} \mid \alpha \in \Delta_0^s, m_{i_1} + \dots + m_{i_k} \cong 1 \pmod{2} \} \end{aligned}$$

and $\Delta_{\text{im}} = \{ \delta_{\underline{m}} \mid \underline{m} \in \mathbb{Z}^n \}$. Let $\Delta = \Delta_{\text{re}} \cup \Delta_{\text{im}}$. For $\alpha \in \Delta$, let

$$\tau_\alpha = \{ x \in \tau(\mu) \mid [h, x] = \alpha(h)x \text{ for all } h \in \tilde{\mathfrak{h}}(\mu) \}.$$

Then clearly $\tau(\mu) = \bigoplus_{\alpha \in \Delta} \tau_\alpha$ is the root space decomposition of $\tau(\mu)$ with respect to $\tilde{\mathfrak{h}}(\mu)$.

Define

$$\Delta_{\text{re}}^+ = \{\alpha + \delta_{\underline{m}} \mid \alpha > 0, m_n \geq 0 \text{ or } \alpha < 0, m_n \geq 1\}, \quad \Delta_{\text{im}}^+ = \{\delta_{\underline{m}} \mid m_n \geq 1\},$$

$$\Delta^0 = \{\delta_{\underline{m}} \mid m_n = 0\}.$$

Then $\Delta^+ = \Delta_{\text{re}}^+ \cup \Delta_{\text{im}}^+$ and $\Delta^- = -\Delta^+$. Further $\Delta = \Delta^+ \cup \Delta^0 \cup \Delta^-$. Let $\alpha \in \Delta$. Then $\alpha \in \Delta_{\text{re}}$ if and only if $\langle \alpha, \alpha \rangle \neq 0$.

Coroots. Let $\alpha \in \Delta_0$. Then

$$\alpha^\vee = \sum_{i=1}^l n_i (|\alpha_i|^2 / |\alpha|^2) \alpha_i^\vee \quad \text{for } \alpha = \sum_{i=1}^l n_i \alpha_i.$$

It is easy to check that $\alpha(\alpha^\vee) = 2$. Suppose $\alpha \in \Delta_{\text{re}}$ and $\alpha = \beta + \delta_{\underline{m}}$. Then define

$$\alpha^\vee = \beta^\vee + (2/|\beta|^2) \sum m_i K_i.$$

Weyl group of $\tau(\mu)$. For $\alpha \in \Delta_{\text{re}}$, we define a reflection

$$r_\alpha : \tilde{\mathfrak{h}}(\mu)^* \rightarrow \tilde{\mathfrak{h}}(\mu)^*, \quad \lambda \mapsto \lambda - \lambda(\alpha^\vee)\alpha.$$

Let W be the group generated by $r_\alpha, \alpha \in \Delta_{\text{re}}$. Then it is easy to prove that $\langle w\lambda, w\mu \rangle = \langle \lambda, \mu \rangle$ for all $w \in W$. W is called the Weyl group of $\tau(\mu)$.

Definition 1.2. A module V of $\tau(\mu)$ is said to be integrable if

- $V = \bigoplus_{\lambda \in \tilde{\mathfrak{h}}(\mu)^*} V_\lambda$, where $V_\lambda = \{v \in V \mid hv = \lambda(h)v \text{ for all } h \in \tilde{\mathfrak{h}}(\mu)\}$;
- For all $\alpha + \delta_{\underline{m}} \in \Delta_{\text{re}}$ and $v \in V$, there exists an integer $k = k(\alpha, \underline{m}, v)$ such that $(X_\alpha \otimes t^{\underline{m}})^k v = 0$, where X_α is the root vector corresponding to root α .

Lemma 1.3. Suppose V is an irreducible integrable module for $\tau(\mu)$ with finite-dimensional weight spaces. Let $P(V)$ denote the set of all weights of V . Then

- (i) $P(V)$ is W -invariant;
- (ii) $\dim V_\lambda = \dim V_{w\lambda}$ for $w \in W$ and $\lambda \in P(V)$;
- (iii) if $\alpha \in \Delta_{\text{re}}$ and $\lambda \in P(V)$, then $\lambda(\alpha^\vee) \in \mathbb{Z}$;
- (iv) if $\alpha \in \Delta_{\text{re}}$ and $\lambda \in P(V)$, and if $\lambda(\alpha^\vee) > 0$, then $\lambda - \alpha \in P(V)$.
- (v) $\lambda(K_i)$ is a constant integer for all $\lambda \in P(V)$, for $i = 1, 2, \dots, n$.

Proof. The proof is standard. □

From now on we will consider $\tau(\mu)$ up to a change of coordinates. Recall that μ leaves the algebra $B = \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$ invariant. Let B_μ be the μ -fixed points of B . Now B can be seen as a group algebra of \mathbb{Z}^{n-1} . Similarly, B_μ can be seen as a group algebra of subgroup Γ of \mathbb{Z}^{n-1} of index 2. But by Remark 1.1, one can change the basis so that $\Gamma = \underline{2e_1}\mathbb{Z} \oplus \underline{2e_2}\mathbb{Z} \oplus \dots \oplus \underline{2e_{n-1}}\mathbb{Z}$. Thus we conclude that it

is possible to change coordinates so that $S = \{1, n\}$. In this case when $m = 2$, we have for $X \in \mathfrak{g}$ that

$$\begin{aligned} \mu(X \otimes t_1^{2m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n}) &= \mu(X) \otimes t_1^{2m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n}, \\ \mu(X \otimes t_1^{2m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n}) &= -\mu(X) \otimes t_1^{2m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n}, \\ \mu(X \otimes t_1^{2m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n+1}) &= -\mu(X) \otimes t_1^{2m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n+1}, \\ \mu(X \otimes t_1^{2m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n+1}) &= \mu(X) \otimes t_1^{2m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} t_n^{2m_n+1}. \end{aligned}$$

So

$$\begin{aligned} \tau(\mu) = \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m} B_\mu \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m+1} t_1 B_\mu \\ \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m} t_1 B_\mu \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m+1} B_\mu \bigoplus (\Omega_A/d_A)^0 \bigoplus D, \end{aligned}$$

where $B_\mu = \mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$.

Let Q be the \mathbb{Z} -lattice spanned by $\alpha_1, \dots, \alpha_l, \delta_n$. Recall $\mathfrak{g}_0 \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}K_n$ is an affine Lie algebra, which need not be a subalgebra of $\tau(\mu)$. Let β be the highest root of the finite root system Δ_0 . Let $\alpha_0 = -\beta + \delta_n$. Then it is well known that $\{\alpha_1, \dots, \alpha_l, \alpha_0\}$ forms a system of simple roots for $\mathfrak{g}_0 \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}K_n$.

Let $Q(\mu)$ be the root lattice of $\tau(\mu)$. This is a \mathbb{Z} -lattice spanned by $\alpha_1, \dots, \alpha_l, \delta_1, \dots, \delta_n$. For $\lambda, \mu \in Q(\mu)$, we say $\lambda \geq \mu$ if

$$\lambda - \mu = \sum_{i=0}^l n_i \alpha_i \quad \text{for } 0 \leq n_i \in \mathbb{Z}.$$

Hence $\lambda > 0$ means $\lambda \geq 0$ and $\lambda \neq 0$. Now consider the twisted affine Lie algebra, $\mathfrak{g}_0 \otimes \mathbb{C}[t_n^2, t_n^{-2}] \oplus \mathfrak{g}_1 \otimes t_n \mathbb{C}[t_n^2, t_n^{-2}] \oplus \mathbb{C}K_n$, which is also a subalgebra of $\tau(\mu)$. Its positive roots are also positive in the above sense.

2. Central operators

The purpose of this paper is to classify irreducible integrable modules for $\tau(\mu)$ with finite-dimensional weight spaces with respect to the Cartan subalgebra $\tilde{\mathfrak{h}}(\mu)$, where the center acts nontrivially. We first get information on a (fixed) irreducible module V for $\tau(\mu)$ with finite-dimensional weight spaces.

Definition 2.1. A linear map $Z : V \rightarrow V$ is called a central operator of degree $\underline{m} \in \mathbb{Z}^n$ if Z commutes with the $\tau_0(\mu)$ action and

$$d_i Z - Z d_i = m_i Z \quad \text{for all } i.$$

For example, $t^{\underline{m}} K_i$ is a central operator of degree \underline{m} and $t^{\underline{r}} K_i t^{\underline{s}} K_j$ is a central operator of degree $\underline{r} + \underline{s}$.

Lemma 2.2. • Let Z be a central operator of degree \underline{m} such that $Zv \neq 0$ for some $v \in V$. Then $Zw \neq 0$ for every nonzero $w \in V$.

- Let Z be a nonzero central operator of degree \underline{m} . Then there exists a central operator T on V of degree $-\underline{m}$ such that $ZT = TZ = \text{Id}$.
- Let Z_1 and Z_2 be nonzero central operators of degree \underline{m} on V . Then there exists a scalar λ such that $Z_1 = \lambda Z_2$.

Proof. The proofs are easy and given in [Rao 2004]. □

Theorem 2.3. Let V be an irreducible module for $\tau(\mu)$ with finite-dimensional weight spaces with respect to the Cartan subalgebra $\tilde{\mathfrak{h}}(\mu)$. Let

$$L = \{\underline{m} \in \mathbb{Z}^n \mid t^{\underline{m}} K_i \neq 0 \text{ on } V \text{ for some } i\}.$$

Let $\langle L \rangle$ be the subgroup of \mathbb{Z}^n generated by L . Suppose $\text{rank}\langle L \rangle = k \leq n$. Then up to a coordinate change,

- (i) there exist nonzero positive integers k_1, \dots, k_k and nonzero central operators Z_1, Z_2, \dots, Z_k of degrees $(k_1, 0, \dots, 0), \dots, (0, \dots, k_k, 0, \dots, 0)$, respectively;
- (ii) $k < n$;
- (iii) $t^{\underline{r}} K_i \neq 0$ on V implies that $i \geq k + 1$ and $r_{k+1} = \dots = r_n = 0$;
- (iv) there exists a proper submodule W of $\tau_0(\mu) \oplus D_k$, where D_k is the linear span of $\{d_{k+1}, \dots, d_n\}$, such that V/W has finite-dimensional weight spaces with respect to $\mathfrak{h}(\mu) \oplus \sum_{i=1}^n \mathbb{C}K_i \oplus D_k$.

Proof. The proofs are exactly the same as those of [Rao 2004, Theorem 4.5]. □

Theorem 2.4. With the notation above, suppose there is an element in $(\Omega_A/d_A)^0$ that acts nontrivially on V . Then $k = n - 1$.

Proof. The proof is the same as that of [Rao 2004, Proposition 4.8]. □

Proposition 2.5. Let $\bar{\tau}(\mu)$ be the quotient of $\tau(\mu)$ by all nonzero-degree central operators. Suppose \bar{V} is a module for $\bar{\tau}(\mu)$ with finite-dimensional weight spaces. Then the zero-degree central operators K_i for $1 \leq i \leq n$ act trivially on \bar{V} .

Proof. The proof follows from [Rao 2004]. □

Let V be an irreducible integrable module for $\tau(\mu)$ with finite-dimensional weight space. Choose coordinates using Theorems 2.3 and 2.4, so that $t^{\underline{m}} K_i \neq 0$ on V implies $m_n = 0$ and $i = n$.

Proposition 2.6. Let V be as above. Suppose K_n acts as positive integer. Then there exists $\lambda \in P(V)$ such that $\lambda + \eta + \delta_{\underline{m}'} \notin P(V)$ for all $\eta > 0$, where $\delta_{\underline{m}'} = \sum_{i=1}^{n-1} m_i \delta_i$, and for all $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$.

Proof. The proof uses lemmas from [Rao 2004]; these we enclose in brackets.

Notice that $\mathfrak{g}_{0,\text{aff}} = \mathfrak{g}_0 \otimes \mathbb{C}[t_n^2, t_n^{-2}] \oplus \mathbb{C}K_n$ is an affine subalgebra of $\tau(\mu)$. Then V is a module for $\mathfrak{g}_{0,\text{aff}}$. Now, arguing as in [Lemma 2.8], we conclude that

(*) there exists a $\lambda \in P(V)$ such that $\lambda + \eta \notin P(V)$ for all $\eta > 0$.

Next we prove that

(**) $\lambda + \eta + \delta_1 \notin P(V)$ for all $\eta > 0$ or $\lambda + \eta - \delta_1 \notin P(V)$ for all $\eta > 0$.

For proving (**), first we get $\lambda \in P(V)$ such that $\lambda + \eta \notin P(V)$ for all $\eta > 0$. Suppose $\lambda + \eta + \delta_1 \in P(V)$ for some $\eta > 0$. Then, arguing as in [Lemma 2.8], we get $\eta' \geq 0$ such that $\lambda + \eta + \delta_1 + \eta' \in P(V)$ and $\lambda + \eta + \delta_1 + \eta' + \eta'' \notin P(V)$ for all $\eta'' > 0$. Also for all $\eta'' > 0$, $\lambda + \eta + \delta_1 + \eta' + \eta'' - \delta_1 = \lambda + \eta + \eta' + \eta'' \notin P(V)$, since $\eta + \eta' + \eta'' > 0$.

Case 2.6a. $m = 2$.

Suppose that there exists a $\lambda \in P(V)$ such that $\lambda + \eta \notin P(V)$ and $\lambda + \eta + \delta_1 \notin P(V)$ for all $\eta > 0$. If $\lambda + \alpha + \delta_m \notin P(V)$ for all $\alpha + \delta_m \in \Delta_{\text{re}}^+$, then we are done. Suppose $\lambda + \alpha + \delta_m \in P(V)$ for some $\alpha + \delta_m \in \Delta_{\text{re}}^+$. Let $\mu = \lambda + \alpha + \delta_m$.

We claim $\mu + \beta + \delta_s \notin P(V)$ for all $\beta + \delta_s \in \Delta_{\text{re}}^+$. Supposing it is false, let $\mu + \beta + \delta_s \in P(V)$ for some $\beta + \delta_s \in \Delta_{\text{re}}^+$.

Case 2.6a1. $\langle \alpha + \beta + \delta_m + \delta_s, \alpha + \delta_m \rangle > 0$.

Let $\alpha = \alpha' + \delta_{m'}$, where $\delta_{m'} = \sum_{i=1}^{n-1} m_i \delta_i$, and $\beta = \beta' + \delta_{s'}$, where $\delta_{s'} = \sum_{i=1}^{n-1} s_i \delta_i$, where $\alpha', \beta' > 0$ and $\alpha', \beta' \in \mathfrak{g}_{0,\text{aff}}$.

Suppose $\alpha + \delta_{s'} \in \Delta_{\text{re}}$. Then by [Lemma 1.4(2)] we have $\lambda + \alpha + \beta + \delta_m + \delta_s((\alpha + \delta_{s'})^\vee) > 0$, since $\lambda(\alpha + \delta_{s'})^\vee > 0$.

By Lemma 1.3(iv), $\lambda + \alpha + \beta + \delta_m + \delta_s - (\alpha + \delta_{s'}) \in P(V)$. This implies $\lambda + \beta' + \delta_m + \delta_s \in P(V)$. We note that $\beta' + \delta_m + \delta_s > 0$. Since $\alpha + \delta_m$ and $\beta + \delta_s$ are in Δ_{re}^+ , the n -th component m_n of \underline{m} and the n -th component s_n of \underline{s} are ≥ 0 . Hence $\beta' + \delta_m + \delta_s > 0$. So $\lambda + \beta' + \delta_m + \delta_s \in P(V)$, which contradicts (*).

Suppose $\alpha + \delta_{s'} \notin \Delta_{\text{re}}$. Then $\alpha + \delta_{s'} - \delta_1 \in \Delta_{\text{re}}$. Then by [Lemma 1.4(2)] we have $\lambda + \alpha + \beta + \delta_m + \delta_s((\alpha + \delta_{s'} - \delta_1)^\vee) > 0$. Notice that $\lambda(\delta_i) = 0$ for every $1 \leq i \leq n-1$. Thus by Lemma 1.3(iv), $\lambda + \alpha + \beta + \delta_m + \delta_s - \alpha - \delta_{s'} + \delta_1 = \lambda + \beta' + \delta_m + \delta_s + \delta_1 \in P(V)$. Again as above $\beta' + \delta_m + \delta_s > 0$. So $\lambda + \beta' + \delta_m + \delta_s + \delta_1 \in P(V)$, which contradicts (**).

Case 2.6a2. $\langle \alpha + \beta + \delta_m + \delta_s, \beta + \delta_s \rangle > 0$.

This can be done as in Case 2.6a1.

Case 2.6a3. $\langle \alpha + \beta + \delta_m + \delta_s, \alpha + \beta + \delta_m + \delta_s \rangle \geq 0$.

Note that $\langle \alpha + \beta + \delta_m + \delta_s, \alpha + \beta + \delta_m + \delta_s \rangle = \langle \alpha' + \beta', \alpha' + \beta' \rangle$. Thus the cases $\langle \alpha' + \beta', \alpha' \rangle > 0$ and $\langle \alpha' + \beta', \beta' \rangle > 0$ are already done. Note that α', β' are in affine root system and $\langle \cdot, \cdot \rangle$ is a semipositive definite form in the affine root lattice.

So we are left with the case $\langle \alpha' + \beta', \alpha' + \beta' \rangle = 0$. Thus by standard arguments, $\alpha' + \beta' = l\delta_n$ for some $l > 0$, and further $\langle \alpha' + \beta', \alpha' \rangle = \langle \alpha' + \beta', \beta' \rangle = 0$. Now either α' or $\alpha' + \delta_1 \in \Delta_{\text{re}}^+$. In any case, $\lambda(\alpha'^{\vee}) \geq 0$. (Since $\lambda + \alpha', \lambda + \alpha' + \delta_1 \notin P(V)$, by $(*)$ and $(**)$.)

Suppose $\lambda(\alpha'^{\vee}) > 0$.

Subcase 1. Suppose $\alpha + \delta_{\underline{s}'} \in \Delta_{\text{re}}$. Then by [Lemma 1.4(2)] we have $\lambda + \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}}((\alpha + \delta_{\underline{s}'})^{\vee}) > 0$, since $\langle \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}}, \alpha + \delta_{\underline{s}'} \rangle = \langle \alpha' + \beta', \alpha' \rangle = 0$. By Lemma 1.3(iv), $\lambda + \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}} - \alpha - \delta_{\underline{s}'} = \lambda + \beta' + \delta_{\underline{m}} + \delta_{\underline{s}} \in P(V)$, which contradicts $(*)$ since $\beta' + \delta_{\underline{m}} + \delta_{\underline{s}} > 0$.

Subcase 2. Suppose $\alpha + \delta_{\underline{s}'} - \delta_1 \in \Delta_{\text{re}}$. Then by [Lemma 1.4(2)] we have $\lambda + \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}}((\alpha + \delta_{\underline{s}'} - \delta_1)^{\vee}) > 0$. By Lemma 1.3(iv), $\lambda + \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}} - \alpha - \delta_{\underline{s}'} + \delta_1 = \lambda + \beta' + \delta_{\underline{m}} + \delta_{\underline{s}} + \delta_1 \in P(V)$, which contradicts $(**)$ since $\beta' + \delta_{\underline{m}} + \delta_{\underline{s}} > 0$.

Suppose $\lambda(\alpha'^{\vee}) = 0$.

Subcase 1. Suppose $\alpha + \delta_{\underline{s}'} \in \Delta_{\text{re}}$. Then $\lambda((-\beta' + l\delta_n + \delta_{\underline{m}'} + \delta_{\underline{s}'})^{\vee}) = l\lambda(\delta_n) > 0$ since $l > 0$ and $\lambda(\delta_n) = K_n > 0$. Now $\lambda + \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}}((-\beta' + l\delta_n + \delta_{\underline{m}'} + \delta_{\underline{s}'})^{\vee}) > 0$. So by Lemma 1.3(iv), $\lambda + \alpha + \beta + \delta_{\underline{m}} + \delta_{\underline{s}} - \alpha' - \delta_{\underline{m}'} - \delta_{\underline{s}'} = \lambda + \beta' + \delta_{\underline{m}} + \delta_{\underline{s}} \in P(V)$, which contradicts $(*)$ since $\beta' + \delta_{\underline{m}} + \delta_{\underline{s}} > 0$.

Subcase 2. $\alpha + \delta_{\underline{s}'} - \delta_1 \in \Delta_{\text{re}}$ can be handled similarly.

This concludes the proof of Proposition 2.6. \square

3. Classification of integrable $\tau(\mu)$ -modules

Let $\mathfrak{g}_{\text{aff}} = \mathfrak{g} \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}K_n$. Consider $\mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}] \oplus D$ and consider the subalgebra $(\mathfrak{h} \oplus \mathbb{C}K_n) \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$. Let

$$H = \tau(\mu) \cap ((\mathfrak{h} \oplus \mathbb{C}K_n) \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]).$$

Let $\tilde{H} = H \oplus D$. We have proved in Proposition 2.6 that there exists a $\lambda \in P(V)$ such that V_{λ} is a ‘‘highest weight’’ space. In fact it is easy to see that $\bigoplus_{\underline{m}'} V_{\lambda + \delta_{\underline{m}'}}$ is also a highest weight space. Let $\bigoplus_{\underline{m}'} V_{\lambda + \delta_{\underline{m}'}} = V^{\lambda}$. Note that H is a graded abelian Lie algebra and V^{λ} is a graded H -module. Since V is irreducible, it follows by weight arguments that V^{λ} is a graded irreducible H module. From known facts [Rao 2004], it follows that V^{λ} is isomorphic to a subalgebra E of $\mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$ such that each nonzero homogeneous element of E is invertible. Then the H module action on V^{λ} can be described by a \mathbb{Z}^{n-1} graded map from H to E , which we call $\bar{\psi}$. Now consider the evaluation map $E(1) : E \rightarrow \mathbb{C}$ defined by $t^m \mapsto 1$. Denote the composition map $E(1) \circ \bar{\psi}$ as ψ . In what follows, we prove that for such $\bar{\psi}$ and ψ there exist an irreducible module $V(\bar{\psi})$ and a non-graded irreducible module $V(\psi)$, and we prove that $V \cong V(\bar{\psi})$. We will also indicate how to get $V(\psi)$

from $V(\bar{\psi})$, and conversely. Thus our problem reduces to classifying $V(\psi)$, and it suffices to describe ψ , which is a map from $H \rightarrow \mathbb{C}$.

Nongraded and graded modules for $\tau(\mu)$. We recall from Section 1 that $\Delta = \Delta_{\text{re}} \cup \Delta_{\text{im}}$, and $\Delta = \Delta^+ \cup \Delta^0 \cup \Delta^-$. Let $\tau_0(\mu) = J = (\mathfrak{g} \otimes A)_\mu \oplus (\Omega_A/d_A)^0$. Then $J = J^+ \oplus J^0 \oplus J^-$, where

$$J^+ = \bigoplus_{\alpha \in \Delta^+} J_\alpha, \quad J^- = \bigoplus_{\alpha \in \Delta^-} J_\alpha, \quad J^0 = \bigoplus_{\alpha \in \Delta^0} J_\alpha.$$

Note that $H = J^0$. For any Lie algebra \mathfrak{g} , let $U(\mathfrak{g})$ be the universal enveloping algebra.

Let $\psi : J^0 \rightarrow \mathbb{C}$ be a linear map. We make the one-dimensional vector space $\mathbb{C}v$ a $J^+ \oplus J^0 \oplus \mathbb{C}d_n$ -module by defining

$$h.v = \psi(h)v \quad \text{for } h \in J^0, \quad Xv = 0 \quad \text{for } X \in J^+, \quad d_n v = dv \quad \text{for some } d \in \mathbb{C}.$$

We consider the induced Verma module

$$M(\psi) = U(J^-) \otimes_{J^+ \oplus J^0 \oplus \mathbb{C}d_n} \mathbb{C}v.$$

From standard arguments, $M(\psi)$ has a proper maximal submodule and hence a unique irreducible quotient $V(\psi)$, which is the non-graded module for $\tau(\mu)$.

We define

$$\bar{\psi} : J^0 \rightarrow \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}] = A_{n-1}.$$

Let $\underline{m} = (m_1, \dots, m_n) \in \mathbb{Z}^n$, and let $\underline{m}' = (m_1, \dots, m_{n-1}) \in \mathbb{Z}^{n-1}$. Define

$$\bar{\psi}(h \otimes t^{\underline{m}}) = \psi(h \otimes t^{\underline{m}})t^{\underline{m}'}$$

Let $E = \text{image } \bar{\psi}$. Let $\underline{b} = (b_1, b_2, \dots, b_{n-1}) \in \mathbb{C}^{n-1}$. Define a $\tau(\mu)$ -module structure on $V(\psi) \otimes A_{n-1}$ by the action

$$\begin{aligned} X \otimes t^{\underline{m}}(v \otimes t^{\underline{s}}) &= (X \otimes t^{\underline{m}}v) \otimes t^{\underline{s} + \underline{m}'} && \text{for every } v \in V(\psi), \\ & && X \otimes t^{\underline{m}} \in \tau_0(\mu), \quad t^{\underline{s}} \in A_{n-1}, \\ d_i.(v \otimes t^{\underline{s}}) &= (b_i + s_i)v \otimes t^{\underline{s}} && \text{for } 1 \leq i \leq (n-1), \\ d_n.(v \otimes t^{\underline{s}}) &= (d_n v) \otimes t^{\underline{s}}. \end{aligned}$$

Lemma 3.1. *E is an irreducible $J^0 \oplus D$ -module if and only if each nonzero homogeneous element of E is invertible.*

Proof. This follows from methods from [Rao 2004]. □

Proposition 3.2. *Let E, ψ and $\bar{\psi}$ be as above. Suppose every nonzero homogeneous element of E is invertible. Then $V(\psi) \otimes A_{n-1}$ is completely reducible as a $\tau(\mu)$ -module, and all components are isomorphic up to a grade shift.*

Proof. This follows from methods from [Rao 1995]. □

We recall that $\tau(\mu) = J \oplus D$. Let ψ , $\bar{\psi}$ and E be as above. Make E a $J^+ \oplus J^0 \oplus D$ -module by defining that J^+ acts trivially on E and

$$\begin{aligned} h \otimes t^m(t^s) &= \psi(h \otimes t^m)t^{s+m'} && \text{for } t^s \in E, h \otimes t^m \in J^0, \\ d_i.(t^s) &= (b_i + s_i)t^s && \text{for } 1 \leq i \leq n-1, \\ d_n.(t^s) &= 0. \end{aligned}$$

Let $M(\bar{\psi}) = U(J^-) \otimes_{J^+ \oplus J^0 \oplus D} E$. By standard arguments, let $V(\bar{\psi})$ be the irreducible quotient of $M(\bar{\psi})$. $V(\bar{\psi})$ is the graded module for $\tau(\mu)$.

Proposition 3.3. *Suppose E is a $J^0 \oplus D$ -irreducible module. Then a component of $V(\psi) \otimes A_{n-1}$ is isomorphic to $V(\bar{\psi})$ as a $\tau(\mu)$ -module for some suitable \underline{b} .*

Proof. The proof is same as in [Rao 2004, Proposition 3.5]. \square

The following is the main result of this paper.

Theorem 3.4. *Any irreducible, integrable module for $\tau(\mu)$ with finite-dimensional weight spaces, on which the center acts nontrivially, is an irreducible component of $V(\psi) \otimes A_{n-1}$, where $V(\psi)$ is the non-graded module for $\tau(\mu)$.*

Outline of the proof. By Proposition 3.3, there is a one-to-one correspondence between graded and non-graded modules for $\tau(\mu)$. So it suffices to classify the non-graded modules for $\tau(\mu)$. We will prove this theorem in Section 5.

4. Going to the affine Kac–Moody Lie algebra

Let V be an irreducible integrable module for $\tau(\mu)$ with finite-dimensional weight spaces. Also we know that if $t^m K_i \neq 0$ on V , then $m_n = 0$ and $i = n$. We have

$$\begin{aligned} \tau_0(\mu) &= \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m} B_\mu \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m+1} t_1 B_\mu \\ &\quad \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m} t_1 B_\mu \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m+1} B_\mu \bigoplus (\Omega_A/d_A)^0. \end{aligned}$$

Let $\mathfrak{g}_{\text{aff}} = \mathfrak{g} \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}K_n$. We define a map

$$\theta : \tau \rightarrow \mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}] \oplus D$$

by

$$\begin{aligned} X \otimes t^m &\mapsto (X t_n^{m_n}) \otimes t^{\underline{m}'}, \quad \text{where } \underline{m}' = (m_1, \dots, m_{n-1}), \\ t^m K_i &\mapsto \begin{cases} 0 & \text{if } i \neq n \text{ and } m_n = 0, \\ K_n \otimes t^m & \text{if } i = n \text{ and } m_n = 0, \\ 0 & \text{if } m_n \neq 0, \end{cases} \\ d_i &\mapsto d_i \quad \text{for } 1 \leq i \leq n. \end{aligned}$$

It is easy to check that θ defines a Lie algebra homomorphism and that $\ker \theta \cap \tau(\mu)$ is zero on V . Below we give a surjective map ϕ from $\mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$ to $\bigoplus_N \mathfrak{g}_{\text{aff}}$ (the direct sum of N copies of $\mathfrak{g}_{\text{aff}}$) and prove the conditions under which $\phi(\mu) = \phi|_{\tau_0(\mu)}$ is a surjective map from $\tau_0(\mu)$ to $\bigoplus_N \mathfrak{g}_{\text{aff}}$. We also prove in Proposition 5.3 that the non-graded module $V(\psi)$ is zero on the kernel of $\phi(\mu)$. Thus $V(\psi)$ is a module for $\bigoplus_N \mathfrak{g}_{\text{aff}}$, which is an affine Kac–Moody Lie algebra on which the integrable irreducible modules are tensor products of irreducible highest weight modules [Rao 2005].

Let $n \geq 2$ be a positive integer. For each i in $1 \leq i \leq (n - 1)$, let N_i be a positive integer. Let $\underline{a}_i = (a_{i1}, a_{i2}, \dots, a_{iN_i})$ be distinct nonzero complex numbers. Let $I = (i_1, i_2, \dots, i_{n-1})$, where $1 \leq i_j \leq N_j$. Let $\underline{m}' = (m_1, m_2, \dots, m_{n-1}) \in \mathbb{Z}^{n-1}$. Let ϕ be a Lie algebra homomorphism defined by

$$\phi : \mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, \dots, t_{n-1}^{\pm 1}] \rightarrow \left(\bigoplus_{N=N_1 N_2 \dots N_{n-1}\text{-copies}} \mathfrak{g}_{\text{aff}} \right) = \mathfrak{g}_{\text{aff} N},$$

$$X t_1^{m_1} \dots t_{n-1}^{m_{n-1}} \mapsto (a_{1i_1}^{m_1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}} X)_{1 \leq i_1 \leq N_1; \dots; 1 \leq i_{n-1} \leq N_{n-1}},$$

where $X \in \mathfrak{g}_{\text{aff}}$.

Following the same proof as in [Rao 2004, Lemma 3.11(a)], we see that ϕ is surjective. Let $T \subset \{11, 12, \dots, 1N_1\}$ be a maximal set such that $a_{1i}^2 \neq a_{1j}^2$ whenever $1i, 1j \in T$ and $1 \leq i, j \leq N_1$.

Proposition 4.1. *Let $\phi(\mu) = \phi|_{\tau_0(\mu)}$. If the cardinality of T is N_1 , then $\phi(\mu)$ is a surjective Lie algebra homomorphism from $\tau_0(\mu)$ to $\mathfrak{g}_{\text{aff} N}$.*

Proof. We already know that if $m = 2$, then $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$. Now there are five cases.

Case 4.1a. Fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_0$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{2m_n} t_1^{2m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n}, Y_2 t_n^{2m_n}, \dots, Y_N t_n^{2m_n}),$$

where the b_i are complex numbers, $X_i t_n^{2m_n} t_1^{2m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$ and $0 \leq m_j \leq N_j - 1$ for $j = 1, \dots, n - 1$. So, we have to solve the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{2m_n} a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n}, Y_2 t_n^{2m_n}, \dots, Y_N t_n^{2m_n}).$$

Writing this in matrix form, we must prove that the matrix

$$M = (a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq n - 1$, determines

the columns. Observe that M can be written as a tensor product of other matrices as

$$M = (a_{1i_1}^{2m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1} \otimes M_2 \otimes \cdots \otimes M_{n-1},$$

where

$$(1) \quad M_j := (a_{ji_j}^{m_j})_{1 \leq i_j \leq N_j, 0 \leq m_j \leq N_j-1} \quad \text{for } j = 2, \dots, n-1.$$

Let $M_1 = (a_{1i_1}^{2m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. In other upcoming cases, M_1 will be defined differently, but the other M_j will be the same. If the cardinality of T is N_1 , which means the $a_{1i_1}^2$ are all distinct for all $1 \leq i_1 \leq N_1$, then M_1 is a Vandermonde matrix, so it is invertible. Similarly M_2 is a Vandermonde matrix because $a_{2i} \neq a_{2j}$ for all $1 \leq i, j \leq N_2$. So M_2 is invertible. The other M_j for $3 \leq j \leq (n-1)$ are likewise invertible. According to a standard result in linear algebra, the determinant of a tensor product of invertible matrices is nonzero. So M is invertible.

Case 4.1b. Fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_0$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{2m_n+1} t_1^{2m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n+1}, Y_2 t_n^{2m_n+1}, \dots, Y_N t_n^{2m_n+1}),$$

where the b_i are complex numbers and $X_i t_n^{2m_n+1} t_1^{2m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$ and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n-1)$. So we have to solve the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{2m_n+1} a_{1i_1}^{2m_1+1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n+1}, Y_2 t_n^{2m_n+1}, \dots, Y_N t_n^{2m_n+1}).$$

Writing this in the matrix form, we have to prove that the matrix

$$M = (a_{1i_1}^{2m_1+1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$, $1 \leq i_j \leq N_j$ determines rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n-1)$, determines the columns. The matrix M is a tensor product of other matrices:

$$M = (a_{1i_1}^{2m_1+1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1} \otimes M_2 \otimes \cdots \otimes M_{n-1}.$$

Let $M_1 = (a_{1i_1}^{2m_1+1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. Again as above, if the cardinality of T is N_1 , then M_1 is product of two matrices: one is diagonal matrix whose diagonal entries are $a_{11}, a_{12}, \dots, a_{1N_1}$ and other is $(a_{1i_1}^{2m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. So M_1 is invertible because the diagonal matrix is invertible and because the other matrix is a Vandermonde matrix. Similarly the M_j for $2 \leq j \leq (n-1)$ are invertible since they are Vandermonde matrices. Hence M is invertible.

Case 4.1c. Fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_1$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{2m_n} t_1^{2m_1+1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n}, Y_2 t_n^{2m_n}, \dots, Y_N t_n^{2m_n}),$$

where the b_i are complex numbers and $X_i t_n^{2m_n} t_1^{2m_1+1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$ and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n - 1)$. As above we must solve the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{2m_n} a_{1i_1}^{2m_1+1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n}, Y_2 t_n^{2m_n}, \dots, Y_N t_n^{2m_n}).$$

For this we must prove that the matrix

$$M = (a_{1i_1}^{2m_1+1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}}),$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n - 1)$, determines the columns. Again by Case 4.1b, M is invertible if the cardinality of T is N_1 .

Case 4.1d. Fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_1$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{2m_n+1} t_1^{2m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n+1}, Y_2 t_n^{2m_n+1}, \dots, Y_N t_n^{2m_n+1}),$$

where the b_i are complex numbers and $X_i t_n^{2m_n+1} t_1^{2m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$ and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n - 1)$. So we have to solve the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{2m_n+1} a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{2m_n+1}, Y_2 t_n^{2m_n+1}, \dots, Y_N t_n^{2m_n+1}).$$

For this we must prove the matrix

$$M = (a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n - 1)$, determines the columns. Again, by Case 4.1a, M is invertible if the cardinality of T is N_1 .

Case 4.1e. Let $c_1, c_2, \dots, c_N \in \mathbb{C}$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i K_n t_1^{2m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \right) = (c_1 K_n, c_2 K_n, \dots, c_N K_n),$$

where the b_i are complex numbers and $K_n t_1^{2m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$ and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n - 1)$. So we must solve

$$\left(\sum_{i=1}^N b_i K_n a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (c_1 K_n, c_2 K_n, \dots, c_N K_n).$$

So, we have to prove that the matrix

$$M = (a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n - 1)$, determines the columns. Again by Case 4.1a, M is invertible if the cardinality of T is N_1 .

So if the cardinality of T is N_1 , then $\phi(\mu)$ is a surjective Lie algebra homomorphism from $\tau_0(\mu)$ to $\mathfrak{g}_{\text{aff}N}$. This completes the proof of Proposition 4.1. \square

We now define polynomials $P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1})$ by

$$P_1(t_1^2) = \prod_{i=1}^{N_1} (t_1^2 - a_{1i}^2), \quad P_j(t_j) = \prod_{l=1}^{N_j} (t_j - a_{jl}) \quad \text{for } 2 \leq j \leq (n - 1).$$

Notation. Denote by $\langle P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle$ the ideal generated by the polynomials $P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1})$ inside $\mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$.

Lemma 4.2. *If the cardinality of T is N_1 , then $\ker \phi(\mu)$ is given by $I_1 + I_2 + I_3 + I_4 + I_5$, where, for some fixed $m \in \mathbb{Z}$,*

$$\begin{aligned} I_1 &= \mathfrak{g}_0 \otimes t_n^{2m} \langle P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_2 &= \mathfrak{g}_0 \otimes t_n^{2m+1} \langle t_1 P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_3 &= \mathfrak{g}_1 \otimes t_n^{2m} \langle t_1 P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_4 &= \mathfrak{g}_1 \otimes t_n^{2m+1} \langle P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_5 &= K_n \langle P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle. \end{aligned}$$

Proof. We consider the map

$$\begin{aligned} \phi_1 : \mathfrak{g}_0 \otimes t_n^{2m} t^{m'} &\rightarrow \bigoplus_{N=N_1 N_2 \cdots N_{n-1}} \mathfrak{g}_0 \otimes t_n^{2m}, \quad \text{where } t^{m'} \in B_\mu, \\ X t_n^{2m} t^{m'} &\mapsto (X t_n^{2m} a_{1i_1}^{2m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}})_{1 \leq i_1 \leq N_1; \dots; 1 \leq i_{n-1} \leq N_{n-1}}. \end{aligned}$$

Let $M = (N = N_1 N_2 \cdots N_{n-1})$ (the dimension of \mathfrak{g}_0). As proved in Proposition 4.1, one can show that ϕ_1 is surjective. Since ϕ_1 is nothing but the evaluation

map at the roots of $P_1(t_1^2), P_2(t_2), \dots, P_{n-1}(t_{n-1})$, we have $I_1 \subset \ker \phi_1$. Therefore $\dim(\mathfrak{g}_0 \otimes t_n^{2m} t^{m'} / I_1) \geq M$. Consider the space

$$R = \{ \mathfrak{g}_0 \otimes t_n^{2m} t_1^{2m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \mid 0 \leq m_i \leq N_i - 1 \text{ for } 1 \leq i \leq (n-1) \}.$$

Note that ϕ_1 is injective on R since the corresponding matrix is invertible. Also note that any element of $\mathfrak{g}_0 \otimes t_n^{2m} t^{m'}$ can be written as linear combination of elements of R modulo I_1 . Thus $\dim(\mathfrak{g}_0 \otimes t_n^{2m} t^{m'} / I_1) \leq M$, and so $\dim(\mathfrak{g}_0 \otimes t_n^{2m} t^{m'} / I_1) = M$. Therefore I_1 is the kernel of ϕ_1 . Arguing similarly on the other components of $\tau(\mu)$, we get that $I_1 + I_2 + I_3 + I_4 + I_5$ is the kernel of $\phi(\mu)$ if the cardinality of T is N_1 . \square

5. Proof of the main theorem and propositions

Proof of Theorem 3.4. Let V be an irreducible integrable module for $\tau(\mu)$ with finite-dimensional weight spaces. Then, by Theorems 2.3 and 2.4, there exist up to a coordinate change nonzero central operators Z_1, \dots, Z_{n-1} of respective degrees $(k_1, 0, \dots, 0), \dots, (0, \dots, k_{n-1}, 0)$. Let $W = \{Z_i v - v \mid v \in V \text{ for } 1 \leq i \leq (n-1)\}$. We claim that W is a maximal proper $J \oplus \mathbb{C}d_n$ -submodule of V .

Consider $W_1 = \{Z_1 v - v \mid v \in V\}$. W_1 is a proper $J \oplus D_1$ -module. Note that if $i \neq 1$, then d_i commutes with Z_1 , and hence W_1 is d_i -invariant. Clearly W_1 is J -invariant. To see that W_1 is proper, just note that W_1 cannot contain d_1 weight vectors. Now consider $W_2 = \{Z_2 v - v \mid v \in V / W_1\}$. By the above argument, we can see that W_2 is a proper $J \oplus D_2$ -module. Continuing this process, we see that $W = \{Z_i v - v \mid v \in V \text{ for } 1 \leq i \leq (n-1)\}$ is a maximal proper $J \oplus \mathbb{C}d_n$ -submodule of V .

Thus we have proved that W is a maximal proper $J \oplus \mathbb{C}d_n$ -submodule of V and V/W has finite-dimensional weight spaces with respect to $\mathfrak{h}(\mu) \oplus \sum_{i=1}^n \mathbb{C}K_i \oplus \mathbb{C}d_n$. Further, we proved in Proposition 2.6 that there exists a $\lambda \in P(V)$ such that $\lambda + \eta + \delta_{\underline{m}'} \notin P(V)$ for every $\eta > 0$ and for every $\underline{m}' \in \mathbb{Z}^{n-1}$.

Let $\bigoplus_{\underline{m}'} V_{\lambda + \delta_{\underline{m}'}} = V^\lambda$. Then it follows that

$$X_\alpha \otimes t^\underline{s} V^\lambda = 0 \quad \text{for all } \alpha + \delta_{\underline{s}} \in \Delta_{\text{re}}^+.$$

Let \bar{V}^λ be the image of V^λ in V/W , which is known to be finite-dimensional. J^0 leaves \bar{V}^λ invariant. Notice that J^0 is an abelian Lie algebra. Thus J^0 has a one-dimensional invariant subspace in \bar{V}^λ . Let W^λ be the maximal proper invariant subspace of \bar{V}^λ . Then it is easy to check that $\bar{V}^\lambda / W^\lambda$ is one-dimensional by Lie's theorem; otherwise there will be a contradiction to the fact that W^λ is a maximal proper subspace. Let $\bar{V}^\lambda / W^\lambda = \mathbb{C}v$. Let $X \otimes t^m v = \psi(X \otimes t^m)v$ for $X \otimes t^m \in J^0$, where $\psi : J^0 \rightarrow \mathbb{C}$ is a linear map. Now consider the irreducible module $V(\psi)$ as described in Section 3.

We claim that $V(\psi)$ is a $\tau_0(\mu) \oplus \mathbb{C}d_n$ -irreducible quotient of V/W . This follows from the fact that the module generated by W^λ is proper and maximal. Thus we have an irreducible non-graded module $V(\psi)$ for $\tau_0(\mu) \oplus \mathbb{C}d_n$. Now $V(\psi) \otimes A_{n-1}$ can be made as a $\tau(\mu)$ -module as described in Section 3. Now by Propositions 3.2 and 3.3, it follows that any irreducible integrable module for $\tau(\mu)$ with finite-dimensional weight spaces is an irreducible component of $V(\psi) \otimes A_{n-1}$. This completes the proof of Theorem 3.4. \square

In case $m = 2$, we have $B_\mu = \mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$ and $\Gamma = 2e_1\mathbb{Z} \oplus e_2\mathbb{Z} \oplus \dots \oplus e_{n-1}\mathbb{Z}$. Notice that $t^{\underline{m}'} \in B_\mu$ if and only if $\underline{m}' \in \Gamma$. Let

$$\begin{aligned} T_1 &= \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m} B_\mu, & T_2 &= \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m+1} t_1 B_\mu, \\ T_3 &= \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m} t_1 B_\mu, & T_4 &= \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m+1} B_\mu. \end{aligned}$$

So $\tau(\mu) = T_1 \oplus T_2 \oplus T_3 \oplus T_4 \oplus (\Omega_A/d_A)^0 \oplus D$. Let $T_i^+ = T_i \cap (\bigoplus_{\alpha \in \Delta^+} \tau_\alpha)$. Let $\alpha_1, \dots, \alpha_l$ be all simple roots of \mathfrak{g}_0 , and let $\alpha_{i_1}, \dots, \alpha_{i_s}$ be all positive short roots of \mathfrak{g}_1 . Let

$$\begin{aligned} S = \{ & \alpha_1 + \delta_{\underline{m}'}, \dots, \alpha_l + \delta_{\underline{m}'}, \underline{m}' \in \Gamma, \\ & -\alpha_1 + \delta_n + \delta_1 + \delta_{\underline{m}'}, \dots, -\alpha_l + \delta_n + \delta_1 + \delta_{\underline{m}'}, \delta_n + \delta_1 + \delta_{\underline{m}'}, \underline{m}' \in \Gamma, \\ & \alpha_{i_1} + \delta_1 + \delta_{\underline{m}'}, \dots, \alpha_{i_s} + \delta_1 + \delta_{\underline{m}'}, \underline{m}' \in \Gamma, \\ & -\alpha_{i_1} + \delta_n + \delta_{\underline{m}'}, \dots, -\alpha_{i_s} + \delta_n + \delta_{\underline{m}'}, \underline{m}' \in \Gamma \}. \end{aligned}$$

Proposition 5.1. *Let $\langle S \rangle$ be the Lie algebra generated by the root spaces corresponding to root $\alpha \in S$. Then*

$$\langle S \rangle = T_1^+ \oplus T_2^+ \oplus T_3^+ \oplus T_4^+ = \bigoplus_{\alpha \in \Delta^+} \tau_\alpha.$$

Proof. Let

$$\tau_1(\mu) = \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{2m} \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{2m+1} \bigoplus \mathbb{C}K_n,$$

which is a twisted affine Lie algebra. Let β be the weight corresponding to the highest weight vector of the \mathfrak{g}_0 -module \mathfrak{g}_1 as in [Kac 1990]. Then it is known from [Kac 1990] that $\tau_1(\mu)$ is generated as a Lie algebra by the root spaces corresponding to the root basis $\{\alpha_1, \dots, \alpha_l, -\beta + \delta_n\}$. Thus from this we conclude that $\langle S \rangle$ contains $T_1^+ \oplus T_4^+$, which is the Lie algebra generated by the root spaces corresponding to the roots $\alpha_1 + \delta_{\underline{m}'}, \dots, \alpha_l + \delta_{\underline{m}'}, -\beta + \delta_n + \delta_{\underline{m}'}$ for $\underline{m}' \in \Gamma$. Also

$$[\mathfrak{h}(\mu)t_1t_n, T_1^+] = T_2^+ \quad \text{and} \quad [\mathfrak{h}(\mu)t_1t_n, T_4^+] = t_n^2 T_3^+,$$

and the root spaces corresponding to the roots $\alpha_{i_1} + \delta_1 + \delta_{\underline{m}'}, \dots, \alpha_{i_s} + \delta_1 + \delta_{\underline{m}'}$ for $\underline{m}' \in \Gamma$ are in $\langle S \rangle$ by definition. Thus $\langle S \rangle = T_1^+ \oplus T_2^+ \oplus T_3^+ \oplus T_4^+$. \square

We will now define the concept of the height ht of a positive root. All roots of \mathfrak{g}_0 are given the usual height. Thus if $\alpha = \sum n_i \alpha_i$, then $\text{ht } \alpha = \sum n_i$. Also $\text{ht } \delta_n = N + 1$, where N is the height of the highest root of the finite root system Δ_0 of \mathfrak{g}_0 . Let $\alpha \in \Delta^+$. Let $\alpha = \alpha' + \delta_{\underline{m}'}$ and $m_n = 0$. Then we define $\text{ht } \alpha = \text{ht } \alpha'$. Let

$$\bar{S} = \{\alpha_1, \dots, \alpha_l, -\alpha_1 + \delta_n + \delta_1, \dots, -\alpha_l + \delta_n + \delta_1, \delta_1 + \delta_n, \alpha_{i_1} + \delta_1, \dots, \alpha_{i_s} + \delta_1, -\alpha_{i_1} + \delta_n, \dots, -\alpha_{i_s} + \delta_n\}.$$

Clearly $\text{ht } \gamma \leq N + 1$ for every $\gamma \in \bar{S}$. It is easy to see that for $\gamma' \in \Delta^+$, there exists a $\gamma \in \bar{S}$ such that $\gamma' = \gamma + \delta_{\underline{m}'}$ for $\underline{m}' \in \Gamma$ and $\text{ht } \gamma \leq N + 1$.

Lemma 5.2. *Suppose $V(\psi)$ has finite-dimensional weight spaces. Then $X_\alpha \otimes IV(\psi) = 0$ for some cofinite ideal I of $\mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1} \dots t_{n-1}^{\pm 1}]$ and for all $\alpha \in \Delta$, including $\alpha = 0$.*

Proof. Let X_α be a root vector of root $\alpha \in -\bar{S}$. For $2 \leq i \leq (n - 1)$, consider the set $\{X_\alpha \otimes t_i^k v \mid k \in \mathbb{Z}\}$, which is contained in the same weight space. Since weight spaces are finite-dimensional, there exists a nonzero polynomial $P_{i,\alpha}(t_i) = \sum_j a_j t_i^j$ such that

$$X_\alpha \otimes P_{i,\alpha}(t_i)v = 0, \quad \text{where } X_\alpha \otimes P_{i,\alpha}(t_i) = \sum_j a_j X_\alpha \otimes t_i^j.$$

Denote by (P) the ideal generated by the polynomial P in $\mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$.

Claim 5.2a. $X_\alpha \otimes (P_{i,\alpha}(t_i))v = 0$ for $2 \leq i \leq n - 1$. Let $\underline{m}' \in B_\mu$. Then

$$h \otimes t^{\underline{m}'}(X_\alpha \otimes P_{i,\alpha}(t_i))v = X_\alpha \otimes P_{i,\alpha}(t_i)h \otimes t^{\underline{m}'}v + \alpha(h)X_\alpha \otimes t^{\underline{m}'}P_{i,\alpha}(t_i)v = 0.$$

The first term is zero since $h \otimes t^{\underline{m}'}$ acts as scalar. Thus

$$X_\alpha \otimes t^{\underline{m}'}(P_{i,\alpha}(t_i))v = 0 \quad \text{for } 2 \leq i \leq n - 1.$$

Let $P_i(t_i) = \prod_{-\alpha \in \bar{S}} P_{i,\alpha}(t_i)$. We note that $X_\alpha \otimes B_\mu(P_i(t_i))v = 0$ for $2 \leq i \leq n - 1$. Similarly for t_1 , consider the set $\{X_\alpha \otimes t_1^{2k} v \mid k \in \mathbb{Z}\}$, which is contained in the same weight space. Since weight spaces are finite-dimensional, there exists a nonzero polynomial $P_{1,\alpha}(t_1) = \sum_j a_{2j} t_1^{2j}$ such that $X_\alpha \otimes P_{1,\alpha}(t_1)v = 0$. On similar lines as above, we prove that

$$X_\alpha \otimes t^{\underline{m}'}(P_{1,\alpha}(t_1))v = 0 \quad \text{for } \underline{m}' \in \Gamma.$$

So we have proved that $X_\alpha \otimes B_\mu(P_i(t_i))v = 0$ for $1 \leq i \leq n - 1$.

Claim 5.2b. For $1 \leq i \leq n-1$,

$$X_\gamma \otimes (P_i(t_i))v = 0 \quad \text{for every } \gamma \in -\Delta^+, \quad \text{where } \gamma = \eta + \delta_{\underline{m}'} \text{ and } m_n = 0.$$

We will prove this by induction on the height of a positive root. We have already seen this by Claim 5.2a for all $\gamma \in -\bar{S}$ such that $\text{ht } \gamma \leq \text{ht } \delta_n$. Let $\gamma \in -\Delta^+$, where $\gamma = \eta + \delta_{\underline{m}'}$ and $m_n = 0$, and consider $X_\gamma \otimes (P_i(t_i))v$ for $1 \leq i \leq (n-1)$. We will prove that $X_\gamma \otimes (P_i(t_i))v$ is killed by X_β for every $\beta \in S$. Then by Proposition 5.1, it follows that $X_\gamma \otimes (P_i(t_i))v$ is killed by X_β for every $\beta \in \Delta^+$, since S generates all the positive root spaces. So $X_\gamma \otimes (P_i(t_i))v$ is a highest weight vector. Since $V(\psi)$ is an irreducible highest weight module, it follows that $X_\gamma \otimes (P_i(t_i))v = 0$ for every $\gamma \in -\Delta^+$, where $\gamma = \eta + \delta_{\underline{m}'}$ and $m_n = 0$.

Let $\beta \in S$, and let $\beta = \alpha + \delta_{\underline{m}'}$ for $\alpha \in \bar{S}$ and $\underline{m}' \in \Gamma$. Then

$$X_\alpha \otimes t^{\underline{m}'}.X_\gamma \otimes (P_i(t_i))v = X_\gamma \otimes (P_i(t_i)).X_\alpha \otimes t^{\underline{m}'}v + [X_\alpha, X_\gamma] \otimes t^{\underline{m}'}(P_i(t_i))v.$$

The first component is zero since v is a highest weight vector. The second is zero by induction on the height of a positive root. This proves Claim 5.2b.

Claim 5.2c. $b \otimes (P_i(t_i))v = 0$ for every $b \in J^0$ and for $1 \leq i \leq (n-1)$.

For any $\alpha \in -\Delta_{\text{re}}^+$, consider $-\alpha^\vee \otimes (P_i(t_i))v = X_{-\alpha}X_\alpha \otimes (P_i(t_i))v - X_\alpha \otimes (P_i(t_i))X_{-\alpha}v = 0$ by Claim 5.2b. Since α^\vee for $\alpha \in -\Delta_{\text{re}}^+$ will cover all of J^0 , this proves Claim 5.2c.

Let I be the ideal $P_1(t_1), \dots, P_{n-1}(t_{n-1})$ generates inside $\mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1} \dots t_{n-1}^{\pm 1}]$. It is easy to see that $\mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1} \dots t_{n-1}^{\pm 1}]/I$ is finite-dimensional. We consider

$$W = \{w \in V(\psi) \mid X_\alpha \otimes Iw = 0 \text{ for all } \alpha, \text{ including } \alpha = 0\},$$

which is nonempty by the above. It is easy to see that W is a submodule of $V(\psi)$. Since $V(\psi)$ is irreducible, $W = V(\psi)$. This completes the proof of Lemma 5.2. \square

Let I' be the ideal generated by the polynomials $P_1(t_1^2), \dots, P_{n-1}(t_{n-1})$ inside $\mathbb{C}[t_1^{\pm 2}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$. That is, let

$$I' = \left\langle P_1(t_1^2) = \prod_{i=1}^{N_1} (t_1^2 - a_{1i}^2), P_j(t_j) = \prod_{l=1}^{N_j} (t_j - a_{jl}) \quad \text{for } 2 \leq j \leq (n-1) \right\rangle.$$

We observe that the polynomials $P_1(t_1^2), \dots, P_{n-1}(t_{n-1})$ have distinct roots. So $I \subset I'$. Our aim is to prove that $X_\alpha \otimes I'v = 0$ for all α , including $\alpha = 0$.

Let α be a simple root of \mathfrak{g}_0 .

Case 5.2d. Suppose α is a long root.

Let $X_\alpha \in (\mathfrak{g}_0)_\alpha$ and $Y_\alpha \in (\mathfrak{g}_0)_{-\alpha}$ such that $h_\alpha = [X_\alpha, Y_\alpha]$ is an \mathfrak{sl}_2 -copy. Let G_α be the space spanned by $X_\alpha \otimes I'$, $Y_\alpha \otimes I'$, $h_\alpha \otimes I'$, and $K_n \otimes I'$.

Case 5.2e. Suppose α is a short root.

Let $X_\alpha \in (\mathfrak{g}_0)_\alpha$ and $Y_\alpha \in (\mathfrak{g}_0)_{-\alpha}$ such that $h_\alpha = [X_\alpha, Y_\alpha]$ is an \mathfrak{sl}_2 -copy. Let $X'_\alpha \in (\mathfrak{g}_1)_\alpha$ and $Y'_\alpha \in (\mathfrak{g}_1)_{-\alpha}$ such that $h'_\alpha = [X'_\alpha, Y'_\alpha]$ is an \mathfrak{sl}_2 -copy. Let G_α be the space spanned by

$$\begin{aligned} X_\alpha \otimes I', & \quad Y_\alpha \otimes I', & \quad h_\alpha \otimes I', & \quad X'_\alpha \otimes t_1 I', \\ Y'_\alpha \otimes t_1^{-1} I', & \quad h'_\alpha \otimes I', & \quad [X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes I', & \quad [X'_\alpha \otimes t_1, Y_\alpha] \otimes I'. \end{aligned}$$

Case 5.2f. Suppose β is the highest root of the finite root system Δ_0 of \mathfrak{g}_0 .

Consider

$$\alpha = -\beta + \delta_n, \quad X_\alpha = X_{-\beta} \otimes t_n, \quad Y_\alpha = X_\beta \otimes t_n^{-1}, \quad h_\alpha = [X_\alpha, Y_\alpha] = h_\beta + K_n.$$

Let G_α be the space spanned by $X_\alpha \otimes I'$, $Y_\alpha \otimes I'$, $h_\alpha \otimes I'$.

Proposition 5.3. *Let α be as in Cases 5.2d, 5.2e, and 5.2f. Then $G_\alpha v = 0$.*

Proof. If we take Lie brackets in G_α , then the result will contain higher powers of I' . But $I \subset I^m$ for large n . So G_α/I is solvable. In fact it is nilpotent. Since I'/I is finite-dimensional, so is G_α/I' . By Lie's theorem, there exists nonzero w in $V(\psi)$ such that $g \otimes p(t)w = \lambda(g, p(t))w$ for some scalar $\lambda(g, p(t))$, $g \in G_\alpha$, and $p(t) \in I'$.

Claim 5.3a. $G_\alpha w = 0$.

We will prove the proposition for Case 5.2e. For Cases 5.2d and 5.2f, the proof is based on similar arguments. So suppose α is a short root and let $X_\alpha \in (\mathfrak{g}_0)_\alpha$. Then let $X_\alpha \otimes P(t)w = \lambda(X_\alpha, P(t))w$ for $P(t) \in I'$. This implies $(X_\alpha \otimes P(t))^m w = \lambda(X_\alpha, P(t))^m w$. Since V is integrable, $(X_\alpha \otimes P(t))^m w = 0$ for large m , and therefore $\lambda(X_\alpha, P(t)) = 0$. Hence $X_\alpha \otimes P(t)w = 0$. Similarly we can prove that $Y_\alpha \otimes P(t)w = 0$. We claim that $h_\alpha \otimes P(t)w = 0$, which follows from the arguments in [Rao 2001, Proposition 2.1], except that we replace the finite-dimensional V by integrable V .

Now let $X'_\alpha \otimes Q(t)t_1 w = \lambda(X'_\alpha, Q(t)t_1)w$ for $Q(t) \in I'$. Then

$$(X'_\alpha \otimes Q(t)t_1)^m w = \lambda(X'_\alpha, Q(t)t_1)^m w.$$

Since V is integrable, $(X'_\alpha \otimes Q(t)t_1)^m w = 0$. Hence $\lambda(X'_\alpha, Q(t)t_1)^m = 0$, and therefore $\lambda(X'_\alpha, Q(t)t_1) = 0$. This implies $X'_\alpha \otimes Q(t)t_1 w = 0$. Similarly we can prove that $Y'_\alpha \otimes Q(t)t_1^{-1} w = 0$.

Claim 5.3b. $h'_\alpha \otimes Q(t)w = 0$.

Let $h'_\alpha \otimes Q(t)w = \lambda(h'_\alpha, Q(t))w$ and let $\lambda = \lambda(h'_\alpha, Q(t))$.

Subclaim 1. $h'_\alpha \otimes Q(t)(Y'_\alpha \otimes t_1^{-1})^m w = \lambda(Y'_\alpha \otimes t_1^{-1})^m w$.

We prove this by induction on m . Let $m = 1$ and consider

$$h'_\alpha \otimes Q(t)(Y'_\alpha \otimes t_1^{-1})w = (Y'_\alpha \otimes t_1^{-1})h'_\alpha \otimes Q(t)w - 2Y'_\alpha \otimes t_1^{-1}Q(t)w = \lambda(Y'_\alpha \otimes t_1^{-1})w.$$

Now we assume the claim for m and consider

$$\begin{aligned} & h'_\alpha \otimes Q(t)(Y'_\alpha \otimes t_1^{-1})^{m+1}w \\ &= (Y'_\alpha \otimes t_1^{-1})(h'_\alpha \otimes Q(t))(Y'_\alpha \otimes t_1^{-1})^m w + [h'_\alpha \otimes Q(t), Y'_\alpha \otimes t_1^{-1}](Y'_\alpha \otimes t_1^{-1})^m w \\ &= \lambda(Y'_\alpha \otimes t_1^{-1})^{m+1}w - 2Y'_\alpha \otimes Q(t)t_1^{-1}(Y'_\alpha \otimes t_1^{-1})^m w \\ &= \lambda(Y'_\alpha \otimes t_1^{-1})^{m+1}w, \end{aligned}$$

which follows by induction and the fact that $Y'_\alpha \otimes Q(t)t_1^{-1}w = 0$.

Subclaim 2. $X'_\alpha \otimes Q(t)t_1(Y'_\alpha \otimes t_1^{-1})^m w = m\lambda(Y'_\alpha \otimes t_1^{-1})^{m-1}w$.

The proof is by induction on m . Let $m = 1$. Then

$$X'_\alpha \otimes Q(t)t_1(Y'_\alpha \otimes t_1^{-1})w = Y'_\alpha \otimes t_1^{-1}.X'_\alpha \otimes Q(t)t_1w + h'_\alpha \otimes Q(t)w = \lambda w.$$

Now assume the claim for m , and consider

$$\begin{aligned} & (X'_\alpha \otimes Q(t)t_1)(Y'_\alpha \otimes t_1^{-1})^{m+1}w \\ &= (Y'_\alpha \otimes t_1^{-1})(X'_\alpha \otimes Q(t)t_1)(Y'_\alpha \otimes t_1^{-1})^m w + h'_\alpha \otimes Q(t)(Y'_\alpha \otimes t_1^{-1})^m w \\ &= m\lambda(Y'_\alpha \otimes t_1^{-1})^m w + \lambda(Y'_\alpha \otimes t_1^{-1})^m w \\ &= (m+1)\lambda(Y'_\alpha \otimes t_1^{-1})^m w. \end{aligned}$$

Since V is integrable, there exists an $n_0 > 0$ such that $(Y'_\alpha \otimes t_1^{-1})^{n_0}w = 0$, and $(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \neq 0$. Then

$$\begin{aligned} \lambda(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w &= h'_\alpha \otimes Q(t)(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \\ &= [X'_\alpha \otimes Q(t)t_1, Y'_\alpha \otimes t_1^{-1}](Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \\ &= -(n_0 - 1)\lambda(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w. \end{aligned}$$

This implies that either $n_0 = 0$ or $\lambda = 0$. But, by choice, $n_0 \neq 0$, and hence $\lambda = 0$.

This proves that $h'_\alpha \otimes Q(t)w = 0$.

Claim 5.3c. $[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)w = 0$ for $Q'(t) \in I'$.

$$\text{Let } [X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)w = \lambda w.$$

Subclaim 3. $[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})^m w = \lambda(Y'_\alpha \otimes t_1^{-1})^m w$.

We prove this for $m = 1$:

$$\begin{aligned} & [X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})w \\ &= (Y'_\alpha \otimes t_1^{-1})[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)w + [[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}] \\ &= \lambda(Y'_\alpha \otimes t_1^{-1})w. \end{aligned}$$

This follows by the assumption and the fact that

$$[[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}] = 0.$$

In fact $X_\alpha \in \mathfrak{g}_0$ and $Y'_\alpha \in \mathfrak{g}_1$ implies that $[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t) \in \mathfrak{g}_{1,0} \otimes t_1^{-1}Q'(t)$. So $[[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}] \in \mathfrak{g}_{0,\alpha} \otimes t_1^{-2}Q'(t)$. Since $\dim \mathfrak{g}_{0,\alpha} = 1$, we have $[[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}] \in X_\alpha \otimes t_1^{-2}Q'(t) = 0$, because $X_\alpha \otimes I' = 0$. Now assume the claim for m and consider

$$\begin{aligned} & [X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})^{m+1}w \\ &= (Y'_\alpha \otimes t_1^{-1})[X_\alpha, Y'_\alpha \otimes t_1^{-1}]Q'(t)(Y'_\alpha \otimes t_1^{-1})^m w \\ &\quad + [[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}](Y'_\alpha \otimes t_1^{-1})^m w \\ &= \lambda(Y'_\alpha \otimes t_1^{-1})^{m+1}w, \end{aligned}$$

because $[[X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}] = 0$, as proved above.

Subclaim 4. $(X_\alpha \otimes Q'(t))(Y'_\alpha \otimes t_1^{-1})^m w = m\lambda(Y'_\alpha \otimes t_1^{-1})^{m-1}w$.

We prove this by induction on m . Let $m = 1$, and consider

$$X_\alpha \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})w = Y'_\alpha \otimes t_1^{-1}X_\alpha \otimes Q'(t)w + [X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)w = \lambda w,$$

because $X_\alpha \otimes Q'(t)w = 0$.

Let Subclaim 4 be true for m and consider

$$\begin{aligned} & X_\alpha \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})^{m+1}w \\ &= (Y'_\alpha \otimes t_1^{-1})(X_\alpha \otimes Q'(t))(Y'_\alpha \otimes t_1^{-1})^m w + [X_\alpha \otimes Q'(t), Y'_\alpha \otimes t_1^{-1}](Y'_\alpha \otimes t_1^{-1})^m w \\ &= m\lambda(Y'_\alpha \otimes t_1^{-1})^m w + \lambda(Y'_\alpha \otimes t_1^{-1})^m w \\ &= (m+1)\lambda(Y'_\alpha \otimes t_1^{-1})^m w \quad \text{(by Subclaim 3)}. \end{aligned}$$

Since V is integrable, there exists an $n_0 > 0$ such that $(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \neq 0$ but $(Y'_\alpha \otimes t_1^{-1})^{n_0}w = 0$. Now

$$\begin{aligned} \lambda(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w &= [X_\alpha, Y'_\alpha \otimes t_1^{-1}] \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \\ &= X_\alpha \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \\ &\quad - (Y'_\alpha \otimes t_1^{-1})X_\alpha \otimes Q'(t)(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w \\ &= -(n_0 - 1)\lambda(Y'_\alpha \otimes t_1^{-1})^{n_0-1}w, \end{aligned}$$

which means either $n_0 = 0$ or $\lambda = 0$. But by our choice $n_0 \neq 0$, and hence $\lambda = 0$. This proves Claim 5.3c. Similarly we can prove that $[X'_\alpha \otimes t_1, Y_\alpha] \otimes Q'(t)w = 0$ for $Q'(t) \in I'$.

Now consider the nonzero submodule $W = \{w \in V(\psi) \mid G_\alpha w = 0\}$ of $V(\psi)$. Since $V(\psi)$ is irreducible, we have $W = V(\psi)$. Thus the module $V(\psi)$ is zero on I' . Let

$$I'' = \left\langle P_1(t_1^2) = t_1 \prod_{i=1}^{N_1} (t_1^2 - a_{1i}^2), P_j(t_j) = \prod_{l=1}^{N_j} (t_j - a_{jl}) \quad \text{for } 2 \leq j \leq (n-1) \right\rangle.$$

In a similar way as above, we can prove that the module $V(\psi)$ is zero on I'' . So the module $V(\psi)$ is zero on kernel $\phi(\mu)$ by Lemma 4.2. Thus $V(\psi)$ is a module for $\bigoplus_N \mathfrak{g}_{\text{aff}}$. \square

6. When the diagram automorphism is of order 3

We now consider the case when μ is a diagram automorphism of \mathfrak{g} of order 3. We will prove analogues of Proposition 4.1 and Lemma 4.2 in this case.

In this case also, μ leaves \mathfrak{h} invariant. Let $\mathfrak{h}(\mu)$ be the μ -fixed points of \mathfrak{h} , and let ϵ be a primitive third root of unity. In this case $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_{-1}$, where \mathfrak{g}_i is the eigenspace of μ for the eigenvalue ϵ^i for $i \in \{0, 1, -1\}$. Suppose $(X, Y) \neq 0$ implies $X, Y \in \mathfrak{g}_0$ or $X \in \mathfrak{g}_1$ and $Y \in \mathfrak{g}_{-1}$. Because of this, the center in the twisted toroidal Lie algebra, which we denote by $(\Omega_A/d_A)^0$ will be

$$(\Omega_A/d_A)^0 = \text{span}\{t^m K_i \mid m_{i_1} + \dots + m_{i_k} \cong 0 \pmod{3}\}.$$

Also μ leaves $\mathfrak{g} \otimes A$ invariant. Let $(\mathfrak{g} \otimes A)_\mu$ be the μ -fixed points of $\mathfrak{g} \otimes A$. Then $\tau(\mu) = (\mathfrak{g} \otimes A)_\mu \oplus (\Omega_A/d_A)^0 \oplus D$. It is known from [Berman and Krylyuk 1995] that $(\mathfrak{g} \otimes A)_\mu \oplus (\Omega_A/d_A)^0$ is the universal central extension of $(\mathfrak{g} \otimes A)_\mu$. In this case also we choose the coordinates in such a way that $S = \{1, n\}$. Here

$$\begin{aligned} \tau(\mu) = & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{3m+1} t_1^{3m_1+2} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \\ & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{3m+2} t_1^{3m_1+1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{3m} t_1^{3m_1+2} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \\ & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{3m+1} t_1^{3m_1+1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{3m+2} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \\ & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_{-1} \otimes t_n^{3m} t_1^{3m_1+1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_{-1} \otimes t_n^{3m+1} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \\ & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_{-1} \otimes t_n^{3m+2} t_1^{3m_1+2} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} (\Omega_A/d_A)^0 \bigoplus D. \end{aligned}$$

Also let

$$\begin{aligned}
 \tau_0(\mu) = & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{3m+1} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \\
 & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_0 \otimes t_n^{3m+2} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{3m} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \\
 & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{3m+1} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_1 \otimes t_n^{3m+2} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \\
 & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_{-1} \otimes t_n^{3m} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_{-1} \otimes t_n^{3m+1} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \\
 & \bigoplus_{m \in \mathbb{Z}} \mathfrak{g}_{-1} \otimes t_n^{3m+2} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} & \bigoplus_{m \in \mathbb{Z}} (\Omega_A/d_A)^0.
 \end{aligned}$$

Root system for $\tau(\mu)$. Let $m = 3$. Then

$$\begin{aligned}
 \tau(\mu) = & \mathfrak{g}_0 \otimes \sum_{m_{i_1} + \cdots + m_{i_k} \cong 0 \pmod{3}} \mathbb{C} t^m \bigoplus \mathfrak{g}_1 \otimes \sum_{m_{i_1} + \cdots + m_{i_k} \cong 1 \pmod{3}} \mathbb{C} t^m \\
 & \bigoplus \mathfrak{g}_{-1} \otimes \sum_{m_{i_1} + \cdots + m_{i_k} \cong 1 \pmod{3}} \mathbb{C} t^m \bigoplus (\Omega_A/d_A)^0 \bigoplus D.
 \end{aligned}$$

Let Δ_0 be the roots of \mathfrak{g}_0 and Δ_0^s be the short roots of \mathfrak{g}_0 . Define

$$\begin{aligned}
 \Delta_{\text{re}} = & \{\alpha + \delta_{\underline{m}} \mid \alpha \in \Delta_0, m_{i_1} + \cdots + m_{i_k} \cong 0 \pmod{3}\} \\
 & \bigcup \{\alpha + \delta_{\underline{m}} \mid \alpha \in \Delta_0^s, m_{i_1} + \cdots + m_{i_k} \cong 1 \pmod{3}\},
 \end{aligned}$$

and let $\Delta_{\text{im}} = \{\delta_{\underline{m}} \mid \underline{m} \in \mathbb{Z}^n\}$. Let $\Delta = \Delta_{\text{re}} \cup \Delta_{\text{im}}$.

Define

$$\Delta_{\text{re}}^+ = \{\alpha + \delta_{\underline{m}} \mid \alpha > 0, m_n \geq 0 \text{ or } \alpha < 0, m_n \geq 1\}, \quad \Delta_{\text{im}}^+ = \{\delta_{\underline{m}} \mid m_n \geq 1\},$$

and $\Delta^0 = \{\delta_{\underline{m}} \mid m_n = 0\}$. Then $\Delta^+ = \Delta_{\text{re}}^+ \cup \Delta_{\text{im}}^+$ and $\Delta^- = -\Delta^+$. Further $\Delta = \Delta^+ \cup \Delta^0 \cup \Delta^-$. Let $\alpha \in \Delta$. Then $\alpha \in \Delta_{\text{re}}$ if and only if $\langle \alpha, \alpha \rangle \neq 0$.

Let $n \geq 2$ be a positive integer. For each i in $1 \leq i \leq (n-1)$, let N_i be a positive integer. Let $\underline{a}_i = (a_{i1}, a_{i2}, \dots, a_{iN_i})$ be nonzero distinct complex numbers. Let $I = (i_1, i_2, \dots, i_{n-1})$, where $1 \leq i_j \leq N_{i_j}$. Let $\underline{m}' = (m_1, m_2, \dots, m_{n-1}) \in \mathbb{Z}^{n-1}$. Let $\mathfrak{g}_{\text{aff}} = \mathfrak{g} \otimes \mathbb{C}[t_n, t_n^{-1}] \oplus \mathbb{C}K_n$. Let ϕ be the Lie algebra homomorphism

$$\begin{aligned}
 \phi : \mathfrak{g}_{\text{aff}} \otimes \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}] & \rightarrow \left(\bigoplus_{N=N_1 N_2 \cdots N_{n-1} \text{-copies}} \mathfrak{g}_{\text{aff}} \right) = \mathfrak{g}_{\text{aff}N}, \\
 X t_1^{m_1} \cdots t_{n-1}^{m_{n-1}} & \mapsto (a_{1i_1}^{m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} X)_{1 \leq i_1 \leq N_1; \dots; 1 \leq i_{n-1} \leq N_{n-1}},
 \end{aligned}$$

where $X \in \mathfrak{g}_{\text{aff}}$.

Following the proof of [Rao 2004, Lemma 3.11(a)], we see that ϕ is surjective. Let $T \subset \{11, 12, \dots, 1N_1\}$ be a maximal set such that $a_{1i}^3 \neq a_{1j}^3$ whenever $1i, 1j \in T$ and $1 \leq i, j \leq N_1$.

Proposition 6.1. *Let $\phi(\mu) = \phi|_{\tau_0(\mu)}$. If the cardinality of T is N_1 , then $\phi(\mu)$ is a surjective Lie algebra homomorphism from $\tau_0(\mu)$ to $\mathfrak{g}_{\text{aff}N}$.*

Proof. We know that if $m = 3$, then $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_{-1}$. Now there are ten cases.

Case 6.1a. We fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_0$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{3m_n} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{3m_n}, Y_2 t_n^{3m_n}, \dots, Y_N t_n^{3m_n}),$$

where the b_i are complex numbers and $X_i t_n^{3m_n} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$ and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n-1)$. So, we must solve the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{3m_n} a_{1i_1}^{3m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{3m_n}, Y_2 t_n^{3m_n}, \dots, Y_N t_n^{3m_n}).$$

Writing this in matrix form, we must prove that the matrix

$$M = (a_{1i_1}^{3m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$ where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n-1)$, determines the columns. The matrix M is a tensor product of other matrices:

$$M = (a_{1i_1}^{3m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1 - 1} \otimes M_2 \otimes \cdots \otimes M_{n-1},$$

where the M_j for $2 \leq j \leq (n-1)$ defined as in Equation (1). Let

$$M_1 = (a_{1i_1}^{3m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1 - 1}.$$

If the cardinality of T is N_1 , which means the $a_{1i_1}^3$ are all distinct for all $1 \leq i_1 \leq N_1$, then M_1 is a Vandermonde matrix and is therefore invertible. Similarly M_2 is a Vandermonde matrix because $a_{2i} \neq a_{2j}$ for all $1 \leq i, j \leq N_2$ with $i \neq j$. Similarly M_j is also invertible for $3 \leq j \leq (n-1)$. According to a standard result of linear algebra, the determinant of a tensor product of invertible matrices is nonzero. So matrix M is invertible.

Case 6.1b. We fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_0$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{3m_n+1} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{3m_n+1}, Y_2 t_n^{3m_n+1}, \dots, Y_N t_n^{3m_n+1}),$$

where the b_i are complex numbers, $X_i t_n^{3m_n+1} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$, and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n-1)$. So, we must solve the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{3m_n+1} a_{1i_1}^{3m_1+2} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{3m_n+1}, Y_2 t_n^{3m_n+1}, \dots, Y_N t_n^{3m_n+1}).$$

Writing this in matrix form, we must prove that the matrix

$$M = (a_{1i_1}^{3m_1+2} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n-1)$, determines the columns. The matrix M is a tensor product of other matrices

$$M = (a_{1i_1}^{3m_1+2})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1} \otimes M_2 \otimes \cdots \otimes M_{n-1}.$$

Let $M_1 = (a_{1i_1}^{3m_1+2})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. Again, as above if the cardinality of T is N_1 , then M_1 is product of two matrices: one is the diagonal matrix whose diagonal entries are $a_{11}^2, a_{12}^2, \dots, a_{1N_1}^2$, and the other is $(a_{1i_1}^{3m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. Since $a_{11} \neq a_{12} \neq \cdots \neq a_{1N_1}$, we have $a_{11}^2 \neq a_{12}^2 \neq \cdots \neq a_{1N_1}^2$. Hence the diagonal matrix is invertible. The other matrix is a Vandermonde matrix and hence is invertible. So M_1 is invertible. Similarly the M_j for $2 \leq j \leq (n-1)$ are invertible since they are Vandermonde matrices. Hence M is invertible.

Case 6.1c. We fix $m_n \in \mathbb{Z}$, and let $Y_1, Y_2, \dots, Y_N \in \mathfrak{g}_0$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i X_i t_n^{3m_n+2} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \right) = (Y_1 t_n^{3m_n+2}, Y_2 t_n^{3m_n+2}, \dots, Y_N t_n^{3m_n+2}),$$

where the b_i are complex numbers, $X_i t_n^{3m_n+2} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$, and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n-1)$. So, solving the equation

$$\left(\sum_{i=1}^N b_i X_i t_n^{3m_n+2} a_{1i_1}^{3m_1+1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (Y_1 t_n^{3m_n+2}, Y_2 t_n^{3m_n+2}, \dots, Y_N t_n^{3m_n+2}),$$

we must prove that the matrix

$$M = (a_{1i_1}^{3m_1+1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1} \otimes M_2 \otimes \cdots \otimes M_{n-1},$$

is invertible. Let $M_1 = (a_{1i_1}^{3m_1+1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. Again, if the cardinality of T is N_1 , then M_1 is product of two matrices: one is the diagonal matrix whose diagonal entries are $a_{11}, a_{12}, \dots, a_{1N_1}$, and the other is $(a_{1i_1}^{3m_1})_{1 \leq i_1 \leq N_1, 0 \leq m_1 \leq N_1-1}$. Since $a_{11} \neq a_{12} \neq \cdots \neq a_{1N_1}$, the diagonal matrix is invertible. And the other matrix is a Vandermonde matrix and hence invertible. So M_1 is invertible. Similarly the

M_j for $2 \leq j \leq (n-1)$ are invertible as they are Vandermonde matrices. Hence M is invertible.

Case 6.1d. If we consider $(X_i t_n^{3m_n} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}}) \in \tau_0(\mu)$, where $X_i \in \mathfrak{g}_1$, then the proof of this case is similar to that of Case 6.1b.

Case 6.1e. If we consider $(X_i t_n^{3m_n+1} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}}) \in \tau_0(\mu)$, where $X_i \in \mathfrak{g}_1$, then the proof of this case is similar to that of Case 6.1c.

Case 6.1f. If we consider $(X_i t_n^{3m_n+2} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}}) \in \tau_0(\mu)$, where $X_i \in \mathfrak{g}_1$, then the proof of this case is similar to that of Case 6.1a.

Case 6.1g. If we consider $(X_i t_n^{3m_n} t_1^{3m_1+1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}}) \in \tau_0(\mu)$, where $X_i \in \mathfrak{g}_{-1}$, then the proof of this case is similar to that of Case 6.1c.

Case 6.1h. If we consider $(X_i t_n^{3m_n+1} t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}}) \in \tau_0(\mu)$, where $X_i \in \mathfrak{g}_{-1}$, then the proof of this case is similar to that of Case 6.1a.

Case 6.1i. If we consider $(X_i t_n^{3m_n+2} t_1^{3m_1+2} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}}) \in \tau_0(\mu)$, where $X_i \in \mathfrak{g}_{-1}$, then the proof of this case is similar to that of Case 6.1b.

Case 6.1j. Let $c_1, c_2, \dots, c_N \in \mathbb{C}$. Let

$$\phi(\mu) \left(\sum_{i=1}^N b_i K_n t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \right) = (c_1 K_n, c_2 K_n, \dots, c_N K_n),$$

where the b_i are complex numbers, $K_n t_1^{3m_1} t_2^{m_2} \cdots t_{n-1}^{m_{n-1}} \in \tau_0(\mu)$, and $0 \leq m_j \leq N_j - 1$ for $1 \leq j \leq (n-1)$. So, solving the equation

$$\left(\sum_{i=1}^N b_i K_n a_{1i_1}^{3m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}} \right) = (c_1 K_n, c_2 K_n, \dots, c_N K_n),$$

we must prove that the matrix

$$M = (a_{1i_1}^{3m_1} a_{2i_2}^{m_2} \cdots a_{(n-1)i_{n-1}}^{m_{n-1}})$$

is invertible, where $(i_1, i_2, \dots, i_{n-1})$ for $1 \leq i_j \leq N_j$ determines the rows and $\underline{m}' = (m_1, m_2, \dots, m_{n-1})$, where $0 \leq m_j \leq N_j - 1$ for all $1 \leq j \leq (n-1)$, determines the columns. Again by Case 6.1a, M is invertible if the cardinality of T is N_1 .

So if the cardinality of T is N_1 , then $\phi(\mu)$ is a surjective Lie algebra homomorphism from $\tau_0(\mu)$ to $\mathfrak{g}_{\text{aff}N}$. \square

We now define polynomials $P_1(t_1^3)$ and $P_2(t_2), \dots, P_{n-1}(t_{n-1})$ by

$$P_1(t_1^3) = \prod_{i=1}^{N_1} (t_1^3 - a_{1i}^3) \quad \text{and} \quad P_j(t_j) = \prod_{l=1}^{N_j} (t_j - a_{jl}) \quad \text{for } 2 \leq j \leq (n-1).$$

Notation $\langle P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle$ denotes the ideal generated by the polynomials $P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1})$ inside $\mathbb{C}[t_1^{\pm 3}, t_2^{\pm 1}, \dots, t_{n-1}^{\pm 1}]$.

Lemma 6.2. *If the cardinality of T is N_1 , then $\ker \phi(\mu)$ is given by $I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 + I_8 + I_9 + I_{10}$, where for some fixed $m \in \mathbb{Z}$,*

$$\begin{aligned} I_1 &= \mathfrak{g}_0 \otimes t_n^{3m} \quad \langle P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_2 &= \mathfrak{g}_0 \otimes t_n^{3m+1} \quad \langle t_1^2 P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_3 &= \mathfrak{g}_0 \otimes t_n^{3m+2} \quad \langle t_1 P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_4 &= \mathfrak{g}_1 \otimes t_n^{3m} \quad \langle t_1^2 P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_5 &= \mathfrak{g}_1 \otimes t_n^{3m+1} \quad \langle t_1 P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_6 &= \mathfrak{g}_1 \otimes t_n^{3m+2} \quad \langle P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_7 &= \mathfrak{g}_{-1} \otimes t_n^{3m} \quad \langle t_1 P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_8 &= \mathfrak{g}_{-1} \otimes t_n^{3m+1} \quad \langle P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_9 &= \mathfrak{g}_{-1} \otimes t_n^{3m+2} \quad \langle t_1^2 P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle, \\ I_{10} &= K_n \otimes \quad \langle P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1}) \rangle. \end{aligned}$$

Proof. We consider the map ϕ_1 defined by

$$\begin{aligned} \mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} &\rightarrow \bigoplus_{N=N_1 N_2 \dots N_{n-1}} \mathfrak{g}_0 \otimes t_n^{3m}, \\ X t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} &\mapsto (X t_n^{3m} a_{1i_1}^{3m_1} a_{2i_2}^{m_2} \dots a_{(n-1)i_{n-1}}^{m_{n-1}})_{1 \leq i_1 \leq N_1; \dots; 1 \leq i_{n-1} \leq N_{n-1}}. \end{aligned}$$

Let $M = (N = N_1 N_2 \dots N_{n-1})$ (the dimension of \mathfrak{g}_0). As in Proposition 6.1, one can show that ϕ_1 is surjective. Since ϕ_1 is nothing but the evaluation map at the roots of $P_1(t_1^3), P_2(t_2), \dots, P_{n-1}(t_{n-1})$, we have $I_1 \subset \ker \phi_1$. So

$$\dim(\mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} / I_1) \geq M.$$

Consider the space

$$R = \{ \mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} \mid 0 \leq m_i \leq N_i - 1 \text{ for } 1 \leq i \leq (n-1) \}.$$

Any element of $\mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}}$ can be written as linear combination of elements of R modulo I_1 . Hence ϕ_1 is injective on R modulo I_1 , since the corresponding matrix is invertible. So R is a spanning set for

$$(\mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} / I_1),$$

and $\dim(\mathfrak{g}_0 \otimes t_n^{3m} t_1^{3m_1} t_2^{m_2} \dots t_{n-1}^{m_{n-1}} / I_1) \leq M$. So I_1 is the kernel of ϕ_1 . Similarly arguing on the other components of $\tau_0(\mu)$, we get that $\sum_{j=1}^{10} I_j$ is the kernel of $\phi(\mu)$ if the cardinality of T is N_1 . This proves Lemma 6.2. \square

Remark 6.3. The results of Section 5 can be proved similarly in case $m = 3$.

Note 6.4. The proof of Proposition 2.6 will remain same in case $m = 3$ except that for $\lambda, \lambda + \delta_1, \lambda + 2\delta_1$ or $\lambda, \lambda + \delta_1, \lambda - \delta_1$ or $\lambda, \lambda - \delta_1, \lambda - 2\delta_1$ will have the properties of the first paragraph of the proof. We can assume $\lambda + \eta, \lambda + \delta_1 + \eta, \lambda + 2\delta_1 + \eta \notin P(V)$ for all $\eta > 0$.

Note 6.5. Let V be an integrable irreducible module for the twisted toroidal Lie algebra $\tau(\mu)$ with finite-dimensional weight spaces. Let k be as in Theorem 2.3. Suppose $k \geq 1$ and $K_i = 0$ for all i . Then such a module V does not exist. This result can be proved in a way similar to the proof of [Rao 2004, Proposition 4.13].

Acknowledgments

We thank the referee for helpful comments and suggestions.

References

- [Batra 2004] P. Batra, “Representations of twisted multi-loop Lie algebras”, *J. Algebra* **272**:1 (2004), 404–416. MR 2004k:17011 Zbl 1077.17005
- [Berman and Billig 1999] S. Berman and Y. Billig, “Irreducible representations for toroidal Lie algebras”, *J. Algebra* **221**:1 (1999), 188–231. MR 2000k:17004 Zbl 0942.17016
- [Berman and Krylyuk 1995] S. Berman and Y. Krylyuk, “Universal central extensions of twisted and untwisted Lie algebras extended over commutative rings”, *J. Algebra* **173**:2 (1995), 302–347. MR 96e:17066 Zbl 0852.17020
- [Kac 1990] V. G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, 1990. MR 92k:17038 Zbl 0716.17022
- [Morita and Yoshii 1985] J. Morita and Y. Yoshii, “Universal central extensions of Chevalley algebras over Laurent polynomial rings and GIM Lie algebras”, *Proc. Japan Acad. Ser. A Math. Sci.* **61**:6 (1985), 179–181. MR 87c:17028 Zbl 0581.17010
- [Rao 1993] S. E. Rao, “On representations of loop algebras”, *Comm. Algebra* **21**:6 (1993), 2131–2153. MR 95c:17039 Zbl 0777.17019
- [Rao 1995] S. E. Rao, “Iterated loop modules and a filtration for vertex representation of toroidal Lie algebras”, *Pacific J. Math.* **171**:2 (1995), 511–528. MR 97c:17034 Zbl 0867.17017
- [Rao 2001] S. E. Rao, “Classification of irreducible integrable modules for multi-loop algebras with finite-dimensional weight spaces”, *J. Algebra* **246**:1 (2001), 215–225. MR 2003c:17010 Zbl 0994.17002
- [Rao 2004] S. E. Rao, “Classification of irreducible integrable modules for toroidal Lie algebras with finite dimensional weight spaces”, *J. Algebra* **277**:1 (2004), 318–348. MR 2005d:17011 Zbl 02102481
- [Rao 2005] S. E. Rao, “Irreducible representations for toroidal Lie-algebras”, *J. Pure Appl. Algebra* **202**:1-3 (2005), 102–117. MR 2006e:17011 Zbl 02217995
- [Rao and Moody 1994] S. E. Rao and R. V. Moody, “Vertex representations for n -toroidal Lie algebras and a generalization of the Virasoro algebra”, *Comm. Math. Phys.* **159**:2 (1994), 239–264. MR 94m:17028 Zbl 0808.17018

Received November 11, 2007. Revised May 6, 2008.

S. ESWARA RAO
SCHOOL OF MATHEMATICS
TATA INSTITUTE OF FUNDAMENTAL RESEARCH
HOMI BHABHA ROAD
MUMBAI 400 005
INDIA
senapati@math.tifr.res.in

PUNITA BATRA
HARISH CHANDRA RESEARCH INSTITUTE
CHHATNAG ROAD
JHUNSI
ALLAHABAD 211 019
INDIA
batra@mri.ernet.in

