

*Pacific
Journal of
Mathematics*

THICK TRIANGULATIONS OF HYPERBOLIC n -MANIFOLDS

WILLIAM BRESLIN

Volume 241 No. 2

June 2009

THICK TRIANGULATIONS OF HYPERBOLIC n -MANIFOLDS

WILLIAM BRESLIN

We show that a complete hyperbolic n -manifold has a geodesic triangulation such that the tetrahedra contained in the thick part are L -bilipschitz diffeomorphic to the standard Euclidean n -simplex, for some constant L depending only on the dimension and the constant used to define the thick-thin decomposition of M .

A geodesic triangulation of a complete hyperbolic n -manifold M may be forced by the geometry of M to have simplices which are either small or flat. Big simplices without small dihedral angles cannot live in the thin part of M . We show that a complete hyperbolic n -manifold M has a geodesic triangulation such that the simplices contained in the thick part of M are L -bilipschitz diffeomorphic to the standard Euclidean n -simplex, for some constant L depending only on the dimension n and the constant μ used to define the thick-thin decomposition of M . We call such a triangulation a (μ, L) -thick geodesic triangulation of M .

Theorem. *Let $n \geq 2$. Let μ be a Margulis constant for \mathbb{H}^n . There exists a constant $L := L(n, \mu)$ such that every complete hyperbolic n -manifold has a (μ, L) -thick geodesic triangulation.*

Existence of thick geodesic triangulations implies that any hyperbolic n -manifold M has a geodesic triangulation such that the simplices contained in the thick part of M come from a fixed compact set of simplices which does not depend on the manifold. In [Breslin 2006], we used this compactness to prove existence of bounds on the curvatures of surfaces embedded in hyperbolic 3-manifolds. In particular, it is shown that there exists a fixed constant $\omega > 0$ such that if S is an incompressible surface or a strongly irreducible Heegaard surface in a complete orientable hyperbolic 3-manifold, then S is isotopic to a surface whose principal curvatures are bounded in absolute value by ω . The constant ω depends on neither the hyperbolic 3-manifold nor the surface. Another interesting application of thick triangulations can be found in [Kapovich 2007], which explains how to use thick triangulations of hyperbolic n -manifolds to obtain an inequality between the

MSC2000: primary 30F40, 53A35; secondary 57M50.

Keywords: hyperbolic manifold, triangulation, thick.

This work was partially supported by the NSF grant DMS-0135345.

relative homological dimension of a Kleinian group $\Gamma \subset \text{Isom } \mathbb{H}^n$ and its critical exponent.

To prove existence of thick geodesic triangulations, we examine Delaunay triangulations of well-spaced point sets in hyperbolic n -space and the problem of eliminating flat simplices (i.e. simplices with small dihedral angles). The corresponding question for Euclidean space has been well-studied. The only tetrahedra in such a triangulation which can have small dihedral angles are called *slivers*, and it was a problem to show how to remove them without creating new ones. Several techniques for removing slivers have been developed in the Euclidean setting; see [Edelsbrunner et al. 2000; Miller et al. 1996; Li 2000; 2003]. We adapt the technique of perturbing vertices of a Delaunay triangulation, introduced in the first of these references, in order to remove slivers to the hyperbolic setting.

Emil Saucan [2006a; 2006b; 2005] has shown that hyperbolic n -orbifolds have triangulations whose simplices are uniformly round, called *fat triangulations*, and he uses this to prove existence of quasimeromorphic maps which are automorphic with respect to the corresponding Kleinian group.

Let M be a complete hyperbolic n -manifold.

Definition (thick-thin decomposition). Let $\mu > 0$. The μ -*thick part* of M , denoted by $M_{[\mu, \infty)}$, is the set of points where the injectivity radius is at least $\mu/2$. The μ -*thin part* of M , denoted by $M_{(0, \mu]}$, is the closure of the complement of $M_{[\mu, \infty)}$.

Definition (thick triangulation). Let $\mu > 0$, $L > 0$. A triangulation T of a complete hyperbolic n -manifold M is (μ, L) -*thick* if every n -simplex of T which is contained in the μ -thick part of M is L -bilipschitz diffeomorphic to the standard Euclidean n -simplex. Once we have fixed μ and L , we will refer a *thick* triangulation.

Definition (Delaunay triangulation). Let \mathcal{S} be a generic set of points in M such that for any $p \in M$ the ball $B(p, \text{inj}(M, p)/5)$ centered at p with radius $\text{inj}(M, p)/5$ contains a point of \mathcal{S} in its interior. The *Delaunay triangulation* of \mathcal{S} is the geodesic triangulation of M determined as follows: A set, $\{p_0, \dots, p_n\}$, of $n + 1$ vertices in \mathcal{S} determines an n -simplex in \mathcal{T} if and only if the minimal radius circumscribing sphere contains no points of \mathcal{S} in its interior.

See [Leibon and Letscher 2000] for existence of Delaunay triangulations in Riemannian manifolds.

We want to find triangulations such that the simplices in the thick part of M are neither too big nor too small, and which do not have small dihedral angles. It is not difficult to find triangulations such that the simplices in the thick part are neither too big nor too small. Let $\mu > 0$ be a Margulis constant for \mathbb{H}^n . Let $\varepsilon := \mu/100$. Let \mathcal{S} be a generic set of points in M such that for any $p \in M$ the ball

$B(p, \text{inj}(M, p)/5)$ centered at p with radius $\text{inj}(M, p)/5$ contains a point of \mathcal{S} in its interior. Also assume that the set \mathcal{S} is maximal with respect to the condition that each point in $\mathcal{S} \cap M_{[\mu, \infty)}$ is no closer than ε to another point of \mathcal{S} . Let T be the Delaunay triangulation of \mathcal{S} . Any simplex of T in the μ -thick part of M has edge lengths in the interval $[\varepsilon, 2\varepsilon]$. In fact, this triangulation is not very far from the one we want. We will show that each vertex of $\mathcal{S} \cap M_{[\mu, \infty)}$ can be moved a small distance so that the Delaunay triangulation of the new set of points is (μ, L) -thick.

Definition (altitude). The *altitude* of a vertex v of a geodesic n -simplex in \mathbb{H}^n is the distance from v to the hyperplane of \mathbb{H}^n containing the other vertices.

Definition (good simplices). For $2 \leq k \leq n$, $0 < a < b$, and $0 < d$, a geodesic k -dimensional simplex S in hyperbolic n -space \mathbb{H}^n is (a, b, d) -good if the lengths of the edges of S are contained in the interval $[a, b]$ and the altitude of each vertex of S is at least d . When a, b, d are understood, we will refer to *good simplices*. We say S is *bad* if it is not good.

Remark 1. If a geodesic n -simplex S in \mathbb{H}^n has edge lengths in $[a, b]$, then there are two ways that it can be (a, b, d) -bad for a small number $d > 0$. Either S has big circumradius or the vertices of S all lie near a hyperbolic $(n - 2)$ -sphere. In the triangulation T described above the simplices in $T \cap M_{[\mu, \infty)}$ have bounded circumradii, so that the vertices of any (a, b, d) -bad simplex in $T \cap M_{[\mu, \infty)}$ must all lie close to a hyperbolic $(n - 2)$ -sphere. The vertices get closer to an $(n - 2)$ -sphere as $d \rightarrow 0$.

Remark 2. Let $b > a > 0$ and $d > 0$. Consider the set of compact hyperbolic n -simplices in \mathbb{H}^n up to isometry. The set of geodesic (a, b, d) -good simplices is a compact subset. Hence:

Lemma 1. For each $n \geq 2$, $b > a > 0$, and $d > 0$, there exists a constant $L := L(n, a, b, d)$ such that each (a, b, d) -good simplex is L -bilipschitz diffeomorphic to the standard Euclidean n -simplex.

Definition (good perturbation). Let $\delta > 0$. A δ -good perturbation of \mathcal{S} is a collection of points \mathcal{S}' in M such that there exists a bijection $\phi : \mathcal{S} \rightarrow \mathcal{S}'$ with $d(p, \phi(p)) \leq \delta$ for every $p \in \mathcal{S}$. Denote $\phi(p)$ by p' . If \mathcal{T} and \mathcal{T}' are the Delaunay triangulations of \mathcal{S} and \mathcal{S}' , then we will say that \mathcal{T}' is a δ -good perturbation of \mathcal{T} . When δ is understood, we will refer to a *good perturbation*.

Definition (bad region). Let $S = [v_1, \dots, v_k]$ be a geodesic $(k - 1)$ -simplex in \mathbb{H}^n . Let $b > a > 0$, $c > 0$, $d > 0$. The (a, b, c, d) -bad region of S is the set of points p in \mathbb{H}^3 such that $[p, v_1, \dots, v_k]$ has edge lengths in $[a, b]$, circumradius at most c , and the distance from p to the hyperplane containing the opposite face is less than d .

The next lemma shows if a bad simplex has good proper subsimplices, then each vertex is close to the plane containing the other vertices.

Lemma 2. *Let $S = [v_0, \dots, v_k]$ be a geodesic k -simplex in \mathbb{H}^k with edge lengths in $[a, b]$. If every proper subcomplex of S is (a, b, d_0) -good and S is (a, b, d) -bad, then the distance from each vertex of S to the hyperplane containing the opposite face is at most a constant $D := D(b, d_0, d)$ such that $D(b, d_0, d) \rightarrow 0$ as $d \rightarrow 0$ and b and d_0 remain fixed.*

Proof. Since S is (a, b, d) -bad and the edge lengths are in $[a, b]$, the distance from some vertex, say v_0 , to the hyperplane in \mathbb{H}^k containing the opposite face $[v_1, \dots, v_k]$ is less than d . Let P_0 be the hyperplane containing v_0, \dots, v_{k-1} . Let P_k be the hyperplane containing v_1, \dots, v_k . Let α be the angle between P_0 and P_k . Let v'_0 be the orthogonal projection of v_0 to $P_0 \cap P_k$ and let v''_0 be the orthogonal projection of v_0 to P_k . The angle of the hyperbolic triangle $[v_0, v'_0, v''_0]$ at v'_0 is α . See Figure 1(a). The hyperbolic law of sines [Fenchel 1989] gives us

$$\sin \alpha = \frac{\sinh \|[v_0, v''_0]\|}{\sinh \|[v_0, v'_0]\|}.$$

Let v'_k be the orthogonal projection of v_k to $P_0 \cap P_k$ and let v''_k be the orthogonal projection of v_k to P_0 . The angle the hyperbolic triangle $[v_k, v'_k, v''_k]$ at v''_k is also α . See Figure 1(b). Using the hyperbolic law of sines again we get

$$\sinh \|[v_k, v''_k]\| = \sin \alpha \cdot \sinh \|[v_k, v'_k]\| = \frac{\sinh \|[v_0, v''_0]\|}{\sinh \|[v_0, v'_0]\|} \cdot \sinh \|[v_k, v'_k]\|.$$




Figure 1. If every proper subcomplex of S is good and S is bad, then each vertex is close to the plane containing the opposite face.




Figure 2. The distance to the circumsphere A of $[v_1, \dots, v_k]$ from v_0 is small if every proper subcomplex of S is good, but S is bad.

Now $\|[v_0, v_0']\| \geq d_0$ and $\|[v_k, v_k']\| \leq b$ since $\|[v_0, \dots, v_{k-1}]\|$ and $\|[v_1, \dots, v_k]\|$ are (a, b, d_0) -good $(k-1)$ -simplices. Also, $\|[v_0, v_0']\| < d$ by our assumption. Thus

$$\sinh \|[v_k, v_k']\| \leq \frac{\sinh d}{\sinh d_0} \cdot \sinh b.$$

We have shown that the distance to the hyperplane containing $[v_0, \dots, v_{k-1}]$ from v_k is at most

$$D(b, d_0, d) := \operatorname{arcsinh} \left(\frac{\sinh d}{\sinh d_0} \cdot \sinh b \right).$$

A similar argument shows the distance from each vertex to the hyperplane containing the opposite face is at most

$$D(b, d_0, d). \quad \square$$

Next we show that if a bad k -simplex has bounded circumradius and good proper simplices, then the vertices lie near a hyperbolic $(k-2)$ -sphere.

Lemma 3. *Let $S = [v_0, \dots, v_k]$ be a geodesic k -simplex in \mathbb{H}^k with edge lengths in $[a, b]$ and circumradius at most c . If every proper subcomplex of S is (a, b, d_0) -good and S is (a, b, d) -bad, then the distance from each vertex to the circumsphere of the opposite face is at most a constant $R := R(a, b, c, d_0, d)$ such that $R \rightarrow 0$ as $d \rightarrow 0$ and a, b, c, d_0 remain fixed.*

Proof. Since every proper subcomplex of S is (a, b, d_0) -good and S is (a, b, d) -bad and has edge lengths in $[a, b]$ and circumradius at most c , the distance from

each vertex to the hyperplane containing the opposite face is at most the constant $D := D(b, d_0, d)$ provided by Lemma 2.

Let A be the circumsphere of $[v_1, \dots, v_k]$. Let B be the circumsphere of S . Now A is the intersection of B with some hyperplane P . Let v'_0 be the orthogonal projection of v_0 to P . Let v''_0 be the point on A which is closest to v_0 . Let Q be the 2-dimensional hyperbolic plane which contains v_0, v'_0 , and v''_0 . Now $B \cap Q$ is a hyperbolic circle which intersects the hyperbolic line $P \cap Q$ (see Figure 2). Since the radii of A and B are in the interval $[a/2, c]$, the angle between $P \cap Q$ and the tangent of $B \cap Q$ at the two points in $(P \cap Q) \cap (B \cap Q)$ is bounded from below by a positive constant $\alpha_0 := \alpha_0(a, c)$. Thus the angle α between $[v'_0, v_0]$ and $[v''_0, v'_0]$ is bounded from below by α_0 . Let β be the angle between $[v_0, v'_0]$ and $[v_0, v''_0]$. By the hyperbolic law of sines we have

$$\sinh \|[v'_0, v''_0]\| = \frac{\sinh \|[v_0, v'_0]\|}{\sin \alpha} \cdot \sin \beta \leq \frac{\sinh D}{\sin \alpha_0}.$$

The triangle inequality now gives us

$$\begin{aligned} \|[v_0, v''_0]\| &\leq \|[v_0, v'_0]\| + \|[v'_0, v''_0]\| \\ &\leq D(b, d_0, d) + \operatorname{arcsinh} \frac{\sinh D(b, d_0, d)}{\sin \alpha_0(a, c)}. \end{aligned}$$

Now take

$$R(a, b, c, d_0, d) := D(b, d_0, d) + \operatorname{arcsinh} \frac{\sinh D(b, d_0, d)}{\sin \alpha_0(a, c)}. \quad \square$$

We can now bound the volume of the bad region of a simplex with bounded circumradius and good proper subsimplices.

Lemma 4. *Let $b > a > 0$, $c > 0$, $d_0 > 0$, $d > 0$. Let $n \geq 3$ and $k < n$. Let S be a geodesic k -simplex in \mathbb{H}^n such that the circumradius of S is at most c and every proper subsimplex of S is (a, b, d_0) -good. The volume of the (a, b, c, d) -bad region of S is at most a constant $V_k := V_k(n, a, b, c, d_0, d)$ such that $V_k \rightarrow 0$ as $d \rightarrow 0$ and a, b, c, d_0 remain fixed.*

Proof. The (a, b, c, d) -bad region of S is contained in the R -neighborhood of the circumsphere B of S , where $R := R(a, b, c, d_0, d)$ is the constant provided by Lemma 3. Since the radius of S is at most c and $R(a, b, c, d_0, d) \rightarrow 0$ as $d \rightarrow 0$, we can let V_k be the volume of the R -neighborhood of a $(k - 1)$ -dimensional hyperbolic sphere of radius c in \mathbb{H}^n . \square

Let $\delta = \varepsilon/10$. Given a point p in the set $\mathcal{S} \cap M_{[\mu, \infty)}$, the following Lemma bounds the number of k -tuples of points which might form a k -simplex with p' in a δ -good perturbation T' of the triangulation T .




Figure 3. Left: if the radius r_0 is fixed, then moving q and r closer to p makes the altitude from p smaller. Right: both $\|[p, x]\|$ and h are bounded from below in terms of a, b , and R .

Lemma 5. *Let $p \in \mathcal{S} \cap M_{[\mu, \infty)}$. For each $k = 3, \dots, n$, the number of k -tuples $\{v_1, \dots, v_k\}$ such that $[p', v'_1, \dots, v'_k]$ is a k -simplex in some δ -good perturbation of \mathcal{T} is bounded by a constant $N := N(n, k, \mu)$.*

Proof. The $(\frac{\epsilon}{2} - \delta)$ -balls centered at the points of $\mathcal{S} \cap M_{[\mu, \infty)}$ are mutually disjoint since no two points of $\mathcal{S} \cap M_{[\mu, \infty)}$ are closer than $\epsilon - 2\delta$ to each other. If p' and q' are vertices of a k -simplex in a δ -good perturbation \mathcal{T}' of \mathcal{T} , then $d(p', q') \leq 2\epsilon + 2\delta$, so that the $(\frac{\epsilon}{2} - \delta)$ -ball centered at q' is contained in the $(2\epsilon + 2\delta)$ -ball centered at p' . There can be at most

$$m := m(n, \mu) = \left\lfloor \frac{\text{Vol}_{\mathbb{H}^n} B(2\epsilon + 2\delta)}{\text{Vol}_{\mathbb{H}^n} B(\frac{\epsilon}{2} - \delta)} \right\rfloor$$

mutually disjoint $(\frac{\epsilon}{2} - \delta)$ -balls contained in a $(2\epsilon + 2\delta)$ -ball, where $\lfloor w \rfloor$ is the integer part of w . One of these is centered at p' . So there are at most $m - 1$ vertices in \mathcal{S} which may be the vertex of a k -simplex in \mathcal{T}' which also has p' as a vertex. Thus the number of k -tuples $\{v_1, \dots, v_k\}$ of points in \mathcal{S} such that $[p', v'_1, \dots, v'_k]$ is a k -simplex in some good perturbation of \mathcal{T} is at most $\binom{m}{k}$. Let $N(n, k, \mu) := \binom{m(n, \mu)}{k}$. □

Lemma 6. *A geodesic triangle in \mathbb{H}^2 with edge lengths in $[a, b]$ and circumradius at most R has altitudes bounded from below by a positive constant $h_0 := h_0(a, b, R)$.*

Proof. Since the sum of the angles of t are less than π there are at least two angles of t which are less than $\pi/2$. Let p be the vertex opposite these angles. The orthogonal projection of p onto the line containing the opposite edge $[q, r]$ is contained in the interior of $[q, r]$. Now suppose we have fixed the circumradius $r_0 \in [a/2, R]$ of t and consider all triangles with edges of length at least a such that p projects to the interior of $[q, r]$. The triangle with the shortest altitude at p is an isosceles triangle which lies on a hyperbolic circle of radius r_0 (see Figure 3). Let c be the center of the hyperbolic circle containing p, q, r . Let x be the intersection of $[p, c]$ and $[q, r]$. Let $\alpha = \angle qpx$. Let $\beta = \angle pqx$.

By the hyperbolic law of cosines, we have

$$\cos \alpha = \frac{\cosh \|[p, q]\| \cdot \cosh \|[p, c]\| - \cosh \|[q, c]\|}{\sinh \|[p, q]\| \cdot \sinh \|[p, c]\|} = \frac{\cosh a \cdot \cosh r_0 - \cosh r_0}{\sinh a \cdot \sinh r_0}.$$

By the hyperbolic law of sines we have $\sinh \|[q, x]\| = \sinh \|[p, q]\| \cdot \sin \alpha$. Now the altitude of $[p, q, r]$ from p is $\|[p, x]\|$. Using the law of cosines again, we get

$$\begin{aligned} \cosh \|[p, x]\| &= \cosh a \cosh \|[q, x]\| - \sinh a \sinh \|[q, x]\| \cos \beta \\ &\geq \cosh a \cosh \|[q, x]\| - \sinh a \sinh \|[q, x]\|, \end{aligned}$$

Define $h_1(a, r_0) = \operatorname{arccosh}(\cosh a \cosh \|[q, x]\| - \sinh a \sinh \|[q, x]\|)$. So far we have shown that the altitude from a vertex which projects to the interior of the opposite face is at least $h_1(a, r_0)$ if the circumradius of $[p, q, r]$ is r_0 . Since $h_1(a, r_0)$ decreases as r_0 increases, we have that $h_1(a, R)$ is a lower bound on the altitude from a vertex which projects to the interior of the opposite face for triangles satisfying the hypotheses of the lemma. Let h' be the altitude from r .

We have

$$\sin \beta = \frac{\sinh \|[p, x]\|}{\sinh \|[p, q]\|} \geq \frac{\sinh h_1(a, R)}{\sinh b}.$$

Also,

$$\sinh h = \sinh \|[q, r]\| \cdot \sin \beta \geq \sinh a \cdot \frac{\sinh h_1(a, R)}{\sinh b},$$

so

$$h \geq \operatorname{arcsinh}\left(\frac{\sinh a}{\sinh b} \cdot \sinh h_1(a, R)\right).$$

A similar argument works for the altitude from q . Let

$$h_0(a, b, R) = \operatorname{arcsinh}\left(\frac{\sinh a}{\sinh b} \cdot \sinh h_1(a, R)\right). \quad \square$$

Proof of the Theorem. The idea of the proof is to show that there is a δ -good perturbation T' of the triangulation T such that each tetrahedron of T' contained in the thick part of M is (a, b, d) -good for fixed constants a, b , and d . We know

each k -simplex of T contained in the thick part of M has edge lengths in the interval $[\varepsilon, 2\varepsilon]$ and circumradius at most ε (for $k = 1, \dots, n$). Note that if t is a k -simplex in the Delaunay triangulation of a δ -good perturbation of \mathcal{S} which is contained in $M_{[\mu, \infty)}$, then t has edge lengths between $\varepsilon - 2\delta$ and $2\varepsilon + 2\delta$, and circumradius no more than $\varepsilon + \delta$.

We will remove the bad simplices one dimension at a time. Let $a := \varepsilon - 2\delta$, $b := \varepsilon + 2\delta$, $c := \varepsilon + \delta$. We will proceed by induction on the dimension of the simplices.

Since any 2-simplex in T has edge lengths in $[a, b]$ and circumradius at most c , we conclude from Lemma 6 that each 2-simplex in T is $(a, b, h_0(a, b, c))$ -good.

Assume that T_k is a $\frac{\delta}{100 \cdot 2^k}$ -good perturbation of T such that every simplex of dimension at most k which is contained in the μ -thick part of M is (a, b, d_k) -good for some positive constant $d_k \leq d_2$. Let

$$\delta_{k+1} := \frac{\delta}{100 \cdot 2^{k+1}}.$$

Let $d_{k+1} \leq d_k$ be a positive constant to be determined later.

Take $p_1 \in \mathcal{S} \cap M_{[\mu, \infty)}$. Let \mathcal{U}_1 be the set of simplices $[v_0, \dots, v_l] \in \mathcal{T}_k$, with $l \leq k$, satisfying the following property: There exists a δ_{k+1} -good perturbation \mathcal{T}' of \mathcal{T} obtained by perturbing only the point p_1 and such that $[p'_1, v_0, \dots, v_l] \in \mathcal{T}'$.

By Lemma 4 and Lemma 5, the total volume of the (a, b, c, d_{k+1}) -bad regions of the l -simplices in \mathcal{U}_1 is bounded by $V_l(a, b, c, d_2, d_{k+1}) \cdot N(n, l, \mu)$, so that the total volume of the (a, b, c, d_{k+1}) -bad regions of all simplices in \mathcal{U}_1 is bounded by $\sum_{l=1}^{k+1} V_l(a, b, c, d_2, d_{k+1}) \cdot N(n, l, \mu)$. Let $B(p, \delta_{k+1})$ be the ball of radius δ_{k+1} centered at p . If we choose d_{k+1} so small that $\sum_{l=1}^{k+1} V_l(a, b, c, d_2, d_{k+1}) \cdot N(n, l, \mu) \leq \text{vol}(B(p_1, \delta_{k+1}))$, then the (a, b, c, d_{k+1}) -bad regions of the simplices in \mathcal{U}_1 cannot cover $B(p_1, \delta_{k+1})$. Now choose p'_1 in $B(p_1, \delta_{k+1})$ (so that the perturbation is δ_{k+1} -good) and outside the (a, b, c, d_{k+1}) -bad region of every simplex in \mathcal{U}_1 . Call the new set of points \mathcal{S}_1 and the new triangulation \mathcal{T}_1 .

Assume we have perturbed the points p_1, \dots, p_s to p'_1, \dots, p'_s and now have a set of points \mathcal{S}_s and a triangulation \mathcal{T}_s such that none of p'_1, \dots, p'_s is the vertex of a (a, b, c, d_{k+1}) -bad simplex of dimension less than $k + 2$. Let p_{s+1} be a point in $[\mathcal{S}_s \cap M_{[\mu, \infty)}] - \{p'_1, \dots, p'_s\}$. Let \mathcal{U}_{s+1} be the set of simplices $[v_0, \dots, v_l] \in \mathcal{T}_s$ of dimension at most k such that there exists a δ_{k+1} -good perturbation \mathcal{T}'_s of \mathcal{T}_s which is obtained by perturbing only the point p_{s+1} and such that $[p'_{s+1}, v_0, \dots, v_k] \in \mathcal{T}'_s$. Since d_{k+1} is so small, we can choose a point p'_{s+1} in the ball of radius δ_{k+1} centered at p_{s+1} and outside the (a, b, c, d_{k+1}) -bad region of every simplex in \mathcal{U}_{s+1} .

Assume that M has finite volume. Let \mathcal{T}' be the triangulation we get after perturbing every point of $\mathcal{S} \cap M_{[\mu, \infty)}$ once and only once (There are only finitely

many since M has finite volume). Let $[p', v'_1, \dots, v'_l] \in \mathcal{T}'$ be a simplex of dimension at most $k + 1$. Suppose that p' was the last point perturbed among these $l + 1$ vertices. We chose p' to be outside the (a, b, c, d_{k+1}) -bad region of $[v'_1, \dots, v'_l]$, so that $[p', v'_1, \dots, v'_l]$ is (a, b, d_{k+1}) -good. Thus any $(k + 1)$ -simplex of \mathcal{T}' contained in $M_{[\mu, \infty)}$ is (a, b, d_{k+1}) -good.

If M has infinite volume, then for each positive integer m the above procedure can be used to perturb the vertices contained in an m -ball centered at some fixed point x_0 , giving us a geodesic triangulation $\mathcal{T}(m)$ of M such that any tetrahedron contained in $M_{[\mu, \infty)} \cap B(x_0, m)$ is (a, b, d_{k+1}) -good. Suppose we want to define the final triangulation on a ball $B(x_0, N)$ for some positive integer N . Since the triangulations $\mathcal{T}(m)$ agree on the ball $B(x_0, N)$ for $m \geq 100N$, we can use the triangulation $\mathcal{T}(100N)$ to define the triangulation inside $B(x_0, N)$.

We have shown that M has a geodesic triangulation such that every simplex of dimension at most $k + 1$ which is contained in the thick part of M is (a, b, c, d_{k+1}) -good, completing the induction. When $k = n - 1$, we get a geodesic triangulation of M such that every simplex of dimension at most n which is contained in the thick part of M is (a, b, d_n) -good. Thus the triangulation is L -thick for a constant L depending only on a, b , and d_n , which depend only on μ and n . \square

References

- [Breslin 2006] W. G. Breslin, “Curvature bounds for surfaces in hyperbolic 3-manifolds”, preprint, 2006. arXiv math.GT/0703208
- [Edelsbrunner et al. 2000] H. Edelsbrunner, X.-Y. Li, G. Miller, A. Stathopoulos, D. Talmor, S.-H. Teng, A. Üngör, and N. Walkington, “Smoothing and cleaning up slivers”, pp. 273–277 in *Proceedings of the Thirty-Second Annual ACM Symposium on Theory of Computing* (Portland, OR), ACM, New York, 2000. MR 2114541
- [Fenchel 1989] W. Fenchel, *Elementary geometry in hyperbolic space*, Studies in Mathematics **11**, de Gruyter, Berlin, 1989. MR 91a:51009 Zbl 0674.51001
- [Kapovich 2007] M. Kapovich, “Homological dimension and critical exponent of Kleinian groups”, preprint, 2007. arXiv math/0701797
- [Leibon and Letscher 2000] G. Leibon and D. Letscher, “Delaunay triangulations and Voronoi diagrams for Riemannian manifolds”, pp. 341–349 in *Proceedings of the Sixteenth Annual Symposium on Computational Geometry* (Hong Kong, 2000), ACM, New York, 2000. MR 2001i:53011
- [Li 2000] X.-Y. Li, “Spacing control and sliver-free Delaunay mesh”, pp. 295–306 in *Proceedings, 9th International Meshing Roundtable* (New Orleans, 2000), Sandia National Laboratories, Albuquerque, NM, 2000.
- [Li 2003] X.-Y. Li, “Generating well-shaped d -dimensional Delaunay meshes”, *Theoret. Comput. Sci.* **296**:1 (2003), 145–165. MR 2004b:68168 Zbl 1044.68153
- [Miller et al. 1996] G. L. Miller, D. Talmor, S. H. Teng, N. Walkington, and H. Wang, “Control volume meshes using sphere packing: generation refinement, and coarsening”, pp. 47–61 in *Proceedings, 5th International Meshing Roundtable* (Pittsburgh, PA, 1996), Sandia National Laboratories, Albuquerque, NM, 1996.

- [Saucan 2005] E. Saucan, “Note on a theorem of Munkres”, *Mediterr. J. Math.* **2**:2 (2005), 215–229. MR 2006g:57048 Zbl 05058627
- [Saucan 2006a] E. Saucan, “The existence of quasimeromorphic mappings”, *Ann. Acad. Sci. Fenn. Math.* **31**:1 (2006), 131–142. MR 2006k:30023 Zbl 1100.30020
- [Saucan 2006b] E. Saucan, “The existence of quasimeromorphic mappings in dimension 3”, *Conform. Geom. Dyn.* **10** (2006), 21–40. MR 2006k:30024 Zbl 1118.30016

Received August 27, 2008.

WILLIAM BRESLIN
UNIVERSITY OF MICHIGAN
DEPARTMENT OF MATHEMATICS
530 CHURCH STREET
ANN ARBOR, MI 48109-1043
UNITED STATES

breslin@umich.edu

<http://www.math.lsa.umich.edu/people/facultyDetail.php?unique=breslin>

