

*Pacific
Journal of
Mathematics*

**FOUR-DIMENSIONAL OSSERMAN METRICS
OF NEUTRAL SIGNATURE**

EDUARDO GARCÍA-RÍO, PETER B. GILKEY,
M. ELENA VÁZQUEZ-ABAL AND RAMÓN VÁZQUEZ-LORENZO

FOUR-DIMENSIONAL OSSERMAN METRICS OF NEUTRAL SIGNATURE

EDUARDO GARCÍA-RÍO, PETER B. GILKEY,
M. ELENA VÁZQUEZ-ABAL AND RAMÓN VÁZQUEZ-LORENZO

In the algebraic context, we show null Osserman, spacelike Osserman, and timelike Osserman are equivalent conditions for a model of signature $(2, 2)$. We also classify the null Jordan Osserman models of signature $(2, 2)$. In the geometric context, we show that a pseudo-Riemannian manifold with this signature is null Jordan Osserman if and only if either it has constant sectional curvature or it is locally a complex space form.

1. Introduction

Let $\mathcal{M} := (M, g)$ be a pseudo-Riemannian manifold. We say a tangent vector v is *spacelike*, *timelike*, or *null* if $g(v, v) > 0$, if $g(v, v) < 0$, or if $g(v, v) = 0$, respectively. Geometric properties derived from conditions on spacelike, timelike, and null vectors can have quite different meanings. For instance, the conditions of spacelike, timelike, and null geodesic completeness are nonequivalent and independent. Although spacelike and timelike conditions can sometimes become equivalent (for example, as they concern boundedness conditions on the sectional curvature), they can be quite different from similar null conditions, which are sometimes related to the conformal geometry of the manifold.

Let $R(x, y) := \nabla_x \nabla_y - \nabla_y \nabla_x - \nabla_{[x, y]}$ be the curvature operator of \mathcal{M} . The associated *Jacobi operator* $\mathcal{F}_R(x) : y \rightarrow R(y, x)x$ encodes much of the manifold's geometric information. The rescaling property $\mathcal{F}_R(\lambda v) = \lambda^2 \mathcal{F}_R(v)$ plays a crucial role. Let $S^\pm(\mathcal{M})$ be the unit sphere bundles of spacelike and timelike unit tangent vectors in M , and let $N(\mathcal{M})$ be the null cone of nonzero null vectors. One says that \mathcal{M} is spacelike Osserman if the eigenvalues of \mathcal{F}_R are constant on $S^+(\mathcal{M})$; one says instead timelike if they are constant on $S^-(\mathcal{M})$. Normalizing the length

MSC2000: primary 53C20, 53C50; secondary 53B30.

Keywords: spacelike, timelike and null Jacobi operator, Osserman and Jordan Osserman metric, neutral signature $(2, 2)$.

García-Río, Vázquez-Abal, and Vázquez-Lorenzo are supported by projects MTM2006-01432 and PGIDIT06PXIB207054PR (Spain). Gilkey is partially supported by the Max Planck Institute in the Mathematical Sciences (Leipzig, Germany) and by project MTM2006-01432 (Spain).

of the tangent vector to be ± 1 takes into account the above scaling of the Jacobi operator. Perhaps surprisingly, spacelike Osserman and timelike Osserman are equivalent conditions [García-Río et al. 1999; Gilkey 2001].

We say that \mathcal{M} is *null Osserman* if the eigenvalues of \mathcal{F}_R are constant on the null cone $N(\mathcal{M})$; with this definition, if \mathcal{M} is null Osserman, then necessarily $\mathcal{F}_R(v)$ is nilpotent if $v \in N(\mathcal{M})$ and $\mathcal{F}_R(v)$ has only the eigenvalue 0. Any spacelike or timelike Osserman manifold is necessarily null Osserman; the converse can fail in general — see for example [García-Río et al. 1997] in the Lorentzian setting.

The Jordan normal form plays a crucial role in the higher signature setting — a self-adjoint linear transformation need not be determined by its eigenvalues if the metric in question is indefinite. One says that \mathcal{M} is spacelike, timelike, or null Jordan Osserman if the Jordan normal form of $\mathcal{F}_R(\cdot)$ is constant on $S^+(\mathcal{M})$, on $S^-(\mathcal{M})$, or on $N(\mathcal{M})$, respectively. It is known from [Gilkey 2001; Gilkey and Ivanova 2002; 2001] that spacelike and timelike Jordan Osserman are inequivalent conditions; further neither necessarily implies the null Jordan Osserman condition.

In this paper, we concentrate on the 4-dimensional setting. Chi [1988] showed that any Riemannian Osserman 4-manifold is locally isometric to a 2-point homogeneous space; from later work [Blažić et al. 1997; García-Río et al. 1997], it follows that any Lorentzian 4-manifold has constant sectional curvature. However the situation is much more complicated in neutral signature $(2, 2)$; there exist many examples of nonsymmetric Osserman pseudo-Riemannian manifolds of neutral signature — see [Díaz-Ramos et al. 2006b] and [García-Río et al. 1998]. Despite the results of [Alekseevsky et al. 1999; Blažić et al. 2001; Díaz-Ramos et al. 2006a; García-Río and Vázquez-Lorenzo 2001], it is still an open problem to completely describe 4-dimensional Osserman metrics of neutral signature.

It is convenient to work algebraically. Let V be a finite-dimensional real vector space that is equipped with a nondegenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ of signature (p, q) . Let $A \in \otimes^4(V^*)$ be an algebraic curvature tensor on V , that is, a tensor that has the symmetries of the Riemann curvature tensor:

$$\begin{aligned} A(x, y, z, v) &= -A(y, x, z, v) = A(z, v, x, y), \\ A(x, y, z, v) + A(y, z, x, v) + A(z, x, y, v) &= 0. \end{aligned}$$

This defines a model $\mathfrak{M} := (V, \langle \cdot, \cdot \rangle, A)$. We often prove results on the algebraic level (that is, for models), and then obtain corresponding conclusions in the geometric context. The notions spacelike unit vector, timelike unit vector, null vector, Jacobi operator, and so on extend naturally to this setting.

1.1. Null Osserman algebraic curvature tensors. Henceforth, suppose $\langle \cdot, \cdot \rangle$ is an inner product of signature $(2, 2)$ on a 4-dimensional real vector space V . Fix

an orientation of V , and let $\mathcal{B} = \{e_1, e_2, e_3, e_4\}$ be an oriented orthonormal basis for V , where e_1 and e_2 are timelike and where e_3 and e_4 are spacelike.

At the algebraic level, in signature $(2, 2)$ the conditions spacelike Osserman, timelike Osserman, spacelike Jordan Osserman and timelike Jordan Osserman are equivalent to the condition that \mathfrak{M} is Einstein and self-dual with respect to a suitably chosen local orientation [Alekseevsky et al. 1999; García-Río et al. 2002]. In Section 2, we will establish Theorem 1.2, which shows that these conditions are also equivalent to null Osserman:

Theorem 1.2. *Let \mathfrak{M} be a model of neutral signature $(2, 2)$. Then the following conditions are equivalent:*

- (1) \mathfrak{M} is spacelike Osserman.
- (2) \mathfrak{M} is timelike Osserman.
- (3) \mathfrak{M} is spacelike Jordan Osserman.
- (4) \mathfrak{M} is timelike Jordan Osserman.
- (5) \mathfrak{M} is Einstein and self-dual for a suitably chosen local orientation.
- (6) \mathfrak{M} is null Osserman.

Remark 1.3. The action of homothety on the null vectors is a central one in this subject. With our definition, it is immediate that $\mathfrak{M} = (V, \langle \cdot, \cdot \rangle, A)$ is null Osserman implies that 0 is the only eigenvalue of \mathcal{F}_A on $N(V, \langle \cdot, \cdot \rangle)$. There is, although, an alternative, and different, formulation. One says that \mathfrak{M} is *projectively null Osserman* if either \mathfrak{M} is null Osserman or if given $0 \neq n_1, n_2 \in N(V, \langle \cdot, \cdot \rangle)$, there is a nonzero constant λ such that $\text{Spec}(\mathcal{F}_A(n_1)) = \lambda \text{Spec}(\mathcal{F}_A(n_2))$. We refer to [Brozos-Vázquez et al. 2008] for related work; we only introduce this concept for the sake of completeness as it plays no role in our development.

1.4. Null Jordan Osserman algebraic curvature tensors. Two algebraic curvature tensors will play a distinguished role. If Ψ is an skew-adjoint endomorphism of V , define the associated algebraic curvature tensor A^Ψ by setting

$$(1-1) \quad A^\Psi(x, y, z, v) := \langle \Psi y, z \rangle \langle \Psi x, v \rangle - \langle \Psi x, z \rangle \langle \Psi y, v \rangle - 2 \langle \Psi x, y \rangle \langle \Psi z, v \rangle.$$

Such tensors span the linear space of all algebraic curvature tensors [Fiedler 2002].

The sectional curvature of a nondegenerate 2-plane $\pi = \text{Span}\{x, y\}$ is given by

$$K_A(\pi) := \frac{A(x, y, y, x)}{\langle x, x \rangle \langle y, y \rangle - \langle x, y \rangle \langle x, y \rangle};$$

A has *constant sectional curvature* κ_0 if and only if $A = \kappa_0 A^0$, where A^0 is the algebraic curvature tensor of constant sectional curvature $+1$ defined by

$$(1-2) \quad A^0(x, y, z, v) := \langle y, z \rangle \langle x, v \rangle - \langle x, z \rangle \langle y, v \rangle.$$

We note that (1-1) and (1-2) imply that

$$(1-3) \quad \mathcal{F}_{A^\Psi}(x) : y \rightarrow 3\langle y, \Psi x \rangle \Psi x \quad \text{and} \quad \mathcal{F}_{A^0}(x) : y \rightarrow \langle x, x \rangle y - \langle x, y \rangle x.$$

Assume that Ψ is skew-adjoint. We say Ψ is an *orthogonal complex structure* if $\Psi^2 = -\text{id}$ and say Ψ is an *adapted paracomplex structure* if $\Psi^2 = \text{id}$. We say that a triple of skew-adjoint operators $\{\Psi_1, \Psi_2, \Psi_3\}$ is a *paraquaternionic structure* if $\Psi_1^2 = -\text{id}$, $\Psi_2^2 = \text{id}$, $\Psi_3^2 = \text{id}$, and if $\Psi_i \Psi_j + \Psi_j \Psi_i = 0$ for $i \neq j$. We can define a paraquaternionic structure by setting

$$(1-4) \quad \begin{aligned} \Psi_1 e_1 &= -e_2, & \Psi_1 e_2 &= e_1, & \Psi_1 e_3 &= e_4, & \Psi_1 e_4 &= -e_3, \\ \Psi_2 e_1 &= e_3, & \Psi_2 e_2 &= e_4, & \Psi_2 e_3 &= e_1, & \Psi_2 e_4 &= e_2, \\ \Psi_3 e_1 &= e_4, & \Psi_3 e_2 &= -e_3, & \Psi_3 e_3 &= -e_2, & \Psi_3 e_4 &= e_1. \end{aligned}$$

Note that $\Psi_3 = \Psi_1 \Psi_2$. If $\{\bar{\Psi}_1, \bar{\Psi}_2, \bar{\Psi}_3\}$ is another paraquaternionic structure on V , there is an isometry ϕ of V such that $\phi^* \bar{\Psi}_1 = \Psi_1$, $\phi^* \bar{\Psi}_2 = \Psi_2$, and $\phi^* \bar{\Psi}_3 = \pm \Psi_3$; this slight sign ambiguity plays no role in our constructions.

Let x be a spacelike or timelike vector. Then there is an orthogonal direct sum decomposition $V = \mathbb{R}x \oplus x^\perp$. Since $\mathcal{F}_A(x)x = 0$, $\mathcal{F}_A(x)$ preserves x^\perp . There are four different possibilities that describe the Jordan normal form of $\mathcal{F}_A(x)$ restricted to x^\perp (see [Blažić et al. 2001; García-Río et al. 2002] for further details):

$$(1-5) \quad \begin{array}{cccc} \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{pmatrix}, & \begin{pmatrix} \alpha & -\beta & 0 \\ \beta & \alpha & 0 \\ 0 & 0 & \gamma \end{pmatrix}, & \begin{pmatrix} \beta & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 1 & \alpha \end{pmatrix}, & \begin{pmatrix} \alpha & 0 & 0 \\ 1 & \alpha & 0 \\ 0 & 1 & \alpha \end{pmatrix}. \\ \text{Type Ia} & \text{Type Ib} & \text{Type II} & \text{Type III} \end{array}$$

Type Ia corresponds to a diagonalizable operator, **Type Ib** to an operator with a complex eigenvalue and **Type II** (respectively **Type III**) to a double (respectively triple) root of the minimal polynomial of the operator. If \mathfrak{M} is spacelike, timelike, or null Osserman, then the Jordan normal form of \mathcal{F}_A is constant on the spacelike and timelike unit vectors, and we classify A according to the four types above. In **Section 3**, we construct, up to isomorphism, all the spacelike Jordan Osserman algebraic curvature tensors and perform the analysis necessary to establish the following classification result:

Theorem 1.5. *Let $\mathfrak{M} := (V, \langle \cdot, \cdot \rangle, A)$ be a model of signature $(2, 2)$. Then \mathfrak{M} is null Jordan Osserman if and only if A is of **Type Ia** and one of the following holds:*

- (1) *There exists a constant κ_0 such that $A = \kappa_0 A^0$.*
- (2) *There exists constants κ_0 and κ_J with $\kappa_J \neq 0$ such that $A = \kappa_0 A^0 + \kappa_J A^J$, where J is an orthogonal complex structure on V .*
- (3) *There exists a constant $\kappa_P \neq 0$ such that $A = \kappa_P A^P$, where P is an adapted paracomplex structure on V .*

- (4) *There exist constants $\kappa_1, \kappa_2, \kappa_3$ such that $\kappa_2\kappa_3(\kappa_2 + \kappa_1)(\kappa_3 + \kappa_1) > 0$, such that the associated eigenvalues $\{3\kappa_1, -3\kappa_2, -3\kappa_3\}$ are all distinct, and such that $A = \kappa_1 A^{\Psi_1} + \kappa_2 A^{\Psi_2} + \kappa_3 A^{\Psi_3}$, where (Ψ_1, Ψ_2, Ψ_3) is a paraquaternionic structure on V .*

Remark 1.6. The inequality $\kappa_2\kappa_3(\kappa_2 + \kappa_1)(\kappa_3 + \kappa_1) > 0$ is equivalent to the cross ratio satisfying

$$(0, \kappa_1, -\kappa_3, -\kappa_2) = \frac{\kappa_3(\kappa_2 + \kappa_1)}{\kappa_2(\kappa_3 + \kappa_1)} > 0.$$

Let \mathbb{S}^2 be the unit sphere in \mathbb{R}^3 . This inequality is equivalent to the fact that the set of points $(0, -\kappa_3, -\kappa_2)$ and $(\kappa_1, -\kappa_3, -\kappa_2)$ give (via the stereographic projection) the corresponding circles in \mathbb{S}^2 the same orientation [Marden 2007].

1.7. Null Jordan Osserman manifolds. We characterize those neutral signature 4-manifolds that are null Jordan Osserman; null Osserman and null Jordan Osserman are not equivalent conditions, as the analysis of Section 3.7 shows. We say that \mathcal{M} is locally a complex space form if it is an indefinite Kähler manifold of constant holomorphic sectional curvature. In Section 4, we will use Theorem 1.5 to establish the following geometric result:

Theorem 1.8. *If \mathcal{M} is a connected pseudo-Riemannian manifold of neutral signature $(2, 2)$, then \mathcal{M} is null Jordan Osserman if and only if either \mathcal{M} has constant sectional curvature or \mathcal{M} is locally a complex space form.*

Remark 1.9. There is another family of Osserman 4-manifolds with diagonalizable Jacobi operator, namely, the paracomplex space forms [Blažić et al. 2001]. Although the geometry of complex and paracomplex space forms is very similar, the Jordan–Osserman condition distinguishes them. To our knowledge, this is the first algebraic curvature condition that distinguishes these two geometries.

2. Null Osserman models of signature $(2, 2)$

We work in the algebraic context to prove Theorem 1.2. Here is a brief outline to this section. Previous work establishes that parts (1)–(5) are equivalent. In Section 2.1, we introduce notation and show that spacelike Osserman models are null Osserman and that null Osserman models are Einstein. Thus to complete the proof, it suffices to show null Osserman models are self-dual or anti-self-dual. In Section 2.3, we examine Einstein models. Lemma 2.4 describes the Weyl curvature operators in that setting, and Lemma 2.5 gives an alternative characterization of self-duality for an Einstein model. We use Lemma 2.5 to complete the proof of Theorem 1.2 in Section 2.6.

2.1. Notation. Let $\mathfrak{M} := (V, \langle \cdot, \cdot \rangle, A)$ be a neutral signature 4-dimensional model. We use the inner product to raise indices and to define an associated Jacobi operator \mathcal{F}_A , which is characterized by the identity $\langle \mathcal{F}_A(x)y, z \rangle = A(y, x, x, z)$. Let $\mathcal{B} = \{e_1, e_2, e_3, e_4\}$ be an oriented orthonormal basis for V as in [Section 1.1](#). Let $g_{ij} := \langle e_i, e_j \rangle$, and let g^{ij} be the inverse matrix. The associated *Ricci tensor* ρ_A , the *scalar curvature* τ_A , and the *Weyl tensor* W_A are then defined by setting

$$\begin{aligned} \rho_A(x, y) &:= \sum_{i,j=1}^4 g^{ij} A(e_i, x, y, e_j), & \tau_A &:= \sum_{i,j=1}^4 g^{ij} \rho_A(e_i, e_j), \\ W_A(x, y, z, v) &:= A(x, y, z, v) + \frac{1}{6} \tau_A (\langle y, z \rangle \langle x, v \rangle - \langle x, z \rangle \langle y, v \rangle) \\ &\quad - \frac{1}{2} (\rho_A(y, z) \langle x, v \rangle - \rho_A(x, z) \langle y, v \rangle \\ &\quad + \rho_A(x, v) \langle y, z \rangle - \rho_A(y, v) \langle x, z \rangle). \end{aligned}$$

Let $A_{ijkl} = A_{ijkl}^{\mathcal{B}} := A(e_i, e_j, e_k, e_l)$ denote the components of A with respect to \mathcal{B} , where $1 \leq i, j, k, l \leq 4$; we drop the dependence on \mathcal{B} from the notation when there is no danger of confusion. Let $\{e^1, \dots, e^4\}$ be the dual basis for V^* . The Hodge operator $\star : \Lambda^p(V^*) \rightarrow \Lambda^{4-p}(V^*)$ is characterized by the identity

$$\phi_p \wedge \star \theta_p = \langle \phi_p, \theta_p \rangle e^1 \wedge e^2 \wedge e^3 \wedge e^4.$$

Thus,

$$\begin{aligned} \star(e^1 \wedge e^2) &= e^3 \wedge e^4, & \star(e^1 \wedge e^3) &= e^2 \wedge e^4, & \star(e^1 \wedge e^4) &= -e^2 \wedge e^3, \\ \star(e^2 \wedge e^3) &= -e^1 \wedge e^4, & \star(e^2 \wedge e^4) &= e^1 \wedge e^3, & \star(e^3 \wedge e^4) &= e^1 \wedge e^2. \end{aligned}$$

A crucial feature of 4-dimensional geometry now enters. Since $\star^2 = \text{id}$, the Hodge star induces a splitting $\Lambda^2(V^*) = \Lambda^+ \oplus \Lambda^-$ of the space of 2-forms, where

$$\Lambda^+ = \{\alpha \in \Lambda^2 : \star \alpha = \alpha\} \quad \text{and} \quad \Lambda^- = \{\alpha \in \Lambda^2 : \star \alpha = -\alpha\}$$

denote the spaces of *self-dual* and *anti-self-dual* two-forms. We have orthonormal bases $\{E_1^\mp, E_2^\mp, E_3^\mp\}$ for Λ^\mp that are given by

$$\begin{aligned} E_1^\mp &= \frac{1}{\sqrt{2}}(e^1 \wedge e^2 \mp e^3 \wedge e^4), & E_2^\mp &= \frac{1}{\sqrt{2}}(e^1 \wedge e^3 \mp e^2 \wedge e^4), \\ E_3^\mp &= \frac{1}{\sqrt{2}}(e^1 \wedge e^4 \pm e^2 \wedge e^3), \end{aligned}$$

where the induced inner product on Λ^\mp has signature (2, 1):

$$\langle E_1^\mp, E_1^\mp \rangle = 1, \quad \langle E_2^\mp, E_2^\mp \rangle = -1, \quad \langle E_3^\mp, E_3^\mp \rangle = -1.$$

Let W_A^\mp be the restriction of W_A to the spaces Λ^\mp , that is, $W_A^\mp : \Lambda^\mp \rightarrow \Lambda^\mp$, where W_A also stands for the associated Weyl curvature operator on Λ^2 . We say \mathfrak{M} is *self-dual* if $W_A^- = 0$ and *anti-self-dual* if $W_A^+ = 0$.

Lemma 2.2. *Let $\mathfrak{M} = (V, \langle \cdot, \cdot \rangle, A)$ be a model of signature (2, 2).*

- (1) If \mathfrak{M} is spacelike Osserman, then \mathfrak{M} is null Osserman.
 (2) If \mathfrak{M} is null Osserman, then \mathfrak{M} is Einstein.

Proof. Suppose first \mathfrak{M} is spacelike Osserman. Set $T_j(v) := \text{Tr}\{\mathcal{F}_A(v)^j\}$. Since the eigenvalues of \mathcal{F}_A are constant on $S^+(V, \langle \cdot, \cdot \rangle)$, there are constants c_j such that $T_j(v) = c_j$ for $v \in S^+(V, \langle \cdot, \cdot \rangle)$. It follows since $T_j(\lambda v) = \lambda^{2j} T_j(v)$ that $T_j(v) = c_j \langle v, v \rangle^j$ for v spacelike. Since the spacelike vectors form an open subset of V , this polynomial identity holds for all $v \in V$. Thus, in particular, $T_j(v) = 0$ if $v \in N(V, \langle \cdot, \cdot \rangle)$. This implies that 0 is the only eigenvalue of $\mathcal{F}_A(v)$ and shows \mathfrak{M} is null Osserman.

Now suppose \mathfrak{M} is null Osserman. Let s_1 and s_2 be spacelike unit vectors. We may choose a unit timelike vector t that is perpendicular to s_1 and s_2 . Let $n_i^\pm := s_i \pm t$ be null vectors. Thus $0 = \text{Tr}(\mathcal{F}_A(n_i^\pm)) = \rho_A(n_i^\pm, n_i^\pm)$, and

$$0 = \rho_A(s_i \pm t, s_i \pm t) = \rho_A(s_i, s_i) + \rho_A(t, t) \pm 2\rho_A(s_i, t).$$

This implies $\rho_A(s_i, t) = 0$ and $\rho_A(s_i, s_i) + \rho_A(t, t) = 0$; in particular, one has $\rho_A(s_1, s_1) = -\rho_A(t, t) = \rho_A(s_2, s_2)$. Therefore, after rescaling, there is a constant c such that $\rho_A(s, s) = c\langle s, s \rangle$ for every spacelike vector s ; this polynomial identity then continues to hold for all $s \in V$. Polarizing this identity then yields $\rho_A = c\langle \cdot, \cdot \rangle$, and hence \mathfrak{M} is Einstein. \square

2.3. The Weyl tensor for an Einstein algebraic curvature tensor. Let

$$\begin{aligned} \sigma_1 &= 2A_{1212} + 3\varepsilon A_{1234} + A_{1313} + A_{1414}, \\ \sigma_2 &= A_{1212} + 2A_{1313} + 3\varepsilon A_{1324} - A_{1414}, \\ \sigma_3 &= A_{1212} + 3\varepsilon A_{1234} - A_{1313} - 3\varepsilon A_{1324} + 2A_{1414}. \end{aligned}$$

Then we have an immediate lemma:

Lemma 2.4. *If \mathfrak{M} is Einstein, then the self-dual Weyl curvature operator W_A^+ (in which $\varepsilon = 1$) and the anti-self-dual Weyl curvature operator W_A^- (in which $\varepsilon = -1$) are given by*

$$\begin{pmatrix} \sigma_1/3 & A_{1213} + \varepsilon A_{1224} & A_{1214} - \varepsilon A_{1223} \\ -A_{1213} - \varepsilon A_{1224} & -\sigma_2/3 & -A_{1314} + \varepsilon A_{1323} \\ -A_{1214} + \varepsilon A_{1223} & -A_{1314} + \varepsilon A_{1323} & -\sigma_3/3 \end{pmatrix}.$$

The next observation is of interest in its own right:

Lemma 2.5. *If \mathfrak{M} is Einstein, then the model \mathfrak{M} is anti-self-dual if and only if $A_{1214}^{\mathfrak{B}} - A_{1223}^{\mathfrak{B}} = 0$ for every oriented orthonormal frame \mathfrak{B} .*

Proof. If \mathfrak{M} is anti-self-dual, we set $\varepsilon = 1$ in [Lemma 2.4](#) to see $A_{1214}^{\mathfrak{B}} - A_{1223}^{\mathfrak{B}} = 0$. Conversely, suppose $A_{1214}^{\mathfrak{B}} - A_{1223}^{\mathfrak{B}} = 0$ for every \mathfrak{B} . Define a new basis $\tilde{\mathfrak{B}}$ by

setting $\tilde{e}_1 = e_1$, $\tilde{e}_2 = e_2$, $\tilde{e}_3 = e_4$, and $\tilde{e}_4 = -e_3$. We then have

$$0 = -A_{1214}^{\tilde{\mathfrak{B}}} + A_{1223}^{\tilde{\mathfrak{B}}} = A_{1213}^{\mathfrak{B}} + A_{1224}^{\mathfrak{B}}.$$

Next, define $\tilde{\mathfrak{B}}$ by setting $\tilde{e}_1 = e_1$, $\tilde{e}_2 = \cosh \theta e_2 + \sinh \theta e_3$, $\tilde{e}_3 = \sinh \theta e_2 + \cosh \theta e_3$, and $\tilde{e}_4 = e_4$. This yields the relation

$$0 = -A_{1214}^{\tilde{\mathfrak{B}}} + A_{1223}^{\tilde{\mathfrak{B}}} = \cosh \theta (-A_{1214}^{\mathfrak{B}} + A_{1223}^{\mathfrak{B}}) + \sinh \theta (-A_{1314}^{\mathfrak{B}} + A_{1323}^{\mathfrak{B}}).$$

This shows $-A_{1314}^{\mathfrak{B}} + A_{1323}^{\mathfrak{B}} = 0$. Thus, by [Lemma 2.4](#),

$$W_A^+ = \frac{1}{3} \begin{pmatrix} \sigma_1^{\mathfrak{B}} & 0 & 0 \\ 0 & -\sigma_2^{\mathfrak{B}} & 0 \\ 0 & 0 & -\sigma_3^{\mathfrak{B}} \end{pmatrix}.$$

Setting the \tilde{e}_i as before yields bases for Λ^\pm in the form

$$\tilde{E}_1^\pm = \cosh \theta E_1^\pm + \sinh \theta E_2^\pm, \quad \tilde{E}_2^\pm = \cosh \theta E_2^\pm + \sinh \theta E_1^\pm, \quad \tilde{E}_3^\pm = E_3^\pm.$$

We may compute

$$\begin{aligned} W_A^+ \tilde{E}_1^+ &= \sigma_1^{\tilde{\mathfrak{B}}} \tilde{E}_1^+ = \sigma_1^{\tilde{\mathfrak{B}}} (\cosh \theta E_1^+ + \sinh \theta E_2^+) \\ &= W_A^+ (\cosh \theta E_1^+ + \sinh \theta E_2^+) = \sigma_1^{\mathfrak{B}} \cosh \theta E_1^+ - \sigma_2^{\mathfrak{B}} \sinh \theta E_2^+. \end{aligned}$$

This shows $\sigma_1^{\tilde{\mathfrak{B}}} = \sigma_1^{\mathfrak{B}} = -\sigma_2^{\mathfrak{B}}$. A similar argument applied to the basis $\tilde{e}_1 = e_1$, $\tilde{e}_2 = \cosh \theta e_2 + \sinh \theta e_4$, $\tilde{e}_3 = e_3$, and $\tilde{e}_4 = \sinh \theta e_2 + \cosh \theta e_4$ yields $\sigma_1^{\tilde{\mathfrak{B}}} = -\sigma_3^{\mathfrak{B}}$. Since $\sigma_1^{\mathfrak{B}} - \sigma_2^{\mathfrak{B}} - \sigma_3^{\mathfrak{B}} = 0$, it now follows that $W_A^+ = 0$. \square

2.6. Proof of Theorem 1.2. Let \mathfrak{M} be a null Osserman model. By [Lemma 2.2](#), \mathfrak{M} is Einstein. We complete the proof of [Theorem 1.2](#) by showing \mathfrak{M} is self-dual or anti-self-dual. Suppose the contrary and argue for a contradiction. As \mathfrak{M} is null Osserman, \mathcal{F}_A is nilpotent, so the characteristic polynomial has $p_\lambda(\mathcal{F}_A(u)) = \lambda^4$. Let

$$\begin{aligned} \mathcal{E}_1 &:= A_{1212} + 2A_{1214} - 2A_{1223} + 2A_{1234} - A_{1324} + A_{1414}, \\ Q(a, b) &:= (A_{1212} - 2A_{1214} - 2A_{1223} - 2A_{1234} + A_{1324} + A_{1414})a^4 \\ &\quad + (A_{1212} + 2A_{1214} + 2A_{1223} - 2A_{1234} + A_{1324} + A_{1414})b^4 \\ &\quad + 2(A_{1212} + 2A_{1313} - 3A_{1324} - A_{1414})a^2b^2 \\ &\quad + 4(A_{1213} - A_{1224} - A_{1314} - A_{1323})a^3b \\ &\quad + 4(A_{1213} - A_{1224} + A_{1314} + A_{1323})ab^3. \end{aligned}$$

If we take $u = ae_1 + be_2 + ae_3 + be_4$, then $\lambda^4 = p_\lambda(\mathcal{F}_A(u)) = \lambda^2(\lambda^2 - Q(a, b)\mathcal{E}_1)$. As $p_\lambda(\mathcal{F}_A(u)) = \lambda^4$, either $Q(a, b) = 0$ or $\mathcal{E}_1 = 0$. If we suppose that $\mathcal{E}_1 \neq 0$, then

$Q(a, b)$ vanishes identically for all a, b . This leads to the relations

$$\begin{aligned} A_{1213} - A_{1224} &= 0, & A_{1214} + A_{1223} &= 0, & A_{1314} + A_{1323} &= 0, \\ A_{1234} + A_{1313} - 2A_{1324} - A_{1414} &= 0, & A_{1212} + 2A_{1313} - 3A_{1324} - A_{1414} &= 0. \end{aligned}$$

From this, we see that the matrix in [Lemma 2.4](#) vanishes for $\varepsilon = -1$. This means that the anti-self-dual Weyl curvature operator W_A^- vanishes, so \mathfrak{M} is self-dual, which is a contradiction. Thus for *any* oriented orthonormal frame, we have

$$(2-1) \quad 0 = A_{1212} + 2A_{1214} - 2A_{1223} + 2A_{1234} - A_{1324} + A_{1414}.$$

Setting $\tilde{e}_1 = -e_1$, $\tilde{e}_2 = e_2$, $\tilde{e}_3 = e_3$, and $\tilde{e}_4 = -e_4$ yields

$$(2-2) \quad 0 = A_{1212} - 2A_{1214} + 2A_{1223} + 2A_{1234} - A_{1324} + A_{1414}.$$

Subtracting (2-2) from (2-1) then yields the relation $0 = -A_{1214} + A_{1223}$. We may now use [Lemma 2.5](#) to complete the proof of [Theorem 1.2](#). \square

3. Proof of [Theorem 1.5](#)

Here is a brief outline of this section. In [Section 3.1](#), we construct, up to isomorphism, all spacelike Jordan Osserman models of signature $(2, 2)$. In the remainder of [Section 3](#), we analyze each possible Jordan normal form in some detail using the classification of (1-5). Sections [3.5–3.8](#) deal with [Type Ia](#) models. In [Section 3.5](#) we study the case when all the eigenvalues are equal; this gives rise to [Theorem 1.5\(1\)](#). In [Section 3.6](#), we study the case of two equal spacelike eigenvalues, and in [Section 3.7](#), we study equal timelike and spacelike eigenvalues; these involve parts (2) and (3) of [Theorem 1.5](#), respectively. In [Section 3.8](#), we study [Type Ia](#) models with distinct eigenvalues; this leads to [Theorem 1.5\(4\)](#). We complete the proof of [Theorem 1.5](#) by showing the remaining types do not give rise to null Jordan Osserman models. We study [Type Ib](#) models in [Section 3.9](#), [Type II](#) models in [Section 3.10](#), and [Type III](#) models in [Section 3.11](#).

3.1. Spacelike Jordan Osserman models. We use the ansatz from [[Gilkey and Ivanova 2001](#)]. Let $\{\Psi_1, \Psi_2, \Psi_3\}$ be the paraquaternionic structure given in (1-4). Let $\xi_{ij} \in \mathbb{R}$ for $1 \leq i \leq j \leq 3$, and let $\kappa_0 \in \mathbb{R}$ be given. Let

$$(3-1) \quad \begin{aligned} A_{\kappa_0, \xi} &:= \kappa_0 A^0 + \frac{1}{3} \xi_{11} A^{\Psi_1} + \frac{1}{3} \xi_{22} A^{\Psi_2} + \frac{1}{3} \xi_{33} A^{\Psi_3} \\ &\quad + \frac{1}{3} \xi_{12} A^{\Psi_1 + \Psi_2} + \frac{1}{3} \xi_{13} A^{\Psi_1 + \Psi_3} + \frac{1}{3} \xi_{23} A^{\Psi_2 + \Psi_3}, \\ \mathcal{F}_{\kappa_0, \xi} &:= \kappa_0 \text{id} + \begin{pmatrix} \xi_{11} + \xi_{12} + \xi_{13} & & -\xi_{12} & & -\xi_{13} \\ & \xi_{12} & -\xi_{22} - \xi_{12} & -\xi_{23} & -\xi_{23} \\ & & \xi_{13} & -\xi_{23} & -\xi_{33} - \xi_{13} - \xi_{23} \end{pmatrix}. \end{aligned}$$

Lemma 3.2. *Adopt the notation established above. Let $\mathfrak{M}_{\kappa_0, \xi} := (V, \langle \cdot, \cdot \rangle, A_{\kappa_0, \xi})$.*

- (1) If $x \in S^\pm(V, \langle \cdot, \cdot \rangle)$, then $\mathcal{F}_{A_{\kappa_0, \zeta}}(x)$ is conjugate to the matrix $\pm \mathcal{F}_{\kappa_0, \zeta}$.
- (2) The model $\mathfrak{M}_{\kappa_0, \zeta}$ is spacelike and timelike Jordan Osserman.
- (3) Let $\mathfrak{M}_i = (V, \langle \cdot, \cdot \rangle, A_i)$ be spacelike Osserman models of signature $(2, 2)$. If $\mathcal{F}_{A_1}(x)$ is conjugate to $\mathcal{F}_{A_2}(x)$ for some $x \in S^\pm(V, \langle \cdot, \cdot \rangle)$, then there exists an isometry ϕ of $(V, \langle \cdot, \cdot \rangle)$ such that $\phi^* A_2 = A_1$.

Remark 3.3. Any self-adjoint map of a signature $(2, 1)$ vector space is conjugate to $\mathcal{F}_{\kappa_0, \zeta}$ for some $\{\kappa_0, \zeta\}$, so every spacelike Osserman model of signature $(2, 2)$ is isomorphic to one given by (3-1).

Proof. We suppose x is a spacelike unit vector; the timelike case is similar. Let $f_1 := \Psi_1 x$, $f_2 := \Psi_2 x$, and $f_3 := \Psi_3 x$. Then $\{f_1, f_2, f_3\}$ is an orthonormal basis of signature $(+, -, -)$ for x^\perp . Let $\mathcal{F} := \mathcal{F}_{A_{\kappa_0, \zeta}}(x)$. We use (1-3) to see that

$$\begin{aligned}\mathcal{F}f_1 &= (\kappa_0 + \zeta_{11} + \zeta_{12} + \zeta_{13})f_1 + \zeta_{12}f_2 + \zeta_{13}f_3, \\ \mathcal{F}f_2 &= -\zeta_{12}f_1 + (\kappa_0 - \zeta_{22} - \zeta_{12} - \zeta_{23})f_2 - \zeta_{23}f_3, \\ \mathcal{F}f_3 &= -\zeta_{13}f_1 - \zeta_{23}f_2 + (\kappa_0 - \zeta_{33} - \zeta_{13} - \zeta_{23})f_3.\end{aligned}$$

Part (1) now follows; part (2) follows from part (1). Suppose that \mathfrak{M} is a **Type Ia** spacelike Osserman model, so $\mathcal{F}_A(x) = \text{diag}[\alpha, \beta, \gamma]$ for any x in $S^+(V, \langle \cdot, \cdot \rangle)$; choose the notation so $\text{Ker}(\mathcal{F}_A(x) - \alpha \text{id})$ is spacelike. It then follows from the discussion in [Blažić et al. 2001; García-Río et al. 2002] that there is an orthonormal basis \mathcal{B} such that the nonzero components of the curvature tensor are

$$\begin{aligned}A_{1221} &= A_{4334} = \alpha, & A_{1331} &= A_{2442} = -\beta, \\ A_{1441} &= A_{3223} = -\gamma, & A_{1234} &= (-2\alpha + \beta + \gamma)/3, \\ A_{1423} &= (\alpha + \beta - 2\gamma)/3, & A_{1342} &= (\alpha - 2\beta + \gamma)/3.\end{aligned}$$

Similar forms exist for the other types of (1-5). Thus the Jordan normal form of $\mathcal{F}_A(x)$ determines A up to the action of $O(2, 2)$. Part (3) follows. \square

We immediately have this:

Lemma 3.4. *A null Osserman model \mathfrak{M} of signature $(2, 2)$ is null Jordan Osserman if and only if the functions $\text{Rank}\{\mathcal{F}_A(\cdot)\}$ and $\text{Rank}\{\mathcal{F}_A(\cdot)^2\}$ are constant on $N(V, \langle \cdot, \cdot \rangle)$.*

3.5. Type Ia with all eigenvalues equal: $\alpha = \beta = \gamma$. We set $A = \kappa_0 A^0$. By Lemma 3.2, the Jordan normal form is given by $\text{diag}[\kappa_0, \kappa_0, \kappa_0]$. If v belongs to $N(V, \langle \cdot, \cdot \rangle)$, then $\mathcal{F}_A(v)y = -\kappa_0 \langle v, y \rangle v$, and hence \mathfrak{M} is null Jordan Osserman.

3.6. Type Ia with two equal spacelike eigenvalues: $\beta = \gamma$ and $\alpha \neq \beta$. Let J be an orthogonal almost complex structure on V , and let $A = \kappa_0 A^0 + \kappa_J A^J$. The Jordan normal form is then given by $\text{diag}[\kappa_0 + 3\kappa_J, \kappa_0, \kappa_0]$, which has the desired form for suitably chosen κ_0 and κ_J with $\kappa_J \neq 0$. Let $v \in N(V, \langle \cdot, \cdot \rangle)$. We have

$$\mathcal{F}_A(v)y = -\kappa_0 \langle v, y \rangle v + 3\kappa_J \langle y, Jv \rangle Jv.$$

Because $J^2 = -\text{id}$, the vectors v and Jv are linearly independent. We note that $\langle v, v \rangle = \langle v, Jv \rangle = \langle Jv, Jv \rangle = 0$. Consequently $\mathcal{F}_A(v)v = \mathcal{F}_A(v)Jv = 0$. Since v^\perp and Jv^\perp are distinct 3-dimensional subspaces, we can choose y so $\langle v, y \rangle = 1$ and $\langle Jv, y \rangle = 0$. It now follows that $\mathcal{F}_A(v)y = -\kappa_0 v$, while $\mathcal{F}_A(v)Jy = 3\kappa_J Jv$. Thus $\mathcal{F}_A(v)$ has rank 2 and $\mathcal{F}_A(v)^2 = 0$. This implies A is null Jordan Osserman.

3.7. Type Ia with equal timelike and spacelike eigenvalues: $\alpha = \beta$ and $\beta \neq \gamma$.

Let $A = \kappa_0 A^0 + \kappa_P A^P$, where $\kappa_P \neq 0$ and where P is an adapted paracomplex structure; the Jordan normal form is then given by $\text{diag}[\kappa_0, \kappa_0 - 3\kappa_P, \kappa_0]$, which has the desired form for suitably chosen parameters. If $v \in N(V, \langle \cdot, \cdot \rangle)$, then

$$\mathcal{F}_A(v)y = -\kappa_0 \langle v, y \rangle v + 3\kappa_P \langle y, Pv \rangle Pv.$$

If $\kappa_0 = 0$, \mathfrak{M} is null Jordan Osserman. Suppose $\kappa_0 \neq 0$. If $v = e_1 + Pe_1$, then $Pv = v$, so $\text{Rank}\{\mathcal{F}_A(v)\} \leq 1$. On the other hand, if $v = e_1 + e_4$, then v and Pv are linearly independent, so $\text{Rank}\{\mathcal{F}_A(v)\} = 2$ and \mathfrak{M} is not null Jordan Osserman.

3.8. Type Ia with three distinct eigenvalues. We set $A := \sum_i \kappa_i A^{\Psi_i}$, where the triple $\{\Psi_1, \Psi_2, \Psi_3\}$ is the paraquaternionic structure of (1-4); the Jordan normal form is given by $\text{diag}[3\kappa_1, -3\kappa_2, -3\kappa_3]$, which has the desired form for suitably chosen parameters with

$$\kappa_1 + \kappa_2 \neq 0, \quad \kappa_1 + \kappa_3 \neq 0, \quad \kappa_2 - \kappa_3 \neq 0.$$

Let $\tilde{e} \in S^+(V, \langle \cdot, \cdot \rangle)$, let $V_+ := \text{Span}\{\tilde{e}, \Psi^1 \tilde{e}\}$, and let $V_- = V_+^\perp = \text{Span}\{\Psi_2 \tilde{e}, \Psi_3 \tilde{e}\}$. We then have an orthogonal direct sum decomposition $V = V_- \oplus V_+$, where V_+ is spacelike and V_- is timelike. Decompose $v \in N(V, \langle \cdot, \cdot \rangle)$ as $v = \lambda(e_+ + e_-)$, where $e_\pm \in V_\pm$. Let \mathfrak{M} be spacelike Osserman. We have $\mathcal{F}_A(v) = \lambda^2 \mathcal{F}_A(e_+ + e_-)$. Since $\mathcal{F}_A(v)$ is nilpotent, $\mathcal{F}_A(v)$ and $\mathcal{F}_A(e_+ + e_-)$ have the same Jordan normal form. Thus we may safely take $\lambda = 1$, so $v = e_+ + e_-$. Set $e = e_+$ and expand $e_- = \cos \theta \Psi_2 e + \sin \theta \Psi_3 e$. This expresses

$$v = e + \cos \theta \Psi_2 e + \sin \theta \Psi_3 e \quad \text{for } e \in S^+(V, \langle \cdot, \cdot \rangle).$$

We use the relations $\Psi_1\Psi_2 = \Psi_3$, $\Psi_1\Psi_3 = -\Psi_2$, and $\Psi_2\Psi_3 = -\Psi_1$ to see that

$$(3-2) \quad \begin{aligned} \Psi_1 v &= 0 + \Psi_1 e - \sin\theta\Psi_2 e + \cos\theta\Psi_3 e, \\ \Psi_2 v &= \cos\theta e - \sin\theta\Psi_1 e + \Psi_2 e + 0, \\ \Psi_3 v &= \sin\theta e + \cos\theta\Psi_1 e + 0 + \Psi_3 e, \end{aligned}$$

so that $0 = \Psi_1 v + \sin\theta\Psi_2 v - \cos\theta\Psi_3 v$. Thus the vectors $\{\Psi_1 v, \Psi_2 v, \Psi_3 v\}$ span a 2-dimensional subspace. Since $\langle\Psi_i v, \Psi_j v\rangle = 0$, $\text{Span}\{\Psi_i v\} \subset \text{Ker}\{\mathcal{F}_A(v)\}$. Since $\text{Range}\{\mathcal{F}_A(v)\} \subset \text{Span}\{\Psi_i v\}$,

$$\text{Rank}\{\mathcal{F}_A(v)\} \leq 2 \quad \text{and} \quad \mathcal{F}_A(v)^2 = 0.$$

Note that $\{e, \Psi_1 e, \Psi_2 v, \Psi_3 v\}$ is a basis for V . Let π_+ denote orthogonal projection on $V_+ = \text{Span}\{e, \Psi_1 e\}$. Since π_+ is injective on $\text{Range}\{\mathcal{F}_A(v)\} \subset \text{Span}\{\Psi_2 v, \Psi_3 v\}$,

$$r(v) := \dim \text{Range}\{\mathcal{F}_A(v)\} = \dim(\text{Span}\{\pi_+\mathcal{F}_A(v)e, \pi_+\mathcal{F}_A(v)\Psi_1 e\}).$$

By (3-2) and the linear dependency it contains,

$$\begin{aligned} \mathcal{F}_A(v)e &= 3\kappa_2 \cos\theta\Psi_2 v + 3\kappa_3 \sin\theta\Psi_3 v, \\ \mathcal{F}_A(v)\Psi_1 e &= 3\kappa_1\Psi_1 v - 3\kappa_2 \sin\theta\Psi_2 v + 3\kappa_3 \cos\theta\Psi_3 v, \\ \pi_+\mathcal{F}_A(v)e &= 3(\kappa_2 \cos\theta(\cos\theta) + \kappa_3 \sin\theta(\sin\theta))e \\ &\quad + 3(\kappa_2 \cos\theta(-\sin\theta) + \kappa_3 \sin\theta(\cos\theta))\Psi_1 e, \\ \pi_+\mathcal{F}_A(v)\Psi_1 e &= 3(-\kappa_2 \sin\theta(\cos\theta) + \kappa_3 \cos\theta(\sin\theta))e \\ &\quad + 3(\kappa_1 - \kappa_2 \sin\theta(-\sin\theta) + \kappa_3 \cos\theta(\cos\theta))\Psi_1 e. \end{aligned}$$

This leads to a coefficient matrix for $\pi_+\mathcal{F}_A(v)$ on V_+ given by

$$\mathcal{C}_A(\theta) = 3 \begin{pmatrix} \kappa_2 \cos^2\theta + \kappa_3 \sin^2\theta & (-\kappa_2 + \kappa_3) \sin\theta \cos\theta \\ (-\kappa_2 + \kappa_3) \sin\theta \cos\theta & \kappa_1 + \kappa_2 \sin^2\theta + \kappa_3 \cos^2\theta \end{pmatrix}.$$

We compute

$$\begin{aligned} \frac{1}{9} \det(\mathcal{C}_A)(\theta) &= \kappa_1\kappa_2 \cos^2\theta + \kappa_2^2 \cos^2\theta \sin^2\theta + \kappa_2\kappa_3 \cos^4\theta \\ &\quad + \kappa_1\kappa_3 \sin^2\theta + \kappa_2\kappa_3 \sin^4\theta + \kappa_3^2 \sin^2\theta \cos^2\theta \\ &\quad - \kappa_2^2 \sin^2\theta \cos^2\theta - \kappa_3^2 \sin^2\theta \cos^2\theta + 2\kappa_2\kappa_3 \sin^2\theta \cos^2\theta \\ &= \kappa_1\kappa_2 \cos^2\theta + \kappa_1\kappa_3 \sin^2\theta + \kappa_2\kappa_3 \\ &= (\kappa_1 + \kappa_3)\kappa_2 \cos^2\theta + (\kappa_1 + \kappa_2)\kappa_3 \sin^2\theta. \end{aligned}$$

Observe that $\kappa_2\kappa_3 = 0$ implies that $\det(\mathcal{C}_A)(\theta)$ vanishes for some θ , and thus \mathfrak{M} is not null Jordan Osserman. Hence, since $(\kappa_1 + \kappa_3)\kappa_2$ and $(\kappa_1 + \kappa_2)\kappa_3$ are nonzero, $\det(\mathcal{C}_A)(\theta)$ never vanishes, or equivalently \mathfrak{M} is null Jordan Osserman, if and only if these two real numbers have the same sign, that is, $\kappa_2\kappa_3(\kappa_1 + \kappa_3)(\kappa_1 + \kappa_2) > 0$.

3.9. Type Ib models. Let $b \neq 0$. We take a curvature tensor of the form

$$A = \frac{1}{3}((a-b)A^{\Psi_1} + (-b-a)A^{\Psi_2} + bA^{\Psi_1+\Psi_2} + cA^{\Psi_3}).$$

Proceeding as in the previous case, we have for any $e \in S^+(V, \langle \cdot, \cdot \rangle)$ that

$$\begin{aligned} \mathcal{F}_A(x)y &= \langle (a\Psi_1 + b\Psi_2)x, y \rangle \Psi_1x + \langle (b\Psi_1 - a\Psi_2)x, y \rangle \Psi_2x + c\langle \Psi_3x, y \rangle \Psi_3x, \\ \mathcal{F}_A(e)\Psi_1e &= a\Psi_1e + b\Psi_2e, \quad \mathcal{F}_A(e)\Psi_2e = -b\Psi_1e + a\Psi_2e, \quad \mathcal{F}_A(e)\Psi_3e = -c\Psi_3e. \end{aligned}$$

Thus $\mathfrak{M} := (V, \langle \cdot, \cdot \rangle, A)$ is **Type Ib** and any **Type Ib** model is isomorphic to \mathfrak{M} for suitably chosen parameters. As in [Section 3.8](#), put $v = e + \cos\theta\Psi_2e + \sin\theta\Psi_3e$. We compute

$$\begin{aligned} \mathcal{F}_A(v)e &= b\cos\theta\Psi_1v - a\cos\theta\Psi_2v + c\sin\theta\Psi_3v, \\ \mathcal{F}_A(v)\Psi_1e &= (a - b\sin\theta)\Psi_1v + (b + a\sin\theta)\Psi_2v + c\cos\theta\Psi_3v, \\ \pi_+\mathcal{F}_A(v)e &= (-a\cos\theta(\cos\theta) + c\sin\theta(\sin\theta))e \\ &\quad + (b\cos\theta - a\cos\theta(-\sin\theta) + c\sin\theta(\cos\theta))\Psi_1e, \\ \pi_+\mathcal{F}_A(v)\Psi_1e &= ((b + a\sin\theta)(\cos\theta) + c\cos\theta(\sin\theta))e \\ &\quad + ((a - b\sin\theta) + (b + a\sin\theta)(-\sin\theta) + c\cos\theta(\cos\theta))\Psi_1e. \end{aligned}$$

The coefficient matrix for $\pi_+\mathcal{F}_A(v)$ on V_+ is then given by

$$\mathcal{C}_A(\theta) = \begin{pmatrix} -a\cos^2\theta + c\sin^2\theta & b\cos\theta + (a+c)\sin\theta\cos\theta \\ b\cos\theta + (a+c)\sin\theta\cos\theta & -2b\sin\theta + (a+c)\cos^2\theta \end{pmatrix}.$$

We have $\det(\mathcal{C}_A)(\pi/2) = -2bc$ and $\det(\mathcal{C}_A)(-\pi/2) = 2bc$. If $c \neq 0$, then these signs differ and hence $\det(\mathcal{C}_A)(\theta) = 0$ for some $-\pi/2 < \theta < \pi/2$ and \mathfrak{M} is not null Jordan Osserman. If $c = 0$, then $\det(\mathcal{C}_A)(\pi/2) = 0$ and $\det(\mathcal{C}_A)(0) = -a^2 - b^2 \neq 0$ and again \mathfrak{M} is not null Jordan Osserman.

3.10. Type II models. We approach this case directly. Let $\mathfrak{M} = (V, \langle \cdot, \cdot \rangle, A)$ be a model of signature $(2, 2)$, where A is a **Type II** algebraic curvature tensor. Then the analysis of [[Blažić et al. 2001](#); [García-Río et al. 2002](#)] shows there exists an orthonormal basis $\{e_1, e_2, e_3, e_4\}$ for V such that the nonvanishing components of A are

$$\begin{aligned} A_{1221} &= A_{4334} = \pm(\alpha - \frac{1}{2}), & A_{1224} &= A_{1334} = \pm\frac{1}{2}, \\ A_{1331} &= A_{4224} = \mp(\alpha + \frac{1}{2}), & A_{2113} &= A_{2443} = \mp\frac{1}{2}, \\ A_{1234} &= (\pm(-\alpha + \frac{3}{2}) + \beta)/3, & A_{1423} &= 2(\pm\alpha - \beta)/3, \\ A_{1342} &= (\pm(-\alpha - \frac{3}{2}) + \beta)/3, & A_{1441} &= A_{3223} = -\beta. \end{aligned}$$

Let $u = e_2 - e_3$ and let $v = e_2 + e_3$. Then

$$\mathcal{F}_A(u) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \beta & \beta & 0 \\ 0 & -\beta & -\beta & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{F}_A(v) = \begin{pmatrix} \pm 2 & 0 & 0 & \mp 2 \\ 0 & \beta & -\beta & 0 \\ 0 & \beta & -\beta & 0 \\ \pm 2 & 0 & 0 & \mp 2 \end{pmatrix}.$$

If $\beta = 0$, then $r(u) = 0$ and $r(v) = 1$; if $\beta \neq 0$, then $r(u) = 1$ and $r(v) = 2$. Thus \mathfrak{M} is not null Jordan Osserman.

3.11. Type III models. For \mathfrak{M} of this type, there exists by [Blažić et al. 2001; García-Río et al. 2002] an orthonormal basis $\{e_1, e_2, e_3, e_4\}$ for V such that the nonvanishing components of A are

$$\begin{aligned} A_{1221} &= A_{4334} = \alpha, & A_{1331} &= A_{4224} = -\alpha, \\ A_{1441} &= A_{3223} = -\alpha, \\ A_{2114} &= A_{2334} = -\sqrt{2}/2, & A_{3114} &= -A_{3224} = \sqrt{2}/2, \\ A_{1223} &= A_{1443} = A_{1332} = -A_{1442} = \sqrt{2}/2. \end{aligned}$$

Let $u = e_2 - e_3$ and $v = e_2 + e_3$. Then

$$\mathcal{F}_A(u) = \begin{pmatrix} 0 & -\sqrt{2} & -\sqrt{2} & 0 \\ -\sqrt{2} & \alpha & \alpha & \sqrt{2} \\ \sqrt{2} & -\alpha & -\alpha & -\sqrt{2} \\ 0 & -\sqrt{2} & -\sqrt{2} & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{F}_A(v) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \alpha & -\alpha & 0 \\ 0 & \alpha & -\alpha & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

It now follows that $r(u) = 2$ while $r(v) \leq 1$ and hence \mathfrak{M} is not null Jordan Osserman. This completes the proof of [Theorem 1.5](#). \square

4. The proof of [Theorem 1.8](#)

Let \mathcal{M} be a null Jordan Osserman manifold of signature $(2, 2)$. First note that, by [Theorem 1.5](#), \mathcal{M} has [Type Ia](#). Results of [Blažić et al. 2001] then show that \mathcal{M} either has constant sectional curvature, is locally isometric to a complex space form, or is locally isometric to a paracomplex space form. Since the curvature tensor of a paracomplex space form of constant paraholomorphic sectional curvature κ satisfies

$$R(x, y)z = \frac{1}{4}\kappa(R^0(x, y)z - R^J(x, y)z),$$

this is ruled out by [Theorem 1.5](#), thus proving [Theorem 1.8](#). \square

References

- [Alekseevsky et al. 1999] D. Alekseevsky, N. Blažić, N. Bokan, and Z. Rakić, “Self-duality and pointwise Osserman manifolds”, *Arch. Math. (Brno)* **35**:3 (1999), 193–201. [MR 2000j:53094](#) [Zbl 1054.53082](#)
- [Blažić et al. 1997] N. Blažić, N. Bokan, and P. Gilkey, “A note on Osserman Lorentzian manifolds”, *Bull. London Math. Soc.* **29**:2 (1997), 227–230. [MR 97m:53111](#) [Zbl 0865.53018](#)
- [Blažić et al. 2001] N. Blažić, N. Bokan, and Z. Rakić, “Osserman pseudo-Riemannian manifolds of signature $(2, 2)$ ”, *J. Aust. Math. Soc.* **71**:3 (2001), 367–395. [MR 2002g:53119](#) [Zbl 1011.53049](#)
- [Brozos-Vázquez et al. 2008] M. Brozos-Vázquez, P. Gilkey, S. Nikčević, and U. Simon, “Projectively Osserman manifolds”, *Publ. Math. Debrecen* **72**:3-4 (2008), 359–370. [MR 2406926](#) [Zbl 05367462](#)
- [Chi 1988] Q.-S. Chi, “A curvature characterization of certain locally rank-one symmetric spaces”, *J. Differential Geom.* **28**:2 (1988), 187–202. [MR 90a:53060](#) [Zbl 0654.53053](#)
- [Díaz-Ramos et al. 2006a] J. C. Díaz-Ramos, E. García-Río, and R. Vázquez-Lorenzo, “Four-dimensional Osserman metrics with nondiagonalizable Jacobi operators”, *J. Geom. Anal.* **16**:1 (2006), 39–52. [MR 2006m:53107](#) [Zbl 1105.53052](#)
- [Díaz-Ramos et al. 2006b] J. C. Díaz-Ramos, E. García-Río, and R. Vázquez-Lorenzo, “New examples of Osserman metrics with nondiagonalizable Jacobi operators”, *Differential Geom. Appl.* **24**:4 (2006), 433–442. [MR 2007h:53106](#) [Zbl 1099.53047](#)
- [Fiedler 2002] B. Fiedler, “Determination of the structure of algebraic curvature tensors by means of Young symmetrizers”, *Sém. Lothar. Combin.* **48** (2002), Art. B48d. [MR 2004d:53018](#)
- [García-Río and Vázquez-Lorenzo 2001] E. García-Río and R. Vázquez-Lorenzo, “Four-dimensional Osserman symmetric spaces”, *Geom. Dedicata* **88**:1-3 (2001), 147–151. [MR 2002k:53137](#) [Zbl 1008.53023](#)
- [García-Río et al. 1997] E. García-Río, D. N. Kupeli, and M. E. Vázquez-Abal, “On a problem of Osserman in Lorentzian geometry”, *Differential Geom. Appl.* **7**:1 (1997), 85–100. [MR 98a:53099](#) [Zbl 0880.53017](#)
- [García-Río et al. 1998] E. García-Río, M. E. Vázquez-Abal, and R. Vázquez-Lorenzo, “Nonsymmetric Osserman pseudo-Riemannian manifolds”, *Proc. Amer. Math. Soc.* **126**:9 (1998), 2771–2778. [MR 98m:53089](#) [Zbl 0939.53014](#)
- [García-Río et al. 1999] E. García-Río, D. N. Kupeli, M. E. Vázquez-Abal, and R. Vázquez-Lorenzo, “Affine Osserman connections and their Riemann extensions”, *Differential Geom. Appl.* **11**:2 (1999), 145–153. [MR 2000i:53030](#) [Zbl 0940.53017](#)
- [García-Río et al. 2002] E. García-Río, D. N. Kupeli, and R. Vázquez-Lorenzo, *Osserman manifolds in semi-Riemannian geometry*, Lecture Notes in Mathematics **1777**, Springer, Berlin, 2002. [MR 2003e:53052](#) [Zbl 1005.53040](#)
- [Gilkey 2001] P. B. Gilkey, *Geometric properties of natural operators defined by the Riemann curvature tensor*, World Scientific Publishing, River Edge, NJ, 2001. [MR 2002k:53052](#) [Zbl 1007.53001](#)
- [Gilkey and Ivanova 2001] P. Gilkey and R. Ivanova, “The Jordan normal form of Osserman algebraic curvature tensors”, *Results Math.* **40**:1-4 (2001), 192–204. [MR 2002j:53093](#) [Zbl 0999.53014](#)
- [Gilkey and Ivanova 2002] P. B. Gilkey and R. Ivanova, “Spacelike Jordan Osserman algebraic curvature tensors in the higher signature setting”, pp. 179–186 in *Differential geometry* (Valencia, 2001), edited by O. Gil-Medrano and V. Miquel, World Scientific Publishing, River Edge, NJ, 2002. [MR 2003f:53125](#) [Zbl 1031.53037](#)

[Marden 2007] A. Marden, *Outer circles: An introduction to hyperbolic 3-manifolds*, Cambridge University Press, 2007. MR 2008i:57001 Zbl 1149.57030

Received April 3, 2008. Revised September 15, 2009.

EDUARDO GARCÍA-RÍO
DEPARTMENT OF GEOMETRY AND TOPOLOGY
FACULTY OF MATHEMATICS
UNIVERSITY OF SANTIAGO DE COMPOSTELA
15782 SANTIAGO DE COMPOSTELA
SPAIN

eduardo.garcia.rio@usc.es

PETER B. GILKEY
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OREGON
EUGENE, OR 97403
UNITED STATES

gilkey@uoregon.edu

M. ELENA VÁZQUEZ-ABAL
DEPARTMENT OF GEOMETRY AND TOPOLOGY
FACULTY OF MATHEMATICS
UNIVERSITY OF SANTIAGO DE COMPOSTELA
15782 SANTIAGO DE COMPOSTELA
SPAIN

elena.vazquez.abal@usc.es

RAMÓN VÁZQUEZ-LORENZO
DEPARTMENT OF GEOMETRY AND TOPOLOGY
FACULTY OF MATHEMATICS
UNIVERSITY OF SANTIAGO DE COMPOSTELA
15782 SANTIAGO DE COMPOSTELA
SPAIN

ravazlor@edu.xunta.es