

*Pacific
Journal of
Mathematics*

**A NEW PROOF OF
REIFENBERG'S TOPOLOGICAL DISC THEOREM**

GUANGHAO HONG AND LIHE WANG

Volume 246 No. 2

June 2010

A NEW PROOF OF REIFENBERG'S TOPOLOGICAL DISC THEOREM

GUANGHAO HONG AND LIHE WANG

We offer a new proof to the classical topological disk theorem of Reifenberg. The novelty of our method is that we construct the approximating surfaces globally, which makes our proof rather simple and direct.

1. Introduction

Definition 1.1 (Reifenberg flat set). We say that a compact set $K \subseteq \bar{B}_1^n \subseteq \mathbb{R}^n$ is an m -dimensional (δ, R) -Reifenberg flat set if for every $a \in K$ and $r \in (0, R]$ with $|a| + r \leq 1$, there exists an m -plane $T_{a,r} \in \text{GL}(m, n)$ such that

$$HD(K \cap B_r^n(a), T_{a,r} \cap B_r^n(a)) \leq \delta r,$$

where $HD(A, B) = \max(\sup\{\text{dist}(a, B) : a \in A\}, \sup\{\text{dist}(b, A) : b \in B\})$ is the Hausdorff distance.

The definition is only meaningful for small $\delta > 0$.

Theorem 1.2 (Reifenberg's topological disk theorem). *For K as above, if δ is sufficiently small, then $K \cap B_{1/2}^n$ is a C^α -topological m -dimensional disk.*

This theorem is adopted from the book [Lin and Yang 2002, Chapter 2, page 58]. The original and its proof is due to Reifenberg [1960, Chapter 4]. In that paper, Reifenberg investigated the higher-dimensional Plateau problem. He started with this kind of very weak “surfaces”; the C^α topological disk property was used in a fundamental way. Both his original proof and our new one are based on a sequence of approximating surfaces converging to K . The first such surface is just a standard m -dimensional disk, and there exist uniform bi-Lipschitz maps between any two adjacent surfaces. The composition of these bi-Lipschitz maps gives a bi-Hölder map from B_1^m to K . What is crucial is the actual construction of these surfaces. Reifenberg did the work locally, using several averaging processes and complicated estimates, which made his proof horribly difficult and messy. Recently, the authors had the idea of constructing these surfaces globally by mollifying the distance function to K , which made it possible to write down a much simpler proof

MSC2000: primary 49Q15; secondary 28A15.

Keywords: topological disk, Reifenberg flat, mollification.

of this beautiful theorem. In fact, we define the bi-Lipschitz maps between two adjacent level sets of the mollified distance functions to be the gradient vector of the mollified distance functions. These level sets will converge to K , but they are codimension 1 surfaces. It is easy to extract an m -dimensional surface from the first of these level sets, and we track its images under these bi-Lipschitz maps step by step. Then we obtain a sequence of m -dimensional surfaces converging to K . It is crucial in this process to show that at every scale and every neighborhood the corresponding surface is almost parallel to K . The uniform bound $C(m, n)\delta$ for the error angles guarantees the uniform bi-Lipschitz property of the restricted maps.

In [Hong and Wang 2007], we could only treat the codimension 1 case, when we took the level sets at $1/2$ of the mollified characteristic functions of Ω (here $\partial\Omega$ is the Reifenberg flat set) as the approximating surfaces. The method of this paper works for any codimension.

2. Proof of Theorem 1.2

We will use standard notations. Let e_1, \dots, e_n be the standard basis in \mathbb{R}^n . Let O be the origin of \mathbb{R}^n . We use the same notation in another coordinate system if no confusion arises. For simplicity, we assume $T_{O,1} = \text{span}\{e_1, \dots, e_m\}$. For fixed $m < n$, we write $x = (x', x'') \in \mathbb{R}^n$ with $x' = (x_1, \dots, x_m)$ and $x'' = (x_{m+1}, \dots, x_n)$, or $x = (x', x''', x_n)$ with $x''' = (x_{m+1}, \dots, x_{n-1})$. Then $T_{O,1} = \{x \in \mathbb{R}^n : |x''| = 0\}$. We denote the distance function from K by d and the distance functions from $T_{a,10r}$ by $d_{a,r}$; then $d_{O,0.1}(x) = |x''|$.

2.1. Derivative estimates on the mollified distance functions. Our mollifier η is defined so that $\eta(x) = C_1 \exp(-1/(1 - |x|^2))$ for $x \in B_1$ and $\eta(x) = 0$ for x outside of B_1 , where C_1 is chosen so that $\int \eta = 1$. Through the paper, C (with any subscript) denotes a constant only depending on dimensions m and n . Define $\eta_\epsilon(x) := \epsilon^{-n} \eta(\epsilon^{-1}x)$. Then $d_\epsilon := d * \eta_\epsilon$ are the mollified distance functions.

We define $h_{a,r} := d - d_{a,r}$; then $HD(K \cap B_{10r}(a), T_{a,10r} \cap B_{10r}(a)) \leq 10\delta r$ implies $|h_{a,r}| \leq 10\delta r$ in $B_{9.9r}(a)$. Thus $d_\epsilon = d_{a,r} * \eta_\epsilon + h_{a,r} * \eta_\epsilon$; in particular

$$d_{0.01r} = d_{a,r} * \eta_{0.01r} + h_{a,r} * \eta_{0.01r} := \tilde{d}_{a,r} + \tilde{h}_{a,r}.$$

The following two lemmas give derivative estimates on $\tilde{d}_{a,r}$ and $\tilde{h}_{a,r}$ near point a .

Lemma 2.1. *For any unit vector v perpendicular to $T_{a,10r}$, any $b \in T_{a,10r}$ and t satisfying $0.4r < t$, we have $\nabla \tilde{d}_{a,r}(b + tv) = \tau v$ with $0.95 < \tau < 1.05$; moreover, if we chose a coordinate system such that $b = O$, $T_{a,10r} = \text{span}\{e_1, \dots, e_m\}$ and $v = e_n$, then*

$$\begin{aligned} |\partial_i \partial_j \tilde{d}_{a,r}(b + tv)| &< 3/r \quad \text{if } m < i, j \leq n, \\ |\partial_i \partial_j \tilde{d}_{a,r}(b + tv)| &= 0 \quad \text{if } \min(i, j) \leq m. \end{aligned}$$

Proof. In the coordinate system above, $d_{a,r}(x) = |x''|$. Then for $x \notin T_{a,r}$, we have $\nabla d_{a,r}(x) = (0, x''/|x''|)$ and $\partial_i \partial_j d_{a,r}(x) = \delta_{ij}/|x''| - x_i x_j / |x''|^3$ when $\min(i, j) > m$ and $\partial_i \partial_j d_{a,r}(x) = 0$ when $\min(i, j) \leq m$. Therefore,

$$\begin{aligned} \nabla \tilde{d}_{a,r}(tv) &= \int_{B_{0.01r}} \frac{(0, (te_n - y)'')}{|(te_n - y)''|} \cdot \eta_{0.01r}(y) dy \\ &= \int_{B_1} \frac{(0, (te_n - 0.01ry)'')}{|(te_n - 0.01ry)''|} \cdot \eta(y) dy \\ &= \int_{B_1} \frac{(0, (e_n - 0.01(r/t)y)'')}{|(e_n - 0.01(r/t)y)''|} \cdot \eta(y) dy. \end{aligned}$$

Now $\partial_i \tilde{d}_{a,r}(tv) = 0$ for $1 \leq i \leq m$, whereas

$$\partial_i \tilde{d}_{a,r}(tv) = \int_{B_1} \frac{-0.01(r/t)y_i}{|(e_n - 0.01(r/t)y)''|} \cdot \eta(y) dy = 0 \quad \text{for } m + 1 \leq i \leq n - 1$$

by symmetry properties of the functions about the hyperplane $\{x_i = 0\}$; finally

$$\partial_n \tilde{d}_{a,r}(tv) = \int_{B_1} \frac{(1 - 0.01(r/t)y_n)}{|(e_n - 0.01(r/t)y)''|} \cdot \eta(y) dy.$$

Both $1 - 0.01(r/t)y_n$ and $|(e_n - 0.01(r/t)y)''|$ are between 0.975 and 1.025 for $y \in B_1$, which implies the first part of the conclusion. The estimates of the second order derivatives can be proved by the same way. \square

Lemma 2.2. *In $B_{9,8r}(a)$, we have*

$$|\tilde{h}_{a,r}| \leq 10\delta r, \quad |\nabla \tilde{h}_{a,r}| \leq C_2\delta, \quad |\nabla^2 \tilde{h}_{a,r}| \leq C_2\delta/r.$$

Proof. Put the derivatives onto $\eta_{0.01r}$ and use $|h_{a,r}| \leq 10\delta r$ in $B_{9,9r}(a)$. \square

2.2. Construction of approximating surfaces. We let $r_k := 0.01 \times 2^{-k}$ and define $L_k := \{x \in B_1 \mid d_{r_k/100} = r_k\}$ for $k = 1, 2, \dots$. The gradient estimation of $d_{r_k/100}$ and the implicit function theorem tell us that every L_k is a smooth hypersurface. Let $\bar{L}_k := \{x \in B_1 \mid d = r_k\}$. Then $|d - d_{r_k/100}| < r_k/100$ implies $HD(L_k, \bar{L}_k) < r_k/90$, and then $(1 - 1/80)r_k < HD(L_k, K) < (1 + 1/80)r_k$, where we used that δ is small enough.

We define $\pi_k : L_k \rightarrow L_{k+1}$ by $\pi_k(P) = P - \sigma(k, P)\nabla d_{r_k/100}(P)$ with a real number $\sigma(k, P)$ chosen so that $\pi_k(P) \in L_{k+1}$. It is easy to show that if δ is small enough, $0.9r_{k+1} < \sigma(k, P) < 1.1r_{k+1}$. We define

$$K_0 := \{x \in \mathbb{R}^n : |x'| < 0.8, |x''| = 0\}$$

to be a standard m -disk, and let

$$K_1 := L_1 \cap \{x \in \mathbb{R}^n : |x'| < 0.8, |x'''| = 0 \text{ and } x_n > 0\}.$$

Then K_1 can be written as $K_1 = \{x + h(x)e_n : x \in K_0\}$ with $0.98r_1 < h(x) < 1.02r_1$ and $|\nabla h(x)| < 0.1$ by the implicit function theorem. Hence $g_0(x) := x + h(x)e_n$ is a bi-Lipschitz map from K_0 to K_1 .

Then we define $K_{k+1} := \pi_k(K_k)$.

2.3. Uniform bound on the non-tangential angles. We need some notation for angles. If T is an m -plane, \tilde{T} is the m -plane parallel to T and containing the origin, and P is a vector in \mathbb{R}^n , then we define $\angle(P, T) := \angle(P, \tilde{T}) := \angle(P, \hat{P})$, where \hat{P} is the projection of P onto \tilde{T} . If T_1 is another m -plane, then $\angle(T_1, T) := \angle(\tilde{T}_1, \tilde{T}) := \sup\{\angle(P, \tilde{T}) : P \in \tilde{T}_1\}$. Now we are ready to prove the key lemma.

Lemma 2.3. *If δ is small enough, there exists a constant C_3 with $C_3\delta < 10^{-4}$ such that for any k and $P, Q \in L_k \cap B_{0,9}$ with $|P - Q| < 6r_k$, the inequality*

$$\angle(Q - P, \text{span}\{T_{a,10r_k}, P\}) < C_3\delta$$

implies

$$0.9 < |\pi_k(P) - \pi_k(Q)|/|P - Q| < 1.1,$$

$$\angle(\pi_k(Q) - \pi_k(P), \text{span}\{T_{\tilde{a},10r_{k+1}}, \pi_k(P)\}) < C_3\delta.$$

Here a is any of the points in K nearest to P , and \tilde{a} is one of the points in K nearest to $\pi_k(P)$.

Proof. Choose a coordinate system such that $T_{a,10r_k} = \{x \in \mathbb{R}^n : |x''| = 0\}$ and $P = |P|e_n$. Then $\text{span}\{T_a, 10r_k, P\} = \{x \in \mathbb{R}^n : |x'''| = 0\}$. Throughout this proof, we restrict our analysis to the inside of the box

$$U := \{x \in \mathbb{R}^n : |(x - P)'| < 8r_k, |(x - P)'''| < 8C_3\delta r_k, 0.8r_{k+1} < x_n < 1.2r_k\}.$$

Then L_k and L_{k+1} can be represented as the graphs of $(n - 1)$ -variable functions ϕ and ψ , respectively; for example, $L_k \cap U = \{x \in U : x_n = \phi(x', x''')\}$. By the implicit function theorem, the previous two lemmas tell us information about the derivatives of ϕ and ψ :

$$|\phi_i| = \left| \frac{-\partial_i d_{r_k/100}}{\partial_n d_{r_k/100}} \right| < 1.1C_2\delta, \quad |\psi_i| < 1.1C_2\delta \quad \text{for } 1 \leq i \leq m,$$

$$|\phi_i| < 9C_3\delta + 1.1C_2\delta, \quad |\psi_i| < 18C_3\delta + 1.1C_2\delta \quad \text{for } m + 1 \leq i \leq n - 1.$$

Then $|Q_n - |P|| < C_4\delta(|Q'| + |Q''''|)$ by the differential mean value theorem. Our angle assumption reads

$$\arctan \frac{|Q''''|}{\sqrt{|Q'|^2 + (Q_n - |P|)^2}} < C_3\delta,$$

which implies that $|Q''''| < 1.1C_3\delta|Q'|$ if δ is small enough. And $|P - Q| = (|Q'|^2 + |Q''''|^2 + (Q_n - |P|)^2)^{1/2}$, so $|Q'| \leq |P - Q| < 1.01|Q'|$ if δ is small enough.

We rewrite $\pi_k(Q) = Q - \sigma(k, Q)\nabla d_{r_k/100}(Q)$ as

$$\begin{aligned} \pi_k(Q) &= P - \sigma(k, P)\nabla d_{r_k/100}(P) \\ &\quad + (Q', 0, 0) + (0, Q''', Q_n - |P|) \\ &\quad + \sigma(k, Q)(\nabla d_{r_k/100}(P) - \nabla d_{r_k/100}(Q)) \\ &\quad + (\sigma(k, P) - \sigma(k, Q))\nabla d_{r_k/100}(P). \end{aligned}$$

Because of the estimates of the second order derivatives of $d_{r_k/100}$, the norm of the third row won't exceed $C_5\delta|Q'|$. In fact, along the Q' direction, the second order derivative is as small as $C_2\delta/r_k$ and the step length is $|Q'|$; along the $(Q - P)''$ direction, the second order derivative is bounded by $4/r_k$ but the step length is as small as $(1.1C_3 + 2C_4)\delta|Q'|$. The first row is nothing but $\pi_k(P)$. We denote the first three rows by I_1 . We know $d_{r_{k+1}/100}(\pi_k(P)) = r_{k+1}$, hence

$$\begin{aligned} |d_{r_{k+1}/100}(I_1) - r_{k+1}| &< C_2\delta|Q'| + 1.1 \times (1.1C_3 + 2C_4)\delta|Q'| + 1.1 \times C_5\delta|Q'| \\ &= C_6\delta|Q'|. \end{aligned}$$

Here we use $|\partial_i d_{r_{k+1}/100}| = |\partial_i \tilde{h}_{a,r_k}| < C_2\delta$ for $1 \leq i \leq m$ and $|\nabla d_{r_{k+1}/100}| < 1.1$. But $d_{r_{k+1}/100}(\pi_k(Q)) = r_{k+1}$, and $\partial_\zeta d_{r_{k+1}/100} > 0.8$ near $\pi_k(Q)$ with $\zeta := \nabla d_{r_k/100}(P)$. Therefore, we have $|\sigma(k, P) - \sigma(k, Q)| < 1.25C_6\delta|Q'|$.

Now we consider $E := P - \sigma(k, P)\nabla \tilde{d}_{a,r_k}(P)$, which equals $|P|e_n - \sigma(k, P)\tau_1 e_n$ for $0.95 < \tau_1 < 1.05$ by Lemma 2.1, so $E := \tau_2 r_k e_n$ with $0.4 < \tau_2 < 0.6$ if one notes that $|P|$ and $\sigma(k, P)$ are approximately r_k and $r_k/2$ respectively. Now

$$\begin{aligned} F &:= Q - \sigma(k, Q)\nabla \tilde{d}_{a,r_k}(Q) \\ &= (Q', 0) + (0, Q'') - \sigma(k, Q)\nabla \tilde{d}_{a,r_k}((Q', 0) + (0, Q'')). \end{aligned}$$

Since $1.05r_k > |Q''| > 0.95r_k$, by Lemma 2.1 we have

$$\tilde{d}_{a,r_k}((Q', 0) + (0, Q'')) = \tau_3(0, Q'')/|Q''| \quad \text{for } 0.95 < \tau_3 < 1.05.$$

So $F = (Q', 0) + \tau_4(0, Q'')$ with $0.4 < \tau_4 < 0.6$. Therefore

$$\tan \angle(F - E, \text{span}\{T_{a,10r_k}, P\}) = \frac{|(F - E)'''|}{\sqrt{|(F - E)'|^2 + (F - E)_n^2}} = \frac{|\tau_4 Q'''|}{\sqrt{|Q'|^2 + (F - E)_n^2}}.$$

But

$$(F - E)_n = (Q_n - |P|) + (\sigma(k, P) - \sigma(k, Q))\tau_1 + \sigma(k, Q)(\partial_n \tilde{d}_{a,r_k}(P) - \partial_n \tilde{d}_{a,r_k}(Q)),$$

so $|(F - E)_n| < (2C_4 + 2C_6 + C_5)\delta|Q'|$. Then if δ is small enough, we have

$$\tan \angle(F - E, \text{span}\{T_{a,10r_k}, P\}) < 0.61|Q'''|/|Q'| < 0.7C_3\delta,$$

and $|Q'| \leq |F - E| < 1.01|Q'|$.

We know

$$\pi_k(P) = E - \sigma(k, P)\nabla\tilde{h}_{a,r_k}(P) \quad \text{and} \quad \pi_k(Q) = F - \sigma(k, Q)\nabla\tilde{h}_{a,r_k}(Q).$$

So we define

$$G := F - \sigma(k, P)\nabla\tilde{h}_{a,r_k}(P) \quad \text{and} \quad H := G + (\sigma(k, P) - \sigma(k, Q))\nabla\tilde{h}_{a,r_k}(P).$$

Then

$$\pi_k(Q) = H + \sigma(k, Q)(\nabla\tilde{h}_{a,r_k}(P) - \nabla\tilde{h}_{a,r_k}(Q)) \quad \text{and} \quad G - \pi_k(P) = F - E.$$

However, $|H - G| < 1.25C_2C_6\delta^2|Q'| < C_2\delta|Q'|$ (if $C_6\delta < 0.8$) implies

$$\angle(H - \pi_k(P), G - \pi_k(P)) < 2C_2\delta,$$

and $0.99|Q'| < |H - \pi_k(P)| < 1.01|Q'|$ (if $C_2\delta < 10^{-3}$). Moreover

$$|\pi_k(Q) - H| < 0.55r_k \cdot C_2\delta/r_k \cdot 1.01|Q'| < C_2\delta|Q'|$$

implies

$$\angle(\pi_k(Q) - \pi_k(P), H - \pi_k(P)) < 2C_2\delta,$$

and $0.97|Q'| < |\pi_k(Q) - \pi_k(P)| < 1.03|Q'|$, which means

$$0.9 < \frac{|\pi_k(P) - \pi_k(Q)|}{|P - Q|} < 1.1.$$

The three angle inequalities above give

$$\angle(\pi_k(Q) - \pi_k(P), \text{span}\{T_{a,10r_k}, P\}) < (0.7C_3 + 4C_2)\delta.$$

On the other hand,

$$|(\pi_k(P))'''| < |\sigma(k, P)\nabla\tilde{h}_{a,r_k}(P)| < C_2\delta r_k \quad \text{and} \quad (\pi_k(P))_n > 0.4r_k$$

show that $\angle(\text{span}\{T_{a,10r_k}, \pi_k(P)\}, \text{span}\{T_{a,10r_k}, P\}) < 3C_2\delta$. We also know that $\angle(T_{\bar{a},10r_{k+1}}, T_{a,10r_k}) < 4\delta$, because K is close to either of these two m -planes. So finally, we can conclude

$$\angle(\pi_k(Q) - \pi_k(P), \text{span}\{T_{\bar{a},10r_{k+1}}, \pi_k(P)\}) < (0.7C_3 + 7C_2 + 4)\delta < C_3\delta$$

if we let $0.3C_3 > 7C_2 + 4$. □

K_1 satisfies the angle condition in the lemma above if $C_3 > 21$, so all the of K_k satisfy the angle condition and all of the $\pi_k|_{K_k}$ are bi-Lipschitz by induction.

2.4. Bi-Hölder parametrization. Define $g_k : K_1 \rightarrow K_{k+1}$, $x \mapsto \pi_k \circ \dots \circ \pi_1(x)$ and $f : K_1 \rightarrow K$, $x \mapsto \lim_{k \rightarrow \infty} g_k(x)$. We are ready to prove f is bi-Hölder continuous. Given x and $y \in K_1$, we write $\ell := |f(x) - f(y)|$. First notice that $|x - y| > 0$ implies $\ell > 0$ since

$$0.9^k |x - y| \leq |g_k(x) - g_k(y)| \leq 3r_k + \ell \quad \text{for all } k.$$

Now we choose k with $r_{k+1} < 0.01\ell \leq r_k$, which implies $-\log_2 \ell - 1 < k \leq -\log_2 \ell$. The triangle inequality gives

$$0.9\ell < |g_k(x) - g_k(y)| < 1.1\ell.$$

However $0.9^k |x - y| \leq |g_k(x) - g_k(y)| \leq 1.1^k |x - y|$, so we obtain

$$0.5|x - y|^\beta < \ell < 2|x - y|^\alpha \quad \text{with } \alpha = \frac{1}{1 + \log_2 1.1} \text{ and } \beta = \frac{1}{1 + \log_2 0.9}.$$

2.5. Completeness of the approximating surfaces. To show $f(K_1) \supset K \cap B_{1/2}^n$, we have to show K_k is complete enough so that any point of $K \cap B_{1/2}^n$ is close to it. A short topological argument suffices.

Lemma 2.4. *For any k and a given point $P \in K_k$, there exists an m -plane \tilde{T}_P going through P with $\angle(\tilde{T}_P, T_{a, 10r_k}) < 20\delta$, where a is the point in K nearest to P , such that $K_k \cap B_{5.1r_k}^n(P)$ contains the set $\{x + \zeta(x) : x \in \tilde{T}_P \cap B_{5r_k}^n(P)\}$, where $\zeta(x)$ is a vector-valued function defined on $\tilde{T}_P \cap B_{5r_k}^n(P)$ with $\zeta(x) \perp \tilde{T}_P$ and $|\zeta(x)| < 5(C_3 + 1.1C_2)\delta r_k$.*

Proof. We prove the lemma by induction. We know it is true for K_1 , and assume it is for K_k . Given $Q \in K_{k+1}$, define $\tilde{T}_Q := T_{\tilde{a}, 10r_{k+1}} + (Q - \tilde{a})$ with \tilde{a} being the point in K nearest to Q . For any point $y \in \tilde{T}_Q \cap B_{5r_{k+1}}^n(Q)$, we want to show that the $(n - m)$ -plane $N_y := \{x \in \mathbb{R}^n : x - y \perp \tilde{T}_Q\}$ must contain a point of K_{k+1} . Define $P := (\pi_k)^{-1}(Q) \in K_k$. Then we have $\angle(\tilde{T}_Q, \tilde{T}_P) < 23\delta$. Let \tilde{y} be the point in \tilde{T}_P nearest to y , and define $S_{\tilde{y}} := \{x \in \tilde{T}_P : |x - \tilde{y}| = r_k\}$ and $S_y := S_{\tilde{y}} + (y - \tilde{y})$. From Lemma 2.3, we know, for all $x \in S_{\tilde{y}}$, that

$$\begin{aligned} \text{dist}(\pi_k(x + \zeta(x)), S_y) &\leq |\pi_k(x + \zeta(x)) - (x + y - \tilde{y})| \\ &< |\zeta(x)| + |\sigma(k, x + \zeta(x)) \nabla d_{r_k/100}(x + \zeta(x)) - (y - \tilde{y})| \\ &< 0.3r_k. \end{aligned}$$

The $(m - 1)$ -sphere S_y cannot contract to a point without passing through N_y , and neither can the topological $(m - 1)$ -sphere $\{\pi_k(x + \zeta(x)) : x \in S_{\tilde{y}}\}$. So if $N_y \cap K_{k+1} = \emptyset$, then $\{\pi_k(x + \zeta(x)) : x \in S_{\tilde{y}}\}$ cannot contract to a point within K_{k+1} , which contradicts that K_{k+1} is a topological m -disk. Once we have a point $z \in N_y \cap K_{k+1}$, it is easy to see z is unique and $|z - y| < 5(C_3 + 1.1C_2)\delta r_{k+1}$ from Lemma 2.3. \square

Lemma 2.5. *For any point $b \in K \cap B_{1/2}^n$, we have $\text{dist}(b, K_k) < 1.1r_k$ for all k .*

Proof. We prove the lemma by induction. It is true for K_1 , and assume it is for K_k . Then we have $P \in K_k$ with $|b - P| < 1.1r_k$. Let a be the point in K nearest to P . Choose a local coordinate system such that $a = O$ and such that $T_{a,10r_k} = \{x \in \mathbb{R}^n : |x''| = 0\}$. Then $|b'| < r_k$ and $|b''| < 10\delta r_k$. Write $Q := \pi_k(P)$ and $E := (b', Q'')$, and let F be the point in \tilde{T}_Q nearest to E . Then $|Q''| < 1.02r_{k+1}$ and $|Q'| < r_{k+1}$. So $E - Q \in T_{a,10r_k}$ and $\angle(\tilde{T}_Q, T_{a,10r_k}) < 23\delta$ implies $|E - F| < 30\delta r_k$. Therefore,

$$\begin{aligned} \text{dist}(b, K_{k+1}) &\leq |b - (F + \zeta(F))| \\ &\leq |b - E| + |E - F| + |\zeta(F)| \\ &\leq |b''| + |Q''| + 30\delta r_k + 5(C_3 + 1.1C_2)\delta r_{k+1} < 1.1r_{k+1}. \quad \square \end{aligned}$$

Corollary 2.6. $K \cap B_{1/2}^n \subset f(K_1)$.

Proof. Given $b \in K \cap B_{1/2}^n$, for every k we have

$$\text{dist}(b, K_k) < 1.1r_k \quad \text{and} \quad HD(K_k, f(K_1)) < 1.1r_k.$$

Then $\text{dist}(b, f(K_1)) < 2.2r_k$. But $r_k \rightarrow 0$ and $f(K_1) \cap \bar{B}_{1/2}^n$ is closed. \square

References

- [Hong and Wang 2007] G. Hong and L. Wang, “A geometric approach to the topological disk theorem of Reifenberg”, *Pacific J. Math.* **233**:2 (2007), 321–339. MR 2008m:49203 Zbl 1155.49037
- [Lin and Yang 2002] F. Lin and X. Yang, *Geometric measure theory: An introduction*, Advanced Mathematics **1**, Science Press, Beijing and International Press, Boston, 2002. MR 2005a:28001 Zbl 1074.49011
- [Reifenberg 1960] E. R. Reifenberg, “Solution of the Plateau Problem for m -dimensional surfaces of varying topological type”, *Acta Math.* **104** (1960), 1–92. MR 22 #4972 Zbl 0099.08503

Received May 12, 2009. Revised November 16, 2009.

GUANGHAO HONG
XI'AN JIAOTONG UNIVERSITY
DEPARTMENT OF MATHEMATICS
XI'AN, SHAANXI 710049
CHINA
ghhongmath@gmail.com

LIHE WANG
UNIVERSITY OF IOWA
DEPARTMENT OF MATHEMATICS
14 MACLEAN HALL
IOWA CITY, IA 52242-1419
UNITED STATES
lwang@math.uiowa.edu
www.math.uiowa.edu/~lwang

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Paulo Ney de Souza, Production Manager

Silvio Levy, Senior Production Editor

Matthew Cargo, Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2010 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer-review and production is managed by EditFlow™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2010 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 246 No. 2 June 2010

Geometric structures associated to a contact metric (κ, μ) -space BENIAMINO CAPPELLETTI MONTANO and LUIGIA DI TERLIZZI	257
Multilinear singular operators with fractional rank CIPRIAN DEMETER, MALABIKA PRAMANIK and CHRISTOPH THIELE	293
A new proof of Reifenberg's topological disc theorem GUANGHAO HONG and LIHE WANG	325
Global classical solutions to hyperbolic geometric flow on Riemann surfaces FAGUI LIU and YUANZHANG ZHANG	333
An extension of the quintuple product identity and its applications ZHI-GUO LIU	345
A generalization of the Pontryagin–Hill theorems to projective modules over Prüfer domains JORGE MACÍAS-DÍAZ	391
Elliptic pseudodifferential equations and Sobolev spaces over p -adic fields J. J. RODRÍGUEZ-VEGA and W. A. ZÚÑIGA-GALINDO	407
Absolutely isolated singularities of holomorphic maps of \mathbb{C}^n tangent to the identity FENG RONG	421
Pullbacks of Eisenstein series from $\mathrm{GU}(3, 3)$ and critical L -values for $\mathrm{GSp}(4) \times \mathrm{GL}(2)$ ABHISHEK SAHA	435
Isomorphism invariants of restricted enveloping algebras HAMID USEFI	487
Spacelike S-Willmore spheres in Lorentzian space forms PENG WANG	495



0030-8730(201006)246:2;1-B