

*Pacific
Journal of
Mathematics*

**ISOMORPHISM INVARIANTS
OF RESTRICTED ENVELOPING ALGEBRAS**

HAMID USEFI

Volume 246 No. 2

June 2010

ISOMORPHISM INVARIANTS OF RESTRICTED ENVELOPING ALGEBRAS

HAMID USEFI

Let L and H be finite-dimensional restricted Lie algebras over a perfect field \mathbb{F} . Suppose $u(L) \cong u(H)$, where $u(L)$ is the restricted enveloping algebra of L . We prove that $L \cong H$ if L is p -nilpotent and abelian, or if L is abelian and \mathbb{F} is algebraically closed. We use these results to prove our main result, that if L is p -nilpotent, then $L/L'^p + \gamma_3(L) \cong H/H'^p + \gamma_3(H)$.

1. Introduction

Let L be a restricted Lie algebra with restricted enveloping algebra $u(L)$. We say that a particular invariant of L is *determined* by $u(L)$ if every restricted Lie algebra H also has this invariant whenever $u(L)$ and $u(H)$ are isomorphic as associative algebras. The restricted isomorphism problem asks whether the isomorphism type of L is determined by $u(L)$. This problem is motivated by the classical isomorphism problem for group rings: Is every finite group G determined by its integral group ring $\mathbb{Z}G$? The survey article [Sandling 1985] contains most of the development in this area. In the late 1980s, Roggenkamp and Scott [1987] and Weiss [1988] independently settled the group ring problem for finite nilpotent groups.

There are close analogies between restricted Lie algebras and finite p -groups. In particular, the restricted isomorphism problem is the Lie analogue of the modular isomorphism problem that asks, Given finite p -groups G and H with the property that $\mathbb{F}_p G \cong \mathbb{F}_p H$, can we deduce that $G \cong H$? Here, \mathbb{F}_p denotes the field of p elements. There has been intensive investigation on the modular isomorphism problem; however the main problem is rather far from being completely answered. Unfortunately not every technique from finite p -groups can be used for restricted Lie algebras. For example, it is known that the class sums form a basis of the center of $\mathbb{F}_p G$. It then follows that the center of G is determined; see [Sehgal 1978, Theorem 6.6]. Whether or not the center of L is determined by $u(L)$ remains an interesting open question.

MSC2000: primary 17B35, 17B50; secondary 20C05.

Keywords: restricted Lie algebras, enveloping algebras, isomorphism problem.

Research supported by an NSERC Postdoctoral Fellowship.

In analogy with finite p -groups, we consider the class \mathcal{F}_p of restricted Lie algebras that are finite-dimensional and p -nilpotent. Let $L \in \mathcal{F}_p$. It follows from Engel's theorem that L is nilpotent. We shall examine the nilpotence class of L in Corollary 2.2. Whether or not the nilpotence class of G is determined by $\mathbb{F}_p G$ has been considered in recent years; however no major result is reported at this time; see [Bagiński and Konovalov 2007].

We start investigating the restricted isomorphism problem by first considering the abelian case. In Proposition 2.5, we prove that if $L \in \mathcal{F}_p$ is an abelian restricted Lie algebra over a perfect field \mathbb{F} , then the isomorphism type of L is determined by $u(L)$. Furthermore, if \mathbb{F} is algebraically closed, then every abelian restricted Lie algebra is determined by its enveloping algebra; see Corollary 2.8.

It is not clear what the next step is beyond the abelian case in both the modular isomorphism problem and the restricted isomorphism problem. Nevertheless, we have proved in [Usefi 2008] that if $L \in \mathcal{F}_p$ is a metacyclic restricted Lie algebra over a perfect field, then the isomorphism type of L is determined by $u(L)$. The main result of this paper, which will be proved in Section 3, is another contribution in this direction; a similar result for finite p -groups was proved by Sandling [1989]. For a Lie subalgebra $I \subseteq L$, we denote by I^p the restricted Lie subalgebra of L generated by all x^p for $x \in I$. Also, $\gamma_i(L)$ denotes the i -th term of the lower central series of L . Our main result is as follows:

Theorem. *Let $L \in \mathcal{F}_p$ be a restricted Lie algebra over a perfect field. Then the restricted Lie algebra $L/(L^p + \gamma_3(L))$ is determined.*

2. Preliminaries

Let L be a restricted Lie algebra with restricted enveloping algebra $u(L)$ over a field \mathbb{F} . By the Poincaré–Birkhoff–Witt (PBW) theorem (see [Jacobson 1962]), we can view L as a restricted Lie subalgebra of $u(L)$. Denote by $\omega(L)$ the augmentation ideal of $u(L)$ that is the kernel of the augmentation map $\epsilon_L : u(L) \rightarrow \mathbb{F}$ induced by $x \mapsto 0$ for every $x \in L$.

Let H be another restricted Lie algebra such that $\varphi : u(L) \rightarrow u(H)$ is an algebra isomorphism. The map $\eta : L \rightarrow u(H)$ defined by $\eta = \varphi - \epsilon_H \varphi$ is a restricted Lie algebra homomorphism. Therefore, η extends to an algebra homomorphism $\bar{\eta} : u(L) \rightarrow u(H)$. In fact, $\bar{\eta}$ is an isomorphism preserving the augmentation ideals, that is, $\bar{\eta}(\omega(L)) = \omega(H)$; see [Riley and Usefi 2007] for the proof of a similar fact for Lie algebras. So, without loss of generality, we assume that $\varphi : u(L) \rightarrow u(H)$ is an algebra isomorphism that preserves the augmentation ideals.

Recall that L is said to be nilpotent if $\gamma_n(L) = 0$ for some n ; the nilpotence class $\text{cl}(L)$ of L is the minimal integer c such that $\gamma_{c+1}(L) = 0$. We denote by L'_p the restricted subalgebra of L generated by $L' = \gamma_2(L)$. The n -th dimension

subalgebra of L is

$$D_n(L) = L \cap \omega^n(L) = \sum_{ip^j \geq n} \gamma_i(L)^{p^j};$$

see [Riley and Shalev 1995].

Recall that L is said to be in the class \mathcal{F}_p if L is finite-dimensional and p -nilpotent. The *exponent* of $x \in L$, denoted by $\text{exp}(x)$, is the least integer s such that $x^{p^s} = 0$. Whether or not $L \in \mathcal{F}_p$ is determined by the following lemma; see [Riley and Shalev 1995].

Lemma 2.1. *Let L be a restricted Lie algebra. Then $L \in \mathcal{F}_p$ if and only if $\omega(L)$ is nilpotent.*

Now, consider the graded restricted Lie algebra

$$\text{gr}(L) := \bigoplus_{i \geq 1} D_i(L)/D_{i+1}(L),$$

where the Lie bracket and the p -map are induced from L . It is well known that $u(\text{gr}(L)) \cong \text{gr}(u(L))$ as algebras; see [Usefi 2008]. So we may identify $\text{gr}(L)$ as the graded restricted Lie subalgebra of $\text{gr}(u(L))$ generated by $\omega^1(L)/\omega^2(L)$. Thus, $\text{gr}(L)$ is determined. We can now deduce the following:

Corollary 2.2. *Let L and H be restricted Lie algebras such that $u(L) \cong u(H)$. If $L \in \mathcal{F}_p$, then $|\text{cl}(L) - \text{cl}(H)| \leq 1$.*

Proof. Let $c = \text{cl}(L)$. We note that

$$\gamma_n(\text{gr}(L)) = \bigoplus_{i \geq n} \gamma_i(L) + D_{i+1}(L)/D_{i+1}(L) \quad \text{for every } n \geq 1.$$

Since $\text{gr}(L)$ is determined, $\gamma_{c+1}(\text{gr}(H)) = 0$. Hence, $\gamma_{c+1}(H) \subseteq D_{c+2}(H)$. So, $\gamma_{c+2}(H) = \gamma_{c+3}(H)$. Since H is nilpotent, it follows that $\gamma_{c+2}(H) = 0$. \square

Note that $D_n(\text{gr}(L)) = \bigoplus_{i \geq n} D_i(L)/D_{i+1}(L)$. Thus, $D_n(L)/D_{n+1}(L)$ is determined for every $n \geq 1$. The methods of [Ritter and Sehgal 1983] and [Riley and Usefi 2007] can be adapted to prove that $D_n(L)/D_{2n+1}(L)$ and $D_n(L)/D_{n+2}(L)$ are also determined for every $n \geq 1$. In particular, $L/D_3(L)$ is determined. We need the following analogue of [Riley and Usefi 2007, Lemma 5.1].

Lemma 2.3. *If $\varphi : u(L) \rightarrow u(H)$ is an isomorphism, then $\varphi(D_n(L) + \omega^{n+1}(L)) = D_n(H) + \omega^{n+1}(H)$ for every positive integer n .*

Now suppose that L is an abelian restricted Lie algebra. The conditions on the p -map reduce to $(x + y)^p = x^p + y^p$ and $(\alpha x)^p = \alpha^p x^p$ for every $x, y \in L$ and $\alpha \in \mathbb{F}$. Thus the p -map is a semilinear transformation. Let σ be an automorphism of \mathbb{F} . Consider the skew polynomial ring $\mathbb{F}[t; \sigma]$ that consists of polynomials $f(t) \in \mathbb{F}[t]$ with multiplication given by $at^i \beta t^j = \alpha \beta^{\sigma^{-i}} t^{i+j}$. It is well known that $\mathbb{F}[t; \sigma]$

is a PID. Now suppose that \mathbb{F} is perfect and let σ be the automorphism given by $\sigma(\alpha) = \alpha^p$. Note that $\mathbb{F}[t; \sigma]$ acts on L by $x \cdot t = x^p$. Then, by the theory of finitely generated modules over a PID, L decomposes as a direct sum of cyclic $\mathbb{F}[t; \sigma]$ -modules, with a unique number of the summands. We summarize this in the following; see also [Jacobson 1962] or [Bahturin et al. 1992, Section 4.3]. We denote by $\langle x \rangle_p$ the subalgebra generated by x .

Theorem 2.4. *Let L be a finitely generated abelian restricted Lie algebra over a perfect field \mathbb{F} . Then there exist a unique integer n and generators $x_1, \dots, x_n \in L$ such that $L = \langle x_1 \rangle_p \oplus \dots \oplus \langle x_n \rangle_p$.*

Proposition 2.5. *Let $L \in \mathcal{F}_p$ be an abelian restricted Lie algebra over a perfect field \mathbb{F} . If H is a restricted Lie algebra such that $u(L) \cong u(H)$, then $L \cong H$.*

Proof. We argue by induction on $\dim_{\mathbb{F}} L$. Let A be the subalgebra of $\omega(L)$ generated by all u^p , where $u \in \omega(L)$. We observe that $A \cong \omega(L^p)$ as algebras. Thus there is an induced isomorphism $\omega(L^p) \cong \omega(H^p)$. Since $L \in \mathcal{F}_p$, we have $\dim_{\mathbb{F}} L^p < \dim_{\mathbb{F}} L$. Thus, by the induction hypothesis, there exists a restricted Lie algebra isomorphism $\varphi : L^p \cong H^p$. We now lift φ to an isomorphism of L and H . By Theorem 2.4, there exist generators $x_1, \dots, x_n \in L$ such that $L = \langle x_1 \rangle_p \oplus \dots \oplus \langle x_n \rangle_p$. Without loss of generality we assume

$$L^p = \langle x_1^p \rangle_p \oplus \dots \oplus \langle x_m^p \rangle_p \quad \text{for some } m \leq n.$$

Thus, $x_i^p = 0$ for every i in the range $m < i \leq n$. Note that $\dim L = n + \dim L^p$. So, as mentioned in Theorem 2.4, n is determined. Let $y_1, \dots, y_n \in H$ such that $H = \langle y_1 \rangle_p \oplus \dots \oplus \langle y_n \rangle_p$. Then $H^p = \langle y_1^p \rangle_p \oplus \dots \oplus \langle y_m^p \rangle_p$. So, we can assume that $\varphi(x_i^p) = y_i^p$ for every $1 \leq i \leq m$. We can verify that the map induced by $x_i \mapsto y_i$ for every $1 \leq i \leq n$ is a restricted Lie algebra isomorphism between L and H . \square

Corollary 2.6. *Let $L \in \mathcal{F}_p$ be a restricted Lie algebra over a perfect field. Then L/L'_p is determined.*

Proof. Note that $[u(L), u(L)]u(L) = L'_p u(L)$. Also, $u(L/L'_p) \cong u(L)/L'_p u(L)$. Hence, $u(L/L'_p)$ is determined. Since $L/L'_p \in \mathcal{F}_p$, it follows from Proposition 2.5 that L/L'_p is determined. \square

It turns out that stronger results hold over an algebraically closed field. Before we state the next result, we recall a well-known theorem; see [Jacobson 1962] or [Bahturin et al. 1992, Section 4.3]. Let $T_L = \langle x \in L \mid x^p = x \rangle_{\mathbb{F}}$, and denote by $\text{Rad}(L)$ the subalgebra of L spanned by all p -nilpotent elements.

Theorem 2.7. *Let L be a finite-dimensional abelian restricted Lie algebra over an algebraically closed field \mathbb{F} . Then $L = T_L \oplus \text{Rad}(L)$.*

Corollary 2.8. *Let L be a finite-dimensional abelian restricted Lie algebra over an algebraically closed field \mathbb{F} . If H be a restricted Lie algebra such that $u(L) \cong u(H)$, then $L \cong H$.*

Proof. Note that for every $k \geq 1$,

$$\dim_{\mathbb{F}} L/D_{p^k}(L) = \dim_{\mathbb{F}} L/D_p(L) + \dots + \dim_{\mathbb{F}} D_{p^{k-1}}(L)/D_{p^k}(L)$$

is determined. So $\dim_{\mathbb{F}} D_{p^k}(L)$ is determined for every $k \geq 1$. Let t be the least integer such that $\text{Rad}(L)^{p^t} = 0$. It follows that $D_{p^t}(L) = T_L$. Hence, $\dim_{\mathbb{F}} \text{Rad}(L) = \dim_{\mathbb{F}} \text{Rad}(H)$ by Theorem 2.7. Note that $L/T_L \cong \text{Rad}(L)$ as restricted Lie algebras. We claim that $\varphi(u(T_L)) = u(T_H)$. If so, then $\varphi(T_L u(L)) = T_H u(H)$. So

$$u(L/T_L) \cong u(L)/T_L u(L) \cong u(H)/T_H u(H) \cong u(H/T_H).$$

Thus, $u(\text{Rad}(L)) \cong u(\text{Rad}(H))$. Since $\text{Rad}(L), \text{Rad}(H) \in \mathcal{F}_p$, Proposition 2.5 then implies that there exists an isomorphism $\varphi : \text{Rad}(L) \rightarrow \text{Rad}(H)$. Clearly, φ can be extended to an isomorphism of L and H .

Now, we prove the claim. Let z_1, \dots, z_n be a basis of $\text{Rad}(H)$ and y_1, \dots, y_s be a basis of T_H , and assume that every y_i is less than every z_j . Let $x \in T_L$ and express $\varphi(x)$ in terms of PBW monomials in the y_i and z_j . So we have

$$\varphi(x) = u + \sum \alpha y_1^{a_1} \dots y_s^{a_s} z_1^{b_1} \dots z_n^{b_n},$$

where u is a linear combination of PBW monomials in the y_i only and each term in the sum has the property that $b_1 + \dots + b_n \neq 0$. Note that for a large k we have $\varphi(x)^{p^k} = u^{p^k} \in u(T_H)$. But $\varphi(x) = \varphi(x)^{p^k}$. So, $\varphi(x) \in u(T_H)$. Since $u(T_L)$ is generated by L and φ is an algebra homomorphism, we can get $\varphi(u(T_L)) \subseteq u(T_H)$. But $u(T_L)$ and $u(T_H)$ are finite-dimensional. So we get $\varphi(u(T_L)) = u(T_H)$. This proves the claim, completing the proof. \square

3. The quotient $L/L'^p + \gamma_3(L)$

We first record a couple of easy statements.

Lemma 3.1. *Let N be a restricted subalgebra of L . Then*

$$\omega(L)N + N\omega(L) = [N, L] + N\omega(L)$$

Lemma 3.2. *For every restricted subalgebra N of L , we have*

- $L \cap ([N, L] + N\omega(L)) = [N, L] + N^p$ and
- $Nu(L)/\omega(L)N + N\omega(L) \cong N/([N, L] + N^p)$.

Now write $J_L = \omega(L)L' + L'\omega(L) = \omega(L)L'_p + L'_p\omega(L)$. Since both $\omega(L)L'$ and $L'\omega(L)$ are determined, it follows that J_L is determined.

Corollary 3.3. *If $L \in \mathcal{F}_p$, then $\dim_{\mathbb{F}}(L/L^p + \gamma_3(L))$ is determined.*

Proof. Since $L'_p u(L)$ and J_L are determined, $\dim_{\mathbb{F}}(L'_p/L^p + \gamma_3(L))$ is determined, by Lemma 3.2. The result follows since L/L'_p is determined by Corollary 2.6. \square

From now on we assume that $L \in \mathcal{F}_p$ and \mathbb{F} is perfect. By Theorem 2.4, there exist $e_1, \dots, e_n \in L$ such that $L/L'_p = \langle e_1 + L'_p \rangle_p \oplus \dots \oplus \langle e_n + L'_p \rangle_p$. Let \bar{X} be a basis of L/L'_p consisting of $\bar{e}_i^{p^j}$, where $\bar{e}_i = e_i + L'_p$ and $1 \leq i \leq n$. Fix a set X of representatives of \bar{X} . So the elements of X are linearly independent modulo L'_p .

We define the *height* $v(x)$ of an element $x \in L$ to be the largest integer n such that $x \in D_n(L)$ if n exists and to be infinite otherwise. The *weight* of a PBW monomial $x_1^{a_1} \dots x_t^{a_t}$ is defined to be $\sum_{i=1}^t a_i v(x_i)$. We observe that $v(e_i^{p^j}) = p^j$ for every $1 \leq i \leq n$ and every $1 \leq j < \exp(\bar{e}_i)$. Indeed, if $e_i^{p^j} \in D_m(L)$ for some $m > p^j$, then

$$e_i^{p^j} = \sum_{k>j} a_k e_i^{p^k} \pmod{L'_p}.$$

It follows then that $e_i^{\hat{p}} \in L'_p$, where $\hat{p} = p^{\exp(\bar{e}_i)-1}$, which is a contradiction. Let Y be a linearly independent subset of L'_p such that $Z = X \cup Y$ is a basis of L and the set $\{z + D_{v(z)+1} \mid z \in Z\}$ is a basis of $\text{gr}(L)$. One way to construct such a subset Y is to take coset representatives of a basis for

$$\bigoplus_{i \geq 1} D_i(L) \cap (L'_p + \langle X \rangle_{\mathbb{F}}) / D_{i+1}(L).$$

We need the following variant of [Riley and Shalev 1995, Theorem 2.1].

Lemma 3.4. *Let $L \in \mathcal{F}_p$. Let \bar{Z} be a homogeneous basis of $\text{gr}(L)$ with a fixed set of representatives Z . Then the set of all PBW monomials in Z of weight at least k forms a basis for $\omega^k(L)$ for every $k \geq 1$.*

Note that J_L is linearly independent with the set of all PBW monomials in X . Let E denote the vector space spanned by J_L and all PBW monomials in X of degree at least two. The following lemma is easy to see, so we omit the proof.

Lemma 3.5. (1) $\omega(L) = L + E$.

(2) $(L + J_L) \cap E = J_L = E \cap L'_p u(L)$.

(3) $\omega(L)/J_L = L + J_L/J_L \oplus E/J_L$.

Lemma 3.6. *If $L \in \mathcal{F}_p$ then E/J_L is a central restricted Lie ideal of $\omega(L)/J_L$.*

Proof. The fact that E/J_L is a central Lie ideal of $\omega(L)/J_L$ easily follows from the identity $[ab, c] = a[b, c] + [a, c]b$, which holds in any associative algebra. So we have to prove that E/J_L is closed under the p -map. Since J_L is an associative ideal of $\omega(L)$, it is enough to prove that $u^p \in E$ for every PBW monomial u in E . Let $u = e_1^{a_1} \dots e_n^{a_n}$, where each a_i is in the range $0 \leq a_i < p^{\exp \bar{e}_i}$. It is not hard to see that $u^p = e_1^{pa_1} \dots e_n^{pa_n}$ modulo J_L . Since $L \in \mathcal{F}_p$, each \bar{e}_i is p -nilpotent. If

$pa_i < p^{\exp(\bar{e}_i)}$ for every $1 \leq i \leq n$, then u^p is a PBW monomial of degree at least two. Now suppose that $pa_i \geq p^{\exp(\bar{e}_i)}$ for some i . If $pa_i = p^{\exp(\bar{e}_i)}$, then a_i is a power of p . Since u has degree at least two, there exists $j \neq i$ such that $a_j \neq 0$. It now follows that $u^p \in J_L$. If $pa_i > p^{\exp(\bar{e}_i)}$ then $e_i^{pa_i} \in J_L$, and so $u^p \in E$. \square

Lemma 3.7. *We have $H \cap \varphi(E) \subseteq J_H$.*

Proof. We suppose $J_H = 0$ and prove that $H \cap \varphi(E) = 0$. Let $v \in H \cap \varphi(E) \subseteq \omega^2(H)$. Let $u \in E$ such that $\varphi(u) = v$. So, $u \in \omega^2(L)$. We prove by induction that $u \in \omega^n(L)$ for every n . But $\omega(L)$ is nilpotent by Lemma 2.1, and so $u = 0$. Supposing now by induction that $u \in \omega^n(L)$, we prove that $u \in \omega^{n+1}(L)$. So, $v \in H \cap \omega^n(H) = D_n(H)$. Thus, by Lemma 2.3, $u \in (D_n(L) + \omega^{n+1}(L)) \cap E$. But

$$(D_n(L) + \omega^{n+1}(L)) \cap E \subseteq \omega^{n+1}(L).$$

Indeed, let $u = \sum \alpha_i z_i + w$, where each $z_i \in Z$ has height n and $w \in \omega^{n+1}(L)$. By Lemma 3.4, w is a linear combination of PBW monomials in Z of weight at least $n + 1$. Since $u \in E$, it follows by the PBW Theorem that $\alpha_i = 0$ for every i . So $u = w \in \omega^{n+1}(L)$, as required. \square

Lemma 3.8. *We have $\omega(H)/J_H = H + J_H/J_H \oplus \varphi(E)/J_H$.*

Proof. By Lemma 3.7, it is enough to prove $\omega(H)/J_H \subseteq H + J_H/J_H \oplus \varphi(E)/J_H$. Note that both $\omega(H)/J_H$ and $\varphi(E)/J_H$ are determined. Since $\dim_{\mathbb{F}}(H + J_H/J_H) = \dim_{\mathbb{F}}(H/(H')^p + \gamma_3(H))$ is determined by Corollary 3.3, the result follows from Lemma 3.5. \square

Noting that $L + J_L/J_L \cong L/L^p + \gamma_3(L)$ by Lemma 3.2, we can now finish the proof of our main result.

Lemma 3.9. *The restriction of the natural isomorphism $\omega(L)/J_L \rightarrow \omega(H)/J_H$ to $L + J_L/J_L$ induces an isomorphism of $L + J_L/J_L$ and $H + J_H/J_H$.*

Proof. We denote by φ the induced isomorphism $\omega(L)/J_L \rightarrow \omega(H)/J_H$. Let $\varphi|_{L+J_L/J_L} = \varphi_1 + \varphi_2$ denote the restriction of φ to $L + J_L/J_L$, where φ_1 maps from $L + J_L/J_L$ to $H + J_H/J_H$. It is enough to prove φ_1 is a restricted Lie algebra isomorphism. Since E/L is a central Lie ideal of $\omega(L)/J_L$ by Lemma 3.6, $\varphi(E)/J_H$ is a central Lie ideal of $\omega(H)/J_H$. So, for every $x, z \in L$, we have

$$\varphi([x, z] + J_L) = [\varphi(x) + J_H, \varphi(z) + J_H] = [\varphi_1(x), \varphi_1(z)] + J_H.$$

So, φ_1 preserves the Lie brackets. Also,

$$\varphi(x^p + J_L) = \varphi(x)^p + J_H = (\varphi_1(x))^p + (\varphi_2(x))^p + J_H$$

Since $(\varphi_2(x))^p + J_H \in \varphi(E)/J_H$, it follows that φ_1 preserves the p -powers. Also, φ_1 is injective by Lemma 3.5. Since $L + J_L/J_L$ and $H + J_H/J_H$ have the same dimension by Corollary 3.3, it follows that φ_1 is an isomorphism. \square

Acknowledgments

I am grateful to the referee for careful reading of the paper and Luzius Grunenfelder for useful discussions.

References

- [Bagiński and Konovalov 2007] C. Bagiński and A. Konovalov, “The modular isomorphism problem for finite p -groups with a cyclic subgroup of index p^2 ”, pp. 186–193 in *Groups* (St. Andrews 2005), vol. 1, edited by C. M. Campbell et al., London Math. Soc. Lecture Note Ser. **339**, Cambridge Univ. Press, 2007. MR 2008f:20009 Zbl 1117.20003
- [Bahturin et al. 1992] Y. A. Bahturin, A. A. Mikhalev, V. M. Petrogradsky, and M. V. Zaicev, *Infinite-dimensional Lie superalgebras*, Expositions in Mathematics **7**, de Gruyter, Berlin, 1992. MR 94b:17001
- [Jacobson 1962] N. Jacobson, *Lie algebras*, Pure and Applied Mathematics **10**, Wiley, New York, 1962. MR 26 #1345 Zbl 0121.27504
- [Riley and Shalev 1995] D. M. Riley and A. Shalev, “Restricted Lie algebras and their envelopes”, *Canad. J. Math.* **47**:1 (1995), 146–164. MR 96e:16031 Zbl 0833.16024
- [Riley and Usefi 2007] D. Riley and H. Usefi, “The isomorphism problem for universal enveloping algebras of Lie algebras”, *Algebr. Represent. Theory* **10**:6 (2007), 517–532. MR 2008i:17015 Zbl 05265503
- [Ritter and Sehgal 1983] J. Ritter and S. Sehgal, “Isomorphism of group rings”, *Arch. Math. (Basel)* **40**:1 (1983), 32–39. MR 84k:16015 Zbl 0504.20003
- [Roggenkamp and Scott 1987] K. Roggenkamp and L. Scott, “Isomorphisms of p -adic group rings”, *Ann. of Math. (2)* **126**:3 (1987), 593–647. MR 89b:20021 Zbl 0633.20003
- [Sandling 1985] R. Sandling, “The isomorphism problem for group rings: a survey”, pp. 256–288 in *Orders and their applications* (Oberwolfach, 1984), edited by I. Reiner and K. W. Roggenkamp, Lecture Notes in Math. **1142**, Springer, Berlin, 1985. MR 87b:20007 Zbl 0565.20005
- [Sandling 1989] R. Sandling, “The modular group algebra of a central-elementary-by-abelian p -group”, *Arch. Math. (Basel)* **52**:1 (1989), 22–27. MR 90b:20007 Zbl 0632.16011
- [Sehgal 1978] S. K. Sehgal, *Topics in group rings*, Monographs and Textbooks in Pure and Applied Math. **50**, Marcel Dekker, New York, 1978. MR 80j:16001 Zbl 0411.16004
- [Usefi 2008] H. Usefi, “The restricted isomorphism problem for metacyclic restricted Lie algebras”, *Proc. Amer. Math. Soc.* **136**:12 (2008), 4125–4133. MR 2009f:17032 Zbl 1160.17013
- [Weiss 1988] A. Weiss, “Rigidity of p -adic p -torsion”, *Ann. of Math. (2)* **127**:2 (1988), 317–332. MR 89g:20010 Zbl 0647.20007

Received April 7, 2009. Revised December 17, 2009.

HAMID USEFI
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF BRITISH COLUMBIA
 1984 MATHEMATICS ROAD
 VANCOUVER, BC V6T 1Z2
 CANADA
 usefi@math.ubc.ca

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Paulo Ney de Souza, Production Manager

Silvio Levy, Senior Production Editor

Matthew Cargo, Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2010 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer-review and production is managed by EditFlow™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2010 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 246 No. 2 June 2010

Geometric structures associated to a contact metric (κ, μ) -space BENIAMINO CAPPELLETTI MONTANO and LUIGIA DI TERLIZZI	257
Multilinear singular operators with fractional rank CIPRIAN DEMETER, MALABIKA PRAMANIK and CHRISTOPH THIELE	293
A new proof of Reifenberg's topological disc theorem GUANGHAO HONG and LIHE WANG	325
Global classical solutions to hyperbolic geometric flow on Riemann surfaces FAGUI LIU and YUANZHANG ZHANG	333
An extension of the quintuple product identity and its applications ZHI-GUO LIU	345
A generalization of the Pontryagin–Hill theorems to projective modules over Prüfer domains JORGE MACÍAS-DÍAZ	391
Elliptic pseudodifferential equations and Sobolev spaces over p -adic fields J. J. RODRÍGUEZ-VEGA and W. A. ZÚÑIGA-GALINDO	407
Absolutely isolated singularities of holomorphic maps of \mathbb{C}^n tangent to the identity FENG RONG	421
Pullbacks of Eisenstein series from $\mathrm{GU}(3, 3)$ and critical L -values for $\mathrm{GSp}(4) \times \mathrm{GL}(2)$ ABHISHEK SAHA	435
Isomorphism invariants of restricted enveloping algebras HAMID USEFI	487
Spacelike S -Willmore spheres in Lorentzian space forms PENG WANG	495



0030-8730(201006)246:2;1-B