

*Pacific
Journal of
Mathematics*

A CASSON-LIN TYPE INVARIANT FOR LINKS

ERIC HARPER AND NIKOLAI SAVELIEV

Volume 248 No. 1

November 2010

A CASSON–LIN TYPE INVARIANT FOR LINKS

ERIC HARPER AND NIKOLAI SAVELIEV

In 1992, Xiao-Song Lin constructed an invariant $h(K)$ of knots $K \subset S^3$ via a signed count of conjugacy classes of irreducible $SU(2)$ representations of $\pi_1(S^3 - K)$ with trace-free meridians. Lin showed that $h(K)$ equals one half times the knot signature of K . Using methods similar to Lin's, we construct an invariant $h(L)$ of two-component links $L \subset S^3$. Our invariant is a signed count of conjugacy classes of projective $SU(2)$ representations of $\pi_1(S^3 - L)$ with a fixed 2-cocycle and corresponding nontrivial w_2 . We show that $h(L)$ is, up to a sign, the linking number of L .

1. Introduction

One of the characteristic features of the fundamental group of a closed 3-manifold is that its representation variety in a compact Lie group tends to be finite, in a properly understood sense. This has been a guiding principle for defining invariants of 3-manifolds ever since Casson defined his λ -invariant for integral homology 3-spheres via a signed count of the $SU(2)$ representations of the fundamental group, where signs were determined using Heegaard splittings.

Among numerous generalizations of Casson's construction, we will single out the invariant of knots in S^3 defined by Xiao-Song Lin [1992] via a signed count of $SU(2)$ representations of the fundamental group of the knot exterior. The latter is a 3-manifold with nonempty boundary so the finiteness principle above only applies after one imposes a proper boundary condition. Lin's choice of boundary condition, namely, that all of the knot meridians are represented by trace-free $SU(2)$ matrices, resulted in an invariant $h(K)$ of knots $K \subset S^3$. Lin further showed that $h(K)$ equals half the knot signature of K .

The signs in Lin's construction were determined using braid representations for knots. Austin (unpublished) and Heusener and Kroll [1998] extended Lin's construction by letting the meridians of the knot be represented by $SU(2)$ matrices with

MSC2000: primary 57M25; secondary 57M05.

Keywords: link group, braid, projective representation.

Saveliev was partially supported by NSF grant 0305946 and the Max-Planck-Institut für Mathematik in Bonn, Germany.

a fixed trace that need not be zero. Their construction gives a knot invariant that equals, for most choices of the trace, one half times the equivariant knot signature.

In this paper, we extend Lin's construction to two-component links L in S^3 . In essence, we replace the count of $SU(2)$ representations with a count of *projective* $SU(2)$ representations of $\pi_1(S^3 - L)$, in the sense of [Ruberman and Saveliev 2004], with a fixed 2-cocycle representing a nontrivial element in the second group cohomology of $\pi_1(S^3 - L)$. The resulting signed count is denoted by $h(L)$. The two main results of this paper are then as follows.

Theorem 1. *For any two-component link $L \subset S^3$, the integer $h(L)$ is a well-defined invariant of L .*

Theorem 2. *For any two-component link $L = \ell_1 \cup \ell_2$ in S^3 , one has*

$$h(L) = \pm \text{lk}(\ell_1, \ell_2).$$

Our choice of the 2-cocycle imposes Lin's trace-free condition on us. This is in contrast to Lin's construction, where the choice of boundary condition seemed somewhat arbitrary. This also means one should not expect to extend our construction to $SU(2)$ representations with nonzero trace boundary condition.

Shortly after Casson introduced his invariant for homology 3-spheres, Taubes [1990] gave a gauge theoretic description of it in terms of a signed count of flat $SU(2)$ connections. After Lin's work, but before Heusener and Kroll, a gauge theoretic interpretation of the Lin invariant was given by Herald [1997]. He used this interpretation to define an extension of the Lin invariant, now known as the Herald–Lin invariant, to knots in arbitrary homology spheres, with arbitrary fixed-trace (possibly nonzero) boundary condition.

Another attractive feature of the gauge theoretic approach is that it can be used to produce ramified versions of the above invariants. Floer [1988] introduced the instanton homology theory whose Euler characteristic is twice the Casson invariant. We expect that our invariant will have a similar interpretation, perhaps along the lines of the knot instanton homology theory of Kronheimer and Mrowka [2008], which in turn is a variant of the orbifold Floer homology of Collin and Steer [1999]. We hope to discuss this elsewhere, together with possible extensions to links in homology spheres and to links of more than two components.

2. Braids and representations

Let F_n be a free group of rank $n \geq 2$, with a fixed generating set x_1, \dots, x_n . We will follow the conventions of [Long 1989] and define the n -string braid group \mathcal{B}_n to be the subgroup of $\text{Aut}(F_n)$ generated by the automorphisms $\sigma_1, \dots, \sigma_{n-1}$,

where the action of σ_i is given by

$$\begin{aligned}\sigma_i : x_i &\mapsto x_{i+1}, \\ x_{i+1} &\mapsto (x_{i+1})^{-1}x_ix_{i+1}, \\ x_j &\mapsto x_j \quad \text{for } j \neq i, i+1.\end{aligned}$$

The natural homomorphism $\mathcal{B}_n \rightarrow S_n$, $\sigma \mapsto \bar{\sigma}$, onto the symmetric group on n letters maps each generator σ_i to the transposition $\bar{\sigma}_i = (i, i+1)$. A useful observation is that, for any $\sigma \in \text{Aut}(F_n)$, one has

$$(1) \quad \sigma(x_i) = wx_{\bar{\sigma}^{-1}(i)}w^{-1}$$

for some word $w \in F_n$. Also σ preserves the product $x_1 \cdots x_n$, that is,

$$(2) \quad \sigma(x_1 \cdots x_n) = x_1 \cdots x_n.$$

2a. SU(2) representations. Consider the Lie group $\text{SU}(2)$ of unitary two-by-two matrices with determinant one, that is, complex matrices

$$\begin{pmatrix} u & v \\ -\bar{v} & \bar{u} \end{pmatrix}$$

such that $u\bar{u} + v\bar{v} = 1$. We will often identify $\text{SU}(2)$ with the group $\text{Sp}(1)$ of unit quaternions via

$$\begin{pmatrix} u & v \\ -\bar{v} & \bar{u} \end{pmatrix} \mapsto u + vj \in \mathbb{H}.$$

Let $R_n = \text{Hom}(F_n, \text{SU}(2))$ be the space of $\text{SU}(2)$ representations of F_n , and identify it with $\text{SU}(2)^n$ by sending a representation $\alpha : F_n \rightarrow \text{SU}(2)$ to the vector $(\alpha(x_1), \dots, \alpha(x_n))$ of $\text{SU}(2)$ matrices. The above representation $\mathcal{B}_n \rightarrow \text{Aut}(F_n)$ then gives rise to the representation

$$(3) \quad \rho : \mathcal{B}_n \rightarrow \text{Diff}(R_n)$$

via $\rho(\sigma)(\alpha) = \alpha \circ \sigma^{-1}$. We will abbreviate $\rho(\sigma)$ to σ . We will also denote $X = (X_1, \dots, X_n) \in R_n$ and write $\sigma(X) = (\sigma(X)_1, \dots, \sigma(X)_n)$.

Example. For any $(X_1, \dots, X_n) \in R_n$, we have

$$\sigma_1(X_1, X_2, X_3, \dots, X_n) = (X_1X_2X_1^{-1}, X_1, X_3, \dots, X_n).$$

2b. Extension to the wreath product $\mathbb{Z}_2 \wr \mathcal{B}_n$. The wreath product $\mathbb{Z}_2 \wr \mathcal{B}_n$ is the semidirect product of \mathcal{B}_n with $(\mathbb{Z}_2)^n$, where \mathcal{B}_n acts on $(\mathbb{Z}_2)^n$ by permuting the coordinates, $\sigma(\varepsilon_1, \dots, \varepsilon_n) = (\varepsilon_{\bar{\sigma}(1)}, \dots, \varepsilon_{\bar{\sigma}(n)})$. Thus the elements of $\mathbb{Z}_2 \wr \mathcal{B}_n$ are the pairs $(\varepsilon, \sigma) \in (\mathbb{Z}_2)^n \times \mathcal{B}_n$, with the group multiplication law

$$(\varepsilon, \sigma) \cdot (\varepsilon', \sigma') = (\varepsilon\sigma(\varepsilon'), \sigma\sigma').$$

The representation (3) can be extended to a representation

$$(4) \quad \rho : \mathbb{Z}_2 \wr \mathcal{B}_n \rightarrow \text{Diff}(R_n)$$

by defining

$$\rho(\varepsilon, \sigma)(X) = \varepsilon \cdot \sigma(X) = (\varepsilon_1 \sigma(X)_1, \dots, \varepsilon_n \sigma(X)_n),$$

where the ε_i are viewed as elements of the center $\mathbb{Z}_2 = \{\pm 1\}$ of $\text{SU}(2)$. That (4) is a representation follows by a direct calculation after one observes that, by (1),

$$(5) \quad \sigma(X)_i = AX_{\bar{\sigma}(i)}A^{-1} \quad \text{for some } A \in \text{SU}(2).$$

Again, we will abuse notation and write simply $\varepsilon\sigma$ for both (ε, σ) and $\rho(\varepsilon, \sigma)$.

Example. For any $(X_1, \dots, X_n) \in R_n$ and $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) \in (\mathbb{Z}_2)^n$, we have

$$\begin{aligned} (\varepsilon\sigma_1)(X_1, X_2, X_3, \dots, X_n) &= (\varepsilon_1 X_1 X_2 X_1^{-1}, \varepsilon_2 X_1, \varepsilon_3 X_3, \dots, \varepsilon_n X_n), \\ \sigma_1(\varepsilon X) &= \sigma_1(\varepsilon)\sigma_1(X) = (\varepsilon_2 X_1 X_2 X_1^{-1}, \varepsilon_1 X_1, \varepsilon_3 X_3, \dots, \varepsilon_n X_n). \end{aligned}$$

2c. Braids and link groups. The closure $\hat{\sigma}$ of a braid $\sigma \in \mathcal{B}_n$ is a link in S^3 with link group

$$\pi_1(S^3 - \hat{\sigma}) = \langle x_1, \dots, x_n \mid x_i = \sigma(x_i) \text{ for } i = 1, \dots, n \rangle,$$

where each x_i represents a meridian of $\hat{\sigma}$. One can easily see that the fixed points of the diffeomorphism $\sigma : R_n \rightarrow R_n$ are representations $\pi_1(S^3 - \hat{\sigma}) \rightarrow \text{SU}(2)$. This paper grew out of the observation that a fixed point $\alpha = (\alpha(x_1), \dots, \alpha(x_n))$ of the map $\varepsilon\sigma : R_n \rightarrow R_n$ gives rise to a representation $\text{ad } \alpha : \pi_1(S^3 - \hat{\sigma}) \rightarrow \text{SO}(3)$ by composing with the adjoint representation $\text{ad} : \text{SU}(2) \rightarrow \text{SO}(3)$. Depending on ε , the representation $\text{ad } \alpha$ may or may not lift to an $\text{SU}(2)$ representation, the obstruction being the second Stiefel–Whitney class $w_2(\text{ad } \alpha) \in H^2(\pi_1(S^3 - \hat{\sigma}); \mathbb{Z}_2)$.

3. Definition of $h(\varepsilon\sigma)$

Every link in S^3 is the closure $\hat{\sigma}$ of a braid σ ; see [Alexander 1923]. Let σ be a braid whose closure $\hat{\sigma}$ has two components. We will associate with it, for a carefully chosen ε , an integer $h(\varepsilon\sigma)$. We will prove in Section 4 that h is an invariant of the link $\hat{\sigma}$.

3a. Choice of ε . The number of components of the link $\hat{\sigma}$ is exactly the number of cycles in the permutation $\bar{\sigma}$. We will be interested in two component links, that is, the closures of braids σ with

$$(6) \quad \bar{\sigma} = (i_1 \dots i_m)(i_{m+1} \dots i_n) \quad \text{for some } 1 \leq m \leq n-1.$$

Given such a braid σ , choose a vector $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) \in (\mathbb{Z}_2)^n$ such that

$$(7) \quad \varepsilon_{i_1} \cdots \varepsilon_{i_m} = \varepsilon_{i_{m+1}} \cdots \varepsilon_{i_n} = -1.$$

This choice of ε is dictated by the following two considerations. First, we wish to preserve condition (2) in the form

$$(8) \quad (\varepsilon\sigma)(X)_1 \cdots (\varepsilon\sigma)(X)_n = X_1 \cdots X_n,$$

and second, we want the fixed points α of the diffeomorphism $\varepsilon\sigma : R_n \rightarrow R_n$ to have nonzero $w_2(\text{ad } \alpha)$.

Lemma 3.1. *If α is a fixed point of $\varepsilon\sigma : R_n \rightarrow R_n$ with ε as in (7), then $w_2(\text{ad } \alpha) \neq 0$.*

Proof. The class $w_2(\text{ad } \alpha)$ is the obstruction to lifting $\text{ad } \alpha$ to an $\text{SU}(2)$ representation. Extend α arbitrarily to a function $\alpha : \pi_1(S^3 - \hat{\sigma}) \rightarrow \text{SU}(2)$ lifting $\text{ad } \alpha$. Then $w_2(\text{ad } \alpha)$ will vanish if and only if there is a function $\eta : \pi_1(S^3 - \hat{\sigma}) \rightarrow \mathbb{Z}_2 = \{\pm 1\}$ such that $\eta \cdot \alpha$ is a representation. Suppose such a function exists, and write $\eta(x_i) = \eta_i = \pm 1$. Also, assume without loss of generality that $\bar{\sigma} = (1 \dots m)(m + 1 \dots n)$. It follows from (5) that to satisfy the relations $X_i = (\varepsilon\sigma)(X)_i$ we must have $\eta_1 = \varepsilon_1 \eta_2 = \varepsilon_1 \varepsilon_2 \eta_3 = \cdots = \varepsilon_1 \cdots \varepsilon_m \eta_1 = -\eta_1$, a contradiction with $\eta_1 = \pm 1$. \square

This result for $w_2(\text{ad } \alpha)$ can be sharpened using an algebraic topology lemma.

Lemma 3.2. *Let $\hat{\sigma}$ be a link of two components. Then $H^2(\pi_1(S^3 - \hat{\sigma}); \mathbb{Z}_2)$ is equal to \mathbb{Z}_2 if $\hat{\sigma}$ is nonsplit and is zero otherwise.*

Proof. If $\hat{\sigma}$ is nonsplit, then $S^3 - \hat{\sigma}$ is a $K(\pi, 1)$ by the sphere theorem; hence $H^2(\pi_1(S^3 - \hat{\sigma}); \mathbb{Z}_2) = H^2(S^3 - \hat{\sigma}; \mathbb{Z}_2) = \mathbb{Z}_2$. If $\hat{\sigma}$ is split, then $K(\pi_1(S^3 - \hat{\sigma}), 1)$ has the homotopy type of a one-point union of two circles and the result again follows. \square

Corollary 3.3. *Let $\hat{\sigma}$ be a split link of two components, and let ε be chosen as in (7). Then the diffeomorphism $\varepsilon\sigma : R_n \rightarrow R_n$ has no fixed points.*

3b. The zero-trace condition. A naive way to define $h(\varepsilon\sigma)$ would be as the intersection number of the graph of $\varepsilon\sigma : R_n \rightarrow R_n$ with the diagonal in the product $R_n \times R_n$. One can observe though that, in addition to this intersection not being transversal, its points $(X, X) = (X_1, \dots, X_n, X_1, \dots, X_n)$ have the property that $\text{tr } X_1 = \cdots = \text{tr } X_n = 0$. This can be seen as follows.

Assume without loss of generality that $\bar{\sigma} = (1 \dots m)(m + 1 \dots n)$. Then the relations $X = \varepsilon\sigma(X)$ together with (5) imply that

$$\begin{aligned} X_1 &= \varepsilon_1 \sigma(X)_1 = \varepsilon_1 A_1 \cdot X_{\bar{\sigma}(1)} \cdot A_1^{-1} = \varepsilon_1 A_1 X_2 A_1^{-1} \\ &= \varepsilon_1 A_1 \cdot \varepsilon_2 \sigma(X)_2 \cdot A_1^{-1} = \varepsilon_1 \varepsilon_2 A_1 A_2 \cdot X_{\bar{\sigma}(2)} \cdot A_2^{-1} A_1^{-1} = \cdots \\ &= \varepsilon_1 \cdots \varepsilon_m (A_1 \cdots A_m) \cdot X_1 \cdot (A_1 \cdots A_m)^{-1}. \end{aligned}$$

Since trace is conjugation invariant and $\varepsilon_1 \cdots \varepsilon_m = -1$, we conclude that $\text{tr } X_1 = \cdots = \text{tr } X_m = 0$. Similarly, $\text{tr } X_{m+1} = \cdots = \text{tr } X_n = 0$.

Hence, in our definition we will restrict ourselves to the subset of R_n consisting of $X = (X_1, \dots, X_n)$ with $\text{tr } X_1 = \cdots = \text{tr } X_n = 0$. The nontransversality problem will be addressed below by factoring out the conjugation symmetry and lowering the dimension of the ambient manifold.

3c. The definition. The subset of $\text{SU}(2)$ consisting of the matrices with zero trace is a conjugacy class in $\text{SU}(2)$ diffeomorphic to S^2 . Define

$$Q_n = \{(X_1, \dots, X_n) \in R_n \mid \text{tr } X_i = 0\} \subset R_n,$$

so that Q_n is a manifold diffeomorphic to $(S^2)^n$. Also define

$$H_n = \{(X_1, \dots, X_n, Y_1, \dots, Y_n) \in Q_n \times Q_n \mid X_1 \cdots X_n = Y_1 \cdots Y_n\}.$$

This is no longer a manifold due to the presence of *reducibles*. We call a point $(X_1, \dots, X_n, Y_1, \dots, Y_n) \in Q_n \times Q_n$ reducible if all X_i and Y_j commute with each other or, equivalently, if there is a matrix $A \in \text{SU}(2)$ such that the $AX_i A^{-1}$ and $AY_j A^{-1}$ are diagonal matrices for $i = 1, \dots, n$. The subset $S_n \subset Q_n \times Q_n$ of reducibles is closed.

Lemma 3.4. $H_n^* = H_n - S_n$ is an open manifold of dimension $4n - 3$.

Proof. Consider the open manifold $(Q_n \times Q_n)^* = Q_n \times Q_n - S_n$ of dimension $4n$ and the map $f : (Q_n \times Q_n)^* \rightarrow \text{SU}(2)$ given by

$$(9) \quad f(X_1, \dots, X_n, Y_1, \dots, Y_n) = X_1 \cdots X_n Y_n^{-1} \cdots Y_1^{-1}.$$

According to [Lin 1992, Lemma 1.5], this map has $1 \in \text{SU}(2)$ as a regular value. Since $H_n^* = f^{-1}(1)$, the result follows. \square

Because of (5) and the fact that multiplication by $-1 \in \text{SU}(2)$ preserves the zero trace condition, the representation (4) gives rise to a representation

$$(10) \quad \rho : \mathbb{Z}_2 \wr \mathcal{B}_n \rightarrow \text{Diff}(Q_n).$$

Given $\varepsilon\sigma \in \mathbb{Z}_2 \wr \mathcal{B}_n$ such that (6) and (7) are satisfied, consider two submanifolds of $Q_n \times Q_n$: the graph $\Gamma_{\varepsilon\sigma} = \{(X, \varepsilon\sigma(X)) \mid X \in Q_n\}$ of $\varepsilon\sigma : Q_n \rightarrow Q_n$, and the diagonal $\Delta_n = \{(X, X) \mid X \in Q_n\}$. Note that both $\Gamma_{\varepsilon\sigma}$ and Δ_n are subsets of H_n ; this is obvious for Δ_n and follows from (8) for $\Gamma_{\varepsilon\sigma}$.

Proposition 3.5. *The intersection $\Gamma_{\varepsilon\sigma} \cap \Delta_n \subset H_n$ consists of irreducible representations.*

Proof. Assume without loss of generality that $\bar{\sigma} = (1 \dots m)(m+1 \dots n)$, and suppose that $(X, X) = (X_1, \dots, X_n, X_1, \dots, X_n) \in \Gamma_{\varepsilon\sigma} \cap \Delta_n$ is reducible. Then all of the X_i commute with each other, and in particular $\sigma(X) = (X_{\bar{\sigma}(1)}, \dots, X_{\bar{\sigma}(n)})$.

The equality $X = \varepsilon\sigma(X)$ then implies that $X_1 = \varepsilon_1 X_{\bar{\sigma}(1)} = \varepsilon_1 X_2 = \varepsilon_1 \varepsilon_2 X_{\bar{\sigma}(2)} = \dots = \varepsilon_1 \dots \varepsilon_m X_1 = -X_1$, which contradicts that $X_1 \in \text{SU}(2)$. \square

Let $\Gamma_{\varepsilon\sigma}^* = \Gamma_{\varepsilon\sigma} \cap H_n^*$ and $\Delta_n^* = \Delta_n \cap H_n^*$ be the irreducible parts of $\Gamma_{\varepsilon\sigma}$ and Δ_n , respectively. They are both open submanifolds of H_n^* of dimension $2n$.

Corollary 3.6. *The intersection $\Delta_n^* \cap \Gamma_{\varepsilon\sigma}^* \subset H_n^*$ is compact.*

Proof. Proposition 3.5 implies that $\Delta_n^* \cap \Gamma_{\varepsilon\sigma}^* = \Delta_n \cap \Gamma_{\varepsilon\sigma}$, and the latter intersection is obviously compact since it is the intersection of two compact subsets of H_n . \square

The group $\text{SO}(3) = \text{SU}(2)/\{\pm 1\}$ acts freely by conjugation on H_n^* , Δ_n^* , and $\Gamma_{\varepsilon\sigma}^*$. Denote the resulting quotient manifolds by

$$\hat{H}_n = H_n^*/\text{SO}(3), \quad \hat{\Delta}_n = \Delta_n^*/\text{SO}(3), \quad \text{and} \quad \hat{\Gamma}_{\varepsilon\sigma} = \Gamma_{\varepsilon\sigma}^*/\text{SO}(3).$$

The dimension of \hat{H}_n is $4n - 6$, and $\hat{\Delta}_n$ and $\hat{\Gamma}_{\varepsilon\sigma}$ are $(2n - 3)$ -dimensional submanifolds. Since the intersection $\hat{\Delta}_n \cap \hat{\Gamma}_{\varepsilon\sigma}$ is compact, one can isotope $\hat{\Gamma}_{\varepsilon\sigma}$ into a submanifold $\tilde{\Gamma}_{\varepsilon\sigma}$ using an isotopy with compact support so that $\hat{\Delta}_n \cap \tilde{\Gamma}_{\varepsilon\sigma}$ consists of finitely many points. Define

$$h(\varepsilon\sigma) = \#_{\hat{H}_n}(\hat{\Delta}_n \cap \tilde{\Gamma}_{\varepsilon\sigma})$$

as the algebraic intersection number, where the orientations of \hat{H}_n , $\hat{\Delta}_n$, and $\tilde{\Gamma}_{\varepsilon\sigma}$ are described in the next subsection. It is obvious that $h(\varepsilon\sigma)$ does not depend on the perturbation of $\hat{\Gamma}_{\varepsilon\sigma}$, so we will simply write

$$h(\varepsilon\sigma) = \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle_{\hat{H}_n}.$$

3d. Orientations. Choose an arbitrary orientation of the copy of $S^2 \subset \text{SU}(2)$ cut out by the trace zero condition, and endow $Q_n = (S^2)^n$ and $Q_n \times Q_n$ with product orientations. The diagonal Δ_n and the graph $\Gamma_{\varepsilon\sigma}$ are naturally diffeomorphic to Q_n via projection onto the first factor, and they are given the induced orientations. If we reverse the orientation of S^2 , then the orientation of Q_n is reversed if n is odd. Hence the orientations of both Δ_n and $\Gamma_{\varepsilon\sigma}$ are reversed if n is odd, while the orientation of $Q_n \times Q_n = (S^2)^{2n}$ is preserved regardless of the parity of n .

Orient $\text{SU}(2)$ by the standard basis $\{i, j, k\}$ in its Lie algebra $\mathfrak{su}(2)$, and orient $H_n^* = f^{-1}(1)$ by applying the base-fiber rule to the map (9). The adjoint action of $\text{SO}(3)$ on $S^2 \subset \text{SU}(2)$ is orientation preserving; hence the $\text{SO}(3)$ quotients \hat{H}_n , $\hat{\Delta}_n$, and $\hat{\Gamma}_{\varepsilon\sigma}$ are orientable. We orient them using the base-fiber rule. The discussion in the previous paragraph shows that reversing orientation on S^2 may reverse the orientations of $\hat{\Delta}_n$ and $\hat{\Gamma}_{\varepsilon\sigma}$ but that it does not affect the intersection number $\langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle_{\hat{H}_n}$.

4. The link invariant h

In this section, we will prove Theorem 1. This will be accomplished by proving that $h(\varepsilon\sigma)$ is independent first of ε and then of σ .

4a. Independence of ε . We will first show that, for a fixed σ whose closure $\hat{\sigma}$ is a link of two components, $h(\varepsilon\sigma)$ is independent of the choice of ε as long as ε satisfies (7).

Proposition 4.1. *Let ε and ε' be such that (7) is satisfied. Then $h(\varepsilon\sigma) = h(\varepsilon'\sigma)$.*

Proof. Assume without loss of generality that $\bar{\sigma} = (1 \dots m)(m + 1 \dots n)$ and let $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$ and $\varepsilon' = (\varepsilon'_1, \dots, \varepsilon'_n)$. Define $\delta = (\delta_1, \dots, \delta_n)$ as the vector in $(\mathbb{Z}_2)^n$ with coordinates

$$\delta_1 = 1 \quad \text{and} \quad \delta_{k+1} = \delta_k \varepsilon_k \varepsilon'_k \quad \text{for } k = 1, \dots, n - 1,$$

and define the involution $\tau : Q_n \rightarrow Q_n$ by the formula

$$\tau(X) = \delta X = (\delta_1 X_1, \delta_2 X_2, \dots, \delta_n X_n).$$

Recall that $Q_n = (S^2)^n$ so that τ is a diffeomorphism that restricts to each of the factors S^2 as either the identity or the antipodal map. In particular, τ need not be orientation preserving.

The map $\tau \times \tau : Q_n \times Q_n \rightarrow Q_n \times Q_n$ obviously preserves the irreducibility condition and commutes with the $\text{SO}(3)$ action. It gives rise to an orientation preserving automorphism of \hat{H}_n , which will again be called $\tau \times \tau$. It is clear that $(\tau \times \tau)(\hat{\Delta}_n) = \hat{\Delta}_n$. It is also true that $(\tau \times \tau)(\hat{\Gamma}_{\varepsilon\sigma}) = \hat{\Gamma}_{\varepsilon'\sigma}$, which can be seen as follows. Write a pair $(\delta X, \delta\varepsilon\sigma(X))$ whose conjugacy class belongs to $(\tau \times \tau)(\hat{\Gamma}_{\varepsilon\sigma})$ as

$$(\delta X, \delta\varepsilon\sigma(X)) = (\delta X, \delta\varepsilon\sigma(\delta\delta X)) = (\delta X, \delta\varepsilon\sigma(\delta)\sigma(\delta X))$$

using the multiplication law in the group $\mathbb{Z}_2 \wr \mathcal{B}_n$. The conjugacy class of this pair belongs to $\Gamma_{\varepsilon'\sigma}$ if and only if $\delta\varepsilon\sigma(\delta) = \varepsilon'$. That this condition holds can be verified directly from the definition of δ .

Recall that the orientations of $\hat{\Delta}_n$, $\hat{\Gamma}_{\varepsilon\sigma}$, and $\hat{\Gamma}_{\varepsilon'\sigma}$ are induced by the orientation of Q_n . Therefore, the maps $\tau \times \tau : \hat{\Delta}_n \rightarrow \hat{\Delta}_n$ and $\tau \times \tau : \hat{\Gamma}_{\varepsilon\sigma} \rightarrow \hat{\Gamma}_{\varepsilon'\sigma}$ are either both orientation preserving or both orientation reversing depending on whether $\tau : Q_n \rightarrow Q_n$ preserves or reverses orientation. Hence we have

$$\begin{aligned} h(\varepsilon\sigma) &= \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle_{\hat{H}_n} = \langle (\tau \times \tau)(\hat{\Delta}_n), (\tau \times \tau)(\hat{\Gamma}_{\varepsilon\sigma}) \rangle_{(\tau \times \tau)(\hat{H}_n)} \\ &= \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon'\sigma} \rangle_{\hat{H}_n} = h(\varepsilon'\sigma). \end{aligned} \quad \square$$

From now on, we will drop ε from the notation and simply write $h(\sigma)$ for $h(\varepsilon\sigma)$ assuming that a choice of ε satisfying (7) has been made.

4b. Independence of σ . In this section, we will show that $h(\sigma)$ only depends on the link $\hat{\sigma}$, not on a particular choice of braid σ , by verifying that h is preserved under Markov moves. We will follow the proof of [Lin 1992, Theorem 1.8], which goes through with little change once the right ε are chosen.

Recall that two braids $\alpha \in \mathcal{B}_n$ and $\beta \in \mathcal{B}_m$ have isotopic closures $\hat{\alpha}$ and $\hat{\beta}$ if and only if one braid can be obtained from the other by a finite sequence of Markov moves; see for instance [Birman 1974]. A type 1 Markov move replaces $\sigma \in \mathcal{B}_n$ by $\xi^{-1}\sigma\xi \in \mathcal{B}_n$ for any $\xi \in \mathcal{B}_n$. A type 2 Markov move means replacing $\sigma \in \mathcal{B}_n$ by $\sigma_n^{\pm 1}\sigma \in \mathcal{B}_{n+1}$, or the inverse of this operation.

Proposition 4.2. *The invariant $h(\sigma)$ is preserved by type 1 Markov moves.*

Proof. Let $\xi, \sigma \in \mathcal{B}_n$ and assume as usual that $\bar{\sigma} = (1 \dots m)(m + 1 \dots n)$. Then

$$\overline{\xi^{-1}\sigma\xi} = (\bar{\xi}(1) \dots \bar{\xi}(m))(\bar{\xi}(m + 1) \dots \bar{\xi}(n))$$

has the same cycle structure as $\bar{\sigma}$. To compute $h(\xi^{-1}\sigma\xi)$, we will make a choice of $\varepsilon \in (\mathbb{Z}_2)^n$ that satisfies condition (7) with respect to the braid $\xi^{-1}\sigma\xi$, that is, $\varepsilon_{\bar{\xi}(1)} \dots \varepsilon_{\bar{\xi}(m)} = \varepsilon_{\bar{\xi}(m+1)} \dots \varepsilon_{\bar{\xi}(n)} = -1$.

The braid ξ gives rise to the map $\xi : \mathcal{Q}_n \rightarrow \mathcal{Q}_n$. It acts by permutation and conjugation on the S^2 factors in \mathcal{Q}_n ; hence it is orientation preserving (we use the fact that S^2 is even-dimensional). It induces an orientation-preserving map $\xi \times \xi : \mathcal{Q}_n \times \mathcal{Q}_n \rightarrow \mathcal{Q}_n \times \mathcal{Q}_n$, which preserves the irreducibility condition and commutes with the $SO(3)$ action. Equation (2) then ensures that we have a well-defined orientation-preserving automorphism $\xi \times \xi : \hat{H}_n \rightarrow \hat{H}_n$.

That this automorphism preserves the diagonal, $(\xi \times \xi)(\hat{\Delta}_n) = \hat{\Delta}_n$, is obvious. Concerning the graphs, let $(X, \varepsilon\xi^{-1}\sigma\xi(X)) \in \hat{\Gamma}_{\varepsilon\xi^{-1}\sigma\xi}$; then

$$(\xi \times \xi)(X, \varepsilon\xi^{-1}\sigma\xi(X)) = (\xi(X), \xi(\varepsilon\xi^{-1}\sigma\xi(X))) = (\xi(X), \xi(\varepsilon)\sigma(\xi(X))) \in \hat{\Gamma}_{\xi(\varepsilon)\sigma}.$$

Therefore, $(\xi \times \xi)(\hat{\Gamma}_{\varepsilon\xi^{-1}\sigma\xi}) = \hat{\Gamma}_{\xi(\varepsilon)\sigma}$. Since $\xi : \mathcal{Q}_n \rightarrow \mathcal{Q}_n$ is orientation preserving, the identifications above of the diagonals and graphs via $\xi \times \xi$ are also orientation preserving.

Observe that $\xi(\varepsilon)_i = \varepsilon_{\bar{\xi}(i)}$. Hence $\xi(\varepsilon)$ satisfies (7) with respect to σ and thus can be used to compute $h(\sigma)$. The argument is completed by the calculation

$$\begin{aligned} h(\xi^{-1}\sigma\xi) &= \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\xi^{-1}\sigma\xi} \rangle_{\hat{H}_n} = \langle (\xi \times \xi)(\hat{\Delta}_n), (\xi \times \xi)(\hat{\Gamma}_{\varepsilon\xi^{-1}\sigma\xi}) \rangle_{(\xi \times \xi)(\hat{H}_n)} \\ &= \langle \hat{\Delta}_n, \hat{\Gamma}_{\xi(\varepsilon)\sigma} \rangle_{\hat{H}_n} = h(\sigma). \end{aligned} \quad \square$$

Proposition 4.3. *The invariant $h(\sigma)$ is preserved by type 2 Markov moves.*

Proof. Given $\sigma \in \mathcal{B}_n$ and ε satisfying (7), change σ to $\sigma_n \sigma \in \mathcal{B}_{n+1}$ and let $\varepsilon' = \sigma_n(\varepsilon, 1)$. If $X = (X_1, \dots, X_n)$ and $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$, then

$$\begin{aligned} (\sigma_n \sigma)(X, X_{n+1}) &= \sigma_n(\sigma(X), X_{n+1}) \\ &= (\sigma(X)_1, \dots, \sigma(X)_{n-1}, \sigma(X)_n X_{n+1} \sigma(X)_n^{-1}, \sigma(X)_n) \end{aligned}$$

and $\varepsilon' = (\varepsilon_1, \dots, \varepsilon_{n-1}, 1, \varepsilon_n)$. In particular, ε' satisfies (7) with respect to $\sigma_n \sigma$. Consider the embedding $g : \mathcal{Q}_n \times \mathcal{Q}_n \rightarrow \mathcal{Q}_{n+1} \times \mathcal{Q}_{n+1}$ given by

$$g(X_1, \dots, X_n, Y_1, \dots, Y_n) = (X_1, \dots, X_n, Y_n, Y_1, \dots, Y_n, Y_n).$$

One can easily see that $g(H_n) \subset H_{n+1}$ and that g commutes with the conjugation, thus giving rise to an embedding $\hat{g} : \hat{H}_n \rightarrow \hat{H}_{n+1}$. A straightforward calculation using the formulas above for $\sigma_n \sigma$ and ε' then shows that

$$\hat{g}(\hat{\Delta}_n) \subset \hat{\Delta}_{n+1}, \quad \hat{g}(\hat{\Gamma}_{\varepsilon\sigma}) \subset \hat{\Gamma}_{\varepsilon'\sigma_n\sigma}, \quad \text{and} \quad \hat{g}(\hat{\Delta}_n \cap \hat{\Gamma}_{\varepsilon\sigma}) = \hat{\Delta}_{n+1} \cap \hat{\Gamma}_{\varepsilon'\sigma_n\sigma}.$$

Now one can achieve all the necessary transversalities and match the orientations in exactly the same way as in the second half of the proof of [Lin 1992, Theorem 1.8]. This shows that $h(\sigma_n \sigma) = h(\sigma)$. The proof that $h(\sigma_n^{-1} \sigma) = h(\sigma)$ is similar. \square

5. The invariant $h(\sigma)$ as the linking number

In this section we will prove Theorem 2, that is, show that for any link $\hat{\sigma} = \ell_1 \cup \ell_2$ of two components, one has $h(\sigma) = \pm \text{lk}(\ell_1, \ell_2)$. Our strategy will be to show that the invariant $h(\sigma)$ and the linking number $\text{lk}(\ell_1, \ell_2)$ change according to the same rule as we change a crossing between two strands from two different components of $\hat{\sigma} = \ell_1 \cup \ell_2$ (the link $\hat{\sigma}$ will need to be oriented for that, although a particular choice of orientation will not matter). After changing finitely many such crossings, we will arrive at a split link, for which both the invariant $h(\sigma)$ and the linking number $\text{lk}(\ell_1, \ell_2)$ vanish; see Corollary 3.3. The change of crossing as above obviously changes the linking number by ± 1 . To calculate the effect of the crossing change on $h(\sigma)$, we will follow [Lin 1992] and reduce the problem to a calculation in the pillowcase \hat{H}_2 .

5a. The pillowcase. We begin with a geometric description of \hat{H}_2 as a pillowcase; compare with [Lin 1992, Lemma 1.2]. Remember that

$$H_2 = \{(X_1, X_2, Y_1, Y_2) \in \mathcal{Q}_2 \times \mathcal{Q}_2 \mid X_1 X_2 = Y_1 Y_2\}.$$

We will use the identification of $\text{SU}(2)$ with $\text{Sp}(1)$ when convenient. Since X_2 is trace free, we may assume that $X_2 = i$ after conjugation. Conjugating by $e^{i\varphi}$ will

not change X_2 but, for an appropriate choice of φ , will make X_1 into

$$X_1 = \begin{pmatrix} ir & u \\ -u & -ir \end{pmatrix},$$

where both r and u are real, and u is also nonnegative. Since $r^2 + u^2 = 1$, we can write $r = \cos\theta$ and $u = \sin\theta$ for a unique θ such that $0 \leq \theta \leq \pi$. In quaternionic language, $X_1 = ie^{-k\theta}$ with $0 \leq \theta \leq \pi$. Similarly, the condition $\text{tr}(Y_2) = \text{tr}(Y_1^{-1}X_1X_2) = 0$ implies that $Y_1 = ie^{-k\psi}$, this time with $-\pi \leq \psi \leq \pi$. To summarize,

$$X_1 = ie^{-k\theta}, \quad X_2 = i, \quad Y_1 = ie^{-k\psi}, \quad Y_2 = ie^{-k(\psi-\theta)}.$$

Thus \hat{H}_2 is parametrized by the rectangle $[0, \pi] \times [-\pi, \pi]$, with proper identifications along the edges and with the reducibles removed. The reducibles occur when both θ and ψ are multiples of π , and hence \hat{H}_2 is a 2-sphere with the points $A = (0, 0)$, $B = (\pi, 0)$, $A' = (0, \pi)$, and $B' = (\pi, \pi)$ removed; see Figure 1. According to [Lin 1992], the orientation on the front sheet of \hat{H}_2 coincides with the standard orientation on the (θ, ψ) plane.

Example. Let $\sigma = \sigma_1^2$, so that $\hat{\sigma} = \ell_1 \cup \ell_2$ is the Hopf link with $\text{lk}(\ell_1, \ell_2) = \pm 1$. To calculate $h(\sigma)$, we let $\varepsilon = (-1, -1)$, the only available choice satisfying (7), and consider the submanifolds $\hat{\Delta}_2$ and $\hat{\Gamma}_{\varepsilon\sigma}$ of \hat{H}_2 . We have, in quaternionic notation, $\hat{\Delta}_2 = \{(ie^{-k\theta}, i, ie^{-k\theta}, i)\}$, which is the diagonal $\psi = \theta$ in the pillowcase. A straightforward calculation shows that $\hat{\Gamma}_{\varepsilon\sigma} \subset \hat{H}_2$ is given by $\psi = 3\theta - \pi$. As can be seen in Figure 1, the intersection $\hat{\Delta}_2 \cap \hat{\Gamma}_{\varepsilon\sigma}$ consists of one point coming with a sign. Hence $h(\sigma_1^2) = \pm 1$, which is consistent with the fact that $\text{lk}(\ell_1, \ell_2) = \pm 1$.




Figure 1. The pillowcase.

Example. Let $\sigma = \sigma_1^{2n}$. Then arguing as above one can show that $\hat{\Gamma}_{\varepsilon\sigma} \subset \hat{H}_2$ is given by $\psi = (2n+1)\theta - \pi$. In this case there are n intersection points all of which come with the same sign. This shows that $h(\sigma_1^{2n}) = \pm n$, which is again consistent with the fact that $\text{lk}(\ell_1, \ell_2) = \pm n$.

5b. The difference cycle. Fix an orientation on a given two component link $\hat{\sigma}$. A particular choice of orientation will not matter because we are only interested in identifying $h(\sigma)$ with the linking number $\text{lk}(\ell_1, \ell_2)$ up to sign. We wish to change one of the crossings between the two components of $\hat{\sigma}$. Using a sequence of first Markov moves, we may assume that the first two strands of σ belong to two different components of $\hat{\sigma}$, and that the crossing change occurs between these two strands. Furthermore, we may assume that the crossing change makes σ into $\sigma_1^{\pm 2}\sigma$, where the sign depends on the type of the crossing we change. Note that the braid $\sigma_1^{\pm 2}\sigma$ has the same permutation type as σ ; in particular, its closure is a link of two components. In fact, if we let $\sigma' = \sigma_1^{-2}\sigma$, then

$$h(\sigma_1^{-2}\sigma) - h(\sigma) = h(\sigma') - h(\sigma_1^2\sigma') = -(h(\sigma_1^2\sigma') - h(\sigma'));$$

hence we only need to understand the difference $h(\sigma_1^2\sigma) - h(\sigma)$. Let us fix $\varepsilon = (-1, -1, 1, \dots, 1)$. Since σ_1^2 and ε commute, we have

$$\begin{aligned} h(\sigma_1^2\sigma) - h(\sigma) &= \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma_1^2\sigma} \rangle - \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle \\ &= \langle \hat{\Gamma}_{\sigma_1^{-2}}, \hat{\Gamma}_{\varepsilon\sigma} \rangle - \langle \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle = \langle \hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle, \end{aligned}$$

where all intersection numbers are taken in \hat{H}_n . This leads us to consider the *difference cycle* $\hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n$ that is carried by \hat{H}_n . The next step in our argument will be to reduce the analysis of the intersection above to an intersection theory in the pillowcase \hat{H}_2 .

5c. The pillowcase reduction. We consider the subset $V_n \subset H_n$ consisting of all $(X_1, \dots, X_n, Y_1, \dots, Y_n) \in H_n$ such that $X_k = Y_k$ for $k = 3, \dots, n$. Equivalently, V_n consists of all $(X_1, \dots, X_n, Y_1, \dots, Y_n) \in Q_n \times Q_n$ such that $(X_1, X_2, Y_1, Y_2) \in H_2$ and $X_k = Y_k$ for all $k = 3, \dots, n$. Therefore, V_n can be identified as

$$V_n = H_2 \times \Delta_{n-2} \subset (Q_2 \times Q_2) \times (Q_{n-2} \times Q_{n-2}).$$

Lemma 5.1. *The quotient $\hat{V}_n = (H_2^* \times \Delta_{n-2})/\text{SO}(3)$ is a submanifold of \hat{H}_n of dimension $2n - 2$.*

Proof. Since H_2^* and Δ_{n-2} are smooth manifolds of dimensions 5 and $2n - 4$, respectively, and their product contains no reducibles, the statement follows. \square

Lemma 5.2. *The difference cycle $\hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n$ is carried by \hat{V}_n .*

Proof. Observe that neither $\hat{\Gamma}_{\sigma_1^{-2}}$ nor $\hat{\Delta}_n$ are subsets of \hat{V}_n . However, their portions that do not fit in \hat{V}_n ,

$$\hat{\Gamma}_{\sigma_1^{-2}} - (\hat{\Gamma}_{\sigma_1^{-2}} \cap \hat{V}_n) \quad \text{and} \quad \hat{\Delta}_n - (\hat{\Delta}_n \cap \hat{V}_n),$$

are exactly the same. Namely, they consist of the equivalence classes of $2n$ -tuples $(X_1, \dots, X_n, X_1, \dots, X_n)$ such that X_1 commutes with X_2 . These cancel in the difference cycle $\hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n$, thus making it belong to \hat{V}_n . \square

One can isotope $\hat{\Gamma}_{\varepsilon\sigma}$ into $\tilde{\Gamma}_{\varepsilon\sigma}$ using an isotopy with compact support so that $\tilde{\Gamma}_{\varepsilon\sigma}$ is transverse to $\hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n$. The latter means precisely that $\tilde{\Gamma}_{\varepsilon\sigma}$ stays away from $(S_2 \times \Delta_{n-2})/\text{SO}(3)$ and is transverse to both $\hat{\Gamma}_{\sigma_1^{-2}}$ and $\hat{\Delta}_n$; a precise argument can be found in [Heusener and Kroll 1998, page 491]. We further extend this isotopy to make $\tilde{\Gamma}_{\varepsilon\sigma}$ transverse to \hat{V}_n so that their intersection is a naturally oriented 1-dimensional submanifold of \hat{H}_n .

The natural projection $p : V_n \rightarrow H_2$ induces a map $\hat{p} : \hat{V}_n \rightarrow \hat{H}_2$. Use a further small compactly supported isotopy of $\tilde{\Gamma}_{\varepsilon\sigma}$, if necessary, to make $\hat{p}(\hat{V}_n \cap \tilde{\Gamma}_{\varepsilon\sigma})$ into a 1-submanifold of \hat{H}_2 . The proofs of [Lin 1992, Lemmas 2.2 and 2.3] then go through with little change to give us the identity

$$\langle \hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n, \hat{\Gamma}_{\varepsilon\sigma} \rangle_{\hat{H}_n} = \langle \hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_n, \hat{p}(\hat{V}_n \cap \tilde{\Gamma}_{\varepsilon\sigma}) \rangle_{\hat{H}_2}.$$

5d. Computation in the pillowcase. We first study the behavior of $\hat{p}(\hat{V}_n \cap \hat{\Gamma}_{\varepsilon\sigma})$ near the corners of \hat{H}_2 .

Proposition 5.3. *There is a neighborhood around A' in the pillowcase \hat{H}_2 inside which $\hat{p}(\hat{V}_n \cap \hat{\Gamma}_{\varepsilon\sigma})$ is a curve approaching A' .*

Proof. Let us consider the submanifold

$$\Delta'_n = \{(X_1, X_2, X_3, \dots, X_n; Y_1, Y_2, X_3, \dots, X_n)\} \subset Q_n \times Q_n$$

and observe that $V_n \cap \Gamma_{\varepsilon\sigma} = \Delta'_n \cap \Gamma_{\varepsilon\sigma}$. We will show that the intersection of Δ'_n with $\Gamma_{\varepsilon\sigma}$ is transversal at $(i, \varepsilon i) = (i, \dots, i; -i, -i, i, \dots, i)$. This will imply that $\Delta'_n \cap \Gamma_{\varepsilon\sigma}$ is a submanifold of dimension four in a neighborhood of $(i, \varepsilon i)$ and, after factoring out the $\text{SO}(3)$ symmetry, that $\hat{p}(\hat{V}_n \cap \hat{\Gamma}_{\varepsilon\sigma})$ is a curve approaching $A' = p(i, \varepsilon i)$.

Since $\dim \Delta'_n = 2n+4$, the dimension of $T_{(i, \varepsilon i)}(\Delta'_n \cap \Gamma_{\varepsilon\sigma}) = T_{(i, \varepsilon i)}\Delta'_n \cap T_{(i, \varepsilon i)}\Gamma_{\varepsilon\sigma}$ is at least four. Therefore, checking the transversality amounts to showing that this dimension is exactly four. Write

$$\begin{aligned} T_{(i, \varepsilon i)}(\Delta'_n) &= \{(u_1, \dots, u_n; v_1, v_2, u_3, \dots, u_n)\} \subset T_{(i, \varepsilon i)}(Q_n \times Q_n), \\ T_{(i, \varepsilon i)}(\Gamma_{\varepsilon\sigma}) &= \{(u_1, \dots, u_n; d_i(\varepsilon\sigma)(u_1, \dots, u_n))\} \subset T_{(i, \varepsilon i)}(Q_n \times Q_n). \end{aligned}$$

Then $T_{(i,\varepsilon i)}(\Delta'_n) \cap T_{(i,\varepsilon i)}(\Gamma_{\varepsilon\sigma})$ consists of the vectors $(u_1, \dots, u_n) \in T_i Q_n = T_i S^2 \oplus \dots \oplus T_i S^2$ that solve the matrix equation

$$(11) \quad [d_i(\sigma)] \begin{bmatrix} u_1 \\ u_2 \\ u_3 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} * \\ * \\ u_3 \\ \vdots \\ u_n \end{bmatrix};$$

since $\varepsilon = (-1, -1, 1, \dots, 1)$, we can safely replace $[d_i(\varepsilon\sigma)]$ by $[d_i(\sigma)]$. It is shown in [Long 1989] that $[d_i(\sigma)]$ is the Burau matrix of σ with parameter equal to -1 . It is a real matrix acting on $T_i Q_n = \mathbb{C}^n$; hence all we need to show is that the space of $(u_1, \dots, u_n) \in \mathbb{R}^n$ solving (11) has real dimension two. Let us write

$$[d_i(\sigma)] = \begin{bmatrix} A & B \\ C & D \end{bmatrix},$$

where A is a 2×2 matrix and D is an $(n-2) \times (n-2)$ matrix. Equation (11) is equivalent to

$$[C] \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = [1 - D] \begin{bmatrix} u_3 \\ \vdots \\ u_n \end{bmatrix},$$

so the proposition will follow as soon as we show that $1 - D$ is invertible. The invertibility of $1 - D$ is a consequence of the following two lemmas. \square

Lemma 5.4. *Let $\sigma \in \mathcal{B}_n$. Then the Burau matrix of σ with parameter -1 and the permutation matrix of $\bar{\sigma}^{-1}$ are the same modulo 2.*

Proof. According to [Birman 1974], the Burau matrix of σ with parameter t is the matrix

$$\left. \frac{\partial \sigma(x_i)}{\partial x_j} \right|_{x_i=t},$$

where the x_i are generators of the free group and ∂ is the derivative in the Fox free differential calculus; see [Fox 1962]. Applying the Fox calculus we obtain

$$\begin{aligned} \frac{\partial \sigma(x_i)}{\partial x_j} &= \frac{\partial (w x_{\bar{\sigma}^{-1}(i)} w^{-1})}{\partial x_j} = \frac{\partial w}{\partial x_j} + w \left(\frac{\partial (x_{\bar{\sigma}^{-1}(i)} w^{-1})}{\partial x_j} \right) \\ &= \frac{\partial w}{\partial x_j} + w \left(\frac{\partial x_{\bar{\sigma}^{-1}(i)}}{\partial x_j} + x_{\bar{\sigma}^{-1}(i)} \frac{\partial w^{-1}}{\partial x_j} \right) \\ &= \frac{\partial w}{\partial x_j} + w \frac{\partial x_{\bar{\sigma}^{-1}(i)}}{\partial x_j} - w x_{\bar{\sigma}^{-1}(i)} w^{-1} \frac{\partial w}{\partial x_j}, \end{aligned}$$

where w is a word in the x_i . After evaluating at $t = -1$ and reducing modulo 2, the above becomes simply $\partial x_{\bar{\sigma}^{-1}(i)}/\partial x_j$, which is the permutation matrix of $\bar{\sigma}^{-1}$. \square

Lemma 5.5. *Let $\sigma \in \mathcal{B}_n$ be such that $\hat{\sigma}$ is a two component link. Then $1 - D$ is invertible.*

Proof. Our assumption in this section has been that $\bar{\sigma} = (1, \dots)(2, \dots)$. We may further assume that

$$\bar{\sigma} = (1, 3, 4, \dots, k)(2, k + 1, k + 2, \dots, n)$$

by applying a sequence of first Markov moves fixing the first two strands of σ . The matrix $D \pmod{2}$ is obtained by crossing out the first two rows and first two columns in the permutation matrix of $\bar{\sigma}$; see Lemma 5.4. This description implies that $D \pmod{2}$ is upper diagonal, and hence so is $(1 - D) \pmod{2}$. The diagonal elements of the latter matrix are all equal to one; therefore, $\det(1 - D) = 1 \pmod{2}$ so $1 - D$ is invertible. \square

Remark 5.6. The orientation of the component of $\hat{p}(\hat{V}_n \cap \hat{\Gamma}_{\varepsilon\sigma})$ limiting to A' can be read off its description near A' given in the proof of Proposition 5.3. In particular, this orientation is independent of the choice of σ .

Proposition 5.7. *There are neighborhoods around A and B' in the pillowcase \hat{H}_2 that are disjoint from $\hat{p}(\hat{V}_n \cap \hat{\Gamma}_{\varepsilon\sigma})$.*

Proof. The arguments for A and B' are essentially the same so we will only give the proof for A . Assuming the contrary we have a curve in $\hat{V}_n \cap \hat{\Gamma}_{\varepsilon\sigma}$ limiting to a reducible representation in $V_n \cap \Gamma_{\varepsilon\sigma}$. After conjugating if necessary, this representation must have the form

$$(i, i, e^{i\varphi_3}, \dots, e^{i\varphi_n}, i, i, e^{i\varphi_3}, \dots, e^{i\varphi_n}).$$

Using that $\varepsilon = (-1, -1, 1, \dots, 1)$ and arguing as in the proof of Proposition 3.5, we arrive at the contradiction $i = -i$. \square

Proof of Theorem 2. According to Proposition 5.3, the 1-submanifold $\hat{p}(\hat{V}_n \cap \tilde{\Gamma}_{\varepsilon\sigma})$, near A' , is a curve approaching A' . According to Proposition 5.7, the other end of this curve must approach B . Therefore

$$h(\sigma_1^2\sigma) - h(\sigma) = \langle \hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_2, \hat{p}(\hat{V}_n \cap \tilde{\Gamma}_{\varepsilon\sigma}) \rangle_{\hat{H}_2}$$

is the same as the intersection number of an arc going from A' to B with the difference cycle $\hat{\Gamma}_{\sigma_1^{-2}} - \hat{\Delta}_2$. This number is either 1 or -1 but is the same for all σ ; see Remark 5.6. This is sufficient to prove that $h(\sigma)$ is the linking number up to an overall sign. \square

References

- [Alexander 1923] J. Alexander, “A lemma on systems of knotted curves”, *Proc. Nath. Acad. Sci.* **9** (1923), 93–95. JFM 49.0408.03
- [Birman 1974] J. S. Birman, *Braids, links, and mapping class groups*, Annals of Mathematics Studies **82**, Princeton University Press, 1974. MR 51 #11477 Zbl 0305.57013
- [Collin and Steer 1999] O. Collin and B. Steer, “Instanton Floer homology for knots via 3-orbifolds”, *J. Differential Geom.* **51**:1 (1999), 149–202. MR 2000j:57069 Zbl 1025.57034
- [Floer 1988] A. Floer, “An instanton-invariant for 3-manifolds”, *Comm. Math. Phys.* **118**:2 (1988), 215–240. MR 89k:57028 Zbl 0684.53027
- [Fox 1962] R. H. Fox, “A quick trip through knot theory”, pp. 120–167 in *Topology of 3-manifolds and related topics* (Athens, GA, 1961), edited by M. K. Fort, Jr., Prentice-Hall, Englewood Cliffs, NJ, 1962. MR 25 #3522
- [Herald 1997] C. M. Herald, “Flat connections, the Alexander invariant, and Casson’s invariant”, *Comm. Anal. Geom.* **5**:1 (1997), 93–120. MR 98i:57062 Zbl 0893.57003
- [Heusener and Kroll 1998] M. Heusener and J. Kroll, “Deforming abelian $SU(2)$ -representations of knot groups”, *Comment. Math. Helv.* **73**:3 (1998), 480–498. MR 99g:57008 Zbl 0910.57004
- [Kronheimer and Mrowka 2008] P. Kronheimer and T. Mrowka, “Knot homology groups from instantons”, preprint, version 2, 2008. arXiv 0806.1053v2
- [Lin 1992] X.-S. Lin, “A knot invariant via representation spaces”, *J. Differential Geom.* **35**:2 (1992), 337–357. MR 93e:57012 Zbl 0774.57007
- [Long 1989] D. D. Long, “On the linear representation of braid groups”, *Trans. Amer. Math. Soc.* **311**:2 (1989), 535–560. MR 89h:20054 Zbl 0714.20019
- [Ruberman and Saveliev 2004] D. Ruberman and N. Saveliev, “Rohlin’s invariant and gauge theory, I: Homology 3-tori”, *Comment. Math. Helv.* **79**:3 (2004), 618–646. MR 2006e:57020
- [Taubes 1990] C. H. Taubes, “Casson’s invariant and gauge theory”, *J. Differential Geom.* **31**:2 (1990), 547–599. MR 91m:57025 Zbl 0702.53017

Received July 20, 2009.

ERIC HARPER
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF MIAMI
 PO BOX 249085
 CORAL GABLES, FL 33124
 UNITED STATES
 harper@math.miami.edu

NIKOLAI SAVELIEV
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF MIAMI
 PO BOX 249085
 CORAL GABLES, FL 33124
 UNITED STATES
 saveliev@math.miami.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2010 is US \$420/year for the electronic version, and \$485/year for print and electronic.

Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2010 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 248 No. 1 November 2010

An existence theorem of conformal scalar-flat metrics on manifolds with boundary	1
SÉRGIO DE MOURA ALMARAZ	
Parasurface groups	23
KHALID BOU-RABEE	
Expressions for Catalan Kronecker products	31
ANDREW A. H. BROWN, STEPHANIE VAN WILLIGENBURG and MIKE ZABROCKI	
Metric properties of higher-dimensional Thompson's groups	49
JOSÉ BURILLO and SEAN CLEARY	
Solitary waves for the Hartree equation with a slowly varying potential	63
KIRIL DATCHEV and IVAN VENTURA	
Uniquely presented finitely generated commutative monoids	91
PEDRO A. GARCÍA-SÁNCHEZ and IGNACIO OJEDA	
The unitary dual of p -adic $\widetilde{\mathrm{Sp}}(2)$	107
MARCELA HANZER and IVAN MATIĆ	
A Casson–Lin type invariant for links	139
ERIC HARPER and NIKOLAI SAVELIEV	
Semiquandles and flat virtual knots	155
ALLISON HENRICH and SAM NELSON	
Infinitesimal rigidity of polyhedra with vertices in convex position	171
IVAN IZMESTIEV and JEAN-MARC SCHLENKER	
Robust four-manifolds and robust embeddings	191
VYACHESLAV S. KRUSHKAL	
On sections of genus two Lefschetz fibrations	203
SINEM ÇELİK ONARAN	
Biharmonic hypersurfaces in Riemannian manifolds	217
YE-LIN OU	
Singular fibers and 4-dimensional cobordism group	233
OSAMU SAEKI	



0030-8730(201011)248:1;1-C