

*Pacific  
Journal of  
Mathematics*

**THE BRAID GROUP SURJECTS ONTO  $G_2$  TENSOR SPACE**

SCOTT MORRISON

Volume 249 No. 1

January 2011

## THE BRAID GROUP SURJECTS ONTO $G_2$ TENSOR SPACE

SCOTT MORRISON

**Let  $V$  be the 7-dimensional irreducible representation of the quantum group  $U_q(\mathfrak{g}_2)$ . For each  $n$ , there is a map from the braid group  $\mathcal{B}_n$  to the endomorphism algebra of the  $n$ -th tensor power of  $V$ , given by  $\mathcal{R}$  matrices. Extending linearly to the braid group algebra, we get a map**

$$\mathcal{A}\mathcal{B}_n \rightarrow \mathbf{End}_{U_q(\mathfrak{g}_2)}(V^{\otimes n}).$$

**Lehrer and Zhang have proved that map is surjective, as a special case of a more general result.**

**Using Kuperberg's spider for  $G_2$ , we give an elementary diagrammatic proof of this result.**

### 1. Kuperberg's spider for $G_2$

We recall just enough from [Kuperberg 1994; 1996] for our purposes.

We fix the ground ring  $\mathcal{A} = \mathbb{C}(q)$ . Kuperberg's  $q$  is  $q^2$  here, which brings our conventions into agreement with those for quantum groups as presented in [Chari and Pressley 1995; Jantzen 1996; Sawin 2006].

First consider the braided tensor category  $\mathbf{Rep}'(G_2)$ , with objects tensor powers of the 7-dimensional representation of  $U_q(\mathfrak{g}_2)$  and morphisms linear maps commuting with the actions of  $U_q(\mathfrak{g}_2)$ . (The prime in the notation indicates this is just a full subcategory of the actual representation category; we don't allow arbitrary representations, although every representation does appear as a subobject of some tensor power.)

Second consider the category  $\mathcal{T}(G_2)$ , with objects natural numbers and morphisms planar trivalent graphs embedded in a rectangle, modulo certain relations. A morphism from  $n$  to  $m$  should be a graph with  $n$  boundary points along the bottom edge of the rectangle, and  $m$  boundary points along the top edge. Note that we allow embedded circles in the interior of the rectangle, called *loops*. The edges of the graph break the rectangle into faces; those that do not meet the boundary of the rectangle are called *internal* faces. Composition of morphisms is vertical stacking. The category becomes a tensor category by adding natural numbers at

---

*MSC2000:* primary 17B37; secondary 20G42, 17B10, 18D10.

*Keywords:* braid group, spider,  $G_2$ , tensor category, representation theory.

$$\begin{aligned}
 \bigcirc &= q^{10} + q^8 + q^2 + 1 + q^{-2} + q^{-8} + q^{-10} \\
 \bigcirc &= 0 \\
 \bigcirc &= -(q^6 + q^4 + q^2 + q^{-2} + q^{-4} + q^{-6}) \quad | \\
 \triangle &= (q^4 + 1 + q^{-4}) \quad \text{Y} \\
 \square &= -(q^2 + q^{-2}) (\text{Y} + \text{X}) + (q^2 + 1 + q^{-2}) (\text{C} + \text{I}) \\
 \text{pentagon} &= \text{A} + \text{B} + \text{C} + \text{D} + \text{E} \\
 &\quad - \text{F} - \text{G} - \text{H} - \text{I} - \text{J}
 \end{aligned}$$

**Figure 1.** Kuperberg’s [1994; 1996] relations for the  $G_2$  spider, omitting relations involving double edges that we don’t use. Note also that there is a sign error in [1996] but not in [1994].

the level of objects, and juxtaposing graphs side by side at the level of morphisms. The relations from [Kuperberg 1996] are shown in Figure 1; the category  $\mathcal{T}(G_2)$  becomes a braided tensor category with Kuperberg’s formulas for a crossing:

$$(1) \quad \text{X} = \frac{1}{1+q^{-2}} \text{Y} + \frac{1}{1+q^2} \text{X} + \frac{1}{q^2+q^4} \text{C} + \frac{1}{q^{-2}+q^{-4}} \text{I}$$

We now need two theorems from [Kuperberg 1996], one easy, one hard.

**Theorem 1.1** (web bases for  $\mathcal{T}(G_2)$ ). *Each Hom space in  $\mathcal{T}(G_2)$  has a basis given by diagrams with no loops, in which each internal face has at least 6 edges.*

**Theorem 1.2** (isomorphism). *The categories  $\mathbf{Rep}'(G_2)$  and  $\mathcal{T}(G_2)$  are equivalent as braided tensor categories.*

### 2. Diagrammatic proof

Our goal is now to prove that an arbitrary trivalent graph in the rectangle (with equal numbers of boundary points along the top and bottom edges) can be written, modulo the relations above, as a  $\mathcal{A}$ -linear combination of diagrams coming from

braids via equation (1). Using Kuperberg’s Theorems 1.1 and 1.2, this would then give a combinatorial proof of Lehrer and Zhang’s theorem:

**Theorem 2.1** (surjectivity). *The map from the braid group algebra to endomorphisms of tensor powers of the 7-dimensional irreducible representation of  $U_q(\mathfrak{g}_2)$ ,*

$$\mathcal{AB}_n \rightarrow \text{End}_{U_q(\mathfrak{g}_2)}(V^{\otimes n}),$$

*is surjective.*

**Remark.** In fact, their result holds for any “strong multiplicity free” representation of a quantum group.

Beginning over-optimistically, we might guess that any such diagram can in fact be written as a composition of factors

$$|\cdots| \begin{array}{c} \cup \\ \cap \end{array} |\cdots|, \quad |\cdots| \begin{array}{c} \diagup \\ \diagdown \end{array} |\cdots| \quad \text{and} \quad |\cdots| \begin{array}{c} \diagdown \\ \diagup \end{array} |\cdots|$$


and then make use of the (presumably easy) special case of Theorem 2.1 for  $n = 2$ . Even though this is false, it’s the right direction and contains the essential idea. Small counterexamples to this guess are provided by

$$\begin{array}{c} \cup \\ \cap \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \quad \text{and} \quad \begin{array}{c} \cup \\ \cap \end{array} \begin{array}{c} \cup \\ \cap \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \quad \text{in } \text{End}_{U_q(\mathfrak{g}_2)}(V^{\otimes 3}).$$


The correct argument will involve three steps. First, we’ll prove a little lemma allowing us to rearrange connected components of a graph. Second, we’ll prove that the image of the braid group does hit a certain finite list of small graphs (including the examples above). Third, we’ll use Euler measure to inductively rewrite a graph in terms of linear combinations of braids and a graph with fewer vertices, eventually getting down to graphs in our list.

**Lemma 2.2.** *Suppose a graph  $D_1 \otimes D_2$  is a tensor product of two diagrams. Then  $D_1 \otimes D_2$  is in the image of the braid group if and only if  $D_2 \otimes D_1$  is.*


*Proof.* If



is in the image of the braid group, so is



Since the  $G_2$  spider is a braided tensor category, this is exactly the same as



□

**Lemma 2.3.** *The graphs*



are in the image of  $\mathcal{AB}_2$ .

*Proof.* Since all four diagrams have no loops or internal faces, they are basis diagrams. Thus by Theorem 1.1 they are linearly independent. Also, it is easy to see that they are the only basis diagrams with 4 boundary points, and so they span  $\text{End}_{U_q(\mathfrak{g}_2)}(V^{\otimes 2})$ . Moreover, the braiding (hereafter written as  $\sigma$  in formulas) lives in the same space, and has four distinct eigenvalues (see [Morrison, Peters and Snyder 2010]), and so its powers also span  $\text{End}_{U_q(\mathfrak{g}_2)}(V^{\otimes 2})$ .


Purely for reference, we'll give the characteristic equation for the braiding:

$$\sigma^4 = -q^{-16}\sigma^0 + (q^{-18} - q^{-16} - q^{-10} + q^{-4})\sigma^1 + (q^{-18} + q^{-12} - q^{-10} - q^{-6} + q^{-4} + q^2)\sigma^2 + (q^{-12} - q^{-6} - 1 + q^2)\sigma^3.$$

Explicit formulas for these four diagrams as polynomials in  $\sigma$  appear in the appendix, along with instructions for using the short computer program that produces them. (However, these formulas are not actually needed anywhere here.) □

**Remark.** Note that  $\cup$  corresponds to  $q^{10} + q^8 + q^2 + 1 + q^{-2} + q^{-8} + q^{-10}$  (this is the value of a loop, from Figure 1) times the idempotent projecting  $V \otimes V$  onto its trivial submodule. Similarly,  $\cap$  corresponds to  $-(q^6 + q^4 + q^2 + q^{-2} + q^{-4} + q^{-6})$  times the idempotent projecting onto  $V$ .

**Lemma 2.4.** *The graphs*




*are in the image of  $\mathcal{AB}_3$ .*

*Proof.* We just do the computations directly, with the help of a computer, obtaining the formulas that appear in the appendix. In particular, we find 35 braids that provide a basis for  $\text{End}_{U_q(\mathfrak{g}_2)}(V^{\otimes 3})$ .  $\square$


**Remark.** See also [Lehrer and Zhang 2006, 5.2.2(2) and 5.2.2(4)].

**Corollary 2.5.** *The graphs*




*are in the image of the appropriate braid groups.*

*Proof.* We'll show that each of these graphs is actually in the braided tensor subcategory generated by the graphs appearing in the two previous lemmas. (In fact, without even needing to take linear combinations.)



**Proposition 2.6** (base case for the induction). *Any  $\mathcal{T}(G_2)$  basis diagram in which every connected component contains at most one vertex is in the image of the braid group.*

*Proof.* Any such diagram is just a tensor product of factors, each of which is one of the graphs



By Lemma 2.2, we can take this tensor product in any order we like. We claim that any such tensor product is actually a tensor product of the graphs in Lemmas 2.3

and 2.4 and Corollary 2.5, and so the entire diagram is in the image of the braid group. The argument is a tedious but straightforward case-bash. Suppose we have the diagram

$$\cup^{\otimes a} \quad \smile^{\otimes b} \quad \cup^{\otimes c} \quad \cap^{\otimes d} \quad \begin{array}{c} \diagup \\ \diagdown \end{array}^{\otimes e} \quad \begin{array}{c} \diagdown \\ \diagup \end{array}^{\otimes f} .$$

Note that  $2a + 3c + e = 2b + 3d + f$ , by counting boundary points at top and bottom. By splitting off factors of  $\cup$ , we can assume that at most one of  $a$  and  $b$  is nonzero. Similarly, by splitting off  $\begin{array}{c} \diagup \\ \diagdown \end{array}$  or  $\begin{array}{c} \diagdown \\ \diagup \end{array}$ , we see that at most one of  $c$  and  $d$  and at most one of  $e$  and  $f$  are nonzero. Let's assume without loss of generality that  $b = 0$ .

If  $a > 0$ , then either we can split off copies of  $\begin{array}{c} \diagdown \\ \diagup \end{array}$ , or  $f < 2$ . If  $f = 1$ ,  $e$  must be zero, and we have  $2a + 3c = 3d + 1$ , and so  $a \equiv 2 \pmod{3}$ . Thus  $d \geq 1$ , and we can split off a copy of  $\begin{array}{c} \diagdown \\ \diagup \end{array}$ . Otherwise, if  $f = 0$ , we have  $2a + 3c + e = 3d$ , so  $d \geq 1$  and  $c = 0$ . Now, since  $2a + e = 3d \geq 3$ , either  $a \geq 3$ ,  $e \geq 3$ , or  $a, e \geq 1$ . In each of these cases, we can split something off, either  $\begin{array}{c} \diagdown \\ \diagup \end{array}$ ,  $\begin{array}{c} \diagup \\ \diagdown \end{array}$  or  $\begin{array}{c} \diagdown \\ \diagup \end{array}$ , respectively.

On the other hand, if  $a = 0$ , let's further assume without loss of generality that  $c = 0$ . We thus have  $e = 3d + f$ ; since  $e = 0$  would imply we have the empty diagram, we have  $f = 0$  instead, and the entire diagram is just a tensor power of  $\begin{array}{c} \diagup \\ \diagdown \end{array}$ . □

*Proof of Theorem 2.1.* Suppose now we have an arbitrary basis diagram. We will show that a connected component with at least two vertices has either a  $\begin{array}{c} \diagdown \\ \diagup \end{array}$  or a  $\begin{array}{c} \diagup \\ \diagdown \end{array}$  attached along its top or bottom edge. Since these two diagrams are Laurent polynomials in the positive crossing, we can rewrite the basis diagram as the product of something in the image of the braid group and another basis diagram with fewer vertices. Repeating this, we reduce to the case that no connected component contains more than one vertex, at which point we're done by Proposition 2.6.

The Euler measure argument is straightforward. Consider a connected component of the basis diagram with at least two vertices. Assign formal angles of  $2\pi/3$  around trivalent vertices, and of  $\pi/2$  on either side of an edge meeting the boundary. (We *don't* assign angles to the corners of the rectangle, so the total Euler measure will be the Euler measure of the disc,  $+1$ .) Since internal faces of the component have at least 6 edges, by Theorem 1.1, they have nonpositive Euler measure. Let  $b_k$  be the number of faces adjacent to the boundary, and meeting  $k$  edges of the graph. These faces meet the boundary of the disc surrounding the component exactly once, since the component is connected. Thus the Euler measure of a face counted by  $b_k$  is

$$\frac{1}{4} + \frac{1}{4} + \frac{1}{3}(k - 1) - \frac{1}{2}(k + 1) + 1 = \frac{2}{3} - \frac{1}{6}k.$$

The number  $b_1$  is zero (this could only occur if the component were a single strand, but it must have at least two vertices), and so we obtain

$$\sum_{k \geq 2} b_k \left( \frac{2}{3} - \frac{1}{6}k \right) \geq 1,$$

and so  $\frac{1}{3}b_2 + \frac{1}{6}b_3 \geq 1$ , which we soften to  $b_2 + b_3 \geq 3$ . There are thus at least three faces touching either 2 or 3 edges of the component. At least one of these must be attached to the top or bottom of the rectangle, avoiding the sides. If that face touches 3 edges, we're done, as there must be a  $\succ\prec$  adjacent to the boundary. If it only touches 2 edges, the hypothesis that the component is connected and has at least two vertices ensures that there's a  $\searrow\swarrow$  adjacent to the boundary.  $\square$

### 3. Questions

We end with two questions relating Kuperberg's spider and the category of tilting modules at a root of unity.

**Question 3.1.** We can specialize Kuperberg's spider for  $G_2$  to any root  $q$  of unity. The braiding is still defined as long as  $q + q^{-1} \neq 0$ , and the braiding is still surjective as long as

$$t(q^8 - 1)(q^4 - q^2 + 1)(q^6 + q^4 + q^2 + 1) \neq 0.$$

(See the explicit formulas in the appendix.) We can form idempotent completion, and quotient by negligibles to produce a semisimple category. Is this category equivalent to the semisimple quotient of the category of tilting modules for the integral form of  $U_q(G_2)$ ?

More optimistically, we might hope for an affirmative answer to this:

**Question 3.2.** Are these categories equivalent even before we quotient by negligible morphisms on either side?

Results in [Kuperberg 1994] and [Morrison, Peters and Snyder 2010] ensure that there is a functor from the specialization of the spider to the category of tilting modules. I'd hoped to be able to leverage the surjectivity of the braiding map into a proof that this functor was an equivalence, but I still don't see how to do this.

### Appendix: Explicit formulas

The online supplement to this article is a Mathematica notebook. It relies on the QuantumGroups package, which you can download as part of the KnotTheory package from [The Knot Atlas 2007]. After setting the path in the first line to point at your copy of the QuantumGroups package, you can run the remaining lines to produce the formulas appearing below.



Formulas for Lemma 2.3.

$$\begin{aligned}
\text{C} &= ((q^2-1)(q^4-q^2+1)(q^6+q^4+q^2+1))^{-1} \\
&\quad \times (q^{22}\sigma^0 + (-q^{20}+q^{22}+q^{28})\sigma^1 + (-q^{20}-q^{26}-q^{28})\sigma^2 - q^{26}\sigma^3), \\
\text{Y} &= ((q^2-1)(q^4-q^2+1)(q^6+q^4+q^2+1))^{-1} \\
&\quad \times ((-q^8-q^{12})\sigma^0 + (q^6-q^8+q^{10}-q^{12}+q^{20}+q^{24})\sigma^1 \\
&\quad \quad + (q^6+q^{10}-q^{18}+q^{20}-q^{22}+q^{24})\sigma^2 + (-q^{18}-q^{22})\sigma^3), \\
\text{X} &= ((q^2-1)(q^4-q^2+1)(q^6+q^4+q^2+1))^{-1} \\
&\quad \times ((q^2-q^4+2q^6+q^{12}-q^{14}-q^{18})\sigma^0 \\
&\quad \quad + (-q^{-2}-2q^4+2q^6-q^8+q^{10}+q^{12}+q^{16}-2q^{18}-q^{22}-q^{24})\sigma^1 \\
&\quad \quad + (-q^4-q^8+2q^{16}-q^{18}+q^{20}-q^{24})\sigma^2 + (q^{16}+q^{20}+q^{22})\sigma^3).
\end{aligned}$$

Formulas for Lemma 2.4.

$$\begin{aligned}
&((q^2-1)^2(q^2+1)(q^4-q^2+1)^2) \text{U} \\
&= -q^2(q^2-1)(q^{26}-q^{24}+q^{22}+q^{20}-2q^{18} \\
&\quad \quad \quad + 4q^{16}-3q^{14}+2q^{10}-4q^8+q^6-q^4-1)\mathbf{1} \\
&\quad + q^8(-q^{22}+3q^{20}-5q^{18}+4q^{16}-3q^{12}+7q^{10}-6q^8+2q^6+q^4-4q^2+2)\sigma_1 \\
&\quad - q^4(q^2-1)(q^{18}-q^{16}+2q^{14}+2q^8-q^6+q^4+1)\sigma_1^{-1} \\
&\quad + q^2(q^2-1)(q^{22}-q^{20}+q^{18}+3q^{12}-2q^{10}+3q^8+q^6-2q^4+2q^2-1)\sigma_2 \\
&\quad - q^4(q^2-1)(2q^{14}+q^{10}+3q^8-q^6+q^4+1)\sigma_2^{-1} \\
&\quad + q^{10}(q^2-1)^2(q^2+1)(q^{12}-q^{10}+3q^8-q^6+3q^4+1)\sigma_1\sigma_1 \\
&\quad + (-q^{24}+3q^{22}-3q^{20}+q^{18}+q^{16}-5q^{14}+5q^{12}-4q^{10}+q^8+2q^6-2q^4+q^2)\sigma_1\sigma_2 \\
&\quad - q^{10}(q^{10}-2q^8+q^6-q^4-q^2+1)\sigma_1\sigma_2^{-1} + q^4(q^4-q^2+1)(q^8+q^2-1)\sigma_1^{-1}\sigma_2 \\
&\quad - q^6(q^8-q^6+q^4-q^2+1)\sigma_1^{-1}\sigma_2^{-1} \\
&\quad + (-q^{24}+3q^{22}-3q^{20}+q^{18}+q^{16}-5q^{14}+5q^{12}-4q^{10}+q^8+2q^6-2q^4+q^2)\sigma_2\sigma_1 \\
&\quad + q^4(q^4-q^2+1)(q^8+q^2-1)\sigma_2\sigma_1^{-1} + (q^{22}-q^{24})\sigma_2\sigma_2 \\
&\quad - q^{10}(q^{10}-2q^8+q^6-q^4-q^2+1)\sigma_2^{-1}\sigma_1 - q^6(q^8-q^6+q^4-q^2+1)\sigma_2^{-1}\sigma_1^{-1} \\
&\quad + 0\sigma_1\sigma_1\sigma_2 + q^{16}(q^2-1)\sigma_1\sigma_1\sigma_2^{-1} \\
&\quad + (3q^{22}-6q^{20}+4q^{18}-q^{16}-2q^{14}+5q^{12}-3q^{10}+3q^8-q^4+q^2)\sigma_1\sigma_2\sigma_1 \\
&\quad + q^4(q^{12}-2q^{10}+q^8-q^6-q^4+q^2-1)\sigma_1\sigma_2\sigma_1^{-1} \\
&\quad + q^{20}(q^2-1)\sigma_1\sigma_2\sigma_2 - q^{12}\sigma_1\sigma_2^{-1}\sigma_1
\end{aligned}$$

$$\begin{aligned}
& +0\sigma_1\sigma_2^{-1}\sigma_1^{-1}+q^4(q^{12}-2q^{10}+q^8-q^6-q^4+q^2-1)\sigma_1^{-1}\sigma_2\sigma_1 \\
& +(q^{10}-q^8+q^6)\sigma_1^{-1}\sigma_2\sigma_1^{-1}+0\sigma_1^{-1}\sigma_2\sigma_2+0\sigma_1^{-1}\sigma_2^{-1}\sigma_1 \\
& +q^8(q^4-q^2+1)\sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1}+0\sigma_2\sigma_1\sigma_1+q^{20}(q^2-1)\sigma_2\sigma_2\sigma_1 \\
& +0\sigma_2\sigma_2\sigma_1^{-1}+q^{16}(q^2-1)\sigma_2^{-1}\sigma_1\sigma_1+(q^{18}-q^{20})\sigma_1\sigma_1\sigma_2\sigma_1 \\
& +(q^{18}-q^{20})\sigma_1\sigma_2\sigma_1\sigma_1+(q^{18}-q^{20})\sigma_1\sigma_2\sigma_2\sigma_1+0\sigma_2\sigma_1\sigma_1\sigma_2, \\
& ((q^2-1)^2(q^2+1)(q^4+1)(q^4-q^2+1)^2) \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \\
= & -q^2(q^2-1)(q^4+1)(q^{24}-q^{22}+2q^{18}-2q^{16}+q^{12}-2q^{10}-q^4+q^2-1)\mathbf{1} \\
& -q^2(q^{30}-3q^{28}+5q^{26}-4q^{24}+2q^{20}-3q^{18}+q^{16} \\
& \qquad \qquad \qquad +2q^{14}-q^{12}+2q^{10}-2q^6+2q^4-2q^2+1)\sigma_1 \\
& +(-q^{26}+2q^{24}-2q^{22}+q^{20}+q^{18}-q^{16}+q^{14}-q^{10}+q^8-q^6+q^4)\sigma_1^{-1} \\
& +q^2(q^{26}-2q^{24}+2q^{22}-4q^{18}+8q^{16}-9q^{14}+5q^{12}-5q^8+6q^6-5q^4+3q^2-1)\sigma_2 \\
& -q^4(q^{18}-q^{16}+q^{12}-2q^{10}+q^6-q^4+q^2-1)\sigma_2^{-1} \\
& +(q^{30}-2q^{28}+3q^{26}-2q^{24}-q^{18}+q^{12})\sigma_1\sigma_1 \\
& -q^2(q^{24}-3q^{22}+3q^{20}+q^{18}-5q^{16}+9q^{14} \\
& \qquad \qquad \qquad -7q^{12}+3q^{10}-3q^6+3q^4-2q^2+1)\sigma_1\sigma_2 \\
& +(q^{18}-q^{10}+q^8-q^6+q^4)\sigma_1\sigma_2^{-1} \\
& +q^4(q^4-q^2+1)(q^{14}-q^{12}+q^{10}-q^6+q^4-q^2+1)\sigma_1^{-1}\sigma_2-q^6\sigma_1^{-1}\sigma_2^{-1} \\
& -q^6(q^{20}-2q^{18}+2q^{16}-q^{12}+3q^{10}-2q^8+q^4-2q^2+1)\sigma_2\sigma_1 \\
& +q^6(q^4-q^2+1)(q^8+q^2-1)\sigma_2\sigma_1^{-1}-q^{14}(q^2-1)(q^4+1)(q^6-q^2+1)\sigma_2\sigma_2 \\
& -q^{12}(q^{10}-2q^8+q^6-q^4-q^2+1)\sigma_2^{-1}\sigma_1 \\
& -q^2(q^{14}-q^{10}+q^8+q^6-2q^4+2q^2-1)\sigma_2^{-1}\sigma_1^{-1}+q^{14}(q^4-q^2+1)\sigma_1\sigma_1\sigma_2 \\
& -q^{16}\sigma_1\sigma_1\sigma_2^{-1}+q^6(2q^{18}-4q^{16}+2q^{14}+q^{12}-3q^{10}+5q^8-2q^6+q^4-1)\sigma_1\sigma_2\sigma_1 \\
& +q^6(q^4+1)(q^8-2q^6+q^2-1)\sigma_1\sigma_2\sigma_1^{-1}+q^{14}(q^{10}-q^8+q^4-q^2+1)\sigma_1\sigma_2\sigma_2 \\
& -q^{14}\sigma_1\sigma_2^{-1}\sigma_1+(q^2+1)(q^5-q^3+q)^2\sigma_1\sigma_2^{-1}\sigma_1^{-1}-q^8(q^2-1)(q^4+1)\sigma_1^{-1}\sigma_2\sigma_1 \\
& +q^8(q^4-q^2+1)\sigma_1^{-1}\sigma_2\sigma_1^{-1}-q^{16}(q^4-q^2+1)\sigma_1^{-1}\sigma_2\sigma_2+0\sigma_1^{-1}\sigma_2^{-1}\sigma_1 \\
& +(-q^8+q^6-q^4)\sigma_1^{-1}\sigma_2^{-1}\sigma_1^{-1}+0\sigma_2\sigma_1\sigma_1+q^{22}(q^2-1)\sigma_2\sigma_2\sigma_1 \\
& +0\sigma_2\sigma_2\sigma_1^{-1}+q^{18}(q^2-1)\sigma_2^{-1}\sigma_1\sigma_1+q^{12}(q^2-1)(q^4+1)\sigma_1\sigma_1\sigma_2\sigma_1 \\
& +(q^{20}-q^{22})\sigma_1\sigma_2\sigma_1\sigma_1+(q^{20}-q^{22})\sigma_1\sigma_2\sigma_2\sigma_1+0\sigma_2\sigma_1\sigma_1\sigma_2.
\end{aligned}$$

## References

- [Chari and Pressley 1995] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge University Press, 1995. MR 96h:17014 Zbl 0839.17010
- [Jantzen 1996] J. C. Jantzen, *Lectures on quantum groups*, Graduate Studies in Mathematics **6**, American Mathematical Society, Providence, RI, 1996. MR 96m:17029 Zbl 0842.17012
- [Kuperberg 1994] G. Kuperberg, “The quantum  $G_2$  link invariant”, *Internat. J. Math.* **5**:1 (1994), 61–85. MR 95g:57013 Zbl 0797.57008
- [Kuperberg 1996] G. Kuperberg, “Spiders for rank 2 Lie algebras”, *Comm. Math. Phys.* **180**:1 (1996), 109–151. MR 97f:17005 Zbl 0870.17005
- [Lehrer and Zhang 2006] G. I. Lehrer and R. B. Zhang, “Strongly multiplicity free modules for Lie algebras and quantum groups”, *J. Algebra* **306** (2006), 138–174. MR 2007g:17008 Zbl 1169.17003
- [Morrison, Peters and Snyder 2010] S. Morrison, E. Peters, and N. Snyder, “Knot polynomial identities and quantum group coincidences”, preprint, 2010. To appear in *Quantum Topology*. arXiv 1003.0022
- [Sawin 2006] S. F. Sawin, “Quantum groups at roots of unity and modularity”, *J. Knot Theory Ramifications* **15**:10 (2006), 1245–1277. MR 2008f:17032 Zbl 1117.17006
- [The Knot Atlas 2007] D. Bar-Natan and S. Morrison (editors), “The Knot Atlas”, wiki, 2007, available at <http://katlas.org/>.

Received September 22, 2009. Revised January 4, 2010.

SCOTT MORRISON  
MICROSOFT STATION Q  
CNSI BUILDING  
UNIVERSITY OF CALIFORNIA  
SANTA BARBARA, CA 93106-6105  
scott@math.berkeley.edu  
<http://tqft.net/>

# PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by  
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [www.pjmath.org](http://www.pjmath.org) for submission instructions.

---

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 249    No. 1    January 2011

---

Metabelian $SL(n, \mathbb{C})$ representations of knot groups, II: Fixed points	1
HANS U. BODEN and STEFAN FRIEDL	
Lewis–Zagier correspondence for higher-order forms	11
ANTON DEITMAR	
Topology of positively curved 8-dimensional manifolds with symmetry	23
ANAND DESSAI	
Strong Kähler with torsion structures from almost contact manifolds	49
MARISA FERNÁNDEZ, ANNA FINO, LUIS UGARTE and RAQUEL VILLACAMPA	
Connections between Floer-type invariants and Morse-type invariants of Legendrian knots	77
MICHAEL B. HENRY	
A functional calculus for unbounded generalized scalar operators on Banach spaces	135
DRAGOLJUB KEČKIĆ and ĐORĐE KRINIĆ	
Geometric formality of homogeneous spaces and of biquotients	157
D. KOTSCHICK and S. TERZIĆ	
Positive solutions for a nonlinear third order multipoint boundary value problem	177
YANG LIU, ZHANG WEIGUO, LIU XIPING, SHEN CHUNFANG and CHEN HUA	
The braid group surjects onto $G_2$ tensor space	189
SCOTT MORRISON	
Analogues of the Wiener Tauberian and Schwartz theorems for radial functions on symmetric spaces	199
E. K. NARAYANAN and ALLADI SITARAM	
Semidirect products of representations up to homotopy	211
YUNHE SHENG and CHENCHANG ZHU	
Homology sequence and excision theorem for Euler class group	237
YONG YANG	
Acknowledgement	255



0030-8730(201101)249:1;1-D