

*Pacific  
Journal of  
Mathematics*

**NECESSARY AND SUFFICIENT CONDITIONS FOR UNIT  
GRAPHS TO BE HAMILTONIAN**

H. R. MAIMANI, M. R. POURNAKI AND S. YASSEMI

## NECESSARY AND SUFFICIENT CONDITIONS FOR UNIT GRAPHS TO BE HAMILTONIAN

H. R. MAIMANI, M. R. POURNAKI AND S. YASSEMI

**The unit graph corresponding to an associative ring  $R$  is the graph obtained by setting all the elements of  $R$  to be the vertices and defining distinct vertices  $x$  and  $y$  to be adjacent if and only if  $x + y$  is a unit of  $R$ . By a constructive method, we derive necessary and sufficient conditions for unit graphs to be Hamiltonian.**

### 1. Introduction

A graph is *Hamiltonian* if it has a cycle that visits every vertex exactly once; such a cycle is called a *Hamiltonian cycle*. In general, the problem of finding a Hamiltonian cycle in a given graph is an *NP*-complete problem and a special case of the traveling salesman problem. It is a problem in combinatorial optimization studied in operations research and theoretical computer science; see [Garey and Johnson 1979]. The only known way to determine whether a given graph has a Hamiltonian cycle is to undertake an exhaustive search, and until now no theorem giving a necessary and sufficient condition for a graph to be Hamiltonian was known. The study of Hamiltonian graphs has long been an important topic. See [Gould 2003] for a survey, updating earlier surveys in this area.

Let  $n$  be a positive integer, and let  $\mathbb{Z}_n$  be the ring of integers modulo  $n$ . Grimaldi [1990] defined a graph  $G(\mathbb{Z}_n)$  based on the elements and units of  $\mathbb{Z}_n$ . The vertices of  $G(\mathbb{Z}_n)$  are the elements of  $\mathbb{Z}_n$ , and distinct vertices  $x$  and  $y$  are defined to be adjacent if and only if  $x + y$  is a unit of  $\mathbb{Z}_n$ . For a positive integer  $m$ , it follows that  $G(\mathbb{Z}_{2m})$  is a  $\varphi(2m)$ -regular graph, where  $\varphi$  is the Euler phi function. In case  $m \geq 2$ , the graph  $G(\mathbb{Z}_{2m})$  can be expressed as the union of  $\varphi(2m)/2$  Hamiltonian cycles. The odd case is not quite so easy, but the structure is clear and the results are similar to the even case. We recall that a *cone* over a graph is obtained by taking

---

The research of H. R. Maimani and S. Yassemi was in part supported by a grant from IPM (numbers 89050211 and 89130213). The research of M. R. Pournaki was in part supported by a grant from the Academy of Sciences for the Developing World (TWAS–UNESCO Associateship—Ref. FR3240126591).

*MSC2000:* primary 05C45; secondary 13M05.

*Keywords:* Hamiltonian cycle, Hamiltonian graph, finite ring.

the categorical product of the graph and a path with a loop at one end, and then identifying all the vertices whose second coordinate is the other end of the path. When  $p$  is an odd prime,  $G(\mathbb{Z}_p)$  can be expressed as a cone over a complete partite graph with  $(p - 1)/2$  partitions of size two. This leads to an explicit formula for the chromatic polynomial of  $G(\mathbb{Z}_p)$ . Grimaldi [1990] also concludes with some properties of the graphs  $G(\mathbb{Z}_{p^m})$ , where  $p$  is a prime number and  $m \geq 2$ . Recently, the authors of this paper generalized  $G(\mathbb{Z}_n)$  to  $G(R)$ , the unit graph of  $R$ , where  $R$  is an arbitrary associative ring with nonzero identity and studied the properties of this graph; see [Ashrafi et al. 2010; Maimani et al. 2010].

By a constructive method, we derive necessary and sufficient conditions for unit graphs to be Hamiltonian.

## 2. Preliminaries and the main result

Throughout the paper, by a graph we mean a finite undirected graph without loops or multiple edges. Also all rings are finite commutative with nonzero identity. For undefined terms and concepts, see [West 1996; Atiyah and Macdonald 1969].


We first start with recalling some notions from graph theory. For a graph  $G$  and for any two vertices  $x$  and  $y$  of  $G$ , we recall that a *walk* between  $x$  and  $y$  is a sequence  $x = v_0, e_1, v_1, \dots, e_k, v_k = y$  of vertices and edges of  $G$ , denoted by

$$x = v_0 \rightarrow v_1 \rightarrow \dots \rightarrow v_k = y,$$


such that for every  $i$  with  $1 \leq i \leq k$ , the edge  $e_i$  has endpoints  $v_{i-1}$  and  $v_i$ . Also a *path* between  $x$  and  $y$  is a walk between  $x$  and  $y$  without repeated vertices. A *cycle* of a graph is a path such that the start and end vertices are the same. Two cycles are considered the same if they consist of the same vertices and edges. The number of edges (counting repeats) in a walk, path or a cycle, is called its *length*. A *Hamiltonian path (cycle)* in  $G$  is a path (cycle) in  $G$  that visits every vertex exactly once. A graph is called *Hamiltonian* if it contains a Hamiltonian cycle. Also a graph  $G$  is called *connected* if for any vertices  $x$  and  $y$  of  $G$  there is a path between  $x$  and  $y$ .

We now define the unit graph corresponding to a ring. Let  $R$  be a ring and  $U(R)$  be the set of unit elements of  $R$ . The *unit graph* of  $R$ , denoted by  $G(R)$ , is the graph obtained by setting all the elements of  $R$  to be the vertices and defining distinct vertices  $x$  and  $y$  to be adjacent if and only if  $x + y \in U(R)$ . The graphs in Figure 1 are the unit graphs of the rings indicated. It is easy to see that, for given rings  $R$  and  $S$ , if  $R \cong S$  as rings, then  $G(R) \cong G(S)$  as graphs. This point is illustrated in Figure 2.

We continue this section by collecting some notions from ring theory. First of all, for a given ring  $R$ , the *Jacobson radical*  $J(R)$  of  $R$  is defined to the intersection



**Figure 1.** Unit graphs of some specific rings.



**Figure 2.** Unit graphs of two isomorphic rings.

of all maximal ideals of  $R$ . Let  $R$  be a ring and let  $k$  be a positive integer. An element  $r \in R$  is said to be  $k$ -good if we may write  $r = u_1 + \dots + u_k$ , where  $u_1, \dots, u_k \in U(R)$ . The ring  $R$  is said to be  $k$ -good if every element of  $R$  is  $k$ -good. Following [Goldsmith et al. 1998], we now define an invariant of a ring, called the unit sum number, which expresses in a fairly precise way how the units generate the ring. The *unit sum number*  $u(R)$  of  $R$  is given by

- $\min\{k \mid R \text{ is } k\text{-good}\}$  if  $R$  is  $k$ -good for some  $k \geq 1$ ,
- $\omega$  if  $R$  is not  $k$ -good for every  $k$ , but every element of  $R$  is  $k$ -good for some  $k$  (that is, when at least  $U(R)$  generates  $R$  additively), and
- $\infty$  otherwise (that is, when  $U(R)$  does not generate  $R$  additively).

For example, let  $D$  be a division ring. If  $|D| \geq 3$ , then  $u(D) = 2$ ; whereas if  $|D| = 2$ , that is,  $D = \mathbb{Z}_2$ , the field of two elements, then  $u(\mathbb{Z}_2) = \omega$ . We have also  $u(\mathbb{Z}_2 \times \mathbb{Z}_2) = \infty$  — see [Ashrafi and Vámos 2005] for unit sum numbers of some other rings. The topic of unit sum numbers seems to have arisen with a paper by Zelinsky [1954], in which he shows that if  $V$  is any finite- or infinite-dimensional vector space over a division ring  $D$ , then every linear transformation is the sum of two automorphisms unless  $\dim V = 1$  and  $D$  is the field of two elements. Interest in this topic increased recently after Goldsmith, Pabst and Scott [1998] defined the

unit sum number. For additional historical background, see [Vámos 2005], which also contains references to recent work in this area.

We are now ready to state the main result of this paper. The proof is given in Section 3 by a sequence of lemmas and propositions.

**Theorem 2.1.** *Let  $R$  be a ring such that  $R \not\cong \mathbb{Z}_2$  and  $R \not\cong \mathbb{Z}_3$ . Then the following statements are equivalent:*

- (a) *The unit graph  $G(R)$  is Hamiltonian.*
- (b) *The ring  $R$  cannot have  $\mathbb{Z}_2 \times \mathbb{Z}_2$  as a quotient.*
- (c) *The ring  $R$  is generated by its units.*
- (d) *The unit sum number of  $R$  is less than or equal to  $\omega$ .*
- (e) *The unit graph  $G(R)$  is connected.*

### 3. The proofs

In this section we state and prove some lemmas that will be used in the proof of Theorem 2.1. For the convenience of the reader we state without proof a few known results in the form of propositions that will be used in the proofs. We also recall some definitions and notations for later use.

A *bipartite* graph is one whose vertex-set is partitioned into two (not necessarily nonempty) disjoint subsets so that the two end vertices for each edge lie in distinct partitions. Among bipartite graphs, a *complete bipartite* graph is one in which each vertex is joined to every vertex that is not in the same partition. The complete bipartite graph with two partitions of size  $m$  and  $n$  is denoted by  $K_{m,n}$ .

The following result characterizes the complete bipartite unit graphs of rings.

**Proposition 3.1** [Ashrafi et al. 2010, Theorem 3.5]. *Let  $R$  be a ring and  $\mathfrak{m}$  be a maximal ideal of  $R$  such that  $|R/\mathfrak{m}| = 2$ . Then  $G(R)$  is a bipartite graph. The unit graph  $G(R)$  is a complete bipartite graph if and only if  $R$  is a local ring.*

The degrees of all vertices of a unit graph is given by the following result. For a graph  $G$  and for a vertex  $x$  of  $G$ , the *degree*  $\deg(x)$  of  $x$  is the number of edges of  $G$  incident with  $x$ .

**Proposition 3.2** [Ashrafi et al. 2010, Proposition 2.4]. *Let  $R$  be a ring. Then the following statements hold for the unit graph of  $R$ :*

- (1) *If  $2 \notin U(R)$ , then  $\deg(x) = |U(R)|$  for every  $x \in R$ .*
- (2) *If  $2 \in U(R)$ , then  $\deg(x) = |U(R)| - 1$  for every  $x \in U(R)$  and  $\deg(x) = |U(R)|$  for every  $x \in R \setminus U(R)$ .*

We also need the following well known result due to Dirac, which initiated the study of Hamiltonian graphs. This work was continued by Ore [1960].

**Proposition 3.3** [Dirac 1952, Theorem 3]. *If  $G$  is a graph with  $n$  vertices,  $n \geq 3$ , and every vertex has degree at least  $n/2$ , then  $G$  is Hamiltonian.*

**Lemma 3.4.** *Let  $R$  be a local ring with  $|R| \geq 4$ . Then the unit graph  $G(R)$  is Hamiltonian.*

*Proof.* Suppose  $\mathfrak{m}$  is the unique maximal ideal of  $R$ . There are two possibilities: either  $|R/\mathfrak{m}| = 2$  or  $|R/\mathfrak{m}| > 2$ .

First, suppose that  $|R/\mathfrak{m}| = 2$ . In this case, Proposition 3.1 implies that the unit graph  $G(R)$  is a complete bipartite graph. Moreover, its proof shows that  $\mathfrak{m}$  and  $R \setminus \mathfrak{m}$  are the partite sets of  $G(R)$ . Since  $|R/\mathfrak{m}| = 2$ , we conclude that  $|\mathfrak{m}| = |R \setminus \mathfrak{m}|$  and so  $G(R) \cong K_{|\mathfrak{m}|, |\mathfrak{m}|}$ . The assumptions  $|R| \geq 4$  and  $|R/\mathfrak{m}| = 2$  imply that  $|\mathfrak{m}| \geq 2$  and thus  $G(R)$  is Hamiltonian.

Second, suppose that  $|R/\mathfrak{m}| > 2$ . In this case, Proposition 3.2 implies that  $\deg(x) \geq |U(R)| - 1$  for all  $x \in R$ . We claim that  $|U(R)| - 1 \geq |R|/2$ . To show this, note that  $R$  is a local ring with  $|R| \geq 4$ . If  $|R| = 4$ , then the assumption  $|R/\mathfrak{m}| > 2$  implies that  $|\mathfrak{m}| < 2$  and so  $\mathfrak{m} = 0$ . Therefore  $R$  is a field and so  $|U(R)| = 3$ . Thus  $|U(R)| - 1 = 2 = |R|/2$ . If  $|R| = 5$ , then  $R$  is again a field and so  $|U(R)| = 4$ . Thus  $|U(R)| - 1 = 3 > 2.5 = |R|/2$ . If  $|R| \geq 6$ , then since  $R$  is local with  $|R/\mathfrak{m}| > 2$ , we conclude that  $|U(R)| \geq 2|R|/3$ . Therefore  $|U(R)| - 1 \geq (2|R|/3) - 1 \geq |R|/2$ . Thus the claim holds and so  $\deg(x) \geq |R|/2$  for every  $x \in R$ . Therefore Proposition 3.3 implies that  $G(R)$  is Hamiltonian.  $\square$

The following result gives us information about the existence of a Hamiltonian cycle in unit graphs of the direct product of a ring and a field.

**Lemma 3.5.** *Let  $T$  be a ring with Hamiltonian unit graph and let  $F$  be a field. If  $F \not\cong \mathbb{Z}_2$ , then the unit graph  $G(T \times F)$  is Hamiltonian.*

*Proof.* Since the unit graph  $G(T)$  is Hamiltonian, there is a Hamiltonian cycle with length  $n = |T|$  in  $G(T)$ , say

$$0 = a_1 \rightarrow a_2 \rightarrow \dots \rightarrow a_{n-1} \rightarrow a_n \rightarrow a_{n+1} = 0.$$

Either the characteristic of  $F$  is equal to 2 or it is not.

First, suppose the latter. In this case we may assume that

$$F = \{0, x_1, \dots, x_{(|F|-1)/2}, -x_1, \dots, -x_{(|F|-1)/2}\}.$$

If  $n$  is even and  $|F| \geq 5$ , then  $x_2 \neq -x_1$  and so  $x_1 + x_2$  is a unit element of  $F$ . Now consider the following paths in the unit graph  $G(T \times F)$ :

$$\begin{aligned} P_0 &: (0, 0) \rightarrow (a_2, x_1) \rightarrow (a_3, 0) \rightarrow (a_4, x_1) \rightarrow \dots \rightarrow (a_n, x_1), \\ P_1 &: (0, x_2) \rightarrow (a_2, 0) \rightarrow (a_3, x_2) \rightarrow \dots \rightarrow (a_{n-1}, x_2) \rightarrow (a_n, 0), \\ P_2 &: (0, x_1) \rightarrow (a_2, x_2) \rightarrow (a_3, x_1) \rightarrow \dots \rightarrow (a_n, x_2). \end{aligned}$$

Also for every  $i$  with  $3 \leq i \leq (|F| - 1)/2$ , consider the path

$$P_i : (0, x_i) \rightarrow (a_2, x_i) \rightarrow (a_3, x_i) \rightarrow \cdots \rightarrow (a_n, x_i),$$

and for every  $i$  with  $1 \leq i \leq (|F| - 1)/2$ , consider the path

$$P'_i : (0, -x_i) \rightarrow (a_2, -x_i) \rightarrow \cdots \rightarrow (a_n, -x_i).$$

It is easy to see that  $P_{i-1}$  is adjacent to  $P_i$  for every  $i$  with  $1 \leq i \leq (|F| - 1)/2$  and  $P'_{i-1}$  is adjacent to  $P'_i$  for every  $i$  with  $2 \leq i \leq (|F| - 1)/2$ , and  $P_{(|F|-1)/2}$  is adjacent to  $P'_1$ . Therefore  $P_0 P_1 P_2 P_3 \cdots P_{(|F|-1)/2} P'_1 \cdots P'_{(|F|-1)/2} (0, 0)$  is a Hamiltonian cycle in the unit graph  $G(T \times F)$ , which shows that it is Hamiltonian. If  $n$  is even and  $|F| = 3$ , then  $F \cong \mathbb{Z}_3$  and thus the cycle

$$\begin{aligned} (a_1, 1) &\rightarrow (a_2, 0) \rightarrow (a_3, 2) \rightarrow (a_4, 2) \rightarrow (a_3, 0) \\ &\rightarrow (a_2, 1) \rightarrow (a_3, 1) \rightarrow \cdots \rightarrow (a_{n-2}, 1) \rightarrow (a_{n-1}, 1) \\ &\rightarrow (a_1, 2) \rightarrow (a_2, 2) \rightarrow (a_1, 0) \rightarrow (a_n, 1) \rightarrow (a_1, 1), \end{aligned}$$

is a Hamiltonian cycle in the unit graph  $G(T \times F)$ , and thus it is Hamiltonian.

If  $n$  is odd and  $|F| \geq 5$ , consider the path

$$P_0 : (a_1, 0) \rightarrow (a_2, x_1) \rightarrow \cdots \rightarrow (a_n, 0) \rightarrow (a_1, x_1) \rightarrow (a_2, 0) \rightarrow \cdots \rightarrow (a_n, x_1),$$

and for  $1 \leq i \leq (|F| - 1)/2$  consider the paths

$$\begin{aligned} P_i &: (a_1, x_i) \rightarrow (a_2, x_i) \rightarrow \cdots \rightarrow (a_n, x_i), \\ P'_i &: (a_1, -x_i) \rightarrow (a_2, -x_i) \rightarrow \cdots \rightarrow (a_n, -x_i). \end{aligned}$$

It is easy to see that  $P_0 P_1 \cdots P_{(|F|-1)/2} P'_1 \cdots P'_{(|F|-1)/2} (a_1, 0)$  is a Hamiltonian cycle in the unit graph  $G(T \times F)$  and thus it is Hamiltonian. If  $n$  is odd and  $|F| = 3$ , we may obtain a Hamiltonian cycle in the unit graph  $G(T \times F)$  by replacing the eleven end-vertices in the cycle above with

$$\begin{aligned} (a_{n-3}, 1) &\rightarrow (a_{n-2}, 1) \rightarrow (a_{n-1}, 0) \rightarrow (a_n, 2) \rightarrow (a_{n-1}, 1) \\ &\rightarrow (a_n, 1) \rightarrow (a_1, 0) \rightarrow (a_2, 2) \rightarrow (a_1, 2) \rightarrow (a_n, 0) \rightarrow (a_1, 1). \end{aligned}$$

This shows that the unit graph  $G(T \times F)$  is Hamiltonian.

Second, suppose that characteristic of  $F$  is equal to 2. Therefore we have  $|F| \geq 4$ . In this case we may assume that

$$F = \{x_1, \dots, x_{2^m}\} = \{x_{2i-1}, x_{2i} \mid 1 \leq i \leq 2^{m-1}\}.$$

If  $n$  is even, then for every  $i$  with  $1 \leq i \leq 2^{m-1}$ , consider the following paths in the unit graph  $G(T \times F)$ :

$$\begin{aligned} P_i &: (a_1, x_{2i-1}) \rightarrow (a_2, x_{2i}) \rightarrow \cdots \rightarrow (a_n, x_{2i}), \\ P'_i &: (a_1, x_{2i}) \rightarrow (a_2, x_{2i-1}) \rightarrow \cdots \rightarrow (a_n, x_{2i-1}). \end{aligned}$$

Since  $|F| \geq 4$ , it is clear that  $P_1 P'_{2^{m-1}} P_2 P'_{2^{m-1}-1} \cdots P_{2^{m-1}} P'_1(0, x_1)$  is a Hamiltonian cycle in the unit graph  $G(T \times F)$  and thus it is Hamiltonian.

If  $n$  is odd, then consider the path

$$P_i : (a_1, x_{2i-1}) \rightarrow (a_2, x_{2i}) \rightarrow \cdots \rightarrow (a_{n-1}, x_{2i}) \rightarrow (a_n, x_{2i-1}) \rightarrow \cdots \rightarrow (a_n, x_{2i}).$$

Therefore  $P_1 P_2 \cdots P_{2^{m-1}}(a_1, x_1)$  is a Hamiltonian cycle in the unit graph  $G(T \times F)$  and thus it is Hamiltonian.  $\square$

In the sequel we need Lemmas 3.7, 3.8 and 3.10. But first, we state the following proposition, which is useful in the proof of Lemma 3.7. Recall that a *clique* of a graph  $G$  is a complete subgraph of  $G$ . Also a *coclique* (also called an *independent set of vertices*) in a graph  $G$  is a set of pairwise nonadjacent vertices.

**Proposition 3.6** [Ashrafi et al. 2010, Lemma 2.7]. *Let  $R$  be a ring and suppose that  $J(R)$  denotes the Jacobson radical of  $R$ . Suppose  $x, y \in R$ .*

- (a) *If  $x + J(R)$  and  $y + J(R)$  are adjacent in the unit graph  $G(R/J(R))$ , then every element of  $x + J(R)$  is adjacent to every element of  $y + J(R)$  in the unit graph  $G(R)$ .*
- (b) *If  $2x \in U(R)$ , then  $x + J(R)$  is a clique in the unit graph  $G(R)$ .*
- (c) *If  $2x \notin U(R)$ , then  $x + J(R)$  is a coclique in the unit graph  $G(R)$ .*

**Lemma 3.7.** *Let  $T$  be a ring and let  $R$  be a local ring with unique maximal ideal  $\mathfrak{m}$ . If the unit graph  $G(T \times R/\mathfrak{m})$  is Hamiltonian, then the unit graph  $G(T \times R)$  is Hamiltonian.*

*Proof.* Since the unit graph  $G(T \times R/\mathfrak{m})$  is Hamiltonian, there is a Hamiltonian cycle in  $G(T \times R/\mathfrak{m})$ , say

$$(a_1, y_1 + \mathfrak{m}) \rightarrow \cdots \rightarrow (a_n, y_n + \mathfrak{m}) \rightarrow (a_1, y_1 + \mathfrak{m}),$$

where  $n = |T \times R/\mathfrak{m}|$ . Let  $\mathfrak{m} = \{x_1, \dots, x_t\}$ . Therefore for every  $i$  with  $1 \leq i \leq t$ , we have  $y_i + \mathfrak{m} = \{y_i + x_1, \dots, y_i + x_t\}$  and so  $T \times R = \bigcup_{i=1}^n M_i$ , where  $M_i = \{(a_i, y_i + x_j) \mid 1 \leq j \leq t\}$ . It is easy to see that for every  $r$  with  $1 \leq r \leq n - 1$ , every element of  $M_r$  is adjacent to every element of  $M_{r+1}$ . Also every element of  $M_n$  is adjacent to every element of  $M_1$ . Let  $S_r$  for  $1 \leq r \leq n - 1$  be a subgraph of the unit graph  $G(T \times R)$  with vertex-set  $M_r \cup M_{r+1}$  and edge-set  $\{(a_r, y_r + x_j) \rightarrow (a_{r+1}, y_{r+1} + x_\ell) \mid 1 \leq j, \ell \leq t\}$ . Also let  $S_n$  be a subgraph of the unit graph  $G(T \times R)$  with vertex-set  $M_n \cup M_1$  and edge-set  $\{(a_n, y_n + x_j) \rightarrow (a_1, y_1 + x_\ell) \mid 1 \leq j, \ell \leq t\}$ . It is easy to see that  $S_r$  for  $1 \leq r \leq n$  is a Hamiltonian complete bipartite subgraph of the unit graph  $G(T \times R)$ . For every  $r$  with  $1 \leq r \leq n - 1$ , let  $P_r$  be a Hamiltonian path of  $S_r$  with initial vertex  $(a_r, y_r + x_1)$  and end point  $(a_{r+1}, y_{r+1} + x_1)$ . Also let  $P_n$  be a Hamiltonian path of  $S_n$  with initial vertex  $(a_n, y_n + x_1)$  and end point  $(a_1, y_1 + x_1)$ . Now we consider the following two cases:



Case 1:  $n$  is even. In this case, the cycle

$$P_1 \rightarrow P_3 \rightarrow \dots \rightarrow P_{n-1} \rightarrow (a_1, y_1 + x_1)$$

is a Hamiltonian cycle in the unit graph  $G(T \times R)$  and thus it is Hamiltonian.

Case 2:  $n$  is odd. In this case, since  $|T \times R/\mathfrak{m}|$  is odd,  $|R/\mathfrak{m}|$  is odd. This implies that  $|R|$  is odd and so  $2 \in U(R)$ . We may assume that  $y_1 + \mathfrak{m} = \mathfrak{m}$ . Therefore  $y_n + \mathfrak{m} \neq \mathfrak{m}$ . Now Proposition 3.6 implies that the subgraph induced by  $M_n$  is a clique. Therefore the cycle

$$P_1 \rightarrow P_3 \rightarrow \dots \rightarrow P_{n-2} \rightarrow (a_n, y_n + x_1) \rightarrow \dots \rightarrow (a_n, y_n + x_t) \rightarrow (a_1, y_1 + x_1)$$

is a Hamiltonian cycle in the unit graph  $G(T \times R)$  and thus it is Hamiltonian.  $\square$

**Lemma 3.8.** *Let  $R \cong R_1 \times \dots \times R_n$ , where every  $R_i$  is a local ring with maximal ideal  $\mathfrak{m}_i$ . Suppose that  $R \not\cong \mathbb{Z}_3$  and for every  $i$  with  $1 \leq i \leq n$ , we have  $R_i/\mathfrak{m}_i \not\cong \mathbb{Z}_2$ . Then the unit graph  $G(R)$  is Hamiltonian.*

*Proof.* We prove the lemma by induction on  $n$ . If  $n = 1$ , then  $R$  is local and assumptions imply that  $|R| \geq 4$ . Therefore by using Lemma 3.4 we conclude that the unit graph  $G(R)$  is Hamiltonian. Now suppose that the lemma holds true for  $n - 1$ . Consider  $T = R_1 \times \dots \times R_{n-1}$  and  $F = R_n/\mathfrak{m}_n$ . There are two possibilities: either  $T \cong \mathbb{Z}_3$  or  $T \not\cong \mathbb{Z}_3$ .

First, suppose that  $T \cong \mathbb{Z}_3$ . If  $|R_n| \geq 4$ , then by Lemma 3.5 the unit graph  $G(R) \cong G(\mathbb{Z}_3 \times R_n)$  is Hamiltonian. If  $|R_n| = 3$ , then  $R_n \cong \mathbb{Z}_3$  and so  $R \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ . Therefore the cycle

$$\begin{aligned} (0, 0) \rightarrow (1, 1) \rightarrow (0, 1) \rightarrow (2, 1) \rightarrow (2, 0) \rightarrow (2, 2) \\ \rightarrow (0, 2) \rightarrow (1, 0) \rightarrow (1, 2) \rightarrow (0, 0), \end{aligned}$$

is a Hamiltonian cycle in the unit graph  $G(R) \cong G(\mathbb{Z}_3 \times \mathbb{Z}_3)$  and thus it is Hamiltonian.

Second, suppose that  $T \not\cong \mathbb{Z}_3$ . In this case the induction hypothesis implies that the unit graph  $G(T)$  is Hamiltonian. On the other hand,  $F \cong R_n/\mathfrak{m}_n$  is a field with  $|F| \geq 3$ . Therefore Lemma 3.5 implies that the unit graph  $G(T \times F)$  is Hamiltonian. Therefore by applying Lemma 3.7, we conclude that the unit graph  $G(R)$  is Hamiltonian.  $\square$

We need the following result to give a proof of Lemma 3.10.

**Proposition 3.9** [Chartrand and Oellermann 1993, Theorem 8.6]. *Let  $G$  be a bipartite graph with partite sets  $X$  and  $Y$  such that  $|X| = |Y| = n \geq 2$ . If  $\deg(x) > n/2$  for every vertex  $x$  of  $G$ , then  $G$  is Hamiltonian.*

**Lemma 3.10.** *Let  $R \cong R_1 \times \dots \times R_n \times \mathbb{Z}_2$ , where every  $R_i$  is a local ring with maximal ideal  $\mathfrak{m}_i$ . If  $R_i/\mathfrak{m}_i \not\cong \mathbb{Z}_2$  for every  $i$  with  $1 \leq i \leq n$ , then the unit graph  $G(R)$  is Hamiltonian.*

*Proof.* We prove the lemma by induction on  $n$ . If  $n = 1$ , then  $R \cong R_1 \times \mathbb{Z}_2$ . In this case, it is easy to see that the unit graph  $G(R)$  is a bipartite graph with partite sets  $X = R_1 \times \{0\}$  and  $Y = R_1 \times \{1\}$ . On the other hand, by Proposition 3.2(1), we have  $\deg(x) = |U(R)| = |U(R_1)| > |U(R_1)|/2 \geq |R|/4$  for every vertex  $x$  in  $G(R)$ . Therefore, by Proposition 3.9, the unit graph  $G(R)$  is Hamiltonian.

Now suppose that the lemma holds for  $n - 1$ . The induction hypothesis implies that the unit graph  $G(R_1 \times \cdots \times R_{n-1} \times \mathbb{Z}_2)$  is Hamiltonian. On the other hand,  $F \cong R_n/\mathfrak{m}_n$  is a field with  $|F| \geq 3$ . Therefore Lemma 3.5 implies that the unit graph  $G(R_1 \times \cdots \times R_{n-1} \times \mathbb{Z}_2 \times F)$  is Hamiltonian and so by applying Lemma 3.7 we conclude that the unit graph  $G(R)$  is Hamiltonian.  $\square$

A *cycle graph* is a graph that consists of a single cycle. The following result characterizes the unit graphs of rings that are cycle graphs.

**Proposition 3.11** [Ashrafi et al. 2010, Theorem 3.2]. *Let  $R$  be a ring. Then the unit graph  $G(R)$  is a cycle graph if and only if  $R$  is isomorphic to either*

- (a)  $\mathbb{Z}_4$ ,
- (b)  $\mathbb{Z}_6$ , or
- (c)  $\left\{ \begin{bmatrix} a & b \\ 0 & a \end{bmatrix} \mid a, b \in \mathbb{Z}_2 \right\}$ .

The next result gives a sufficient condition for a unit graph to be Hamiltonian.

**Lemma 3.12.** *Let  $R$  be a ring such that  $R \not\cong \mathbb{Z}_2$  and  $R \not\cong \mathbb{Z}_3$ . If  $R$  cannot have  $\mathbb{Z}_2 \times \mathbb{Z}_2$  as a quotient, then the unit graph  $G(R)$  is Hamiltonian.*

*Proof.* Every ring is isomorphic to a direct product of local rings; see [McDonald 1974, page 95]. Therefore we may write  $R \cong R_1 \times \cdots \times R_n$ , where every  $R_i$  is a local ring with maximal ideal  $\mathfrak{m}_i$ . We claim that  $|U(R)| \geq 2$ . To show this, suppose to the contrary that  $|U(R)| = 1$ . This implies that  $|J(R)| = 1$ , where  $J(R)$  denotes the Jacobson radical of  $R$ . Therefore  $|\mathfrak{m}_1 \times \cdots \times \mathfrak{m}_n| = 1$  and so  $|\mathfrak{m}_i| = 1$  for every  $i$  with  $1 \leq i \leq n$ . Therefore  $R_i$  for  $1 \leq i \leq n$  is a field and thus  $R \cong \mathbb{Z}_2 \times \cdots \times \mathbb{Z}_2$ , where  $\mathbb{Z}_2$  occurs  $n$  times in the product. Now the assumption implies that  $R \cong \mathbb{Z}_2$ , a contradiction. Thus the claim holds and we have  $|U(R)| \geq 2$ .

First, suppose  $|U(R)| = 2$ . In this case, by Proposition 3.2, the unit graph  $G(R)$  is a 2-regular connected graph and so is a cycle graph. Hence by Proposition 3.11,  $R$  is isomorphic to either  $\mathbb{Z}_4$ ,  $\mathbb{Z}_6$ , or  $\left\{ \begin{bmatrix} a & b \\ 0 & a \end{bmatrix} \mid a, b \in \mathbb{Z}_2 \right\}$ . It is easy to see that the unit graph of each of them is Hamiltonian and therefore so is the unit graph  $G(R)$ .

Second, suppose that  $|U(R)| \geq 3$ . By the assumption,  $R_i/\mathfrak{m}_i \not\cong \mathbb{Z}_2$  for every  $i$ , except for possibly at most one  $i$ . If  $R_i/\mathfrak{m}_i \not\cong \mathbb{Z}_2$  for every  $i$ , then by Lemma 3.8 the unit graph  $G(R)$  is Hamiltonian. If for one  $i$ , say  $n$ , we have  $R_n/\mathfrak{m}_n \cong \mathbb{Z}_2$ , then by Lemma 3.10 the unit graph  $G(R_1 \times \cdots \times R_n \times \mathbb{Z}_2)$  is Hamiltonian. Now by applying Lemma 3.7 we conclude that the unit graph  $G(R)$  is Hamiltonian.  $\square$

*Proof of Theorem 2.1.* (a) implies (b): By assumption, the unit graph  $G(R)$  is Hamiltonian and so it is obviously connected. Therefore, by [Ashrafi et al. 2010, Theorem 4.3], we have  $u(R) \leq \omega$ . This means that the ring  $R$  is generated by its units and thus by [Raphael 1974, Corollary 7] it cannot have  $\mathbb{Z}_2 \times \mathbb{Z}_2$  as a quotient.

(b) implies (a): This holds by Lemma 3.12.

(b) is equivalent to (c): This holds by [Raphael 1974, Corollary 7].

(c) is equivalent to (d): This is true by definition.

(d) is equivalent to (e): This holds by [Ashrafi et al. 2010, Theorem 4.3].  $\square$

### Acknowledgments

The authors would like to thank the referee for carefully reading the paper and for his/her suggestions. Part of this work was done while M. R. Pournaki visited the Delhi Center of the Indian Statistical Institute (ISID). He would like to thank the Academy of Sciences for the Developing World (TWAS) and ISID for sponsoring his visits to New Delhi in July and August 2007 and January 2010. Especially he thanks Professor Rajendra Bhatia for the hospitality enjoyed at ISID.

### References

- [Ashrafi and Vámos 2005] N. Ashrafi and P. Vámos, “On the unit sum number of some rings”, *Q. J. Math.* **56**:1 (2005), 1–12. MR 2005k:11220 Zbl 1100.11036
- [Ashrafi et al. 2010] N. Ashrafi, H. R. Maimani, M. R. Pournaki, and S. Yassemi, “Unit graphs associated with rings”, *Comm. Algebra* **38**:8 (2010), 2851–2871. MR 2730284 Zbl 05803773
- [Atiyah and Macdonald 1969] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley, Reading, MA, 1969. MR 39 #4129 Zbl 0175.03601
- [Chartrand and Oellermann 1993] G. Chartrand and O. R. Oellermann, *Applied and algorithmic graph theory*, McGraw-Hill, New York, 1993. MR 1211413
- [Dirac 1952] G. A. Dirac, “Some theorems on abstract graphs”, *Proc. London Math. Soc.* (3) **2** (1952), 69–81. MR 13,856e Zbl 0047.17001
- [Garey and Johnson 1979] M. R. Garey and D. S. Johnson, *Computers and intractability, A guide to the theory of NP-completeness*, W. H. Freeman, San Francisco, CA, 1979. MR 80g:68056 Zbl 0411.68039
- [Goldsmith et al. 1998] B. Goldsmith, S. Pabst, and A. Scott, “Unit sum numbers of rings and modules”, *Quart. J. Math. Oxford Ser. (2)* **49**:195 (1998), 331–344. MR 99i:16060 Zbl 0933.16035
- [Gould 2003] R. J. Gould, “Advances on the Hamiltonian problem—a survey”, *Graphs Combin.* **19**:1 (2003), 7–52. MR 2004a:05092 Zbl 1024.05057
- [Grimaldi 1990] R. P. Grimaldi, “Graphs from rings”, pp. 95–103 in *Proceedings of the Twentieth Southeastern Conference on Combinatorics, Graph Theory, and Computing* (Boca Raton, FL, 1989), vol. 71, edited by F. Hoffman et al., 1990. MR 90m:05122 Zbl 0747.05091
- [Maimani et al. 2010] H. R. Maimani, M. R. Pournaki, and S. Yassemi, “Weakly perfect graphs arising from rings”, *Glasg. Math. J.* **52**:3 (2010), 417–425. MR 2679902 Zbl 05799531
- [McDonald 1974] B. R. McDonald, *Finite rings with identity*, Pure and Applied Mathematics **28**, Marcel Dekker, New York, 1974. MR 50 #7245 Zbl 0294.16012
- [Ore 1960] O. Ore, “Note on Hamilton circuits”, *Amer. Math. Monthly* **67** (1960), 55. MR 22 #9454 Zbl 0089.39505

- [Raphael 1974] R. Raphael, "Rings which are generated by their units", *J. Algebra* **28** (1974), 199–205. MR 49 #7300 Zbl 0271.16013
- [Vámos 2005] P. Vámos, "2-good rings", *Q. J. Math.* **56**:3 (2005), 417–430. MR 2006e:16055 Zbl 1156.16303
- [West 1996] D. B. West, *Introduction to graph theory*, Prentice Hall, Upper Saddle River, NJ, 1996. MR 96i:05001 Zbl 0845.05001
- [Zelinsky 1954] D. Zelinsky, "Every linear transformation is a sum of nonsingular ones", *Proc. Amer. Math. Soc.* **5** (1954), 627–630. MR 16,8c Zbl 0056.11002

Received December 28, 2009. Revised July 14, 2010.

H. R. MAIMANI

MATHEMATICS SECTION, DEPARTMENT OF BASIC SCIENCES  
SHAHID RAJAEI TEACHER TRAINING UNIVERSITY  
P.O. BOX 16785-163  
TEHRAN  
IRAN

and

SCHOOL OF MATHEMATICS  
INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM)  
P.O. BOX 19395-5746  
TEHRAN  
IRAN  
maimani@ipm.ir

M. R. POURNAKI

DEPARTMENT OF MATHEMATICAL SCIENCES  
SHARIF UNIVERSITY OF TECHNOLOGY  
P.O. BOX 11155-9415  
TEHRAN  
IRAN

pournaki@ipm.ir

<http://math.ipm.ac.ir/pournaki/>

S. YASSEMI

SCHOOL OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE  
COLLEGE OF SCIENCE  
UNIVERSITY OF TEHRAN  
TEHRAN  
IRAN

and

SCHOOL OF MATHEMATICS  
INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM)  
P.O. BOX 19395-5746  
TEHRAN  
IRAN

yassemi@ipm.ir

<http://math.ipm.ac.ir/yassemi/>

# PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
chari@math.ucr.edu

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
long@math.ucsb.edu

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
popa@math.ucla.edu

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
finn@math.stanford.edu

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
jhlu@maths.hku.hk

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
qing@cats.ucsc.edu

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
liu@math.ucla.edu

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
merkurev@math.ucla.edu

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
jonr@math.ucla.edu

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [www.pjmath.org](http://www.pjmath.org) for submission instructions.

---

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 249 No. 2 February 2011

---

A gluing construction for prescribed mean curvature ADRIAN BUTSCHER	257
Large eigenvalues and concentration BRUNO COLBOIS and ALESSANDRO SAVO	271
Sur les conditions d'existence des faisceaux semi-stables sur les courbes multiples primitives JEAN-MARC DRÉZET	291
A quantitative estimate for quasiintegral points in orbits LIANG-CHUNG HSIA and JOSEPH H. SILVERMAN	321
Möbius isoparametric hypersurfaces with three distinct principal curvatures, II ZEJUN HU and SHUJIE ZHAI	343
Discrete Morse theory and Hopf bundles DMITRY N. KOZLOV	371
Regularity of canonical and deficiency modules for monomial ideals MANOJ KUMMINI and SATOSHI MURAI	377
$SL_2(\mathbb{C})$ -character variety of a hyperbolic link and regulator WEIPING LI and QINGXUE WANG	385
Hypergeometric evaluation identities and supercongruences LING LONG	405
Necessary and sufficient conditions for unit graphs to be Hamiltonian H. R. MAIMANI, M. R. POURNAKI and S. YASSEMI	419
Instability of the geodesic flow for the energy functional DOMENICO PERRONE	431
String structures and canonical 3-forms CORBETT REDDEN	447
Dual pairs and contragredients of irreducible representations BINYONG SUN	485
On the number of pairs of positive integers $x_1, x_2 \leq H$ such that $x_1x_2$ is a $k$ -th power DOYCHIN I. TOLEV	495
Correction to the article A Floer homology for exact contact embeddings KAI CIELIEBAK and URS ADRIAN FRAUENFELDER	509



0030-8730(201102)249:2;1-B