

*Pacific  
Journal of  
Mathematics*

**BURGHELEA–HALLER ANALYTIC TORSION  
FOR TWISTED DE RHAM COMPLEXES**

GUANGXIANG SU

Volume 250 No. 2

April 2011

## BURGHELEA–HALLER ANALYTIC TORSION FOR TWISTED DE RHAM COMPLEXES

GUANGXIANG SU

**We extend the Burghelea–Haller analytic torsion to the twisted de Rham complexes, and compare it with the twisted refined analytic torsion defined by Huang. Finally, we briefly discuss the Cappell–Miller analytic torsion.**

### 1. Introduction

Let  $E$  be a unitary flat vector bundle on a closed Riemannian manifold  $M$ . Ray and Singer [1971] defined an analytic torsion associated to  $(M, E)$  and proved that it does not depend on the Riemannian metric on  $M$ . Moreover, they conjectured that this analytic torsion coincides with the classical Reidemeister torsion defined using a triangulation on  $M$  (see [Milnor 1966]). This conjecture was later proved in two celebrated papers [Cheeger 1979; Müller 1978]. Müller [1993] generalized this result to the case when  $E$  is a unimodular flat vector bundle on  $M$ . Inspired by the considerations of Quillen [1985], Bismut and Zhang [1992] reformulated the above Cheeger–Müller theorem as an equality between the Reidemeister and Ray–Singer metrics defined on the determinant of cohomology, and proved an extension of it to the case of general flat vector bundle over  $M$ . The method used by Bismut and Zhang is different from that of Cheeger and Müller in that it makes use of a deformation by Morse functions introduced by Witten [1982] on the de Rham complex.

Braverman and Kappeler [2007b; 2007c; 2008] defined the refined analytic torsion for a flat vector bundle over an odd dimensional manifold and showed that it equals the Turaev torsion [1989] (see also [Farber and Turaev 2000]) up to multiplication by a complex number of absolute value one. Burghelea and Haller [2007; 2008], following a suggestion of Müller, defined a generalized analytic torsion associated to a nondegenerate symmetric bilinear form on a flat vector bundle over an arbitrary dimensional manifold and make an explicit conjecture between this generalized analytic torsion and the Turaev torsion. This conjecture was proved up to sign in [Burghelea and Haller 2010] and in full generality in [Su and Zhang 2008]. Cappell and Miller [2010] used non-self-adjoint Laplace operators to define

---

*MSC2000:* primary 58J52; secondary 19K56.

*Keywords:* analytic torsion, symmetric bilinear form, de Rham complex.

another complex-valued analytic torsion and used the method in [Su and Zhang 2008] to prove an extension of the Cheeger–Müller theorem.

Mathai and Wu [2008; 2010b] generalized the classical Ray–Singer analytic torsion to the twisted de Rham complex with an odd degree closed differential form  $H$ . In [Mathai and Wu 2010a], they defined and studied analytic torsion of  $\mathbb{Z}_2$ -graded elliptic complexes. Huang [2010a] generalized Braverman and Kappeler’s refined analytic torsion to the twisted de Rham complex, proved a duality theorem and compared it with the twisted Ray–Singer metric.

In this paper, supposing there exists a nondegenerate symmetric bilinear form on the flat vector bundle  $E$ , we generalize the Burghelea–Haller analytic torsion to the twisted de Rham complex. For the odd dimensional manifold, we also compare it with the twisted refined analytic torsion and the twisted Ray–Singer metric.

The rest of this paper is organized as follows. In Section 2, supposing there exists a  $\mathbb{Z}_2$ -graded nondegenerate symmetric bilinear form on a  $\mathbb{Z}_2$ -graded finite dimensional complex, we define a symmetric bilinear torsion on it. In Section 3, we generalize the Burghelea–Haller analytic torsion to the twisted de Rham complex. In Section 4, when the dimension of the manifold is odd, we show that the twisted Burghelea–Haller analytic torsion is independent of the Riemannian metric  $g$ , the symmetric bilinear form  $b$  and the representative  $H$  in the cohomology class  $[H]$ . In Section 5, we compare this new torsion with the twisted refined analytic torsion. In Section 6, we briefly discuss the Cappell–Miller analytic torsion on the twisted de Rham complex of an odd dimensional manifold.

**2. Symmetric bilinear torsion on a finite dimensional  $\mathbb{Z}_2$ -graded complex**

Consider a cochain complex

$$0 \longrightarrow C^0 \xrightarrow{d_0} C^1 \xrightarrow{d_1} \dots \xrightarrow{d_{n-1}} C^n \longrightarrow 0$$

of finite dimensional complex vector space. Set

$$C^{\bar{k}} = \bigoplus_{i = k \bmod 2} C^i, \quad k = 0, 1.$$

Let

$$(2-1) \quad (C^\bullet, d) : \dots \xrightarrow{d_1} C^{\bar{0}} \xrightarrow{d_{\bar{0}}} C^{\bar{1}} \xrightarrow{d_{\bar{1}}} C^{\bar{0}} \xrightarrow{d_{\bar{0}}} \dots$$

be a  $\mathbb{Z}_2$ -graded cochain complex. Denote its cohomology by  $H^{\bar{k}}$ ,  $k = 0, 1$ . Set

$$\det(C^\bullet, d) = \det C^{\bar{0}} \otimes (\det C^{\bar{1}})^{-1}, \quad \det(H^\bullet, d) = \det H^{\bar{0}} \otimes (\det H^{\bar{1}})^{-1}.$$

Then we have a canonical isomorphism between the determinant lines

$$(2-2) \quad \phi : \det(C^\bullet, d) \rightarrow \det(H^\bullet, d).$$

Suppose that there is a nondegenerate symmetric bilinear form on  $C^{\bar{k}}$ ,  $k = 0, 1$ . Then it induces a nondegenerate symmetric bilinear form  $b_{\det H^\bullet(C^\bullet, d)}$  on the determinant line  $\det(H^\bullet, d)$  via the isomorphism (2-2). Let  $d_k^\#$  be the adjoint of  $d_{\bar{k}}$  with respect to the nondegenerate symmetric bilinear form and define

$$\Delta_{b, \bar{k}} = d_{\bar{k}}^\# d_{\bar{k}} + d_{\bar{k}+1} d_{\bar{k}+1}^\#.$$

Let  $\lambda$  be the generalized eigenvalue of  $\Delta_{b, \bar{k}}$  and let  $C_b^{\bar{k}}(\lambda)$  be the generalized  $\lambda$ -eigenspace of  $\Delta_{b, \bar{k}}$ . Then we have a  $b$ -orthogonal decomposition

$$(2-3) \quad C^{\bar{k}} = \bigoplus_{\lambda} C_b^{\bar{k}}(\lambda)$$

and the inclusion  $C_b^{\bar{k}}(0) \rightarrow C^{\bar{k}}$  induces an isomorphism in cohomology. Particularly, we obtain a canonical isomorphism

$$(2-4) \quad \det H^\bullet(C_b^\bullet(0)) \cong \det H^\bullet(C^\bullet).$$

**Proposition 2.1.** *The following identity holds:*

$$(2-5) \quad b_{\det H^\bullet(C^\bullet, d)} = b_{\det H^\bullet(C_b^\bullet(0), d)} \cdot \det(d_0^\# d_0 |_{C_b^{\bar{0}, \perp}(0) \cap \text{im } d_0^\#})^{-1} \cdot \det(d_1^\# d_1 |_{C_b^{\bar{1}, \perp}(0) \cap \text{im } d_1^\#}),$$

where  $C_b^{\bar{k}, \perp}(0) = \bigoplus_{\lambda \neq 0} C_b^{\bar{k}}(\lambda)$ ,  $k = 0, 1$ .

*Proof.* Same as [Burghilea and Haller 2007, Lemma 3.3]. Suppose  $(C_1^\bullet, b_1)$  and  $(C_2^\bullet, b_2)$  are finite-dimensional  $\mathbb{Z}_2$ -graded complexes equipped with  $\mathbb{Z}_2$ -graded nondegenerate symmetric bilinear forms. Clearly,  $H^\bullet(C_1^\bullet \oplus C_2^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$  and we obtain a canonical isomorphism of determinant lines

$$\det H^\bullet(C_1^\bullet \oplus C_2^\bullet) = \det H^\bullet(C_1^\bullet) \otimes \det H^\bullet(C_2^\bullet).$$

Then we have

$$b_{\det H^\bullet(C_1^\bullet \oplus C_2^\bullet)} = b_{\det H^\bullet(C_1^\bullet)} \otimes b_{\det H^\bullet(C_2^\bullet)}.$$

In view of the  $b$ -orthogonal decomposition (2-3) we may therefore without loss of generality assume  $\ker \Delta_{b, \bar{k}} = 0$ ,  $k = 0, 1$ . Then by the lemma just cited we have

$$C^{\bar{k}} = \text{im } d_{\bar{k}+1} \oplus \text{im } d_{\bar{k}}^\#.$$

This decomposition is  $b$ -orthogonal and invariant under  $\Delta_b$ . Thus we have the exact complexes

$$\begin{aligned} 0 \longrightarrow C^{\bar{0}} \cap \text{im } d_0^\# &\xrightarrow{d_0} C^{\bar{1}} \cap \text{im } d_0 \longrightarrow 0, \\ 0 \longrightarrow C^{\bar{1}} \cap \text{im } d_1^\# &\xrightarrow{d_1} C^{\bar{0}} \cap \text{im } d_1 \longrightarrow 0. \end{aligned}$$

Then from [Burghilea and Haller 2007, Example 3.2], we get the proposition.  $\square$

### 3. Symmetric bilinear torsion on the twisted de Rham complexes

In this section, we suppose that there is a fiberwise nondegenerate symmetric bilinear form on  $E$ . Then we define a symmetric bilinear torsion on the determinant line of the twisted de Rham complex.

**Twisted de Rham complexes.** In this section, we review the twisted de Rham complexes from [Mathai and Wu 2008].

Let  $M$  be a closed Riemannian manifold and  $E \rightarrow M$  be a complex flat vector bundle with flat connection  $\nabla$ . Let  $H$  be an odd-degree closed differential form on  $M$ . We set  $\Omega^{\bar{0}} = \Omega^{\text{even}}(M, E)$ ,  $\Omega^{\bar{1}} = \Omega^{\text{odd}}(M, E)$  and  $\nabla^H = \nabla + H \wedge$ . We define the twisted de Rham cohomology groups as

$$H^{\bar{k}}(M, E, H) = \frac{\ker(\nabla^H : \Omega^{\bar{k}}(M, E) \rightarrow \Omega^{\overline{k+1}}(M, E))}{\text{im}(\nabla^H : \Omega^{\overline{k+1}}(M, E) \rightarrow \Omega^{\bar{k}}(M, E))}, \quad k = 0, 1.$$

Suppose  $H$  is replaced by  $H' = H - dB$  for some  $B \in \Omega^{\bar{0}}(M)$ , then there is an isomorphism  $\varepsilon_B = e^B \wedge \cdot : \Omega^\bullet(M, E) \rightarrow \Omega^\bullet(M, E)$  satisfying

$$\varepsilon_B \circ \nabla^H = \nabla^{H'} \circ \varepsilon_B.$$

Therefore  $\varepsilon_B$  induces an isomorphism

$$\varepsilon_B : H^\bullet(M, E, H) \rightarrow H^\bullet(M, E, H')$$

on the twisted de Rham cohomology.

**The construction of the symmetric bilinear torsion.** Suppose that there exists a nondegenerate symmetric bilinear form on  $E$ . To simplify notation, let  $C^{\bar{k}} = \Omega^{\bar{k}}(X, E)$  and let  $d_{\bar{k}} = d_{\bar{k}}^{E, H}$  be the operator  $\nabla^H$  acting on  $C^{\bar{k}}$  ( $k = 0, 1$ ). Then  $d_{\bar{1}}d_{\bar{0}} = d_{\bar{0}}d_{\bar{1}} = 0$  and we have a complex

$$(3-1) \quad \dots \xrightarrow{d_{\bar{1}}} C^{\bar{0}} \xrightarrow{d_{\bar{0}}} C^{\bar{1}} \xrightarrow{d_{\bar{1}}} C^{\bar{0}} \xrightarrow{d_{\bar{0}}} \dots$$

The metric  $g^M$  and the symmetric bilinear form  $b$  determine together a symmetric bilinear form on  $\Omega^\bullet(M, E)$  such that if  $u = \alpha f$ ,  $v = \beta g \in \Omega^\bullet(M, E)$  such that  $\alpha, \beta \in \Omega^\bullet(M)$ ,  $f, g \in \Gamma(E)$ , then

$$(3-2) \quad \beta_{g,b}(u, v) = \int_M (\alpha \wedge * \beta) b(f, g),$$

where  $*$  is the Hodge star operator. Denote by  $d_{\bar{k}}^\#$  the adjoint of  $d_{\bar{k}}$  with respect to the nondegenerate symmetric bilinear form (3-2). Then we define the Laplacians

$$\Delta_{b, \bar{k}} = d_{\bar{k}}^\# d_{\bar{k}} + d_{\overline{k+1}} d_{\overline{k+1}}^\#, \quad k = 0, 1.$$

If  $\lambda$  is in the spectrum of  $\Delta_{b,\bar{k}}$ , then the image of the associated spectral projection is finite dimensional and contains smooth forms only. Referring to this image as the (generalized)  $\lambda$ -eigenspace of  $\Delta_{b,\bar{k}}$  and denoting it by  $\Omega_{\{\lambda\}}^{\bar{k}}(M, E)$ , there exists  $N_\lambda \in \mathbb{N}$  such that

$$(\Delta_{b,\bar{k}} - \lambda)^{N_\lambda} |_{\Omega_{\{\lambda\}}^{\bar{k}}(M, E)} = 0.$$

Therefore for different generalized eigenvalues  $\lambda, \mu$ , the spaces  $\Omega_{\{\lambda\}}^{\bar{k}}(M, E)$  and  $\Omega_{\{\mu\}}^{\bar{k}}(M, E)$  are  $\beta_{g,b}$ -orthogonal.

For any  $a \geq 0$ , set

$$\Omega_{[0,a]}^{\bar{k}}(M, E) = \bigoplus_{0 \leq |\lambda| \leq a} \Omega_{\{\lambda\}}^{\bar{k}}(M, E).$$

Then  $\Omega_{[0,a]}^{\bar{k}}(M, E)$  is finite dimensional and one gets a nondegenerate symmetric bilinear form

$$b_{\det H^\bullet(\Omega_{[0,a]}^\bullet, d)} \quad \text{on} \quad \det H^\bullet(\Omega_{[0,a]}^\bullet, d).$$

Let  $\Omega_{(a,+\infty)}^{\bar{k}}(M, E)$  denote the  $\beta_{g,b}$ -orthogonal complement to  $\Omega_{[0,a]}^{\bar{k}}(M, E)$ .

For the subcomplexes  $(\Omega_{(a,+\infty)}^{\bar{k}+1}(M, E), d)$ , since the operators  $d_{\bar{k}} d_{\bar{k}}^\#$  and  $\Delta_{b,\overline{k+1}}$  are equal and invertible on  $\text{im}(d_{\bar{k}}) \cap \Omega_{(a,+\infty)}^{\bar{k}+1}(M, E)$ , we have

$$(3-3) \quad P_{\bar{k}} := d_{\bar{k}}^\# (d_{\bar{k}} d_{\bar{k}}^\#)^{-1} d_{\bar{k}} = d_{\bar{k}}^\# (\Delta_{b,\overline{k+1}})^{-1} d_{\bar{k}}$$

is a pseudodifferential operator of order 0 and satisfies

$$P_{\bar{k}}^2 = P_{\bar{k}}.$$

By definition we have

$$(3-4) \quad \begin{aligned} \zeta(s, d_{\bar{k}}^\# d_{\bar{k}} |_{\text{im } d_{\bar{k}}^\# \cap \Omega_{(a,+\infty)}^{\bar{k}}(M, E)}) &= \text{Tr}(\Delta_{b,\bar{k}}^{-s} P_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M, E)}) \\ &= \text{Tr}(P_{\bar{k}} \Delta_{b,\bar{k}}^{-s} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M, E)}). \end{aligned}$$

Then  $\zeta(s, d_{\bar{k}}^\# d_{\bar{k}} |_{\text{im } d_{\bar{k}}^\# \cap \Omega_{(a,+\infty)}^{\bar{k}}(M, E)})$  has a meromorphic extension to the whole complex plane and, by [Wodzicki 1984, Section 7], it is regular at 0. So by [Wodzicki 1984; Grubb and Seeley 1995], we have the following analogue of [Mathai and Wu 2008, Theorem 2.1].

**Theorem 3.1.** *For  $k = 0, 1$ ,  $\zeta(s, d_{\bar{k}}^\# d_{\bar{k}} |_{\text{im } d_{\bar{k}}^\# \cap \Omega_{(a,+\infty)}^\bullet(M, E)})$  is holomorphic in the half plane for  $\text{Re}(s) > n/2$  and extends meromorphically to  $\mathbb{C}$  with possible poles at  $\{(n-l)/2, l = 0, 1, 2, \dots\}$  only, and is holomorphic at  $s = 0$ .*

Then for  $k = 0, 1$  and any  $a \geq 0$ , the regularized zeta determinant

$$(3-5) \quad \det'(d_{\bar{k}}^\# d_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M, E)}) := \exp(-\zeta'(0, d_{\bar{k}}^\# d_{\bar{k}} |_{\text{im } d_{\bar{k}}^\# \cap \Omega_{(a,+\infty)}^{\bar{k}}(M, E)}).$$

is well defined.

**Proposition 3.2.** *The symmetric bilinear form on  $\det H^\bullet(\Omega^\bullet(M, E, H), d)$  given by*

$$(3-6) \quad b_{\det H^\bullet(\Omega^\bullet_{[0,a]}(M, E), d)} \cdot \det'(d_0^\# d_0^\# |_{\Omega^\bullet_{(a,+\infty)}(M, E)})^{-1} \cdot (\det'(d_1^\# d_1^\# |_{\Omega^\bullet_{(a,+\infty)}(M, E)}))$$

*is independent of the choice of  $a \geq 0$ .*

*Proof.* Let  $0 \leq a < c < \infty$ . We have

$$(3-7) \quad (\Omega^\bullet_{[0,c]}(M, E), d_{\bar{k}}) = (\Omega^\bullet_{[0,a]}(M, E), d_{\bar{k}}) \oplus (\Omega^\bullet_{(a,c]}(M, E), d_{\bar{k}}),$$

$$(3-8) \quad (\Omega^\bullet_{(a,+\infty)}(M, E), d_{\bar{k}}) = (\Omega^\bullet_{(a,c]}(M, E), d_{\bar{k}}) \oplus (\Omega^\bullet_{(c,+\infty)}(M, E), d_{\bar{k}}).$$

By the definition of the determinant,

$$(3-9) \quad \det'(d_{\bar{k}}^\# d_{\bar{k}}^\# |_{\Omega^\bullet_{(a,+\infty)}(M, E)}) = \det'(d_{\bar{k}}^\# d_{\bar{k}}^\# |_{\Omega^\bullet_{(a,c]}(M, E)}) \cdot \det'(d_{\bar{k}}^\# d_{\bar{k}}^\# |_{\Omega^\bullet_{(c,+\infty)}(M, E)}).$$

Applying Proposition 2.1 to (3-7),

$$b_{\det H^\bullet(\Omega^\bullet_{[0,c]})} = b_{\det H^\bullet(\Omega^\bullet_{[0,a]})} \cdot \det'(d_0^\# d_0^\# |_{\Omega^\bullet_{(a,c]}(M, E)})^{-1} \cdot (\det'(d_1^\# d_1^\# |_{\Omega^\bullet_{(a,c]}(M, E)})).$$

Then we get the proposition.  $\square$

**Definition 3.3.** The symmetric bilinear form defined by (3-6) is called the Ray–Singer symmetric bilinear torsion on  $\det H^\bullet(\Omega^\bullet(M, E, H), d)$  and is denoted by  $\tau_{b, \nabla, H}$ .

#### 4. Symmetric bilinear torsion under metric and flux deformations

In this section, we will use the methods in [Mathai and Wu 2008] to study the dependence of the torsion on the metric  $g$ , the symmetric bilinear form  $b$  and the flux  $H$ .

**Variation of the torsion with respect to the metric and symmetric bilinear form.**

We assume that  $M$  is a closed compact oriented manifold of odd dimension. Suppose the pair  $(g_u, b_u)$  is deformed smoothly along a one-parameter family with parameter  $u \in \mathbb{R}$ . Let  $Q_{\bar{k}}$  be the spectral projection onto  $\Omega^\bullet_{[0,a]}(M, E)$  and  $\Pi_{\bar{k}} = 1 - Q_{\bar{k}}$  be the spectral projection onto  $\Omega^\bullet_{(a,+\infty)}(M, E)$ . Let

$$\alpha = *_u^{-1} \frac{\partial *_u}{\partial u} + b_u^{-1} \frac{\partial b_u}{\partial u}.$$

**Lemma 4.1.** *Under the assumptions above,*

$$(4-1) \quad \frac{\partial}{\partial u} \log \left( \det'(d_0^\# d_0^\# |_{\Omega^\bullet_{(a,+\infty)}(M, E)})^{-1} \cdot (\det'(d_1^\# d_1^\# |_{\Omega^\bullet_{(a,+\infty)}(M, E)})) \right) \\ = - \sum_{k=0,1} (-1)^k \text{Tr}(\alpha Q_{\bar{k}}).$$

*Proof.* While  $d_{\bar{k}}$  is independent of  $u$ , we have

$$\frac{\partial d_{\bar{k}}^{\#}}{\partial u} = -[\alpha, d_{\bar{k}}^{\#}].$$

Using  $P_{\bar{k}} d_{\bar{k}}^{\#} = d_{\bar{k}}^{\#}$ ,  $d_{\bar{k}} P_{\bar{k}} = d_{\bar{k}}$  and  $P_{\bar{k}}^2 = P_{\bar{k}}$ , we get  $d_{\bar{k}}^{\#} d_{\bar{k}} P_{\bar{k}} = P_{\bar{k}} d_{\bar{k}}^{\#} d_{\bar{k}} = d_{\bar{k}}^{\#} d_{\bar{k}}$  and

$$P_{\bar{k}} \frac{\partial P_{\bar{k}}}{\partial u} P_{\bar{k}} = 0.$$

Following the  $\mathbb{Z}$ -graded case, we set

$$\begin{aligned} (4-2) \quad f(s, u) &= \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^{s-1} \operatorname{Tr}(e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}}^{\#} P_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M,E)}) dt \\ &= \Gamma(s) \sum_{k=0,1} (-1)^k \zeta(s, d_{\bar{k}}^{\#} d_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M,E)}). \end{aligned}$$

Using the above identities on  $P_{\bar{k}}$ , the trace property and by an application of Duhamel’s principal, we get

$$\begin{aligned} (4-3) \quad \frac{\partial f}{\partial u} &= \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^{s-1} \operatorname{Tr} \left( t[\alpha, d_{\bar{k}}^{\#}] d_{\bar{k}} e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}} \Pi_{\bar{k}} + e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}} \frac{\partial P_{\bar{k}}}{\partial u} P_{\bar{k}} \Pi_{\bar{k}} \right) dt \\ &= \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^{s-1} \operatorname{Tr} \left( t\alpha [d_{\bar{k}}^{\#}, d_{\bar{k}} e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}}] \Pi_{\bar{k}} + P_{\bar{k}} e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}} \frac{\partial P_{\bar{k}}}{\partial u} \Pi_{\bar{k}} \right) dt \\ &= \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^{s-1} \operatorname{Tr} \left( t\alpha (e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}} d_{\bar{k}}^{\#} d_{\bar{k}} - e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}} d_{\bar{k}}^{\#}) \Pi_{\bar{k}} \right. \\ &\quad \left. + e^{-t d_{\bar{k}}^{\#}} d_{\bar{k}} P_{\bar{k}} \frac{\partial P_{\bar{k}}}{\partial u} \Pi_{\bar{k}} \right) dt \\ &= \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^s \operatorname{Tr}(\alpha e^{-t \Delta_{b,\bar{k}}} \Delta_{b,\bar{k}} \Pi_{\bar{k}}) dt \\ &= - \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^s \frac{\partial}{\partial t} \operatorname{Tr}(\alpha (e^{-t \Delta_{b,\bar{k}}} \Pi_{\bar{k}})) dt. \end{aligned}$$

Integrating by parts, we have

$$\begin{aligned} (4-4) \quad \frac{\partial f}{\partial u} &= s \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^{s-1} \operatorname{Tr}(\alpha (e^{-t \Delta_{b,\bar{k}}} \Pi_{\bar{k}})) dt \\ &= s \sum_{k=0,1} (-1)^k \left( \int_0^1 + \int_1^{+\infty} \right) t^{s-1} \operatorname{Tr}(\alpha e^{-t \Delta_{b,\bar{k}}} (1 - Q_{\bar{k}})) dt. \end{aligned}$$

Since  $\alpha$  is a smooth tensor and  $n$  is odd, the asymptotic expansion as  $t \downarrow 0$  for  $\operatorname{Tr}(\alpha e^{-t \Delta_{b,\bar{k}}})$  does not contain a constant term. Therefore  $\int_0^1 t^{s-1} \operatorname{Tr}(\alpha e^{-t \Delta_{b,\bar{k}}}) dt$



does not have a pole at  $s = 0$ . On the other hand, because of the exponential decay of  $\text{Tr}(\alpha e^{-t\Delta_{b,\bar{k}}}\Pi_{\bar{k}})$  for large  $t$ ,

$$\int_1^{+\infty} t^{s-1} \text{Tr}(\alpha e^{-t\Delta_{b,\bar{k}}}\Pi_{\bar{k}})$$

is an entire function in  $s$ . So

$$(4-5) \quad \left. \frac{\partial f}{\partial u} \right|_{s=0} = -s \sum_{k=0,1} (-1)^k \int_0^1 t^{s-1} \text{Tr}(\alpha Q_{\bar{k}}) dt \Big|_{s=0} = - \sum_{k=0,1} (-1)^k \text{Tr}(\alpha Q_{\bar{k}})$$

and hence

$$(4-6) \quad \frac{\partial}{\partial u} \sum_{k=0,1} (-1)^k \zeta(0, d_{\bar{k}}^{\#} d_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}}(M, E)) = 0.$$

Finally, from (4-5), (4-6), we have

$$(4-7) \quad \det'(d_0^{\#} d_0 |_{\Omega_{(a,+\infty)}^{\bar{0}}}(M, E))^{-1} \cdot (\det'(d_1^{\#} d_1 |_{\Omega_{(a,+\infty)}^{\bar{1}}}(M, E))) \\ = \exp\left(\lim_{s \rightarrow 0} \left( f(s, u) - \frac{1}{s} \sum_{k=0,1} (-1)^k \zeta(0, d_{\bar{k}}^{\#} d_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}}(M, E)) \right)\right),$$

and the result follows. □

**Lemma 4.2.** *Under the same assumptions, along any one-parameter deformation of  $(g_u, b_u)$ , we have*

$$(4-8) \quad \left. \frac{\partial}{\partial w} \right|_u \left( \frac{b_{w, \det H^{\bullet}(\Omega_{[0,a]}^{\bullet}(M, E), d)}}{b_{u, \det H^{\bullet}(\Omega_{[0,a]}^{\bullet}(M, E), d)}} \right) = \sum_{k=0,1} (-1)^k \text{Tr}(\alpha Q_{\bar{k}}).$$

*Proof.* For sufficiently small  $w - u$ , the restriction of the spectral projection

$$Q_{\bar{k}} |_{\Omega_{u, [0,a]}^{\bar{k}}}(M, E) : \Omega_{u, [0,a]}^{\bar{k}}(M, E) \rightarrow \Omega_{w, [0,a]}^{\bar{k}}(M, E)$$

is an isomorphism of complexes. Then for sufficiently small  $w - u$ , we have

$$(4-9) \quad \frac{b_{w, \det H^{\bullet}(\Omega_{[0,a]}^{\bullet}(M, E), d)}}{b_{u, \det H^{\bullet}(\Omega_{[0,a]}^{\bullet}(M, E), d)}} \\ = \det((\beta_{g_u, b_u} |_{\Omega_{u, [0,a]}^{\bar{0}}}(M, E))^{-1} (Q_{\bar{0}} |_{\Omega_{u, [0,a]}^{\bar{0}}}(M, E))^* \\ \cdot (\beta_{g_w, b_w} |_{\Omega_{w, [0,a]}^{\bar{0}}}(M, E))) \cdot \det((\beta_{g_u, b_u} |_{\Omega_{u, [0,a]}^{\bar{1}}}(M, E))^{-1} (Q_{\bar{1}} |_{\Omega_{u, [0,a]}^{\bar{1}}}(M, E))^* \\ \cdot (\beta_{g_w, b_w} |_{\Omega_{w, [0,a]}^{\bar{1}}}(M, E)))^{-1}.$$

Then similarly to [Burghlea and Haller 2007], we get (4-8). □

Combining Lemma 4.1 and Lemma 4.2, we have:

**Theorem 4.3.** *Let  $M$  be a closed, compact manifold of odd dimension,  $E$  be a flat vector bundle over  $M$ , and  $H$  be a closed differential form on  $M$  of odd degree. Then the symmetric bilinear torsion  $\tau_{b,\nabla,H}$  on the twisted de Rham complex does not depend on the choices of the Riemannian metric on  $M$  and the symmetric bilinear form  $b$  in a same homotopy class of nondegenerate symmetric bilinear forms on  $E$ .*

**Variation of analytic torsion with respect to the flux in a cohomology class.** We continue to assume that  $\dim M$  is odd and use the same notation as above. Suppose the (real) flux form  $H$  is deformed smoothly along a one-parameter family with parameter  $v \in \mathbb{R}$  in such a way that the cohomology class  $[H] \in H^1(M, \mathbb{R})$  is fixed. Then  $\partial H/\partial v = -dB$  for some form  $B \in \Omega^0(M)$  that depends smoothly on  $v$ ; let

$$\beta = B \wedge \cdot.$$

**Lemma 4.4.** *Under the above assumptions,*

$$(4-10) \quad \frac{\partial}{\partial v} \log \left( \det'(d_0^\# d_0 |_{\Omega_{(a,+\infty)}^0(M,E)})^{-1} \cdot \left( \det'(d_1^\# d_1 |_{\Omega_{(a,+\infty)}^1(M,E)}) \right) \right) = -2 \sum_{k=0,1} (-1)^k \operatorname{Tr}(\beta Q_{\bar{k}}).$$

*Proof.* As in the proof of Lemma 4.1, we set

$$f(s, v) = \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^{s-1} \operatorname{Tr}(e^{-td_k^\# d_k} P_{\bar{k}} |_{\Omega_{(a,+\infty)}^k(M,E)}) dt.$$

We note that  $B$ , hence  $\beta$  is real. Using

$$\frac{\partial d_{\bar{k}}}{\partial v} = [\beta, d_{\bar{k}}], \quad \frac{\partial d_{\bar{k}}^\#}{\partial v} = -[\beta^\#, d_{\bar{k}}^\#], \quad P_{\bar{k}}^2 = P_{\bar{k}} = P_{\bar{k}}^\#, \quad P_{\bar{k}} \frac{\partial P_{\bar{k}}}{\partial v} P_{\bar{k}} = 0$$

and Duhamel’s principle, similarly to [Mathai and Wu 2008, Lemma 3.5], we get

$$(4-11) \quad \frac{\partial f}{\partial v} = -2 \sum_{k=0,1} (-1)^k \int_0^{+\infty} t^s \frac{\partial}{\partial t} \operatorname{Tr}(\beta e^{-t\Delta_{b,\bar{k}}} \Pi_{\bar{k}}) dt.$$

The rest is similar to the proof of Lemma 4.1. □

**Lemma 4.5.** *Under the same assumptions, along any one-parameter deformation of  $H$  that fixes the cohomology class  $[H]$ , we have*

$$(4-12) \quad \frac{\partial}{\partial w} \Big|_v \left( \frac{b_{\det H^\bullet(\Omega_{[0,a]}^\bullet(M,E,H^w),d)}}{b_{\det H^\bullet(\Omega_{[0,a]}^\bullet(M,E,H^v),d)}} \right) = 2 \sum_{k=0,1} (-1)^k \operatorname{Tr}(\beta Q_{\bar{k}}),$$

where we identify  $\det H^\bullet(M, E, H)$  along the deformation.

*Proof.* For sufficiently small  $w - v$ , we have

$$Q_{\bar{k}}\varepsilon_B : \Omega_{[0,a]}^{\bar{k}}(M, E, H^v) \rightarrow \Omega_{[0,a]}^{\bar{k}}(M, E, H^w)$$

is an isomorphism of complexes and the induced symmetric bilinear form on the determinant line  $\det H^\bullet(\Omega_{[0,a]}^\bullet(M, E, H^v), d)$  is

$$(4-13) \quad \left( (\det(Q_{\bar{k}}\varepsilon_B))^* b_{\det H^\bullet(\Omega_{[0,a]}^\bullet(M, E, H^w), d)} \right) (\cdot, \cdot) \\ = b_{\det H^\bullet(\Omega_{[0,a]}^\bullet(M, E, H^w), d)} (\det(Q_{\bar{k}}\varepsilon_B) \cdot, \det(Q_{\bar{k}}\varepsilon_B) \cdot),$$

where

$$\det(Q_{\bar{k}}\varepsilon_B) : \det H^\bullet(\Omega_{[0,a]}^*(M, E, H^v)) \rightarrow \det H^\bullet(\Omega_{[0,a]}^*(M, E, H^w))$$

is the induced isomorphism on the determinant lines. Then we can compare it with  $b_{\det H^\bullet(\Omega_{[0,a]}^\bullet(M, E, H^w), d)}$ , and similarly to [Mathai and Wu 2008, Lemma 3.7], we get (4-12).  $\square$

Combining Lemma 4.4 and Lemma 4.5, we have:

**Theorem 4.6.** *Let  $M$  be a closed, compact manifold of odd dimension,  $E$  be a flat vector bundle over  $M$ . Suppose  $H$  and  $H'$  are closed differential forms on  $M$  of odd degrees representing the same de Rham cohomology class, and let  $B$  be an even form so that  $H' = H - dB$ . Then the symmetric bilinear torsion satisfies  $(\det \varepsilon_B)^* \tau_{b, \nabla, H'} = \tau_{b, \nabla, H}$ .*

### 5. Compare with the refined analytic torsion

In this section, we will compare the symmetric bilinear torsion  $\tau_{b, \nabla, H}$  with the refined analytic torsion  $\rho_{\text{an}}(\nabla^H)$  defined in [Huang 2010a]. The main theorem of this section is the following.

**Theorem 5.1.** *Let  $M$  be a closed odd dimensional manifold,  $E$  be a complex vector bundle over  $M$  with connection  $\nabla$ ,  $H$  be a closed odd-degree differential form on  $M$ . Suppose there exists a nondegenerate symmetric bilinear form on  $E$ . Then*

$$(5-1) \quad \tau_{b, \nabla, H}(\rho_{\text{an}}(\nabla^H)) = \pm e^{-2\pi i(\eta(\nabla^H) - \text{rank } E \cdot \eta_{\text{trivial}})}.$$

(Here  $\eta(\nabla^H)$  and  $\eta_{\text{trivial}}$  are defined in [Huang 2010a].)

We will use the method in [Braverman and Kappeler 2007a] to prove the theorem and the proof will be given later.

Let  $h$  be a Hermitian metric on  $E$ . One can construct the Ray–Singer analytic torsion as an inner product on  $\det H^\bullet(M, E, H)$ , or equivalently as a metric on the determinant line; see [Huang 2010a, (6.13)]. We denote the resulting inner product by  $\tau_{h, \nabla, H}$ . Then by our Theorem 5.1 and Theorem 6.2 of the same reference, we get:

**Corollary 5.2.** *If  $\dim M$  is odd,  $\left| \frac{\tau_{b,\nabla,H}}{\tau_{h,\nabla,H}} \right| = 1$ .*

**The dual connection.** Let  $M$  be an odd dimensional closed manifold and  $E$  be a flat vector bundle over  $M$ , with flat connection  $\nabla$ . Assume that there exists a nondegenerate symmetric bilinear form  $b$  on  $E$ . The dual connection  $\nabla'$  to  $\nabla$  on  $E$  with respect to the form  $b$  is defined by the formula

$$db(u, v) = b(\nabla u, v) + b(u, \nabla' v), \quad u, v \in \Gamma(M, E).$$

We denote by  $E'$  the flat vector bundle  $(E, \nabla')$ .

**Choices of the metric and the spectral cut.** Until the end of this section we fix a Riemannian metric  $g$  on  $M$  and set  $\mathcal{B}^H = \mathcal{B}(\nabla^H, g) = \Gamma \nabla^H + \nabla^H \Gamma$  and  $\mathcal{B}'^H = \mathcal{B}'(\nabla'^H, g) = \Gamma \nabla'^H + \nabla'^H \Gamma$ , where  $\Gamma : \Omega^\bullet(M, E) \rightarrow \Omega^\bullet(M, E)$  is the chirality operator defined by

$$\Gamma \omega = i^{(n+1)/2} * (-1)^{q(q+1)/2} \omega, \quad \omega \in \Omega^q(M, E).$$

We also fix  $\theta \in (-\pi/2, 0)$  such that both  $\theta$  and  $\theta + \pi$  are Agmon angles for the odd signature operator  $\mathcal{B}^H$ . One easily checks that

$$(5-2) \quad (\nabla^H)^\# = \Gamma \nabla'^H \Gamma, \quad (\nabla'^H)^\# = \Gamma \nabla^H \Gamma, \quad \text{and} \quad (\mathcal{B}^H)^\# = \mathcal{B}'^H.$$

As  $\mathcal{B}^H$  and  $(\mathcal{B}^H)^\#$  have the same spectrum it then follows that

$$(5-3) \quad \eta(\mathcal{B}'^H) = \eta(\mathcal{B}^H) \quad \text{and} \quad \text{Det}_{\text{gr},\theta}(\mathcal{B}'^H) = \text{Det}_{\text{gr},\theta}(\mathcal{B}^H).$$

*Proof of Theorem 5.1.* The symmetric bilinear form  $\beta_{g,b}$  induces a nondegenerate symmetric bilinear form

$$H^j(M, E') \otimes H^{n-j}(M, E) \rightarrow \mathbb{C}, \quad j = 0, \dots, n,$$

and, hence, identifies  $H^j(M, E')$  with the dual space of  $H^{n-j}(M, E)$ . Using the construction of [Huang 2010a, Section 5.1] (with  $\tau : \mathbb{C} \rightarrow \mathbb{C}$  be the identity map) we obtain a linear isomorphism

$$(5-4) \quad \alpha : \det H^\bullet(M, E, H) \rightarrow \det H^\bullet(M, E', H).$$

**Lemma 5.3.** *Let  $E \rightarrow M$  be a complex vector bundle over a closed oriented odd dimensional manifold  $M$  endowed with a nondegenerate bilinear form  $b$  and let  $\nabla$  be a flat connection on  $E$ . Let  $\nabla'$  denote the connection dual to  $\nabla$  with respect to  $b$ . Let  $H$  be a closed odd-degree differential form on  $M$ . Then*

$$(5-5) \quad \alpha(\rho_{\text{an}}(\nabla^H)) = \rho_{\text{an}}(\nabla'^H).$$

The proof is that of [Huang 2010a, Theorem 5.3] and will be omitted. (Actually, it is simpler, since  $\mathcal{B}^H$  and  $\mathcal{B}'^H$  have the same spectrum, so there is no complex conjugation involved.)

For simplicity, we set

$$\tau_{b, \nabla, H, (a, +\infty)} = \det'(d_0^\# d_0^- |_{\Omega_{(a, +\infty)}^{\bar{0}}(M, E)})^{-1} \cdot (\det'(d_1^\# d_1^- |_{\Omega_{(a, +\infty)}^{\bar{1}}(M, E)})).$$

Setting  $\Delta'^H = (\nabla'^H)^\# \nabla'^H + \nabla'^H (\nabla'^H)^\#$ , we then have

$$\Delta'^H = \Gamma \Delta^H \Gamma.$$

**Lemma 5.4.**  $\tau_{b, \nabla, H, (a, +\infty)} = \tau_{b, \nabla', H, (a, +\infty)}$ .

*Proof.* Applying (5-2) and using the fact that

$$\nabla'^H : \Omega_{(a, +\infty)}^{\bar{k}}(M, E, H) \cap \text{im}(\nabla'^H)^\# \rightarrow \Omega_{(a, +\infty)}^{\overline{k+1}}(M, E, H) \cap \text{im} \nabla'^H$$

is an isomorphism, we get

$$\begin{aligned} (5-6) \quad \tau_{b, \nabla, H, (a, +\infty)} &= \prod_{k=0,1} \det'((\nabla^H)^\# \nabla^H |_{\Omega_{(a, +\infty)}^{\bar{k}}(M, E, H)})^{(-1)^{k+1}} \\ &= \prod_{k=0,1} \det'(\Gamma \nabla'^H (\nabla'^H)^\# \Gamma |_{\Omega_{(a, +\infty)}^{\bar{k}}(M, E, H)})^{(-1)^{k+1}} \\ &= \prod_{k=0,1} \det'(\nabla'^H (\nabla'^H)^\# |_{\Omega_{(a, +\infty)}^{\bar{k}}(M, E, H)})^{(-1)^k} \\ &= \prod_{k=0,1} \det'((\nabla'^H)^\# \nabla'^H |_{\Omega_{(a, +\infty)}^{\bar{k}}(M, E, H)})^{(-1)^{k+1}} = \tau_{b, \nabla', H}, \end{aligned}$$

which completes the proof. □

Then for any  $h \in \det H^\bullet(M, E, H)$ , we have

$$(5-7) \quad \tau_{b, \nabla, H}(h) = \tau_{b, \nabla', H}(\alpha(h)).$$

Hence, by (5-5) and (5-7),

$$(5-8) \quad \tau_{b, \nabla, H}(\rho_{\text{an}}(\nabla^H)) = \tau_{b, \nabla', H}(\rho_{\text{an}} \nabla'^H).$$

Let

$$\tilde{\nabla} = \begin{pmatrix} \nabla & 0 \\ 0 & \nabla' \end{pmatrix}, \quad \tilde{\nabla}^H = \begin{pmatrix} \nabla^H & 0 \\ 0 & \nabla'^H \end{pmatrix}.$$

Then, for any  $a \geq 0$ ,

$$\begin{aligned} \tau_{b, \tilde{\nabla}, H, (a, +\infty)} &= \tau_{b, \nabla, H, (a, +\infty)} \cdot \tau_{b, \nabla', H, (a, +\infty)}, \\ \tau_{b, \tilde{\nabla}, H}(\rho_{\text{an}}(\tilde{\nabla}^H)) &= \tau_{b, \nabla, H}(\rho_{\text{an}}(\nabla^H)) \cdot \tau_{b, \nabla', H}(\rho_{\text{an}}(\nabla'^H)). \end{aligned}$$

Combining the latter equality with (5-8) shows that

$$\tau_{b, \tilde{\nabla}, H}(\rho_{\text{an}}(\tilde{\nabla}^H)) = \tau_{b, \nabla, H}(\rho_{\text{an}}(\nabla^H))^2.$$

Hence, (5-1) is equivalent to the equality

$$(5-9) \quad \tau_{b, \tilde{\nabla}, H}(\rho_{\text{an}}(\tilde{\nabla}^H)) = e^{-4\pi i(\eta(\nabla^H) - \text{rank } E \cdot \eta_{\text{trivial}})}.$$

By a slight modification of the deformation argument in [Braverman and Kappeler 2007a, Section 4.7] where the untwisted case was treated, we obtain (5-9). This concludes the proof of Theorem 5.1.  $\square$

### 6. On the Cappell–Miller analytic torsion

In this section, we briefly discuss the extension of the Cappell–Miller analytic torsion to the twisted de Rham complexes. Let  $\dim M$  be odd.

In the notation above, we have the twisted de Rham complex  $\nabla^H : \Omega^{\bar{k}}(M, E) \rightarrow \Omega^{\bar{k}+1}(M, E)$  and the chirality operator  $\Gamma : \Omega^{\bar{k}}(M, E) \rightarrow \Omega^{\bar{k}+1}(M, E)$ ,  $k = 0, 1$ . Define

$$d_{\bar{k}}^b = \Gamma d_{\bar{k}} \Gamma : \Omega^{\bar{k}}(M, E) \rightarrow \Omega^{\bar{k}+1}(M, E).$$

Then consider the non-self-adjoint Laplacian

$$\Delta_{\bar{k}}^b = (d_{\bar{k}}^b + d_{\bar{k}}^b)^2 : \Omega^{\bar{k}}(M, E) \rightarrow \Omega^{\bar{k}}(M, E).$$

For any  $a \geq 0$ , let  $\Omega_{[0,a]}^{b,\bar{k}}(M, E)$  ( $\Omega_{(a,+\infty)}^{b,\bar{k}}(M, E)$ ) denote the span in  $\Omega^{\bar{k}}(M, E)$  of the generalized eigensolutions of  $\Delta_{\bar{k}}^b$  with generalized eigenvalues with absolute value in  $[0, a]$  ( $(a, +\infty)$ ). Then we have the decomposition of the complex

$$(\Omega^\bullet(M, E), d) = (\Omega_{[0,a]}^{b,\bullet}(M, E), d) \oplus (\Omega_{(a,+\infty)}^{b,\bullet}(M, E), d).$$

The subcomplex  $(\Omega_{[0,a]}^{b,\bullet}(M, E), d)$  is a  $\mathbb{Z}_2$ -graded finite dimensional complex. Then we can define the torsion element  $\rho_{\Gamma_{[0,a]}}^b \otimes \rho_{\Gamma_{[0,a]}}^b \in \det H^\bullet(\Omega_{[0,a]}^{b,\bullet}(M, E), d)^2 \cong \det H^\bullet(M, E, H)^2$ , where  $\rho_{\Gamma_{[0,a]}}^b$  defined by [Huang 2010a, (2.22)]. On the other hand, for the subcomplex  $(\Omega_{(a,+\infty)}^{b,\bullet}(M, E), d)$ , the following zeta-regularized determinant is well defined (see (3-5)):

$$(6-1) \quad \det'(d_{\bar{k}}^b d_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M, E)}) := \exp(-\zeta'(0, d_{\bar{k}}^b d_{\bar{k}} |_{\lim d_{\bar{k}}^b \cap \Omega_{(a,+\infty)}^{b,\bar{k}}(M, E)})).$$

Considering the square of the graded determinant defined in [Huang 2010a, (2.38)], for the  $\mathbb{Z}_2$ -graded finite dimensional complex  $\Omega_{(a,c]}^{b,\bullet}(M, E)$ ,  $0 \leq a < c < \infty$ , we find that

$$\det'(d_0^b d_0^b |_{\Omega_{(a,c]}^{b,0}(M, E)}) \cdot \det'(d_1^b d_1^b |_{\Omega_{(a,c]}^{b,1}(M, E)})^{-1} = (\text{Det}_{\text{gr}}(\mathcal{B}_0 |_{\Omega_{(a,c]}^{b,\bullet}(M, E)}))^2.$$

Then by [Huang 2010a, Proposition 2.7], we easily get:

**Proposition 6.1.** *The torsion element defined by*

$$(6-2) \quad \rho_{\Gamma_{[0,a]}}^b \otimes \rho_{\Gamma_{[0,a]}}^b \cdot \prod_{k=0,1} (\det'(d_{\bar{k}}^b d_{\bar{k}} |_{\Omega_{(a,+\infty)}^{\bar{k}}(M,E)}))^{(-1)^k} \in \det H^\bullet(M, E, H)^2$$

*is independent of the choice of  $a \geq 0$ .*

**Definition 6.2.** The torsion element in  $\det H^\bullet(M, E, H)^2$  defined by (6-2) is called the twisted Cappell–Miller analytic torsion for the twisted de Rham complex and is denoted by  $\tau_{\nabla,H}$ .

Next we study the torsion  $\tau_{\nabla,H}$  under metric and flux deformations. Since the methods are the same as the cases in the twisted refined analytic torsion [Huang 2010a] and the twisted Burghelea–Haller analytic torsion above, we only briefly outline the results.

**Theorem 6.3** (Metric independence). *Let  $M$  be a closed odd dimensional manifold,  $E$  be a complex vector bundle over  $M$  with flat connection  $\nabla$  and  $H$  be a closed odd-degree differential form on  $M$ . Then the torsion  $\tau_{\nabla,H}$  is independent of the choice of the Riemannian metric  $g$ .*

*Proof.* By the definition of  $\tau_{\nabla,H}$  and the observation on the determinants, this theorem follows easily from Proposition 2.4 and Equations (3.18) and (4.14) of [Huang 2010a]. □

**Theorem 6.4** (Flux representative independence). *Let  $M$  be a closed odd dimensional manifold and  $E$  be a complex vector bundle over  $M$  with flat connection  $\nabla$ . Suppose  $H$  and  $H'$  are closed differential forms on  $M$  of odd degrees representing the same de Rham cohomology class, and let  $B$  be an even form so that  $H' = H - dB$ . Then we have  $\tau_{\nabla,H'} = \det(\varepsilon_B) \tau_{\nabla,H}$ .*

*Proof.* From the above observation, this follows easily from Lemmas 4.6 and 4.7 of [Huang 2010a]. □

From the definition in (6-2), we see that the twisted Cappell–Miller analytic torsion is closely related to the twisted refined analytic torsion  $\rho_{\text{an}}(\nabla^H)$ . Explicitly:

**Theorem 6.5** (compare [Huang 2010b, Theorem 4.5]). *In  $\det H^\bullet(M, E, H)^2$ ,*

$$(6-3) \quad \rho_{\text{an}}(\nabla^H) \otimes \rho_{\text{an}}(\nabla^H) = \tau_{\nabla,H} e^{-2\pi i(\eta(\nabla^H) - \text{rank } E \cdot \eta_{\text{trivial}})}.$$

*Proof.* The twisted refined analytic torsion [Huang 2010a, (4.15)] is defined by

$$\rho_{\text{an}}(\nabla^H) = \text{Det}_{\text{gr},\theta}(\mathcal{B}_{0,(\lambda,\infty)}^H) \cdot \rho_{\Gamma_{[0,\lambda]}} \cdot e^{i\pi(\text{rank } E)\eta_{\text{trivial}}}.$$

By [Huang 2010a, (5.31)], we have

$$(6-4) \quad \rho_{\text{an}}(\nabla^H) \otimes \rho_{\text{an}}(\nabla^H) \\ = \rho_{\Gamma_{[0,\lambda]}} \otimes \rho_{\Gamma_{[0,\lambda]}} \cdot \exp(2\xi_\lambda(\nabla^H, g^M, \theta)) \\ \cdot \exp\left(-2i\pi\eta_\lambda(\nabla^H) - i\pi \sum_{k=0,1} (-1)^k d_{\bar{k},\lambda}^- + 2i\pi(\text{rank } E)\eta_{\text{trivial}}\right),$$

where  $\eta_\lambda(\nabla^H)$ ,  $\xi_\lambda(\nabla^H, g^M, \theta)$ , and  $d_{\bar{k},\lambda}^-$  are defined in equations (3.17), (3.18), and (3.19) of [Huang 2010a]. By (6-2) and (6-4), we find that

$$(6-5) \quad \rho_{\text{an}}(\nabla^H) \otimes \rho_{\text{an}}(\nabla^H) \\ = \tau_{\nabla, H} \exp(-2i\pi\eta_\lambda(\nabla^H) - i\pi \sum_{k=0,1} (-1)^k d_{\bar{k},\lambda}^- + 2i\pi(\text{rank } E)\eta_{\text{trivial}}).$$

From [Huang 2010a, (5.28)], we get

$$(6-6) \quad 2\eta_\lambda(\nabla^H) + \sum_{k=0,1} (-1)^k d_{\bar{k},\lambda}^- \equiv 2\eta(\nabla^H) \pmod{2\mathbb{Z}}.$$

Then (6-5) and (6-6) imply (6-3).  $\square$

Theorem 5.1 and Theorem 6.5 give the relation between the twisted Burghelea–Haller analytic torsion  $\tau_{b,\nabla,H}$  and the twisted Cappell–Miller analytic torsion  $\tau_{\nabla,H}$  if there is a nondegenerate symmetric bilinear form on the bundle  $E$ .

**Corollary 6.6.** *If there is a nondegenerate symmetric bilinear form on  $E$  and  $\dim M$  is odd, we have*

$$\tau_{b,\nabla,H}(\tau_{\nabla,H}) = \pm 1.$$

**Remark 6.7.** Almost at the same time of the preprint [Su 2010] of this paper, Huang [2010b] defined and studied the twisted Cappell–Miller torsion both for holomorphic and analytic cases.

## References

- [Bismut and Zhang 1992] J.-M. Bismut and W. Zhang, *An extension of a theorem by Cheeger and Müller*, *Astérisque* **205**, 1992. With an appendix by F. Laudenbach. MR 93j:58138 Zbl 0781.58039
- [Braverman and Kappeler 2007a] M. Braverman and T. Kappeler, “Comparison of the refined analytic and the Burghelea–Haller torsions”, *Ann. Inst. Fourier (Grenoble)* **57**:7 (2007), 2361–2387. Festival Yves Colin de Verdière. MR 2009f:58051 Zbl 1147.58033
- [Braverman and Kappeler 2007b] M. Braverman and T. Kappeler, “Ray–Singer type theorem for the refined analytic torsion”, *J. Funct. Anal.* **243**:1 (2007), 232–256. MR 2008e:58042 Zbl 1122.58018
- [Braverman and Kappeler 2007c] M. Braverman and T. Kappeler, “Refined analytic torsion as an element of the determinant line”, *Geom. Topol.* **11** (2007), 139–213. MR 2008a:58031 Zbl 1135.58014
- [Braverman and Kappeler 2008] M. Braverman and T. Kappeler, “Refined analytic torsion”, *J. Differential Geom.* **78**:2 (2008), 193–267. MR 2009a:58041 Zbl 1147.58034



- [Burghelena and Haller 2007] D. Burghelena and S. Haller, “Complex-valued Ray–Singer torsion”, *J. Funct. Anal.* **248**:1 (2007), 27–78. MR 2008b:58035 Zbl 1131.58020
- [Burghelena and Haller 2008] D. Burghelena and S. Haller, “Torsion, as a function on the space of representations”, pp. 41–66 in *C\*-algebras and elliptic theory II*, edited by D. Burghelena et al., Trends Math., Birkhäuser, Basel, 2008. MR 2009i:58045 Zbl 1147.58035
- [Burghelena and Haller 2010] D. Burghelena and S. Haller, “Complex valued Ray–Singer torsion II”, *Math. Nachr.* **283**:10 (2010), 1372–1402. MR 2744135 Zbl 05815198
- [Cappell and Miller 2010] S. E. Cappell and E. Y. Miller, “Complex-valued analytic torsion for flat bundles and for holomorphic bundles with (1, 1) connections”, *Comm. Pure Appl. Math.* **63**:2 (2010), 133–202. MR 2011a:58061 Zbl 1201.58021
- [Cheeger 1979] J. Cheeger, “Analytic torsion and the heat equation”, *Ann. of Math. (2)* **109**:2 (1979), 259–322. MR 80j:58065a Zbl 0412.58026
- [Farber and Turaev 2000] M. Farber and V. Turaev, “Poincaré–Reidemeister metric, Euler structures, and torsion”, *J. Reine Angew. Math.* **520** (2000), 195–225. MR 2001m:58069 Zbl 0938.57020
- [Grubb and Seeley 1995] G. Grubb and R. T. Seeley, “Weakly parametric pseudodifferential operators and Atiyah–Patodi–Singer boundary problems”, *Invent. Math.* **121**:3 (1995), 481–529. MR 96k:58216 Zbl 0851.58043
- [Huang 2010a] R.-T. Huang, “Refined analytic torsion for twisted de Rham complexes”, preprint, 2010. arXiv 1001.0654
- [Huang 2010b] R.-T. Huang, “Twisted Cappell–Miller holomorphic and analytic torsions”, preprint, 2010. arXiv 1001.3917
- [Mathai and Wu 2008] V. Mathai and S. Wu, “Analytic torsion for twisted de Rham complexes”, preprint, 2008. arXiv 0810.4204
- [Mathai and Wu 2010a] V. Mathai and S. Wu, “Analytic torsion of  $\mathbb{Z}_2$ -graded elliptic complexes”, preprint, 2010. arXiv 1001.3212
- [Mathai and Wu 2010b] V. Mathai and S. Wu, “Twisted analytic torsion”, *Sci. China Math.* **53**:3 (2010), 555–563. MR 2608312 Zbl 1202.58019
- [Milnor 1966] J. Milnor, “Whitehead torsion”, *Bull. Amer. Math. Soc.* **72** (1966), 358–426. MR 33 #4922 Zbl 0147.23104
- [Müller 1978] W. Müller, “Analytic torsion and  $R$ -torsion of Riemannian manifolds”, *Adv. in Math.* **28**:3 (1978), 233–305. MR 80j:58065b Zbl 0395.57011
- [Müller 1993] W. Müller, “Analytic torsion and  $R$ -torsion for unimodular representations”, *J. Amer. Math. Soc.* **6**:3 (1993), 721–753. MR 93m:58119 Zbl 0789.58071
- [Quillen 1985] D. Quillen, “Determinants of Cauchy–Riemann operators on Riemann surfaces”, *Funct. Anal. Appl.* **14**:1 (1985), 31–34. MR 86g:32035 Zbl 0603.32016
- [Ray and Singer 1971] D. B. Ray and I. M. Singer, “ $R$ -torsion and the Laplacian on Riemannian manifolds”, *Advances in Math.* **7** (1971), 145–210. MR 45 #4447 Zbl 0239.58014
- [Su 2010] G. Su, “Burghelena–Haller analytic torsion for twisted de Rham complexes”, preprint, 2010. arXiv 1001.2728
- [Su and Zhang 2008] G. Su and W. Zhang, “A Cheeger–Müller theorem for symmetric bilinear torsions”, *Chin. Ann. Math. Ser. B* **29**:4 (2008), 385–424. MR 2009j:58053 Zbl 1152.58028
- [Turaev 1989] V. G. Turaev, “Euler structures, nonsingular vector fields, and Reidemeister-type torsions”, *Izv. Akad. Nauk SSSR Ser. Mat.* **53**:3 (1989), 607–643, 672. In Russian. MR 90m:57021 Zbl 0692.57015

[Witten 1982] E. Witten, “Supersymmetry and Morse theory”, *J. Differential Geom.* **17**:4 (1982), 661–692 (1983). MR 84b:58111 Zbl 0499.53056

[Wodzicki 1984] M. Wodzicki, “Local invariants of spectral asymmetry”, *Invent. Math.* **75**:1 (1984), 143–177. MR 85g:58089 Zbl 0538.58038

Received January 25, 2010. Revised September 23, 2010.

GUANGXIANG SU  
CHERN INSTITUTE OF MATHEMATICS AND LPMC  
NANKAI UNIVERSITY  
TIANJIN 300071  
CHINA  
guangxiangsu@nankai.edu.cn

# PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
chari@math.ucr.edu

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
long@math.ucsb.edu

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
popa@math.ucla.edu

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
finn@math.stanford.edu

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
jhlu@maths.hku.hk

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
qing@cats.ucsc.edu

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
liu@math.ucla.edu

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
merkurev@math.ucla.edu

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
jonr@math.ucla.edu

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [www.pjmath.org](http://www.pjmath.org) for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic.

Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 250    No. 2    April 2011

---

Realizing profinite reduced special groups	257
VINCENT ASTIER and HUGO MARIANO	
On fibered commensurability	287
DANNY CALEGARI, HONGBIN SUN and SHICHENG WANG	
On an overdetermined elliptic problem	319
LAURENT HAUSWIRTH, FRÉDÉRIC HÉLEIN and FRANK PACARD	
Minimal sets of a recurrent discrete flow	335
HATTAB HAWETE	
Trace-positive polynomials	339
IGOR KLEP	
Remarks on the product of harmonic forms	353
LIVIU ORNEA and MIHAELA PILCA	
Steinberg representation of $\mathrm{GSp}(4)$ : Bessel models and integral representation of $L$ -functions	365
AMEYA PITALE	
An integral expression of the first nontrivial one-cocycle of the space of long knots in $\mathbb{R}^3$	407
KEIICHI SAKAI	
Burghelea–Haller analytic torsion for twisted de Rham complexes	421
GUANGXIANG SU	
$K(n)$ -localization of the $K(n+1)$ -local $E_{n+1}$ -Adams spectral sequences	439
TAKESHI TORII	
Thompson’s group is distorted in the Thompson–Stein groups	473
CLAIRE WLADIS	
Parabolic meromorphic functions	487
ZHENG JIAN-HUA	



0030-8730(201104)250:2;1-F