

*Pacific
Journal of
Mathematics*

**INEQUALITIES FOR THE NAVIER AND DIRICHLET
EIGENVALUES OF ELLIPTIC OPERATORS**

QIAOLING WANG AND CHANGYU XIA

Volume 251 No. 1

May 2011

INEQUALITIES FOR THE NAVIER AND DIRICHLET EIGENVALUES OF ELLIPTIC OPERATORS

QIAOLING WANG AND CHANGYU XIA

This paper studies eigenvalues of elliptic operators on a bounded domain in a Euclidean space. We obtain lower bounds for the eigenvalues of elliptic operators of higher orders with Navier boundary condition. We also prove lower bounds and universal inequalities of Payne–Pólya–Weinberger–Yang type for the eigenvalues of second order elliptic equations in divergence form with Dirichlet boundary condition.

1. Introduction

Let Ω be a bounded domain in an n -dimensional Euclidean space \mathbb{R}^n ($n \geq 2$) with smooth boundary $\partial\Omega$. Let Δ be the Laplacian of \mathbb{R}^n and consider the fixed membrane or Dirichlet eigenvalue problem

$$(1-1) \quad \begin{cases} \Delta u = -\lambda u & \text{in } \Omega, \\ u|_{\partial\Omega} = 0. \end{cases}$$

Let

$$0 < \lambda_1 < \lambda_2 \leq \dots \rightarrow \infty$$

denote the eigenvalues (repeated with multiplicity) of the problem (1-1). Weyl's asymptotic formula [1912] tells us that

$$(1-2) \quad \lambda_k \sim C(n) \left(\frac{k}{|\Omega|} \right)^{2/n} \quad \text{as } k \rightarrow \infty,$$

where $|\Omega|$ is the volume of Ω and $C(n) = (2\pi)^2 \omega_n^{-2/n}$ with ω_n being the volume of the unit ball in \mathbb{R}^n . Pólya [1961] showed that for any “plane covering domain” Ω in \mathbb{R}^2 (those that tile \mathbb{R}^2) this asymptotic relation is a one-sided inequality (his proof

Wang was partially supported by CNPq. Xia was partially supported by CNPq and FAPDF.

MSC2000: primary 35P15, 53C20, 53C42; secondary 35P15, 53C42.

Keywords: eigenvalue, lower bound, universal inequality, elliptic operator, Navier boundary condition.

also works for \mathbb{R}^n -covering domains) and conjectured, for any domain $\Omega \subset \mathbb{R}^n$, the inequality

$$(1-3) \quad \lambda_k \geq C(n) \left(\frac{k}{|\Omega|} \right)^{2/n} \quad \text{for all } k.$$

Li and Yau [1983] showed the lower bound

$$(1-4) \quad \sum_{i=1}^k \lambda_i \geq \frac{nkC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n},$$

which yields an individual lower bound on λ_k in the form

$$(1-5) \quad \lambda_k \geq \frac{nC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n}.$$

Similar bounds for eigenvalues with Neumann boundary condition were proved in [Kröger 1992; 1994; Laptev 1997]. It was pointed out in [Laptev and Weidl 2000] that (1-4) also follows from an earlier result of Berezin [1972] by using the Legendre transformation. Melas [2003] gave an improvement of (1-4):

$$(1-6) \quad \sum_{i=1}^k \lambda_i \geq \frac{nkC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n} + d_n k \frac{|\Omega|}{I(\Omega)},$$

where the constant d_n depends only on the dimension and

$$I(\Omega) = \min_{a \in \mathbb{R}^n} \int_{\Omega} |x - a|^2 dx$$

is the “moment of inertia” of Ω .

In this paper, we study eigenvalues of elliptic operators of higher orders with Navier boundary condition and of second order elliptic equations in divergence form with Dirichlet boundary condition and prove lower bounds for them which are similar to the inequality (1-6). We will also prove universal inequalities of Yang type for the Dirichlet eigenvalues of second order equations in divergence form. The first two results concern eigenvalues with Navier boundary condition.

Theorem 1.1. *Let Ω be a bounded domain in \mathbb{R}^n and let l be a positive integer. Consider the eigenvalue problem*

$$(1-7) \quad \begin{cases} (-\Delta)^l u = \lambda u & \text{in } \Omega, \quad u \in C^\infty(\Omega), \\ u|_{\partial\Omega} = \Delta u|_{\partial\Omega} = \cdots = \Delta^{l-1} u|_{\partial\Omega} = 0. \end{cases}$$

Let

$$0 < \lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_k \leq \cdots \rightarrow \infty$$

be the eigenvalues of (1-7) and denote by μ_1, \dots, μ_n the first n nonzero eigenvalues of the Neumann problem

$$(1-8) \quad \begin{cases} -\Delta v = \mu v & \text{in } \Omega, \\ (\partial v / \partial \nu)|_{\partial \Omega} = 0, \end{cases}$$

where ν is the unit outward normal vector field along $\partial \Omega$. Then

$$(1-9) \quad \sum_{j=1}^k \lambda_j^{1/l} \geq \frac{nkC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n} + \frac{d(n)k}{\sum_{i=1}^n \mu_i^{-1}}.$$

Here $d(n)$ is a positive constant depending only on n .

Theorem 1.2. Let Ω be a bounded domain in \mathbb{R}^n and let l be a fixed positive integer. Let L be the elliptic operator given by

$$Lu = \sum_{m=1}^l a_m (-\Delta)^m u, \quad u \in C^\infty(\Omega),$$

where the a_m are constants with $a_m \geq 0$, $1 \leq m \leq l$, and $a_l = 1$. Consider the eigenvalue problem

$$(1-10) \quad \begin{cases} Lu = \Lambda u & \text{in } \Omega, \\ u|_{\partial \Omega} = \Delta u|_{\partial \Omega} = \dots = \Delta^{l-1} u|_{\partial \Omega} = 0. \end{cases}$$

Let

$$0 < \Lambda_1 \leq \Lambda_2 \leq \dots \leq \Lambda_k \leq \dots \rightarrow \infty$$

be the eigenvalues of (1-10). Then

$$(1-11) \quad \Lambda_k \geq \sum_{m=1}^l a_m \left(\frac{nC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n} + \frac{d(n)}{\sum_{i=1}^n \mu_i^{-1}} \right)^m,$$

where μ_1, \dots, μ_n are the first n nonzero Neumann eigenvalues of Ω and $d(n)$ is a positive constant depending on n .

Our next results are about second order equations in divergence form with Dirichlet boundary condition. Firstly, we have a Li-Yau type inequality.

Theorem 1.3. Let Ω be a bounded domain in \mathbb{R}^n and let V be a nonnegative continuous function on Ω . Consider the eigenvalue problem

$$(1-12) \quad \begin{cases} -\sum_{\alpha, \beta=1}^n \frac{\partial}{\partial x_\alpha} \left(a_{\alpha\beta}(x) \frac{\partial u}{\partial x_\beta} \right) + V(x)u = \lambda u & \text{in } \Omega, \\ u|_{\partial \Omega} = 0. \end{cases}$$

Assume that there is a positive number ξ_0 such that the symmetric matrix $[a_{\alpha\beta}]$ satisfies $[a_{\alpha\beta}] \geq \xi_0 I$ in the sense of quadratic forms throughout Ω , where I is the identity matrix of order n . Let

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots \rightarrow \infty$$

be the eigenvalues of (1-12). Then

$$(1-13) \quad \sum_{j=1}^k \lambda_j \geq \xi_0 k \left(\frac{nC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n} + \frac{d(n)}{\sum_{i=1}^n \mu_i^{-1}} + \frac{V_0}{\xi_0} \right).$$

Here $V_0 = \inf_{x \in \Omega} V(x)$, μ_1, \dots, μ_n and $d(n)$ are as in Theorem 1.1.

We then prove a universal inequality of Payne–Pólya–Weinberger–Yang type [Payne et al. 1956; Yang 1991] for an eigenvalue problem more general than (1-12).

Theorem 1.4. Let Ω be a connected bounded domain in \mathbb{R}^n and let V be a non-negative continuous function on Ω with $V_0 = \inf_{x \in \Omega} V(x)$. Let ρ be a continuous function on Ω satisfying $\rho_1 \leq \rho(x) \leq \rho_2$ for all $x \in \Omega$, for some positive constants ρ_1 and ρ_2 . Assume also that there are positive numbers ξ_1 and ξ_2 such that the symmetric matrix $[a_{\alpha\beta}]$ satisfies $[a_{\alpha\beta}] \geq \xi_1 I$ in the sense of quadratic forms and $\sum_{\alpha=1}^n a_{\alpha\alpha} \leq n\xi_2$ throughout Ω . Consider the eigenvalue problem

$$(1-14) \quad \begin{cases} -\sum_{\alpha,\beta=1}^n \frac{\partial}{\partial x_\alpha} \left(a_{\alpha\beta}(x) \frac{\partial u}{\partial x_\beta} \right) + V(x)u = \lambda \rho u & \text{in } \Omega, \\ u|_{\partial\Omega} = 0. \end{cases}$$

Let

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots \rightarrow \infty$$

be the eigenvalues of (1-14). Then

$$(1-15) \quad \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \leq \frac{4\xi_2 \rho_2^2}{n\xi_1 \rho_1^2} \sum_{i=1}^k (\lambda_{k+1} - \lambda_i) \left(\lambda_i - \frac{V_0}{\rho_2} \right).$$

Remark 1.5. Universal inequalities of Payne–Pólya–Weinberger–Yang type for eigenvalues of elliptic operators on Riemannian manifolds have been studied recently by many mathematicians. One can find various interesting results in this direction, for example, in [Ashbaugh 1999; 2002; Ashbaugh and Benguria 1993a; 1993b; Ashbaugh and Hermi 2004; Cheng and Yang 2005; 2006a; 2006b; 2006c; 2007; El Soufi et al. 2007; Harrell 1993; Harrell and Michel 1994; Harrell and Stubbe 1997; Harrell and Yıldırım Yolcu 2009; Hile and Protter 1980; Hook 1990; Laptev 1997; Levitin and Parnovski 2002; Sun et al. 2008; Wang and Xia 2007a; 2007b; 2008; 2010a; 2010b; 2010c; 2011].

2. An auxiliary result

Before proving 1.1–1.3, we show the following fact.

Theorem 2.1. *Let Ω be a bounded domain in \mathbb{R}^n and let $w_1, \dots, w_k : \Omega \rightarrow \mathbb{R}$ be smooth functions satisfying*

$$(2-1) \quad w_i|_{\partial\Omega} = 0 \quad \text{and} \quad \int_{\Omega} w_i(x)w_j(x) dx = \delta_{ij} \quad \text{for } i, j = 1, \dots, k.$$

Then

$$(2-2) \quad \sum_{j=1}^k \int_{\Omega} |\nabla w_j(x)|^2 dx \geq \frac{nkC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n} + \frac{d(n)k}{\sum_{i=1}^n \mu_i^{-1}},$$

where $d(n)$ is a computational positive constant depending only on n and the μ_i are the first n nonzero Neumann eigenvalues of the Laplacian of Ω .

Proof. Let v_1, \dots, v_n be orthonormal eigenfunctions corresponding to μ_1, \dots, μ_n :

$$-\Delta v_i = \mu_i v_i \quad \text{in } \Omega, \quad \frac{\partial v_i}{\partial \nu} \Big|_{\partial\Omega} = 0, \quad \int_{\Omega} v_i v_j = \delta_{ij} \quad \text{for } i, j = 1, \dots, n.$$

By a translation of the origin and a suitable rotation of axes, we can assume, using [Ashbaugh and Benguria 1993b, p. 563], that

$$\int_{\Omega} x_i dx = 0 \quad \text{for } i = 1, \dots, n,$$

$$\int_{\Omega} x_j v_i dx = 0 \quad \text{for } j = 2, \dots, n, i = 1, \dots, j - 1.$$

It then follows from inequality (2.8) in the same paper that

$$(2-3) \quad \sum_{i=1}^n \frac{1}{\mu_i} \geq \frac{\int_{\Omega} |x|^2 dx}{|\Omega|}.$$

By a simple rearrangement argument, we have

$$(2-4) \quad \int_{\Omega} |x|^2 dx \geq \frac{n}{n+2} |\Omega| \left(\frac{|\Omega|}{\omega_n} \right)^{2/n}.$$

Extend each w_i to \mathbb{R}^n by letting $w_i(x) = 0$ for $x \in \mathbb{R}^n \setminus \Omega$. For a function g on \mathbb{R}^n , we will denote by $\mathcal{F}(g)$ the Fourier transformation of g . For any $z \in \mathbb{R}^n$, we have, by definition,

$$(2-5) \quad \mathcal{F}(w_j)(z) = (2\pi)^{-n/2} \int_{\Omega} e^{-i\langle x, z \rangle} w_j(x) dx.$$

Since $\{w_j\}_{j=1}^k$ is an orthonormal set in $L^2(\Omega)$, the Bessel inequality gives

$$(2-6) \quad \sum_{j=1}^k |\mathcal{F}(w_j)(z)|^2 \leq (2\pi)^{-n} \int_{\Omega} |e^{-i\langle x, z \rangle}|^2 dx = (2\pi)^{-n} |\Omega|.$$

For each $q = 1, \dots, n$ and $j = 1, \dots, k$, since w_j vanishes on $\partial\Omega$, one gets from the divergence theorem that

$$(2-7) \quad \begin{aligned} z_q \mathcal{F}(w_j)(z) &= i(2\pi)^{-n/2} \int_{\Omega} \frac{\partial e^{-i\langle x, z \rangle}}{\partial x_q} w_j(x) dx \\ &= -i(2\pi)^{-n/2} \int_{\Omega} \frac{\partial w_j(x)}{\partial x_q} e^{-i\langle x, z \rangle} dx = -i \mathcal{F}\left(\frac{\partial w_j}{\partial z_q}\right)(z). \end{aligned}$$

It then follows from the Plancherel formula that

$$(2-8) \quad \begin{aligned} \int_{\mathbb{R}^n} |z|^2 |\mathcal{F}(w_j)(z)|^2 dz &= \int_{\mathbb{R}^n} \sum_{q=1}^n \left| \mathcal{F}\left(\frac{\partial w_j}{\partial z_q}\right)(z) \right|^2 dz \\ &= \int_{\Omega} \sum_{q=1}^n \left(\frac{\partial w_j}{\partial x_q} \right)^2 dx = \int_{\Omega} |\nabla w_j(x)|^2 dx. \end{aligned}$$

Since

$$\nabla(\mathcal{F}(w_j))(z) = (2\pi)^{-n/2} \int_{\Omega} (-ix e^{-i\langle x, z \rangle} w_j(x)) dx,$$

we have

$$(2-9) \quad \sum_{j=1}^k |\nabla(\mathcal{F}(w_j))(z)|^2 \leq (2\pi)^{-n} \int_{\Omega} |ix e^{-i\langle x, z \rangle}|^2 dx = (2\pi)^{-n} \int_{\Omega} |x|^2 dx.$$

Set

$$G(z) = \sum_{j=1}^k |\mathcal{F}(w_j)(z)|^2.$$

Then $0 \leq G(z) \leq (2\pi)^{-n} |\Omega|$ and

$$(2-10) \quad \begin{aligned} |\nabla G(z)| &\leq 2 \left(\sum_{j=1}^k |\mathcal{F}(w_j)(z)|^2 \right)^{1/2} \left(\sum_{j=1}^k |\nabla(\mathcal{F}(w_j))(z)|^2 \right)^{1/2} \\ &\leq 2(2\pi)^{-n} \left(|\Omega| \int_{\Omega} |x|^2 dx \right)^{1/2} \end{aligned}$$

for every $z \in \mathbb{R}^n$. We also have

$$(2-11) \quad \int_{\mathbb{R}^n} G(z) dz = \sum_{j=1}^k \int_{\Omega} w_j(x)^2 dx = k.$$

Let $G^*(z) = g(|z|)$ be the decreasing spherical rearrangement of G . By approximation, we may assume that $g : [0, +\infty) \rightarrow [0, (2\pi)^{-n}|\Omega|]$ is absolutely continuous. Setting $\alpha(t) = |\{G^* > t\}| = |\{G > t\}|$, we have

$$(2-12) \quad \alpha(g(s)) = \omega_n s^n,$$

which implies that $n\omega_n s^{n-1} = \alpha'(g(s))g'(s)$ for almost every s . The coarea formula [Chavel 1984] tells us that

$$(2-13) \quad \alpha'(t) = - \int_{\{G=t\}} \frac{1}{|\nabla G|} dA_t.$$

Set $\eta = 2(2\pi)^{-n}(|\Omega| \int_{\Omega} |x|^2 dx)^{1/2}$; then one infers from (2-10) and the isoperimetric inequality that

$$(2-14) \quad -\alpha'(g(s)) \geq \eta^{-1} \text{area}(\{G = g(s)\}) \geq \eta^{-1} n\omega_n s^{n-1},$$

and so

$$(2-15) \quad -\eta \leq g'(s) \leq 0$$

for almost every s . It follows from (2-11) that

$$(2-16) \quad k = \int_{\mathbb{R}^n} G(z) dz = \int_{\mathbb{R}^n} G^*(z) dz = n\omega_n \int_0^\infty s^{n-1} g(s) ds$$

and, by (2-8),

$$(2-17) \quad \begin{aligned} \sum_{j=1}^k \int_{\Omega} |\nabla w_j|^2 &= \int_{\mathbb{R}^n} |z|^2 G(z) dz \geq \int_{\mathbb{R}^n} |z|^2 G^*(z) dz \\ &= n\omega_n \int_0^\infty s^{n+1} g(s) ds, \end{aligned}$$

since $z \rightarrow |z|^2$ is radial and increasing.

We next apply the following lemma to the function g , with $A = \frac{k}{n\omega_n}$ and

$$\eta = 2(2\pi)^{-n} \left(|\Omega| \int_{\Omega} |x|^2 dx \right)^{1/2} :$$

Lemma [Melas 2003]. *Let $n \geq 1$ and $\eta, A > 0$ and let $h : [0, +\infty) \rightarrow [0, +\infty)$ be a decreasing and absolutely continuous function such that*

$$-\eta \leq h'(s) \leq 0 \quad \text{and} \quad \int_0^\infty s^{n-1} h(s) ds = A.$$

Then

$$(2-18) \quad \int_0^\infty s^{n+1} h(s) ds \geq \frac{(nA)^{(n+2)/n}}{n+2} h(0)^{-2/n} + \frac{Ah(0)^2}{6(n+2)\eta^2}.$$

After applying the lemma and using (2-17), we infer that

$$(2-19) \quad \begin{aligned} \sum_{j=1}^k \int_{\Omega} |\nabla w_j(x)|^2 dx &\geq \frac{n}{n+2} \omega_n^{-2/n} k^{1+2/n} g(0)^{-2/n} + \frac{kg(0)^2}{6(n+2)\eta^2} \\ &\geq \frac{n}{n+2} \omega_n^{-2/n} k^{1+2/n} g(0)^{-2/n} + \frac{\tau kg(0)^2}{(n+2)\eta^2}, \end{aligned}$$

where τ is any constant with $0 < \tau \leq \frac{1}{6}$. From (2-4) we know that

$$(2-20) \quad \eta \geq 2(2\pi)^{-n} \left(\frac{n}{n+2} \right)^{1/2} \omega_n^{-1/n} |\Omega|^{(n+1)/n}.$$

Observe that $0 < g(0) \leq (2\pi)^{-n} |\Omega|$. Let $\tau = \tau(n)$ be the constant given by

$$\tau = \min \left\{ \frac{1}{6}, \frac{16\pi^2 n}{(n+2)\omega_n^{4/n}} \right\}.$$

Then one can see by using (2-20) that the function

$$\beta(t) = \frac{n}{n+2} \omega_n^{-2/n} k^{1+2/n} t^{-2/n} + \frac{\tau k t^2}{(n+2)\eta^2}$$

satisfies

$$\beta'((2\pi)^{-n} |\Omega|) \leq 0,$$

and so β is decreasing on $(0, (2\pi)^{-n} |\Omega|]$. Hence, choosing $d(n) = \frac{\tau}{4(n+2)}$, we have

$$(2-21) \quad \begin{aligned} \sum_{j=1}^k \int_{\Omega} |\nabla w_j(x)|^2 dx &\geq \beta(g(0)) \geq \beta((2\pi)^{-n} |\Omega|) \\ &= \frac{n}{n+2} \omega_n^{-2/n} k^{1+2/n} ((2\pi)^{-n} |\Omega|)^{-2/n} + \frac{\tau k ((2\pi)^{-n} |\Omega|)^2}{(n+2)\eta^2} \\ &= \frac{n}{n+2} \left(\frac{2\pi}{\omega_n^{1/n}} \right)^2 k^{1+2/n} |\Omega|^{-2/n} + \frac{d(n)k|\Omega|}{\int_{\Omega} |x|^2 dx}. \end{aligned}$$

Substituting (2-3) into (2-21), one gets (2-2). completing the proof of Theorem 2.1. \square

3. Proof of the main results

Proof of Theorem 1.1. Let $\{u_i\}_{i=1}^k$ be a set of orthonormal eigenfunctions corresponding to $\{\lambda_i\}_{i=1}^k$:

$$(-\Delta)^l u_i = \lambda_i u_i \quad \text{in } \Omega, \quad u_i|_{\partial\Omega} = \Delta u_i|_{\partial\Omega} = \cdots = \Delta^{l-1} u_i|_{\partial\Omega} = 0,$$

$$\int_{\Omega} u_i u_j = \delta_{ij} \quad \text{for } i, j = 1, \dots, k.$$

We show that for each $s = 1, \dots, l$ and $i = 1, \dots, k$,

$$(3-1) \quad 0 \leq \int_{\Omega} u_i (-\Delta)^s u_i \leq \lambda_i^{s/l}.$$

When $l = 1$, (3-1) holds trivially, so assume $l > 1$. when $s \in \{1, \dots, l\}$ is even, we have from the divergence theorem that

$$\int_{\Omega} u_i (-\Delta)^s u_i = \int_{\Omega} u_i \Delta^s u_i = \int_{\Omega} (\Delta^{s/2} u_i)^2 \geq 0.$$

On the other hand, if $s \in \{1, \dots, l\}$ is odd,

$$\begin{aligned} \int_{\Omega} u_i (-\Delta)^s u_i &= - \int_{\Omega} u_i \Delta^s u_i \\ &= - \int_{\Omega} \Delta^{(s-1)/2} u_i \Delta(\Delta^{(s-1)/2} u_i) = \int_{\Omega} |\nabla(\Delta^{(s-1)/2} u_i)|^2 \geq 0. \end{aligned}$$

Thus the first inequality in (3-1) holds.

We claim now that for any $s = 1, \dots, l - 1$,

$$(3-2) \quad \left(\int_{\Omega} u_i (-\Delta)^s u_i \right)^{s+1} \leq \left(\int_{\Omega} u_i (-\Delta)^{s+1} u_i \right)^s.$$

Since

$$\left(\int_{\Omega} u_i \Delta u_i \right)^2 \leq \int_{\Omega} u_i^2 \int_{\Omega} (\Delta u_i)^2 = \int_{\Omega} u_i \Delta^2 u_i,$$

we know that (3-2) holds when $s = 1$.

Suppose that (3-2) is true for $s - 1$, that is,

$$(3-3) \quad \left(\int_{\Omega} u_i (-\Delta)^{s-1} u_i \right)^s \leq \left(\int_{\Omega} u_i (-\Delta)^s u_i \right)^{s-1}.$$

When s is even, we have

$$\begin{aligned} \int_{\Omega} u_i (-\Delta)^s u_i &= \int_{\Omega} \Delta^{s/2-1} u_i \Delta(\Delta^{s/2} u_i) = - \int_{\Omega} \nabla(\Delta^{s/2-1} u_i) \nabla(\Delta^{s/2} u_i) \\ &\leq \left(\int_{\Omega} |\nabla(\Delta^{s/2-1} u_i)|^2 \right)^{1/2} \left(\int_{\Omega} |\nabla(\Delta^{s/2} u_i)|^2 \right)^{1/2} \\ &= \left(- \int_{\Omega} \Delta^{s/2-1} u_i \Delta^{s/2} u_i \right)^{1/2} \left(- \int_{\Omega} \Delta^{s/2} u_i \Delta^{s/2+1} u_i \right)^{1/2} \\ &= \left(\int_{\Omega} u_i (-\Delta)^{s-1} u_i \right)^{1/2} \left(\int_{\Omega} u_i (-\Delta)^{s+1} u_i \right)^{1/2}. \end{aligned}$$

On the other hand, when s is odd,

$$\begin{aligned} \int_{\Omega} u_i (-\Delta)^s u_i &= \int_{\Omega} (-\Delta)^{(s-1)/2} u_i (-\Delta)^{(s+1)/2} u_i \\ &\leq \left(\int_{\Omega} ((-\Delta)^{(s-1)/2} u_i)^2 \right)^{1/2} \left(\int_{\Omega} ((-\Delta)^{(s+1)/2} u_i)^2 \right)^{1/2} \\ &= \left(\int_{\Omega} u_i (-\Delta)^{s-1} u_i \right)^{1/2} \left(\int_{\Omega} u_i (-\Delta)^{s+1} u_i \right)^{1/2}. \end{aligned}$$

Thus we always have

$$(3-4) \quad \int_{\Omega} u_i (-\Delta)^s u_i \leq \left(\int_{\Omega} u_i (-\Delta)^{s-1} u_i \right)^{1/2} \left(\int_{\Omega} u_i (-\Delta)^{s+1} u_i \right)^{1/2}.$$

Substituting (3-3) into (3-4), we know that (3-2) is true for s . Using (3-2) repeatedly, we get

$$\int_{\Omega} u_i (-\Delta)^s u_i \leq \left(\int_{\Omega} u_i (-\Delta)^{s+1} u_i \right)^{s/(s+1)} \leq \cdots \leq \left(\int_{\Omega} u_i (-\Delta)^l u_i \right)^{s/l} = \lambda_i^{s/l}.$$

Thus the second inequality in (3-1) is also true. Consequently,

$$(3-5) \quad \sum_{j=1}^k \int_{\Omega} |\nabla u_j|^2 = \sum_{j=1}^k \int_{\Omega} u_j (-\Delta u_j) \leq \sum_{j=1}^k \lambda_j^{1/l}$$

which implies (1-9) by applying Theorem 2.1 to the functions u_1, \dots, u_k . This completes the proof of Theorem 1.1. \square

Proof of Theorem 1.2. Let $\{u_i\}_{i=1}^k$ be a set of orthonormal eigenfunctions of the problem (1-11) corresponding to $\{\lambda_i\}_{i=1}^k$:

$$\begin{aligned} Lu_i &= \Lambda u_i \quad \text{in } \Omega, & u_i|_{\partial\Omega} &= \Delta u_i|_{\partial\Omega} = \cdots = \Delta^{l-1} u_i|_{\partial\Omega} = 0, \\ \int_{\Omega} u_i u_j &= \delta_{ij} \quad \text{for } i, j = 1, \dots, k. \end{aligned}$$

Denote by $\{\lambda_i\}_{i=1}^k$ the first k fixed membrane eigenvalues of the Laplacian of Ω corresponding to the orthonormal eigenfunctions $\{v_i\}_{i=1}^k$:

$$-\Delta v_i = \lambda_i v_i \quad \text{in } \Omega, \quad v_i|_{\partial\Omega} = 0, \quad \int_{\Omega} v_i v_j = \delta_{ij} \quad \text{for } i, j = 1, \dots, k.$$

Let $w = \sum_{j=1}^k \alpha_j u_j \neq 0$ be such that

$$(3-6) \quad \int_{\Omega} w v_i = 0 \quad \text{for } i = 1, \dots, k-1.$$

Such an element w exists because $\{\alpha_j \mid 1 \leq j \leq k\}$ is a nontrivial solution of a system of $k-1$ linear equations

$$(3-7) \quad \sum_{j=1}^k \alpha_j \int_{\Omega} u_j v_i = 0, \quad 1 \leq i \leq k-1,$$

in k unknowns. Also assume without loss of generality that

$$(3-8) \quad \int_{\Omega} w^2 = 1.$$

Then it follows from the Rayleigh–Ritz inequality that

$$(3-9) \quad \lambda_k \leq \int_{\Omega} w(-\Delta w).$$

By using the arguments similar to those in the proof of (3-2), we have

$$(3-10) \quad \left(\int_{\Omega} w(-\Delta)^j w \right)^{j+1} \leq \left(\int_{\Omega} w(-\Delta)^{j+1} w \right)^j, \quad j = 1, \dots, l-1.$$

Hence

$$(3-11) \quad \int_{\Omega} w(-\Delta w) \leq \left(\int_{\Omega} w(-\Delta)^s w \right)^{1/s}, \quad s = 1, \dots, l,$$

which, combined with (3-9), gives

$$\lambda_k^s \leq \int_M w(-\Delta)^s w, \quad s = 1, 2, \dots, l.$$

Thus we have

$$(3-12) \quad \begin{aligned} & a_1 \lambda_k + a_2 \lambda_k^2 + \dots + a_{l-1} \lambda_k^{l-1} + \lambda_k^l \\ & \leq \int_{\Omega} w(a_1(-\Delta) + a_2(-\Delta)^2 + \dots + a_{l-1}(-\Delta)^{l-1} + (-\Delta)^l)w \\ & = \int_{\Omega} wLw = \sum_{i,j=1}^k \alpha_i \alpha_j \int_{\Omega} u_i L u_j = \sum_{i,j=1}^k \alpha_i \alpha_j \int_{\Omega} u_i \Lambda_j u_j \\ & = \sum_{i,j=1}^k \alpha_i \alpha_j \Lambda_j \delta_{ij} = \sum_{i=1}^k \alpha_i^2 \Lambda_i \leq \Lambda_k, \end{aligned}$$

where, in the last equality, we have used the fact that

$$\sum_{i=1}^k \alpha_i^2 = \int_{\Omega} w^2 = 1.$$

It is easy to see by taking $l = 1$ in (1-9) that

$$(3-13) \quad \lambda_k \geq \frac{nC(n)}{n+2} \left(\frac{k}{|\Omega|} \right)^{2/n} + \frac{d(n)}{\sum_{i=1}^n \mu_i^{-1}}.$$

Substituting (3-13) into (3-12), we get (1-11). Theorem 1.2 follows. □

Proof of Theorem 1.3. Let $\{u_i\}_{i=1}^k$ be a set of orthonormal eigenfunctions of the problem (1-12) corresponding to $\{\lambda_i\}_{i=1}^k$:

$$(3-14) \quad - \sum_{\alpha, \beta=1}^n \frac{\partial}{\partial x_\alpha} \left(a_{\alpha\beta}(x) \frac{\partial u_i}{\partial x_\beta} \right) + V(x)u_i = \lambda_i u_i \quad \text{in } \Omega,$$

$$(3-15) \quad u_i|_{\partial\Omega} = 0, \quad \int_{\Omega} u_i u_j = \delta_{ij} \quad \text{for } i, j = 1, \dots, k.$$

Multiplying (3-14) by u_i , integrating over Ω , and using the divergence theorem and the inequalities $V \geq V_0$ and $[a_{\alpha\beta}] \geq \xi_0 I$, we obtain

$$\begin{aligned} \lambda_i &= \int_{\Omega} \left(\sum_{\alpha, \beta=1}^n a_{\alpha\beta}(x) \frac{\partial u_i}{\partial x_\alpha} \frac{\partial u_i}{\partial x_\beta} + V(x)u_i^2 \right) \\ &\geq \int_{\Omega} \xi_0 \sum_{\alpha=1}^n \left(\frac{\partial u_i}{\partial x_\alpha} \right)^2 + V_0 \int_{\Omega} u_i^2 = \xi_0 \int_{\Omega} |\nabla u_i|^2 + V_0, \end{aligned}$$

which gives

$$(3-16) \quad \sum_{i=1}^k \int_{\Omega} |\nabla u_i|^2 \leq \frac{1}{\xi_0} \left(\sum_{i=1}^k \lambda_i - kV_0 \right).$$

Observing (3-15), one gets (1-13) by using Theorem 2.1 applied to u_1, \dots, u_k . Theorem 1.3 follows. □

Proof of Theorem 1.4. Let x_1, \dots, x_n be the coordinate functions on \mathbb{R}^n . For a function $f : \Omega \rightarrow \mathbb{R}$, set $f_{,\alpha} = \partial f / \partial x_\alpha$, $\alpha = 1, \dots, n$. Let $\{u_i\}_{i=1}^k$ be a set of orthonormal eigenfunctions of the problem (1-14) corresponding to $\{\lambda_i\}_{i=1}^k$:

$$\begin{aligned} - \sum_{\alpha, \beta=1}^n \frac{\partial}{\partial x_\alpha} \left(a_{\alpha\beta}(x) \frac{\partial u_i}{\partial x_\beta} \right) + V(x)u_i &= \lambda_i \rho u_i \quad \text{in } \Omega, \\ u_i|_{\partial\Omega} &= 0, \quad \int_{\Omega} u_i u_j = \delta_{ij} \quad \text{for } i, j = 1, \dots, k. \end{aligned}$$

For each $\alpha = 1, \dots, n$ and $i = 1, \dots, k$, following [Payne et al. 1956], consider the functions $\phi_{\alpha i} : \Omega \rightarrow \mathbb{R}$ given by

$$(3-17) \quad \phi_{\alpha i} = x_\alpha u_i - \sum_{j=1}^k r_{\alpha ij} u_j,$$

where

$$(3-18) \quad r_{\alpha ij} = \int_{\Omega} \rho x_{\alpha} u_i u_j.$$

Since $\phi_{\alpha i}|_{\partial\Omega} = 0$ and

$$\int_{\Omega} \rho u_j \phi_{\alpha i} = 0 \quad \text{for } i, j = 1, \dots, k \text{ and } \alpha = 1, \dots, n,$$

it follows from the Rayleigh–Ritz inequality that

$$(3-19) \quad \begin{aligned} \lambda_{k+1} \int_{\Omega} \rho \phi_{\alpha i}^2 &\leq \int_{\Omega} \phi_{\alpha i} \left(- \sum_{\beta, \gamma=1}^n (a_{\beta\gamma} \phi_{\alpha i, \gamma})_{, \beta} + V \phi_{\alpha i} \right) \\ &= \int_{\Omega} \phi_{\alpha i} \left(- \sum_{\beta, \gamma=1}^n (a_{\beta\gamma} (x_{\alpha} u_i)_{, \gamma})_{, \beta} + V x_{\alpha} u_i - \sum_{j=1}^k r_{\alpha ij} \lambda_j \rho u_j \right) \\ &= \int_{\Omega} \phi_{\alpha i} \left(- \sum_{\beta, \gamma=1}^n (a_{\beta\gamma} (x_{\alpha} u_i)_{, \gamma})_{, \beta} + V x_{\alpha} u_i \right) \\ &= \int_{\Omega} \phi_{\alpha i} \left(\lambda_i \rho x_{\alpha} u_i - \sum_{\beta=1}^n ((a_{\alpha\beta} u_i)_{, \beta} + a_{\alpha\beta} u_{i, \beta}) \right) \\ &= \lambda_i \int_{\Omega} \rho \phi_{\alpha i}^2 - \int_{\Omega} \phi_{\alpha i} \left(\sum_{\beta=1}^n ((a_{\alpha\beta} u_i)_{, \beta} + a_{\alpha\beta} u_{i, \beta}) \right) \\ &= \lambda_i \int_{\Omega} \rho \phi_{\alpha i}^2 - \int_{\Omega} x_{\alpha} u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta} u_i)_{, \beta} + a_{\alpha\beta} u_{i, \beta}) \right) + \sum_{j=1}^k r_{\alpha ij} s_{\alpha ij}, \end{aligned}$$

where

$$s_{\alpha ij} = \int_{\Omega} \left(\sum_{\beta=1}^n ((a_{\alpha\beta} u_i)_{, \beta} + a_{\alpha\beta} u_{i, \beta}) \right) u_j.$$

Multiplying the equation

$$(3-20) \quad - \sum_{\beta, \gamma=1}^n (a_{\beta\gamma} u_{j, \beta})_{, \gamma} + V u_j = \lambda_j \rho u_j$$

by $x_{\alpha} u_i$, we have

$$(3-21) \quad - \sum_{\beta, \gamma=1}^n (a_{\beta\gamma} u_{j, \beta})_{, \gamma} x_{\alpha} u_i + V x_{\alpha} u_i u_j = \lambda_j \rho x_{\alpha} u_i u_j.$$

Interchanging the roles of i and j , we get

$$(3-22) \quad - \sum_{\beta\gamma=1}^n (a_{\beta\gamma}u_{i,\beta})_{,\gamma}x_{\alpha}u_j + Vx_{\alpha}u_iu_j = \lambda_i\rho x_{\alpha}u_iu_j.$$

Subtracting (3-21) from (3-22) and integrating the resulted equation on Ω , we get by using the divergence theorem that

$$(3-23) \quad \begin{aligned} (\lambda_i - \lambda_j)r_{\alpha ij} &= \sum_{\beta,\gamma=1}^n \int_{\Omega} ((a_{\beta\gamma}u_{j,\beta})_{,\gamma}x_{\alpha}u_i - (a_{\beta\gamma}u_{i,\beta})_{,\gamma}x_{\alpha}u_j) \\ &= \sum_{\beta,\gamma=1}^n \int_{\Omega} (-a_{\beta\gamma}u_{j,\beta}(x_{\alpha}u_i)_{,\gamma} + a_{\beta\gamma}u_{i,\beta}(x_{\alpha}u_j)_{,\gamma}) \\ &= \sum_{\beta=1}^n \int_{\Omega} (-a_{\alpha\beta}u_{j,\beta}u_i + a_{\alpha\beta}u_{i,\beta}u_j) \\ &= \sum_{\beta=1}^n \int_{\Omega} ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta})u_j = s_{\alpha ij}, \end{aligned}$$

which, combined with (3-19), gives

$$(3-24) \quad \begin{aligned} (\lambda_{k+1} - \lambda_i) \int_{\Omega} \rho \phi_{\alpha i}^2 \\ \leq - \int_{\Omega} \phi_{\alpha i} \left(\sum_{\beta=1}^n ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta}) \right) \\ = - \int_{\Omega} x_{\alpha}u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta}) \right) + \sum_{j=1}^k (\lambda_i - \lambda_j)r_{\alpha ij}^2. \end{aligned}$$

Set

$$t_{\alpha ij} = \int_{\Omega} u_j u_{i,\alpha};$$

then $t_{\alpha ij} + t_{\alpha ji} = 0$ and

$$(3-25) \quad \int_{\Omega} (-2)\phi_{\alpha i}u_{i,\alpha} = -2 \int_{\Omega} x_{\alpha}u_iu_{i,\alpha} + 2 \sum_{j=1}^k r_{\alpha ij}t_{\alpha ij} = \|u_i\|^2 + 2 \sum_{j=1}^k r_{\alpha ij}t_{\alpha ij}.$$

Multiplying (3-25) by $(\lambda_{k+1} - \lambda_i)^2$ and using the Schwarz inequality and (3-24), we get

$$(\lambda_{k+1} - \lambda_i)^2 \left(\|u_i\|^2 + 2 \sum_{j=1}^k r_{\alpha ij}t_{\alpha ij} \right)$$

$$\begin{aligned}
 &= (\lambda_{k+1} - \lambda_i)^2 \int_{\Omega} (-2)\sqrt{\rho}\phi_{\alpha i} \left(\frac{1}{\sqrt{\rho}}u_{i,\alpha} - \sum_{j=1}^k t_{\alpha ij}\sqrt{\rho}u_j \right) \\
 &\leq \delta(\lambda_{k+1} - \lambda_i)^3 \|\sqrt{\rho}\phi_{\alpha i}\|^2 + \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \int_{\Omega} \left| \frac{1}{\sqrt{\rho}}u_{i,\alpha} - \sum_{j=1}^k t_{\alpha ij}\sqrt{\rho}u_j \right|^2 \\
 &= \delta(\lambda_{k+1} - \lambda_i)^3 \|\sqrt{\rho}\phi_{\alpha i}\|^2 + \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \left(\left\| \frac{1}{\sqrt{\rho}}u_{i,\alpha} \right\|^2 - \sum_{j=1}^k t_{\alpha ij}^2 \right) \\
 &\leq \delta(\lambda_{k+1} - \lambda_i)^2 \left(- \int_{\Omega} x_{\alpha}u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta}) \right) + \sum_{j=1}^k (\lambda_i - \lambda_j)r_{\alpha ij}^2 \right) \\
 &\qquad\qquad\qquad + \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \left(\left\| \frac{1}{\sqrt{\rho}}u_{i,\alpha} \right\|^2 - \sum_{j=1}^k t_{\alpha ij}^2 \right),
 \end{aligned}$$

where δ is any positive constant. Summing over i and noticing that $r_{\alpha ij} = r_{\alpha ji}$ and $t_{\alpha ij} = -t_{\alpha ji}$, we infer that

$$\begin{aligned}
 &\sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \|u_i\|^2 - 2 \sum_{i,j=1}^k (\lambda_{k+1} - \lambda_i)(\lambda_i - \lambda_j)r_{\alpha ij}t_{\alpha ij} \\
 &\leq \delta \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \left(- \int_{\Omega} x_{\alpha}u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta}) \right) \right) \\
 &\quad + \sum_{i=1}^k \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \left\| \frac{1}{\sqrt{\rho}}u_{i,\alpha} \right\|^2 - \sum_{i,j=1}^k (\lambda_{k+1} - \lambda_i)\delta(\lambda_i - \lambda_j)^2 r_{\alpha ij}^2 - \sum_{i,j=1}^k \frac{(\lambda_{k+1} - \lambda_i)}{\delta} t_{\alpha ij}^2.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \|u_i\|^2 &\leq \delta \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \left(- \int_{\Omega} x_{\alpha}u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta}) \right) \right) \\
 &\quad + \sum_{i=1}^k \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \left\| \frac{1}{\sqrt{\rho}}u_{i,\alpha} \right\|^2.
 \end{aligned}$$

Summing over α , we infer

$$\begin{aligned}
 (3-26) \quad n \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \|u_i\|^2 \\
 \leq \delta \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \left(- \sum_{\alpha=1}^n \int_{\Omega} x_{\alpha}u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta}u_i)_{,\beta} + a_{\alpha\beta}u_{i,\beta}) \right) \right) \\
 \quad + \sum_{i=1}^k \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \left\| \frac{1}{\sqrt{\rho}}|\nabla u_i| \right\|^2.
 \end{aligned}$$

Since $\int_{\Omega} \rho u_i^2 = 1$ and $\rho_1 \leq \rho(x) \leq \rho_2$ for $x \in \Omega$, we have

$$(3-27) \quad \frac{1}{\rho_2} \leq \|u_i\|^2 \leq \frac{1}{\rho_1}.$$

One gets from the divergence theorem that

$$(3-28) \quad \begin{aligned} & - \sum_{\alpha=1}^n \int_{\Omega} x_{\alpha} u_i \left(\sum_{\beta=1}^n ((a_{\alpha\beta} u_i)_{,\beta} + a_{\alpha\beta} u_{i,\beta}) \right) \\ & = \int_{\Omega} \left(\sum_{\alpha,\beta=1}^n (a_{\alpha\beta} u_i (x_{\alpha} u_i)_{,\beta} - a_{\alpha\beta} u_{i,\beta} x_{\alpha} u_i) \right) = \int_{\Omega} \left(\sum_{\alpha=1}^n a_{\alpha\alpha} \right) u_i^2 \\ & \leq n \xi_2 \int_{\Omega} u_i^2 \leq \frac{n \xi_2}{\rho_1}. \end{aligned}$$

Multiplying the equation $-\sum_{\alpha,\beta=1}^n (a_{\alpha\beta} u_i)_{,\alpha} + V(x) u_i = \lambda_i \rho u_i$ by u_i and integrating over Ω , we get

$$(3-29) \quad \lambda_i = \int_{\Omega} \left(\sum_{\alpha,\beta=1}^n a_{\alpha\beta}(x) u_{i,\alpha} u_{i,\beta} + V(x) u_i^2 \right) \geq \int_{\Omega} \xi_1 |\nabla u_i|^2 + \frac{V_0}{\rho_2},$$

which gives

$$(3-30) \quad \left\| \frac{1}{\sqrt{\rho}} |\nabla u_i| \right\|^2 \leq \frac{1}{\rho_1} \int_{\Omega} |\nabla u_i|^2 \leq \frac{1}{\rho_1 \xi_1} \left(\lambda_i - \frac{V_0}{\rho_2} \right).$$

Substituting (3-27), (3-28) and (3-30) into (3-26), we infer

$$\frac{n}{\rho_2} \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \leq \delta \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \cdot \frac{n \xi_2}{\rho_1} + \sum_{i=1}^k \frac{(\lambda_{k+1} - \lambda_i)}{\delta} \cdot \frac{1}{\rho_1 \xi_1} \left(\lambda_i - \frac{V_0}{\rho_2} \right).$$

Taking

$$(3-31) \quad \delta = \frac{\left(\sum_{i=1}^k (\lambda_{k+1} - \lambda_i) \left(\lambda_i - \frac{V_0}{\rho_2} \right) \right)^{1/2}}{\left(\sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 n \xi_1 \xi_2 \right)^{1/2}},$$

we get (1-15). This completes the proof of Theorem 1.4. □

References

[Ashbaugh 1999] M. S. Ashbaugh, "Isoperimetric and universal inequalities for eigenvalues", pp. 95–139 in *Spectral theory and geometry* (Edinburgh, 1998), edited by B. Davies and Y. Safarov, London Math. Soc. Lecture Note Ser. **273**, Cambridge Univ. Press, Cambridge, 1999. MR 2001a:35131 Zbl 0937.35114

- [Ashbaugh 2002] M. S. Ashbaugh, “The universal eigenvalue bounds of Payne–Pólya–Weinberger, Hile–Protter, and H. C. Yang”, pp. 3–30 in *Spectral and inverse spectral theory* (Goa, 2000), edited by P. D. Hislop and M. Krishna, Proc. Indian Acad. Sci. Math. Sci. **112**, 2002. MR 2004c:35302 Zbl 1199.35261
- [Ashbaugh and Benguria 1993a] M. S. Ashbaugh and R. D. Benguria, “More bounds on eigenvalue ratios for Dirichlet Laplacians in n dimensions”, *SIAM J. Math. Anal.* **24**:6 (1993), 1622–1651. MR 94i:35139 Zbl 0809.35067
- [Ashbaugh and Benguria 1993b] M. S. Ashbaugh and R. D. Benguria, “Universal bounds for the low eigenvalues of Neumann Laplacians in n dimensions”, *SIAM J. Math. Anal.* **24**:3 (1993), 557–570. MR 94b:35191 Zbl 0796.35122
- [Ashbaugh and Hermi 2004] M. S. Ashbaugh and L. Hermi, “A unified approach to universal inequalities for eigenvalues of elliptic operators”, *Pacific J. Math.* **217**:2 (2004), 201–219. MR 2005k:35305 Zbl 1078.35080
- [Berezin 1972] F. A. Berezin, “Covariant and contravariant symbols of operators”, *Izv. Akad. Nauk SSSR Ser. Mat.* **36** (1972), 1134–1167. In Russian. MR 50 #2996 Zbl 0259.47004
- [Chavel 1984] I. Chavel, *Eigenvalues in Riemannian geometry*, Pure and Applied Math. **115**, Academic Press, Orlando, FL, 1984. MR 86g:58140 Zbl 0551.53001
- [Cheng and Yang 2005] Q.-M. Cheng and H. Yang, “Estimates on eigenvalues of Laplacian”, *Math. Ann.* **331**:2 (2005), 445–460. MR 2005i:58038 Zbl 1122.35086
- [Cheng and Yang 2006a] Q.-M. Cheng and H. Yang, “Inequalities for eigenvalues of a clamped plate problem”, *Trans. Amer. Math. Soc.* **358**:6 (2006), 2625–2635. MR 2006m:35263 Zbl 1096.35095
- [Cheng and Yang 2006b] Q.-M. Cheng and H. Yang, “Inequalities for eigenvalues of Laplacian on domains and compact complex hypersurfaces in complex projective spaces”, *J. Math. Soc. Japan* **58**:2 (2006), 545–561. MR 2007k:58051 Zbl 1127.35026
- [Cheng and Yang 2006c] Q.-M. Cheng and H. Yang, “Universal bounds for eigenvalues of a buckling problem”, *Comm. Math. Phys.* **262**:3 (2006), 663–675. MR 2007f:35056 Zbl 1170.35379
- [Cheng and Yang 2007] Q.-M. Cheng and H. Yang, “Bounds on eigenvalues of Dirichlet Laplacian”, *Math. Ann.* **337**:1 (2007), 159–175. MR 2007k:35064 Zbl 1110.35052
- [El Soufi et al. 2007] A. El Soufi, E. M. Harrell, and S. Ilias, “Universal inequalities for the eigenvalues of Laplace and Schrödinger operators on submanifolds”, preprint, 2007. to appear in *Trans. Amer. Math. Soc.* arXiv 0706.0910
- [Harrell 1993] E. M. Harrell, II, “Some geometric bounds on eigenvalue gaps”, *Comm. Partial Differential Equations* **18**:1-2 (1993), 179–198. MR 94c:35135 Zbl 0810.35067
- [Harrell and Michel 1994] E. M. Harrell, II and P. L. Michel, “Commutator bounds for eigenvalues, with applications to spectral geometry”, *Comm. Partial Differential Equations* **19**:11-12 (1994), 2037–2055. MR 95i:58182 Zbl 0815.35078
- [Harrell and Stubbe 1997] E. M. Harrell, II and J. Stubbe, “On trace identities and universal eigenvalue estimates for some partial differential operators”, *Trans. Amer. Math. Soc.* **349**:5 (1997), 1797–1809. MR 97i:35129 Zbl 0887.35111
- [Harrell and Yıldırım Yolcu 2009] E. M. Harrell, II and S. Yıldırım Yolcu, “Eigenvalue inequalities for Klein–Gordon operators”, *J. Funct. Anal.* **256**:12 (2009), 3977–3995. MR 2010e:35198 Zbl 05572175
- [Hile and Protter 1980] G. N. Hile and M. H. Protter, “Inequalities for eigenvalues of the Laplacian”, *Indiana Univ. Math. J.* **29**:4 (1980), 523–538. MR 82c:35052 Zbl 0454.35064
- [Hook 1990] S. M. Hook, “Domain-independent upper bounds for eigenvalues of elliptic operators”, *Trans. Amer. Math. Soc.* **318**:2 (1990), 615–642. MR 90h:35075 Zbl 0727.35096

- [Kröger 1992] P. Kröger, “Upper bounds for the Neumann eigenvalues on a bounded domain in Euclidean space”, *J. Funct. Anal.* **106**:2 (1992), 353–357. MR 93d:47091 Zbl 0777.35044
- [Kröger 1994] P. Kröger, “Estimates for sums of eigenvalues of the Laplacian”, *J. Funct. Anal.* **126**:1 (1994), 217–227. MR 95j:58173 Zbl 0817.35066
- [Laptev 1997] A. Laptev, “Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces”, *J. Funct. Anal.* **151**:2 (1997), 531–545. MR 99a:35027 Zbl 0892.35115
- [Laptev and Weidl 2000] A. Laptev and T. Weidl, “Recent results on Lieb–Thirring inequalities”, pp. XX–1–14 in *Journées “Équations aux Dérivées Partielles”* (La Chapelle sur Erdre, 2000), Univ. Nantes, Nantes, 2000. MR 2001j:81064 Zbl 1135.81337
- [Levitin and Parnowski 2002] M. Levitin and L. Parnowski, “Commutators, spectral trace identities, and universal estimates for eigenvalues”, *J. Funct. Anal.* **192**:2 (2002), 425–445. MR 2003g:47040 Zbl 1058.47022
- [Li and Yau 1983] P. Li and S. T. Yau, “On the Schrödinger equation and the eigenvalue problem”, *Comm. Math. Phys.* **88**:3 (1983), 309–318. MR 84k:58225 Zbl 0554.35029
- [Melas 2003] A. D. Melas, “A lower bound for sums of eigenvalues of the Laplacian”, *Proc. Amer. Math. Soc.* **131**:2 (2003), 631–636. MR 2003i:35218 Zbl 1015.58011
- [Payne et al. 1956] L. E. Payne, G. Pólya, and H. F. Weinberger, “On the ratio of consecutive eigenvalues”, *J. Math. and Phys.* **35** (1956), 289–298. MR 18,905c Zbl 0073.08203
- [Pólya 1961] G. Pólya, “On the eigenvalues of vibrating membranes”, *Proc. London Math. Soc.* (3) **11** (1961), 419–433. MR 23 #B2256 Zbl 0107.41805
- [Sun et al. 2008] H. Sun, Q.-M. Cheng, and H. Yang, “Lower order eigenvalues of Dirichlet Laplacian”, *Manuscripta Math.* **125**:2 (2008), 139–156. MR 2009i:58042 Zbl 1137.35050
- [Wang and Xia 2007a] Q. Wang and C. Xia, “Universal bounds for eigenvalues of the biharmonic operator on Riemannian manifolds”, *J. Funct. Anal.* **245**:1 (2007), 334–352. MR 2008e:58033 Zbl 1113.58013
- [Wang and Xia 2007b] Q. Wang and C. Xia, “Universal inequalities for eigenvalues of the buckling problem on spherical domains”, *Comm. Math. Phys.* **270**:3 (2007), 759–775. MR 2007m:35180 Zbl 1112.74017
- [Wang and Xia 2008] Q. Wang and C. Xia, “Universal bounds for eigenvalues of Schrödinger operator on Riemannian manifolds”, *Ann. Acad. Sci. Fenn. Math.* **33** (2008), 319–336. MR 2009c:35333 Zbl 1171.35091
- [Wang and Xia 2010a] Q. Wang and C. Xia, “Inequalities for eigenvalues of the biharmonic operator with weight on Riemannian manifolds”, *J. Math. Soc. Japan* **62**:2 (2010), 597–622. MR 2662854 Zbl 1200.53042
- [Wang and Xia 2010b] Q. Wang and C. Xia, “Isoperimetric bounds for the first eigenvalue of the Laplacian”, *Z. Angew. Math. Phys.* **61**:1 (2010), 171–175. MR 2011e:35246 Zbl 1192.35125
- [Wang and Xia 2010c] Q. Wang and C. Xia, “Universal bounds for eigenvalues of the biharmonic operator”, *J. Math. Anal. Appl.* **364**:1 (2010), 1–17. MR 2010k:35334 Zbl 1189.35208
- [Wang and Xia 2011] Q. Wang and C. Xia, “Inequalities for eigenvalues of a clamped plate problem”, *Calc. Var. Partial Differential Equations* **40** (2011), 273–289. MR 2745203 Zbl 1205.35175
- [Weyl 1912] H. Weyl, “Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung)”, *Math. Ann.* **71**:4 (1912), 441–479. MR 1511670
- [Yang 1991] H. C. Yang, “An estimate of the difference between consecutive eigenvalues”, preprint IC/91/60, ICTP, Trieste, 1991.

Received April 22, 2009. Revised October 20, 2009.

QIAOLING WANG
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BRASÍLIA
ASA NORTE
70910-900 BRASÍLIA
BRAZIL
wang@impa.br

CHANGYU XIA
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BRASÍLIA
ASA NORTE
70910-900 BRASÍLIA
BRAZIL
xia@pq.cnpq.br

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic.

Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 251 No. 1 May 2011

An analogue of the Cartan decomposition for p -adic symmetric spaces of split p -adic reductive groups	1
PATRICK DELORME and VINCENT SÉCHERRE	
Unital quadratic quasi-Jordan algebras	23
RAÚL FELIPE	
The Dirichlet problem for constant mean curvature graphs in $\mathbb{H} \times \mathbb{R}$ over unbounded domains	37
ABIGAIL FOLHA and SOFIA MELO	
Osgood–Hartogs-type properties of power series and smooth functions	67
BUMA L. FRIDMAN and DAOWEI MA	
Twisted Cappell–Miller holomorphic and analytic torsions	81
RUNG-TZUNG HUANG	
Generalizations of Agol’s inequality and nonexistence of tight laminations	109
THILO KUESSNER	
Chern numbers and the indices of some elliptic differential operators	173
PING LI	
Blocks of the category of cuspidal \mathfrak{sp}_{2n} -modules	183
VOLODYMYR MAZORCHUK and CATHARINA STROPPEL	
A constant mean curvature annulus tangent to two identical spheres is Delauney	197
SUNG-HO PARK	
A note on the topology of the complements of fiber-type line arrangements in $\mathbb{C}\mathbb{P}^2$	207
SHENG-LI TAN, STEPHEN S.-T. YAU and FEI YE	
Inequalities for the Navier and Dirichlet eigenvalues of elliptic operators	219
QIAOLING WANG and CHANGYU XIA	
A Beurling–Hörmander theorem associated with the Riemann–Liouville operator	239
XUECHENG WANG	



0030-8730(201105)251:1;1-E