

*Pacific
Journal of
Mathematics*

**REGULARITY OF WEAKLY HARMONIC MAPS FROM A
FINSLER SURFACE INTO AN n -SPHERE**

XIAOHUAN MO AND LIANG ZHAO

Volume 253 No. 1

September 2011

REGULARITY OF WEAKLY HARMONIC MAPS FROM A FINSLER SURFACE INTO AN n -SPHERE

XIAOHUAN MO AND LIANG ZHAO

We study harmonic maps on Finsler surfaces. Using Berwald frames on Finsler surfaces, we prove conformal invariance for the energy of Finsler harmonic maps. As an application, we show that weakly harmonic maps from a Finsler surface to a sphere \mathbb{S}^n are in fact smooth by establishing a new Jacobi structure, generalizing the regularity result previously known for the case of a harmonic map from a Riemannian surface.

1. Introduction

Harmonic maps between Riemannian manifolds are very important both in classical and modern differential geometry. They are defined as the critical points of the energy functionals. One of the fundamental problems in the theory of harmonic maps is to study regularity [Evans 1991; Hélein 1990; 1991b; 1991a]. Harmonic maps on Riemann surfaces are the natural extension of minimal surfaces in Riemannian manifolds. More importantly, the energy of a harmonic map on a two-dimensional domain is a conformal invariant, i.e. the set of harmonic maps from a Riemannian surface M depends only on the conformal structure on M .

Finsler manifolds are just Riemannian manifolds with metrics without the quadratic restriction [Chern 1996]. Many Finslerian geometers have made effort in the investigation of the geometric and analytic properties (such as, existence, stability, etc) of harmonic maps on Finsler manifolds [He and Shen 2005; Mo 2001; Mo and Yang 2005; Shen and Wei 2008; Shen and Zhang 2004].

In this paper, we are going to study the theory of harmonic maps on Finsler surfaces. Let (M, F) be a Finsler manifold and (N, h) a Riemannian manifold. We will say that $\phi : (M, F) \rightarrow (N, h)$ is *Finsler harmonic* if ϕ is a critical point of the energy functional with respect to any compactly supported variation of ϕ [Mo 2001]. First, we obtain conformal invariance for Finsler harmonic maps from surfaces by using the Berwald frames on surfaces (see Theorem 3.1).

This work is supported by the National Natural Science Foundation of China 11071005, 11001268 and PCSIRT.

MSC2000: primary 53C60; secondary 54E40.

Keywords: Finsler surface, conformal invariance, weakly harmonic map, regularity.

As an application, we obtain a regularity result of a weakly Finsler harmonic map on a Finsler surface M , generalizing a theorem of Hélein for the case of M being a Riemannian surface [Hélein 1990]. More precisely:

Theorem 1.1. *For a Finsler surface (M, F) with $\partial M = \emptyset$, if $\phi \in W^{1,2}(M, \mathbb{S}^n)$ be a weakly Finsler harmonic map, then $\phi \in C^\infty(M, \mathbb{S}^n)$.*

Recall that $W^{1,2}(M, \mathbb{R}^l)$ is the usual Sobolev space of L^2 functions on M to \mathbb{R}^l having first derivatives in the sense of distributions in L^2 . A map $\phi \in W^{1,2}(M, N)$ is *weakly Finsler harmonic map* if it is a weak solution to the Euler–Lagrange equation (4-8), where the Sobolev space $W^{1,2}(M, N)$ of maps from M to $N \subset \mathbb{R}^l$ is defined by

$$W^{1,2}(M, N) = \{\varphi \in W^{1,2}(M, \mathbb{R}^l) : \varphi(x) \in N \text{ a.e. } x \in M\}$$

(see Section 4).

It is well-known that the Jacobi (determinant) structure plays an important role in studying the regularity of harmonic maps from Riemannian surfaces to Riemannian spheres [Lin and Wang 2008]. But, in general, we cannot get the Jacobi structure of Finsler harmonic maps directly. So we have to construct a new Jacobi structure by using the existence and regularity theory of elliptic equations (see Lemma 4.1). After obtaining this interesting structure, we can prove Theorem 1.1 successfully.

2. Preliminaries

Finsler harmonic maps. Let $\phi : (M, F) \rightarrow (N, h)$ be a map from an m -dimensional Finsler manifold (M, F) to an n -dimensional Riemannian manifold (N, h) . In natural coordinates, the energy density of ϕ is given by

$$(2-1) \quad e(\phi) = \frac{1}{2} g^{ij} \frac{\partial \phi^\alpha}{\partial x^i} \frac{\partial \phi^\beta}{\partial x^j} h_{\alpha\beta}$$

where $\phi = (\phi^1, \dots, \phi^n)$, $g_{ij} = (\frac{1}{2} F^2)_{y^i y^j}$ and $(g^{ij}) = (g_{ij})^{-1}$. We introduce a dual adapted orthonormal frame e_i on the Riemannian vector bundle $(\pi^* TM, g)$ and coframe ω_i with $\omega_m := (\partial F / \partial y^i) dx^i$ where $\pi : SM \rightarrow M$ is the canonical projection map, SM the projective sphere bundle of M and g is the fundamental tensor of F . Put $\omega_i = v_{ij} dx^j$. Then $\det(v_{ij}) = \sqrt{\det(g_{kl})}$ [Mo 2001, p. 1333] and $g = \sum_{i=1}^m \omega_i^2$.

Taking the exterior derivative of ω_m yields the Chern connection on $\pi^* TM$ described by an $m \times m$ matrix of 1-forms (ω_{ij}) on SM . The coframe ω_i and connection forms $\omega_{m\alpha}$ determine the volume form with respect to the Riemannian metric $G := \sum_{i=1}^m \omega_i^2 + \sum_{\lambda=1}^{m-1} \omega_{m\lambda}^2$ on SM as follows:

$$\Pi := \omega_1 \wedge \dots \wedge \omega_m \wedge \omega_{m1} \wedge \dots \wedge \omega_{m,m-1}.$$

A direct calculation yields

$$(2-2) \quad \begin{aligned} \Pi &= v_{1i_1} dx^{i_1} \wedge \cdots \wedge v_{m i_m} dx^{i_m} \wedge \omega_{m1} \wedge \cdots \wedge \omega_{m, m-1} \\ &= \det(v_{ij}) dx^1 \wedge \cdots \wedge dx^m \wedge \chi = \sqrt{\det(g_{kl})} dx \wedge \chi, \end{aligned}$$

where $dx = dx^1 \wedge \cdots \wedge dx^m$ and

$$(2-3) \quad \chi \equiv \omega_{m1} \wedge \cdots \wedge \omega_{m, m-1} \pmod{dx^j}.$$

If Ω is a compact domain in M , define the energy of $\phi : (\Omega, F) \rightarrow (N, h)$ by

$$(2-4) \quad E(\phi, \Omega) = \frac{1}{c} \int_{S\Omega} e(\phi) \Pi,$$

where c is the volume of the standard $(m-1)$ -dimensional sphere and $S\Omega$ is the projective sphere bundle of Ω . If M is compact, we write $E(\phi) = E(\phi, M)$.

Plugging (2-1) into (2-4) and using (2-2) we have

$$(2-5) \quad E(\phi) = \frac{1}{2c} \int_{SM} g^{ij} \frac{\partial \phi^\alpha}{\partial x^i} \frac{\partial \phi^\beta}{\partial x^j} h_{\alpha\beta} \sqrt{\det(g_{kl})} dx \wedge \chi.$$

Conformal transformations of Finsler metrics. Let F be a Finsler metric on a manifold M and g its fundamental tensor. For two arbitrary nonzero vectors v, y in $T_x M$, the angle $\theta(y, v)$ between y and v is defined by [Antonelli et al. 1993; Bácsó and Cheng 2007]

$$\cos \theta(y, v) := \frac{g_{(x, y)}(y, v)}{\sqrt{g_{(x, y)}(y, y)} \sqrt{g_{(x, y)}(v, v)}}.$$

Let F and \bar{F} be two Finsler metrics on a manifold M . If the angle $\theta(y, v)$ with respect to F is equal to the angle $\bar{\theta}(y, v)$ with respect to \bar{F} for any nonzero vectors $v, y \in T_x M$ and any $x \in M$, then F is called *conformal* to \bar{F} and the transformation $F \rightarrow \bar{F}$ of the metric is called a *conformal transformation*.

Now assume that F is conformal to \bar{F} . From [Antonelli et al. 1993], we have

$$(2-6) \quad \bar{F}(x, y) = e^{c(x)} F(x, y)$$

for some scalar function $c(x)$. Furthermore, we have (see [Matsumoto 2003])

$$(2-7) \quad \bar{g}_{ij}(x, y) = e^{2c(x)} g_{ij}(x, y),$$

$$(2-8) \quad \bar{g}^{ij}(x, y) = e^{-2c(x)} g^{ij}(x, y).$$

3. Conformal invariance

In this section we are going to show an important property for Finsler harmonic maps on two-dimensional domains: the conformal invariance required in the proof of the regularity result.

Theorem 3.1. *The energy functional E on the set of Finsler harmonic maps from a surface M to a Riemannian manifold N depends only on the conformal structures on M .*

Proof. This follows from the conformal invariance of the energy functional E in dimension two. We assume that $\varphi : F \rightarrow \bar{F}$ is given by (2-6) and denote the corresponding objects with respect to \bar{F} by adding a bar $\bar{\cdot}$.

By using (2-7), we obtain

$$(3-1) \quad \sqrt{\det(\bar{g}_{kl})} = \sqrt{\det(e^{2c(x)}g_{kl})} = \sqrt{e^{4c(x)}\det(g_{kl})} = e^{2c(x)}\sqrt{\det(g_{kl})}.$$

The first structure equation for (M, F) can be written as

$$(3-2) \quad d\omega_i = \sum \omega_j \wedge \omega_{ji}, \quad \omega_{ij} + \omega_{ji} = -2 \sum A_{ijk}\omega_{2k},$$

where $A_{ijk} = A(e_i, e_j, e_k)$ and A is the Cartan tensor of F . From [Mo 2006, (3.79)], we have

$$(3-3) \quad \omega_{12} \equiv -u_1^i F_{y^i y^j} dy^j \pmod{dx^i}.$$

By using (2-3), (3-2) and (3-3) we obtain

$$(3-4) \quad \chi \equiv \omega_{21} = -\omega_{12} - 2 \sum A_{21i}\omega_{2i} = -\omega_{12} \equiv u_1^i F_{y^i y^j} dy^j,$$

where the equivalences are mod dx^i and the u_j^i are defined by

$$e_j = u_j^i \frac{\partial}{\partial x^i}.$$

It is not difficult to conclude that the section e_1 is in fact globally defined [Mo 2006]. The frame $\{e_1, e_2\}$ is call the *Berwald frame* [Bao and Chern 1996]. Explicitly,

$$e_1 = \frac{F_{y^2}}{\sqrt{\det(g_{ij})}} \frac{\partial}{\partial x^1} + \frac{-F_{y^1}}{\sqrt{\det(g_{ij})}} \frac{\partial}{\partial x^2}.$$

It follows that

$$u_1^1 = \frac{F_{y^2}}{\sqrt{\det(g_{ij})}}, \quad u_1^2 = \frac{-F_{y^1}}{\sqrt{\det(g_{ij})}}.$$

Under the conformal transformation (2-6), from (3-1) we have

$$\begin{aligned} \bar{u}_1^1 &= \frac{\bar{F}_{y^2}}{\sqrt{\det(\bar{g}_{ij})}} = \frac{e^{c(x)}F_{y^2}}{e^{2c(x)}\sqrt{\det(g_{ij})}} = e^{-c(x)}u_1^1, \\ \bar{u}_1^2 &= -\frac{\bar{F}_{y^1}}{\sqrt{\det(\bar{g}_{ij})}} = e^{-c(x)}u_1^2, \\ \bar{F}_{y^i y^j} &= (e^{c(x)}F)_{y^i y^j} = e^{c(x)}F_{y^i y^j}. \end{aligned}$$

Hence

$$\bar{u}_1^i \bar{F}_{y^i y^j} = u_1^i F_{y^i y^j}.$$

From this together with (3-4) we get

$$(3-5) \quad \bar{\chi} \equiv \chi \pmod{dx^i}.$$

Thus, for any $\phi \in W^{1,2}(M, N)$,

$$\begin{aligned} \bar{E}(\phi) &= \frac{1}{2c} \int_{SM} \bar{g}^{ij} \frac{\partial \phi^\alpha}{\partial x^i} \frac{\partial \phi^\beta}{\partial x^j} h_{\alpha\beta} \sqrt{\det(\bar{g}_{kl})} dx \wedge \bar{\chi} \\ &= \frac{1}{2c} \int_{SM} e^{-2c(x)} g^{ij} \frac{\partial \phi^\alpha}{\partial x^i} \frac{\partial \phi^\beta}{\partial x^j} h_{\alpha\beta} e^{2c(x)} \sqrt{\det(g_{kl})} dx \wedge \chi \\ &= \frac{1}{2c} \int_{SM} g^{ij} \frac{\partial \phi^\alpha}{\partial x^i} \frac{\partial \phi^\beta}{\partial x^j} h_{\alpha\beta} \sqrt{\det(g_{kl})} dx \wedge \chi = E(\phi). \end{aligned} \quad \square$$

Let (M, F) be a Finsler surface and set

$$\begin{aligned} \tilde{g}^{ij}(x) &:= \frac{\int_{S_x M} g^{ij}(x, y) \sqrt{\det(g_{kl}(x, y))} \chi}{\int_{S_x M} \sqrt{\det(g_{kl}(x, y))} \chi}, \\ \tilde{g} &:= \tilde{g}_{ij} dx^i \otimes dx^j, \quad \text{where } (\tilde{g}_{ij}) = (\tilde{g}^{ij})^{-1}. \end{aligned}$$

Then (see [Mo and Yang 2005, Lemma 5.2]) \tilde{g} is a Riemannian metric on M , called the (Riemannian) metric *induced* by F .

Proposition 3.2. *Let F and \bar{F} be two Finsler metrics on a surface. If $\bar{F}(x, y) = e^{c(x)} F(x, y)$, then*

$$\tilde{\bar{g}}(x) = e^{2c(x)} \tilde{g},$$

where $\tilde{\bar{g}}$ and \tilde{g} are the induced metrics by \bar{F} and F respectively.

Proof. Using (2-7), (3-1) and (3-5) we get

$$\begin{aligned} \tilde{\bar{g}}^{ij}(x) &= \frac{\int_{S_x M} \bar{g}^{ij}(x, y) \sqrt{\det(\bar{g}_{kl}(x, y))} \bar{\chi}}{\int_{S_x M} \sqrt{\det(\bar{g}_{kl}(x, y))} \bar{\chi}} \\ &= \frac{\int_{S_x M} e^{-2c(x)} g^{ij}(x, y) e^{2c(x)} \sqrt{\det(g_{kl}(x, y))} \chi}{\int_{S_x M} e^{2c(x)} \sqrt{\det(g_{kl}(x, y))} \chi} = e^{-2c(x)} \tilde{g}^{ij}(x). \end{aligned}$$

It follows that

$$\tilde{\bar{g}}(x) := \tilde{\bar{g}}_{ij}(x) dx^i \otimes dx^j = e^{2c(x)} \tilde{g}_{ij}(x) dx^i \otimes dx^j = e^{2c(x)} \tilde{g}(x). \quad \square$$

Note that locally all Riemannian conformal structures are equivalent [Hélein 2002, Theorem 1.1.3].

Corollary 3.3. *Let (M, F) be a Finsler surface. For each point x_0 , there is a local coordinate system (U, x^i) in M , and a conformal transformation $F \rightarrow \bar{F}$, such that the Riemannian metric induced by \bar{F} is a canonical Euclidean metric on U .*

4. Proof of Theorem 1.1

Before giving the proof proper, we state several lemmas. Throughout this section, D will denote a small disc in \mathbb{R}^2 , but its radius may shrink as we go along if necessary.

Lemma 4.1. *Let $D \subset \mathbb{R}^2$ be a disk. If $\phi \in W^{1,2}(D, \mathbb{S}^n)$ is a weakly Finsler harmonic map, then for any $1 \leq \alpha, \beta \leq n$, the field*

$$V^{\alpha\beta} \equiv \phi^\beta \nabla \phi^\alpha - \phi^\alpha \nabla \phi^\beta - \nabla \phi^{\alpha\beta} \in L^2(D, \mathbb{R}^2),$$

is divergence-free, where $\varphi^{\alpha\beta}$ is a solution of

$$(4-1) \quad \Delta \varphi^{\alpha\beta} = \nabla \psi \cdot (\phi^\beta \nabla \phi^\alpha - \phi^\alpha \nabla \phi^\beta)$$

in which

$$\psi(x) = -\log \sqrt{\det(\tilde{g}_{kl})} - \log \int_{S_x M} \sqrt{\det(g_{kl}(x, y))} \chi.$$

Moreover, ϕ satisfies

$$(4-2) \quad -\Delta(\phi - \eta)^\alpha = \sum_{\beta=1}^{n+1} \nabla \phi^\beta \cdot V^{\alpha\beta}$$

where $\eta = (\eta^1, \dots, \eta^{n+1})$ is a solution of

$$(4-3) \quad -\Delta \eta^\alpha = \sum_{\beta=1}^{n+1} \nabla \phi^\beta \cdot \nabla \phi^{\alpha\beta} - \nabla \psi \cdot \nabla \phi^\alpha.$$

Proof. For a round spherical target manifold, the Euler–Lagrange equation of ϕ is given by $\Delta \phi = |\nabla \phi|^2 \phi + \nabla \psi \cdot \nabla \phi$ [Mo and Yang 2005], equivalently,

$$(4-4) \quad \Delta \phi^\alpha = |\nabla \phi|^2 \phi^\alpha + \nabla \psi \cdot \nabla \phi^\alpha$$

for $\alpha = 1, \dots, n+1$. Set $\Omega^{\alpha\beta} \equiv \phi^\beta \nabla \phi^\alpha - \phi^\alpha \nabla \phi^\beta$. Then

$$\begin{aligned} \operatorname{div}(\Omega^{\alpha\beta}) &= \operatorname{div}(\phi^\beta \nabla \phi^\alpha) - \operatorname{div}(\phi^\alpha \nabla \phi^\beta) \\ &= \phi^\beta \Delta \phi^\alpha + \nabla \phi^\beta \cdot \nabla \phi^\alpha - (\phi^\alpha \Delta \phi^\beta + \nabla \phi^\alpha \cdot \nabla \phi^\beta) \\ &= \phi^\beta \Delta \phi^\alpha - \phi^\alpha \Delta \phi^\beta \\ &= \phi^\beta (|\nabla \phi|^2 \phi^\alpha + \nabla \psi \cdot \nabla \phi^\alpha) - \phi^\alpha (|\nabla \phi|^2 \phi^\beta + \nabla \psi \cdot \nabla \phi^\beta) \\ &= \nabla \psi \cdot (\phi^\beta \nabla \phi^\alpha - \phi^\alpha \nabla \phi^\beta) = \nabla \psi \cdot \Omega^{\alpha\beta}. \end{aligned}$$

Together with (4-1) we obtain

$$\begin{aligned} \operatorname{div}(V^{\alpha\beta}) &= \operatorname{div}(\Omega^{\alpha\beta} - \nabla\varphi^{\alpha\beta}) = \operatorname{div}(\Omega^{\alpha\beta}) - \operatorname{div}(\nabla\varphi^{\alpha\beta}) \\ &= \nabla\psi \cdot \Omega^{\alpha\beta} - \Delta\varphi^{\alpha\beta} = \nabla\psi \cdot (\Omega^{\alpha\beta} - \phi^\beta \nabla\phi^\alpha + \phi^\alpha \nabla\phi^\beta) = 0. \end{aligned}$$

This implies that $V^{\alpha\beta}$ is divergence-free. From (4-1) and the regularity theory of elliptic equations, we obtain $\varphi^{\alpha\beta} \in W^{2,2}(D)$. To see (4-2), observe that

$$(4-5) \quad \sum_{\beta=1}^{n+1} \phi^\beta \nabla\phi^\beta = 0,$$

since $|\phi|^2 = 1$. We rewrite (4-4) as

$$(4-6) \quad -\Delta\phi^\alpha = -K - \nabla\psi \cdot \nabla\phi^\alpha,$$

where

$$\begin{aligned} (4-7) \quad K &= |\nabla\phi|^2\phi^\alpha = \left(\sum_{\beta=1}^{n+1} \nabla\phi^\beta \cdot \nabla\phi^\beta \right) \phi^\alpha = \sum_{\beta=1}^{n+1} \nabla\phi^\beta \cdot (\phi^\alpha \nabla\phi^\beta) \\ &= \sum_{\beta=1}^{n+1} \nabla\phi^\beta \cdot (\phi^\alpha \nabla\phi^\beta - \phi^\beta \nabla\phi^\alpha) = - \sum_{\beta=1}^{n+1} \nabla\phi^\beta \cdot (V^{\alpha\beta} + \nabla\varphi^{\alpha\beta}), \end{aligned}$$

where we used (4-5). Plugging the last expression into (4-6) and using (4-3) yields

$$-\Delta\phi^\alpha = \sum_{\beta=1}^{n+1} \nabla\phi^\beta \cdot (V^{\alpha\beta} + \nabla\varphi^{\alpha\beta}) - \nabla\psi \cdot \nabla\phi^\alpha = \sum_{\beta=1}^{n+1} \nabla\phi^\beta \cdot V^{\alpha\beta} - \Delta\eta^\alpha.$$

Thus $\phi - \eta$ satisfies (4-2). □

Remark. From (4-3) and the regularity theory of elliptic equations, we have $\eta^\alpha \in C^0(D)$.

The right-hand side of (4-2) has a special structure: it is the product of a curl-free vector field and a divergence-free vector field. It follows from Lemma 4.3 below that the product belongs to the Hardy space $\mathcal{H}^1(\mathbb{R}^m)$ [Lin and Wang 2008, Definition 3.2.3]. (The next three lemmas can be found in [Lin and Wang 2008].)

Lemma 4.2 [Stein 1993]. *For $f \in \mathcal{H}^1(\mathbb{R}^m)$, let $u \in L^1(\mathbb{R}^m)$ be a solution of*

$$-\Delta u = f \quad \text{in } \mathbb{R}^m,$$

where Δu is the Laplacian of u . Then all second-order derivatives $\nabla^2 u$ of u satisfy $\nabla^2 u \in L^1(\mathbb{R}^m)$ and

$$\|\nabla^2 u\|_{L^1(\mathbb{R}^m)} \leq C \|f\|_{\mathcal{H}^1(\mathbb{R}^m)}.$$

Lemma 4.3 [Coifman et al. 1989]. *For $1 < p < +\infty$ and $q = p/(p-1)$, suppose $h \in W^{1,p}(\mathbb{R}^m)$ and $G \in L^q(\mathbb{R}^m, \mathbb{R}^m)$ is a divergence-free vector field. Then the function $f := \nabla h \cdot G \in \mathcal{H}^1(\mathbb{R}^m)$. Moreover, there exists a positive constant C_m depending only on m such that*

$$\|f\|_{\mathcal{H}^1(\mathbb{R}^m)} \leq C_m \|h\|_{W^{1,p}(\mathbb{R}^m)} \|G\|_{L^q(\mathbb{R}^m)}.$$

Lemma 4.4 [Hélein 2002, Theorem 3.3.4]. *If $f \in W^{1,2}(\mathbb{R}^2)$ has compact support and the weak derivative ∇f of f belongs to the Lorentz space $L^{(2,1)}(\mathbb{R}^2)$, then $f \in C^0(\mathbb{R}^2)$.*

Proof of Theorem 1.1. Since regularity is a local property, it suffices to show that it is valid in the neighborhood of each point in M . But, since every sufficiently small geodesic ball in (M, \tilde{g}) is conformal equivalent to the unit ball D of \mathbb{R}^2 equipped with the canonical metric via a conformal transformation of corresponding Finsler metric (Corollary 3.3), and Finsler harmonic maps are preserved under Finsler conformal transformation (Theorem 3.1), we assume that $M = D$ is a disc in \mathbb{R}^2 .

The Euler–Lagrange equation for Finsler harmonic map ϕ is given by [Mo and Yang 2005, (5.8)]

$$(4-8) \quad \Delta\phi = A(\phi)(\nabla\phi, \nabla\phi) + \nabla\psi \cdot \nabla\phi,$$

where $A(\phi)$ is the second fundamental form of \mathbb{S}^n in \mathbb{R}^{n+1} . Note that \mathbb{S}^n is a totally umbilical hypersurface in \mathbb{R}^{n+1} with unit normal field ϕ . Hence

$$A(\phi)(\nabla\phi, \nabla\phi) = |\nabla\phi|^2\phi.$$

Plugging this into (4-8) yields (4-4).

For $1 \leq \alpha, \beta \leq n+1$, let $V^{\alpha\beta} \in L^2(D, \mathbb{R}^2)$ be divergence-free vector fields given by Lemma 4.2. Let $\hat{\phi} \in W^{1,2}(\mathbb{R}^2, \mathbb{R}^n)$ be an extension of ϕ such that $\|\nabla\hat{\phi}\|_{L^2(\mathbb{R}^2)} \leq C\|\nabla\phi\|_{L^2(D)}$ and $\hat{V}^{\alpha\beta} \in L^2(\mathbb{R}^2, \mathbb{R}^2)$ be an extension of $V^{\alpha\beta}$ such that

$$(4-9) \quad \operatorname{div}(\hat{V}^{\alpha\beta}) = 0, \quad \|\hat{V}^{\alpha\beta}\|_{L^2(\mathbb{R}^2)} \leq C\|V^{\alpha\beta}\|_{L^2(D)}.$$

The existence of such $\hat{V}^{\alpha\beta}$ can be obtained as follows.

We write $V^{\alpha\beta}$ as

$$(4-10) \quad V^{\alpha\beta} = (V_1^{\alpha\beta}, V_2^{\alpha\beta}).$$

Then

$$0 = \operatorname{div}(V^{\alpha\beta}) = \frac{\partial V_1^{\alpha\beta}}{\partial x^1} + \frac{\partial V_2^{\alpha\beta}}{\partial x^2}.$$

It follows that

$$\frac{\partial V_1^{\alpha\beta}}{\partial x^1} = -\frac{\partial V_2^{\alpha\beta}}{\partial x^2};$$

therefore, there exists $\rho \in W^{1,2}(D)$ such that

$$\frac{\partial \rho}{\partial x^1} = -V_2^{\alpha\beta}, \quad \frac{\partial \rho}{\partial x^2} = V_1^{\alpha\beta}.$$

Substituting these into (4-10) yields

$$V^{\alpha\beta} = \left(\frac{\partial \rho}{\partial x^2}, -\frac{\partial \rho}{\partial x^1} \right)$$

in D . Let $\hat{\rho} \in W^{1,2}(\mathbb{R}^2)$ be an extension of ρ such that

$$\|\nabla \hat{\rho}\|_{L^2(\mathbb{R}^2)} \leq C \|\nabla \rho\|_{L^2(D)}.$$

Then

$$\hat{V}^{\alpha\beta} := \left(\frac{\partial \hat{\rho}}{\partial x^2}, -\frac{\partial \hat{\rho}}{\partial x^1} \right)$$

satisfies (4-9).

For $1 \leq \alpha \leq n+1$, let $\omega^\alpha \in L^1(\mathbb{R}^2)$ be a solution of

$$-\Delta \omega^\alpha = \sum_{\beta=1}^{n+1} \nabla \hat{\phi}^\beta \cdot \hat{V}^{\alpha\beta}$$

in \mathbb{R}^2 . By Lemmas 4.2 and 4.3, we know that $\nabla^2 \omega^\alpha \in L^1(\mathbb{R}^2)$ and

$$\begin{aligned} \|\nabla^2 \omega^\alpha\|_{L^1(\mathbb{R}^2)} &\leq C \sum_{\beta=1}^{n+1} \|\nabla \hat{\phi}^\beta \cdot \hat{V}^{\alpha\beta}\|_{\mathcal{H}^1(\mathbb{R}^2)} \\ &\leq C \sum_{\beta=1}^{n+1} \|\nabla \hat{\phi}^\beta\|_{L^2(\mathbb{R}^2)} \|\hat{V}^{\alpha\beta}\|_{L^2(\mathbb{R}^2)} \\ &\leq C \sum_{\beta=1}^{n+1} \|\nabla \phi^\beta\|_{L^2(D)} \|V^{\alpha\beta}\|_{L^2(D)} \leq C \|\nabla \phi\|_{L^2(D)}^2. \end{aligned}$$

It follows that $\nabla \omega^\alpha \in W^{1,1}(\mathbb{R}^2)$. Note that for $m \geq 2$, $W^{1,1}(\mathbb{R}^m)$ is continuously embedded in $L^{(\frac{m}{m-1}, 1)}(\mathbb{R}^m)$ [Hélein 2002]. This implies that $\nabla \omega^\alpha \in L^{(2,1)}(\mathbb{R}^2)$. Hence by Lemma 4.4 we have $\omega^\alpha \in C_{loc}^0(\mathbb{R}^2)$.

Let $v^\alpha = \phi^\alpha - \eta^\alpha - \omega^\alpha : D \rightarrow \mathbb{R}$. By using (4-2) we have

$$\begin{aligned} -\Delta v^\alpha &= -\Delta(\phi^\alpha - \eta^\alpha - \omega^\alpha) = -\Delta(\phi^\alpha - \eta^\alpha) + \Delta \omega^\alpha \\ &= \sum_{\beta=1}^{n+1} \nabla \phi^\beta \cdot V^{\alpha\beta} - \sum_{\beta=1}^{n+1} \nabla \hat{\phi}^\beta \cdot \hat{V}^{\alpha\beta} \\ &= \sum_{\beta=1}^{n+1} \nabla \phi^\beta \cdot V^{\alpha\beta} - \sum_{\beta=1}^{n+1} \nabla \phi^\beta \cdot V^{\alpha\beta} = 0 \quad \text{in } D, \end{aligned}$$

so $v^\alpha \in C^0(D)$. This implies that $\phi \in C^0(D, \mathbb{S}^n)$ and $\phi \in C^\infty(D, \mathbb{S}^n)$ by higher-order regularity theory for harmonic maps [Jost 2008, Section 8.6]. \square

References

- [Antonelli et al. 1993] P. L. Antonelli, R. S. Ingarden, and M. Matsumoto, *The theory of sprays and Finsler spaces with applications in physics and biology*, Fundamental Theories of Physics **58**, Kluwer, Dordrecht, 1993. MR 95e:53094 Zbl 0821.53001
- [Bácsó and Cheng 2007] S. Bácsó and X. Cheng, “Finsler conformal transformations and the curvature invariances”, *Publ. Math. Debrecen* **70**:1-2 (2007), 221–231. MR 2008a:53013 Zbl 1174.53013
- [Bao and Chern 1996] D. Bao and S. S. Chern, “A note on the Gauss-Bonnet theorem for Finsler spaces”, *Ann. of Math. (2)* **143**:2 (1996), 233–252. MR 97d:53071 Zbl 0849.53046
- [Chern 1996] S.-S. Chern, “Finsler geometry is just Riemannian geometry without the quadratic restriction”, *Notices Amer. Math. Soc.* **43**:9 (1996), 959–963. MR 97e:53129 Zbl 1044.53512
- [Coifman et al. 1989] R. R. Coifman, P.-L. Lions, Y. Meyer, and S. Semmes, “Compacité par compensation et espaces de Hardy”, *C. R. Acad. Sci. Paris Sér. I Math.* **309**:18 (1989), 945–949. MR 91h:46058
- [Evans 1991] L. C. Evans, “Partial regularity for stationary harmonic maps into spheres”, *Arch. Rational Mech. Anal.* **116**:2 (1991), 101–113. MR 93m:58026 Zbl 0754.58007
- [He and Shen 2005] Q. He and Y.-B. Shen, “Some results on harmonic maps for Finsler manifolds”, *Internat. J. Math.* **16**:9 (2005), 1017–1031. MR 2006f:53094 Zbl 1085.58012
- [Hélein 1990] F. Hélein, “Régularité des applications faiblement harmoniques entre une surface et une sphère”, *C. R. Acad. Sci. Paris Sér. I Math.* **311**:9 (1990), 519–524. MR 92a:58034 Zbl 0728.35014
- [Hélein 1991a] F. Hélein, “Régularité des applications faiblement harmoniques entre une surface et une variété riemannienne”, *C. R. Acad. Sci. Paris Sér. I Math.* **312**:8 (1991), 591–596. MR 92e:58055 Zbl 0728.35015
- [Hélein 1991b] F. Hélein, “Regularity of weakly harmonic maps from a surface into a manifold with symmetries”, *Manuscripta Math.* **70**:2 (1991), 203–218. MR 92a:58035 Zbl 0718.58019
- [Hélein 2002] F. Hélein, *Harmonic maps, conservation laws and moving frames*, 2nd ed., Cambridge Tracts in Mathematics **150**, Cambridge University Press, Cambridge, 2002. MR 2003g:58024 Zbl 1010.58010
- [Jost 2008] J. Jost, *Riemannian geometry and geometric analysis*, Fifth ed., Springer, Berlin, 2008. MR 2009g:53036 Zbl 1143.53001
- [Lin and Wang 2008] F. Lin and C. Wang, *The analysis of harmonic maps and their heat flows*, World Scientific, Hackensack, NJ, 2008. MR 2011a:58030 Zbl 1203.58004
- [Matsumoto 2003] M. Matsumoto, “Finsler geometry in the 20th-century”, pp. 557–966 in *Handbook of Finsler geometry. Vol. 1, 2*, Kluwer, Dordrecht, 2003. MR 2005e:53023 Zbl 1105.53019
- [Mo 2001] X. Mo, “Harmonic maps from Finsler manifolds”, *Illinois J. Math.* **45**:4 (2001), 1331–1345. MR 2003c:53091 Zbl 0996.53047
- [Mo 2006] X. Mo, *An introduction to Finsler geometry*, Peking University Series in Mathematics **1**, World Scientific, Hackensack, NJ, 2006. MR 2007f:53092 Zbl 1114.53001
- [Mo and Yang 2005] X. Mo and Y. Yang, “The existence of harmonic maps from Finsler manifolds to Riemannian manifolds”, *Sci. China Ser. A* **48**:1 (2005), 115–130. MR 2006e:58015 Zbl 1127.58011
- [Shen and Wei 2008] Y. Shen and S. W. Wei, “The stability of harmonic maps on Finsler manifolds”, *Houston J. Math.* **34**:4 (2008), 1049–1056. MR 2009j:58023 Zbl 1162.53054

[Shen and Zhang 2004] Y. Shen and Y. Zhang, “Second variation of harmonic maps between Finsler manifolds”, *Sci. China Ser. A* **47**:1 (2004), 39–51. MR 2005b:58021 Zbl 1217.53077

[Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, Princeton, NJ, 1993. MR 95c:42002 Zbl 0821.42001

Received August 13, 2010. Revised June 19, 2011.

XIAOHUAN MO
KEY LABORATORY OF PURE AND APPLIED MATHEMATICS
SCHOOL OF MATHEMATICAL SCIENCES
PEKING UNIVERSITY
BEIJING, 100871
CHINA
moxh@pku.edu.cn

LIANG ZHAO
LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS
BEIJING NORMAL UNIVERSITY
BEIJING, 100875
CHINA
liangzhao@bnu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic.

Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 253 No. 1 September 2011

Singularities of the projective dual variety	1
ROLAND ABUAF	
Eigenvalue estimates for hypersurfaces in $\mathbb{H}^m \times \mathbb{R}$ and applications	19
PIERRE BÉRARD, PHILIPPE CASTILLON and MARCOS CAVALCANTE	
Conformal Invariants associated to a measure: Conformally covariant operators	37
SUN-YUNG A. CHANG, MATTHEW J. GURSKY and PAUL YANG	
Compact symmetric spaces, triangular factorization, and Cayley coordinates	57
DEREK HABERMAS	
Automorphisms of the three-torus preserving a genus-three Heegaard splitting	75
JESSE JOHNSON	
The rationality problem for purely monomial group actions	95
HIDETAKA KITAYAMA	
On a Neumann problem with p -Laplacian and noncoercive resonant nonlinearity	103
SALVATORE A. MARANO and NIKOLAOS S. PAPAGEORGIOU	
Minimal ramification in nilpotent extensions	125
NADYA MARKIN and STEPHEN V. ULLOM	
Regularity of weakly harmonic maps from a Finsler surface into an n -sphere	145
XIAOHUAN MO and LIANG ZHAO	
On the sum of powered distances to certain sets of points on the circle	157
NIKOLAI NIKOLOV and RAFAEL RAFAILOV	
Formal geometric quantization II	169
PAUL-ÉMILE PARADAN	
Embedded constant-curvature curves on convex surfaces	213
HAROLD ROSENBERG and MATTHIAS SCHNEIDER	
A topological construction for all two-row Springer varieties	221
HEATHER M. RUSSELL	



0030-8730(201109)253:1;1-8