

*Pacific
Journal of
Mathematics*

**ON THE SUM OF POWERED DISTANCES
TO CERTAIN SETS OF POINTS ON THE CIRCLE**

NIKOLAI NIKOLOV AND RAFAEL RAFAILOV

ON THE SUM OF POWERED DISTANCES TO CERTAIN SETS OF POINTS ON THE CIRCLE

NIKOLAI NIKOLOV AND RAFAEL RAFAILOV

We consider an extremal problem in geometry. Let λ be a real number and let A , B and C be arbitrary points on the unit circle Γ . We give a full characterization of the extremal behavior of the function $f(M, \lambda) = MA^\lambda + MB^\lambda + MC^\lambda$, where M is a point on the unit circle as well. We also investigate the extremal behavior of $\sum_{i=1}^n XP_i$, where the P_i , for $i = 1, \dots, n$, are the vertices of a regular n -gon and X is a point on Γ , concentric to the circle circumscribed around $P_1 \dots P_n$. We use elementary analytic and purely geometric methods in the proof.

1. Introduction

The question of placing electrical charges on a sphere in such a way that the potential energy of the system obtains its extremal values has long been of importance to physics. Problems of this kind have also been considered in classical potential theory.

The planar case of this question is answered by the general solution of placing n points M_i , $i = 1, \dots, n$ on the unit circle in such a way as to obtain the extreme values of the sum

$$\sum_{0 \leq i < j \leq n} |M_i M_j|^\lambda,$$

where λ is a given real number, the concrete case being $\lambda = -1$. (We denote the distance between two points P and Q by $|PQ|$, or by PQ when there is no danger of confusion.)

There is a growing amount of literature on this problem, which we can regard as a discrete analog of questions studied in classical potential theory. This has led to the development of the problem of placing a point M on the unit circle in such a way to obtain the extremal values of

$$\sum_{i=1}^n MM_i^\lambda$$

MSC2010: 52A40.

Keywords: powered distance, unit circle.

for a given point set M_i , $i = 1, \dots, n$. This has proven to be a difficult question and in general it remains open.

Here we consider the case $n = 3$ and prove:

Theorem 1.1. *Let λ be a real number and let A , B and C be arbitrary points on the unit circle Γ .*

- (1) *If $\lambda < 0$, there is always a point M on Γ such that $f(M, \lambda) \leq 2 + 2^\lambda$.*
- (2) *If $\lambda \in [0; 2]$, there is always a point M on Γ such that $f(M, \lambda) \geq 2 + 2^\lambda$.*
- (3) *If $\lambda \in (2; 4)$, there is always a point M on Γ such that $f(M, \lambda) \geq 2\sqrt{3}^\lambda$.*
- (4) *If $\lambda \geq 4$, there is always a point M on Γ such that $f(M, \lambda) \geq 2 + 2^\lambda$.*

These bounds are sharp if and only if A , B and C are the vertices of an equilateral triangle.

Or, equivalently,

- (1) $\min_{A,B,C \in \Gamma} \max_{M \in \Gamma} f(M, \lambda) = 2 + 2^\lambda$ for $\lambda \in [0; 2] \cup [4; \infty)$.
- (2) $\min_{A,B,C \in \Gamma} \max_{M \in \Gamma} f(M, \lambda) = 2\sqrt{3}^\lambda$ for $\lambda \in (2; 4)$.
- (3) $\max_{A,B,C \in \Gamma} \min_{M \in \Gamma} f(M, \lambda) = 2 + 2^\lambda$ for $\lambda < 0$.

In the last case, the order of the maximum and the minimum is reversed. We are not interested in the maximum of the function $f(M, \lambda)$ when $\lambda < 0$, since it is infinity when M approaches A , B or C .

Prior to the present article, the exact extremal values of $f(M, \lambda)$, established in [Stolarsky 1975], were only known for $\lambda \in [0; 2]$. We also give another (more elementary) proof of the results obtained in that article.

The question of the extremal behavior of the function

$$f_n(M, -2) = \sum_{i=1}^n MM_i^{-2}$$

is considered in [Ambrus and Ball 2011], where it is proved that there always exists a point $M \in \Gamma$ such that

$$f_n(M, -2) \leq \frac{1}{4}n^2.$$

This bound is sharp if and only if the M_i are the vertices of a regular n -gon. This agrees with our results when $n = 3$.

We also consider the case when there are n points P_i , $i = 1, \dots, n$ on the unit circle that are the vertices of a regular n -gon, and a point X on a circle Γ , concentric to the circumscribed circle of $P_1 \dots P_n$. We study the extremal values of

$$R_n(X, \lambda) = \sum_{i=1}^n XP_i^\lambda,$$

where $X \in \Gamma$.

This problem was considered by Stolarsky [1975], who solved it for $0 \leq \lambda < 2n$, where Γ is the circumscribed circle of the polygon, and by Mushkarov [1982], who found λ for which the sum does not depend on the position of X on Γ , again where Γ is circumscribed around $P_1 \dots P_n$, and gave a trigonometric representation for higher powers. In this paper we characterize the extremal behavior of the sum $\sum_{i=1}^n XP_i^\lambda$ and prove:

Theorem 1.2. *Let $P_i, i = 1, \dots, n$ be the vertices of a regular n -gon inscribed in the unit circle. Now let Γ be a circle concentric to the circumscribed circle. Put $B_i = OP_i \cap \Gamma$, where O is the center of the n -gon.*

Let $X \in \Gamma$ and

$$R_n(X, \lambda) = \sum_{i=1}^n XP_i^\lambda.$$

(1) *Suppose $\lambda < 0$. The minimum of $R_n(X, \lambda)$ is achieved when X bisects the arc between consecutive vertices of $B_1 \dots B_n$, and the maximum when $X \equiv B_i$. In the case when Γ is the circumscribed circle around $P_1 \dots P_n$, this function is not bounded when $X \rightarrow B_i$ for some i .*

(2) *Suppose $0 \leq \lambda < 2n$. If λ is an even integer, then $R_n(X, \lambda)$ is independent of the position of X on Γ .*

Otherwise let m be such an integer that $2m \leq \lambda \leq 2m + 2$.

If m is even (resp. odd), then $R_n(X, \lambda)$ is maximal (resp. minimal) if and only if X bisects the arc between consecutive vertices of $B_1 \dots B_n$. Also $R_n(X, \lambda)$ is minimal (resp. maximal) if and only if $M \equiv B_i$.

(3) *$2n \leq \lambda$. If n is even (resp. odd), the maximum (resp. minimum) of $R_n(X, \lambda)$ is obtained when X coincides with one of the vertices of $B_1 \dots B_n$, and the minimum (resp. maximum) is achieved when X bisects the arc between consecutive vertices.*

Remark 1.3. A case of conclusion (2) of the theorem is proved in [Stolarsky 1975], when Γ is the circle circumscribed around $P_1 \dots P_n$. However, it seems possible that the general result of (2) can be proved in the same manner.

It is easy to see that (3) is actually true for $\lambda > 2n - 2$.

We begin by considering the regular n -gon, since we use the result later.

2. The regular n -gon

We say that y is a root of degree k of an equation $f(x) = 0$, where f is k -times differentiable, if $f(y) = 0$ and $f^t(y) = 0$ for $t = 1, \dots, k - 1$ and $f^k(y) \neq 0$, where $f^t(x)$ denotes the t -th derivative of f .

Lemma 2.1. *Let a_1, a_2, \dots, a_n be real numbers and b_1, b_2, \dots, b_n be nonnegative numbers. Then the function*

$$\varphi(\lambda) = \sum_{i=1}^n a_i b_i^\lambda$$

is either identically zero or has at most $n - 1$ real zeros counted with their multiplicities.

Proof. We proceed by induction on the number of summands. For $n = 1$, we have $ab^\lambda = 0$, which does not have solutions if both a and b are nonzero. If either of them is zero, then ab^λ is identically zero. Now assume the statement to be true for all $k < n$. For $k = n$, if either of a_i or b_i is zero, then we use the induction hypothesis.

Now let b_i, a_i be nonzero. Since all of b_i are nonzero, we can divide each term by b_1^λ to get

$$\sum_{i=1}^n a_i \left(\frac{b_i}{b_1}\right)^\lambda = 0.$$

Assume that this equation is not identically zero, and that its solutions are y_1, \dots, y_k with multiplicities t_1, \dots, t_k and $\sum_{i=1}^k t_i > n - 1$.

Differentiating this with respect to λ , we get

$$\sum_{i=2}^n a_i \ln \frac{b_i}{b_1} \left(\frac{b_i}{b_1}\right)^\lambda = 0 = \sum_{i=2}^n a'_i b_i'^\lambda,$$

where $a'_i = a_i \ln(b_i/b_1)$ and $b'_i = b_i/b_1$. Assume that this expression is identically zero; then $\sum_{i=1}^n a_i b_i^\lambda = 0$ must be a constant, and the claim follows. Assume that the derivative does not vanish for all λ . Now by the induction hypothesis, the derivative has at most $n - 2$ zeros. But y_1, \dots, y_k are solutions to the above equation with multiplicities $t_1 - 1, \dots, t_k - 1$; also, by Rolle's theorem, the derivative has at least one root in each interval (y_i, y_{i+1}) , and thus we obtain $k - 1 + \sum_{i=1}^k t_i - 1$ solutions (counted with their multiplicities), which is greater than $n - 2$; a contradiction. It follows that $\sum_{i=1}^k t_i \leq n - 1$. \square

We continue with another problem, which is a part of [Theorem 1.2](#).

Theorem 2.2. *Let P_1, \dots, P_n be the vertices of a regular polygon, given a circle Γ , concentric to the circle circumscribed around $P_1 P_2 \dots P_n$; then $\sum_{i=1}^n PP_i^{2k}$ is independent of the position of $P \in \Gamma$ for $k \in \{1, \dots, n - 1\}$.*

Proof. We use complex numbers. We may assume that the circumscribed circle around $P_1 \dots P_n$ is the unit circle and that the radius of Γ is R . Let us assign to the vertices of the n -gon the complex numbers ξ, ξ^2, \dots, ξ^n , where ξ is a primitive

n -th root of unity. We wish to prove that $\sum_{i=1}^n |x - \xi^i|^{2k} = \text{const}$ for all x with a fixed norm R and all $k \in \{1, \dots, n-1\}$. We have

$$\sum_{i=1}^n |x - \xi^i|^{2k} = \sum_{i=1}^n (x - \xi^i)^k \overline{(x - \xi^i)^k} = \sum_{i=1}^n (x - \xi^i)^k \left(\frac{R}{x} - \frac{1}{\xi^i}\right)^k.$$

After multiplying, out we obtain

$$(x - \xi^i)^k \left(\frac{R}{x} - \frac{1}{\xi^i}\right)^k = \sum_{j=-k}^k c_j \xi^{-ij} x^j = P_i(x)$$

for all x with $|x| = R$. We now have

$$\sum_{i=1}^n |x - \xi^i|^{2k} = \sum_{i=1}^n P_i(x) = \sum_{j=-k}^k \sum_{i=1}^n c_j \xi^{-ij} x^j,$$

but $\sum_{i=1}^n \xi^{-ij} = 0$ for all j except $j = mn$, where m is an integer, so

$$\sum_{i=1}^n |x - \xi^i|^{2k} = nc_0. \quad \square$$

Remark 2.3. One can prove that this is a characteristic property of the regular n -gon. Given n different points in the plane A_1, \dots, A_n and a circle Γ , such that $\sum_{i=1}^n PA_i^{2k}$ is independent of the position of P on Γ for every $k \in \{1, \dots, n-1\}$, then these points are the vertices of a regular n -gon. It is conjectured that this remains true if the condition holds only for $k = 2n - 2$, and this has been verified for $n = 3$ and $n = 4$, but the authors have no proof for higher values of n .

Proof of Theorem 1.2. We may assume $X \in \widehat{B_1 M}$, where M is the midpoint of arc $\widehat{B_1 B_2}$, because of symmetry.

After we position ourselves in a Cartesian coordinate system without loss of generality, we can assume that P_1 has coordinates $(1, 0)$. Thus the coordinates of P_i are $(\cos((i-1)2\pi/n), \sin((i-1)2\pi/n))$, and X has coordinates $(a \cos x, a \sin x)$, where $x \in [0; 2\pi/n]$.

We can now write the sum

$$\begin{aligned} \sum_{i=1}^n |P_i X|^\lambda &= \sum_{i=1}^n \left(\left(a \cos x - \cos \frac{(i-1)2\pi}{n} \right)^2 + \left(a \sin x - \sin \frac{(i-1)2\pi}{n} \right)^2 \right)^{\lambda/2} \\ &= F(x, \lambda). \end{aligned}$$

We differentiate this with respect to x to obtain

$$\frac{\partial F(x, \lambda)}{\partial x} = \sum_{i=1}^n \lambda |P_i X|^{\lambda-1} \frac{d|P_i X|}{dx}.$$

The partial derivative exists for $x \in (0; 2\pi/n)$. Now fix x and consider this as a function of λ . As we are interested only in the sign of the derivative, we can consider only

$$\sum_{i=1}^n |P_i X|^{\lambda-1} \frac{d|P_i X|}{dx}$$

for $\lambda \neq 0$.

As proved earlier, $F(x, \lambda)$ is constant for $\lambda = 2, 4, \dots, 2n-2$, so $\partial F(x, \lambda)/\partial x$ vanishes for these values of λ . But from [Lemma 2.1](#), this expression is either identically zero or has at most $n-1$ solutions for λ , counted with their multiplicities.

We prove that this expression as a function of λ is not identically zero for fixed $x \in (0; 2\pi/n)$. For the sake of contradiction, assume otherwise. Let $x \in (0; 2\pi/n)$. It is easy to see that for the point X corresponding to this x , the distances $|P_i X|$ are all different. Now take i such that $|P_i X|$ is maximized subject to the condition that $d|P_i X|/dx$ is nonzero. If $|P_i X| > 1$, then

$$\lim_{\lambda \rightarrow \infty} \left| \frac{\partial F(x, \lambda)}{\partial x} \right| = \infty.$$

On the other hand, if $|P_i X| \leq 1$, choose i instead so that $|P_i X|$ is minimized subject to the condition that $d|P_i X|/dx$ is nonzero. In this case $|P_i X| < 1$; otherwise there must be two distances $P_i X$ that are equal (or else $n-1$ among the $\widehat{d|P_i X|/dx}$ vanish, which is not possible). Indeed, as X describes the arcs $\widehat{P'_i P''_i}$ and $\widehat{P''_i P'_i}$, where P'_i, P''_i are the two intersections of Γ with the line through O and P_i , the distance $P_i X$ increases along one arc and decreases along the other. It is obvious that P'_i and P''_i either coincide with some of B_i or are midpoints of some arc between consecutive vertices of $B_1 \dots B_n$.

Again considering $\lim_{\lambda \rightarrow -\infty} |\partial F(x, \lambda)/\partial x| = \infty$, we obtain the desired result.

As mentioned, for a fixed x , $\partial F(x, \lambda)/\partial x$ vanishes for every $\lambda = 0, 2, 4, \dots, 2n-2$, and it follows that these are all the solutions for λ and that each of them—except possibly $\lambda = 0$ —must have multiplicity one. Also the derivative changes sign at $\lambda = 0$. For a fixed x , therefore, $\partial F(x, \lambda)/\partial x$ changes sign at $\lambda = 0, 2, 4, \dots, 2n-2$. Now assume that for some $\lambda_0 \neq 0, 2, 4, \dots, 2n-2$, there exist y and z in the interval $(0; 2\pi/n)$ such that

$$\frac{\partial F(y, \lambda_0)}{\partial x} \frac{\partial F(z, \lambda_0)}{\partial x} < 0;$$

then since $\partial F(x, \lambda_0)/\partial x$ is a continuous function of x , there is $t \in (0; 2\pi/n)$ such that $\partial F(t, \lambda_0)/\partial x = 0$, and it follows that $\partial F(t, \lambda)/\partial x = 0$ for all λ , which is a contradiction. Hence the derivative

$$\frac{\partial F(x, \lambda)}{\partial x}$$

does not change when $x \in (0; 2\pi/n)$ for fixed λ ; also, for every fixed $x \in (0, 2\pi/n)$ it changes sign at $\lambda = 0, 2, \dots, 2n-2$. Thus, since $F(x, \lambda)$ is a continuous function of x , we have shown that the minimum and maximum of that function for $x \in [0; 2\pi/n]$ are obtained when $x = 0$, or $x = 2\pi/n$.

Now consider

$$\lim_{\lambda \rightarrow \infty} \frac{F(0, \lambda)}{F(2\pi/n, \lambda)}.$$

Assume that n is even; then $|B_1 P_{n/2+1}| > |MP_i|$ for every i , and the limit is ∞ .

Assume that n is odd; then $|MP_{\lceil n/2 \rceil}| > |B_1 P_i|$ for every i , and the above limit becomes 0. This proves **part (3)** of **Theorem 1.2**. Now, taking into account parity and the above observations for the intervals in which $\partial F(x, \lambda)/\partial x$ changes sign, the conclusion of the theorem easily follows. \square

Remark 2.4. When Γ is the circumcircle of the regular polygon, **Theorem 1.2(1)** is easily proved by the observation that each of the functions $MP_i^\lambda + MP_{n+1-i}^\lambda$ is concave.

3. The case for three base points (proof of **Theorem 1.1**)

Proof of the case $\lambda < 0$. We now consider the case $\lambda < 0$.

Let $\angle C = \max\{\angle A, \angle B, \angle C\}$, and let M be the midpoint of the smaller arc \widehat{AB} . We prove that $f(M, \lambda) \leq 2 + 2^\lambda$. We consider two cases:

Case 1: $\angle C \geq \pi/2$. Then the maximum of the function $MA^\lambda + MB^\lambda + MC^\lambda$ when C travels along the smaller arc \widehat{AB} is achieved when $C \equiv A$ or when $C \equiv B$, since $MD > MA = MB$ for any point D on the smaller arc \widehat{AB} . Now we have $f(M, \lambda) = 3MA^\lambda \leq 3\sqrt{2}^\lambda < 2^\lambda + 2$.

Case 2: $\angle C = x < \pi/2$. Then $\angle C \in [\pi/3; \pi/2)$. Now let C' and C'' be the points for which $\angle ABC'$ and $\angle BAC''$ equal x . It is easy to see that C belongs to the smaller arc $\widehat{C'C''}$, since $\angle C$ is the largest angle of the triangle. Also, the maximum of $f(M, \lambda)$ when C belongs to the arc $\widehat{C'C''}$ is obtained exactly when $C \equiv C'$ or $C \equiv C''$, since $MC'' = MC' \leq MC$ for every C on $\widehat{C'C''}$. Without loss of generality we can assume that $C \equiv C'$. Then we can write

$$f(M, \lambda) = 2(2 \sin \frac{1}{2}x)^\lambda + (2 \sin \frac{3}{2}x)^\lambda = 2^\lambda (2 \sin^\lambda \frac{1}{2}x + \sin^\lambda \frac{3}{2}x) = F(x, \lambda).$$

We differentiate with respect to x to get

$$\frac{\partial F(x, \lambda)}{\partial x} = \lambda 2^\lambda (\sin^{\lambda-1} \frac{1}{2}x \cos \frac{1}{2}x + \frac{3}{2} \sin^{\lambda-1} \frac{3}{2}x \cos \frac{3}{2}x).$$

Clearly, both $\sin^{\lambda-1} \frac{1}{2}x \cos \frac{1}{2}x$ and $\frac{3}{2} \sin^{\lambda-1} \frac{3}{2}x \cos \frac{3}{2}x$ are decreasing functions in the interval $[\pi/3; \pi/2)$, since $\lambda < 0$.

Then $\partial F(x, \lambda)/\partial x$ is an increasing function of x in this interval since $\lambda < 0$, and hence $F(x, \lambda)$ is a convex function of $x \in [\pi/3; \pi/2)$ if $\lambda < 0$.

From here it follows that

$$\begin{aligned} \sup_{x \in [\pi/3; \pi/2)} F(x, \lambda) &= \max\{F(\pi/3, \lambda), \lim_{x \rightarrow \pi/2} F(x, \lambda)\} \\ &= \max\{2 + 2^\lambda, 3\sqrt{2}^\lambda\}, \end{aligned}$$

since $F(x, \lambda)$ is continuous in the interval $[\pi/3; \pi/2]$. Then $\max\{2 + 2^\lambda, 3\sqrt{2}^\lambda\} = 2 + 2^\lambda$, and this bound is achievable only for ABC an equilateral triangle; for all other configurations of the points ABC , the function $MA^\lambda + MB^\lambda + MC^\lambda$ is less than $2 + 2^\lambda$ for the specified point M .

Now using [Theorem 1.2](#), we get that the minimum of $MA^\lambda + MB^\lambda + MC^\lambda$ is obtained when M bisects the arc between consecutive vertices of the triangle, and in this case we have $MA^\lambda + MB^\lambda + MC^\lambda = 2 + 2^\lambda$. This concludes the proof.

Remark 3.1. This case can also be proved using the main approach of [[Ambrus and Ball 2011](#)], which is based on Lemma 1 of that reference, namely, the fact that the local minima on each of the arcs between consecutive base points must be equal for all $\lambda < 0$. In the case of only three points, one can obtain that the equilateral triangle is indeed the extremal case. Assume otherwise. Clearly it is not possible for all of the local minima to be equal when two of the points are closer than $\sqrt{2}$. Assume now that C is not the midpoint of the arc \widehat{AB} . Consider the function

$$f_1 = |MA|^\lambda + |MB|^\lambda + |MC_1|^\lambda,$$

where C_1 is the midpoint of \widehat{AB} . We may assume that C belongs to the shorter arc C_1B . By symmetry, the local minima of f_1 are equal on the short arcs $\widehat{AC_1}$ and $\widehat{BC_1}$. Now we have $\angle C_1OC \leq \pi/4$. Thus $f > f_1$ on $\widehat{AC_2}$ and $f < f_1$ on $\widehat{BC_2}$, where C_2 is the midpoint of the arc $\widehat{CC_1}$. From here we obtain that the local minima of f cannot be equal on the shorter arcs \widehat{AC} and \widehat{BC} .¹

Proof of the case $\lambda > 2$. We first prove that for every three points A, B and C on the unit circle there exists a point M also on the unit circle, such that $f(M, \lambda) \geq \max\{2 + 2^\lambda, 2(\sqrt{3})^\lambda\}$.

Let $AB = \min\{AB, BC, CA\}$; now let the bisector of AB intersect the larger arc AB at M' . Then $\angle BAM' = \angle ABM' = x$ and $\pi/3 \leq x < \pi/2$. Now by the sine rule, $BM' = AM' = 2 \sin x \geq \sqrt{3}$, and we have $f(M', \lambda) \geq 2(\sqrt{3})^\lambda$ with equality only if $x = \pi/3$, or, equivalently, if ABC is an equilateral triangle.

It remains to prove that for every triangle there is a point M' such that $f(M', \lambda) \geq 2 + 2^\lambda$. We consider two cases — when ABC is obtuse-angled and when it is acute-angled.

¹The authors thank the referee for suggesting this approach.

Case 1: Let $\angle C = \max\{\angle A, \angle B, \angle C\} \geq \pi/2$, and let O be the center of Γ . Let $M' = CO \cap \Gamma$. We have $CM' = 2$ and $f(M', \lambda) = 2^\lambda + BM'^\lambda + AM'^\lambda$. Now $\angle BAM', \angle ABM' \leq \pi/2$ and $\angle BAM' + \angle ABM' \geq \pi/2$.

We have $\pi/4 \leq \max\{\angle BAM', \angle ABM'\} \leq \pi/2$, and so

$$BM'^\lambda + AM'^\lambda > (2.1/\sqrt{2})^\lambda \geq 2$$

as $\lambda \geq 2$, so $f(M', \lambda) > 2 + 2^\lambda$. And this bound cannot be achieved for an obtuse-angled triangle.

Case 2: Let $c = \angle C = \max\{\angle A, \angle B, \angle C\} < \pi/2$. Now $M' = CO \cap \Gamma$. We have $CM' = 2$ and $f(M', \lambda) = 2^\lambda + BM'^\lambda + AM'^\lambda$. We prove that $BM'^\lambda + AM'^\lambda \geq 2$. Let $\angle ACM' = x$. By the sine rule, $BM'^\lambda + AM'^\lambda = (2 \sin x)^\lambda + (2 \sin(c - x))^\lambda = f_1(x, \lambda)$. We prove that $\sin^\lambda x + \sin^\lambda(c - x) > 2 \sin^\lambda(c/2)$. We have

$$\begin{aligned} \frac{\partial f_1(x, \lambda)}{\partial x} &= \lambda 2^\lambda (\sin^{\lambda-1} x \cos x - \sin^{\lambda-1}(c - x) \cos(c - x)) \\ &= \lambda 2^{\lambda-1} (\sin^{\lambda-2} x \sin 2x - \sin^{\lambda-2}(c - x) \sin(2c - 2x)). \end{aligned}$$

It is now easy to see that for $x \in [0; c/2)$, we have $\sin^{\lambda-2} x < \sin^{\lambda-2}(c - x)$ (for $\lambda \geq 2$) and $\sin 2x < \sin(2c - 2x)$, and so

$$\frac{\partial f_1(x, \lambda)}{\partial x} < 0.$$

With similar arguments it follows that $\frac{\partial f_1(x, \lambda)}{\partial x} = 0$ for $x = \frac{c}{2}$, and $\frac{\partial f_1(x, \lambda)}{\partial x} > 0$ for $x \in (c/2; c]$.

Then $\min_{x \in [0; c]} (f_1(x, \lambda) + f_1(c - x, \lambda)) = 2f_1(c/2, \lambda) \geq 2f_1(\pi/6, \lambda) = 2$. Equality holds if and only if ABC is equilateral.

We have $2 + 2^\lambda = 2\sqrt{3}^\lambda$ for $\lambda \in \{2, 4\}$; $2 + 2^\lambda < 2\sqrt{3}^\lambda$ for $\lambda \in (2; 4)$; and $2 + 2^\lambda > 2\sqrt{3}^\lambda$ for $\lambda > 4$. We prove that for ABC an equilateral triangle, those bounds are sharp.

Again using [Theorem 1.2](#), we get:

- (1) When $\lambda \in [2; 4]$, the maximum of $MA^\lambda + MB^\lambda + MC^\lambda$ is achieved when M coincides with one of A, B, C and is equal to $2\sqrt{3}^\lambda$.
- (2) When $\lambda > 4$, the maximum of $MA^\lambda + MB^\lambda + MC^\lambda$ is achieved when M bisects the arc between consecutive vertices of the triangle ABC and is equal to $2 + 2^\lambda$.

These bounds are sharp.

We also have the minimum of $f(M, \lambda)$ when ABC is an equilateral triangle, that is, when $\lambda \in [2, 4]$, we have $\min f(M, \lambda) = 2 + 2^\lambda$; and $\min f(M, \lambda) = 2\sqrt{3}^\lambda$ when $\lambda > 4$. This concludes the proof. \square

Proof of the case $\lambda \in [0; 2]$. This case is proved in [Stolarsky 1975], but we give a new, independent proof.

We now prove that for every three points A, B, C on the unit circle and a real number $\lambda \in (0; 2)$, there exists a point M also on the unit circle, such that $f(M, \lambda) = MA^\lambda + MB^\lambda + MC^\lambda \geq 2 + 2^\lambda$; and this bound is sharp. It is only achievable when A, B and C are the vertices of an equilateral triangle.

Again let $\angle C = \max\{\angle A, \angle B, \angle C\} = x$. As before, it is easy to see that when $\angle C \geq \pi/2$ (with M chosen as the midpoint of the arc \widehat{AB}), we have

$$f(M, \lambda) \geq 3\sqrt{2}^\lambda \geq 2 + 2^\lambda$$

for $\lambda \in [0; 2]$. When $\lambda \in \{0, 2\}$, we have that $MA^\lambda + MB^\lambda + MC^\lambda$ is constant. We can assume that the triangle ABC is acute-angled. Then $\angle C \in [\pi/3; \pi/2]$, and again let M be the midpoint of the arc \widehat{AB} . Now let C' and C'' be the points for which $\angle ABC'$ and $\angle BAC''$ respectively equal x . It is easy to see that C belongs to the smaller arc $\widehat{C'C''}$, since C is the largest angle of the triangle. It is also easy to see that the minimum of $f(M, \lambda)$ when C belongs to the arc $\widehat{C'C''}$ is obtained exactly when $C \equiv C'$ or $C \equiv C''$, since $MC'' = MC' \leq MC$ for every C on $\widehat{C'C''}$.

Let $\angle A = x$. Using the sine rule, we now get

$$f(M, \lambda) = MA^\lambda + MB^\lambda + MC^\lambda = 2(2 \sin \frac{1}{2}x)^\lambda + (\sin \frac{3}{2}x)^\lambda = F(x, \lambda).$$

We prove that if $x \in [\pi/3; \pi/2]$, then $F(x, \lambda) \geq 2 + 2^\lambda$.

We consider two cases:

Case 1: $\lambda \in (0; 1)$. After differentiating with respect to x , we get

$$\frac{\partial F(x, \lambda)}{\partial x} = \lambda 2^\lambda (\sin^{\lambda-1} \frac{1}{2}x \cos \frac{1}{2}x + \frac{3}{2} \sin^{\lambda-1} \frac{3}{2}x \cos \frac{3}{2}x).$$

It is now easy to see that both $\sin^{\lambda-1} \frac{1}{2}x \cos \frac{1}{2}x$ and $\sin^{\lambda-1} \frac{3}{2}x \cos \frac{3}{2}x$ are decreasing functions as $\lambda - 1 < 0$. Then $F(x, \lambda)$ is a concave function of x when $\lambda \in (0; 1)$. It follows that

$$\begin{aligned} \min_{x \in [\pi/3; \pi/2]} F(x, \lambda) &= \min \left\{ F(\pi/3, \lambda), \lim_{x \rightarrow \pi/2} F(x, \lambda) \right\} \\ &= F(\pi/3; \lambda) = 2 + 2^\lambda, \end{aligned}$$

and for every $x \neq \pi/3$ we have $F(x, \lambda) > 2 + 2^\lambda$. We shall later prove that when A, B, C are the vertices of an equilateral triangle this bound is sharp.

Case 2: $\lambda \in [1; 2)$. Let $CO \cap \Gamma = M$. We shall prove that for the point M we have $AM^\lambda + BM^\lambda + CM^\lambda \geq 2 + 2^\lambda$. We have $CM = 2^\lambda$. We only need to prove that $BM + CM \geq 2$, since

$$BM^\lambda + AM^\lambda \geq 2 \left(\frac{AM + BM}{2} \right)^\lambda.$$

Lemma 3.2. *Let A, B and C be points on the unit circle Γ with center O . Assume $\angle C = \max\{\angle A, \angle B, \angle C\}$ and $M = CO \cap \Gamma$ then $MA + MB \geq 2$.*

Proof. We have

$$MA + MB = 2(\sin x + \sin(c - x)) =: f(x),$$

where $x = \angle MAC$ and $c = \pi - \angle ACB < \pi/2$. Then $f'(x) = 2(\cos x - \cos(c - x))$; thus $f'(x) > 0$ for $x \in [0; c/2)$, while $f'(x) = 0$ for $x = c/2$ and $f'(x) < 0$ for $x \in (c/2; c]$.

Now let C' and C'' be the points for which $\angle ABC'$ and $\angle BAC''$ respectively equal x . It is easy to see that C belongs to the smaller arc $\widehat{C'C''}$ as C is the largest angle of the triangle. The minimum $\min MA^\lambda + MB^\lambda$ is obtained when $MO \cap \Gamma = C'$ or $MO \cap \Gamma = C''$ as $f(x)$ is concave. Now $\angle ABC = \angle BCA = \gamma$. Then $MA + MB = 2(\sin(\pi/2 - \gamma) + \sin(2\gamma - \pi/2)) = \cos \gamma - \cos 2\gamma = f_1(\gamma)$. Differentiating $f_1(\gamma)$ we get $f_1'(\gamma) = 2 \sin 2\gamma - \sin \gamma$ which is a decreasing function of $\gamma \in [\pi/3; \pi/2)$. This gives us that $f_1(\gamma)$ is a concave function when $\gamma \in [\pi/3; \pi/2)$, and it follows that

$$\min_{\gamma \in [\pi/3; \pi/2)} f_1(\gamma) = \min \left\{ f_1(\pi/3), \lim_{x \rightarrow \pi/2} f_1(x) \right\} = 2. \quad \square$$

Now

$$BM^\lambda + AM^\lambda \geq 2 \left(\frac{AM + BM}{2} \right)^\lambda \geq 2,$$

with equality only when $AM = BM = 1$ which is possible only when A, B and C are the vertices of an equilateral triangle.

In such a way we obtain that when $\lambda \in (0; 2)$ there exists a point M on the unit circle, such that $MA^\lambda + MB^\lambda + MC^\lambda \geq 2 + 2^\lambda$ and this bound is achievable only if A, B and C are the vertices of an equilateral triangle.

Now using again the result of [Theorem 1.2](#) one easily obtains that the maximum of $MA^\lambda + MB^\lambda + MC^\lambda$ is obtained when M is the midpoint of one of the arcs between consecutive vertices and it indeed equals $2 + 2^\lambda$.

Acknowledgement

The authors would like to thank the referee for his comments and remarks, which helped improve the clarity of the paper.

References

- [Ambrus and Ball 2011] G. Ambrus and K. Ball, “Chebyshev constants for the unit circle”, preprint, 2011. [arXiv 1006.5153v2](#)
- [Mushkarov 1982] O. Mushkarov, “Тригонометрични полиноми и правилни многоъгълници”, *Matematika* (Sofia) **8** (1982), 11–16. Available online as a supplement to this paper. An

English summary by J. Tabov can be found in “[Sums of powers of distances in regular polygons and polyhedra](#)”, *James Cook Math. Notes* **5**:50 (1989), page 5209, available at www.maths.ed.ac.uk/cook/iss_no50.PDF.

[Stolarsky 1975] K. B. Stolarsky, “[The sum of the distances to certain pointsets on the unit circle](#)”, *Pacific J. Math.* **59**:1 (1975), 241–251. [MR 52 #9084](#) [Zbl 0313.52006](#)

Received December 27, 2010. Revised July 16, 2011.

NIKOLAI NIKOLOV
INSTITUTE OF MATHEMATICS AND INFORMATICS
BULGARIAN ACADEMY OF SCIENCES
ACAD. G. BONCHEV 8, 1113 SOFIA
BULGARIA

nik@math.bas.bg

RAFAEL RAFAILOV
SOFIA HIGH SCHOOL OF MATHEMATICS
ISKAR 61, 1000 SOFIA
BULGARIA

rafael.rafilov@yahoo.com

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 253 No. 1 September 2011

Singularities of the projective dual variety	1
ROLAND ABUAF	
Eigenvalue estimates for hypersurfaces in $\mathbb{H}^m \times \mathbb{R}$ and applications	19
PIERRE BÉRARD, PHILIPPE CASTILLON and MARCOS CAVALCANTE	
Conformal Invariants associated to a measure: Conformally covariant operators	37
SUN-YUNG A. CHANG, MATTHEW J. GURSKY and PAUL YANG	
Compact symmetric spaces, triangular factorization, and Cayley coordinates	57
DEREK HABERMAS	
Automorphisms of the three-torus preserving a genus-three Heegaard splitting	75
JESSE JOHNSON	
The rationality problem for purely monomial group actions	95
HIDETAKA KITAYAMA	
On a Neumann problem with p -Laplacian and noncoercive resonant nonlinearity	103
SALVATORE A. MARANO and NIKOLAOS S. PAPAGEORGIU	
Minimal ramification in nilpotent extensions	125
NADYA MARKIN and STEPHEN V. ULLOM	
Regularity of weakly harmonic maps from a Finsler surface into an n -sphere	145
XIAOHUAN MO and LIANG ZHAO	
On the sum of powered distances to certain sets of points on the circle	157
NIKOLAI NIKOLOV and RAFAEL RAFAILOV	
Formal geometric quantization II	169
PAUL-ÉMILE PARADAN	
Embedded constant-curvature curves on convex surfaces	213
HAROLD ROSENBERG and MATTHIAS SCHNEIDER	
A topological construction for all two-row Springer varieties	221
HEATHER M. RUSSELL	