

*Pacific
Journal of
Mathematics*

**INTERLACING LOG-CONCAVITY
OF THE BOROS–MOLL POLYNOMIALS**

WILLIAM Y. C. CHEN, LARRY X. W. WANG AND ERNEST X. W. XIA

INTERLACING LOG-CONCAVITY OF THE BOROS–MOLL POLYNOMIALS

WILLIAM Y. C. CHEN, LARRY X. W. WANG AND ERNEST X. W. XIA

We say a sequence $\{P_m(x)\}_{m \geq 0}$ of polynomials of degree m with positive coefficients is interlacingly log-concave if the ratios of consecutive coefficients of $P_m(x)$ interlace the ratios of consecutive coefficients of $P_{m+1}(x)$ for any $m \geq 0$. Interlacing log-concavity of a sequence of polynomials is stronger than log-concavity of the polynomials themselves. We show that the Boros–Moll polynomials are interlacingly log-concave. Furthermore, we give a sufficient condition for interlacing log-concavity which implies that some classical combinatorial polynomials are interlacingly log-concave.

1. Introduction

Let $\{P_m(x)\}_{m \geq 0}$ be a sequence of polynomials, where

$$P_m(x) = \sum_{i=0}^m a_i(m)x^i$$

is a polynomial of degree m . Let

$$r_i(m) = \frac{a_i(m)}{a_{i+1}(m)}.$$

We say that the sequence of polynomials $\{P_m(x)\}_{m \geq 0}$ is *interlacingly log-concave* if the ratios $r_i(m)$ interlace the ratios $r_i(m+1)$, that is,

$$\begin{aligned} r_0(m+1) \leq r_0(m) \leq r_1(m+1) \leq r_1(m) \\ \leq \cdots \leq r_{m-1}(m+1) \leq r_{m-1}(m) \leq r_m(m+1). \end{aligned}$$

Recall that a sequence $\{a_i\}_{0 \leq i \leq m}$ of positive numbers is said to be log-concave if

$$\frac{a_0}{a_1} \leq \frac{a_1}{a_2} \leq \cdots \leq \frac{a_{m-1}}{a_m}.$$

It is obvious that interlacing log-concavity implies log-concavity.

This work was supported by the 973 Project, the PCSIRT Project, the Doctoral Program Fund of the Ministry of Education, and the National Science Foundation of China.

MSC2000: primary 05A20; secondary 33F10.

Keywords: interlacing log-concavity, log-concavity, Boros–Moll polynomial.

The main objective of this paper is to prove the interlacing log-concavity of the Boros–Moll polynomials. For the background on these polynomials, see [Boros and Moll 1999a; 1999b; 1999c; 2001; 2004; Moll 2002; Amdeberhan and Moll 2009]. From now on, we use $P_m(x)$ to denote the Boros–Moll polynomial given by

$$(1) \quad P_m(x) = \sum_{j,k} \binom{2m+1}{2j} \binom{m-j}{k} \binom{2k+2j}{k+j} \frac{(x+1)^j (x-1)^k}{2^{3(k+j)}}.$$

Boros and Moll [1999b] derived the following formula for the coefficient $d_i(m)$ of x^i in $P_m(x)$:

$$(2) \quad d_i(m) = 2^{-2m} \sum_{k=i}^m 2^k \binom{2m-2k}{m-k} \binom{m+k}{k} \binom{k}{i}.$$

In [Boros and Moll 1999c], they showed that the sequence $\{d_i(m)\}_{0 \leq i \leq m}$ is unimodal and that the maximum element appears in the middle. In other words,

$$(3) \quad d_0(m) < d_1(m) < \cdots < d_{\lfloor m/2 \rfloor}(m) > d_{\lfloor m/2 \rfloor - 1}(m) > \cdots > d_m(m).$$

They also established the unimodality by a different approach [Boros and Moll 1999a]; see also [Alvarez et al. 2001].

Moll [2002] conjectured that the sequence $\{d_i(m)\}_{0 \leq i \leq m}$ is log-concave. Kauers and Paule [2007] proved this conjecture based on recurrence relations found by using a computer algebra approach. Chen and Xia [2009] showed that the sequence $\{d_i(m)\}_{0 \leq i \leq m}$ satisfies the ratio monotone property which implies log-concavity and the spiral property. A combinatorial proof of the log-concavity of $P_m(x)$ was found by Chen, Pang and Qu [≥ 2011].

In addition to the Boros–Moll polynomials, we study polynomials whose coefficients satisfy triangular recurrence relations. It is easy to show that the binomial coefficients, the Narayana numbers and the Bessel numbers are interlacingly log-concave. We also give a sufficient condition for the interlacing log-concavity of a sequence of polynomials and prove that the rising factorials, the Bell polynomials and the Whitney polynomials are interlacingly log-concave.

2. The interlacing log-concavity of $d_i(m)$

In this section, we show that for $m \geq 2$, the Boros–Moll polynomials $P_m(x)$ are interlacingly log-concave.

Theorem 2.1. *For $m \geq 2$ and $0 \leq i \leq m$, we have*

$$(4) \quad d_i(m)d_{i+1}(m+1) > d_{i+1}(m)d_i(m+1),$$

$$(5) \quad d_i(m)d_i(m+1) > d_{i-1}(m)d_{i+1}(m+1).$$

The proof relies on recurrence relations derived in [Kauers and Paule 2007]:

$$(6) \quad d_i(m+1) = \frac{m+i}{m+1} d_{i-1}(m) + \frac{(4m+2i+3)}{2(m+1)} d_i(m), \quad 0 \leq i \leq m+1,$$

$$(7) \quad d_i(m+1) = \frac{(4m-2i+3)(m+i+1)}{2(m+1)(m+1-i)} d_i(m) \\ - \frac{i(i+1)}{(m+1)(m+1-i)} d_{i+1}(m), \quad 0 \leq i \leq m,$$

$$(8) \quad d_i(m+2) = \frac{-4i^2+8m^2+24m+19}{2(m+2-i)(m+2)} d_i(m+1) \\ - \frac{(m+i+1)(4m+3)(4m+5)}{4(m+2-i)(m+1)(m+2)} d_i(m), \quad 0 \leq i \leq m+1,$$

and for $0 \leq i \leq m+1$,

$$(9) \quad (m+2-i)(m+i-1)d_{i-2}(m) - (i-1)(2m+1)d_{i-1}(m) + i(i-1)d_i(m) = 0.$$

Moll [2007] independently derived the recurrence relations (6) and (9) from which the other two relations can be easily deduced.

To prove Theorem 2.1(4), we need the following lemma.

Lemma 2.2. *Assume that $m \geq 2$. For $0 \leq i \leq m-2$, we have*

$$(10) \quad \frac{d_i(m)}{d_{i+1}(m)} < \frac{(4m+2i+3)d_{i+1}(m)}{(4m+2i+7)d_{i+2}(m)}.$$

Proof. We proceed by induction on m . When $m = 2$, it is easy to check that the result holds. Assume that the lemma is valid for n , namely,

$$(11) \quad \frac{d_i(n)}{d_{i+1}(n)} < \frac{(4n+2i+3)d_{i+1}(n)}{(4n+2i+7)d_{i+2}(n)}, \quad 0 \leq i \leq n-2.$$

We aim to show that (10) holds for $n+1$, that is,

$$(12) \quad \frac{d_i(n+1)}{d_{i+1}(n+1)} < \frac{(4n+2i+7)d_{i+1}(n+1)}{(4n+2i+11)d_{i+2}(n+1)}, \quad 0 \leq i \leq n-1.$$

From the recurrence relation (6), we deduce that, for $0 \leq i \leq n-1$,

$$(2i+4n+7)d_{i+1}^2(n+1) - (2i+4n+11)d_i(n+1)d_{i+2}(n+1) \\ = (2i+4n+7) \left(\frac{i+n+1}{n+1} d_i(n) + \frac{2i+4n+5}{2(n+1)} d_{i+1}(n) \right)^2 \\ - (2i+4n+11) \left(\frac{i+n+2}{n+1} d_{i+1}(n) + \frac{2i+4n+7}{2(n+1)} d_{i+2}(n) \right) \\ \times \left(\frac{n+i}{n+1} d_{i-1}(n) + \frac{2i+4n+3}{2(n+1)} d_i(n) \right)$$

$$= \frac{A_1(n, i) + A_2(n, i) + A_3(n, i)}{4(n+1)^2},$$

where $A_1(n, i)$, $A_2(n, i)$ and $A_3(n, i)$ are given by

$$A_1(n, i) = 4(2i + 4n + 7)(i + n + 1)^2 d_i^2(n) - 4(n + i)(2i + 4n + 11)(i + n + 2)d_{i+1}(n)d_{i-1}(n),$$

$$A_2(n, i) = (2i + 4n + 7)(2i + 4n + 5)^2 d_{i+1}^2(n) - (2i + 4n + 3)(2i + 4n + 11)(2i + 4n + 7)d_i(n)d_{i+2}(n),$$

$$A_3(n, i) = (8i^3 + 40i^2 + 58i + 32n^3 + 42n + 80n^2 + 120ni + 40i^2n + 64n^2i + 8) \cdot d_{i+1}(n)d_i(n) - 2(n + i)(2i + 4n + 11)(2i + 4n + 7)d_{i+2}(n)d_{i-1}(n).$$

We will show that $A_1(n, i)$, $A_2(n, i)$ and $A_3(n, i)$ are all positive for $0 \leq i \leq n - 2$. By the induction hypothesis (11), we find that for $0 \leq i \leq n - 2$,

$$\begin{aligned} A_1(n, i) &> 4(2i + 4n + 7)(i + n + 1)^2 d_i^2(n) - 4(n + i)(2i + 4n + 11)(i + n + 2) \frac{(4n + 2i + 1)}{(4n + 2i + 5)} d_i^2(n) \\ &= 4 \frac{35 + 96n + 72i + 64ni + 40n^2 + 28i^2}{2i + 4n + 5} d_i^2(n), \end{aligned}$$

$$\begin{aligned} A_2(n, i) &> (2i + 4n + 7)(2i + 4n + 5)^2 d_{i+1}^2(n) - (2i + 4n + 3)(2i + 4n + 11)(2i + 4n + 7) \frac{(4n + 2i + 3)}{(4n + 2i + 7)} d_{i+1}^2(n) \\ &= (40i + 80n + 76) d_{i+1}^2(n), \end{aligned}$$

which are both positive. Also by the induction hypothesis (11), we see that

$$(13) \quad d_i(n)d_{i+1}(n) > \frac{(2i + 4n + 5)(2i + 4n + 7)}{(2i + 4n + 3)(2i + 4n + 1)} d_{i-1}(n)d_{i+2}(n),$$

for $0 \leq i \leq n - 2$. This implies that

$$\begin{aligned} A_3(n, i) &> (8i^3 + 40i^2 + 58i + 32n^3 + 42n + 80n^2 + 120ni + 40i^2n + 64n^2i + 8)d_{i+1}(n)d_i(n) \\ &\quad - 2(n + i)(2i + 4n + 11)(2i + 4n + 7) \frac{(4n + 2i + 3)(4n + 2i + 1)}{(4n + 2i + 5)(4n + 2i + 7)} d_{i+1}(n)d_i(n) \\ &= 8 \frac{5 + 22n + 30i + 44ni + 24n^2 + 16i^2}{2i + 4n + 5} d_{i+1}(n)d_i(n), \end{aligned}$$

which is positive for $0 \leq i \leq n-2$. Hence the inequality (12) holds for $0 \leq i \leq n-2$. It remains to show that (12) is true for $i = n-1$, that is,

$$(14) \quad \frac{d_{n-1}(n+1)}{d_n(n+1)} < \frac{(6n+5)d_n(n+1)}{(6n+9)d_{n+1}(n+1)}.$$

From (2) it follows that

$$(15) \quad d_n(n+1) = 2^{-n-2}(2n+3) \binom{2n+2}{n+1},$$

$$(16) \quad d_{n+1}(n+1) = \frac{1}{2^{n+1}} \binom{2n+2}{n+1},$$

$$(17) \quad d_n(n+2) = \frac{(n+1)(4n^2+18n+21)}{2^{n+4}(2n+3)} \binom{2n+4}{n+2}.$$

Consequently,

$$\frac{d_{n-1}(n+1)}{d_n(n+1)} = \frac{n(4n^2+10n+7)}{2(2n+1)(2n+3)} < \frac{(2n+3)(6n+5)}{2(6n+9)} = \frac{(6n+5)d_n(n+1)}{(6n+9)d_{n+1}(n+1)}.$$

This completes the proof. \square

We can now prove Theorem 2.1(4). In fact, we shall prove a stronger inequality.

Lemma 2.3. *Assume that $m \geq 2$. For $0 \leq i \leq m-1$, we have*

$$(18) \quad \frac{d_i(m)}{d_{i+1}(m)} > \frac{(2i+4m+5)d_i(m+1)}{(2i+4m+3)d_{i+1}(m+1)}.$$

Proof. By Lemma 2.2, we have for $0 \leq i \leq m-1$,

$$(19) \quad d_i^2(m) > \frac{2i+4m+5}{2i+4m+1} d_{i-1}(m)d_{i+1}(m).$$

From (19) and the recurrence relation (6), for $0 \leq i \leq m-1$,

$$\begin{aligned} & d_{i+1}(m+1)d_i(m) - \frac{2i+4m+5}{2i+4m+3} d_{i+1}(m)d_i(m+1) \\ &= \frac{2i+4m+5}{2(m+1)} d_{i+1}(m)d_i(m) + \frac{i+m+1}{m+1} d_i(m)^2 \\ &\quad - \frac{2i+4m+5}{2i+4m+3} \left(\frac{2i+4m+3}{2(m+1)} d_i(m)d_{i+1}(m) + \frac{i+m}{m+1} d_{i-1}(m)d_{i+1}(m) \right) \\ &= \frac{i+m+1}{m+1} d_i^2(m) - \frac{(4m+2i+5)(m+i)}{(4m+2i+3)(m+1)} d_{i-1}(m)d_{i+1}(m) \\ &> \left(\frac{m+1+i}{m+1} - \frac{(4m+2i+1)(m+i)}{(4m+2i+3)(m+1)} \right) d_i^2(m) \\ &= \frac{6m+4i+3}{(4m+2i+3)(m+1)} d_i^2(m) > 0, \end{aligned}$$

which yields (18). \square

We now turn to the proof of Theorem 2.1(5).

Lemma 2.4. *Assume that $m \geq 2$. For $0 \leq i \leq m-1$, we have*

$$(20) \quad \frac{d_i(m)}{d_{i+1}(m)} < \frac{d_{i+1}(m+1)}{d_{i+2}(m+1)}.$$

Proof. We proceed by induction on m . It is easy to check the lemma holds for $m = 2$. Assume that the lemma is true for $n \geq 2$, that is,

$$(21) \quad \frac{d_i(n)}{d_{i+1}(n)} < \frac{d_{i+1}(n+1)}{d_{i+2}(n+1)}, \quad 0 \leq i \leq n-1.$$

It will be shown that the theorem holds for $n+1$, that is,

$$(22) \quad \frac{d_i(n+1)}{d_{i+1}(n+1)} < \frac{d_{i+1}(n+2)}{d_{i+2}(n+2)}, \quad 0 \leq i \leq n.$$

Recall that the sequence $\{d_i(n+1)\}_{0 \leq i \leq n+1}$ is unimodal. Furthermore, from (3) or the ratio monotone property [Chen and Xia 2009], the maximum element appears in the middle, namely, $d_i(n+1) < d_{i+1}(n+1)$ when $0 \leq i \leq [(n+1)/2] - 1$ and $d_i(n+1) > d_{i+1}(n+1)$ when $[(n+1)/2] \leq i \leq n$.

Showing (22) for $0 \leq i \leq n-1$ breaks into two cases.

The first case is $d_i(n+1) < d_{i+1}(n+1)$, namely, $0 \leq i \leq [(n+1)/2] - 1$. From the recurrence relation (6), we find that for $0 \leq i \leq [(n+1)/2] - 1$,

$$\begin{aligned} & d_{i+1}(n+1)d_{i+1}(n+2) - d_{i+2}(n+2)d_i(n+1) \\ &= \frac{2i+4n+9}{2(n+2)} d_{i+1}^2(n+1) + \frac{i+n+2}{n+2} d_i(n+1)d_{i+1}(n+1) \\ &\quad - \frac{2i+4n+11}{2(n+2)} d_i(n+1)d_{i+2}(n+1) - \frac{i+n+3}{n+2} d_i(n+1)d_{i+1}(n+1) \\ &= \frac{2i+4n+9}{2(n+2)} d_{i+1}^2(n+1) - \frac{2i+4n+11}{2(n+2)} d_i(n+1)d_{i+2}(n+1) \\ &\quad - \frac{1}{n+2} d_i(n+1)d_{i+1}(n+1) \\ &> \frac{2i+4n+7}{2(n+2)} d_{i+1}^2(n+1) - \frac{2i+4n+11}{2(n+2)} d_i(n+1)d_{i+2}(n+1), \end{aligned}$$

which is positive by Lemma 2.2. It follows that for $0 \leq i \leq [(n+1)/2] - 1$,

$$(23) \quad d_{i+1}(n+1)d_{i+1}(n+2) - d_{i+2}(n+2)d_i(n+1) > 0.$$

This completes the proof of the first case.

The second case is when $[(n+1)/2] \leq i \leq n-1$. From the recurrence relations (6) and (7), it follows that for $[(n+1)/2] \leq i \leq n-1$,

$$\begin{aligned}
& d_{i+1}(n+2)d_{i+1}(n+1) - d_{i+2}(n+2)d_i(n+1) \\
&= \left(\frac{(4n-2i+5)(n+i+3)}{2(n+2)(n+1-i)} d_{i+1}(n+1) - \frac{(i+1)(i+2)}{(n+2)(n+1-i)} d_{i+2}(n+1) \right) \\
&\quad \times \left(\frac{n+1+i}{n+1} d_i(n) + \frac{4n+2i+5}{2(n+1)} d_{i+1}(n) \right) \\
&\quad - \left(\frac{n+3+i}{n+2} d_{i+1}(n+1) + \frac{4n+2i+11}{2(n+2)} d_{i+2}(n+1) \right) \\
&\quad \times \left(\frac{(4n-2i+3)(n+i+1)}{2(n+1)(n+1-i)} d_i(n) - \frac{i(i+1)}{(n+1)(n+1-i)} d_{i+1}(n) \right) \\
&= B_1(n, i)d_{i+1}(n+1)d_i(n) + B_2(n, i)d_{i+1}(n+1)d_{i+1}(n) \\
&\quad + B_3(n, i)d_{i+2}(n+1)d_i(n) + B_4(n, i)d_{i+2}(n+1)d_{i+1}(n),
\end{aligned}$$

where $B_1(n, i)$, $B_2(n, i)$, $B_3(n, i)$ and $B_4(n, i)$ are given by

$$(24) \quad B_1(n, i) = \frac{(n+i+3)(n+1+i)}{(n+2)(n+1-i)(n+1)},$$

$$(25) \quad B_2(n, i) = \frac{(n+i+3)(16n^2+40n+25+4i)}{4(n+2)(n+1-i)(n+1)},$$

$$(26) \quad B_3(n, i) = -\frac{(n+1+i)(41+16n^2+56n-4i)}{4(n+2)(n+1-i)(n+1)},$$

$$(27) \quad B_4(n, i) = -\frac{(i+1)(4n+5-i)}{(n+2)(n+1-i)(n+1)}.$$

Since $[(n+1)/2] \leq i \leq n-1$, it follows from (3) that $d_{i+1}(n+1) > d_{i+2}(n+1)$ and $d_i(n) > d_{i+1}(n)$. Thus we get

$$(28) \quad d_{i+1}(n+1)d_i(n) > d_{i+1}(n+1)d_{i+1}(n),$$

$$(29) \quad d_{i+1}(n+1)d_{i+1}(n) > d_{i+2}(n+1)d_{i+1}(n).$$

Observe that $B_1(n, i)$ and $B_2(n, i)$ are positive, and $B_3(n, i)$ and $B_4(n, i)$ are negative. By the induction hypothesis (21) and inequalities (28) and (29), we find that, for $[(n+1)/2] \leq i \leq n-1$,

$$\begin{aligned}
(30) \quad & d_{i+1}(n+2)d_{i+1}(n+1) - d_{i+2}(n+2)d_i(n+1) \\
&> (B_1(n, i) + B_2(n, i) + B_3(n, i) + B_4(n, i)) d_{i+1}(n+1)d_{i+1}(n) \\
&= \frac{24n+10n^2-8ni+8i^2+13}{2(n+2)(n+1-i)(n+1)} d_{i+1}(n+1)d_{i+1}(n) > 0.
\end{aligned}$$

From the inequalities (23) and (30), it follows that (22) holds for $0 \leq i \leq n-1$.

It is still necessary to show that (22) is true for $i = n$, that is,

$$(31) \quad \frac{d_n(n+1)}{d_{n+1}(n+1)} < \frac{d_{n+1}(n+2)}{d_{n+2}(n+2)}.$$

For the recurrence relation (9), setting $i = n + 2$, we find that

$$\frac{d_n(n+1)}{d_{n+1}(n+1)} = \frac{2n+3}{2} < \frac{2n+5}{2} = \frac{d_{n+1}(n+2)}{d_{n+2}(n+2)},$$

as desired. Hence the proof is complete by induction. \square

Lemmas 2.3 and 2.4 immediately imply the interlacing log-concavity of the Boros–Moll polynomials.

3. Polynomials with triangular relations on coefficients

Many combinatorial polynomials admit triangular relations on the coefficients. The log-concavity of polynomials of this kind has been extensively studied. We show that many classical polynomials of this kind are also interlacingly log-concave. For example, it is easy to check that the binomial coefficients, the Narayana numbers

$$N(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k+1},$$

and the Bessel numbers

$$B(n, k) = \frac{(2n-k-1)!}{2^k(n-k)!(k-1)!}$$

are interlacingly log-concave. Moreover, we give a criterion that applies to many combinatorial sequences such as the signless Stirling numbers of the first kind, the Stirling numbers of the second kind and the Whitney numbers.

Theorem 3.1. *Suppose that for any $n \geq 0$,*

$$G_n(x) = \sum_{k=0}^n T(n, k)x^k$$

is a polynomial of degree n which has only real zeros, and suppose that the coefficients $T(n, k)$ satisfy a recurrence relation of the form

$$T(n, k) = f(n, k)T(n-1, k) + g(n, k)T(n-1, k-1).$$

If

$$(32) \quad \frac{(n-k)k}{(n-k+1)(k+1)} f(n+1, k+1) \leq f(n+1, k) \leq f(n+1, k+1),$$

$$(33) \quad g(n+1, k+1) \leq g(n+1, k) \leq \frac{(n-k+1)(k+1)}{(n-k)k} g(n+1, k+1),$$

then the polynomials $G_n(x)$ are interlacingly log-concave.

Proof. Since the polynomial $G_n(x)$ has only real zeros, by Newton's inequality,

$$k(n-k)T(n, k)^2 \geq (k+1)(n-k+1)T(n, k-1)T(n, k+1).$$

Hence

$$\begin{aligned} & T(n, k)T(n+1, k+1) - T(n+1, k)T(n, k+1) \\ &= f(n+1, k+1)T(n, k)T(n, k+1) + g(n+1, k+1)T(n, k)^2 \\ &\quad - f(n+1, k)T(n, k)T(n, k+1) - g(n+1, k)T(n, k-1)T(n, k+1) \\ &\geq (f(n+1, k+1) - f(n+1, k))T(n, k)T(n, k+1) \\ &\quad + \left(\frac{(n-k+1)(k+1)}{(n-k)k} g(n+1, k+1) - g(n+1, k) \right) T(n, k-1)T(n, k+1), \end{aligned}$$

which is positive by (32) and (33). It follows that

$$(34) \quad \frac{T(n, k)}{T(n, k+1)} \geq \frac{T(n+1, k)}{T(n+1, k+1)}.$$

On the other hand, we have

$$\begin{aligned} & T(n, k+1)T(n+1, k+1) - T(n, k)T(n+1, k+2) \\ &= f(n+1, k+1)T(n, k+1)^2 + g(n+1, k+1)T(n, k)T(n, k+1) \\ &\quad - f(n+1, k+2)T(n, k)T(n, k+2) - g(n+1, k+2)T(n, k+1)T(n, k) \\ &\geq \left(f(n+1, k+1) - \frac{(n-k-1)(k+1)}{(n-k)(k+2)} f(n+1, k+2) \right) T(n, k+1)^2 \\ &\quad + (g(n+1, k+1) - g(n+1, k+2))T(n, k+1)T(n, k). \end{aligned}$$

It follows from (32) that

$$(35) \quad \frac{T(n, k)}{T(n, k+1)} \leq \frac{T(n+1, k+1)}{T(n+1, k+2)}.$$

This completes the proof. \square

Employing Theorem 3.1, we can show that many combinatorial polynomials which have only real zeros are interlacingly log-concave, for example,

(1) the polynomials

$$x(x+1)(x+2) \cdots (x+n-1),$$

whose coefficients are the signless Stirling numbers of the first kind, which satisfy the recurrence relation

$$c(n, k) = (n-1)c(n-1, k) + c(n-1, k-1);$$

- (2) the Bell polynomials whose coefficients are the Stirling numbers of the second kind $S(n, k)$, which satisfy the recurrence relation

$$S(n, k) = S(n-1, k-1) + kS(n-1, k);$$

- (3) the Whitney polynomials

$$W_n(x) = \sum_{k=0}^n W_m(n, k)x^k,$$

which have only real zeros; see [Benoumhani 1997; 1999]. The coefficients $W_m(n, k)$ satisfy the recurrence relation

$$W_m(n, k) = (1 + mk)W_m(n-1, k) + W_m(n-1, k-1).$$

Acknowledgments

We wish to thank the referee for valuable comments.

References

- [Alvarez et al. 2001] J. Alvarez, M. Amadis, G. Boros, D. Karp, V. H. Moll, and L. Rosales, “An extension of a criterion for unimodality”, *Electron. J. Combin.* **8**:1 (2001), Research Paper 30. MR 2002g:05017 Zbl 0984.05008
- [Amdeberhan and Moll 2009] T. Amdeberhan and V. H. Moll, “A formula for a quartic integral: a survey of old proofs and some new ones”, *Ramanujan J.* **18**:1 (2009), 91–102. MR 2009m:33008 Zbl 1178.33002
- [Benoumhani 1997] M. Benoumhani, “On some numbers related to Whitney numbers of Dowling lattices”, *Adv. in Appl. Math.* **19**:1 (1997), 106–116. MR 98f:05004 Zbl 0876.05001
- [Benoumhani 1999] M. Benoumhani, “Log-concavity of Whitney numbers of Dowling lattices”, *Adv. in Appl. Math.* **22**:2 (1999), 186–189. MR 2000i:05008 Zbl 0918.05003
- [Boros and Moll 1999a] G. Boros and V. H. Moll, “A criterion for unimodality”, *Electron. J. Combin.* **6** (1999), Research Paper 10. MR 99k:05017 Zbl 0911.05004
- [Boros and Moll 1999b] G. Boros and V. H. Moll, “An integral hidden in Gradshteyn and Ryzhik”, *J. Comput. Appl. Math.* **106**:2 (1999), 361–368. MR 2000c:33024 Zbl 0939.33007
- [Boros and Moll 1999c] G. Boros and V. H. Moll, “A sequence of unimodal polynomials”, *J. Math. Anal. Appl.* **237**:1 (1999), 272–287. MR 2000m:33007 Zbl 0944.33009
- [Boros and Moll 2001] G. Boros and V. H. Moll, “The double square root, Jacobi polynomials and Ramanujan’s master theorem”, *J. Comput. Appl. Math.* **130**:1-2 (2001), 337–344. MR 2002d:33030 Zbl 1011.33005
- [Boros and Moll 2004] G. Boros and V. H. Moll, *Irresistible integrals: Symbolics, analysis and experiments in the evaluation of integrals*, Cambridge University Press, 2004. MR 2005b:00001 Zbl 1090.11075
- [Chen and Xia 2009] W. Y. C. Chen and E. X. W. Xia, “The ratio monotonicity of the Boros–Moll polynomials”, *Math. Comp.* **78**:268 (2009), 2269–2282. MR 2010f:33046 Zbl 1221.33036

- [Chen et al. \geq 2011] W. Y. C. Chen, S. X. M. Pang, and E. X. Y. Qu, “A combinatorial proof of the log-concavity of the Boros–Moll polynomials”, to appear in *Ramanujan J.*
- [Kauers and Paule 2007] M. Kauers and P. Paule, “A computer proof of Moll’s log-concavity conjecture”, *Proc. Amer. Math. Soc.* **135**:12 (2007), 3847–3856. MR 2009d:33063 Zbl 1126.33009
- [Moll 2002] V. H. Moll, “The evaluation of integrals: a personal story”, *Notices Amer. Math. Soc.* **49**:3 (2002), 311–317. MR 2002m:11105 Zbl 1126.11347
- [Moll 2007] V. H. Moll, “Combinatorial sequences arising from a rational integral”, *Online J. Anal. Comb.* **2** (2007), Art. 4. MR 2008m:05006 Zbl 1123.05003

Received August 9, 2010. Revised May 12, 2011.

WILLIAM Y. C. CHEN
CENTER FOR COMBINATORICS, LPMC-TJKLC
NANKAI UNIVERSITY
TIANJIN 300071
CHINA
chen@nankai.edu.cn

LARRY X. W. WANG
CENTER FOR COMBINATORICS, LPMC-TJKLC
NANKAI UNIVERSITY
TIANJIN 300071
CHINA
wxw@cfc.nankai.edu.cn

ERNEST X. W. XIA
CENTER FOR COMBINATORICS, LPMC-TJKLC
NANKAI UNIVERSITY
TIANJIN 300071
CHINA
xxwrml@mail.nankai.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 254 No. 1 November 2011

A mean curvature estimate for cylindrically bounded submanifolds	1
LUIS J. ALÍAS and MARCOS DAJCZER	
Weyl group multiple Dirichlet series of type C	11
JENNIFER BEINEKE, BENJAMIN BRUBAKER and SHARON FRECHETTE	
Milnor open books of links of some rational surface singularities	47
MOHAN BHUPAL and BURAK OZBAGCI	
Simple closed curves, word length, and nilpotent quotients of free groups	67
KHALID BOU-RABEE and ASAF HADARI	
Strong submodules of almost projective modules	73
GÁBOR BRAUN and JAN TRLIJAJ	
Interlacing log-concavity of the Boros–Moll polynomials	89
WILLIAM Y. C. CHEN, LARRY X. W. WANG and ERNEST X. W. XIA	
Schwarzian norms and two-point distortion	101
MARTIN CHUAQUI, PETER DUREN, WILLIAM MA, DIEGO MEJÍA, DAVID MINDA and BRAD OSGOOD	
The principle of stationary phase for the Fourier transform of D -modules	117
JIANGXUE FANG	
Monotonicity and uniqueness of a 3D transonic shock solution in a conic nozzle with variable end pressure	129
JUN LI, ZHOUPING XIN and HUICHENG YIN	
Refined open noncommutative Donaldson–Thomas invariants for small crepant resolutions	173
KENTARO NAGAO	
The Dirichlet problem for harmonic functions on compact sets	211
TONY L. PERKINS	
Extension of an analytic disc and domains in \mathbb{C}^2 with noncompact automorphism group	227
MINJU SONG	
Regularity of the first eigenvalue of the p -Laplacian and Yamabe invariant along geometric flows	239
ER-MIN WANG and YU ZHENG	



0030-8730(201111)254:1;1-C