

*Pacific  
Journal of  
Mathematics*

**AN ANALOGUE OF KREIN'S THEOREM  
FOR SEMISIMPLE LIE GROUPS**

SANJOY PUSTI

Volume 254 No. 2

December 2011

## AN ANALOGUE OF KREIN'S THEOREM FOR SEMISIMPLE LIE GROUPS

SANJOY PUSTI

**We give an integral representation of  $K$ -positive definite functions on a real rank  $n$  connected, noncompact, semisimple Lie group with finite centre. Moreover, we characterize the  $\lambda$ 's for which the  $\tau$ -spherical function  $\phi_{\sigma,\lambda}^\tau$  is positive definite for the group  $G = \text{Spin}_e(n, 1)$  and the complex spin representation  $\tau$ .**

### 1. Introduction

A continuous function  $f$  on  $\mathbb{R}$  is said to be *positive definite* if for any real numbers  $x_1, \dots, x_m$  and complex numbers  $\xi_1, \dots, \xi_m$  the following holds:

$$\sum_{k,j=1}^m f(x_j - x_k) \xi_k \overline{\xi_j} \geq 0.$$

This definition is equivalent to

$$\int_{\mathbb{R}} f(x) (\phi * \phi^*)(-x) dx \geq 0 \quad \text{for all } \phi \in C_c^\infty(\mathbb{R}),$$

where  $\phi^*(x) = \overline{\phi(-x)}$ . Also, an even continuous function  $f$  on  $\mathbb{R}$  is said to be *evenly positive definite* if

$$\int_{\mathbb{R}} f(x) (\phi * \phi^*)(-x) dx = \int_{\mathbb{R}} f(x) (\phi * \phi^*)(x) dx \geq 0 \quad \text{for all } \phi \in C_c^\infty(\mathbb{R})_e,$$

where  $C_c^\infty(\mathbb{R})_e$  denotes the set of infinitely differentiable compactly supported even functions on  $\mathbb{R}$ . Then it is clear that the set of even positive definite functions is a subset of the set of evenly positive definite functions. Bochner's theorem and M. G. Krein's theorem respectively give integral representations of positive definite functions and evenly positive definite functions. Precisely, for a positive definite function  $f$  on  $\mathbb{R}$ , there exists a finite positive measure  $\mu$  on  $\mathbb{R}$  such that

$$f(x) = \int_{\mathbb{R}} e^{i\lambda x} d\mu(\lambda).$$

---

*MSC2010:* primary 43A85; secondary 22E30.

*Keywords:* positive definite functions,  $K$ -positive definite functions,  $\tau$ -positive definite functions.

Also for an evenly positive definite function  $f$  on  $\mathbb{R}$ , there exists a finite positive even measure  $\sigma$  on  $\mathbb{R} \cup i\mathbb{R}$  such that

$$f(x) = \int_{\mathbb{R} \cup i\mathbb{R}} e^{i\lambda x} d\sigma(\lambda).$$

From this integral representation it follows that a bounded evenly positive definite function is a positive definite function. We note that the measure  $\sigma$  in the integral representation of an evenly positive definite function is not unique, whereas the measure  $\mu$  in the integral representation of a positive definite function is unique. However, if an evenly positive definite function satisfies a certain restriction on its growth for  $|x| \rightarrow \infty$ , then the integral representation becomes unique [Gelfand and Vilenkin 1964].

Let  $G$  be a connected, noncompact semisimple Lie group with finite centre, and let  $K$  be a fixed maximal compact subgroup of  $G$ . Integral representations of  $K$ -positive definite distributions and  $K$ -positive definite functions have been derived for real rank-one semisimple Lie groups with finite centre in [Sitaram 1978] and [Pusti 2011], respectively.

An analogue of Krein's theorem on  $\mathbb{R}^n$  has been obtained by N. Bopp [1979]. In this case, instead of evenly positive definite functions, one considers functions which are positive definite relative to the action of a finite subgroup of  $O(n)$ . Here too, if we impose a certain growth condition, then the integral representation of these functions is unique. In this paper, using Bopp's result, we derive an integral representation for the  $K$ -positive definite functions on a real rank  $n$  connected, noncompact, semisimple Lie group with finite centre. We observe that the set of positive definite functions is a proper subset of the set of  $K$ -positive definite functions. Next, we consider the  $\tau$ -positive definite functions,  $\tau \in \widehat{K}$ . The  $K$ -positive definite functions are a special instance of the  $\tau$ -positive definite functions (for  $\tau$  equals the trivial representation). We give an example in which the set of  $\tau$ -positive definite functions is same as the set of positive definite functions. That is, the same conclusion (as in  $K$ -positive definite function) is not true for  $\tau$ -positive definite functions. Finally we characterize the  $\lambda$ 's for which the  $\tau$ -spherical function  $\phi_{\sigma,\lambda}^\tau$  is a positive definite function for the group  $G = \text{Spin}_e(n, 1)$  and the complex spin representation  $\tau$ . We note that G. van Dijk and A. Pasquale [1999] studied positive definiteness of  $\phi_{\sigma,\lambda}^\tau$  for the group  $G = \text{Sp}(1, n)$ .

## 2. Preliminaries

Most of our notations are standard and can be found in [Anker 1991]. Let  $G$  be a real rank  $n$  connected, noncompact, semisimple Lie group with finite centre with Lie algebra  $\mathfrak{g}$ , and let  $\mathcal{U}(\mathfrak{g})$  be the universal enveloping algebra of  $\mathfrak{g}$ . Let  $\mathfrak{g} = \mathfrak{k} + \mathfrak{s}$  be a Cartan decomposition of  $\mathfrak{g}$ , and let  $K$  be the maximal compact subgroup of  $G$

with Lie algebra  $\mathfrak{k}$ . We fix a maximal abelian subspace  $\mathfrak{a}$  of  $\mathfrak{s}$ . Since  $G$  is of real rank  $n$ , we have  $\dim \mathfrak{a} = n$ . We denote the real dual of  $\mathfrak{a}$  by  $\mathfrak{a}^*$  and its complex dual by  $\mathfrak{a}_{\mathbb{C}}^*$ . The Killing form of  $\mathfrak{g}$  induces an  $\text{Ad } K$ -invariant scalar product on  $\mathfrak{s}$  and hence a  $G$ -invariant Riemannian metric on  $G/K$  (or  $K \backslash G$ ). With this structure,  $G/K$  is a Riemannian globally symmetric space of the noncompact type. Also, the Killing form of  $\mathfrak{g}$  induces a scalar product on  $\mathfrak{a}$  and hence on  $\mathfrak{a}^*$ . We denote by  $\langle \cdot, \cdot \rangle$  its  $\mathbb{C}$ -bilinear extension to  $\mathfrak{a}_{\mathbb{C}}^*$ .

Let  $\mathfrak{g} = \mathfrak{g}_0 \oplus (\oplus_{\alpha \in \Sigma} \mathfrak{g}_{\alpha})$  be the root space decomposition of  $\mathfrak{g}$ . Here,  $\mathfrak{g}_0 = \mathfrak{m} \oplus \mathfrak{a}$ , where  $\mathfrak{m}$  is the centralizer of  $\mathfrak{a}$  in  $\mathfrak{k}$  and  $\Sigma \subseteq \mathfrak{a}^*$  is the root system of  $(\mathfrak{g}, \mathfrak{a})$ . Let  $W$  be the Weyl group associated to  $\Sigma$ . We choose a set  $\Sigma^+$  of positive roots. Let  $\mathfrak{a}^+ \subseteq \mathfrak{a}$  be the corresponding positive Weyl chamber and let  $\overline{\mathfrak{a}^+}$  be its closure. We denote by  $(\mathfrak{a}^*)^+$  and  $(\overline{\mathfrak{a}^+})^*$  the similar cones in  $\mathfrak{a}^*$ . Let  $\mathfrak{n} = \oplus_{\alpha \in \Sigma^+} \mathfrak{g}_{\alpha}$ . Then  $\mathfrak{n}$  is a nilpotent subalgebra of  $\mathfrak{g}$ . The element  $\rho \in \mathfrak{a}^*$  is defined by

$$\rho(H) = \frac{1}{2} \sum_{\alpha \in \Sigma^+} m_{\alpha} \alpha(H),$$

where  $m_{\alpha} = \dim \mathfrak{g}_{\alpha}$ . Let  $A$  be the analytic subgroup of  $G$  with Lie algebra  $\mathfrak{a}$ . Then  $A$  is a closed subgroup of  $G$  and the exponential map is an isomorphism from  $\mathfrak{a}$  onto  $A$ . We set  $A^+ = \exp \mathfrak{a}^+$ . Its closure is  $\overline{A^+} = \exp \overline{\mathfrak{a}^+}$ . Let  $N$  be the analytic subgroup of  $G$  with Lie algebra  $\mathfrak{n}$ , and let  $M$  be the centralizer of  $A$  in  $K$ .

The group  $G$  can be decomposed as  $G = K \overline{A^+} K$ . It is called the Cartan decomposition of  $G$  and every element  $x$  of  $G$  can be decomposed as  $x = k_1 a k_2$  with  $k_1, k_2 \in K$  and  $a \in \overline{A^+}$ . We let  $x^+$  be the  $\overline{\mathfrak{a}^+}$ -component of  $x \in G$  in the decomposition  $G = K(\exp \overline{\mathfrak{a}^+})K$  and let  $|x| = \|x^+\|$ . Viewed on  $G/K$ ,  $|\cdot|$  is the distance to the origin  $0 = \{K\}$ . Also, the group  $G$  has Iwasawa decomposition  $G = KAN$ . Let  $k(x)$  and  $H(x)$  be the components of  $x \in G$  in  $K$  and  $\mathfrak{a}$ . Then any element  $x \in G$  can be expressed as  $x = k(x) \exp H(x)n$  for some  $n \in N$ .

For  $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ , the elementary spherical function  $\phi_{\lambda}$  on  $G$  is given by

$$\phi_{\lambda}(x) = \int_K e^{-(i\lambda + \rho)H(x^{-1}k)} dk.$$

It satisfies the following properties:

- (1) It is  $K$ -biinvariant, that is,  $\phi_{\lambda}(k_1 x k_2) = \phi_{\lambda}(x)$  for all  $k_1, k_2 \in K$  and  $x \in G$ . Also, it is  $W$ -invariant in  $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ , that is,  $\phi_{w \cdot \lambda}(x) = \phi_{\lambda}(x)$  for all  $w \in W$  and  $x \in G$ .
- (2) The function  $\phi_{\lambda}(x)$  is  $C^{\infty}$  in  $x$  and holomorphic in  $\lambda$ .
- (3) It is a joint eigenfunction for all  $G$ -invariant differential operators on  $G/K$ ; in particular for the Laplacian  $\Delta$  on  $G/K$ ,

$$\Delta \phi_{\lambda} = -(\langle \lambda, \lambda \rangle + \|\rho\|^2) \phi_{\lambda}.$$

A function  $f$  on  $G$  is called  $K$ -biinvariant if  $f(k_1 x k_2) = f(x)$  for all  $k_1, k_2 \in K$  and  $x \in G$ . For a  $K$ -biinvariant function  $f$  on  $G$ , its spherical Fourier transform is defined by

$$\hat{f}(\lambda) = \int_G f(x) \phi_\lambda(x^{-1}) dx$$

for suitable  $\lambda \in \mathfrak{a}_\mathbb{C}^*$ .

The set of infinitely differentiable compactly supported  $K$ -biinvariant functions and infinitely differentiable  $K$ -biinvariant functions are denoted by  $C_c^\infty(G//K)$  and  $C^\infty(G//K)$ , respectively. For  $0 < p \leq 2$  the  $L^p$ -Schwartz space  $\mathcal{C}^p(G//K)$  is the set of all functions  $f \in C^\infty(G//K)$  such that

$$\sup_{x \in G} (1 + |x|)^s \phi_0(x)^{-2/p} |f(D; x; E)| < \infty$$

for any  $D, E \in \mathcal{O}(\mathfrak{g})$  and any integer  $s \geq 0$ . The Schwartz space  $\mathcal{C}^p(G//K)$  is topologized by the seminorms

$$\sigma_{D,E,s}^p(f) = \sup_{x \in G} (1 + |x|)^s \phi_0(x)^{-2/p} |f(D; x; E)|.$$

Then it follows that  $C_c^\infty(G//K)$  is dense in  $\mathcal{C}^p(G//K)$  and  $\mathcal{C}^p(G//K)$  is dense in  $L^p(G//K)$ .

Let  $\mathcal{P}(\mathfrak{a}_\mathbb{C}^*)$  be the space of entire functions on  $\mathfrak{a}_\mathbb{C}^*$ , which are of exponential type and rapidly decreasing. The set of  $W$ -invariant elements in  $\mathcal{P}(\mathfrak{a}_\mathbb{C}^*)$  is denoted by  $\mathcal{P}(\mathfrak{a}_\mathbb{C}^*)^W$ .

For fixed  $\epsilon > 0$ , let  $C^{\epsilon\rho}$  be the convex hull of the set  $W \cdot \epsilon\rho$  in  $\mathfrak{a}^*$ , and let  $\mathfrak{a}_\epsilon^* = \mathfrak{a}^* + iC^{\epsilon\rho}$  be the tube in  $\mathfrak{a}_\mathbb{C}^*$  with base  $C^{\epsilon\rho}$ . For  $\epsilon = 0$ ,  $\mathfrak{a}_\epsilon^*$  reduces to  $\mathfrak{a}^*$ . Let  $\mathbf{S}(\mathfrak{a}^*)$  be the symmetric algebra over  $\mathfrak{a}^*$ . We define the Schwartz space  $\mathcal{S}(\mathfrak{a}_\epsilon^*)$  as the space of all complex valued functions  $h$  such that the following hold true.

- (1)  $h$  is holomorphic in the interior of  $\mathfrak{a}_\epsilon^*$ .
- (2)  $h$  and all its derivatives extend continuously to  $\mathfrak{a}_\epsilon^*$ .
- (3) for any polynomial  $P \in \mathbf{S}(\mathfrak{a}^*)$  and any (integer)  $t \geq 0$ ,

$$\sup_{\lambda \in \mathfrak{a}_\epsilon^*} (1 + \|\lambda\|)^t \left| P\left(\frac{\partial}{\partial \lambda}\right) h(\lambda) \right| < \infty.$$

The space  $\mathcal{S}(\mathfrak{a}_\epsilon^*)$  is topologized by the seminorms

$$\tau_{P,t}^\epsilon(h) = \sup_{\lambda \in \mathfrak{a}_\epsilon^*} (1 + \|\lambda\|)^t \left| P\left(\frac{\partial}{\partial \lambda}\right) h(\lambda) \right|.$$

We denote by  $\mathcal{S}(\mathfrak{a}_\epsilon^*)^W$  the subspace of  $W$ -invariant functions in  $\mathcal{S}(\mathfrak{a}_\epsilon^*)$ . For  $\epsilon = 0$ ,  $\mathcal{S}(\mathfrak{a}_\epsilon^*)$  becomes the classical Schwartz space on  $\mathfrak{a}^*$ . Then for  $\epsilon \geq 0$ ,  $\mathcal{S}(\mathfrak{a}_\epsilon^*)^W$  is a Fréchet algebra (under pointwise multiplication) and  $\mathcal{P}(\mathfrak{a}_\mathbb{C}^*)^W$  is a dense subalgebra of  $\mathcal{S}(\mathfrak{a}_\epsilon^*)^W$ .

We consider the function

$$\cosh_{\epsilon\rho}(H) = \frac{1}{|W|} \sum_{w \in W} e^{w \cdot \epsilon\rho(H)} \quad \text{on } \mathfrak{a}.$$

Then we define the space  $\mathcal{S}_{\epsilon\rho}(\mathfrak{a})$  consisting of all functions  $g \in C^\infty(\mathfrak{a})$  such that

$$\sup_{H \in \mathfrak{a}} (1 + \|H\|)^s \cosh_{\epsilon\rho}(H) \left| P\left(\frac{\partial}{\partial H}\right)g(H) \right| < \infty$$

for any polynomial  $P \in \mathbf{S}(\mathfrak{a})$  (the symmetric algebra over  $\mathfrak{a}$ ) and any  $s \geq 0$ .

**Theorem 2.1** [Anker 1991]. (1) *The spherical Fourier transform  $f \mapsto \hat{f}$  is a topological isomorphism between  $C_c^\infty(G//K)$  and  $\mathcal{P}(\mathfrak{a}_\mathbb{C}^*)^W$  and also between  $\mathcal{C}^p(G//K)$  and  $\mathcal{S}(\mathfrak{a}_\epsilon^*)^W$ , where  $\epsilon = 2/p - 1$ .*

(2) *The Euclidean Fourier transform  $f \mapsto \tilde{f}$  is a topological isomorphism between  $\mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W$  and  $\mathcal{S}(\mathfrak{a}_\epsilon^*)^W$ , where  $\tilde{f}(\lambda) = \int_{\mathfrak{a}} f(H)e^{-i\lambda(H)}dH$ ,  $\lambda \in \mathfrak{a}^*$ .*

For a suitable  $K$ -biinvariant function  $f$  on  $G$ , the Abel transform is defined by

$$\mathcal{A}f(H) = e^{\rho(H)} \int_N f(\exp Hn) dn.$$

It satisfies the relation  $\hat{f}(\lambda) = \widetilde{\mathcal{A}f}(\lambda)$  for a suitable  $K$ -biinvariant function  $f$  on  $G$ . Therefore it follows from Theorem 2.1 that the Abel transform  $f \mapsto \mathcal{A}f$  is a topological isomorphism between  $\mathcal{C}^p(G//K)$  and  $\mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W$  for  $\epsilon = 2/p - 1$ .

### 3. M. G. Krein's theorem

For  $\alpha \geq 0$ , we define

$$\mathcal{S}_\alpha(\mathbb{R}^n) = \{\phi \in C^\infty(\mathbb{R}^n) : \|\phi\|_p < \infty \text{ for any nonnegative integer } p\},$$

where

$$\|\phi\|_p = \max_{|q| \leq p} \sup_{x \in \mathbb{R}^n} (1 + |x|^2)^p e^{\alpha|x|} |D^q \phi(x)|.$$

Then  $S_\alpha(\mathbb{R}^n)$  becomes a Fréchet space and  $C_c^\infty(\mathbb{R}^n)$  is a dense subspace of  $S_\alpha(\mathbb{R}^n)$ . For a finite subgroup  $E$  of  $O(n)$ , let  $\mathcal{S}_\alpha(\mathbb{R}^n)^E$  be the subspace of  $E$ -invariant functions in  $\mathcal{S}_\alpha(\mathbb{R}^n)$ .

**Theorem 3.1** [Bopp 1979]. *Let  $E$  be a finite subgroup of  $O(n)$  and let*

$$T : \mathcal{S}_\alpha(\mathbb{R}^n) \rightarrow \mathbb{C}$$

*be a continuous, linear functional such that*

- (1)  $T(\eta \cdot \phi) = T(\phi)$  for all  $\eta \in E$ ,  $\phi \in \mathcal{S}_\alpha(\mathbb{R}^n)$ .
- (2)  $T(\phi * \phi^*) \geq 0$  for all  $\phi \in \mathcal{S}_\alpha(\mathbb{R}^n)^E$ .

Then there exists a unique positive tempered measure  $\sigma$ , invariant under the  $E$ -action, such that for all  $\phi \in \mathcal{S}_\alpha(\mathbb{R}^n)$ ,

$$T(\phi) = \int_{M \cap T_\alpha} \tilde{\phi}(\xi) d\sigma(\xi),$$

where  $M = \{\xi \in \mathbb{C}^n : \text{there exists } \eta \in E \text{ such that } \eta \cdot \xi = \bar{\xi}\}$  and

$$T_\alpha = \{\xi \in \mathbb{C}^n : |\text{Im } \xi| \leq \alpha\}.$$

Since we have an isomorphism between  $\mathcal{S}_{\epsilon\rho}(\mathbb{R}^n)$  and  $\mathcal{S}_{\epsilon\rho}(\mathfrak{a})$  we can rewrite the theorem above in the following way:

**Theorem 3.2.** Let  $T : \mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W \rightarrow \mathbb{C}$  be a continuous, linear functional such that

$$T(\phi * \phi^*) \geq 0 \quad \text{for all } \phi \in \mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W.$$

Then there exists a unique positive tempered measure  $\sigma$ , invariant under the  $W$ -action, such that for all  $\phi \in \mathcal{S}_{\epsilon\rho}(\mathfrak{a})$ ,

$$T(\phi) = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \tilde{\phi}(\lambda) d\sigma(\lambda),$$

where  $\mathcal{M} = \{\lambda \in \mathfrak{a}_\epsilon^* : \text{there exists } w \in W \text{ such that } w \cdot \lambda = \bar{\lambda}\}$ .

We call a  $K$ -biinvariant continuous function  $f$  on  $G$   $K$ -positive definite if for all  $g \in C_c^\infty(G//K)$ ,

$$\int_G f(x)(g * g^*)(x^{-1}) dx \geq 0,$$

where  $g^*(x) = \overline{g(x^{-1})}$  for all  $x \in G$ . If the equation above is true for every  $g \in C_c^\infty(G)$  we say that  $f$  is a positive definite function. We prove the following analogue of M. G. Krein's theorem for  $K$ -positive definite functions on semisimple Lie groups.

**Theorem 3.3.** For a  $K$ -positive definite function  $f \in \mathcal{C}^p(G//K)'$  ( $0 < p \leq 2$ ), there exists a unique finite positive measure  $\sigma$  on  $\mathcal{M} \cap \mathfrak{a}_\epsilon^*$ , invariant under the Weyl group action, such that for all  $x \in G$

$$f(x) = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \phi_\lambda(x) d\sigma(\lambda),$$

where  $\epsilon = 2/p - 1$ .

*Proof.* We define a linear functional  $T_f : \mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W \rightarrow \mathbb{C}$  by

$$T_f(h) = \int_G f(x)(\mathcal{A}^{-1}h)(x^{-1}) dx.$$

The integral exists and is continuous by the given condition on  $f$  and the isomorphism of the Abel transform on  $\mathcal{C}^p(G//K)$ . Since  $\widehat{f}(\lambda) = \widetilde{\mathcal{A}f}(\lambda)$  for all  $f \in \mathcal{C}^p(G//K)$ , it follows that

$$\widehat{\mathcal{A}^{-1}h} = \widetilde{h} \quad \text{for all } h \in \mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W.$$

Using this, we easily check that

$$\mathcal{A}^{-1}(h_1 * h_2) = \mathcal{A}^{-1}h_1 * \mathcal{A}^{-1}h_2 \quad \text{and} \quad \mathcal{A}^{-1}h_1^* = (\mathcal{A}^{-1}h_1)^*$$

for all  $h_1, h_2 \in \mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W$ . Then

$$T_f(h * h^*) = \int_G f(x) (\mathcal{A}^{-1}h * (\mathcal{A}^{-1}h)^*)(x^{-1}) dx \geq 0,$$

since  $f$  is  $K$ -positive definite. Therefore, by Theorem 3.2, there exists a unique positive tempered measure  $\sigma$  on  $\mathcal{M} \cap \mathfrak{a}_\epsilon^*$ , invariant under the Weyl group action, such that

$$T_f(h) = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \widetilde{h}(\lambda) d\sigma(\lambda) \quad \text{for all } h \in \mathcal{S}_{\epsilon\rho}(\mathfrak{a})^W.$$

This shows that

$$(3-1) \quad \int_G f(x)g(x^{-1}) dx = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \widehat{g}(\lambda) d\sigma(\lambda) \quad \text{for all } g \in \mathcal{C}^p(G//K).$$

Now we show that the measure  $\sigma$  is finite. For this let  $\{g_n\}$  be a Dirac-delta sequence in  $\mathcal{C}^p(G//K)$ . Then  $\{g_n * g_n^*\}$  is also a Dirac-delta sequence in  $\mathcal{C}^p(G//K)$ . Applying this sequence to the previous equation we get

$$\int_G f(x)(g_n * g_n^*)(x^{-1}) dx = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} |\widehat{g}_n(\lambda)|^2 d\sigma(\lambda).$$

Now we take the limit as  $n \rightarrow \infty$  on both sides of the equation and apply Fatou's lemma to get  $\sigma(\mathcal{M} \cap \mathfrak{a}_\epsilon^*) \leq f(e)$ . Therefore the measure  $\sigma$  is finite. From (3-1) we get, using Fubini's theorem,

$$\int_G f(x)g(x^{-1}) dx = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \widehat{g}(\lambda) d\sigma(\lambda) = \int_G g(x^{-1}) \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \phi_\lambda(x) d\sigma(\lambda) dx.$$

This is true for every  $g \in \mathcal{C}^p(G//K)$ . Hence

$$f(x) = \int_{\mathcal{M} \cap \mathfrak{a}_\epsilon^*} \phi_\lambda(x) d\sigma(\lambda) \quad \text{with } \epsilon = \frac{2}{p} - 1. \quad \square$$

We can easily check that a function  $f$  on  $G$  which has an integral representation as in Theorem 3.3 is a  $K$ -positive definite function. That is, the converse of the Theorem 3.3 holds.

A  $K$ -biinvariant distribution  $T$  on  $G$  is called a  $K$ -positive definite distribution if  $T(\phi * \phi^*) \geq 0$  for all  $\phi \in C_c^\infty(G//K)$ . It is a positive definite distribution if the inequality above holds for all  $\phi \in C_c^\infty(G)$ . Barker [1975, p. 201] raised the question whether a  $K$ -positive definite distribution is a positive definite distribution. We shall see that the answer is negative, that is, the set of positive definite distributions is a proper subset of the set of  $K$ -positive definite distributions. For this let us consider  $\lambda_0 \in \mathcal{M} \setminus \mathfrak{a}_1^*$ . Our claim is that  $\phi_{\lambda_0}$  is a  $K$ -positive definite distribution but not a positive definite distribution. By the Helgason–Johnson theorem  $\phi_\lambda$  is bounded if and only if  $\lambda \in \mathfrak{a}_1^*$ . Since  $\lambda_0 \notin \mathfrak{a}_1^*$ ,  $\phi_{\lambda_0}$  is not bounded. Therefore,  $\phi_{\lambda_0}$  is not a positive definite function. Hence  $\phi_{\lambda_0}$  is not a positive definite distribution. Now  $\lambda_0 \in \mathcal{M}$  implies that there exists  $w \in W$  such that  $w.\lambda_0 = \bar{\lambda}_0$ . This shows that  $\phi_{\bar{\lambda}_0} = \phi_{\lambda_0}$ . Therefore, for a suitable  $K$ -biinvariant function  $f$  on  $G$ ,

$$\int_G \phi_{\lambda_0}(x)(f * f^*)(x^{-1}) dx = \hat{f}(\lambda_0)\widehat{f^*}(\lambda_0) = |\hat{f}(\lambda_0)|^2 \geq 0.$$

This proves our claim.

The same example also shows that the set of positive definite functions is a proper subset of the set of  $K$ -positive definite functions.

We now see in the real rank-one case that if we restrict our attention to certain classes of functions, then the set of positive definite functions is same as the set of  $K$ -positive definite functions. Any real rank-one connected, noncompact, semisimple Lie group  $G$  with finite centre can be classified (up to coverings) as

- (1)  $G = \text{SO}_e(1, n)$ , for which  $m_\alpha = n - 1$  and  $m_{2\alpha} = 0$ ,
- (2)  $G = \text{SU}(1, n)$ , for which  $m_\alpha = 2n - 2$  and  $m_{2\alpha} = 1$ ,
- (3)  $G = \text{Sp}(1, n)$ , for which  $m_\alpha = 4n - 4$  and  $m_{2\alpha} = 3$ , or
- (4)  $G = \text{F}_{4(-20)}$ , for which  $m_\alpha = 8$  and  $m_{2\alpha} = 7$ .

Let  $\mathcal{P}_K$  and  $\mathcal{P}$  be the set of  $K$ -positive definite functions and the set of positive definite functions on  $G$  respectively.

**Proposition 3.4.** (1) For the groups  $G = \text{SO}_e(1, n)$  and  $G = \text{SU}(1, n)$ , we have  $\mathcal{P}_K \cap L^\infty(G//K) = \mathcal{P} \cap L^\infty(G//K)$ .

(2) For the group  $G = \text{Sp}(1, n)$ , we have  $\mathcal{P}_K \cap L^r(G//K) = \mathcal{P} \cap L^r(G//K)$ , for any  $2 < r \leq (2n + 1)$ .

(3) For the group  $G = \text{F}_{4(-20)}$ , we have  $\mathcal{P}_K \cap L^r(G//K) = \mathcal{P} \cap L^r(G//K)$ , for any  $2 < r \leq \frac{11}{3}$ .

*Proof.* It is known [Flensted-Jensen and Koornwinder 1979] that  $\phi_\lambda$  is positive definite if and only if  $\lambda \in \mathbb{R}$  or  $\lambda = i\eta$  for  $\eta \in [-s_0, s_0] \cup \{\pm\rho\}$ , where  $s_0 = \rho$  if  $m_{2\alpha} = 0$ , otherwise  $s_0 = \frac{1}{2}m_\alpha + 1$ . Therefore:

- (a) For the groups  $G = SO_e(1, n)$  and  $G = SU(1, n)$ ,  $\phi_\lambda$  is positive definite if and only if  $\lambda \in \mathbb{R}$  or  $\lambda = i\eta$  with  $\eta \in [-\rho, \rho]$ .
- (b) For the group  $G = Sp(1, n)$ ,  $\phi_\lambda$  is positive definite if and only if  $\lambda \in \mathbb{R}$  or  $\lambda = i\eta$  with  $\eta \in [-(2n - 1), (2n - 1)] \cup \{\pm(2n + 1)\}$ .
- (c) For the group  $G = F_{4(-20)}$ ,  $\phi_\lambda$  is positive definite if and only if  $\lambda \in \mathbb{R}$  or  $\lambda = i\eta$  with  $\eta \in [-5, 5] \cup \{\pm 11\}$ .

We know that

$$(3-2) \quad \phi_\lambda \in L^\infty(G//K) \text{ if and only if } \lambda \in S_1,$$

where  $S_r = \{\lambda \in \mathbb{C} : |\text{Im } \lambda| \leq (2/r - 1)\rho\}$ ,  $r > 0$ . Also, for  $r > 1$ ,

$$(3-3) \quad \phi_\lambda \in L^{r'}(G//K) \text{ if and only if } \lambda \in S_r^\circ,$$

where  $1/r + 1/r' = 1$  [Pusti et al. 2011]. Now, from Theorem 3.3, a  $K$ -positive definite function can be expressed as

$$f(x) = \int_{\mathbb{R}} \phi_\lambda(x) d\mu_1(\lambda) + \int_{i\mathbb{R}} \phi_\lambda(x) d\mu_2(\lambda),$$

where  $\mu_1$  is a finite positive measure and  $\mu_2$  is a positive measure such that the integral  $\int_{\mathbb{R}} \phi_{i\lambda}(x) d\mu_2(\lambda)$  exists for every  $x \in G$ .

If the  $K$ -positive definite function  $f$  is in  $L^\infty(G//K)$ , the measure  $\mu_2$  must be supported in  $i[-\rho, \rho]$  by (3-2). Therefore a  $K$ -positive definite function  $f$  which is in  $L^\infty(G//K)$  has an integral form:

$$f(x) = \int_{\mathbb{R}} \phi_\lambda(x) d\mu_1(\lambda) + \int_{i[-\rho, \rho]} \phi_\lambda(x) d\mu_2(\lambda).$$

However, this is a positive definite function for the groups  $G = SO_e(1, n)$  and  $G = SU(1, n)$ , by (a). This proves (1).

To prove (2), note that if the  $K$ -positive definite function  $f$  is in  $L^r(G//K)$ ,  $2 < r \leq (2n + 1)$  by (3-3) it follows that the measure  $\mu_2$  must be supported in

$$i\mathbb{R} \cap S_r^\circ \subseteq i(-(2n - 1), (2n - 1)).$$

This proves that the function  $f$  is positive definite.

The proof of (3) is similar. □

**Remark 3.5.** For the groups  $G = SO_e(1, n)$  and  $G = SU(1, n)$ , it follows from the above that a  $K$ -positive definite function is a positive definite function if and only if  $f \in L^\infty(G//K)$ . However, a similar statement is not true for the groups  $G = Sp(1, n)$  and  $G = F_{4(-20)}$ . In fact, for the group  $G = Sp(1, n)$ , the function  $\phi_{(2n+1)i}$  is  $K$ -positive definite as well as positive definite, but it does not belong to any  $L^r(G//K)$ ,  $2 < r \leq (2n + 1)$ . Similarly, for the group  $G = F_{4(-20)}$ , the function

$\phi_{11i}$  is  $K$ -positive definite as well as positive definite but it does not belong to any  $L^r(G//K)$ ,  $2 < r \leq \frac{11}{3}$ .

Let  $f \in \mathcal{C}^2(G//K)$  be a  $K$ -positive definite function. Then by Theorem 3.3 there exists a finite positive measure  $\sigma$ , invariant under the Weyl group action such that

$$f(x) = \int_{\mathcal{M} \cap \mathfrak{a}^*} \phi_\lambda(x) d\sigma(\lambda).$$

However, this is a positive definite function on  $G$  because  $\phi_\lambda$  is positive definite for  $\lambda \in \mathfrak{a}^*$ . Hence  $\mathcal{P}_K \cap \mathcal{C}^2(G//K) = \mathcal{P} \cap \mathcal{C}^2(G//K)$  (cf. [Bopp 1979] for distributions).

#### 4. $\tau$ -positive definite functions

In this section we give an example in which the set of  $\tau$ -positive definite functions is same as the set of positive definite functions (without imposing any decay condition on functions). For defining the  $\tau$ -positive definite functions we recall some basic facts [Camporesi 1997; Camporesi and Pedon 2001].

**Definition 4.1.** For  $\tau \in \widehat{K}$  a scalar valued function  $f$  on  $G$  is said to be  $\tau$ -radial if  $f(kxk^{-1}) = f(x)$  for all  $k \in K, x \in G$  and if  $d_\tau \bar{\chi}_\tau * f = f = f * d_\tau \bar{\chi}_\tau$ , where  $\chi_\tau$  and  $d_\tau$  are respectively the character and dimension of  $\tau$ .

When  $\tau$  is the trivial representation of  $K$ , a  $\tau$ -radial function is a  $K$ -biinvariant function. We note that the  $\tau$ -radial functions are radial sections of the homogeneous vector bundle over  $G/K$  associated with the representation  $\tau \in \widehat{K}$ . The set of all compactly supported  $\tau$ -radial infinitely differentiable functions and infinitely differentiable  $\tau$ -radial functions are denoted by  $C_{c,\tau}^\infty(G)$  and  $C_\tau^\infty(G)$ , respectively.

**Definition 4.2.** A  $\tau$ -radial continuous function  $f$  on  $G$  is called  $\tau$ -positive definite if

$$\int_G f(x)(g * g^*)(x^{-1}) dx \geq 0, \quad \text{for all } g \in C_{c,\tau}^\infty(G).$$

Let  $G = \text{Spin}_e(n, 1)$ , the identity component of  $\text{Spin}(n, 1)$ . Then, in the notation of the previous section,  $K = \text{Spin}(n)$  and  $M = \text{Spin}(n-1)$ . In the rest of the section we fix these meanings for  $G, K, M$ .

Let  $\tau_n$  be the complex spin representation of  $K$ . The following proposition gives information about the irreducibility of  $\tau_n$ .

**Proposition 4.3** [Camporesi and Pedon 2001]. (1) *If  $n$  is even, then  $\tau_n$  splits into two irreducible components given by the positive and negative half-spin representations  $\tau_n = \tau_n^+ \oplus \tau_n^-$  and  $\tau_n^\pm|_M = \sigma_{n-1}$ , where  $\sigma_{n-1}$  is the spin representation of  $M$ .*

(2) *If  $n$  is odd, then  $\tau_n$  is irreducible and  $\tau_n|_M = \sigma_{n-1}^+ \oplus \sigma_{n-1}^-$ , where  $\sigma_{n-1}^\pm$ , are irreducible components of the spin representation  $\sigma_{n-1}$  of  $M$ .*

It is known that  $(G, K, \tau)$  is a Gelfand triple, that is, the convolution algebra  $C_{c,\tau}^\infty(G)$  is commutative when  $\tau \in \widehat{K}$  is either  $\tau_n^+$  or  $\tau_n^-$  if  $n$  is even and  $\tau_n$  if  $n$  is odd. For  $n$  even the  $\tau_n^\pm$ -spherical function is given by

$$\phi_\lambda^{\tau_n^\pm}(x) = \int_K e^{-(i\lambda+\rho)H(xk)} \chi_{\tau_n^\pm}(kK(xk)^{-1}) dk.$$

Also, it satisfies  $\phi_{-\lambda}^{\tau_n^\pm}(x) = \phi_\lambda^{\tau_n^\pm}(x)$ .

For  $n$  odd the  $\tau_n$ -spherical functions are denoted by  $\phi_{\sigma_{n-1},\lambda}^{\tau_n^+}$  and  $\phi_{\sigma_{n-1},\lambda}^{\tau_n^-}$ . They are given by the integral formula

$$\phi_{\sigma_{n-1},\lambda}^{\tau_n^\pm}(x) = 2d_{\sigma_{n-1}^\pm} \int_K \int_M e^{-(i\lambda+\rho)H(xk)} \chi_{\tau_n}(km^{-1}K(xk)^{-1}) \chi_{\sigma_{n-1}^\pm}(m) dm dk.$$

They satisfy  $\phi_{\sigma_{n-1},-\lambda}^{\tau_n^+}(x) = \phi_{\sigma_{n-1},\lambda}^{\tau_n^-}(x)$ .

From now on by  $\tau \in \widehat{K}$  we will mean either  $\tau = \tau_n^+$  or  $\tau = \tau_n^-$  if  $n$  is even and  $\tau = \tau_n$  if  $n$  is odd. For  $n$  even we shall write the  $\tau_n^\pm$ -spherical functions  $\phi_{\sigma,\lambda}^\tau$  instead of  $\phi_{\sigma_{n-1},\lambda}^{\tau_n^\pm}$ . Also for  $n$  odd we write the  $\tau_n$ -spherical functions

$$\phi_{\sigma^\pm,\lambda}^\tau \quad \text{instead of} \quad \phi_{\sigma_{n-1},\lambda}^{\tau_n}$$

Henceforth while dealing with  $G = \text{Spin}_e(n, 1)$  and  $\tau$  as above we shall simply say *when  $n$  is even* and *when  $n$  is odd* to distinguish between these two cases.

For a  $\tau$ -radial function  $f$  its spherical Fourier transform is defined by

$$\hat{f}(\sigma, \lambda) = \int_G f(x) \phi_{\sigma,\lambda}^\tau(x^{-1}) dx$$

when  $n$  is even. For  $n$  odd it is defined by

$$\hat{f}(\sigma^\pm, \lambda) = \int_G f(x) \phi_{\sigma^\pm,\lambda}^\tau(x^{-1}) dx.$$

**Theorem 4.4** [Gelfand and Vilenkin 1964, Theorem 3, p. 157, Theorem 5, p. 226].

(a) *Let  $T$  be a positive definite distribution on  $\mathbb{R}$ , that is,*

$$T(\phi * \phi^*) \geq 0 \quad \text{for all } \phi \in C_c^\infty(\mathbb{R}).$$

*Then there exists a positive tempered measure  $\mu$  on  $\mathbb{R}$  such that*

$$T(\phi) = \int_{\mathbb{R}} \tilde{\phi}(\lambda) d\mu(\lambda) \quad \text{for all } \phi \in C_c^\infty(\mathbb{R}).$$

(b) *Let  $T$  be an evenly positive definite distribution on  $\mathbb{R}$ , that is,  $T(\phi * \phi^*) \geq 0$  for all  $\phi \in C_c^\infty(\mathbb{R})_e$ . Then there exists positive even measures  $\mu_1$  and  $\mu_2$  such that*

$$T(\phi) = \int_{\mathbb{R}} \tilde{\phi}(\lambda) d\mu_1(\lambda) + \int_{\mathbb{R}} \tilde{\phi}(i\lambda) d\mu_2(\lambda) \quad \text{for all } \phi \in C_c^\infty(\mathbb{R})_e,$$

where  $\mu_1$  is a tempered measure and  $\mu_2$  is such that

$$\int_{\mathbb{R}} e^{a|\lambda|} d\mu_2(\lambda) < \infty \quad \text{for all } a > 0.$$

The next theorem gives integral representations of  $\tau$ -positive definite functions on  $G = \text{Spin}_e(n, 1)$ .

**Theorem 4.5.** *Let  $G = \text{Spin}_e(n, 1)$  and let  $\tau$  denote one of  $\{\tau_n^+, \tau_n^-\}$  when  $n$  is even and  $\tau_n$  when  $n$  is odd.*

(a) *Let  $n$  be even and let  $f$  be a  $\tau$ -positive definite function on  $G$ . Then there exists even positive measures  $\mu_1$  and  $\mu_2$  such that for all  $x \in G$*

$$f(x) = \int_{\mathbb{R}} \phi_{\sigma,\lambda}^\tau(x) d\mu_1(\lambda) + \int_{\mathbb{R}} \phi_{\sigma,i\lambda}^\tau(x) d\mu_2(\lambda),$$

where  $\mu_1$  is finite measure and  $\mu_2$  is such that

$$\int_{\mathbb{R}} e^{a|\lambda|} d\mu_2(\lambda) < \infty \quad \text{for all } a > 0.$$

(b) *Let  $n$  be odd and let  $f$  be a  $\tau$ -positive definite function on  $G$ . Then there exists a finite positive measure  $\mu$  such that for all  $x \in G$*

$$f(x) = \int_{\mathbb{R}} \phi_{\sigma^+,\lambda}^\tau(x) d\mu(\lambda).$$

*Proof.* We shall prove (b). The proof of (a) is similar. Let  $n$  be odd and let  $f$  be a  $\tau$ -positive definite function on  $G$ . We define the linear functional  $T_f$  on  $C_c^\infty(\mathbb{R})$  as follows:

$$T_f(h) = \int_G f(x)(\mathcal{A}^{-1}h)(x^{-1}) dx \quad \text{for all } h \in C_c^\infty(\mathbb{R}).$$

Here  $\mathcal{A}$  is the Abel transform, which is a topological isomorphism between  $C_{c,\tau}^\infty(G)$  and  $C_c^\infty(\mathbb{R})$ . We also have  $\hat{f}(-\lambda) = \widetilde{\mathcal{A}f}(\lambda)$  for all  $f \in C_{c,\tau}^\infty(G)$ . Then it follows that  $\widehat{\mathcal{A}^{-1}h}(-\lambda) = \widetilde{h}(\lambda)$  for all  $h \in C_c^\infty(\mathbb{R})$ . Using this, we easily check that

$$\mathcal{A}^{-1}(h_1 * h_2) = \mathcal{A}^{-1}h_1 * \mathcal{A}^{-1}h_2$$

and  $\mathcal{A}^{-1}h_1^* = (\mathcal{A}^{-1}h_1)^*$  for all  $h_1, h_2 \in C_c^\infty(\mathbb{R})$ . Then

$$T_f(h * h^*) = \int_G f(x)(\mathcal{A}^{-1}h * (\mathcal{A}^{-1}h)^*)(x^{-1}) dx \geq 0$$

as  $f$  is  $\tau$ -positive definite. Therefore, by (a), there exists a positive tempered measure  $\mu$  on  $\mathbb{R}$  such that for all  $h \in C_c^\infty(\mathbb{R})$

$$T_f(h) = \int_{\mathbb{R}} \widetilde{h}(\lambda) d\mu(\lambda).$$

This shows that for all  $g \in C_{c,\tau}^\infty(G)$

$$(4-1) \quad \int_G f(x)g(x^{-1})dx = \int_{\mathbb{R}} \widehat{g}_+(\lambda) d\mu(\lambda).$$

Using approximate identity techniques we can easily prove that the measure  $\mu$  is finite. Then from Equation (4-1), using Fubini's theorem we get

$$\begin{aligned} \int_G f(x)g(x^{-1})dx &= \int_{\mathbb{R}} \int_G g(x)\phi_{\sigma^+,\lambda}^\tau(x^{-1})dx d\lambda \\ &= \int_G g(x^{-1}) \int_{\mathbb{R}} \phi_{\sigma^+,\lambda}^\tau(x) d\mu(\lambda) dx. \end{aligned}$$

Since this is true for every  $g \in C_{c,\tau}^\infty(G)$ , it follows that

$$f(x) = \int_{\mathbb{R}} \phi_{\sigma^+,\lambda}^\tau(x) d\mu(\lambda). \quad \square$$

It is easy to check that the converse of Theorem 4.5 holds true. We get the following corollary from Theorem 4.5(b):

**Corollary 4.6.** *The set of  $\tau$ -positive definite functions is same as the set of positive definite functions when  $\tau = \tau_n$  and  $n$  is odd.*

**Remark 4.7.** We saw after Theorem 3.3 that the function  $\phi_{\lambda_0}, \lambda_0 \in \mathcal{M} \setminus \mathfrak{a}_1^*$  is  $K$ -positive definite but not positive definite. When  $\tau = \tau_n$  and  $n$  is odd, we could try to find a similar example by considering the function  $\phi_{\sigma^+,\lambda_0}^\tau, \lambda_0 \in i\mathbb{R} \setminus i[-1, 1]$ . But  $\phi_{\sigma^+,\lambda_0}^\tau$  is neither a  $\tau$ -positive definite function nor a positive definite function. The argument used in the  $K$ -positive definite case does not work here. Indeed, unlike the case of the spherical functions  $\phi_\lambda$ , which are  $W$ -invariant in  $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ , there is no relation between  $\phi_{\sigma^+,\lambda}^\tau$  and  $\phi_{\sigma^+,-\lambda}^\tau$  when  $\tau = \tau_n$  and  $n$  is odd.

We Now characterize the  $\lambda$ 's for which  $\phi_{\sigma,\lambda}^\tau$  is positive definite for

$$G = \text{Spin}_e(n, 1)$$

when  $\tau$  is the irreducible component of the complex spin representation.

**Theorem 4.8.** *Let  $G = \text{Spin}_e(n, 1)$  and let  $\tau$  denote one of  $\{\tau_n^+, \tau_n^-\}$  when  $n$  is even and  $\tau_n$  when  $n$  is odd. Then*

- (a)  $\phi_{\sigma,\lambda}^\tau$  is positive definite if and only if  $\lambda \in \mathbb{R}$  when  $n(\geq 4)$  is even, and
- (b)  $\phi_{\sigma^\pm,\lambda}^\tau$  are positive definite if and only if  $\lambda \in \mathbb{R}$  when  $n$  is odd.

*Proof.* (a) Let  $n$  be even and  $n \geq 4$ . The  $\tau$ -spherical function  $\phi_{\sigma,\lambda}^\tau$  is positive definite if and only if  $\tau$  is contained in the unitary principal, discrete or complementary series representations. It is well-known that there is no discrete series representation which contains  $\tau$ . Also, by [Knapp and Stein 1971, Proposition 55]

and the Frobenius reciprocity theorem there is no complementary series containing  $\tau$ . Hence  $\phi_{\sigma,\lambda}^\tau$  is positive definite if and only if  $\lambda \in \mathbb{R}$ .

(b) For the case  $n$  odd we prove the result without using representation theory. By Corollary 4.6 the  $\tau$ -spherical function  $\phi_{\sigma^+,\lambda}^\tau$  is positive definite if and only if it is a  $\tau$ -positive definite function. That is equivalent to

$$\int_G (f * f^*)(x) \phi_{\sigma^+,\lambda}^\tau(x^{-1}) dx \geq 0 \quad \text{for all } f \in \mathcal{C}_\tau^2(G),$$

where  $\mathcal{C}_\tau^2(G)$  is the set of  $\tau$ -radial  $L^2$ -Schwartz class functions on  $G$ . That is,

$$(4-2) \quad \hat{f}(\sigma^+, \lambda) \overline{\hat{f}(\sigma^+, \bar{\lambda})} \geq 0 \quad \text{for all } f \in \mathcal{C}_\tau^2(G),$$

since  $\overline{\phi_{\sigma,\lambda}^\tau(x)} = \phi_{\sigma,\bar{\lambda}}^\tau(x^{-1})$ . Let us consider a function

$$f \in \mathcal{C}_\tau^2(G)$$

such that  $\hat{f}(\sigma^+, \lambda) = \lambda e^{-\lambda^2}$ . Such a function exists by the Schwartz space isomorphism theorem [Camporesi and Pedon 2001, Theorem 6.3]. Then (4-2) is true if and only if  $\lambda \in \mathbb{R}$ .  $\square$

### Acknowledgments

We are grateful to Prof. J. Faraut for giving us the problem and many important suggestions.

### References

- [Anker 1991] J.-P. Anker, “The spherical Fourier transform of rapidly decreasing functions. A simple proof of a characterization due to Harish-Chandra, Helgason, Trombi, and Varadarajan”, *J. Funct. Anal.* **96**:2 (1991), 331–349. MR 92d:22008 Zbl 0732.43006
- [Barker 1975] W. H. Barker, “The spherical Bochner theorem on semisimple Lie groups”, *J. Functional Analysis* **20**:3 (1975), 179–207. MR 53 #3197 Zbl 0308.43002
- [Bopp 1979] N. Bopp, “Les distributions de type positif relativement à un groupe fini d’isométries”, pp. 15–50 in *Analyse harmonique sur les groupes de Lie, II* (Sém., Nancy-Strasbourg 1976–1978), edited by P. Eymard et al., Lecture Notes in Math. **739**, Springer, Berlin, 1979. MR 81e:22018 Zbl 0422.46034
- [Camporesi 1997] R. Camporesi, “The spherical transform for homogeneous vector bundles over Riemannian symmetric spaces”, *J. Lie Theory* **7**:1 (1997), 29–60. MR 98j:58114 Zbl 0871.43005
- [Camporesi and Pedon 2001] R. Camporesi and E. Pedon, “Harmonic analysis for spinors on real hyperbolic spaces”, *Colloq. Math.* **87**:2 (2001), 245–286. MR 2002c:43012 Zbl 0968.22009
- [van Dijk and Pasquale 1999] G. van Dijk and A. Pasquale, “Harmonic analysis on vector bundles over  $\mathrm{Sp}(1, n)/\mathrm{Sp}(1) \times \mathrm{Sp}(n)$ ”, *Enseign. Math.* (2) **45**:3-4 (1999), 219–252. MR 2001b:43011 Zbl 0967.43005

- [Flensted-Jensen and Koornwinder 1979] M. Flensted-Jensen and T. H. Koornwinder, "Positive definite spherical functions on a noncompact, rank one symmetric space", pp. 249–282 in *Analyse harmonique sur les groupes de Lie, II* (Sém., Nancy and Strasbourg 1976–1978), edited by P. Eymard et al., Lecture Notes in Math. **739**, Springer, Berlin, 1979. MR 81j:43015 Zbl 0433.43014
- [Gelfand and Vilenkin 1964] I. M. Gelfand and N. Y. Vilenkin, *Applications of harmonic analysis, Generalized functions* **4**, Academic Press, New York, 1964. MR 30 #4152 Zbl 0136.11201
- [Knapp and Stein 1971] A. W. Knap and E. M. Stein, "Intertwining operators for semisimple groups", *Ann. of Math. (2)* **93** (1971), 489–578. MR 57 #536 Zbl 0257.22015
- [Pusti 2011] S. Pusti, "An analogue of Bochner's theorem for Damek–Ricci spaces", preprint, 2011. arXiv 1111.3157
- [Pusti et al. 2011] S. Pusti, S. K. Ray, and R. P. Sarkar, "Wiener–Tauberian type theorems for radial sections of homogenous vector bundles on certain rank one Riemannian symmetric spaces of noncompact type.", *Math. Z.* **269**:1-2 (2011), 555–586. Zbl 05963446
- [Sitaram 1978] A. Sitaram, "Positive definite distributions on  $K|G/K$ . II", *J. Indian Math. Soc. (N.S.)* **42**:1-4 (1978), 95–104 (1979). MR 81e:43024 Zbl 0499.43010

Received July 10, 2011. Revised July 19, 2011.

SANJOY PUSTI  
MATHEMATICS RESEARCH UNIT  
UNIVERSITY OF LUXEMBOURG, CAMPUS KIRCHBERG  
6, RUE RICHARD COUDENHOVE-KALERGI  
1359 LUXEMBOURG  
LUXEMBOURG  
sanjoy.pusti@uni.lu

# PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by  
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [pacificmath.org](http://pacificmath.org) for submission instructions.

---

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 254 No. 2 December 2011

---

Curvatures of spheres in Hilbert geometry	257
ALEXANDER BORISENKO and EUGENE OLIN	
A formula equating open and closed Gromov–Witten invariants and its applications to mirror symmetry	275
KWOKWAI CHAN	
A note on $p$ -harmonic $l$ -forms on complete manifolds	295
LIANG-CHU CHANG and CHIUNG-JUE ANNA SUNG	
The Cheeger constant of curved strips	309
DAVID KREJČIŘÍK and ALDO PRATELLI	
Structure of solutions of 3D axisymmetric Navier–Stokes equations near maximal points	335
ZHEN LEI and QI S. ZHANG	
Local comparison theorems for Kähler manifolds	345
GANG LIU	
Structurable algebras of skew-rank 1 over the affine plane	361
SUSANNE PUMPLÜN	
An analogue of Krein’s theorem for semisimple Lie groups	381
SANJOY PUSTI	
Une remarque de dynamique sur les variétés semi-abéliennes	397
GAËL RÉMOND	
Fourier transforms of semisimple orbital integrals on the Lie algebra of $SL_2$	407
LOREN SPICE	
On noncompact $\tau$ -quasi-Einstein metrics	449
LIN FENG WANG	
Decomposition of de Rham complexes with smooth horizontal coefficients for semistable reductions	465
QIHONG XIE	
A differentiable sphere theorem inspired by rigidity of minimal submanifolds	499
HONG-WEI XU and LING TIAN	



0030-8730(201112)254:2;1-A