

*Pacific
Journal of
Mathematics*

ON NONCOMPACT τ -QUASI-EINSTEIN METRICS

LIN FENG WANG

Volume 254 No. 2

December 2011

ON NONCOMPACT τ -QUASI-EINSTEIN METRICS

LIN FENG WANG

In this paper, we will study the τ -quasi-Einstein metrics on complete non-compact Riemannian manifolds and get a rigid property. We will also obtain lower and upper estimates for scalar curvatures on these metrics by using the maximum principle.

1. Introduction

For a given smooth potential function f , the τ -Bakry–Émery Ricci curvature tensor

$$\text{Ric}_{f,\tau} = \text{Ric} + \text{Hess } f - \frac{\nabla f \otimes \nabla f}{\tau}$$

is always used to replace the Ricci curvature tensor when one tries to study the weighted measure $d\mu = e^{-f} dx$, where $0 < \tau \leq +\infty$ and dx is the Riemann–Lebesgue measure determined by the metric. There has been an active interest in the study of the weighted measure under some conditions about the τ -Bakry–Émery Ricci curvature tensor; see [Li 2005; Wang 2010] and the references therein.

According to [Kim and Kim 2003; Case 2010; Case et al. 2011; Wang 2011], we call a metric g τ -quasi-Einstein with potential function f , if for some constant λ ,

$$(1-1) \quad \text{Ric} + \text{Hess } f - \frac{\nabla f \otimes \nabla f}{\tau} = \lambda g,$$

where $0 < \tau \leq +\infty$. A τ -quasi-Einstein metric becomes an Einstein metric when the potential function f is constant. We note that an ∞ -quasi-Einstein metric indicates a gradient Ricci soliton. As in [Hamilton 1995; Perelman 2002; Cao and Zhu 2006], a gradient Ricci soliton is shrinking, steady or expanding when $\lambda > 0$, $\lambda = 0$ or $\lambda < 0$, respectively.

For a positive integer τ , the τ -quasi-Einstein metric is closely relative to the existence of warped product Einstein manifolds [Besse 1987; Case 2010; Case et al. 2011]. Let (M, g) and (N^τ, h) be two Riemannian manifolds. Then, for

The author was supported in part by the NSF of China(10871070, 10971066, 11171254).

MSC2010: primary 53C21; secondary 53C25.

Keywords: τ -quasi-Einstein metric, rigidity properties, maximum principle, potential function.

some potential function f on M , the warped product manifold $(M \times N, \tilde{g})$ with product metric

$$\tilde{g} = g \oplus \exp\left(-\frac{2f}{\tau}\right)h$$

is Einstein if and only if (N^τ, h) is Einstein and the Ricci curvature tensor of M satisfies the quasi-Einstein equation (1-1) for some constant λ .

It was proved in [Qian 1997; Wei and Wylie 2007] that a manifold with a τ -quasi-Einstein metric (τ is finite) is automatically compact when $\lambda > 0$. It was also proved in [Ivey 1993] that any expanding or steady gradient Ricci solitons on closed manifolds should be trivial. The same rigid properties for the τ -quasi-Einstein metrics on closed manifolds were proved in [Kim and Kim 2003; Wang 2011]. But for the τ -quasi-Einstein metrics on closed manifolds with $\lambda > 0$, the rigid properties rely on the constant μ which appears in the following identity:

$$(1-2) \quad R + \frac{\tau-1}{\tau}|\nabla f|^2 + (\tau-n)\lambda = \mu e^{2f/\tau},$$

where R is the scalar curvature. This identity was proved in [Kim and Kim 2003]. See also [Wang 2011], where the author proved that the quasi-Einstein metrics with $\lambda > 0$ should be trivial when $\mu \leq 0$. In fact, the authors of [Lü et al. 2004] constructed nontrivial τ -quasi-Einstein metrics with $\lambda > 0$ and $\tau > 1$, which also satisfy $\mu > 0$.

In this paper, we will study the τ -quasi-Einstein metrics on complete noncompact Riemannian manifolds with $\lambda \leq 0$. Our first result is Theorem 1.1, which is about the rigidity.

Theorem 1.1. *Let M be a complete noncompact Riemannian manifold and g a τ -quasi-Einstein metric on M with potential function f and $\lambda \leq 0$ a constant. If*

$$(1-3) \quad R_0^{-2} \int_{B_{2R_0} \setminus B_{R_0}} |\nabla f|^2 \exp\left(-\frac{\tau+2}{\tau}f\right) dx \rightarrow 0$$

as $R_0 \rightarrow \infty$, where B_{R_0} denotes the geodesic ball centered at a fixed point $O \in M$ with radius R_0 , then e^f is a harmonic function on M , that is, $\Delta e^f = 0$. Moreover, if $\lambda < 0$, then g is trivial in the sense that f is constant.

The following theorem for gradient Ricci solitons was proved in [Zhang 2009]. In fact, part 1 is a consequence of [Chen 2009, Corollary 2.5].

Theorem 1.2. *Let (M^n, g) be a complete noncompact gradient Ricci soliton with potential function f and soliton constant λ .*

- (1) *If the gradient Ricci soliton is shrinking or steady, then $R \geq 0$.*
- (2) *If the gradient Ricci soliton is expanding, then there exists a positive constant $C(n)$ such that $R \geq C(n)\lambda$.*

Zhang [2011] pointed out that $R \geq n\lambda$ is right in Theorem 1.2(2). The lower bound estimates for scalar curvatures play important roles in the study of geometric properties of gradient Ricci solitons. Based on these estimates, compactness theorems for gradient Ricci solitons were proved in [Zhang 2006] and some results about the volume growth for noncompact gradient Ricci solitons were deduced in [Cao 2009; Cao and Zhou 2010; Munteanu 2009; Zhang 2011].

In [Case et al. 2011], the authors got estimates for R on closed τ -quasi-Einstein metrics. Later, Wang [2011] studied the lower bound estimate for scalar curvature R on complete noncompact τ -quasi-Einstein metrics with $\lambda \leq 0$. We state this result as follows.

Theorem 1.3. *Let M be an n -dimensional complete noncompact Riemannian manifold, metric g is τ -quasi-Einstein with potential function f and constant $\lambda \leq 0$, where $\tau \geq 1$. If $\mu \leq 0$ or $\mu > 0$ and f is bounded from above by a constant C , then*

$$(1-4) \quad R(y) \geq n\lambda$$

for any $y \in M$.

The proof of this theorem in [Wang 2011] relies on a gradient estimate of f , this gradient estimate shows that $|\nabla f|^2$ is bounded from above if $\mu \leq 0$ or $\mu > 0$ and f is bounded from above by a constant C . We will give a nontrivial τ -quasi-Einstein metric with $\lambda < 0$, but f is not bounded from above; see Example 2.1. The second main result of this paper is to improve Theorem 1.3. That is to say, we will show that the lower estimate (1-4) is always right for τ -quasi-Einstein metrics with $\lambda \leq 0$.

Theorem 1.4. *Let M be an n -dimensional complete noncompact Riemannian manifold, g be a τ -quasi-Einstein metric with potential function f and $\lambda \leq 0$ be a constant, where $\tau > 0$. Then (1-4) holds for any $y \in M$.*

Remark 1.5. If $\tau = \infty$, we recover the lower bound estimate for R on a complete noncompact steady or expanding gradient Ricci soliton given in [Zhang 2011].

It remains interesting to find out whether R is bounded from above by a constant for noncompact quasi-Einstein metrics. The following theorem states that the scalar curvature of a quasi-Einstein metric with $\lambda \leq 0$ is bounded from above if $\mu \leq 0$.

Theorem 1.6. *Let g be a τ -quasi-Einstein metric with $\lambda \leq 0$ and $\mu \leq 0$. Then*

$$(1-5) \quad R(y) \leq (n - \max\{\tau, 1\})\lambda$$

for any $y \in M$.

2. Examples of quasi-Einstein metrics

In this section, we assume that $M = \mathbb{R} \times N^{n-1}$ is a warped product manifold with the product metric given by

$$ds_M^2 = dt^2 + \varphi^2(t) ds_N^2,$$

where ds_N^2 is a fixed metric on N and φ is a positive function on \mathbb{R} . Consider the orthonormal coframe $\{\theta_\alpha : 2 \leq \alpha \leq n\}$ on N^{n-1} ; then

$$\{\omega_1 = dt, \omega_\alpha = \varphi(t)\theta_\alpha : 2 \leq \alpha \leq n\}$$

is an orthonormal coframe on M^n . We use $R_{M,ijkl}$ and $R_{N,\alpha\beta\gamma\delta}$ to denote the Riemannian curvature tensors of M and N respectively. After the same calculation as in [O'Neill 1983; Wang 2011], we conclude that

$$(2-1) \quad R_{M,1\alpha ij} = \begin{cases} -(\log \varphi(t))'' - ((\log \varphi(t))')^2 & \text{if } i = 1, j = \alpha, \\ (\log \varphi(t))'' + ((\log \varphi(t))')^2 & \text{if } i = \alpha, j = 1, \\ 0 & \text{otherwise} \end{cases}$$

and

$$(2-2) \quad R_{M,\alpha\beta ij} = \begin{cases} \varphi^{-2}(t)R_{N,\alpha\beta\gamma\theta} + ((\log \varphi(t))')^2(\delta_{\alpha\theta}\delta_{\beta\gamma} - \delta_{\alpha\gamma}\delta_{\beta\theta}) & \text{if } i = \gamma, j = \theta, \\ 0 & \text{otherwise.} \end{cases}$$

If we use $R_{N,\alpha\beta}$ to denote the Ricci curvature tensor on N , by (2-1) and (2-2), the Ricci curvature tensor of M can be expressed as

$$(2-3) \quad R_{M,1i} = -(n-1)((\log \varphi(t))'' + ((\log \varphi(t))')^2)\delta_{1i},$$

$$(2-4) \quad R_{M,\alpha\beta} = \varphi^{-2}(t)R_{N,\alpha\beta} - ((\log \varphi(t))'' + (n-1)((\log \varphi(t))')^2)\delta_{\alpha\beta}.$$

Example 2.1. For $\tau > 0$, we assume that N is a flat manifold with

$$R_{N,\alpha\beta} = 0.$$

Let

$$f(t, x) = f(t) = \tau t, \quad \varphi(t) = e^{-t}.$$

It is easy to testify that

$$(2-5) \quad R_{M,ij} + f_{ij} - \frac{f_i f_j}{\tau} = \lambda g_{ij}$$

for $\lambda = -(n + \tau - 1)$. Hence M is τ -quasi-Einstein with potential function $f = \tau t$ and $\lambda = -(n + \tau - 1)$. Moreover, by (2-3) and (2-4), the scalar curvature of M is

$$R_M = -n(n-1),$$

which means that (1-2) follows with $\mu = 0$. It is easy to see that the potential function f is not bounded from above.

Example 2.2. For $\tau > 0$, we assume that N is an Einstein manifold with

$$R_{N,\alpha\beta} = -(n + \tau - 2)\delta_{\alpha\beta}.$$

Choose

$$f(t, x) = f(t) = -\tau \log \cosh t, \quad \varphi(t) = \cosh t.$$

It is easy to testify that (2-5) holds for $\lambda = -(n + \tau - 1)$. Hence M is τ -quasi-Einstein with potential function $f = -\tau \log \cosh t$ and $\lambda = -(n + \tau - 1)$. Moreover, by (2-3) and (2-4), the scalar curvature of M is

$$R_M = -n(n - 1) - \frac{(n - 1)\tau}{\cosh^2 t},$$

which means that (1-2) follows with

$$\mu = -\tau(\tau + n - 2).$$

It is easy to see that $\mu < 0$ and R_M is bounded from above.

3. Basic formulas

In this section, we will first give some basic formulas for quasi-Einstein metrics in Lemma 3.1. These formulas are well-established in [Case et al. 2011; Kim and Kim 2003; Wang 2011].

Lemma 3.1. *If g is a τ -quasi-Einstein metric with potential function f and λ is a constant, then one can get*

$$\begin{aligned} (3-1) \quad \frac{1}{2}\Delta R - \frac{\tau+2}{2\tau}\nabla f \cdot \nabla R \\ = -\frac{\tau-1}{\tau}\left|\text{Ric} - \frac{1}{n}Rg\right|^2 - \frac{n+\tau-1}{n\tau}(R-n\lambda)\left(R - \frac{n(n-1)}{n+\tau-1}\lambda\right). \end{aligned}$$

Moreover, there exists a constant μ such that

$$(3-2) \quad R + \frac{\tau-1}{\tau}|\nabla f|^2 + (\tau - n)\lambda = \mu e^{2f/\tau}.$$

And also one can get

$$(3-3) \quad \nabla \Delta f \cdot \nabla f = \frac{2}{\tau}\Delta f|\nabla f|^2 - 2\text{Ric}(\nabla f, \nabla f)$$

$$(3-4) \quad \Delta f - |\nabla f|^2 - \tau\lambda + \mu e^{2f/\tau} = 0.$$

In the following, we will calculate the weighted Laplacian of $\varphi(R + 2xe^{2f/\tau})$ by using Lemma 3.1, where $x > 0$ is a constant and φ is a smooth cutoff function.

Lemma 3.2. *Let*

$$(3-5) \quad Q = \varphi(R + 2xe^{2f/\tau}),$$

where $x > 0$ is a constant and φ is a smooth cutoff function. If $\tau > 1$ and $\mu > 0$, then for $\epsilon > 0$,

$$(3-6) \quad \frac{1}{2}\Delta_f Q \leq \frac{\Delta_f \varphi}{2\varphi} Q + \frac{\nabla \varphi \cdot \nabla Q}{\varphi} - \frac{|\nabla \varphi|^2}{\varphi^2} Q + \frac{\varphi^2}{4\epsilon\tau} \left| \frac{\nabla Q}{\varphi} - \frac{Q\nabla \varphi}{\varphi^2} \right|^2 \\ + \frac{4x(n+\tau-1)}{n\tau} Q e^{2f/\tau} - \frac{n+\tau-1}{n\tau\varphi} Q^2 \\ + \left(\frac{2n-2+\tau}{\tau} \lambda - \frac{\epsilon}{(\tau-1)\varphi} \right) Q + \varphi A - \frac{n(n-1)\varphi}{\tau} \lambda^2 + \frac{n-\tau}{\tau-1} \epsilon \lambda$$

holds at $y \in M$ with $\varphi(y) \neq 0$, where

$$(3-7) \quad \Delta_f = \Delta - \nabla f \cdot \nabla$$

and A , depending on $x, n, \tau, \mu, \lambda, \epsilon, \varphi$, is defined in (3-14).

Proof. Let

$$(3-8) \quad G = R + 2xe^{2f/\tau}.$$

It is easy to see that

$$(3-9) \quad \Delta_f e^{2f/\tau} = \Delta e^{2f/\tau} - \nabla e^{2f/\tau} \cdot \nabla f = \left(\frac{4-2\tau}{\tau^2} |\nabla f|^2 + \frac{2}{\tau} \Delta f \right) e^{2f/\tau},$$

which, together with (3-1), shows that, for $\epsilon > 0$,

$$\frac{1}{2}\Delta_f G \\ \leq \frac{1}{\tau} \nabla R \cdot \nabla f + x \left(\frac{4-2\tau}{\tau^2} |\nabla f|^2 + \frac{2}{\tau} \Delta f \right) e^{2f/\tau} - \frac{n+\tau-1}{n\tau} (R-n\lambda) \left(R - \frac{n(n-1)}{n+\tau-1} \lambda \right) \\ = \frac{1}{\tau} \nabla G \cdot \nabla f + x \left(-\frac{2}{\tau} |\nabla f|^2 + \frac{2}{\tau} \Delta f \right) e^{2f/\tau} - \frac{n+\tau-1}{n\tau} (R-n\lambda) \left(R - \frac{n(n-1)}{n+\tau-1} \lambda \right) \\ \leq \frac{\varphi}{4\epsilon\tau} |\nabla G|^2 + \frac{\epsilon}{\tau\varphi} |\nabla f|^2 + x \left(-\frac{2}{\tau} |\nabla f|^2 + \frac{2}{\tau} \Delta f \right) e^{2f/\tau} \\ - \frac{n+\tau-1}{n\tau} (R-n\lambda) \left(R - \frac{n(n-1)}{n+\tau-1} \lambda \right)$$

holds at $y \in M$ when $\varphi(y) \neq 0$. By (3-8) and (3-2), we get

$$(3-10) \quad R = G - 2xe^{2f/\tau}$$

and

$$(3-11) \quad |\nabla f|^2 = -\frac{\tau}{\tau-1} G + \frac{\tau(2x+\mu)}{\tau-1} e^{2f/\tau} + \frac{\tau(n-\tau)}{\tau-1} \lambda.$$

Plugging (3-8), (3-10), (3-11) and (3-4) into 3 yields

$$(3-12) \quad \frac{1}{2} \Delta_f G \leq \frac{\varphi}{4\epsilon\tau} |\nabla G|^2 + \frac{4x(n+\tau-1)}{n\tau} G e^{2f/\tau} - \frac{n+\tau-1}{n\tau} G^2 \\ + \left(\frac{2n-2+\tau}{\tau} \lambda - \frac{\epsilon}{(\tau-1)\varphi} \right) G - \frac{4x^2(n+\tau-1) + 2xn\mu}{n\tau} e^{4f/\tau} \\ - \left(\frac{4x(n-1)}{\tau} \lambda - \frac{\epsilon(2x+\mu)}{(\tau-1)\varphi} \right) e^{2f/\tau} - \frac{n(n-1)}{\tau} \lambda^2 + \frac{n-\tau}{(\tau-1)\varphi} \epsilon \lambda.$$

Since for all $a > 0$,

$$-ax^2 + bx \leq \frac{b^2}{4a},$$

we conclude that

$$(3-13) \quad -\frac{4x^2(n+\tau-1) + 2xn\mu}{n\tau} e^{4f/\tau} - \left(\frac{4x(n-1)}{\tau} \lambda - \frac{\epsilon(2x+\mu)}{(\tau-1)\varphi} \right) e^{2f/\tau} \leq A$$

with

$$(3-14) \quad A = \frac{n\tau}{16x^2(n+\tau-1) + 8nx\mu} \left(\frac{4x(n-1)}{\tau} \lambda - \frac{\epsilon(2x+\mu)}{(\tau-1)\varphi} \right)^2.$$

It is easy to see that

$$\nabla G = \frac{\nabla Q}{\varphi} - \frac{Q \nabla \varphi}{\varphi^2}$$

and

$$\Delta_f Q = \frac{Q}{\varphi} \Delta_f \varphi + 2 \nabla \varphi \cdot \nabla G + \varphi \Delta_f G.$$

Hence

$$(3-15) \quad \Delta_f Q = \frac{\Delta_f \varphi}{\varphi} Q + \frac{2 \nabla \varphi \cdot \nabla Q}{\varphi} - \frac{2 |\nabla \varphi|^2}{\varphi^2} Q + \varphi \Delta_f G.$$

Plugging (3-5), (3-12) and (3-13) into (3-15) yields (3-6). □

4. A rigid property

In this section, we will prove Theorem 1.1, a rigid property of τ -quasi-Einstein metrics with $\lambda \leq 0$ on complete noncompact Riemannian manifolds.

Proof. Consider a smooth function $\theta(t) : [0, +\infty) \rightarrow [0, 1]$:

$$(4-1) \quad \theta(t) = \begin{cases} 1 & \text{if } 0 \leq t \leq 1, \\ 0 & \text{if } t \geq 2, \end{cases}$$

so that

$$(4-2) \quad -10\sqrt{\theta} \leq \theta' \leq 0.$$

For $R_0 > 0$, let

$$\varphi(x) = \theta\left(\frac{r(x)}{R_0}\right)$$

be a cutoff function, where $r(x)$ is the distance function determined by $O \in M$. Then

$$0 \leq \varphi \leq 1, \quad |\nabla\varphi|(x) \leq \frac{C}{R_0}$$

and $\varphi(x) = 1$ on B_{R_0} , $\varphi(x) = 0$ outside of B_{2R_0} . Let

$$\alpha = -\frac{\tau+2}{\tau}.$$

Plugging (1-1) into (3-3) yields

$$(4-3) \quad -\nabla\Delta f \cdot \nabla f + \nabla|\nabla f|^2 \cdot \nabla f = 2\lambda|\nabla f|^2 + \frac{2}{\tau}|\nabla f|^4 - \frac{2}{\tau}\Delta f|\nabla f|^2.$$

Integrating (4-3) on M and using the fact that $\lambda \leq 0$, we obtain

$$(4-4) \quad -\int_M \nabla\Delta f \cdot \nabla f \varphi e^{\alpha f} dx + \int_M \nabla|\nabla f|^2 \cdot \nabla f \varphi e^{\alpha f} dx \\ \leq \frac{2}{\tau} \int_M |\nabla f|^4 \varphi e^{\alpha f} dx - \frac{2}{\tau} \int_M \Delta f |\nabla f|^2 \varphi e^{\alpha f} dx.$$

Integrating by parts yields

$$(4-5) \quad \int_M \nabla|\nabla f|^2 \cdot \nabla f \varphi e^{\alpha f} dx = - \int_M |\nabla f|^2 (\Delta f \varphi + \alpha |\nabla f|^2 \varphi + \nabla f \cdot \nabla \varphi) e^{\alpha f} dx$$

and

$$(4-6) \quad \int_M \nabla\Delta f \cdot \nabla f \varphi e^{\alpha f} dx = - \int_M ((\Delta f)^2 \varphi + \alpha \Delta f |\nabla f|^2 \varphi + \Delta f \nabla f \cdot \nabla \varphi) e^{\alpha f} dx.$$

Taking (4-5) and (4-6) into (4-4) yields

$$\int_M ((\Delta f)^2 - 2\Delta f |\nabla f|^2 + |\nabla f|^4) \varphi e^{\alpha f} dx \\ \leq - \int_M (\Delta f \nabla f \cdot \nabla \varphi - |\nabla f|^2 \nabla f \cdot \nabla \varphi) e^{\alpha f} dx \\ \leq \left(\int_M (\Delta f - |\nabla f|^2)^2 \varphi e^{\alpha f} dx \right)^{1/2} \left(\int_{B_{2R_0} \setminus B_{R_0}} \frac{|\nabla f \cdot \nabla \varphi|^2}{\varphi} e^{\alpha f} dx \right)^{1/2}.$$

Observing that

$$|\nabla f \cdot \nabla \varphi| \leq |\nabla f| |\nabla \varphi| \leq \frac{C}{R_0} |\nabla f|,$$

we get

$$\begin{aligned} \int_{B_{R_0}} (\Delta f - |\nabla f|^2)^2 e^{\alpha f} dx &\leq \int_M (\Delta f - |\nabla f|^2)^2 \varphi e^{\alpha f} dx \\ &\leq C R_0^{-2} \int_{B_{2R_0} \setminus B_{R_0}} |\nabla f|^2 e^{\alpha f} dx. \end{aligned}$$

Letting $R_0 \rightarrow \infty$, by (1-3), we conclude that

$$\int_M (\Delta f - |\nabla f|^2)^2 e^{\alpha f} dx = 0.$$

Hence $\Delta e^f = 0$.

When $\lambda < 0$, we deduce from $\Delta e^f = 0$ that $\Delta f = |\nabla f|^2$. Equation (4-3) is then equivalent to $2\lambda |\nabla f|^2 = 0$, which means that f is constant. \square

5. Lower bound of the scalar curvature

In this section, we will prove Theorem 1.4 and Theorem 1.6 by using the weighted Laplacian comparison theorem and the maximum principle. We first introduce the weighted Laplacian comparison theorem, which can be found in [Lott 2003; Wang 2010].

Lemma 5.1. *Let (M, g) be an n -dimensional complete Riemannian manifold, f a real value smooth function on M and $\Delta_f = \Delta - \nabla f \cdot \nabla$ the weighted Laplacian. Assume that the τ -Bakry-Émery Ricci curvature on M is bounded by*

$$\text{Ric}_{f,\tau} \geq \lambda$$

with constant λ and $r(x) = \text{dist}(O, x)$ is the distance function determined by a fixed point O . If a_λ is a solution to the Riccati equation

$$\frac{\partial a_\lambda}{\partial r} = \lambda - \frac{a_\lambda^2}{n + \tau - 1}, \quad \lim_{r \searrow 0} r a_\lambda = n + \tau - 1,$$

then at $y \notin \text{Cut}(O)$,

$$\Delta_f r \leq a_\lambda(r).$$

In particular, if $\lambda \leq 0$,

$$\Delta_f r \leq \frac{n + \tau - 1}{r} \left(1 + \sqrt{-\frac{\lambda}{n + \tau - 1}} r \right).$$

We need the following estimate, which can be proved by using the maximum principle [Pigola et al. 2005; Schoen and Yau 1994; Yau 1975; Cheng and Yau 1975].

Theorem 5.2. *Let M be an n -dimensional complete noncompact Riemannian manifold, g a τ -quasi-Einstein metric with potential function f and $\lambda \leq 0$ a constant. We also assume that $\tau > 1$ and $\mu > 0$. Then for $x > 0$,*

$$(5-1) \quad R(y) + 2xe^{2f/\tau(y)} \geq \frac{n(2n-2+\tau)+n\sqrt{\Delta}}{2(n+\tau-1)}\lambda$$

holds for any $y \in M$, where

$$(5-2) \quad \Delta = \tau^2 + \frac{8(n+\tau-1)(n-1)^2x}{2x(n+\tau-1)+n\mu}.$$

Proof. Consider a smooth function $\theta(t) : [0, +\infty) \rightarrow [0, 1]$,

$$\theta(t) = \begin{cases} 1 & \text{if } 0 \leq t \leq 1, \\ 0 & \text{if } t \geq 2, \end{cases}$$

so that

$$(5-3) \quad -10\theta^{1/2} \leq \theta' \leq 0, \theta'' \geq -10.$$

For a large enough constant $R_0 > 0$, define the smooth cutoff function $\varphi : M \rightarrow \mathbb{R}$ by

$$\varphi(x, t) = \theta\left(\frac{r(x)}{R_0}\right).$$

Then

$$(5-4) \quad \nabla\varphi = \frac{\theta'\nabla r}{R_0}.$$

By Lemma 5.1, we have that for $y \in B_{2R_0}$,

$$(5-5) \quad \begin{aligned} \Delta_f\varphi(y) &= \Delta\varphi - \nabla\varphi \cdot \nabla f = \frac{\theta''}{R_0^2} + \frac{\theta'\Delta_f r}{R_0} \\ &\geq \frac{\theta''}{R_0^2} + \frac{(n+\tau-1)\theta'(1+\sqrt{K}R_0)}{R_0^2}, \end{aligned}$$

where

$$K = -\frac{\lambda}{n+\tau-1} \geq 0.$$

Let

$$Q = \varphi G = \varphi(R + 2xe^{2f/\tau}).$$

If for any $R_0 > 0$ the minimal value of G on B_{R_0} is not smaller than zero, then Theorem 5.2 holds. Hence we can assume that for some large enough value of $R_0 > 0$, the minimal value of G on B_{R_0} is negative. If we assume that Q achieves its minimal value at x_0 on B_{2R_0} , then

$$Q(x_0) \leq \min_{x \in B_{R_0}} Q(x) = \min_{x \in B_{R_0}} G(x) < 0,$$

which means that x_0 is not on the boundary of B_{2R_0} . Hence $\varphi(x_0) > 0$ and

$$(5-6) \quad \nabla Q = 0,$$

$$(5-7) \quad \Delta_f Q \geq 0$$

hold at x_0 . By (3-6), (5-6) and (5-7), we get that, at x_0 ,

$$(5-8) \quad 0 \leq \frac{\Delta_f \varphi}{2} Q - \frac{|\nabla \varphi|^2}{\varphi} Q + \frac{|\nabla \varphi|^2}{4\epsilon \tau \varphi} Q^2 + \frac{4x(n+\tau-1)\varphi}{n\tau} Q e^{2f/\tau} - \frac{n+\tau-1}{n\tau} Q^2 \\ + \left(\frac{2n-2+\tau}{\tau} \varphi \lambda - \frac{\epsilon}{\tau-1} \right) Q + \varphi^2 A - \frac{n(n-1)\varphi^2}{\tau} \lambda^2 + \frac{n-\tau}{\tau-1} \varphi \epsilon \lambda.$$

Noticing that, at x_0 ,

$$(5-9) \quad \frac{4x(n+\tau-1)\varphi}{n\tau} Q e^{2f/\tau} \leq 0.$$

Taking (5-4), (5-5) and (5-9) into (5-8), and using (5-3), we get that, at x_0 ,

$$(5-10) \quad 0 \leq -\frac{105Q}{R_0^2} - \frac{5(n+\tau-1)(1+\sqrt{K}R_0)Q}{R_0^2} + \frac{25}{\epsilon \tau R_0^2} Q^2 - \frac{n+\tau-1}{n\tau} Q^2 \\ + \left(\frac{2n-2+\tau}{\tau} \varphi \lambda - \frac{\epsilon}{\tau-1} \right) Q + \varphi^2 A - \frac{n(n-1)\varphi^2}{\tau} \lambda^2 + \frac{n-\tau}{\tau-1} \varphi \epsilon \lambda.$$

By (3-14) and the fact that $\lambda \leq 0$, we have that, at x_0 ,

$$(5-11) \quad \varphi^2 A \leq B,$$

where

$$(5-12) \quad B = \frac{n\tau}{16x^2(n+\tau-1)+8nx\mu} \left(\frac{4x(n-1)}{\tau} \lambda - \frac{\epsilon(2x+\mu)}{\tau-1} \right)^2.$$

For a large enough value of $R_0 > 0$, define

$$\sigma(R_0) = \frac{\inf\{G(x) : x \in B(O, R_0)\}}{\inf\{G(x) : x \in B(O, 2R_0)\}}.$$

It is easy to see that

$$Q(x_0) = \varphi(x_0)G(x_0) \leq \inf\{G(x) : x \in B(O, R_0)\}$$

and

$$Q(x_0) = \varphi(x_0)G(x_0) \geq \varphi(x_0) \inf\{G(x) : x \in B(O, 2R_0)\}.$$

Using the assumption that $\inf\{G(x) : x \in B(O, R_0)\} < 0$, we get that

$$(5-13) \quad \sigma(R_0) \leq \varphi(x_0) \leq 1.$$

From (5-10), (5-11) and (5-13), it follows that, at x_0 ,

$$(5-14) \quad 0 \leq -\frac{105Q}{R_0^2} - \frac{5(n+\tau-1)(1+\sqrt{K}R_0)Q}{R_0^2} + \frac{25}{\epsilon\tau R_0^2}Q^2 - \frac{n+\tau-1}{n\tau}Q^2 \\ + \left(\frac{2n-2+\tau}{\tau}\sigma(R_0)\lambda - \frac{\epsilon}{\tau-1}\right)Q + B - \frac{n(n-1)\sigma^2(R_0)\lambda^2}{\tau} - \frac{|n-\tau|}{\tau-1}\epsilon\lambda.$$

Now, assume that R_0 is large enough so that

$$\frac{25}{\epsilon\tau R_0^2} < \frac{n+\tau-1}{n\tau}.$$

Then (5-14) gives us that, at x_0 ,

$$(5-15) \quad Q \geq \frac{-E(R_0) - \sqrt{E^2(R_0) + 4D(R_0)F(R_0)}}{2D(R_0)},$$

where

$$D(R_0) = \frac{n+\tau-1}{n\tau} - \frac{25}{\epsilon\tau R_0^2}, \\ E(R_0) = \frac{105 + 5(n+\tau-1)(1+\sqrt{K}R_0)}{R_0^2} - \frac{(2n-2+\tau)\sigma(R_0)\lambda}{\tau} + \frac{\epsilon}{\tau-1}, \\ F(R_0) = B - \frac{n(n-1)\sigma^2(R_0)\lambda^2}{\tau} - \frac{|n-\tau|}{\tau-1}\epsilon\lambda.$$

Hence, for all $y \in B_{R_0}$,

$$(5-16) \quad G(y) = \varphi(y)G(y) \geq Q(x_0) \geq \frac{-E(R_0) - \sqrt{E^2(R_0) + 4D(R_0)F(R_0)}}{2D(R_0)}.$$

Noting that $0 \leq \sigma(R_0) \leq 1$ and B is a constant independent of R_0 , we deduce from (5-16) that for a large enough value of R_0 , G is bounded from below by a constant independent of R_0 . Hence

$$\lim_{R_0 \rightarrow \infty} \sigma(R_0) = 1,$$

which means that

$$D = \lim_{R_0 \rightarrow \infty} D(R_0) = \frac{n+\tau-1}{n\tau}, \\ E = \lim_{R_0 \rightarrow \infty} E(R_0) = -\frac{(2n-2+\tau)\lambda}{\tau} + \frac{\epsilon}{\tau-1}, \\ F = \lim_{R_0 \rightarrow \infty} F(R_0) = B - \frac{n(n-1)\lambda^2}{\tau} - \frac{|n-\tau|}{\tau-1}\epsilon\lambda.$$

By (5-16), we obtain that, for all $y \in M, x > 0$ and $\epsilon > 0$,

$$(5-17) \quad G(y) \geq \frac{-E - \sqrt{E^2 + 4DF}}{2D}.$$

Note that

$$\lim_{\epsilon \searrow 0} E = -\frac{(2n-2+\tau)\lambda}{\tau}, \quad \lim_{\epsilon \searrow 0} F = \frac{2xn(n-1)^2\lambda^2}{(2x(n+\tau-1)+n\mu)\tau} - \frac{n(n-1)\lambda^2}{\tau}.$$

Letting $\epsilon \searrow 0$ in (5-17) leads to (5-1). □

The following result is useful.

Theorem 5.3. *If $\lambda < 0$ and $\mu < 0$, then for all $y \in M$,*

$$(5-18) \quad f(y) \leq \frac{\tau}{2} \ln \frac{\tau\lambda}{\mu}.$$

Proof. Let φ be the cutoff function defined in the proof of Theorem 5.2 and

$$H = \varphi e^{2f/\tau}.$$

Noting that

$$\nabla H = \nabla \varphi e^{2f/\tau} + \varphi \nabla e^{2f/\tau},$$

by (3-4) and (3-9), we have that

$$\Delta_f e^{2f/\tau} \geq \frac{2}{\tau} \Delta_f f e^{2f/\tau} = -\frac{2\mu}{\tau} e^{4f/\tau} + 2\lambda e^{2f/\tau}.$$

Hence

$$(5-19) \quad \begin{aligned} \Delta_f H &= \varphi \Delta_f e^{2f/\tau} + e^{2f/\tau} \Delta_f \varphi + 2\nabla \varphi \cdot \nabla e^{2f/\tau} \\ &\geq -\frac{2\mu\varphi}{\tau} e^{4f/\tau} + 2\lambda\varphi e^{2f/\tau} + \frac{\Delta_f \varphi}{\varphi} H + 2\frac{\nabla \varphi \cdot \nabla H}{\varphi} - 2\frac{|\nabla \varphi|^2}{\varphi^2} H \end{aligned}$$

holds at $y \in M$ when $\varphi(y) > 0$. We assume that H achieves its maximum at x_0 on B_{2R_0} . If $\varphi(x_0) = 0$, then for all $x \in B_{R_0}$,

$$e^{2f/\tau(x)} = \varphi(x)e^{2f/\tau(x)} = H(x) \leq H(x_0) = 0,$$

which is impossible, so $\varphi(x_0) > 0$. Noting that

$$\Delta_f H \leq 0, \quad \nabla H = 0$$

hold at x_0 , by (5-3), (5-4), (5-5) and (5-19), we get that, at x_0 ,

$$\begin{aligned} 0 &\geq -\frac{2\mu\varphi}{\tau}e^{4f/\tau} + 2\lambda\varphi e^{2f/\tau} + \frac{\Delta f\varphi}{\varphi}H - \frac{2|\nabla\varphi|^2}{\varphi^2}H \\ &\geq -\frac{2\mu}{\tau}H^2 + 2\lambda H + \frac{\theta'' + (n + \tau - 1)(1 + \sqrt{K}R_0)\theta'}{R_0^2\varphi}H - \frac{2|\theta'|^2}{R_0^2\varphi^2}H \\ &\geq -\frac{2\mu}{\tau}H^2 + 2\lambda H - \frac{210 + 10(n + \tau - 1)(1 + \sqrt{K}R_0)\sqrt{\varphi}}{R_0^2\varphi}H. \end{aligned}$$

By using the fact that $0 < \varphi(x_0) \leq 1$, we have that, at x_0 ,

$$\begin{aligned} (5-20) \quad -\frac{2\mu}{\tau}H^2 &\leq -2\lambda\varphi H + \frac{210 + 10(n + \tau - 1)(1 + \sqrt{K}R_0)\sqrt{\varphi}}{R_0^2}H \\ &\leq -2\lambda H + \frac{210 + 10(n + \tau - 1)(1 + \sqrt{K}R_0)}{R_0^2}H \end{aligned}$$

or

$$H(x_0) \leq \frac{\tau\lambda}{\mu} - \frac{105\tau + 5\tau(n + \tau - 1)(1 + \sqrt{K}R_0)}{\mu R_0^2}.$$

Hence, for all $y \in B_{R_0}$,

$$\begin{aligned} e^{2f/\tau(y)} &= \varphi(y)e^{2f/\tau(y)} = H(y) \leq H(x_0)m \\ &\leq \frac{\tau\lambda}{\mu} - \frac{105\tau + 5\tau(n + \tau - 1)(1 + \sqrt{K}R_0)}{\mu R_0^2}. \end{aligned}$$

Letting $R_0 \rightarrow \infty$ yields (5-18). □

Remark 5.4. Equation (1-1) still holds if we shift the function f by a constant. However, from Equation (3-2), the constant μ will change after this shift.

Proof of Theorem 1.4. When $\tau \geq 1$ and $\mu \leq 0$, (1-4) follows from Theorem 1.3. When $\tau > 1$ and $\mu > 0$, (1-4) follows by letting $x \searrow 0$ in (5-1). We only need to consider the case that $0 < \tau \leq 1$. Now if $\mu \geq 0$, (3-2) tells us that

$$R = \mu e^{2f/\tau} + (n - \tau)\lambda + \frac{1 - \tau}{\tau}|\nabla f|^2 \geq (n - \tau)\lambda \geq n\lambda.$$

If $\mu < 0$ and $\lambda < 0$, by Theorem 5.3, we have

$$R \geq \mu e^{2f/\tau} + (n - \tau)\lambda \geq n\lambda.$$

Hence (1-4) follows. If $\mu < 0$ and $\lambda = 0$, [Wang 2011, Theorem 3.2] tells us that f is constant, and Theorem 1.4 follows. □

Proof of Theorem 1.6. When $\mu \leq 0$, (3-2) tells us that

$$R \leq (n - \tau)\lambda + \frac{1 - \tau}{\tau} |\nabla f|^2.$$

If $\tau \geq 1$, then $R \leq (n - \tau)\lambda$. If $0 < \tau < 1$, by the gradient estimate in [Wang 2011], we have $|\nabla f|^2 \leq -\tau\lambda$. Hence $R \leq (n - 1)\lambda$. \square

Acknowledgments

The author thanks the referee for valuable comments and suggestions. The author is also grateful to Professor Shengliang Pan for his constant encouragement.

References

- [Besse 1987] A. L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)] **10**, Springer, Berlin, 1987. MR 88f:53087 Zbl 0613.53001
- [Cao 2009] H. Cao, “Geometry of complete gradient shrinking solitons”, preprint, 2009. arXiv 0903.3927
- [Cao and Zhou 2010] H.-D. Cao and D. Zhou, “On complete gradient shrinking Ricci solitons”, *J. Differential Geom.* **85**:2 (2010), 175–185. MR 2011k:53040 Zbl 05835154
- [Cao and Zhu 2006] H.-D. Cao and X.-P. Zhu, “A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow”, *Asian J. Math.* **10**:2 (2006), 165–492. MR 2008d:53090 Zbl 1200.53057
- [Case 2010] J. S. Case, “The nonexistence of quasi-Einstein metrics”, *Pacific J. Math.* **248**:2 (2010), 277–284. MR 2011m:53053 Zbl 1204.53032
- [Case et al. 2011] J. Case, Y.-J. Shu, and G. Wei, “Rigidity of quasi-Einstein metrics”, *Differential Geom. Appl.* **29**:1 (2011), 93–100. MR 2784291 Zbl 1215.53033
- [Chen 2009] B.-L. Chen, “Strong uniqueness of the Ricci flow”, *J. Differential Geom.* **82**:2 (2009), 363–382. MR 2010h:53095 Zbl 1177.53036
- [Cheng and Yau 1975] S. Y. Cheng and S. T. Yau, “Differential equations on Riemannian manifolds and their geometric applications”, *Comm. Pure Appl. Math.* **28**:3 (1975), 333–354. MR 52 #6608 Zbl 0312.53031
- [Hamilton 1995] R. S. Hamilton, “The formation of singularities in the Ricci flow”, pp. 7–136 in *Surveys in differential geometry, Vol. II* (Cambridge, MA, 1993), edited by S. Yau and C. Hsiung, Int. Press, 1995. MR 97e:53075 Zbl 0867.53030
- [Ivey 1993] T. Ivey, “Ricci solitons on compact three-manifolds”, *Differential Geom. Appl.* **3**:4 (1993), 301–307. MR 94j:53048 Zbl 0788.53034
- [Kim and Kim 2003] D.-S. Kim and Y. H. Kim, “Compact Einstein warped product spaces with nonpositive scalar curvature”, *Proc. Amer. Math. Soc.* **131**:8 (2003), 2573–2576. MR 2004b:53063 Zbl 1029.53027
- [Li 2005] X.-D. Li, “Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds”, *J. Math. Pures Appl.* (9) **84**:10 (2005), 1295–1361. MR 2006f:58046 Zbl 1082.58036
- [Lott 2003] J. Lott, “Some geometric properties of the Bakry–Émery–Ricci tensor”, *Comment. Math. Helv.* **78**:4 (2003), 865–883. MR 2004i:53044 Zbl 1038.53041

- [Lü et al. 2004] H. Lü, D. N. Page, and C. N. Pope, “New inhomogeneous Einstein metrics on sphere bundles over Einstein–Kähler manifolds”, *Phys. Lett. B* **593**:1-4 (2004), 218–226. MR 2005f:53063
- [Munteanu 2009] O. Munteanu, “The volume growth of complete gradient shrinking Ricci solitons”, preprint, 2009. arXiv 0904.0798
- [O’Neill 1983] B. O’Neill, *Semi-Riemannian geometry*, Pure and Applied Mathematics **103**, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1983. With applications to relativity. MR 85f:53002 Zbl 0531.53051
- [Perelman 2002] G. Perelman, “The entropy formula for the Ricci flow and its geometric applications”, preprint, 2002. Zbl 1130.53001 arXiv math/0211159
- [Pigola et al. 2005] S. Pigola, M. Rigoli, and A. G. Setti, “Maximum principles on Riemannian manifolds and applications”, *Mem. Amer. Math. Soc.* **174**:822 (2005), x+99. MR 2006b:53048
- [Qian 1997] Z. Qian, “Estimates for weighted volumes and applications”, *Quart. J. Math. Oxford Ser. (2)* **48**:190 (1997), 235–242. MR 98e:53058 Zbl 0902.53032
- [Schoen and Yau 1994] R. Schoen and S.-T. Yau, *Lectures on differential geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology **1**, International Press, Cambridge, MA, 1994. MR 97d:53001 Zbl 0830.53001
- [Wang 2010] L.-F. Wang, “The upper bound of the L^2_μ spectrum”, *Ann. Global Anal. Geom.* **37**:4 (2010), 393–402. MR 2011d:53067 Zbl 1190.53034
- [Wang 2011] L. F. Wang, “Rigid properties of quasi-Einstein metrics”, *Proc. Amer. Math. Soc.* **139**:10 (2011), 3679–3689. MR 2813397 Zbl 05959506
- [Wei and Wylie 2007] G. Wei and W. Wylie, “Comparison geometry for the smooth metric measure spaces”, pp. 191–202 in *Proceedings of the fourth ICCM, Vol. 2* (Hangzhou, China), edited by S. Yau, 2007.
- [Yau 1975] S. T. Yau, “Harmonic functions on complete Riemannian manifolds”, *Comm. Pure Appl. Math.* **28** (1975), 201–228. MR 55 #4042 Zbl 0291.31002
- [Zhang 2006] X. Zhang, “Compactness theorems for gradient Ricci solitons”, *J. Geom. Phys.* **56**:12 (2006), 2481–2499. MR 2008b:53049 Zbl 1107.53045
- [Zhang 2009] Z.-H. Zhang, “On the completeness of gradient Ricci solitons”, *Proc. Amer. Math. Soc.* **137**:8 (2009), 2755–2759. MR 2010a:53057 Zbl 1176.53046
- [Zhang 2011] S. J. Zhang, “On a sharp volume estimate for gradient Ricci solitons with scalar curvature bounded below”, *Acta Math. Sin. (Engl. Ser.)* **27**:5 (2011), 871–882. MR 2786449 Zbl 1219.53039

Received January 10, 2011. Revised November 7, 2011.

LIN FENG WANG
SCHOOL OF SCIENCE
NANTONG UNIVERSITY
NANTONG 226007, JIANGSU
CHINA
wlf711178@126.com

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 254 No. 2 December 2011

Curvatures of spheres in Hilbert geometry	257
ALEXANDER BORISENKO and EUGENE OLIN	
A formula equating open and closed Gromov–Witten invariants and its applications to mirror symmetry	275
KWOKWAI CHAN	
A note on p -harmonic l -forms on complete manifolds	295
LIANG-CHU CHANG and CHIUNG-JUE ANNA SUNG	
The Cheeger constant of curved strips	309
DAVID KREJČIŘÍK and ALDO PRATELLI	
Structure of solutions of 3D axisymmetric Navier–Stokes equations near maximal points	335
ZHEN LEI and QI S. ZHANG	
Local comparison theorems for Kähler manifolds	345
GANG LIU	
Structurable algebras of skew-rank 1 over the affine plane	361
SUSANNE PUMPLÜN	
An analogue of Krein’s theorem for semisimple Lie groups	381
SANJOY PUSTI	
Une remarque de dynamique sur les variétés semi-abéliennes	397
GAËL RÉMOND	
Fourier transforms of semisimple orbital integrals on the Lie algebra of SL_2	407
LOREN SPICE	
On noncompact τ -quasi-Einstein metrics	449
LIN FENG WANG	
Decomposition of de Rham complexes with smooth horizontal coefficients for semistable reductions	465
QIHONG XIE	
A differentiable sphere theorem inspired by rigidity of minimal submanifolds	499
HONG-WEI XU and LING TIAN	



0030-8730(201112)254:2;1-A