

*Pacific
Journal of
Mathematics*

**A DIFFERENTIABLE SPHERE THEOREM
INSPIRED BY RIGIDITY OF MINIMAL SUBMANIFOLDS**

HONG-WEI XU AND LING TIAN

Volume 254 No. 2

December 2011

A DIFFERENTIABLE SPHERE THEOREM INSPIRED BY RIGIDITY OF MINIMAL SUBMANIFOLDS

HONG-WEI XU AND LING TIAN

We prove a vanishing theorem for the fundamental group of a compact submanifold in a space form, and then present a refined version of Ejiri's rigidity theorem for minimal submanifolds in a sphere. Inspired by the refined Ejiri theorem, we verify a new differentiable sphere theorem for compact submanifolds in space forms. We also show that our differentiable sphere theorem is sharp. We emphasize that our method of Ricci flow in the proof of the sphere theorem seems useful in the study of curvature and topology. Also, we obtain a differentiable pinching theorem for compact submanifolds in a Riemannian manifold.

1. Introduction

The investigation of geometrical and topological structures of manifolds plays an important role in global differential geometry. Since the first sphere theorem for Riemannian manifolds was proved by Rauch in 1951, there has been much progress in this field; see [Berger 2000; Brendle 2010; Shiohama 2000; Xu 2012]. Brendle and Schoen [2008] proved a remarkable classification theorem for compact manifolds with weakly $\frac{1}{4}$ -pinched curvatures in the pointwise sense, implying this:

Theorem A. *Suppose that M is an n -dimensional complete and simply connected Riemannian manifold such that $\frac{1}{4} \leq K_M \leq 1$. Then M is either diffeomorphic to S^n or isometric to a compact rank-one symmetric space (CROSS).*

Petersen and Tao [2009] improved Brendle and Schoen's pinching constant in Theorem A to $\frac{1}{4} - \varepsilon_n$, with ε_n being a positive constant depending only on n . Motivated by Shen's topological sphere theorem [1989] for compact manifolds with positive Ricci curvature and Yau's open problem on the pinching theorem [Yau 1993, Problem 12], Gu and Xu [2011] obtained a differentiable sphere theorem for compact manifolds with positive scalar curvature. In particular, they proved that

Research supported by the NSFC, Grant No. 11071211; the Trans-Century Training Programme Foundation for Talents by the Ministry of Education of China.

MSC2010: 53C20, 53C24, 53C40.

Keywords: Submanifolds, differentiable sphere theorem, Ricci curvature, Ricci flow, stable currents.

if M is an n -dimensional compact Riemannian manifold ($n \geq 3$) whose sectional curvature and scalar curvature satisfy $K_M \leq 1$ and $R > n(n-1)\delta_n$, then M is diffeomorphic to a spherical space form. Here

$$\delta_n = 1 - \frac{20 - 8 \operatorname{sgn}(n-3)}{5n(n-1)}$$

and $\operatorname{sgn}(\cdot)$ is the standard sign function.

In addition, results on sphere theorems for Riemannian submanifolds have been obtained by various authors [Huisken 1987; Lawson and Simons 1973; Shiohama 2000; Shiohama and Xu 1997; Xu and Zhao 2009]. Let M^n be an n -dimensional submanifold in an $(n+p)$ -dimensional Riemannian manifold N^{n+p} . Denote by H and S the mean curvature and the squared length of the second fundamental form of M , respectively. Motivated by Lawson, Simons, Shiohama, and Xu's topological sphere theorem [Lawson and Simons 1973; Shiohama and Xu 1997] for submanifolds, Xu and Zhao [2009] proved that for $n \geq 4$, if M is an n -dimensional complete submanifold in S^{n+p} and $S < 2\sqrt{2}$, then M is diffeomorphic to S^n . Let $F^{n+p}(c)$ be an $(n+p)$ -dimensional simply connected space form with constant curvature c . Xu and Gu [2010] extended Huisken's sphere theorem [1987] for hypersurfaces in spheres to the case of submanifolds in space forms, and proved the following optimal differentiable sphere theorem:

Theorem B. *Suppose that M is an n -dimensional oriented complete submanifold in $F^{n+p}(c)$ with $c \geq 0$. If*

$$\lambda(M) := \sup_M \left(S - \frac{n^2 H^2}{n-1} - 2c \right) < 0,$$

then M is diffeomorphic to S^n .

After the pioneering rigidity theorem for closed minimal submanifolds in a sphere due to Simons [1968], several fundamental rigidity results for minimal submanifolds were proved by various authors [Chern et al. 1970; Ejiri 1979; Lawson 1969; An-Min and Jimin 1992; Yau 1974; 1975]. Ejiri [1979] obtained the following rigidity theorem for minimal submanifolds in spheres:

Theorem C. *Suppose that M is an n -dimensional simply connected compact orientable minimal submanifold in an $(n+p)$ -dimensional unit sphere S^{n+p} , with $n \geq 4$. If the Ricci curvature of M satisfies*

$$\operatorname{Ric}_M \geq n - 2,$$

then M is either the totally geodesic submanifold S^n , the Clifford torus $S^m(\sqrt{1/2}) \times S^m(\sqrt{1/2})$ in S^{n+1} with $n = 2m$, or $\mathbb{C}P^2(\frac{4}{3})$ in S^7 . Here $\mathbb{C}P^2(\frac{4}{3})$ denotes the 2-dimensional complex projective space minimally immersed into S^7 with constant holomorphic sectional curvature $\frac{4}{3}$.

Shen [1992] proved that if M is a 3-dimensional compact minimal submanifold in a unit sphere S^{3+p} , and if the Ricci curvature of M is larger than 1, then M is the totally geodesic submanifold S^3 .

Here we investigate differentiable sphere theorems for compact submanifolds of positive Ricci curvature. Using convergence results of Hamilton [1982] and Brendle [2008] for Ricci flow and the Lawson–Simons–Xin formula for the nonexistence of stable currents [Lawson and Simons 1973; Xin 1984], we prove such a theorem inspired by rigidity of minimal submanifolds. We first prove the following differentiable sphere theorem for submanifolds in a Riemannian manifold.

Theorem 1.1. *Suppose that M is an n -dimensional compact submanifold in an $(n + p)$ -dimensional Riemannian manifold N^{n+p} , with $n \geq 4$. Denote by $\bar{K}(x, \pi)$ the sectional curvature of N for tangent 2-plane $\pi(\subset T_x N)$ at point $x \in N$. Set $\bar{K}_{\max}(x) := \max_{\pi \subset T_x N} \bar{K}(x, \pi)$ and $\bar{K}_{\min}(x) := \min_{\pi \subset T_x N} \bar{K}(x, \pi)$. If*

$$\text{Ric}_M > (n - \frac{2}{3})\bar{K}_{\max} - \frac{4}{3}\bar{K}_{\min} + \frac{1}{8}n^2 H^2,$$

then the normalized Ricci flow with initial metric g_0

$$\frac{\partial}{\partial t} g(t) = -2 \text{Ric}_{g(t)} + \frac{2}{n} r_{g(t)} g(t)$$

exists for all time and converges to a constant curvature metric as $t \rightarrow \infty$. Also, M is diffeomorphic to a spherical space form. In particular, if M is simply connected, then M is diffeomorphic to S^n .

We give a vanishing theorem for the fundamental group of a submanifold.

Theorem 1.2. *Suppose that M is an n -dimensional compact submanifold in an $(n + p)$ -dimensional space form $F^{n+p}(c)$ with $c \geq 0$. If the Ricci curvature of M satisfies*

$$\text{Ric}_M > \frac{1}{2}(n - 1)c + \frac{1}{8}n^2 H^2,$$

then $\pi_1(M) = 0$; that is, M is simply connected.

Applying Theorem 1.2 to Theorem C, we obtain a refined version of the Ejiri rigidity theorem for minimal submanifolds in spheres, without the assumption that M is simply connected. Finally, inspired by the refined Ejiri rigidity theorem for minimal submanifolds, we prove the following differentiable sphere theorem for submanifolds in space forms.

Theorem 1.3 (Main Theorem). *Suppose that M is an n -dimensional compact submanifold in an $(n + p)$ -dimensional space form $F^{n+p}(c)$, with $c \geq 0$ and $n \geq 3$. If the Ricci curvature of M satisfies*

$$\text{Ric}_M > (n - 2)c + \frac{1}{8}n^2 H^2,$$

then M is diffeomorphic to S^n .

We show in Example 3.4 that the pinching condition in Theorem 1.3 is sharp. Our method in the proof of the main theorem seems very useful in the study of curvature and topology.

2. Notation and fundamental tools

Throughout, let M^n (where $n \geq 2$) be an n -dimensional compact submanifold in an $(n + p)$ -dimensional Riemannian manifold N^{n+p} . We use the following conventions on the range of indices:

$$1 \leq A, B, C, \dots \leq n + p, \quad 1 \leq i, j, k, \dots \leq n, \quad n + 1 \leq \alpha, \beta, \gamma, \dots \leq n + p.$$

For an arbitrary fixed point $x \in M \subset N$, we choose an orthonormal local frame field $\{e_A\}$ in N^{n+p} such that e_i 's are tangent to M . Let $\{\omega_A\}$ be the dual frame field of $\{e_A\}$. Let Rm, h and ξ denote the Riemannian curvature tensor, second fundamental form and mean curvature vector of M , respectively, and let \bar{Rm} be the Riemannian curvature tensor of N . Then

$$\begin{aligned} Rm &= \sum_{i,j,k,l} R_{ijkl} \omega_i \otimes \omega_j \otimes \omega_k \otimes \omega_l, \\ \bar{Rm} &= \sum_{A,B,C,D} \bar{R}_{ABCD} \omega_A \otimes \omega_B \otimes \omega_C \otimes \omega_D, \\ h &= \sum_{\alpha,i,j} h_{ij}^\alpha \omega_i \otimes \omega_j \otimes e_\alpha, \quad \xi = \frac{1}{n} \sum_{\alpha,i} h_{ii}^\alpha e_\alpha, \\ (2-1) \quad R_{ijkl} &= \bar{R}_{ijkl} + \sum_{\alpha} (h_{ik}^\alpha h_{jl}^\alpha - h_{il}^\alpha h_{jk}^\alpha), \end{aligned}$$

and the mean curvature H of M is given by $H := |(1/n) \sum_{\alpha,i} h_{ii}^\alpha e_\alpha|$. Denote by $\text{Ric}(u)$ the Ricci curvature of M in the direction of $u \in UM$. From the Gauss equation (2-1), we have

$$\text{Ric}(e_i) = \sum_j \bar{R}_{ijij} + \sum_{\alpha,j} (h_{ii}^\alpha h_{jj}^\alpha - h_{ij}^\alpha h_{ij}^\alpha).$$

Denote by A_α the shape operator of M with respect to e_α . Choose $\{e_\alpha\}$ such that e_{n+1} is parallel to ξ and

$$(2-2) \quad \text{tr } A_{n+1} = nH, \quad \text{tr } A_\alpha = 0, \quad \alpha \neq n + 1.$$

Thus the mean curvature vector ξ is equal to He_{n+1} , and

$$\sum_j h_{jj}^\alpha = \begin{cases} nH, & \alpha = n + 1, \\ 0, & \alpha \neq n + 1. \end{cases}$$

Hence

$$(2-3) \quad \text{Ric}(e_i) = \sum_j \bar{R}_{ijij} + nHh_{ii}^{n+1} - \sum_{\alpha,j} h_{ij}^\alpha h_{ij}^\alpha.$$

Denote by $K(x, \pi)$ the sectional curvature of M for tangent 2-plane $\pi(\subset T_x M)$ at point $x \in M$, and by $\bar{K}(x, \pi)$ the sectional curvature of N for tangent 2-plane $\pi(\subset T_x N)$ at point $x \in N$. Set

$$\bar{K}_{\max}(x) := \max_{\pi \subset T_x N} \bar{K}(x, \pi) \quad \text{and} \quad \bar{K}_{\min}(x) := \min_{\pi \subset T_x N} \bar{K}(x, \pi).$$

By Berger’s inequality [Brendle 2010, Proposition 1.9],

$$(2-4) \quad |\bar{R}_{ABCD}| \leq \frac{2}{3}(\bar{K}_{\max} - \bar{K}_{\min})$$

for all distinct indices A, B, C, D .

Theorem 2.1 [Hamilton 1982]. *Let (M, g_0) be a compact Riemannian manifold of dimension 3. If $\text{Ric}_M > 0$, then the normalized Ricci flow with initial metric g_0*

$$\frac{\partial}{\partial t} g(t) = -2 \text{Ric}_{g(t)} + \frac{2}{3} r_{g(t)} g(t)$$

exists for all time and converges to a constant curvature metric as $t \rightarrow \infty$. Here $r_{g(t)}$ denotes the mean value of the scalar curvature of $g(t)$.

Next we quote an important convergence result for the Ricci flow in higher dimensions.

Theorem 2.2 [Brendle 2008; Brendle and Schoen 2009]. *Suppose (M, g_0) is a compact Riemannian manifold of dimension $n \geq 4$. Assume that*

$$(2-5) \quad R_{1313} + \lambda^2 R_{1414} + R_{2323} + \lambda^2 R_{2424} - 2\lambda R_{1234} > 0$$

for all orthonormal four-frames $\{e_1, e_2, e_3, e_4\}$ and all $\lambda \in [-1, 1]$. Then the normalized Ricci flow with initial metric g_0

$$\frac{\partial}{\partial t} g(t) = -2 \text{Ric}_{g(t)} + \frac{2}{n} r_{g(t)} g(t)$$

exists for all time and converges to a constant curvature metric as $t \rightarrow \infty$. Here $r_{g(t)}$ denotes the mean value of the scalar curvature of $g(t)$.

The following nonexistence theorem for stable currents in a compact Riemannian manifold M isometrically immersed into $F^{n+p}(c)$ was proved for $c > 0$ in [Lawson and Simons 1973] and was extended to $c = 0$ in [Xin 1984]. It is used to eliminate the homology groups $H_q(M; \mathbb{Z})$ for $1 \leq q \leq n - 1$ and the fundamental group $\pi_1(M)$.

Theorem 2.3. *Let M be an n -dimensional compact submanifold in $F^{n+p}(c)$ with $c \geq 0$. Assume that*

$$(2-6) \quad \sum_{k=q+1}^n \sum_{i=1}^q [2|h(e_i, e_k)|^2 - \langle h(e_i, e_i), h(e_k, e_k) \rangle] < q(n-q)c$$

for any orthonormal basis $\{e_i\}$ of $T_x M$ at any point $x \in M$, where q is an integer satisfying $1 \leq q \leq n-1$. Then there do not exist any stable q -currents. Also,

$$H_q(M; \mathbb{Z}) = H_{n-q}(M; \mathbb{Z}) = 0,$$

where $H_i(M; \mathbb{Z})$ is the i -th homology group of M with integer coefficients.

3. Proofs of the theorems

To complete the proof of the main theorem (Theorem 1.3), we need to prove the differentiable pinching theorem for submanifolds (Theorem 1.1) and the vanishing theorem for the fundamental group of a submanifold (Theorem 1.2).

Proof of Theorem 1.1. It is sufficient to show that inequality (2-5) in Theorem 2.2 holds for all $\lambda \in \mathbb{R}$. Suppose that $\{e_1, e_2, e_3, e_4\}$ is an orthonormal four-frame and that $\lambda \in \mathbb{R}$. From the Gauss equation, we have

$$(3-1) \quad \begin{aligned} & R_{1313} + \lambda^2 R_{1414} + R_{2323} + \lambda^2 R_{2424} - 2\lambda R_{1234} \\ &= \bar{R}_{1313} + \lambda^2 \bar{R}_{1414} + \bar{R}_{2323} + \lambda^2 \bar{R}_{2424} - 2\lambda \bar{R}_{1234} \\ &\quad + \sum_{\alpha} (h_{11}^{\alpha} h_{33}^{\alpha} - (h_{13}^{\alpha})^2 + h_{22}^{\alpha} h_{33}^{\alpha} - (h_{23}^{\alpha})^2) \\ &\quad + \lambda^2 \sum_{\alpha} (h_{11}^{\alpha} h_{44}^{\alpha} - (h_{14}^{\alpha})^2 + h_{22}^{\alpha} h_{44}^{\alpha} - (h_{24}^{\alpha})^2) \\ &\quad - 2\lambda \sum_{\alpha} (h_{13}^{\alpha} h_{24}^{\alpha} - h_{14}^{\alpha} h_{23}^{\alpha}) \\ &= \bar{R}_{1313} + \lambda^2 \bar{R}_{1414} + \bar{R}_{2323} + \lambda^2 \bar{R}_{2424} - 2\lambda \bar{R}_{1234} \\ &\quad + \sum_{\alpha} (- (h_{13}^{\alpha})^2 - (h_{23}^{\alpha})^2 - \lambda^2 (h_{14}^{\alpha})^2 - \lambda^2 (h_{24}^{\alpha})^2 \\ &\quad - 2\lambda h_{13}^{\alpha} h_{24}^{\alpha} + 2\lambda h_{14}^{\alpha} h_{23}^{\alpha} + h_{11}^{\alpha} h_{33}^{\alpha} + h_{22}^{\alpha} h_{33}^{\alpha} + \lambda^2 h_{11}^{\alpha} h_{44}^{\alpha} + \lambda^2 h_{22}^{\alpha} h_{44}^{\alpha}). \end{aligned}$$

It follows from Berger's inequality that $\bar{R}_{1234} \leq \frac{2}{3}(\bar{K}_{\max} - \bar{K}_{\min})$. This together with (3-1) implies

$$(3-2) \quad \begin{aligned} & R_{1313} + \lambda^2 R_{1414} + R_{2323} + \lambda^2 R_{2424} - 2\lambda R_{1234} \\ &\geq 2(1 + \lambda^2) \bar{K}_{\min} - \frac{4}{3} |\lambda| (\bar{K}_{\max} - \bar{K}_{\min}) \\ &\quad + \sum_{\alpha} (- (h_{13}^{\alpha})^2 - (h_{23}^{\alpha})^2 - \lambda^2 (h_{14}^{\alpha})^2 - \lambda^2 (h_{24}^{\alpha})^2 \\ &\quad - |\lambda| (h_{13}^{\alpha})^2 - |\lambda| (h_{24}^{\alpha})^2 - |\lambda| (h_{23}^{\alpha})^2 - |\lambda| (h_{14}^{\alpha})^2 \\ &\quad - \frac{1}{2} (1 + \lambda^2) (h_{11}^{\alpha})^2 - \frac{1}{2} (1 + \lambda^2) (h_{22}^{\alpha})^2 - (h_{33}^{\alpha})^2 - \lambda^2 (h_{44}^{\alpha})^2 \\ &\quad + \frac{1}{2} (h_{11}^{\alpha} + h_{33}^{\alpha})^2 + \frac{1}{2} (h_{22}^{\alpha} + h_{33}^{\alpha})^2 + \frac{1}{2} \lambda^2 (h_{11}^{\alpha} + h_{44}^{\alpha})^2 + \frac{1}{2} \lambda^2 (h_{22}^{\alpha} + h_{44}^{\alpha})^2) \end{aligned}$$

$$\begin{aligned}
 &\geq 2(1 + \lambda^2)\bar{K}_{\min} - \frac{2}{3}(1 + \lambda^2)(\bar{K}_{\max} - \bar{K}_{\min}) \\
 &\quad + \sum_{\alpha} \left(-(h_{31}^{\alpha})^2 - (h_{32}^{\alpha})^2 - (h_{33}^{\alpha})^2 - \lambda^2(h_{41}^{\alpha})^2 - \lambda^2(h_{42}^{\alpha})^2 - \lambda^2(h_{44}^{\alpha})^2 \right. \\
 &\quad \quad - \frac{1}{2}(1 + \lambda^2)(h_{11}^{\alpha})^2 - |\lambda|(h_{13}^{\alpha})^2 - |\lambda|(h_{14}^{\alpha})^2 \\
 &\quad \quad - \frac{1}{2}(1 + \lambda^2)(h_{22}^{\alpha})^2 - |\lambda|(h_{23}^{\alpha})^2 - |\lambda|(h_{24}^{\alpha})^2 \\
 &\quad \quad \left. + \frac{1}{2}(h_{11}^{\alpha} + h_{33}^{\alpha})^2 + \frac{1}{2}(h_{22}^{\alpha} + h_{33}^{\alpha})^2 + \frac{1}{2}\lambda^2(h_{11}^{\alpha} + h_{44}^{\alpha})^2 + \frac{1}{2}\lambda^2(h_{22}^{\alpha} + h_{44}^{\alpha})^2 \right) \\
 &\geq 2(1 + \lambda^2)\bar{K}_{\min} - \frac{2}{3}(1 + \lambda^2)(\bar{K}_{\max} - \bar{K}_{\min}) \\
 &\quad + \sum_{\alpha} \left(-\sum_j (h_{3j}^{\alpha})^2 + \lambda^2(h_{4j}^{\alpha})^2 + \frac{1}{2}(1 + \lambda^2)(h_{1j}^{\alpha})^2 + \frac{1}{2}(1 + \lambda^2)(h_{2j}^{\alpha})^2 \right. \\
 &\quad \quad \left. + \frac{1}{2}(h_{11}^{\alpha} + h_{33}^{\alpha})^2 + \frac{1}{2}(h_{22}^{\alpha} + h_{33}^{\alpha})^2 + \frac{1}{2}\lambda^2(h_{11}^{\alpha} + h_{44}^{\alpha})^2 + \frac{1}{2}\lambda^2(h_{22}^{\alpha} + h_{44}^{\alpha})^2 \right).
 \end{aligned}$$

Substituting (2-3) into (3-2), we get

$$\begin{aligned}
 &R_{1313} + \lambda^2 R_{1414} + R_{2323} + \lambda^2 R_{2424} - 2\lambda R_{1234} \\
 &\geq 2(1 + \lambda^2)\bar{K}_{\min} - \frac{2}{3}(1 + \lambda^2)(\bar{K}_{\max} - \bar{K}_{\min}) \\
 &\quad + \text{Ric}(e_3) - nHh_{33}^{n+1} - \sum_j \bar{R}_{3j3j} \\
 &\quad + \lambda^2(\text{Ric}(e_4) - nHh_{44}^{n+1} - \sum_j \bar{R}_{4j4j}) \\
 &\quad + \frac{1}{2}(1 + \lambda^2)(\text{Ric}(e_1) - nHh_{11}^{n+1} - \sum_j \bar{R}_{1j1j}) \\
 &\quad + \frac{1}{2}(1 + \lambda^2)(\text{Ric}(e_2) - nHh_{22}^{n+1} - \sum_j \bar{R}_{2j2j}) \\
 &\quad + \frac{1}{2}(h_{11}^{n+1} + h_{33}^{n+1})^2 + \frac{1}{2}(h_{22}^{n+1} + h_{33}^{n+1})^2 \\
 &\quad + \frac{1}{2}\lambda^2(h_{11}^{n+1} + h_{44}^{n+1})^2 + \frac{1}{2}\lambda^2(h_{22}^{n+1} + h_{44}^{n+1})^2 \\
 (3-3) \quad &\geq 2(1 + \lambda^2)\bar{K}_{\min} - \frac{2}{3}(1 + \lambda^2)(\bar{K}_{\max} - \bar{K}_{\min}) - 2(1 + \lambda^2)(n - 1)\bar{K}_{\max} \\
 &\quad + \text{Ric}(e_3) + \lambda^2 \text{Ric}(e_4) + \frac{1}{2}(1 + \lambda^2) \text{Ric}(e_1) + \frac{1}{2}(1 + \lambda^2) \text{Ric}(e_2) \\
 &\quad + \frac{1}{2}(h_{11}^{n+1} + h_{33}^{n+1})^2 - \frac{1}{2}nH(h_{11}^{n+1} + h_{33}^{n+1}) \\
 &\quad + \frac{1}{2}(h_{22}^{n+1} + h_{33}^{n+1})^2 - \frac{1}{2}nH(h_{22}^{n+1} + h_{33}^{n+1}) \\
 &\quad + \frac{1}{2}\lambda^2(h_{11}^{n+1} + h_{44}^{n+1})^2 - \frac{1}{2}nH\lambda^2(h_{11}^{n+1} + h_{44}^{n+1}) \\
 &\quad + \frac{1}{2}\lambda^2(h_{22}^{n+1} + h_{44}^{n+1})^2 - \frac{1}{2}nH\lambda^2(h_{22}^{n+1} + h_{44}^{n+1}) \\
 &= 2(1 + \lambda^2)\bar{K}_{\min} - \frac{2}{3}(1 + \lambda^2)(\bar{K}_{\max} - \bar{K}_{\min}) - 2(1 + \lambda^2)(n - 1)\bar{K}_{\max} \\
 &\quad + \text{Ric}(e_3) + \lambda^2 \text{Ric}(e_4) + \frac{1}{2}(1 + \lambda^2) \text{Ric}(e_1) + \frac{1}{2}(1 + \lambda^2) \text{Ric}(e_2) \\
 &\quad + \frac{1}{2}(h_{11}^{n+1} + h_{33}^{n+1} - \frac{1}{2}nH)^2 + \frac{1}{2}(h_{22}^{n+1} + h_{33}^{n+1} - \frac{1}{2}nH)^2 \\
 &\quad + \frac{1}{2}\lambda^2(h_{11}^{n+1} + h_{44}^{n+1} - \frac{1}{2}nH)^2 + \frac{1}{2}\lambda^2(h_{22}^{n+1} + h_{44}^{n+1} - \frac{1}{2}nH)^2 \\
 &\quad - \frac{1}{4}(1 + \lambda^2)n^2H^2.
 \end{aligned}$$

By assumption, we have

$$(3-4) \quad \text{Ric}(e_i) > \left((n - \frac{2}{3})\bar{K}_{\max} - \frac{4}{3}\bar{K}_{\min} \right) + \frac{1}{8}n^2H^2.$$

Substituting (3-4) into (3-3), we have

$$(3-5) \quad \begin{aligned} &R_{1313} + \lambda^2 R_{1414} + R_{2323} + \lambda^2 R_{2424} - 2\lambda R_{1234} \\ &> 2(1 + \lambda^2)\bar{K}_{\min} - \frac{2}{3}(1 + \lambda^2)(\bar{K}_{\max} - \bar{K}_{\min}) - 2(1 + \lambda^2)(n - 1)\bar{K}_{\max} \\ &\quad + 2(1 + \lambda^2)\left((n - \frac{2}{3})\bar{K}_{\max} - \frac{4}{3}\bar{K}_{\min} + \frac{1}{8}n^2H^2 \right) - (1 + \lambda^2)\frac{1}{4}n^2H^2 \\ &= 2(1 + \lambda^2)\left(- (n - \frac{2}{3}) + (n - \frac{2}{3}) \right)\bar{K}_{\max} \\ &\quad + 2(1 + \lambda^2)\left(1 + \frac{1}{3} - \frac{4}{3} \right)\bar{K}_{\min} = 0. \end{aligned}$$

It follows from Theorem 2.2 that the normalized Ricci flow with initial metric g_0 ,

$$\frac{\partial}{\partial t} g(t) = -2\text{Ric}_{g(t)} + \frac{2}{n}r_{g(t)}g(t),$$

exists for all time and converges to a constant curvature metric as $t \rightarrow \infty$. Also, M is diffeomorphic to a spherical space form. In particular, if M is simply connected, then M is diffeomorphic to S^n . This completes the proof of Theorem 1.1. \square

Corollary 3.1. *Suppose M is an n -dimensional compact submanifold ($n \geq 4$) in an $(n + p)$ -dimensional pinched Riemannian manifold N^{n+p} whose sectional curvature satisfies $b \leq \bar{K}_N \leq c$. If the Ricci curvature of M satisfies*

$$\text{Ric}_M > \left(n - \frac{2}{3} \right)c - \frac{4}{3}b + \frac{1}{8}n^2H^2,$$

then M is diffeomorphic to a spherical space form. In particular, if M is simply connected, then M is diffeomorphic to S^n .

Corollary 3.2. *Suppose M is an n -dimensional compact submanifold ($n \geq 4$) in an $(n + p)$ -dimensional pointwise δ -pinched Riemannian manifold N^{n+p} , where $\delta > 0$. If the Ricci curvature of M satisfies*

$$\text{Ric}_M > \left(n - \frac{2}{3} - \frac{4}{3}\delta \right)\bar{K}_{\max} + \frac{1}{8}n^2H^2,$$

then M is diffeomorphic to a spherical space form. In particular, if M is simply connected, then M is diffeomorphic to S^n .

Proof of Theorem 1.2. It follows from the Gauss equation (2-1) that

$$(3-6) \quad \text{Ric}(e_i) = (n - 1)c + \sum_{\alpha,k} (h_{i\alpha}^\alpha h_{kk}^\alpha - (h_{ik}^\alpha)^2).$$

This, together with the assumption, implies that

$$\begin{aligned}
 (3-7) \quad & \sum_{k=2}^n (2|h(e_1, e_k)|^2 - \langle h(e_1, e_1), h(e_k, e_k) \rangle) \\
 &= 2 \sum_{\alpha} \sum_{k=2}^n (h_{1k}^{\alpha})^2 - \sum_{\alpha} \sum_{k=2}^n h_{11}^{\alpha} h_{kk}^{\alpha} \\
 &= -2 \operatorname{Ric}(e_1) + 2(n-1)c - \sum_{\alpha} (h_{11}^{\alpha})^2 + nHh_{11}^{n+1} \\
 &= -2 \operatorname{Ric}(e_1) + 2(n-1)c - \sum_{\alpha \neq n+1} (h_{11}^{\alpha})^2 - (h_{11}^{n+1} - \frac{1}{2}nH)^2 + \frac{1}{4}n^2H^2 \\
 &\leq -2 \operatorname{Ric}(e_1) + 2(n-1)c + \frac{1}{4}n^2H^2 < (n-1)c,
 \end{aligned}$$

for any orthonormal basis $\{e_i\}$ of T_xM at any point $x \in M$.

Suppose that $\pi_1(M) \neq 0$. Because M is compact, it follows from a classical theorem due to Cartan and Hadamard that there exists a minimal closed geodesic in any nontrivial homotopy class in $\pi_1(M)$. However, by Theorem 2.3, we know that there do not exist any stable integral 1-currents on M . This contradicts the hypothesis. Therefore, $\pi_1(M) = 0$. This proves Theorem 1.2. \square

Applying Theorem 1.2 to Theorem C, we get a refined version of Ejiri’s rigidity theorem.

Theorem 3.3. *Suppose M is an n -dimensional compact orientable minimal submanifold in an $(n + p)$ -dimensional unit sphere S^{n+p} , where $n \geq 4$. If the Ricci curvature of M satisfies*

$$\operatorname{Ric}_M \geq n - 2,$$

then M is either

- the totally geodesic submanifold S^n , or
- the Clifford torus $S^m(\sqrt{1/2}) \times S^m(\sqrt{1/2})$ in S^{n+1} with $n = 2m$, or
- $\mathbb{C}P^2(\frac{4}{3})$, the 2-dimensional complex projective space minimally immersed in S^7 with constant holomorphic sectional curvature $\frac{4}{3}$.

Proof of Theorem 1.3. When $n = 3$, we have

$$(3-8) \quad \operatorname{Ric}_M > c + \frac{9}{8}H^2 \geq 0.$$

From Hamilton’s convergence theorem [1982] for Ricci flow in dimension 3, it follows that M is diffeomorphic to a 3-dimensional spherical space form. When $n \geq 4$, it follows from Theorem 1.1 that M is diffeomorphic to a spherical space form.

On the other hand, it follows from the assumption that

$$(3-9) \quad \operatorname{Ric}_M > (n - 2)c + \frac{1}{8}n^2H^2 \geq \frac{1}{2}(n - 1)c + \frac{1}{8}n^2H^2 \quad \text{for } n \geq 3.$$

This together with Theorem 1.2 implies that M is simply connected. Therefore, M is diffeomorphic to S^n . This completes the proof of Theorem 1.3. \square

The following example shows that the pinching condition in Theorem 1.3 is sharp.

Example 3.4. (i) Let $M = \mathbb{C}P^2(\frac{4}{3}(c + H^2))$ be a 2-dimensional complex projective space minimally immersed into $S^7(1/\sqrt{c + H^2})$ with constant holomorphic sectional curvature $\frac{4}{3}(c + H^2)$, and let $S^7(1/\sqrt{c + H^2})$ be the totally umbilical sphere in $F^{4+p}(c)$. Here H is a nonnegative constant and $c + H^2 > 0$. Then M is a compact submanifold in $F^{4+p}(c)$ with constant mean curvature H and constant Ricci curvature $\text{Ric}_M \equiv 2c + 2H^2$. M is not a topological sphere.

(ii) Let $M = S^2(1/\sqrt{2(c + H^2)}) \times S^2(1/\sqrt{2(c + H^2)})$ be a minimal Clifford hypersurface in $S^5(1/\sqrt{c + H^2})$, and let $S^5(1/\sqrt{c + H^2})$ be the totally umbilical sphere in $F^{4+p}(c)$. Here H is a nonnegative constant and $c + H^2 > 0$. Then M is a compact submanifold in $F^{4+p}(c)$ with constant mean curvature H and constant Ricci curvature $\text{Ric}_M \equiv 2c + 2H^2$ that is not a topological sphere.

(iii) Let $M = S^m(1/\sqrt{2c}) \times S^m(1/\sqrt{2c})$ be a minimal Clifford hypersurface in $F^{n+1}(c)$ with $c > 0$ and $n = 2m \geq 6$, and let $F^{n+1}(c)$ be the totally geodesic submanifold in $F^{n+p}(c)$. Then M is a compact minimal submanifold in $F^{n+p}(c)$ with $\text{Ric}_M \equiv (n - 2)c$, and is not homeomorphic to S^n .

Remark 3.5. Using the nonexistence theorem for stable currents on submanifolds in hyperbolic spaces [Fu and Xu 2008] and Theorem 1.1, one can also extend Theorem 1.3 to the case of compact submanifolds in hyperbolic spaces.

Motivated by Theorem 1.3 and the convergence results for mean curvature flow in higher codimensions [Andrews and Baker 2010; Liu et al. 2011], we would like to propose the following conjecture on mean curvature flow in higher codimensions.

Conjecture 3.6. *Let $M_0 = F_0(M)$ be an n -dimensional compact submanifold in an $(n + p)$ -dimensional space form $F^{n+p}(c)$, with $c + H^2 > 0$. If the Ricci curvature of M_0 satisfies*

$$\text{Ric}_{M_0} > (n - 2)c + \frac{1}{8}n^2 H^2,$$

then there exists for the mean curvature flow

$$(3-10) \quad \begin{cases} \frac{\partial}{\partial t} F(x, t) = n\xi(x, t), & x \in M, \quad t \geq 0, \\ F(\cdot, 0) = F_0(\cdot) \end{cases}$$

a unique smooth solution $F_t(\cdot)$, and either $F_t(\cdot)$ converges to a round point in finite time, or $c > 0$ and $F_t(\cdot)$ converges to a totally geodesic sphere as $t \rightarrow \infty$.

Acknowledgement

The authors thank Dr. Juan-Ru Gu and Dr. En-Tao Zhao for their helpful discussions. They also thank the referee for valuable suggestions.

References

- [An-Min and Jimin 1992] L. An-Min and L. Jimin, “An intrinsic rigidity theorem for minimal submanifolds in a sphere”, *Arch. Math. (Basel)* **58**:6 (1992), 582–594. MR 93b:53050 Zbl 0767.53042
- [Andrews and Baker 2010] B. Andrews and C. Baker, “Mean curvature flow of pinched submanifolds to spheres”, *J. Differential Geom.* **85**:3 (2010), 357–395. MR 2012a:53122 Zbl 05842818
- [Berger 2000] M. Berger, *Riemannian geometry during the second half of the twentieth century*, University Lecture Series **17**, American Mathematical Society, Providence, RI, 2000. MR 2000h:53002 Zbl 0944.53001
- [Brendle 2008] S. Brendle, “A general convergence result for the Ricci flow in higher dimensions”, *Duke Math. J.* **145**:3 (2008), 585–601. MR 2010a:53132 Zbl 1161.53052
- [Brendle 2010] S. Brendle, *Ricci flow and the sphere theorem*, Graduate Studies in Mathematics **111**, American Mathematical Society, Providence, RI, 2010. MR 2011e:53035 Zbl 1196.53001
- [Brendle and Schoen 2008] S. Brendle and R. M. Schoen, “Classification of manifolds with weakly $1/4$ -pinched curvatures”, *Acta Math.* **200**:1 (2008), 1–13. MR 2009h:53087 Zbl 1157.53020
- [Brendle and Schoen 2009] S. Brendle and R. Schoen, “Manifolds with $1/4$ -pinched curvature are space forms”, *J. Amer. Math. Soc.* **22**:1 (2009), 287–307. MR 2010a:53045 Zbl 05859406
- [Chern et al. 1970] S. S. Chern, M. do Carmo, and S. Kobayashi, “Minimal submanifolds of a sphere with second fundamental form of constant length”, pp. 59–75 in *Functional Analysis and Related Fields*, edited by F. E. Browder, Springer, New York, 1970. MR 42 #8424 Zbl 0216.44001
- [Ejiri 1979] N. Ejiri, “Compact minimal submanifolds of a sphere with positive Ricci curvature”, *J. Math. Soc. Japan* **31**:2 (1979), 251–256. MR 80h:53052 Zbl 0396.53026
- [Fu and Xu 2008] H. Fu and H. Xu, “Vanishing and topological sphere theorems for submanifolds in a hyperbolic space”, *Internat. J. Math.* **19**:7 (2008), 811–822. MR 2009f:53075 Zbl 1191.49045
- [Gu and Xu 2011] J. R. Gu and H. W. Xu, “The sphere theorems for manifolds with positive scalar curvature”, preprint, 2011. arXiv 1102.2424v1
- [Hamilton 1982] R. S. Hamilton, “Three-manifolds with positive Ricci curvature”, *J. Differential Geom.* **17**:2 (1982), 255–306. MR 84a:53050 Zbl 0504.53034
- [Huisken 1987] G. Huisken, “Deforming hypersurfaces of the sphere by their mean curvature”, *Math. Z.* **195**:2 (1987), 205–219. MR 88d:53058 Zbl 0626.53039
- [Lawson 1969] H. B. Lawson, Jr., “Local rigidity theorems for minimal hypersurfaces”, *Ann. of Math. (2)* **89** (1969), 187–197. MR 38 #6505 Zbl 0174.24901
- [Lawson and Simons 1973] H. B. Lawson, Jr. and J. Simons, “On stable currents and their application to global problems in real and complex geometry”, *Ann. of Math. (2)* **98** (1973), 427–450. MR 48 #2881 Zbl 0283.53049
- [Liu et al. 2011] K. F. Liu, H. W. Xu, F. Ye, and E. T. Zhao, “The extension and convergence of mean curvature flow in higher codimension”, preprint, 2011. arXiv 1104.0971v1
- [Petersen and Tao 2009] P. Petersen and T. Tao, “Classification of almost quarter-pinched manifolds”, *Proc. Amer. Math. Soc.* **137**:7 (2009), 2437–2440. MR 2010m:53046 Zbl 1168.53020

- [Shen 1989] Z. M. Shen, “A sphere theorem for manifolds of positive Ricci curvature”, *Indiana Univ. Math. J.* **38**:1 (1989), 229–233. MR 90h:53044 Zbl 0678.53032
- [Shen 1992] Y. B. Shen, “Curvature pinching for three-dimensional minimal submanifolds in a sphere”, *Proc. Amer. Math. Soc.* **115**:3 (1992), 791–795. MR 92i:53058 Zbl 0760.53038
- [Shiohama 2000] K. Shiohama, “Sphere theorems”, pp. 865–903 in *Handbook of differential geometry*, vol. 1, edited by F. J. E. Dillen and L. C. A. Verstraelen, North-Holland, Amsterdam, 2000. MR 2001c:53051 Zbl 0968.53003
- [Shiohama and Xu 1997] K. Shiohama and H. Xu, “The topological sphere theorem for complete submanifolds”, *Compositio Math.* **107**:2 (1997), 221–232. MR 98i:53080 Zbl 0905.53038
- [Simons 1968] J. Simons, “Minimal varieties in riemannian manifolds”, *Ann. of Math. (2)* **88** (1968), 62–105. MR 38 #1617 Zbl 0181.49702
- [Xin 1984] Y. L. Xin, “An application of integral currents to the vanishing theorems”, *Sci. Sinica Ser. A* **27**:3 (1984), 233–241. MR 86b:49060 Zbl 0554.49021
- [Xu 2012] H.-W. Xu, “Recent developments in differentiable sphere theorems”, pp. 415–430 in *Fifth International Congress of Chinese Mathematicians*, vol. 1, edited by L. Ji et al., AMS/IP Studies in Advanced Mathematics **51**, Amer. Math. Soc., Providence, RI, 2012.
- [Xu and Gu 2010] H.-W. Xu and J.-R. Gu, “An optimal differentiable sphere theorem for complete manifolds”, *Math. Res. Lett.* **17**:6 (2010), 1111–1124. MR 2012a:53054 Zbl 1221.53068
- [Xu and Zhao 2009] H.-W. Xu and E.-T. Zhao, “Topological and differentiable sphere theorems for complete submanifolds”, *Comm. Anal. Geom.* **17**:3 (2009), 565–585. MR 2010k:53067 Zbl 1185.53033
- [Yau 1974] S. T. Yau, “Submanifolds with constant mean curvature, I”, *Amer. J. Math.* **96**:2 (1974), 346–366. MR 51 #6670 Zbl 0304.53041
- [Yau 1975] S. T. Yau, “Submanifolds with constant mean curvature, II”, *Amer. J. Math.* **97**:1 (1975), 76–100. MR 51 #6670 Zbl 0304.53042
- [Yau 1993] S.-T. Yau, “Open problems in geometry”, pp. 1–28 in *Differential geometry: partial differential equations on manifolds* (Los Angeles, 1990), edited by R. Greene and S.-T. Yau, Proc. Sympos. Pure Math. **54**, Amer. Math. Soc., Providence, RI, 1993. MR 94k:53001 Zbl 0801.53001

Received April 21, 2011. Revised July 29, 2011.

HONG-WEI XU
CENTER OF MATHEMATICAL SCIENCES
ZHEJIANG UNIVERSITY
HANGZHOU, 310027
CHINA
xuhw@cms.zju.edu.cn

LING TIAN
CENTER OF MATHEMATICAL SCIENCES
ZHEJIANG UNIVERSITY
HANGZHOU, 310027
CHINA
tianling@cms.zju.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 254 No. 2 December 2011

Curvatures of spheres in Hilbert geometry	257
ALEXANDER BORISENKO and EUGENE OLIN	
A formula equating open and closed Gromov–Witten invariants and its applications to mirror symmetry	275
KWOKWAI CHAN	
A note on p -harmonic l -forms on complete manifolds	295
LIANG-CHU CHANG and CHIUNG-JUE ANNA SUNG	
The Cheeger constant of curved strips	309
DAVID KREJČIŘÍK and ALDO PRATELLI	
Structure of solutions of 3D axisymmetric Navier–Stokes equations near maximal points	335
ZHEN LEI and QI S. ZHANG	
Local comparison theorems for Kähler manifolds	345
GANG LIU	
Structurable algebras of skew-rank 1 over the affine plane	361
SUSANNE PUMPLÜN	
An analogue of Krein’s theorem for semisimple Lie groups	381
SANJOY PUSTI	
Une remarque de dynamique sur les variétés semi-abéliennes	397
GAËL RÉMOND	
Fourier transforms of semisimple orbital integrals on the Lie algebra of SL_2	407
LOREN SPICE	
On noncompact τ -quasi-Einstein metrics	449
LIN FENG WANG	
Decomposition of de Rham complexes with smooth horizontal coefficients for semistable reductions	465
QIHONG XIE	
A differentiable sphere theorem inspired by rigidity of minimal submanifolds	499
HONG-WEI XU and LING TIAN	



0030-8730(201112)254:2;1-A