

*Pacific
Journal of
Mathematics*

**DIRICHLET-FORD DOMAINS AND ARITHMETIC
REFLECTION GROUPS**

GRANT S. LAKELAND

DIRICHLET–FORD DOMAINS AND ARITHMETIC REFLECTION GROUPS

GRANT S. LAKELAND

We show that a Fuchsian group, acting on the upper half-plane model for \mathbb{H}^2 , admits a Ford domain which is also a Dirichlet domain, for some center, if and only if it is an index 2 subgroup of a reflection group. This is used to exhibit an example of a maximal arithmetic hyperbolic reflection group which is not congruence. Analogous results, and counterexamples, are given in the case of Kleinian groups.

1. Introduction

The action of the modular group $\mathrm{PSL}_2(\mathbb{Z})$ on the upper half-plane model for hyperbolic 2-space \mathbb{H}^2 has been extensively studied. It is well known that a fundamental domain for this action is given by the triangle T with vertices at $\rho = \frac{1}{2} + \frac{\sqrt{-3}}{2}$, $-\bar{\rho}$ and ∞ . This domain is an example of two common constructions of fundamental domains for Fuchsian groups: it is both a Ford domain and a Dirichlet domain for the action of the modular group. Furthermore, it arises from more than one distinct choice of Dirichlet center, as taking the Dirichlet domain centered at any $z_0 = iy$, for $y > 1$, gives rise to T . One expects the Dirichlet domain to change along with the choice of center [Díaz and Ushijima 2009], so in this sense the modular group exhibits some atypical properties.

It is also well known that $\mathrm{PSL}_2(\mathbb{Z})$ is the orientation-preserving index 2 subgroup of the group generated by reflections in an ideal triangle in \mathbb{H}^2 with angles $\frac{\pi}{2}$, $\frac{\pi}{3}$ and 0, located at i , ρ and ∞ respectively. More generally, a *hyperbolic reflection group* is a subgroup of $\mathrm{Isom}(\mathbb{H}^2)$ generated by reflections in the sides of a polygon $Q \subset \mathbb{H}^2$. Such a group is discrete if and only if each angle of Q is either 0 or an integer submultiple of π .

The purpose of this paper is to determine exactly which Fuchsian groups admit a fundamental domain that is both a Dirichlet domain and a Ford domain (which we will call a *DF domain*) or a Dirichlet domain for multiple centers (a *double Dirichlet domain*).

It turns out that the above properties of $\mathrm{PSL}_2(\mathbb{Z})$ are very much related to that;

MSC2010: primary 20H10; secondary 19B37.

Keywords: fundamental domain, arithmetic reflection groups, congruence.

we give in Theorems 5.3 and 5.4 the following characterization for such groups:

Theorem. *A finitely generated, finite coarea Fuchsian group Γ admits a DF domain (or a double Dirichlet domain) P if and only if Γ is an index 2 subgroup of the discrete group G of reflections in a hyperbolic polygon Q .*

The condition given by this result provides a method of checking whether a given Fuchsian group is the index 2 orientation-preserving subgroup of a hyperbolic reflection group. This is particularly useful in the context of *maximal arithmetic* reflection groups. A noncompact hyperbolic reflection group is *arithmetic* if it is commensurable with $\mathrm{PSL}_2(\mathbb{Z})$. A maximal arithmetic reflection group is then an arithmetic reflection group which is not properly contained in another arithmetic reflection group. As an application of the above theorem, we give:

Corollary. *There exists a maximal arithmetic hyperbolic reflection group which is not congruence.*

This answers a question raised by Agol, Belolipetsky, Storm, and Whyte in [Agol et al. 2008] (see also [Belolipetsky 2011]).

Any group Γ satisfying the theorem must have *genus zero* [Long et al. 2006], as well as a certain symmetrical property. Having a double Dirichlet or DF domain therefore also gives an obstruction to the group having *nontrivial cuspidal cohomology* [Grunewald and Schwermer 1981].

Motivated by this, one may ask whether there exists a similar obstruction to nontrivial cuspidal cohomology for Kleinian groups. We will show that this is not the case: in Section 7 we exhibit a Kleinian group which possesses both nontrivial cuspidal cohomology and a DF domain. However, the condition of having such a domain does impose some restrictions on Γ ; perhaps the most striking is that the group possesses a generating set, all of whose elements have real trace (Theorem 7.3).

This paper is organized as follows. After the preliminaries of Section 2, Section 3 will examine Fuchsian groups with DF domains, and show that such domains are symmetrical and give rise to punctured spheres. The more general case of the double Dirichlet domain is discussed in Section 4. In Section 5, it will be shown that the main theorem follows from the previous sections and standard results on reflection groups. An example of a noncongruence maximal arithmetic hyperbolic reflection group can be found in Section 6. Section 7 contains a discussion of these domains in the setting of Kleinian groups.

2. Preliminaries

We will work in the upper half-plane model for the hyperbolic plane. The group of conformal, orientation-preserving isometries (or linear fractional transformations)

of \mathbb{H}^2 can be identified with $\mathrm{PSL}_2(\mathbb{R})$ via the correspondence

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \longleftrightarrow z \mapsto \frac{az + b}{cz + d}.$$

A *Fuchsian group* Γ is a subgroup of $\mathrm{PSL}_2(\mathbb{R})$, discrete with respect to the topology induced by regarding that group as a subset of \mathbb{R}^4 . The *Dirichlet domain* for Γ centered at z_0 is defined to be

$$\{x \in \mathbb{H}^2 \mid d(x, z_0) \leq d(x, \alpha(z_0)) \text{ for all } \alpha \in \Gamma \text{ distinct from } 1\}.$$

It is an intersection of closed half-spaces.

Beardon [1983, Section 9.5] demonstrates an alternative definition, in terms of reflections, which allows us to define a *generalized Dirichlet domain* by taking our center to be on the boundary $\partial\mathbb{H}^2$. We will typically conjugate Γ in $\mathrm{PSL}_2(\mathbb{R})$ so that this center is placed at ∞ in the upper half-plane. We suppose Γ is *zonal*, or that ∞ is a parabolic fixed point, and so the reflections given are not uniquely determined for any parabolic isometry fixing ∞ . To account for this, we define a *Ford domain* [Ford 1951] to be the intersection of the region exterior to all isometric circles with a fundamental domain for the action of the parabolic subgroup stabilizing ∞ , $\Gamma_\infty < \Gamma$, which is a vertical strip.

For a given finitely generated Fuchsian group Γ , the *signature*

$$(g; n_1, \dots, n_t; m; f)$$

of Γ records the topology of the quotient space \mathbb{H}^2/Γ , where g is the genus, t is the number of cone points of orders n_1, \dots, n_t respectively, m is the number of cusps, and f is the number of infinite area funnels. If Γ is the orientation-preserving index 2 subgroup of a reflection group, then \mathbb{H}^2/Γ is a sphere with cusps and/or cone points, and thus in this case we have $g = 0$. If additionally Γ has finite coarea, then we also have that $f = 0$.

The group of orientation-preserving isometries of the upper half-space model of \mathbb{H}^3 can likewise be identified with $\mathrm{PSL}_2(\mathbb{C})$. A *Kleinian group* is a discrete subgroup of this isometry group. The definitions of Dirichlet domain and Ford domain carry over to this situation, with one small modification: instead of Γ being zonal, we assume that Γ_∞ contains a copy of \mathbb{Z}^2 .

Throughout, we will assume that Γ is finitely generated, and hence that all fundamental domains we encounter have a finite number of sides. For simplicity, we will also suppose that $f = 0$ and that Γ has finite covolume (and thus that all fundamental domains have finite volume; that is, finitely many ideal vertices, each adjacent to two sides), although many of the arguments should extend to the case where Γ does not have finite covolume.

3. DF domains

Suppose Γ contains a nontrivial parabolic subgroup Γ_∞ fixing ∞ . In \mathbb{H}^2 , Γ_∞ must be cyclic, and after conjugation, we may take it to be generated by

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Theorem 3.1. *If Γ admits a DF domain, then the quotient space \mathbb{H}^2/Γ is a punctured sphere, possibly with cone points.*

Before commencing the proof of this, we will prove two elementary but important lemmas. The first is given as Exercise 2 in Section 9.6 of [Beardon 1983].

Lemma 3.2. *Any vertex cycle on the boundary of a Ford domain P is contained within a horocycle based at ∞ .*

Proof. Fix a vertex v . By construction of P , v lies on or exterior to all isometric circles, and necessarily lies on at least one. We first consider a $\gamma \in \Gamma$ such that $v \notin S_\gamma$. Then v lies exterior to S_γ . It follows that γ sends v into the interior of $S_{\gamma^{-1}}$. Thus $\gamma(v)$ cannot be a vertex of P . Now suppose that $v \in S_\gamma$. Then γ is the composition of reflection in S_γ , which fixes v , and reflection in a vertical line. It therefore necessarily preserves the imaginary part of v , proving the lemma. \square

Remark. The lemma holds for any point on the boundary of the Ford domain P . For our purposes, it will be enough to have it for the vertices of P .

Lemma 3.3 [Greenberg 1977, p. 203]. *Let P be a Dirichlet domain for Γ with center z_0 . Let $1 \neq \gamma$ be an element of Γ and suppose that $z, \gamma(z) \in \partial P \cap \mathbb{H}^2$. Then $d_{\mathbb{H}}(z, z_0) = d_{\mathbb{H}}(\gamma(z), z_0)$.*

Proof. This is an application of the definition of a Dirichlet domain stated above. Specifically, setting $x = z$ and $\alpha = \gamma^{-1}$ yields the inequality

$$d(z, z_0) \leq d(z, \gamma^{-1}(z_0)) = d(\gamma(z), z_0),$$

the latter equality holding because γ is an isometry. Setting $x = \gamma(z)$ and $\alpha = \gamma$ now gives

$$d(\gamma(z), z_0) \leq d(\gamma(z), \gamma(z_0)) = d(z, z_0).$$

Combining these two inequalities gives the required equality. \square

Proof of Theorem 3.1. Suppose we are given a DF domain P for Γ . Since P is a Ford domain, it is contained in a fundamental region for Γ_∞ , which is a vertical strip

$$\{z \in \mathbb{H}^2 \mid x_0 \leq \operatorname{Re}(z) \leq x_0 + 1\}$$




Figure 1. $\gamma(v) = v^*$.

for some $x_0 \in \mathbb{R}$. Shimizu’s Lemma (see [Maskit 1988], p. 18) tells us that the radii of the isometric circles S_γ cannot exceed 1. Thus we may consider a point $z = x_0 + iy \in \partial P$, where $y > 1$. Choosing $\gamma = T$, and applying Lemma 3.3 to z and $\gamma(z)$, we find that $\text{Re}(z_0) = x_0 + \frac{1}{2}$.

Next suppose that $v \in \mathbb{H}^2$ is a vertex of P , and $\gamma \in \Gamma$ a side pairing such that $\gamma(v)$ is another vertex of P . Then, by Lemma 3.2, $\text{Im}(\gamma(v)) = \text{Im}(v)$, and by Lemma 3.3, $d_{\mathbb{H}}(\gamma(v), z_0) = d_{\mathbb{H}}(v, z_0)$. Consider the two sets $\{z \in \mathbb{H}^2 \mid \text{Im}(z) = \text{Im}(v)\}$ and $\{z \in \mathbb{H}^2 \mid d_{\mathbb{H}}(z, z_0) = d_{\mathbb{H}}(v, z_0)\}$. The former is the horizontal line through v , and the latter a circle with Euclidean center located vertically above z_0 (see Figure 1). In particular, the picture is symmetrical in the vertical line $\{\text{Re}(z) = x_0 + \frac{1}{2}\}$. Either $\gamma(v) = v$ or $\gamma(v) = v^*$, where v^* is the reflection of v in the line $\{\text{Re}(z) = x_0 + \frac{1}{2}\}$.

Suppose that $\gamma(v) = v$. Then, by considering a point $w \in \partial P$ close to v , the fact that $d(w, z_0) = d(\gamma(w), z_0)$ means that v necessarily lies directly below the Dirichlet center z_0 . The contrapositive of this states that if $\text{Re}(v) \neq x_0 + \frac{1}{2}$, then any side pairing γ pairing v with a vertex of P must send v to v^* .

Suppose now that $v \in \partial \mathbb{H}^2$ is a vertex of P . Then two isometric circles meet at v . Fix one such circle S . S is the isometric circle S_γ of some element $\gamma \in \Gamma$. S_γ contains a side of P adjacent to v , and we pick two points of S_γ , $w_1, w_2 \in \partial P \cap \mathbb{H}^2$. By Lemma 3.3, γ must send both w_1 and w_2 to points the same respective distances from z_0 . For each i , the point w_i is either fixed or sent to its reflection in the line $\{\text{Re}(z) = \text{Re}(z_0)\}$. If w_1 were fixed, w_2 would neither be fixed nor sent to its reflection, and vice versa if w_2 were fixed. Thus we conclude that γ sends points of S to their reflections in the line $\{\text{Re}(z) = \text{Re}(z_0)\}$.

We can now show that \mathbb{H}^2/Γ is a punctured sphere. We first identify the two vertical sides of P , creating the cusp at ∞ and a circle awaiting identification. Consider the side of P adjacent to the side contained in the vertical line $\text{Re}(z) = x_0$.

This side lies on some isometric circle S_γ . We see that γ must identify our side with a side adjacent to the side of P contained in the line $\text{Re}(z) = x_0 + 1$. Working inwards toward the center and applying this argument repeatedly, we see that all sides must pair up symmetrically. In particular, there can not exist two hyperbolic generators whose axes intersect precisely once. Thus we conclude that the quotient space has genus zero. \square

Remarks. (1) We may take the Dirichlet center of P to be any point of the interior of P on this vertical line $\{\text{Re}(z) = x_0 + \frac{1}{2}\}$. To see this, let z_0 be any such point, and $\gamma \in \Gamma \setminus \Gamma_\infty$ a side pairing of P . Since $\gamma(S_\gamma) = S_{\gamma^{-1}}$, and this pair is arranged symmetrically with respect to the line $\{\text{Re}(z) = x_0 + \frac{1}{2}\}$, both of these isometric circles are geodesics of the form used to construct the Dirichlet polygon centered at z_0 .

(2) The converse of [Theorem 3.1](#) is false. The symmetrical nature of P implies a certain symmetry in the quotient space \mathbb{H}^2/Γ , namely that the surface admits an orientation-reversing involution of order 2. This is not the case for a generic punctured sphere.

4. Double Dirichlet domains

We now suppose that the same fundamental domain P is obtained when we construct the Dirichlet domains P_0 and P_1 centered at z_0 and $z_1 \in \mathbb{H}^2$ respectively. For comparison with the previous section, we will assume that we have conjugated Γ in $\text{PSL}_2(\mathbb{R})$ so that the geodesic line L containing z_0 and z_1 is vertical.

Theorem 4.1. *If the Dirichlet domains P_0 and P_1 for Γ , centered at $z_0 \neq z_1 \in \mathbb{H}^2$ respectively, coincide, then the quotient space \mathbb{H}^2/Γ is a sphere, with cone points and/or punctures.*

Proof. Much of the work in [Section 3](#) was concerned with showing precisely how the sides of P were identified. This follows relatively swiftly here, once we have cleared up one technical point. We often think of a fundamental domain as a subset of \mathbb{H}^2 combined with a set of side pairings identifying its sides. We only assume that the sets P_0 and P_1 are equal, and thus we must make sure that Γ identifies their sides the same way each time.

Lemma 4.2. *If $P = P_0 = P_1$ is the Dirichlet domain centered at z_0 and at z_1 , then the sides of P are identified the same way in each case.*

Proof. Suppose, for the sake of contradiction, that this is not the case. Any side of a Dirichlet domain bisects the domain's center and its image under some isometry. Here, we have a side A of P which is the bisector of both the pair z_0 and $\gamma_0^{-1}(z_0)$ and the pair z_1 and $\gamma_1^{-1}(z_1)$, where $\gamma_0 \neq \gamma_1$ are the isometries defining that side of P . It follows that γ_0 pairs A with some side B , and γ_1 pairs A with some other

side $C \neq B$. Let $d := d(z_0, z_1)$ be the distance between the two centers z_0 and z_1 . Since $\gamma_0^{-1}(z_0)$ and $\gamma_1^{-1}(z_1)$ are the reflections of each in A , we see that

$$d(\gamma_0^{-1}(z_0), \gamma_1^{-1}(z_1)) = d.$$

Applying the isometry γ_1 to both points, this gives that

$$d(\gamma_1(\gamma_0^{-1}(z_0)), z_1) = d.$$

Now, if $\gamma_1(\gamma_0^{-1}(z_0)) = z_0$, then the isometries γ_0 and γ_1 both send $\gamma_0^{-1}(z_0)$ to z_0 and $\gamma_1^{-1}(z_1)$ to z_1 . Since they also both preserve orientation, this implies that $\gamma_0 = \gamma_1$, which is a contradiction. Thus $\gamma_1(\gamma_0^{-1}(z_0)) \neq z_0$. But then $\gamma_1(\gamma_0^{-1}(z_0))$ is a point in the orbit of z_0 , and thus the construction of P_0 involves the half-space $\{x \in \mathbb{H}^2 \mid d(x, z_0) \leq d(x, \gamma_1(\gamma_0^{-1}(z_0)))\}$. As we saw above,

$$d(\gamma_1(\gamma_0^{-1}(z_0)), z_1) = d(z_0, z_1) = d.$$

Hence z_1 is equidistant from z_0 and $\gamma_1(\gamma_0^{-1}(z_0))$. Thus z_1 cannot be in the interior of P_0 , contradicting the assumption that $P_0 = P_1$. □

The following result will allow us to conclude the proof of [Theorem 4.1](#).

Lemma 4.3. *Each side of P (and each point of ∂P) is identified with its reflection in the line L .*

Proof. Given a point $v \in \partial P$, v is sent to a point of ∂P the same distance away from z_0 . Put another way, v is sent somewhere on the hyperbolic circle of center z_0 and radius $d(v, z_0)$. But v is also sent to a point on the hyperbolic circle of center z_1 and radius $d(v, z_1)$. Thus we see a picture similar to [Figure 1](#), except instead of a horizontal line, we have a second circle, centered vertically above or below z_0 . These two circles intersect only at v if $v \in L$, and at v and v^* , the reflection of v in L , if $v \notin L$. If $v \in L$ then v is necessarily an elliptic fixed point and a vertex of P , and the two sides adjacent to v are identified with one another. If $v \notin L$, it suffices to show that v cannot be fixed by a side pairing, and thus must be identified with v^* . Let v be a nonvertex point, and γ the side pairing associated to the side containing v in its interior. Since v is not a vertex, it cannot be an elliptic fixed point, and so γ must identify v with v^* . From this, it follows that each vertex is also identified with its reflection. □

So we now know that our domain P has the same symmetrical property that we saw DF domains possess. If the line $L \cap \mathring{P}$ extends vertically to ∞ , then the argument from the proof of [Theorem 3.1](#) applies directly, and we are done. If the line terminates at a boundary point of P , then we observe that the two sides adjacent to this vertex are identified symmetrically, creating a cone point instead of a cusp. This creates a circle awaiting identification as in the proof of [Theorem 3.1](#), and the rest of the argument applies from there. □

Remarks. (1) The first remark at the end of [Section 3](#) applies here as well. That is, if we take any point $z \in L \cap \overset{\circ}{P}$ as our Dirichlet center, we will obtain the Dirichlet domain P . Thus we see that a Fuchsian group which admits a DF domain is simply one that admits Dirichlet domain with a line of centers and a cusp on the line of symmetry.

(2) The same discussion can also be used to show that these are the only Dirichlet centers giving rise to P . The Dirichlet center must be equidistant from a point of ∂P and its destination under its side pairing; in this setup, the locus of such points is always precisely L . Thus, it is impossible to find a Fuchsian group with a triangle of Dirichlet centers all giving rise to the same domain.

5. Reflection groups

The goal of this section is to prove the main theorem. As a corollary, we will show that given the signature of any sphere which can be obtained as a quotient of \mathbb{H}^2 , then we may exhibit a Fuchsian group Γ that admits a double Dirichlet domain (and a DF domain if there is at least one puncture) and gives rise to a quotient space of the given signature.

We first recall the following results regarding reflection groups (see [\[Ratcliffe 2006\]](#), Section 7.1).

Theorem 5.1. *Let G be a discrete reflection group with respect to the polygon Q . Then all the angles of Q are submultiples of π , and if g_S and g_T are reflections in the adjacent sides S and T of Q with $\theta(S, T) = \pi/k$, then $g_S \circ g_T$ has order k .*

Theorem 5.2. *Let Q be a finite sided convex hyperbolic polygon of finite volume, all of whose angles are submultiples of π . Then the group G , generated by reflections of \mathbb{H}^2 in the sides of Q , is a discrete reflection group with respect to the polygon Q .*

We will appeal to these results, as well as to the results of [Sections 3](#) and [4](#), in the following discussion.

Theorem 5.3. *If the finitely generated, orientation-preserving, finite coarea Fuchsian group Γ admits a double Dirichlet domain, or a DF domain, P , then Γ is an index 2 subgroup of the discrete group G of reflections in a hyperbolic polygon Q .*

Proof. Suppose first that Γ admits a DF domain P . We know that P has reflectional symmetry about a vertical axis L which bisects P . Since P is a fundamental domain for Γ , the side pairings of P generate Γ . Each side pairing, with the exception of the parabolic element pairing the vertical sides, has the form $\sigma_L \circ \sigma_i$, where σ_L denotes reflection in L and σ_i is reflection in the isometric circle S_i , $1 \leq i \leq m$, where S_i contains a side of P . Furthermore, since each side is paired with its mirror image in L , it suffices to consider the σ_i corresponding to sides in

one half of P . The parabolic side pairing can be written $\sigma_L \circ \sigma_K$, where σ_K is reflection in K , the vertical side of P in the same half as the S_i . Thus we have a generating set for Γ of the form

$$\{\sigma_L \circ \sigma_1, \dots, \sigma_L \circ \sigma_m, \sigma_L \circ \sigma_K\}$$

for some $m \in \mathbb{N}$. Consider the group G obtained by adding the reflection σ_L to this generating set. The set becomes

$$\{\sigma_L, \sigma_L \circ \sigma_1, \dots, \sigma_L \circ \sigma_m, \sigma_L \circ \sigma_K\}$$

and because $\sigma_L = \sigma_L^{-1}$ has order 2, it follows that we can replace the generator $\sigma_L \circ \sigma_i$ with the element σ_i and still have a generating set. The generating set

$$\{\sigma_L, \sigma_1, \dots, \sigma_m, \sigma_K\}$$

is precisely the set of reflections in the sides of a polygon Q with sides on K, L and $S_i, 1 \leq i \leq m$. To prove that all of the angles of Q are submultiples of π , it suffices to observe that the vertices of P are paired symmetrically, and that the Poincaré polyhedron theorem gives that the sum of the angles in each cycle is $2\pi/s$, for $s \in \mathbb{N}$. Now [Theorem 5.2](#) allows us to reach the desired conclusion.

To prove the result for the case where $L \cap \mathring{P}$ does not extend to $\partial\mathbb{H}^2$, we simply observe that, in this case, every side pairing generator of Γ can be written $\sigma_L \circ \sigma_i$, since here there are no vertical sides. Instead of the cusp at ∞ we have another finite vertex of P , but since this vertex lies on the line L , it must also be an elliptic fixed point, and the paragraph above applies. □

We now turn to the converse of [Theorem 5.3](#).

Theorem 5.4. *If G is a discrete group of reflections in a polygon $Q \subset \mathbb{H}^2$, then G contains an index 2 subgroup of orientation-preserving isometries that admits a double Dirichlet domain (and a DF domain if Q has an ideal vertex at ∞).*

Proof. Let Q be such a polygon. If necessary, rotate Q so that one of its sides is vertical. Call this side L . By [Theorem 5.1](#), all angles of Q are submultiples of π . Denote by σ_L reflection in the vertical side L of Q . If Q has another vertical side (and hence an ideal vertex at ∞), call this side K and denote reflection in K by σ_K . Denote by σ_i reflection in the (nonvertical) line S_i containing a side of Q . By definition, these reflections constitute a generating set for G . Let $\Gamma < G$ be the subgroup generated by elements of the form $\sigma_2 \circ \sigma_1$ where σ_1 and σ_2 are reflections in the generating set for G . Then Γ is a Fuchsian group. Since $\sigma_L \notin \Gamma$, we see that the set $P := Q \cup \sigma_L Q$ is contained within a fundamental domain for Γ . We will show that P is itself a fundamental domain for Γ .

To see this, denote by $T_i := \sigma_L(S_i)$ the geodesic obtained by reflecting S_i in L . Then T_i contains a side of P . Also denote $\sigma_L(K)$ by M . Then K is paired

with M by the element $\sigma_L \circ \sigma_K \in \Gamma$, and S_i is paired with T_i by $\sigma_L \circ \sigma_i$. Thus the sides of P are paired by generators of Γ . To see that these side pairings generate Γ themselves, consider a generating element $\sigma_2 \circ \sigma_1 \in \Gamma$. We may write

$$\sigma_2 \circ \sigma_1 = \sigma_2 \circ (\sigma_L \circ \sigma_L) \circ \sigma_1 = (\sigma_2 \circ \sigma_L) \circ (\sigma_L \circ \sigma_1) = (\sigma_L \circ \sigma_2)^{-1} \circ (\sigma_L \circ \sigma_1),$$

which shows that together, the elements $\sigma_L \circ \sigma_i$ and $\sigma_L \circ \sigma_K$ generate Γ . We therefore have that Γ has index 2 in G , and that P is a fundamental domain for Γ .

To see that Γ admits a fundamental domain of the required type, it will suffice to check that P is one. Let z_0 be any point on the line L which lies in the interior of P . If there is a second vertical side K , then it is the line bisecting z_0 and $\sigma_K(z_0)$, so $\sigma_L(K) = M$ bisects $\sigma_L(z_0) = z_0$ and $\sigma_L(\sigma_K(z_0))$. Thus M is a line of the form found in the definition on a Dirichlet domain centered at z_0 . A similar argument applied to $(\sigma_L \circ \sigma_K)^{-1} = \sigma_K \circ \sigma_L$ shows that K is also such a line. Now S_i is the line bisecting z_0 and $\sigma_i(z_0)$, so $\sigma_L(S_i) = T_i$ bisects $\sigma_L(z_0) = z_0$ and $\sigma_L(\sigma_i(z_0))$. This shows that T_i is a line of the form found in the definition on a Dirichlet domain centered at z_0 . A similar argument shows that the same is true of S_i , and thus we see that P must contain a Dirichlet domain centered at z_0 . But we know that P is itself a fundamental domain for Γ , so that P is a Dirichlet domain for any center $z_0 \in L \cap \mathring{P}$.

If there is a second vertical side K , we must also check that P is a Ford domain. S_i is the isometric circle of the generator $\sigma_L \circ \sigma_i$, and $T_i = \sigma_L(S_i)$ is the isometric circle of the inverse element. Since $\sigma_L \circ \sigma_K$ pairs the two vertical sides of P and generates Γ_∞ , it follows that P must contain a Ford domain for Γ . But P is itself a fundamental domain, so this Ford domain cannot be a proper subset, and hence is equal to P . \square

We now show that Fuchsian groups with this symmetrical property, though they necessarily have genus zero, have no other restrictions on their signature.

Corollary 5.5. *Given the signature $(0; n_1, \dots, n_t; m)$ of a (nontrivial, hyperbolic) sphere with $m \geq 0$ punctures and $t \geq 0$ cone points of orders $n_i \in \mathbb{N}$, for $1 \leq i \leq t$, there exists a Fuchsian group Γ such that Γ admits a double Dirichlet domain (and a DF domain if $m > 0$) and \mathbb{H}^2/Γ is a sphere of the given signature.*

Proof. Suppose $m > 0$. Construct Q by placing one vertex at ∞ , t vertices in \mathbb{H}^2 of angles π/n_i ($n_i \geq 2$) for $1 \leq i \leq t$, and $m-1$ ideal vertices in \mathbb{R} . If $m = 0$, construct a compact t -gon with angles $\pi/n_1, \dots, \pi/n_t$. Now let G be the group of hyperbolic isometries generated by reflections in the sides of Q . By [Theorem 5.4](#), Γ admits a DF domain (or double Dirichlet domain if $m = 0$) $P = Q \cup \sigma Q$, where σ denotes reflection in one of the vertical sides L of Q . The symmetrical identifications, combined with the Poincaré polyhedron theorem, give that the quotient surface has the required signature. \square

Remark. If $m > 0$ above, then there is a certain amount of freedom in our choice of the polygon Q . For example, we do not necessarily have to place one of the ideal vertices of Q at ∞ . We do so in order to ensure that we obtain a DF domain for Γ . Instead, we could have all of the ideal vertices lie in \mathbb{R} , thereby placing the line of symmetry L away from any of the ideal vertices. Similarly, if $m > 1$, we could construct Q so that L meets only one of the m ideal vertices, instead of 2 in the construction above. We also do not have to construct Q so that each angle is bisected by a vertical line; we only do so in order to demonstrate that it is possible to find the required polygon.

6. A noncongruence arithmetic maximal reflection group

In this section, we will prove explicitly that there exists a noncongruence arithmetic maximal hyperbolic reflection group. Recall that a noncompact hyperbolic reflection group $\Gamma_{\text{ref}} < \text{Isom}(\mathbb{H}^2)$ is called *arithmetic* if and only if it is commensurable with $\text{PSL}_2(\mathbb{Z})$. Such a group is then called *congruence* if it contains some principal congruence subgroup

$$\Gamma(n) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a \equiv d \equiv \pm 1, b \equiv c \equiv 0 \pmod n \right\} \subset \text{PSL}_2(\mathbb{Z}).$$

Consider the group $\Gamma < \text{PSL}_2(\mathbb{R})$ generated by the matrices

$$\begin{aligned} \gamma_1 &= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \gamma_2 = \begin{pmatrix} 0 & -\frac{1}{\sqrt{11}} \\ \sqrt{11} & 0 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} \sqrt{11} & \frac{5}{\sqrt{11}} \\ 2\sqrt{11} & \sqrt{11} \end{pmatrix}, \\ \gamma_4 &= \begin{pmatrix} 10 & 3 \\ 33 & 10 \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} 23 & 8 \\ 66 & 23 \end{pmatrix}. \end{aligned}$$

We first wish to show that Γ is discrete. Consider the group

$$\Gamma_0(11) = \left\{ \begin{pmatrix} a & b \\ 11c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}, ad - 11bc = 1 \right\} \subset \text{PSL}_2(\mathbb{Z}).$$

It is well known that the normalizer $N(\Gamma_0(11))$ of $\Gamma_0(11)$ in $\text{PSL}_2(\mathbb{R})$ is a (maximal arithmetic) Fuchsian group generated by $\Gamma_0(11)$ and

$$\begin{pmatrix} 0 & -\frac{1}{\sqrt{11}} \\ \sqrt{11} & 0 \end{pmatrix},$$

which is $\gamma_2 \in \Gamma$ [Maclachlan 1981; Long et al. 2006]. We see then that

$$\begin{pmatrix} 0 & -\frac{1}{\sqrt{11}} \\ \sqrt{11} & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ -11 & -5 \end{pmatrix} = \begin{pmatrix} \sqrt{11} & \frac{5}{\sqrt{11}} \\ 2\sqrt{11} & \sqrt{11} \end{pmatrix} = \gamma_3 \in \Gamma,$$

and since $\gamma_1, \gamma_4, \gamma_5 \in \Gamma_0(11)$, we have that $\Gamma < N(\Gamma_0(11))$, and so Γ is discrete. We next wish to construct a Ford domain for Γ . In [Figure 2](#) we see the isometric circles corresponding to the generators listed above and their inverses.




Figure 2. A Ford domain for Γ .

The claim is that this polygon is in fact the required Ford domain. To see this, observe that each generator γ_i can be decomposed into the product of two reflections $\gamma_i = \sigma_L \circ \sigma_i$, where σ_i is reflection in the isometric circle S_i of γ_i , σ_1 is reflection in the line $x = -\frac{1}{2}$, and σ_L is reflection in the line $x = 0$. Thus the elements of the generating set for Γ pair the sides of P , and each pushes \mathring{P} completely off itself. This shows that P is a fundamental domain for Γ ; by its construction, it is a Ford domain.

Thus we see that the quotient space \mathbb{H}^2/Γ is a sphere of signature $(0; 2, 2, 2, 2; 2)$ and area 4π . Further, P is a DF domain, as each of these generators pairs one side S_i of P with its reflection $\sigma_L(S_i)$ in the line $x = 0$. Thus, by [Theorem 5.3](#), we see that Γ is the index 2 orientation-preserving subgroup of the group Γ_{ref} of reflections in a hyperbolic hexagon Q with angles $(0, \frac{\pi}{2}, \frac{\pi}{2}, 0, \frac{\pi}{2}, \frac{\pi}{2})$. The claim is that this hyperbolic reflection group Γ_{ref} is arithmetic, maximal (as an arithmetic reflection group), and noncongruence.

Claim 1. Γ_{ref} is arithmetic.

Proof. Note that the finite area of \mathbb{H}^2/Γ implies that the index $[N(\Gamma_0(11)) : \Gamma]$ is finite. Hence, since $N(\Gamma_0(11))$ is arithmetic, we see that Γ is also arithmetic, from which it follows that Γ_{ref} is arithmetic. □

Claim 2. Γ_{ref} is a maximal reflection group.

Proof. If Γ_{ref} were not maximal, it would be properly contained in another reflection group H_{ref} , which is therefore also arithmetic. Let $H < H_{\text{ref}}$ denote the orientation-preserving index 2 subgroup. Note that then we have $\Gamma < H$. Since Γ and H are both arithmetic Fuchsian groups of genus zero, they are contained in a common maximal, arithmetic, genus zero Fuchsian group M from the appropriate list in [Long et al. 2006]. As we saw above, Γ is contained in the normalizer $N(\Gamma_0(11))$, and by area considerations we find that $[N(\Gamma_0(11)) : \Gamma] = 2$. Further, Γ cannot be contained in any other of these maximal arithmetic groups; to see this, observe that if $n \neq 11$ then, if we pick some nonzero integer b coprime to n , we may find integers a, d such that $\begin{pmatrix} a & b \\ n & d \end{pmatrix} \in \Gamma_0(n)$. We then have

$$\gamma_2 \begin{pmatrix} a & b \\ n & d \end{pmatrix} \gamma_2 = \begin{pmatrix} 0 & -\frac{1}{\sqrt{11}} \\ \sqrt{11} & 0 \end{pmatrix} \begin{pmatrix} a & b \\ n & d \end{pmatrix} \begin{pmatrix} 0 & -\frac{1}{\sqrt{11}} \\ \sqrt{11} & 0 \end{pmatrix} = \begin{pmatrix} -d & \frac{n}{11} \\ 11b & -a \end{pmatrix}.$$

We wish to show that this does not belong to $\Gamma_0(n)$. If n is not divisible by 11 this is clear, so suppose $n \geq 22$ is a multiple of 11. Then, by construction, b is coprime to 11, and so $11b$ is not divisible by n . This shows that γ_2 cannot belong to any normalizer $N(\Gamma_0(n))$ except $N(\Gamma_0(11))$.

It remains to verify that we cannot have $H = M = N(\Gamma_0(11))$. Construction of the Ford domain for $N(\Gamma_0(11))$ yields the generating set

$$\begin{aligned} \delta_1 &= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \delta_2 = \begin{pmatrix} 0 & \frac{-1}{\sqrt{11}} \\ \sqrt{11} & 0 \end{pmatrix}, \quad \delta_3 = \begin{pmatrix} \sqrt{11} & \frac{5}{\sqrt{11}} \\ 2\sqrt{11} & \sqrt{11} \end{pmatrix}, \\ \delta_4 &= \begin{pmatrix} \sqrt{11} & \frac{-4}{\sqrt{11}} \\ 3\sqrt{11} & -\sqrt{11} \end{pmatrix}, \quad \delta'_4 = \begin{pmatrix} -\sqrt{11} & \frac{-4}{\sqrt{11}} \\ 3\sqrt{11} & \sqrt{11} \end{pmatrix}. \end{aligned}$$

The Ford domain corresponding to these generators is shown in Figure 3.




Figure 3. A Ford domain for $N(\Gamma_0(11))$. All sides except those marked are paired with their opposites.

The fact that three of the generating elements are involutions, which pair adjacent sides of the Ford domain, precludes $N(\Gamma_0(11))$ from possessing a DF domain. By [Theorem 5.4](#), this also precludes it from being an index 2 subgroup of a reflection group. Thus Γ_{ref} is maximal. \square

Remark. Since Γ_{ref} is an arithmetic maximal hyperbolic reflection group, one would expect to find it in existing lists of such groups. This example appears to be the lattice 2-fill($L_{26,1}$) in Allcock’s enumeration [2010] of rank-3 reflective Lorentzian lattices, which would correspond to the case $N = 26$ in [Nikulin 2000, Table 1]. If one could show Γ_{ref} is indeed this lattice, this would provide an alternative proof that it is maximal arithmetic; however, we omit this at present, as the proofs given above suffice for our purposes.

Claim 3. Γ_{ref} is not congruence.

Proof. Suppose Γ_{ref} is congruence. Then it contains some principal congruence subgroup $\Gamma(n)$. These groups all belong to the modular group, so $\Gamma(n)$ is contained in $G = \Gamma \cap \text{PSL}_2(\mathbb{Z})$. By Wohlfahrt’s theorem (see [Newman 1972], p. 149), G contains $\Gamma(n)$ for n equal to the level of G , that is, the smallest natural number such that G contains the normal closure of

$$T^n = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^n = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$$

in $\text{PSL}_2(\mathbb{Z})$.

Subclaim. The level of G is 11.

Proof of subclaim. The group $G = \Gamma \cap \text{PSL}_2(\mathbb{Z}) = \Gamma \cap \Gamma_0(11)$ is not equal to Γ , by the presence of the nonintegral elements γ_2 and γ_3 . However, it contains the matrices

$$\beta_1 = \gamma_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \beta_2 = \gamma_2 \gamma_1^{-1} \gamma_2 = \begin{pmatrix} 1 & 0 \\ 11 & 1 \end{pmatrix}, \quad \beta_3 = \gamma_2 \gamma_3 = \begin{pmatrix} -2 & -1 \\ 11 & 5 \end{pmatrix},$$

$$\beta_4 = \gamma_2 \gamma_3^{-1} = \begin{pmatrix} 2 & -1 \\ 11 & -5 \end{pmatrix}, \quad \beta_5 = \gamma_4 = \begin{pmatrix} 10 & 3 \\ 33 & 10 \end{pmatrix}, \quad \beta_6 = \gamma_5 = \begin{pmatrix} 23 & 8 \\ 66 & 23 \end{pmatrix}.$$

The isometric circles of these elements and their inverses are shown in [Figure 4](#). Notice that the isometric circles centered at $\frac{2}{11}$ and $-\frac{2}{11}$ are paired with those at $\frac{5}{11}$ and $-\frac{5}{11}$ respectively; with these four circles excepted, each other side is paired with its reflection in the line $x = 0$. There are four equivalence classes of ideal points: these classes are $\{\infty\}$, $\{0\}$, $\{\frac{1}{3}, -\frac{1}{3}\}$, $\{\frac{4}{11}, \frac{3}{11}, -\frac{3}{11}, -\frac{4}{11}\}$. All four finite vertices belong to the same cycle, and their angles are $\pi/3$ at $x = \pm\frac{1}{2}$, and $2\pi/3$ at $x = \pm\frac{3}{22}$, giving angle sum 2π .




Figure 4. A Ford domain for G .

The region P_G of \mathbb{H}^2 bounded by these circles and the lines $x = -\frac{1}{2}$ and $x = \frac{1}{2}$ has area 8π and contains a Ford domain for G . This is enough for us to conclude that it is a Ford domain for G : since G is a proper subgroup of Γ , of finite index due to the finite area of P_G , G must have coarea a multiple $4m\pi$ of 4π , where $m = [\Gamma : G] > 1$. That the area of P_G is 8π tells us that $m \leq 2$, and hence that in fact $m = 2$. So G has index 2 in Γ and index 24 in $\text{PSL}_2(\mathbb{Z})$, and the list above is a generating set.

To prove the Subclaim, we need to show that given any $\varphi \in \text{PSL}_2(\mathbb{Z})$, we have that $\varphi T^{11} \varphi^{-1} \in G$. If φ fixes ∞ this is clear, so suppose $\varphi(\infty) \neq \infty$. Topologically, \mathbb{H}^2/G is a torus with four cusps, with the cusp orbits in $\mathbb{Q} \cup \{\infty\}$ represented by $0, \infty, \frac{1}{3}$ and $\frac{3}{11}$. Therefore $\varphi(\infty)$ is G -equivalent to exactly one of these four points; let $g \in G$ be such that $g^{-1}\varphi(\infty)$ is this point. We observe that $T^{11} \in G$; we also find that

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 11 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -11 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 11 & 1 \end{pmatrix}^{-1} \in G$$

is a parabolic element fixing 0, that

$$\begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} 1 & 11 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} = \begin{pmatrix} -32 & 11 \\ -99 & 34 \end{pmatrix} = \begin{pmatrix} 10 & 3 \\ 33 & 10 \end{pmatrix} \begin{pmatrix} -23 & 8 \\ 66 & -23 \end{pmatrix} \in G$$

is a parabolic element fixing $\frac{1}{3}$, and that

$$\begin{aligned} \begin{pmatrix} 3 & 1 \\ 11 & 4 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -4 & 1 \\ 11 & -3 \end{pmatrix} &= \begin{pmatrix} -32 & 9 \\ -121 & 34 \end{pmatrix} \\ &= \begin{pmatrix} 2 & -1 \\ 11 & -5 \end{pmatrix} \begin{pmatrix} 23 & 8 \\ 66 & 23 \end{pmatrix} \begin{pmatrix} -5 & -1 \\ 11 & -2 \end{pmatrix} \begin{pmatrix} -10 & 3 \\ 33 & -10 \end{pmatrix} \in G \end{aligned}$$

is a parabolic element fixing $\frac{3}{11}$. Note that in this last case, G also contains a conjugate of T^{11} fixing $\frac{3}{11}$, by taking the 11th power of the given element. Thus there exists a conjugate α of T^{11} such that $\alpha \in G$ and α fixes $g^{-1}\varphi(\infty)$. The element $g.\alpha.g^{-1} \in G$ is therefore a parabolic element, conjugate in $\mathrm{PSL}_2(\mathbb{Z})$ to T^{11} , with parabolic fixed point at $\varphi(\infty)$. We wish to show that $g.\alpha.g^{-1} = \varphi T^{11}\varphi^{-1}$. Since the former element is known to be a conjugate of T^{11} , we may alternatively write it as $\psi T^{11}\psi^{-1}$ for some $\psi \in \mathrm{PSL}_2(\mathbb{Z})$ with $\psi(\infty) = \varphi(\infty)$. Now $\psi^{-1}\varphi \in \mathrm{PSL}_2(\mathbb{Z})$ fixes ∞ and so must be a power of T ; in particular, $\psi^{-1}\varphi$ commutes with T . It follows that $\psi^{-1}\varphi T^{11}\varphi^{-1}\psi = T^{11}$ and therefore

$$g.\alpha.g^{-1} = \psi T^{11}\psi^{-1} = \varphi T^{11}\varphi^{-1}$$

as required. Thus G contains all elements of the form $\varphi T^{11}\varphi^{-1}$, and so the level of G is at most 11. To see that it is not smaller, observe that G does not contain any element of the form $\begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}$ for $t = 1, 2, \dots, 10$. \square

To complete the proof of [Claim 3](#), we note that by the Subclaim, G must contain $\Gamma(11)$. Computation in GAP Version 4.4.12 [[GAP 2008](#)] reveals that the core of G in $\mathrm{PSL}_2(\mathbb{Z})$, the largest normal subgroup of $\mathrm{PSL}_2(\mathbb{Z})$ contained in G , has index $k = 1351680 = 2^{13} \cdot 3 \cdot 5 \cdot 11$ in $\mathrm{PSL}_2(\mathbb{Z})$. Thus G cannot contain a normal subgroup of $\mathrm{PSL}_2(\mathbb{Z})$ of index (in $\mathrm{PSL}_2(\mathbb{Z})$) smaller than this constant. But all principal congruence subgroups are normal, and $[\mathrm{PSL}_2(\mathbb{Z}) : \Gamma(11)] = 660 < k$. From this contradiction we conclude that Γ_{ref} is not congruence. \square

Remark. [Hsu \[1996\]](#) gives a congruence test which can be applied to G . Since G has index 24 in $\mathrm{PSL}_2(\mathbb{Z})$, we obtain a representation in the symmetric group S_{24} . After expressing the known generators for G in terms of $L = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $R = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, we find

$$L = (2\ 4\ 9\ 15\ 8\ 5\ 11\ 13\ 7\ 3\ 6)(10\ 17\ 21\ 23\ 22\ 19\ 14\ 12\ 18\ 20\ 16)$$

and

$$R = (1\ 2\ 5\ 12\ 14\ 7\ 4\ 10\ 16\ 8\ 3)(9\ 17\ 19\ 13\ 11\ 18\ 21\ 24\ 22\ 20\ 15)$$

are both of order 11, also giving that the level of G is 11. Hsu's test is then that G is a congruence subgroup if and only if $(R^2 L^{-\frac{1}{2}})^3 = 1$, where $\frac{1}{2}$ is the multiplicative inverse of 2 mod 11, in this case equal to 6. Here we find that $R^2 L^{-6}$ has order 6, and so G is noncongruence.

Remark. The example given above is not the only arithmetic maximal reflection group which is not congruence. Of the 23 noncompact arithmetic maximal reflection groups which belong to $\mathrm{Isom}(\mathbb{H}^2)$, eight are not congruence. Furthermore, there exists a noncongruence example in $\mathrm{Isom}(\mathbb{H}^3)$. Further details will appear in the author's Ph.D. thesis.

7. Kleinian groups and DF domains

In this section, it will be shown that only one direction (the analogue of 5.4) of the main theorem holds when we consider Kleinian groups in the place of Fuchsian groups. This is because the added dimension gives new possibilities for the shape of the domains in question; in particular, they no longer have to glue up in a completely symmetrical way, although some symmetry remains. Examples will be given to demonstrate this flexibility, which extends as far as having nontrivial cuspidal cohomology. The discussion will be restricted to DF domains; as the above work demonstrates, it is not unreasonable to suppose that double Dirichlet domains share many similar properties.

Theorem 7.1. *Let $Q \subset \mathbb{H}^3$ be a finite sided, convex hyperbolic polyhedron with all dihedral angles integer submultiples of π , and let G be the discrete group of reflections in Q . Then G contains an index 2 Kleinian subgroup that admits a double Dirichlet domain (and a DF domain if Q has an ideal vertex).*

Proof. Suppose that Q is placed in upper half-space \mathbb{H}^3 such that one of its faces L is contained in a vertical plane. Let

$$G = \langle \tau_1, \dots, \tau_m, \tau_L \rangle$$

be a generating set for G . Let

$$\Gamma = \langle \tau_L \circ \tau_1, \dots, \tau_L \circ \tau_m \rangle$$

be the index 2 subgroup. Let $P = Q \cup \tau_L Q$. Let $w_0 = x_0 + y_0i + z_0j \in \dot{L}$, for $z_0 > 0$. The claim is that w_0 is a Dirichlet center for Γ . Fix a generator $\gamma_i = \tau_L \circ \tau_i$. Then the plane P_i fixed by τ_i bisects w_0 and $\tau_i(w_0)$, and so $\tau_L(P_i)$, which by construction is a face of P , bisects w_0 and $\gamma_i(w_0)$. □

The next result provides the first counterexamples of Theorem 5.3 by exhibiting Kleinian groups that admit DF domains and do not have index 2 in a reflection group.

Proposition 7.2. *Let Q be an all right hyperbolic polyhedron, with a vertex at ∞ , and all vertices ideal. Let G be the group of reflections in Q . Then G contains a subgroup of index 4 that admits a DF domain.*

Proof. Since Q is all right, the link of each vertex is a rectangle. Rotate Q in \mathbb{H}^3 so that the four faces adjacent to the vertex at ∞ , each lie above x - or y -lines in \mathbb{C} , where x -lines are parallel to the real axis, and y -lines are parallel to the imaginary axis. Let H be a side above a y -line, V a side above an x -line, and τ_H and τ_V the respective reflections. Let $P = (Q \cup \tau_H Q) \cup \tau_V(Q \cup \tau_H Q)$. Then P is the union of 4 copies of Q . Looking down from ∞ on the floor of P , label by A the nonvertical face adjacent to the top left vertex and to the vertical face opposite H . Label any

nonvertical faces adjacent to this face B . Proceed to label every nonvertical face A or B , with no two adjacent faces sharing the same label. The symmetry of P implies that this labeling is symmetric in both the x - and y -directions. Define the subgroup Γ as follows. Given a nonvertical side P_i of P , if P_i has label A , let the element $\tau_H \circ \tau_i$ belong to Γ ; if P_i has label B , let $\tau_V \circ \tau_i$ belong to Γ . If H' is the face of Q opposite H , and V' the face of Q opposite V , let $\tau_H \circ \tau_{H'}$ and $\tau_V \circ \tau_{V'}$ belong to Γ . Then P is a DF domain for Γ . \square

Remark. Given a group Γ constructed as in the above proof, note that Γ is not an index 2 subgroup of the group of reflections in the polyhedron $(Q \cup \tau_H Q)$. This is because the reflection τ_H will be absent from this group, preventing the construction of elements of Γ of the form $\tau_H \circ \tau_i$. The same is valid for the group of reflections in the polyhedron $(Q \cup \tau_V Q)$.

Since there is no direct analogue of [Theorem 5.3](#) for Kleinian groups, the question arises as to what, if anything, is implied about a Kleinian group by it having a DF domain. For example, one might ask whether such groups must have trivial cuspidal cohomology. The following example gives a Kleinian group which admits a DF domain, but which has nontrivial cuspidal cohomology; that is, there exists a nonperipheral homology class of infinite order in the first homology of the quotient space.

Example. Let $\Gamma < \text{PSL}_2(\mathbb{C})$ be generated by the matrices

$$\begin{pmatrix} 1 & 5 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 5i \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & \frac{-1}{\sqrt{2}} \\ \sqrt{2} & 0 \end{pmatrix}, \quad \begin{pmatrix} -\sqrt{2} & \frac{i}{\sqrt{2}} \\ -i\sqrt{2} & -\sqrt{2} \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & \frac{-1}{\sqrt{2}} \\ \sqrt{2} & 0 \end{pmatrix} \begin{pmatrix} 1 & \bar{a} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \sqrt{2}a & \sqrt{2}a\bar{a} - \frac{1}{\sqrt{2}} \\ \sqrt{2} & \sqrt{2}\bar{a} \end{pmatrix}$$

for each $a \in \{1, 2, 1+i, 2+i, 2i, 1+2i, 2+2i, 1-i, 2-i, -2i, 1-2i, 2-2i\}$, where \bar{a} is the complex conjugate of a . Then the isometric spheres of these matrices have centers at the Gaussian integers $\{x + iy \mid x, y \in \mathbb{Z}\}$ and radius $1/\sqrt{2}$. The square with vertices at $\pm \frac{5}{2} \pm \frac{5}{2}i$ is a Dirichlet domain for the action of Γ_∞ . Let P be the intersection of the exterior of all these isometric spheres with the chimney above the given rectangle. Then P is a DF domain for Γ , with Dirichlet center any point of $\overset{\circ}{P}$ above 0. Every dihedral angle of P is $\pi/2$. The quotient space \mathbb{H}^3/Γ has 14 boundary components; the cusp at ∞ gives a boundary torus, and each of the 13 cusp cycles in \mathbb{C} gives a $(2, 2, 2, 2)$ or a $(2, 4, 4)$ sphere. Thus the peripheral homology has rank 1. Computation using GAP gives that $H_1(\mathbb{H}^3/\Gamma)$ has rank 2, so there is infinite nonperipheral homology.

Remarks. (1) The cuspidal cohomology of this example has rank 1, but it can be modified to give examples where this rank is arbitrarily high.

(2) This example is arithmetic. To see this, consider the Picard group $\mathrm{PSL}_2(\mathbb{C}_1)$. This group contains as a finite index subgroup the congruence subgroup $\Gamma_0(2)$, where the lower left entry is a member of the ideal in \mathbb{C}_1 generated by 2. This congruence subgroup is normalized by the element

$$\gamma = \begin{pmatrix} 0 & \frac{-1}{\sqrt{2}} \\ \sqrt{2} & 0 \end{pmatrix}$$

and, since γ has order 2, $\Gamma_0(2)$ is an index 2 subgroup of the group H obtained by adding γ . The group Γ given in the example is a subgroup of H , of finite index due to the finite volume of the DF domain. In turn, H is commensurable with $\mathrm{PSL}_2(\mathbb{C}_1)$, as both share $\Gamma_0(2)$ as a subgroup of finite index.

(3) The quotient space of \mathbb{H}^3 by this group is not a manifold, so one can thus ask whether there exists another example which has nontrivial cuspidal cohomology, and which is additionally torsion free.

Although there does not appear to be a specific condition for a Kleinian group which is equivalent to having a DF domain, we can say something about a group that admits a DF domain. We cannot always decompose an orientation-preserving isometry of \mathbb{H}^3 into the composition of two reflections, but [Carathéodory \[1954\]](#) shows that we need at most four. If $\gamma \notin \Gamma_\infty$, these can be taken to be $\gamma = \gamma_4 \circ \gamma_3 \circ \gamma_2 \circ \gamma_1$, where γ_1 is reflection in the isometric sphere S_γ , γ_2 in the vertical plane R_γ bisecting S_γ and $S_{\gamma^{-1}}$, and $\gamma_4 \circ \gamma_3$ is rotation around the vertical axis through the north pole of $S_{\gamma^{-1}}$.

Theorem 7.3. *Suppose the Kleinian group Γ admits a DF domain P . Then the planes R_γ , for side pairings $\gamma \in \Gamma \setminus \Gamma_\infty$ of P , all intersect in a vertical axis. Furthermore, for each such γ , $\gamma_4 \circ \gamma_3 = 1$, and so each element of the corresponding generating set for Γ has real trace.*

Proof. Let P be a Ford domain. Suppose there is some side pairing γ such that $\gamma_4 \circ \gamma_3 \neq 1$. By considering the north pole of S_γ and its image, the north pole of $S_{\gamma^{-1}}$, we see that if P were a Dirichlet domain, its center w_0 would have to be in the plane R_γ . But given any such choice of w_0 , one can find a point $w \in P \cap S_\gamma$ such that $d(w_0, w) \neq d(w_0, \gamma(w))$. Thus P is not a Dirichlet domain. Since each $\gamma \in \Gamma \setminus \Gamma_\infty$ is then simply the composition of two reflections, it is the conjugate in $\mathrm{PSL}_2(\mathbb{C})$ of an element of $\mathrm{PSL}_2(\mathbb{R})$. It thus has real trace. Since it is assumed that any element of Γ_∞ is parabolic, these too have real trace.

Next suppose that the planes R_γ do not have a common intersection. Since we know that $\gamma_4 \circ \gamma_3 = 1$, for a given γ , the plane R_γ represents the set of potential

Dirichlet centers. If there is no common such center, P is not a Dirichlet domain. Thus if P is a DF domain, the planes R_γ have a common intersection. \square

The examples given earlier in this section give a flavor of the particular case with only two distinct, perpendicular planes R_γ . It is therefore possible for DF domains to be more complicated than this. This theorem provides a useful criterion for having a DF domain, which can be used to check known Ford domains. Observe that the vertical axis of intersection of the planes R_γ must correspond to a Dirichlet center for the action of Γ_∞ . Thus we see that the figure-8 knot group [Riley 1975], as well as the Whitehead link group and the group of the Borromean rings [Wielenberg 1978] do not admit DF domains. Furthermore, the groups obtained from a standard Ford domain in [Wielenberg 1981] cannot admit DF domains. Although in some cases, with the right choice of Ford domain, one can generate congruence subgroups of Bianchi groups using elements of real trace, the sides of the domain are identified in a way similar to the corresponding Fuchsian congruence subgroup, and so these groups seldom admit a DF domain.

Acknowledgments

I would like to express my gratitude to my advisor, Alan Reid, for his tremendous support, encouragement and guidance. I wish also to thank Daniel Allcock, Misha Belolipetsky and Hossein Namazi for additional valuable conversation and correspondence, and the referee for helpful comments and suggestions.

References

- [Agol et al. 2008] I. Agol, M. Belolipetsky, P. Storm, and K. Whyte, “Finiteness of arithmetic hyperbolic reflection groups”, *Groups, Geometry, and Dynamics* **2**:4 (2008), 481–498. [MR 2009m:20054](#) [Zbl 1194.22011](#)
- [Allcock 2010] D. Allcock, “The reflective Lorentzian lattices of rank 3”, preprint, 2010. To appear in *Mem. Amer. Math. Soc.* [arXiv 1010.0486](#)
- [Beardon 1983] A. F. Beardon, *The geometry of discrete groups*, Graduate Texts in Mathematics **91**, Springer, New York, 1983. [MR 85d:22026](#) [Zbl 0528.30001](#)
- [Belolipetsky 2011] M. Belolipetsky, “Finiteness theorems for congruence reflection groups”, *Transform. Groups* **16**:4 (2011), 939–954. [arXiv 1008.0934](#)
- [Carathéodory 1954] C. Carathéodory, *Theory of functions of a complex variable, I*, Chelsea, New York, 1954. [MR 15,612b](#) [Zbl 0055.30301](#)
- [Díaz and Ushijima 2009] R. Díaz and A. Ushijima, “On the properness of some algebraic equations appearing in Fuchsian groups”, *Topology Proceedings* **33** (2009), 81–106. [MR 2010i:30056](#) [Zbl 1179.30039](#)
- [Ford 1951] L. R. Ford, *Automorphic functions*, Chelsea, 1951. [Zbl 55.0810.04](#)
- [GAP 2008] *GAP - groups, algorithms, and programming*, Version 4.4.12, The GAP Group, St. Andrews, Fife, 2008, Available at <http://www.gap-system.org>.

- [Greenberg 1977] L. Greenberg, “Finiteness theorems for Fuchsian and Kleinian groups”, pp. 199–257 in *Discrete groups and automorphic functions* (Cambridge, 1975), edited by W. J. Harvey, Academic Press, London, 1977. [MR 58 #28483](#)
- [Grunewald and Schwermer 1981] F. Grunewald and J. Schwermer, “A nonvanishing theorem for the cuspidal cohomology of SL_2 over imaginary quadratic integers”, *Math. Ann.* **258**:2 (1981), 183–200. [MR 83c:10032](#) [Zbl 0458.20041](#)
- [Hsu 1996] T. Hsu, “Identifying congruence subgroups of the modular group”, *Proc. Amer. Math. Soc.* **124**:5 (1996), 1351–1359. [MR 96k:20100](#) [Zbl 0846.20051](#)
- [Long et al. 2006] D. D. Long, C. Maclachlan, and A. W. Reid, “Arithmetic Fuchsian groups of genus zero”, *Pure Appl. Math. Q.* **2**:2 (2006), 569–599. [MR 2007f:20092](#) [Zbl 1107.20037](#)
- [Maclachlan 1981] C. Maclachlan, “Groups of units of zero ternary quadratic forms”, *Proc. Roy. Soc. Edinburgh Sect. A* **88**:1–2 (1981), 141–157. [MR 83a:10030](#) [Zbl 0467.10018](#)
- [Maskit 1988] B. Maskit, *Kleinian groups*, Grundlehren der Mathematischen Wissenschaften **287**, Springer, Berlin, 1988. [MR 90a:30132](#) [Zbl 0627.30039](#)
- [Newman 1972] M. Newman, *Integral matrices*, Pure and Applied Mathematics **45**, Academic Press, New York, 1972. [MR 49 #5038](#) [Zbl 0254.15009](#)
- [Nikulin 2000] V. V. Nikulin, “О классификации гиперболических систем корней ранга три”, *Tr. Mat. Inst. Steklova* **230** (2000), 3–255. Translated as “On the classification of hyperbolic root systems of rank three” in *Proc. Steklov Inst. Math.* **230**:3 (2000), 1–241. [MR 2001m:17034](#) [Zbl 0997.17014](#)
- [Ratcliffe 2006] J. G. Ratcliffe, *Foundations of hyperbolic manifolds*, 2nd ed., Graduate Texts in Mathematics **149**, Springer, New York, 2006. [MR 2007d:57029](#) [Zbl 1106.51009](#)
- [Riley 1975] R. Riley, “A quadratic parabolic group”, *Math. Proc. Cambridge Philos. Soc.* **77**:2 (1975), 281–288. [MR 54 #542](#) [Zbl 0309.55002](#)
- [Wielenberg 1978] N. J. Wielenberg, “The structure of certain subgroups of the Picard group”, *Math. Proc. Cambridge Philos. Soc.* **84**:3 (1978), 427–436. [MR 80b:57010](#) [Zbl 0399.57005](#)
- [Wielenberg 1981] N. J. Wielenberg, “Hyperbolic 3-manifolds which share a fundamental polyhedron”, pp. 505–513 in *Riemann surfaces and related topics* (Stony Brook, NY, 1978), edited by I. Kra and B. Maskit, Ann. of Math. Stud. **97**, Princeton University Press, Princeton, NJ, 1981. [MR 82i:57012](#) [Zbl 0465.30039](#)

Received March 8, 2011.

GRANT S. LAKELAND
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TEXAS
AUSTIN, TX 78712
UNITED STATES

glakeland@math.utexas.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)

Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from [Periodicals Service Company](#), 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 255 No. 2 February 2012

On the local Langlands correspondences of DeBacker–Reeder and Reeder for $GL(\ell, F)$, where ℓ is prime	257
MOSHE ADRIAN	
<i>R</i> -groups and parameters	281
DUBRAVKA BAN and DAVID GOLDBERG	
Finite-volume complex-hyperbolic surfaces, their toroidal compactifications, and geometric applications	305
LUCA FABRIZIO DI CERBO	
Character analogues of Ramanujan-type integrals involving the Riemann ζ -function	317
ATUL DIXIT	
Spectral theory for linear relations via linear operators	349
DANA GHEORGHE and FLORIAN-HORIA VASILESCU	
Homogeneous links and the Seifert matrix	373
PEDRO M. GONZÁLEZ MANCHÓN	
Quantum affine algebras, canonical bases, and q -deformation of arithmetical functions	393
HENRY H. KIM and KYU-HWAN LEE	
Dirichlet–Ford domains and arithmetic reflection groups	417
GRANT S. LAKELAND	
Formal equivalence of Poisson structures around Poisson submanifolds	439
IOAN MĂRCUȚ	
A regularity theorem for graphic spacelike mean curvature flows	463
BENJAMIN STUART THORPE	
Analogues of level- N Eisenstein series	489
HIROFUMI TSUMURA	