

*Pacific
Journal of
Mathematics*

**RESONANT SOLUTIONS AND TURNING POINTS
IN AN ELLIPTIC PROBLEM
WITH OSCILLATORY BOUNDARY CONDITIONS**

ALFONSO CASTRO AND ROSA PARDO

Volume 257 No. 1

May 2012

RESONANT SOLUTIONS AND TURNING POINTS IN AN ELLIPTIC PROBLEM WITH OSCILLATORY BOUNDARY CONDITIONS

ALFONSO CASTRO AND ROSA PARDO

We consider the elliptic equation $-\Delta u + u = 0$ with nonlinear boundary conditions $\partial u / \partial n = \lambda u + g(\lambda, x, u)$, where the nonlinear term g is oscillatory and satisfies $g(\lambda, x, s)/s \rightarrow 0$ as $|s| \rightarrow 0$. We provide sufficient conditions on g for the existence of sequences of resonant solutions and turning points accumulating to zero.

1. Introduction

This work complements the study initiated in [Arrieta et al. 2010] and [Castro and Pardo 2011] on the positive solutions to the following boundary-value problem

$$(1-1) \quad \begin{cases} -\Delta u + u = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = \lambda u + g(\lambda, x, u) & \text{on } \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$ is a bounded and sufficiently smooth domain, $N \geq 2$, λ is a real parameter, $g(\lambda, x, s) = o(s)$ as $s \rightarrow 0$ and g is oscillatory. A typical example of such a g is

$$(1-2) \quad g(x, s) := s^\alpha \left(\sin \left| \frac{s}{\Phi_1(x)} \right|^\beta + C \right) \quad \text{with } \alpha + \beta > 1, \quad \beta < 0,$$

where Φ_1 stands for the first eigenfunction of the Steklov eigenvalue problem

$$(1-3) \quad \begin{cases} -\Delta \Phi + \Phi = 0 & \text{in } \Omega, \\ \frac{\partial \Phi}{\partial n} = \sigma \Phi & \text{on } \partial\Omega. \end{cases}$$

Pardo is supported by the Spanish Ministerio de Ciencia e Innovación (MICINN) under Project MTM2009-07540, by UCM-BSCH, Spain, GR58/08, Grupo 920894 and also by the Programa Becas Complutense del Amo. This work was carried out during Pardo's sabbatical visit to the Department of Mathematics, Harvey Mudd College, whose hospitality she thanks.

MSC2010: primary 35B32, 35B34, 35B35, 35J25, 58J55; secondary 35J60, 35J65.

Keywords: turning points, resonance, stability, instability, multiplicity, Steklov eigenvalues, bifurcation, sublinear oscillating boundary conditions.




Figure 1. Bifurcation diagram of subcritical and supercritical solutions, containing infinitely many turning points and infinitely many resonant solutions. In all cases, $\beta = -0.35$.

The first eigenvalue σ_1 is simple and, due to Hopf's lemma, we may assume its eigenfunction Φ_1 to be strictly positive in $\overline{\Omega}$ and we take $\|\Phi_1\|_{L^\infty(\partial\Omega)} = 1$.

The case $\alpha + \beta < 1$, $\beta > 0$ was treated in [Arrieta et al. 2010; Castro and Pardo 2011]. Here we focus on the case $\alpha + \beta > 1$, $\beta < 0$, inside of the complementary range. The case with $\alpha < 1$ corresponds to a *bifurcation from infinity* phenomenon; see [Arrieta et al. 2007; 2009; 2010; Castro and Pardo 2011; Rabinowitz 1973]. In contrast, the case with $\alpha > 1$ corresponds to a *bifurcation from zero* phenomenon; see [Arrieta et al. 2007; Crandall and Rabinowitz 1971; Rabinowitz 1971].

The oscillatory situation is in principle more complex than the monotone one, since order techniques such as sub- and supersolutions are not applicable.

One novelty in problem (1-1) is that the parameter appears explicitly in the boundary condition. With respect to this parameter, we perform an analysis of the local bifurcation diagram of nonnegative solutions to (1-1), which turns out to be different from the case $\alpha < 1$ (see Figure 1 for $\alpha > 1$ and Figure 2 for $\alpha < 1$).

Throughout this paper we make the following assumptions:

- (H1) $g : \mathbb{R} \times \partial\Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function (i.e. $g = g(\lambda, x, s)$ is measurable in $x \in \Omega$, and continuous with respect to $(\lambda, s) \in \mathbb{R} \times \mathbb{R}$). Moreover, there exist $G_1 \in L^r(\partial\Omega)$ with $r > N - 1$ and continuous functions $\Lambda : \mathbb{R} \rightarrow \mathbb{R}^+$, and $U : \mathbb{R} \rightarrow \mathbb{R}^+$, satisfying

$$\begin{cases} |g(\lambda, x, s)| \leq \Lambda(\lambda)G_1(x)U(s) & \text{for all } (\lambda, x, s) \in \mathbb{R} \times \partial\Omega \times \mathbb{R}, \\ \limsup_{|s| \rightarrow 0} \frac{U(s)}{|s|^\alpha} < +\infty & \text{for some } \alpha > 1. \end{cases}$$




Figure 2. Bifurcation diagram in the case $\alpha = 0.5$, $\beta = -0.35$.

(H2) The partial derivative $g_s(\lambda, \cdot, \cdot)$ (where $g_s := \partial g / \partial s$) belongs to $C(\partial\Omega \times \bar{\Omega})$; moreover, $g_s(\cdot, \cdot, 0) = 0$ and there exist $F_1 \in L^r(\partial\Omega)$, with $r > N - 1$ and $\rho > 1$ such that

$$\frac{|g(\lambda, x, s) - sg_s(\lambda, x, s)|}{|s|^\rho} \leq F_1(x) \quad \text{as } \lambda \rightarrow \sigma_1,$$

for $x \in \partial\Omega$ and $s \leq \epsilon$ small enough.

Throughout this paper, by solutions to (1-1) we mean elements $u \in H^1(\Omega)$ such that

$$(1-4) \quad \int_{\Omega} (\nabla u \cdot \nabla v + uv) \, dx = \lambda \int_{\partial\Omega} uv \, d\sigma + \int_{\partial\Omega} g(\lambda, x, u)v \, d\sigma \quad \text{for all } v \in H^1(\Omega).$$

As proven in [Arrieta et al. 2007, Proposition 2.3], all such solutions are in the Holder space $C^\beta(\bar{\Omega})$ for some $\beta > 0$. Moreover, there exists a connected set of positive solutions of (1-1) known as a *branch bifurcating from zero*; see [Arrieta et al. 2007, Theorem 8.1]. We denote it by $\mathcal{C}^+ \subset \mathbb{R} \times C(\bar{\Omega})$, and recall that for $(\lambda, u_\lambda) \in \mathcal{C}^+$

$$u = s\Phi_1 + w, \quad \text{with } w = o(|s|) \quad \text{and} \quad |\sigma_1 - \lambda| = o(1) \quad \text{as } |s| \rightarrow 0.$$

Definition 1.1. A solution (λ^*, u^*) of (1-1) in the branch of solutions $\mathcal{C}^+ \subset \mathbb{R} \times C(\bar{\Omega})$ is called a *turning point* if there is a neighborhood W of (λ^*, u^*) in $\mathbb{R} \times C(\bar{\Omega})$ such that, either $W \cap \mathcal{C}^+ \subset [\lambda^*, \infty) \times C(\bar{\Omega})$ or $W \cap \mathcal{C}^+ \subset (-\infty, \lambda^*] \times C(\bar{\Omega})$.

Our goal is to give conditions on the nonlinear oscillatory term g that guarantee the existence of sequences accumulating to zero of *subcritical* solutions (i.e., for

values of the parameter $\lambda < \sigma_1$), *supercritical* solutions (i.e., for $\lambda > \sigma_1$), *resonant* solutions (i.e., for $\lambda = \sigma_1$), and turning points.

Our main result, Theorem 1.3 below, is exemplified by the case in which g is given by (1-2). In fact we have:

Theorem 1.2. *Assume that g is given by (1-2) with $\beta < 0$. If*

$$|C| < 1 \quad \text{and} \quad \alpha + \beta > 1,$$

then in any neighborhood of the bifurcation point $(\sigma_1, 0)$ in $\mathbb{R} \times C(\overline{\Omega})$, the branch \mathcal{C}^+ of positive solutions of (1-1) contains a sequence of subcritical solutions, a sequence of supercritical solutions, a sequence of turning points, and a sequence of resonant solutions.

The proof of this follows directly from the next theorem.

Theorem 1.3. *Assume the nonlinearity g satisfies hypotheses (H1) and (H2). Assume also that*

$$(1-5) \quad \left| \frac{g(\lambda, x, s) - g(\sigma_1, x, s)}{|s|^\alpha} \right| \rightarrow 0 \quad \text{as} \quad \lambda \rightarrow \sigma_1, \quad s \rightarrow 0$$

pointwise in x .

Let $G : \mathbb{R} \times C(\overline{\Omega}) \rightarrow \mathbb{R}$ be defined by

$$(1-6) \quad G(\lambda, u) := \int_{\partial\Omega} \frac{ug(\lambda, \cdot, u)}{|u|^{1+\alpha}} \Phi_1^{1+\alpha}.$$

If there exist sequences $\{s_n\}, \{s'_n\}$ converging to 0^+ , such that

$$(1-7) \quad \lim_{n \rightarrow +\infty} G(\sigma_1, s'_n \Phi_1) < 0 < \lim_{n \rightarrow +\infty} G(\sigma_1, s_n \Phi_1),$$

then:

(i) *For sufficiently large $n \gg 1$, if (λ, u) is a solution of (1-1) with*

$$P(u) := \frac{\int_{\partial\Omega} u \Phi_1}{\int_{\partial\Omega} \Phi_1^2} = s_n,$$

then (λ, u) is subcritical. Similarly, if $P(u) = s'_n$ it is supercritical. Consequently, there exist two sequences of solutions of (1-1), $\{(\lambda_n, u_n)\}$ and $\{(\lambda'_n, u'_n)\}$ converging to $(\sigma_1, 0)$ as $n \rightarrow \infty$, one of them subcritical, $\lambda_n < \sigma_1$, and the other supercritical, $\lambda'_n > \sigma_1$.

(ii) *There is a sequence converging to zero of turning points $\{(\lambda_n^*, u_n^*)\}$ such that*

$$\lambda_n^* \rightarrow \sigma_1 \quad \text{and} \quad \|u_n^*\|_{L^\infty(\partial\Omega)} \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty.$$

In fact, we can always choose two subsequences of turning points, one of them subcritical, $\lambda_{2n+1}^ < \sigma_1$, and the other supercritical, $\lambda_{2n}^* > \sigma_1$.*

(iii) *There is a sequence converging to zero of resonant solutions; i.e., there are infinitely many solutions $\{(\sigma_1, \tilde{u}_n)\}$ of (1-1) with $\|\tilde{u}_n\|_{L^\infty(\partial\Omega)} \rightarrow 0$.*

The behavior of positive solutions to (1-1) bifurcating from $(\sigma_1, 0)$ described in Theorems 1.2 and 1.3 is similar to that of the solutions bifurcating from (σ_1, ∞) for the sublinear problem; see [Arrieta et al. 2010] for details.

The complex nature of the nonlinearity in (1-2), makes an exhaustive analysis of the global bifurcation diagram outside the scope of this work.

In [Korman 2008] the author considers in the case $\alpha = 1$ $\beta = 1$. He assumes either $N = 1$ or Ω to be a ball and the nonlinearity to be bounded by a constant small enough. He obtains what he calls an oscillatory bifurcation. We refer the reader to [García-Melián et al. 2009] for related problems with nonlinear boundary conditions.

Organization of the paper. Section 2 contains the proof of our main result, giving sufficient conditions for having subcritical, supercritical, and resonant solutions. Section 3 presents two examples; explicit resonant solutions for the oscillatory nonlinearity (1-2) and the one-dimensional case.

2. Subcritical, supercritical and resonant solutions

In this section we give sufficient conditions for the existence of a branch of solutions to (1-1) bifurcating from zero which is neither *subcritical* ($\lambda < \sigma_1$), nor *supercritical*, ($\lambda < \sigma_1$). From this, we conclude the existence of infinitely many *turning points*, see Definition 1.1, and an infinite number of solutions for the resonant problem, i.e. for $\lambda = \sigma_1$. This is achieved in Theorem 1.3

At this step, we analyze when the parameter may cross the first Steklov eigenvalue. To do that, we look at the asymptotic growth rate of the nonlinear term

$$(2-1) \quad \underline{\mathbf{G}}_{0^+} := \int_{\partial\Omega} \liminf_{(\lambda, s) \rightarrow (\sigma_1, 0)} \frac{sg(\lambda, \cdot, s)}{|s|^{1+\alpha}} \Phi_1^{1+\alpha}$$

for $\alpha > 1$. Changing \liminf to \limsup we define the number $\overline{\mathbf{G}}_{0^+}$. If $\underline{\mathbf{G}}_{0^+} > 0$ then \mathcal{C}^+ is subcritical, and if $\overline{\mathbf{G}}_{0^+} < 0$ then \mathcal{C}^+ is supercritical in a neighborhood of $(\sigma_1, 0)$ See [Arrieta et al. 2009, Theorems 3.4 and 3.5] for the bifurcation from infinity case. In this paper we consider nonlinearities for which

$$\underline{\mathbf{G}}_{0^+} < 0 < \overline{\mathbf{G}}_{0^+}.$$

We shall argue as in [Arrieta et al. 2010] for the bifurcation from infinity case. To determine whether a sequence of solutions (λ_n, u_n) is subcritical or supercritical,

one must check the sign of

$$(2-2) \quad \liminf_{n \rightarrow \infty} G(\lambda_n, u_n) \quad \text{and} \quad \limsup_{n \rightarrow \infty} G(\lambda_n, u_n),$$

where G is defined by (1-6). This is done in Lemma 2.3.

In Proposition 2.2, it is proved that when g is such that

$$|g(\lambda, x, s)| = O(|s|^\alpha) \quad \text{as } |s| \rightarrow 0 \text{ for some } \alpha > 1,$$

then the solutions in \mathcal{C}^\pm can be described as

$$u_n = s_n \Phi_1 + w_n, \quad \text{where} \quad \int_{\partial\Omega} w_n \Phi_1 = 0 \quad \text{and} \quad w_n = O(|s_n|^\alpha) \text{ as } n \rightarrow 0.$$

We unveil the signs of the expressions in (2-2) by just looking at the signs of the expressions in (2-2) at $\lambda_n = \sigma_1$ and $u_n = s_n \Phi_1$. This is achieved in Lemma 2.4.

For this we first consider a family of linear Steklov problems with a variable nonhomogeneous term at the boundary h depending on the parameter λ

$$(2-3) \quad \begin{cases} -\Delta u + u = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = \lambda u + h(\lambda, x) & \text{on } \partial\Omega, \end{cases}$$

where $h(\lambda, \cdot) \in L^r(\partial\Omega)$, $r > N - 1$ and $\lambda \in (-\infty, \sigma_2)$.

We use the decomposition

$$L^r(\partial\Omega) = \text{span}[\Phi_1] \oplus \text{span}[\Phi_1]^\perp,$$

where

$$\text{span}[\Phi_1]^\perp := \left\{ u \in L^r(\partial\Omega) : \int_{\partial\Omega} u \Phi_1 = 0 \right\}.$$

For $h(\lambda, \cdot) \in L^r(\partial\Omega)$, with $r > N - 1$, we write

$$(2-4) \quad h(\lambda, \cdot) = a_1(\lambda) \Phi_1 + h_1(\lambda, \cdot),$$

with

$$a_1(\lambda) = \frac{\int_{\partial\Omega} h(\lambda, \cdot) \Phi_1}{\int_{\partial\Omega} \Phi_1^2}, \quad \int_{\partial\Omega} h_1(\lambda, \cdot) \Phi_1 = 0.$$

For $\lambda \neq \sigma_1$ the solution $u = u(\lambda)$ of (2-3) has a unique decomposition

$$(2-5) \quad u = \frac{a_1(\lambda)}{\sigma_1 - \lambda} \Phi_1 + w, \quad \text{where} \quad \int_{\partial\Omega} w \Phi_1 = 0,$$

and $w = w(\lambda) \in \text{span}[\Phi_1]^\perp$ solves the problem

$$(2-6) \quad \begin{cases} -\Delta w + w = 0 & \text{in } \Omega, \\ \frac{\partial w}{\partial n} = \lambda w + h_1(\lambda, x) & \text{on } \partial\Omega. \end{cases}$$

Note that in (2-6) $w(\lambda) \in \text{span}[\Phi_1]^\perp$ is also well defined for $\lambda = \sigma_1$. Moreover:

Lemma 2.1. *For each compact set $K \subset (-\infty, \sigma_2) \subset \mathbb{R}$ there exists a constant $C = C(K)$, independent of λ , such that*

$$\|w(\lambda)\|_{L^\infty(\partial\Omega)} \leq C \|h_1(\lambda, \cdot)\|_{L^r(\partial\Omega)} \quad \text{for any } \lambda \in K,$$

where $w \in \text{span}[\Phi_1]^\perp$ is the solution of (2-6) and $h_1 \in \text{span}[\Phi_1]^\perp$ is defined in (2-4).

Proof. See Lemma 3.1 of [Arrieta et al. 2010]. □

Now we turn our attention to the nonlinear problem (1-1). Recall that for solutions (λ, u) close to the bifurcation point $(\sigma_1, 0)$ we have

$$(2-7) \quad u = s\Phi_1 + w, \quad \text{where } w \in \text{span}[\Phi_1]^\perp$$

satisfies

$$(2-8) \quad w = o(s) \quad \text{as } s \rightarrow 0.$$

We define

$$(2-9) \quad P(u) := \frac{\int_{\partial\Omega} u(\cdot) \Phi_1}{\int_{\partial\Omega} \Phi_1^2}.$$

Next, we give sufficient conditions on the nonlinear term g in (1-1), for $w = O(|s|^\alpha)$ as $s \rightarrow 0$; compare (2-8). We restrict ourselves below to the branch of positive solutions; a completely analogous result holds for the branch of negative solutions. The next result is essentially Proposition 3.2 in [Arrieta et al. 2010] rewritten for $s \rightarrow 0$; we include the proof for completeness.

Proposition 2.2. *Assume g satisfies hypotheses (H1) and (H2). There exists an open set $\mathcal{O} \subset \mathbb{R} \times C(\bar{\Omega})$ of the form $\mathcal{O} = \{(\lambda, u) : |\lambda - \sigma_1| < \delta_0, \|u\|_{L^\infty(\Omega)} < s_0\}$, for some δ_0 and s_0 , satisfying these conditions:*

- (i) *There exists a constant C_1 independent of λ such that, if $(\lambda, u) \in \mathcal{C}^+ \cap \mathcal{O}$ and $(\lambda, u) \neq (\sigma_1, 0)$ then $u = s\Phi_1 + w$, where $s > 0$, $w \in \text{span}[\Phi_1]^\perp$ and*

$$\|w\|_{L^\infty(\partial\Omega)} \leq C_1 \|G_1\|_{L^r(\partial\Omega)} |s|^\alpha \quad \text{as } |s| \rightarrow 0.$$

- (ii) *There exists a constant $S_0 > 0$ such that for all $|s| \leq S_0$ there exists (λ, u) in $\mathcal{C}^+ \cap \mathcal{O}$ satisfying $u = s\Phi_1 + w$, with $w \in \text{span}[\Phi_1]^\perp$.*

- (iii) *Moreover, for any $(\lambda, u) \in \mathcal{C}^+ \cap \mathcal{O}$, $u = s\Phi_1 + w$, with $w \in \text{span}[\Phi_1]^\perp$,*

$$|\sigma_1 - \lambda| \leq C_2 |s|^{\alpha-1} \quad \text{as } |s| \rightarrow 0,$$

with C_2 independent of λ ; in fact,

$$C_2 = \frac{2\|G_1\|_{L^1(\partial\Omega)}}{\int_{\partial\Omega} \Phi_1^2}.$$

Proof. From (2-7) and (2-8), we have $\Phi_1 + w/s \rightarrow \Phi_1$ as $s \rightarrow 0$ in $L^\infty(\partial\Omega)$. Together with (H1) and Lemma 2.1, this implies that $\|w\|_{L^\infty(\partial\Omega)} \leq C|s|^\alpha$ as $s \rightarrow 0$. This proves part (i).

To prove part (ii) note that $\mathcal{C}^+ \cap \mathcal{C}$ is connected. Hence, using the decomposition in (2-7), we have $u = s\Phi_1 + w$ with $w \in \text{span}[\Phi_1]^\perp$. Since the projection P is continuous, by (2-9), the set

$$\{s \in \mathbb{R} : (1-1) \text{ has a solution of the form } u = s\Phi_1 + w \text{ and } w \in [\text{span}[\Phi_1]^\perp]\}$$

contains an interval in \mathbb{R} containing zero.

To prove part (iii) we observe that if (λ, u) is a solution of (1-1), $u = s\Phi_1 + w$, with $w \in \text{span}[\Phi_1]^\perp$, multiplying the equation by the first Steklov eigenfunction $\Phi_1 > 0$ and integrating by parts we obtain,

$$(\sigma_1 - \lambda)s \int_{\partial\Omega} \Phi_1^2 = \int_{\partial\Omega} g(\lambda, x, s\Phi_1 + w)\Phi_1.$$

Taking into account that

$$\frac{|g(\lambda, x, s\Phi_1 + w)|}{|s|} = \frac{|g(\lambda, x, s\Phi_1 + w)|}{|s\Phi_1 + w|} \left| \Phi_1 + \frac{w}{s} \right| \rightarrow 0 \quad \text{as } s \rightarrow 0$$

we get $\lambda \rightarrow \sigma_1$ as $s \rightarrow 0$.

Moreover, from (H1), we obtain that

$$\begin{aligned} |g(\lambda, x, s\Phi_1 + w)| &= |s|^\alpha \frac{|g(\lambda, x, s\Phi_1 + w)|}{|s\Phi_1 + w|^\alpha} \left| \Phi_1 + \frac{w}{s} \right|^\alpha \\ &\leq C|s|^\alpha G_1(x) \left| \Phi_1 + \frac{w}{s} \right|^\alpha, \end{aligned}$$

and therefore

$$|\sigma_1 - \lambda| \leq C \frac{|s|^{\alpha-1}}{\int_{\partial\Omega} \Phi_1^2} \int_{\partial\Omega} G_1(x) \left| \Phi_1 + \frac{w}{s} \right|^\alpha \Phi_1 \leq C\|G_1\|_{L^r(\partial\Omega)} |s|^{\alpha-1},$$

which ends the proof. \square

Our next result is essentially Lemma 3.1 in [Arrieta et al. 2009] rewritten for $s \rightarrow 0$. It allows us to estimate $\sigma_1 - \lambda_n$ as λ_n converges σ_1 .

Lemma 2.3. *Assume the nonlinearity g satisfies hypotheses (H1) and (H2). Let (λ_n, u_n) be a sequence of solutions of (1-1) with $\lambda_n \rightarrow \sigma_1$ and $\|u_n\|_{L^\infty(\partial\Omega)} \rightarrow 0$.*

If $u_n > 0$ then

$$\begin{aligned}
 (2-10) \quad \frac{\underline{G}_{0+}}{\int_{\partial\Omega} \Phi_1^2} &\leq \frac{1}{\int_{\partial\Omega} \Phi_1^2} \liminf_{n \rightarrow \infty} G(\lambda_n, u_n) \\
 &\leq \liminf_{n \rightarrow \infty} \frac{\sigma_1 - \lambda_n}{\|u_n\|_{L^\infty(\partial\Omega)}^{\alpha-1}} \leq \limsup_{n \rightarrow \infty} \frac{\sigma_1 - \lambda_n}{\|u_n\|_{L^\infty(\partial\Omega)}^{\alpha-1}} \\
 &\leq \frac{1}{\int_{\partial\Omega} \Phi_1^2} \limsup_{n \rightarrow \infty} G(\lambda_n, u_n) \leq \frac{\overline{G}_{0+}}{\int_{\partial\Omega} \Phi_1^2}.
 \end{aligned}$$

A similar statement is obtained for the case $u_n < 0$, just replacing \underline{G}_{0+} by \underline{G}_{0-} and \overline{G}_{0+} by \overline{G}_{0-} .

Proof. We show that $u_n > 0$; the other case has a similar proof. Consider a family of solutions u_n of (1-1) for $\lambda = \lambda_n$ with $\lambda_n \rightarrow \sigma_1$ and $0 < u_n \rightarrow 0$. Multiplying (1-1) by Φ_1 and integrating by parts, we get

$$(2-11) \quad (\sigma_1 - \lambda_n) \int_{\partial\Omega} u_n \Phi_1 = \int_{\partial\Omega} g(\lambda_n, x, u_n) \Phi_1.$$

But

$$\int_{\partial\Omega} g(\lambda_n, x, u_n) \Phi_1 = \|u_n\|_{L^\infty(\partial\Omega)}^\alpha \int_{\partial\Omega} \frac{g(\lambda_n, x, u_n)}{u_n^\alpha} \left(\frac{u_n}{\|u_n\|_{L^\infty(\partial\Omega)}} \right)^\alpha \Phi_1.$$

Taking into account the definition of $G(\lambda, u)$ in (1-6), we can write

$$\begin{aligned}
 \int_{\partial\Omega} \frac{g(\lambda_n, x, u_n)}{u_n^\alpha} \left(\frac{u_n}{\|u_n\|_{L^\infty(\partial\Omega)}} \right)^\alpha \Phi_1 \\
 = \int_{\partial\Omega} \frac{g(\lambda_n, x, u_n)}{u_n^\alpha} \left[\left(\frac{u_n}{\|u_n\|_{L^\infty(\partial\Omega)}} \right)^\alpha - \Phi_1^\alpha \right] \Phi_1 + G(\lambda_n, u_n).
 \end{aligned}$$

Moreover,

$$\int_{\partial\Omega} \frac{g(\lambda_n, x, u_n)}{u_n^\alpha} \left[\left(\frac{u_n}{\|u_n\|_{L^\infty(\partial\Omega)}} \right)^\alpha - \Phi_1^\alpha \right] \Phi_1 \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

because $u_n/\|u_n\|_{L^\infty(\partial\Omega)} \rightarrow \Phi_1$ uniformly in $\partial\Omega$.

But, firstly from the above, secondly from Fatou's lemma, and thirdly from definition of \underline{G}_{0+} ,

$$\begin{aligned}
 (2-12) \quad \liminf_{n \rightarrow \infty} \int_{\partial\Omega} \frac{g(\lambda_n, x, u_n)}{u_n^\alpha} \left(\frac{u_n}{\|u_n\|_{L^\infty(\partial\Omega)}} \right)^\alpha \Phi_1 \\
 \geq \liminf_{n \rightarrow \infty} G(\lambda_n, u_n) \geq \int_{\partial\Omega} \liminf_{n \rightarrow \infty} \frac{g(\lambda_n, x, u_n)}{u_n^\alpha} \Phi_1^{1+\alpha} \\
 \geq \underline{G}_{0+}.
 \end{aligned}$$

Dividing both sides of (2-11) by $\|u_n\|_{L^\infty(\partial\Omega)}^\alpha$ and passing to the limit we obtain the first two inequalities in the chain (2-10). The third inequality in the chain is trivial and the last two are obtained in a similar manner. \square

Let $\{s_n\}$ and $\{s'_n\}$ satisfy

$$(2-13) \quad -\infty < \lim_{n \rightarrow +\infty} G(\sigma_1, s'_n \Phi_1) < 0 < \lim_{n \rightarrow +\infty} G(\sigma_1, s_n \Phi_1) < \infty.$$

In order to prove Theorem 1.3, we show that the signs in (2-2) can be deduced from those of (2-13). This is stated in the following result, which is a slight variation of [Arrieta et al. 2010, Lemma 3.3].

Lemma 2.4. *Assume that g satisfies hypotheses (H1), (H2), and (1-5).*

If $(\lambda_n, s_n) \rightarrow (\sigma_1, 0)$ and there exists a constant C such that $\|w_n\|_{L^\infty(\partial\Omega)} \leq C|s_n|^\alpha$ for all $n \rightarrow 0$, then

$$\liminf_{n \rightarrow +\infty} G(\lambda_n, s_n \Phi_1 + w_n) \geq \liminf_{n \rightarrow +\infty} G(\sigma_1, s_n \Phi_1),$$

where G is given by (1-6). Similarly,

$$\limsup_{n \rightarrow +\infty} G(\lambda_n, s_n \Phi_1 + w_n) \leq \limsup_{n \rightarrow +\infty} G(\sigma_1, s_n \Phi_1).$$

Proof. Throughout this proof, C denotes several constants depending only on (Ω, g) . Given $\varepsilon > 0$, assume that $|(\lambda_n, s_n) - (\sigma_1, 0)| < \varepsilon$.

By the mean value theorem we have

$$(2-14) \quad \begin{aligned} g(\lambda_n, x, s_n \Phi_1 + w_n) - g(\lambda_n, x, s_n \Phi_1) &= w_n \int_0^1 g_s(\lambda_n, \cdot, s_n \Phi_1 + \tau w_n) d\tau \\ &\leq \|w_n\|_{L^\infty(\partial\Omega)} \sup_{\tau \in [0,1]} \|g_s(\lambda_n, \cdot, s_n \Phi_1 + \tau w_n)\|_{L^\infty(\partial\Omega)}. \end{aligned}$$

Therefore

$$(2-15) \quad \begin{aligned} \int_{\partial\Omega} \left| g(\lambda_n, x, s_n \Phi_1 + w_n) - g(\lambda_n, x, s_n \Phi_1) \right| \Phi_1 dx &\leq \|w_n\|_{L^\infty(\partial\Omega)} \int_{\partial\Omega} \sup_{\tau \in [0,1]} \|g_s(\lambda_n, \cdot, s_n \Phi_1 + \tau w_n)\|_{L^\infty(\partial\Omega)} \Phi_1 dx \\ &\leq |\partial\Omega| \|w_n\|_{L^\infty(\partial\Omega)} \sup_{\tau \in [0,1]} \|g_s(\lambda_n, \cdot, s_n \Phi_1 + \tau w_n)\|_{L^\infty(\partial\Omega)}. \end{aligned}$$

By hypotheses (H1) and (H2), for all $x \in \partial\Omega$,

$$(2-16) \quad \begin{aligned} \frac{|g_s(\lambda_n, x, s)|}{|s|^{\gamma-1}} &\leq |s|^{\rho-\gamma} F_1(x) + C|s|^{\alpha-\gamma} G_1(x) \max\{\Lambda(\lambda_n), n \geq 1\} =: D_1(x), \end{aligned}$$

for n large, and $\gamma = \min\{\rho, \alpha\} > 1$. Hence, $D_1 \in L^r(\partial\Omega)$ with $r > N - 1$ and

$$(2-17) \quad \sup_{|s| \leq 1/n} |g_s(\lambda_n, x, s)| \leq D_1(x) \left(\frac{1}{n}\right)^{\gamma-1}, \quad \text{with } \gamma > 1.$$

Since $\|w_n\|_{L^\infty(\partial\Omega)} = O(|s_n|^\alpha)$, we obtain from (2-15) and (2-17)

$$(2-18) \quad \int_{\partial\Omega} \frac{|g(\lambda_n, \cdot, s_n \Phi_1 + w_n) - g(\lambda_n, \cdot, s_n \Phi_1)|}{|s_n|^\alpha} \Phi_1$$

$$\leq C \sup_{\tau \in [0,1]} \|g_s(\lambda_n, \cdot, s_n \Phi_1 + \tau w_n)\|_{L^\infty(\partial\Omega)}$$

$$\leq C \sup_{|s| \leq 1/n} \|g_s(\lambda_n, \cdot, s)\|_{L^\infty(\partial\Omega)},$$

which tends to 0 as $n \rightarrow \infty$.

Therefore

$$\liminf_{n \rightarrow +\infty} \int_{\partial\Omega} \frac{s_n g(\lambda_n, \cdot, s_n \Phi_1 + w_n)}{|s_n|^{1+\alpha}} \Phi_1$$

$$\geq \lim_{n \rightarrow \infty} \int_{\partial\Omega} \frac{s_n g(\lambda_n, \cdot, s_n \Phi_1 + w_n) - s_n g(\lambda_n, \cdot, s_n \Phi_1)}{|s_n|^{1+\alpha}} \Phi_1$$

$$+ \liminf_{n \rightarrow +\infty} \int_{\partial\Omega} \frac{s_n g(\lambda_n, \cdot, s_n \Phi_1)}{|s_n|^{1+\alpha}} \Phi_1$$

$$= \liminf_{n \rightarrow +\infty} \int_{\partial\Omega} \frac{s_n g(\lambda_n, \cdot, s_n \Phi_1)}{|s_n|^{1+\alpha}} \Phi_1$$

$$= \liminf_{n \rightarrow +\infty} \int_{\partial\Omega} \frac{s_n g(\sigma_1, \cdot, s_n \Phi_1)}{|s_n|^{1+\alpha}} \Phi_1,$$

where we have used (2-18) and (1-5) respectively.

Now note that, multiplying and dividing by $|\Phi_1 + w_n/s_n|^\alpha$ the integrand of the left hand side above can be written as

$$\frac{s_n g(\lambda_n, \cdot, s_n \Phi_1 + w_n)}{|s_n|^{1+\alpha}} \Phi_1 = \frac{(s_n \Phi_1 + w_n) g(\lambda_n, \cdot, s_n \Phi_1 + w_n)}{|s_n \Phi_1 + w_n|^{1+\alpha}} \left| \Phi_1 + \frac{w_n}{s_n} \right|^\alpha.$$

Then, (H2) and the fact that $\Phi_1 + w_n/s_n \rightarrow \Phi_1$ in $L^\infty(\partial\Omega)$ concludes the proof. \square

Now we prove the first main result in this paper. Roughly speaking, it states that if there are a sequence of subcritical solutions and another of supercritical solutions, since the solution set is connected, there are infinitely many turning points and infinitely many resonant solutions. We prove the result for the positive branch. The same conclusions can be attained for the connected branch of negative solutions bifurcating from zero.

Proof of Theorem 1.3. From Proposition 2.2(ii), consider any two sequences of solutions of (1-1), such that $(\lambda_n, u_n) \rightarrow (\sigma_1, 0)$ and $(\lambda'_n, u'_n) \rightarrow (\sigma_1, 0)$ in \mathcal{C}^+ with

$$P(u_n) = \frac{\int_{\partial\Omega} u_n \Phi_1}{\int_{\partial\Omega} \Phi_1^2} = s_n \quad \text{and} \quad P(u'_n) = \frac{\int_{\partial\Omega} u'_n \Phi_1}{\int_{\partial\Omega} \Phi_1^2} = s'_n.$$

Writing $u_n = s_n \Phi_1 + w_n$, with $w_n \in \text{span}[\Phi_1]^\perp$, from Proposition 2.2(i), we have $\|w_n\|_{L^\infty(\partial\Omega)} = O(|s_n|^\alpha)$. From Lemmata 2.3, and 2.4, hypotheses (1-5) and (1-7) we get that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{\sigma_1 - \lambda_n}{\|u_n\|_{L^\infty(\partial\Omega)}^{\alpha-1}} &\geq \frac{1}{\int_{\partial\Omega} \Phi_1^2} \liminf_{n \rightarrow \infty} \int_{\partial\Omega} \frac{(s_n \Phi_1 + w_n) g(\lambda_n, \cdot, s_n \Phi_1 + w_n)}{|s_n \Phi_1 + w_n|^{1+\alpha}} \Phi_1^{1+\alpha} \\ &\geq \frac{1}{\int_{\partial\Omega} \Phi_1^2} \liminf_{n \rightarrow +\infty} \int_{\partial\Omega} \frac{s_n g(\lambda_n, \cdot, s_n \Phi_1)}{|s_n|^{1+\alpha}} \Phi_1 \\ &= \frac{1}{\int_{\partial\Omega} \Phi_1^2} \liminf_{n \rightarrow +\infty} \int_{\partial\Omega} \frac{s_n g(\sigma_1, \cdot, s_n \Phi_1)}{|s_n|^{1+\alpha}} \Phi_1 > 0, \end{aligned}$$

and therefore $\lambda_n < \sigma_1$.

Analogously, for (λ'_n, u'_n) we get $\lambda'_n > \sigma_1$. Hence (i) is proved.

To prove (ii), assume, by choosing subsequences if necessary, that $s_n > s'_n > s_{n+1}$ for all $n \geq 0$ and that $0 < s_n, s'_n \leq S_0$ where S_0 is the one from Proposition 2.2(ii). In particular, by parts (i) and (ii) of Proposition 2.2, if $(\lambda, u) \in \mathcal{C}^+$ and $P(u) = s < S_0$ then $\|u\|_{L^\infty(\partial\Omega)} \leq (1 + C_1 \|G_1\|_{L^r(\partial\Omega)} |S_0|^{\alpha-1})s$. Taking S_0 small enough we may assume that $\|u\|_{L^\infty(\partial\Omega)} \leq 2s$.

Let

$$(2-19) \quad K_n = \{(\lambda, u) \in \mathcal{C}^+ : P(u) = s \text{ and } s_n \geq s \geq s_{n+1}\}.$$

Let us see that, for each $n \in \mathbb{N}$, K_n is a compact subset of $\mathbb{R} \times C(\bar{\Omega})$. Let $\{(\mu_k, v_k)\}$ be a sequence in K_n . Without loss of generality we may assume that $\{\mu_k\}$ converges to μ^* . Since $v_k = t_k \Phi_1 + w_k$ with $w_k = O(|t_k|^\alpha)$ and $s_n \geq t_k =: P(v_k) \geq s_{n+1}$, for all k , we have $\|v_k\|_{C(\partial\Omega)} \leq t_k + \|w_k\|_{C(\partial\Omega)} \leq C$ with C independent of k . This together with Proposition 2.3 of [Arrieta et al. 2007] yields

$$(2-20) \quad \|v_k\|_{C(\bar{\Omega})} \leq C_1(1 + \|v_k\|_{C(\partial\Omega)}) \leq C,$$

where, again, C is independent of k . Since the embedding $C^\gamma(\bar{\Omega}) \rightarrow C^{\gamma'}(\bar{\Omega})$ is compact for $0 < \gamma' < \gamma$ we may further assume that the sequence $\{v_k\}$ converges to some $u^* \in C^{\gamma'}(\bar{\Omega})$. This, hypothesis (H1) and the dominated convergence theorem imply that $\{g(\mu_k, \cdot, v_k)\}$ converges to $g(\mu^*, \cdot, u^*)$ in $L^r(\partial\Omega)$. Therefore, since

$$(2-21) \quad \begin{cases} -\Delta v_k + v_k = 0 & \text{in } \Omega \\ \frac{\partial v_k}{\partial n} = \mu_k v_k + g(\mu_k, x, v_k) & \text{on } \partial\Omega, \end{cases}$$

passing to the limit in the weak sense we have

$$(2-22) \quad \begin{cases} -\Delta u^* + u^* = 0 & \text{in } \Omega, \\ \frac{\partial u^*}{\partial n} = \mu^* u^* + g(\lambda^*, x, u^*) & \text{on } \partial\Omega. \end{cases}$$

By the continuity of the projection operator we also have $s_n \geq s^* = P(u^*) = \lim_{k \rightarrow \infty} P(v_k) \geq s_{n+1}$. Hence $(\mu^*, u^*) \in K_n$, which proves that K_n is compact.

Since $s_n > s'_n > s_{n+1}$ there exists $(\lambda, u) \in K_n$ with $\lambda > \sigma_1$. Hence, if we define

$$(2-23) \quad \lambda_n^* = \sup\{\lambda : (\lambda, u) \in K_n\},$$

then $\lambda_n^* \geq \lambda'_n > \sigma_1$ see part (i). From the compactness of K_n there exists u_n^* such that $(\lambda_n^*, u_n^*) \in K_n$. From the definition of λ_n^* if (λ, u) is a solution of (1-1) with $s_n > P(u_n) > s_{n+1}$, then $\lambda \leq \lambda_n^*$ which proves that (λ_n^*, u_n^*) is a (supercritical) turning point.

With a completely symmetric argument, using the sets

$$K'_n = \{(\lambda, u) \in \mathcal{C}^+ : P(u) = s \text{ and } s'_n \geq s \geq s'_{n+1}\}$$

and defining $\lambda_n'^* = \inf\{\lambda : (\lambda, u) \in K'_n\}$, we show the existence of u_* such that $(\lambda_n'^*, u_n'^*) \in K'_n$ is a (subcritical) turning point.

In order to prove the existence of resonant solutions, we now show that there exists $n_0 \in \mathbb{N}$ such that for each $n \geq n_0$ both sets K_n and K'_n contain resonant solutions, that is, solutions of the form (σ_1, u) .

We use a *reductio ad absurdum* argument for the sets K_n . If this is not the case, then there will exist a sequence of integers numbers $n_j \rightarrow +\infty$ such that K_{n_j} does not contain any resonant solution. This implies that the compact sets $K_{n_j}^+ = \{(\lambda, u) \in K_{n_j} : \lambda \geq \sigma_1\}$ can be written as

$$K_{n_j}^+ = \mathcal{C}^+ \cap \{(\lambda, u) \in \mathbb{R} \times C(\partial\Omega) : \lambda > \sigma_1, s_{n_j} > P(u) > s_{n_j+1}\};$$

therefore $K_{n_j}^+$ contains at least a connected component of \mathcal{C}^+ . Moreover it is nonempty since we know that there exists at least one solution (λ, u) with $P(u) = s'_{n_j} \in (s_{n_j}, s_{n_j+1})$ and therefore $\lambda > \sigma_1$. The fact that we can construct a sequence of connected components of \mathcal{C}^+ contradicts the fact that \mathcal{C}^+ is a connected near $(\sigma_1, 0) \in \mathbb{R} \times C(\bar{\Omega})$.

A completely symmetric argument can be applied to the sets K'_n . □

3. Two examples

3.1. Resonant solutions for the oscillatory nonlinearity (1-2). In [Arrieta et al. 2007, Theorem 8.1] it is proved that if $\alpha > 1$, for any $\beta \in \mathbb{R}$, and $C \in \mathbb{R}$ there is an unbounded branch of positive solutions. Assume from now that $\beta < 0$.

Taking $|C| \leq 1$ it is not difficult to see that

$$u_k(x) := [\sin(-C) + k\pi]^{1/\beta} \Phi_1(x), \quad k \geq 0,$$

defines a sequence of resonant solutions to (1-1) such that $u_k(x) \rightarrow 0$ as $k \rightarrow \infty$.

3.2. A one-dimensional example. Now we consider the one-dimensional version of (1-1), where most computations can be made explicit.

Let $\{\sigma_i\}$ denote the sequence of *Steklov* eigenvalues of the problem (1-3). For $N > 1$ the Steklov eigenvalues form an increasing sequence of real numbers, $\{\sigma_i\}_{i=1}^{\infty}$ while for $N = 1$ there are only two Steklov eigenvalues as we made explicit below.

Observe that Equation (1-1) in the one-dimensional domain $\Omega = (0, 1)$ reads

$$\begin{cases} -u_{xx} + u = 0 & \text{in } (0, 1), \\ -u_x(0) = \lambda u + g(\lambda, 0, u(0)), \\ u_x(1) = \lambda u + g(\lambda, 1, u(1)). \end{cases}$$

The general solution of the differential equation is $u(x) = ae^x + be^{-x}$ and therefore the nonlinear boundary conditions provide two nonlinear equations in terms of two constants a and b . The function $u = ae^x + be^{-x}$ is a solution if (λ, a, b) satisfy

$$\begin{pmatrix} -(1+\lambda) & (1-\lambda) \\ (1-\lambda)e & -(1+\lambda)e^{-1} \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} g(\lambda, 0, a+b) \\ g(\lambda, 1, ae+be^{-1}) \end{pmatrix}.$$

In this case we only have two Steklov eigenvalues,

$$\sigma_1 = \frac{e-1}{e+1} < \sigma_2 = \frac{1}{\sigma_1} = \frac{e+1}{e-1}.$$

Restricting the analysis to symmetric solutions $u_s(x) = s(e^x + e^{1-x})$, with $s \in \mathbb{R}$, and choosing $g(\lambda, x, s) = g(s)$, it is easy to prove that $u_s(x)$ is a solution if and only if λ satisfies

$$(3-1) \quad \lambda(s) = \sigma_1 - \frac{g(s(e+1))}{s(e+1)}, \quad s > 0.$$

Therefore, whenever $g(u) = o(u)$ at zero, there is a branch of solutions $(\lambda(s), u_s)$ converging to $(\sigma_1, 0)$ as $s \rightarrow 0$.

Now fix $g(s) = s^\alpha \sin(s^\beta)$ for an arbitrary $\alpha > 1$, $\beta < 0$. From the definition in (2-1) we can write

$$\underline{\mathbf{G}}_{0^+} := \int_{\partial\Omega} \liminf_{s \rightarrow 0^+} \frac{sg(s)}{|s|^{1+\alpha}} \Phi^{1+\alpha} = \int_{\partial\Omega} \liminf_{s \rightarrow 0^+} \sin(s^\beta) \Phi^{1+\alpha} = - \int_{\partial\Omega} \Phi^{1+\alpha} < 0,$$

$$\overline{\mathbf{G}}_{0^+} := \int_{\partial\Omega} \limsup_{s \rightarrow 0^+} \frac{sg(s)}{|s|^{1+\alpha}} \Phi^{1+\alpha} = \int_{\partial\Omega} \limsup_{s \rightarrow 0^+} \sin(s^\beta) \Phi^{1+\alpha} = \int_{\partial\Omega} \Phi^{1+\alpha} > 0,$$

and then $\underline{G}_{0^+} < 0 < \overline{G}_{0^+}$.

Moreover, by looking in (3-1) at the values of $s \in \mathbb{R}$ such that $\lambda(s) = \sigma_1$ it is easy to check that (σ_1, u_k) is a solution for any $k \in \mathbb{Z}$, where

$$u_k(x) := \frac{(k\pi)^{1/\beta}}{e+1} (e^x + e^{1-x});$$

that is, there is a sequence of solutions of the resonant problem converging to zero, as shown in Figure 3.

Moreover, computing in (3-1) the local maxima and minima of $\lambda(s)$ it is not difficult to check that (λ_k^*, u_k^*) is a sequence of turning points converging to zero, where

$$\lambda_k^* := \sigma_1 - t_k^{(\alpha-1)/\beta} \sin(t_k), \quad u_k^*(x) := t_k^{1/\beta} (e^x + e^{1-x})$$

and where t_k is such that

$$\tan(t_k) = -\frac{\beta}{\alpha-1} t_k, \quad t_k \in [-\pi/2 + k\pi, \pi/2 + k\pi]$$

with $t_k \rightarrow \infty$ and $t_k^{1/\beta} \rightarrow 0$ as $k \rightarrow \infty$ thanks to $\beta < 0$.

Let us observe that the bifurcation from zero phenomena occurs whenever $\alpha > 1$ for any β and that whenever $\alpha + \beta < 1$ the number of oscillations grows up faster than the number of oscillations of multiples of the eigenfunction and cannot be controlled; compare the two parts of Figure 3.




Figure 3. Bifurcation diagram in the case $\alpha = 1.4$, for two values of β .

References

- [Arrieta et al. 2007] J. M. Arrieta, R. Pardo, and A. Rodríguez-Bernal, “Bifurcation and stability of equilibria with asymptotically linear boundary conditions at infinity”, *Proc. Roy. Soc. Edinburgh Sect. A* **137**:2 (2007), 225–252. MR 2009d:35194 Zbl 1202.35119
- [Arrieta et al. 2009] J. M. Arrieta, R. Pardo, and A. Rodríguez-Bernal, “Equilibria and global dynamics of a problem with bifurcation from infinity”, *J. Differential Equations* **246**:5 (2009), 2055–2080. MR 2010c:35016 Zbl 1195.35040
- [Arrieta et al. 2010] J. M. Arrieta, R. Pardo, and A. Rodríguez-Bernal, “Infinite resonant solutions and turning points in a problem with unbounded bifurcation”, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **20**:9 (2010), 2885–2896. MR 2011m:35107 Zbl 1202.35085
- [Castro and Pardo 2011] A. Castro and R. Pardo, “Infinitely many stability switches in a problem with sublinear oscillatory boundary conditions”, preprint MA-UCM 2011-2, Univ. Complutense de Madrid, 2011, available at <http://www.ucm.es/centros/cont/descargas/documento23873.pdf>.
- [Crandall and Rabinowitz 1971] M. G. Crandall and P. H. Rabinowitz, “Bifurcation from simple eigenvalues”, *J. Functional Analysis* **8** (1971), 321–340. MR 44 #5836 Zbl 0219.46015
- [García-Melián et al. 2009] J. García-Melián, J. D. Rossi, and J. C. Sabina de Lis, “An elliptic system with bifurcation parameters on the boundary conditions”, *J. Differential Equations* **247**:3 (2009), 779–810. MR 2010f:35073 Zbl 1171.35037
- [Korman 2008] P. Korman, “An oscillatory bifurcation from infinity, and from zero”, *NoDEA Non-linear Differential Equations Appl.* **15**:3 (2008), 335–345. MR 2009h:35137 Zbl 1166.34010
- [Rabinowitz 1971] P. H. Rabinowitz, “Some global results for nonlinear eigenvalue problems”, *J. Functional Analysis* **7** (1971), 487–513. MR 46 #745 Zbl 0212.16504
- [Rabinowitz 1973] P. H. Rabinowitz, “On bifurcation from infinity”, *J. Differential Equations* **14** (1973), 462–475. MR 48 #7047 Zbl 0272.35017

Received June 3, 2011. Revised November 22, 2011.

ALFONSO CASTRO
DEPARTMENT OF MATHEMATICS
HARVEY MUDD COLLEGE
301 PLATT BOULEVARD
CLAREMONT, CA 91711
UNITED STATES
castro@math.hmc.edu

ROSA PARDO
DEPARTAMENTO DE MATEMÁTICA APLICADA
UNIVERSIDAD COMPLUTENSE DE MADRID
AVENIDA COMPLUTENSE S/N
28040 MADRID
SPAIN
rpardo@mat.ucm.es

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 257 No. 1 May 2012

Energy and volume of vector fields on spherical domains	1
FABIANO G. B. BRITO, ANDRÉ O. GOMES and GIOVANNI S. NUNES	
Maps on 3-manifolds given by surgery	9
BOLDIZSÁR KALMÁR and ANDRÁS I. STIPSICZ	
Strong solutions to the compressible liquid crystal system	37
YU-MING CHU, XIAN-GAO LIU and XIAO LIU	
Presentations for the higher-dimensional Thompson groups nV	53
JOHANNA HENNIG and FRANCESCO MATUCCI	
Resonant solutions and turning points in an elliptic problem with oscillatory boundary conditions	75
ALFONSO CASTRO and ROSA PARDO	
Relative measure homology and continuous bounded cohomology of topological pairs	91
ROBERTO FRIGERIO and CRISTINA PAGLIANTINI	
Normal enveloping algebras	131
ALEXANDRE N. GRISHKOV, MARINA RASSKAZOVA and SALVATORE SICILIANO	
Bounded and unbounded capillary surfaces in a cusp domain	143
YASUNORI AOKI and DAVID SIEGEL	
On orthogonal polynomials with respect to certain discrete Sobolev inner product	167
FRANCISCO MARCELLÁN, RAMADAN ZEJNULLAHU, BUJAR FEJZULLAHU and EDMUNDO HUERTAS	
Green versus Lempert functions: A minimal example	189
PASCAL THOMAS	
Differential Harnack inequalities for nonlinear heat equations with potentials under the Ricci flow	199
JIA-YONG WU	
On overtwisted, right-veering open books	219
PAOLO LISCA	
Weakly Krull domains and the composite numerical semigroup ring $D + E[\Gamma^*]$	227
JUNG WOOK LIM	
Arithmeticity of complex hyperbolic triangle groups	243
MATTHEW STOVER	



0030-8730(201205)257:1;1-7