

*Pacific
Journal of
Mathematics*

**ON ORTHOGONAL POLYNOMIALS WITH RESPECT TO
CERTAIN DISCRETE SOBOLEV INNER PRODUCT**

FRANCISCO MARCELLÁN, RAMADAN ZEJNULLAHU,
BUJAR FEJZULLAHU AND EDMUNDO HUERTAS

ON ORTHOGONAL POLYNOMIALS WITH RESPECT TO CERTAIN DISCRETE SOBOLEV INNER PRODUCT

FRANCISCO MARCELLÁN, RAMADAN ZEJNULLAHU,
BUJAR FEJZULLAHU AND EDMUNDO HUERTAS

In this paper we deal with sequences of polynomials orthogonal with respect to the discrete Sobolev inner product

$$\langle f, g \rangle_S = \int_0^\infty \omega(x) f(x) g(x) dx + Mf(\xi)g(\xi) + Nf'(\xi)g'(\xi),$$

where ω is a weight function, $\xi \leq 0$, and $M, N \geq 0$. The location of the zeros of discrete Sobolev orthogonal polynomials is given in terms of the zeros of standard polynomials orthogonal with respect to the weight function ω . In particular, for $\omega(x) = x^\alpha e^{-x}$ we obtain the asymptotics for discrete Laguerre–Sobolev orthogonal polynomials.

1. Introduction

Polynomials orthogonal with respect to an inner product

$$(1) \quad \langle f, g \rangle = \int_E \omega(x) f(x) g(x) dx + Mf(\xi)g(\xi) + Nf'(\xi)g'(\xi),$$

where ξ is a real number and $d\mu$ is a positive Borel measure supported on an infinite subset E of the real line have been considered by several authors (see, for instance, [Alfaro et al. 1992; López et al. 1995; Marcellán and Ronveaux 1990; Marcellán and Van Assche 1993] and the references therein). They are known in the literature as Sobolev-type or discrete Sobolev orthogonal polynomials. Special attention has been paid to their algebraic and analytic properties of these polynomials, in particular, the distribution of their zeros taking into account the location of the point ξ with respect to the set E .

When E is the interval $[0, +\infty)$ and $\xi = 0$, Meijer [1993a] analyzed some analytic properties of the zeros of the so called discrete Sobolev orthogonal polynomials (1). Some results of [Meijer 1993a] are direct generalizations of the results of [Koekoek and Meijer 1993], where the weight function is the Laguerre

MSC2010: primary 33C47; secondary 42C05.

Keywords: orthogonal polynomials, discrete Sobolev polynomials, Laguerre polynomials, asymptotics.

weight $\omega(x) = x^\alpha e^{-x}$. Koekoek and Meijer established properties of the discrete Laguerre–Sobolev polynomials such as their representation as a hypergeometric series, an holonomic second order linear differential equation associated with them, properties of the zeros, and a higher-order recurrence relation that such polynomials satisfy. The asymptotic properties of these discrete Laguerre–Sobolev polynomials have been studied in [Álvarez-Nodarse and Moreno-Balcázar 2004; Marcellán and Moreno-Balcázar 2006], while the analysis of convergence of the Fourier expansions in terms of such polynomials was done in [Fejzullahu and Marcellán 2009].

In this paper we consider the discrete Sobolev polynomials $\{\hat{S}_n\}_{n \geq 0}$ orthogonal with respect to (1) where $E = [0, +\infty)$ and $\xi < 0$. We show that these polynomials can be expressed as

$$\hat{S}_n(x) = \hat{P}_n(x) + A_{n,1}(x - \xi)\hat{P}_{n-1}^{[2]}(x) + A_{n,2}(x - \xi)^2\hat{P}_{n-2}^{[4]}(x),$$

where $\{\hat{P}_n\}_{n \geq 0}$ and $\{\hat{P}_n^{[k]}\}_{n \geq 0}$, $k \in \mathbb{N}$, are the sequences of monic polynomials orthogonal with respect to the weight functions $\omega(\cdot)$ and $(\cdot - \xi)^k \omega(\cdot)$, respectively. Moreover, the behavior of the coefficients $A_{n,1}$ and $A_{n,2}$ is studied in more detail. In particular, when ω is the Laguerre weight, we obtain some asymptotic properties for the sequence of discrete Laguerre–Sobolev orthogonal polynomials.

The structure of the manuscript is as follows. In Section 2 we give some basic background concerning polynomial perturbations of a measure as well as interlacing properties for the zeros of the corresponding orthogonal polynomials. We point out that the results presented therein are of independent interest in terms of the core of our contribution. Indeed, they constitute an alternative approach in the subject. In Section 3, a representation of monic polynomials orthogonal with respect to the inner product (1) is given in terms of polynomial orthogonal with respect to polynomial perturbations of the weight function. Some results about their zeros are deduced. In Section 4 we focus our attention on the asymptotics of discrete Laguerre–Sobolev orthogonal polynomials. More precisely, we obtain outer relative asymptotics, a Mehler–Heine formula and the Plancherel–Rotach outer asymptotics for such orthogonal polynomials.

Throughout this paper positive constants are denoted by c, c_1, \dots , and they may vary at every occurrence. The notation $u_n \cong v_n$ means that the sequence $\{u_n/v_n\}_n$ converges to 1. We will denote by $k(\pi_n)$ the leading coefficient of any polynomial π_n and $\hat{\pi}_n(x) = (k(\pi_n))^{-1} \pi_n(x)$.

2. Auxiliary results

Let ω denote a weight function on $(0, \infty)$, i.e., $\omega(x) \geq 0$ and all moments

$$c_n = \int_0^\infty \omega(x)x^n dx, \quad n = 0, 1, \dots$$

exist. Let $\{\hat{P}_n\}_{n \geq 0}$ denote the sequence of monic polynomials orthogonal (SMOP, in short) with respect to the standard inner product

$$\langle f, g \rangle = \int_0^\infty \omega(x) f(x) g(x) dx.$$

In particular, from the moments we get an explicit expression of the SMOP. Indeed, we get

$$\hat{P}_0(x) = 1$$

and

$$(2) \quad \hat{P}_n(x) = \frac{1}{\Delta_{n-1}} \begin{vmatrix} c_0 & c_1 & c_2 & \dots & c_n \\ c_1 & c_2 & c_3 & \dots & c_{n+1} \\ c_2 & c_3 & c_4 & \dots & c_{n+2} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ c_{n-1} & c_n & c_{n+1} & \dots & c_{2n-1} \\ 1 & x & x^2 & \dots & x^n \end{vmatrix}, \quad n \geq 1,$$

where

$$\Delta_{n-1} = \begin{vmatrix} c_0 & c_1 & c_2 & \dots & c_{n-1} \\ c_1 & c_2 & c_3 & \dots & c_n \\ c_2 & c_3 & c_4 & \dots & c_{n+1} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ c_{n-1} & c_n & c_{n+1} & \dots & c_{2n-2} \end{vmatrix}, \quad n \geq 1,$$

are the Gram determinants.

The n -th reproducing kernel for ω is

$$K_n(x, y) = \sum_{k=0}^n \frac{\hat{P}_k(x) \hat{P}_k(y)}{\|\hat{P}_k\|_\omega^2}.$$

Here, $\|\hat{P}_n\|_\omega^2 = \langle \hat{P}_n, \hat{P}_n \rangle$. Because of the Christoffel–Darboux formula, it may also be expressed as

$$K_n(x, y) = \frac{1}{\|\hat{P}_n\|_\omega^2} \frac{\hat{P}_{n+1}(x) \hat{P}_n(y) - \hat{P}_n(x) \hat{P}_{n+1}(y)}{x - y}.$$

The confluent formula reads as

$$(3) \quad K_n(x, x) = \sum_{k=0}^n \frac{(\hat{P}_k(x))^2}{\|\hat{P}_k\|_\omega^2} = \frac{1}{\|\hat{P}_n\|_\omega^2} (\hat{P}'_{n+1}(x) \hat{P}_n(x) - \hat{P}'_n(x) \hat{P}_{n+1}(x)).$$

In the same way we can describe the SMOP $\{\hat{P}_n^{[k]}\}_{n \geq 0}$, orthogonal with respect to the inner product

$$\langle f, g \rangle_k = \int_0^\infty (x - \xi)^k \omega(x) f(x) g(x) dx,$$

where $\xi \leq 0$. For $n \geq 1$, they are given by the determinant (2) where c_i is replaced by d_i^k , $k \in \mathbb{N}$, where

$$(4) \quad d_n^k = \int_0^\infty (x - \xi)^k \omega(x) x^n dx = d_{n+1}^{k-1} - \xi d_n^{k-1}, \quad n = 0, 1, \dots,$$

and $c_n = d_n^0$. In the sequel, we will set

$$\|\hat{P}_n^{[k]}\|_{\omega, k}^2 = \int_0^\infty (x - \xi)^k \omega(x) (\hat{P}_n^{[k]}(x))^2 dx.$$

Proposition 1. Let $D_{n-1}^k = \det[a_{ij}^k]_{0 \leq i, j \leq n-1}$, where $a_{ij}^k = d_{i+j}^k$, $k \in \mathbb{N}$. Then

$$(5) \quad D_{n-1}^k = (-1)^n D_{n-1}^{k-1} \hat{P}_n^{[k-1]}(\xi),$$

with $D_{n-1}^0 = \Delta_{n-1}$.

Proof. For $n \geq 1$ and $k \in \mathbb{N}$,

$$(6) \quad \hat{P}_n^{[k-1]}(x) = \frac{1}{D_{n-1}^{[k-1]}} \begin{vmatrix} d_0^{k-1} & d_1^{k-1} & \dots & d_n^{k-1} \\ d_1^{k-1} & d_2^{k-1} & \dots & d_{n+1}^{k-1} \\ \cdot & \cdot & \dots & \cdot \\ d_{n-1}^{k-1} & d_n^{k-1} & \dots & d_{2n-1}^{k-1} \\ 1 & x & \dots & x^n \end{vmatrix},$$

with $\hat{P}_n = \hat{P}_n^{[0]}$. The determinant in (6) becomes [Szegő 1975, Formula (2.2.9)]

$$\hat{P}_n^{[k-1]}(x) = \frac{(-1)^n}{D_{n-1}^{k-1}} \begin{vmatrix} d_1^{k-1} - d_0^{k-1}x & d_2^{k-1} - d_1^{k-1}x & \dots & d_n^{k-1} - d_{n-1}^{k-1}x \\ d_2^{k-1} - d_1^{k-1}x & d_3^{k-1} - d_2^{k-1}x & \dots & d_{n+1}^{k-1} - d_n^{k-1}x \\ \cdot & \cdot & \dots & \cdot \\ d_n^{k-1} - d_{n-1}^{k-1}x & d_{n+1}^{k-1} - d_n^{k-1}x & \dots & d_{2n-1}^{k-1} - d_{2n-2}^{k-1}x \end{vmatrix}.$$

Now, by using (4), (5) follows. □

Next we will compute some integrals involving the polynomials $\hat{P}_n^{[k]}$.

Proposition 2. (i) The integral $\int_0^\infty (x - \xi)^{k-1} \omega(x) \hat{P}_n^{[k]}(x) dx$ is given by

$$\frac{\|\hat{P}_n^{[k-1]}\|_{\omega, k-1}^2}{\hat{P}_n^{[k-1]}(\xi)} = \begin{cases} \frac{\|\hat{P}_n\|_{\omega}^2}{\hat{P}_n(\xi)} & \text{if } k = 1, \\ \frac{(-1)^{k-1}}{\hat{P}_n^{[k-1]}(\xi)} \prod_{i=1}^{k-1} \frac{\hat{P}_{n+1}^{[i-1]}(\xi)}{\hat{P}_n^{[i-1]}(\xi)} \|\hat{P}_n\|_{\omega}^2 & \text{if } k \geq 2. \end{cases}$$

(ii) The integral $\int_0^\infty (x - \xi)^{k-2} \omega(x) \hat{P}_n^{[k]}(x) dx$ is given by

$$\frac{(\hat{P}_{n+1}^{[k-2]}(x))'_{x=\xi} \|\hat{P}_n^{[k-2]}\|_{\omega, k-2}^2}{\hat{P}_n^{[k-1]}(\xi) \hat{P}_n^{[k-2]}(\xi)} = \begin{cases} \frac{(\hat{P}_{n+1}(x))'_{x=\xi} \|\hat{P}_n\|_{\omega}^2}{\hat{P}_n(\xi) \hat{P}_n^{[1]}(\xi)} & \text{if } k = 2, \\ \frac{(-1)^k (\hat{P}_{n+1}^{[k-2]}(x))'_{x=\xi}}{\hat{P}_n^{[k-1]}(\xi) \hat{P}_n^{[k-2]}(\xi)} \prod_{i=1}^{k-2} \frac{\hat{P}_{n+1}^{[i-1]}(\xi)}{\hat{P}_n^{[i-1]}(\xi)} \|\hat{P}_n\|_{\omega}^2 & \text{if } k \geq 3. \end{cases}$$

Proof. (i) Using (4) recursively as well as properties of determinants, we have

$$\begin{aligned} D_{n-1}^k \int_0^\infty (x - \xi)^{k-1} \omega(x) \hat{P}_n^{[k]}(x) dx &= \begin{vmatrix} d_0^k & d_1^k & d_2^k & \dots & d_n^k \\ d_1^k & d_2^k & d_3^k & \dots & d_{n+1}^k \\ \cdot & \cdot & \cdot & \dots & \cdot \\ d_{n-1}^k & d_n^k & d_{n+1}^k & \dots & d_{2n-1}^k \\ d_0^{k-1} & d_1^{k-1} & d_2^{k-1} & \dots & d_n^{k-1} \end{vmatrix} \\ &= \begin{vmatrix} d_1^{k-1} & d_2^{k-1} & d_3^{k-1} & \dots & d_{n+1}^{k-1} \\ d_2^{k-1} & d_3^{k-1} & d_4^{k-1} & \dots & d_{n+2}^{k-1} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ d_n^{k-1} & d_{n+1}^{k-1} & d_{n+2}^{k-1} & \dots & d_{2n}^{k-1} \\ d_0^{k-1} & d_1^{k-1} & d_2^{k-1} & \dots & d_n^{k-1} \end{vmatrix} \\ &= (-1)^n D_n^{k-1}. \end{aligned}$$

On the other hand,

$$\|\hat{P}_n^{[k-1]}\|_{\omega, k-1}^2 = \int_0^\infty (x - \xi)^{k-1} \omega(x) x^n \hat{P}_n^{[k-1]}(x) dx = \frac{D_n^{k-1}}{D_{n-1}^{k-1}},$$

and by using (5) we get

$$(7) \quad \int_0^\infty (x - \xi)^{k-1} \omega(x) \hat{P}_n^{[k]}(x) dx = \frac{(-1)^n D_{n-1}^{k-1} \|\hat{P}_n^{[k-1]}\|_{\omega, k-1}^2}{D_{n-1}^k} = \frac{\|\hat{P}_n^{[k-1]}\|_{\omega, k-1}^2}{\hat{P}_n^{[k-1]}(\xi)}.$$

On the other hand, we have from [Szegő 1975, Theorem 2.5]

$$(8) \quad (x - \xi) \hat{P}_n^{[k]}(x) = \hat{P}_{n+1}^{[k-1]}(x) - \frac{\hat{P}_{n+1}^{[k-1]}(\xi)}{\hat{P}_n^{[k-1]}(\xi)} \hat{P}_n^{[k-1]}(x).$$

Therefore,

$$\|\hat{P}_n^{[k]}\|_{\omega,k}^2 = -\frac{\hat{P}_{n+1}^{[k-1]}(\xi)}{\hat{P}_n^{[k-1]}(\xi)} \|\hat{P}_n^{[k-1]}\|_{\omega,k-1}^2.$$

Using this relation recursively we obtain

$$(9) \quad \|\hat{P}_n^{[k]}\|_{\omega,k}^2 = (-1)^k \prod_{i=1}^k \frac{\hat{P}_{n+1}^{[i-1]}(\xi)}{\hat{P}_n^{[i-1]}(\xi)} \|\hat{P}_n\|_{\omega}^2, \quad k \geq 2.$$

Combining (7) and (9), our statement follows.

(ii) We have

$$(10) \quad (\hat{P}_{n+1}^{[k-2]}(x))' = \frac{1}{D_n^{k-2}} \begin{vmatrix} d_0^{k-2} & d_1^{k-2} & d_2^{k-2} & \dots & d_{n+1}^{k-2} \\ d_1^{k-2} & d_2^{k-2} & d_3^{k-2} & \dots & d_{n+2}^{k-2} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ d_n^{k-2} & d_{n+1}^{k-2} & d_{n+2}^{k-2} & \dots & d_{2n+1}^{k-2} \\ 0 & 1 & 2x & \dots & nx^{n-1} \end{vmatrix}, \quad n \geq 0.$$

Now, adding to the last column the n -th and $(n-1)$ -th columns multiplied by $-2x$ and x^2 , respectively, and repeating this operation for each of the preceding columns, we obtain

$$(11) \quad (\hat{P}_{n+1}^{[k-2]}(x))' = \frac{1}{D_n^{k-2}} \begin{vmatrix} d_0^{k-2} & d_1^{k-2} & d_2^{k-2} - 2xd_1^{k-2} + x^2d_0^{k-2} & \dots & d_{n+1}^{k-2} - 2xd_n^{k-2} + x^2d_{n-1}^{k-2} \\ d_1^{k-2} & d_2^{k-2} & d_3^{k-2} - 2xd_2^{k-2} + x^2d_1^{k-2} & \dots & d_{n+2}^{k-2} - 2xd_{n+1}^{k-2} + x^2d_n^{k-2} \\ \cdot & \cdot & \cdot & \dots & \cdot \\ d_n^{k-2} & d_{n+1}^{k-2} & d_{n+2}^{k-2} - 2xd_{n+1}^{k-2} + x^2d_n^{k-2} & \dots & d_{2n+1}^{k-2} - 2xd_{2n}^{k-2} + x^2d_{2n-1}^{k-2} \\ 0 & 1 & 0 & \dots & 0 \end{vmatrix} \\ = \frac{1}{D_n^{k-2}} \begin{vmatrix} d_2^{k-2} - 2xd_1^{k-2} + x^2d_0^{k-2} & d_3^{k-2} - 2xd_2^{k-2} + x^2d_1^{k-2} & \dots & d_{n+2}^{k-2} - 2xd_{n+1}^{k-2} + x^2d_n^{k-2} \\ d_3^{k-2} - 2xd_2^{k-2} + x^2d_1^{k-2} & d_4^{k-2} - 2xd_3^{k-2} + x^2d_2^{k-2} & \dots & d_{n+3}^{k-2} - 2xd_{n+2}^{k-2} + x^2d_{n+1}^{k-2} \\ \cdot & \cdot & \dots & \cdot \\ d_{n+1}^{k-2} - 2xd_n^{k-2} + x^2d_{n-1}^{k-2} & d_{n+2}^{k-2} - 2xd_{n+1}^{k-2} + x^2d_n^{k-2} & \dots & d_{2n+1}^{k-2} - 2xd_{2n}^{k-2} + x^2d_{2n-1}^{k-2} \\ \cdot & d_0^{k-2} & \dots & d_1^{k-2} \end{vmatrix}.$$

On the other hand,

$$D_{n-1}^k \int_0^\infty (x - \xi)^{k-2} \omega(x) \hat{P}_n^{[k]}(x) dx = \begin{vmatrix} d_0^k & d_1^k & \dots & d_n^k \\ d_1^k & d_2^k & \dots & d_{n+1}^k \\ \cdot & \cdot & \dots & \cdot \\ d_{n-1}^k & d_n^k & \dots & d_{2n-1}^k \\ d_0^{k-2} & d_1^{k-2} & \dots & d_n^{k-2} \end{vmatrix},$$

and by using (5), (9), and (11) we get

$$\begin{aligned} \int_0^\infty (x - \xi)^{k-2} \omega(x) \hat{P}_n^{[k]}(x) dx &= \frac{D_n^{k-2} (\hat{P}_{n+1}^{[k-2]}(x))'_{x=\xi}}{D_{n-1}^{k-2} \hat{P}_n^{[k-1]}(\xi) \hat{P}_n^{[k-2]}(\xi)} = \frac{(\hat{P}_{n+1}^{[k-2]}(x))'_{x=\xi} \|\hat{P}_n^{[k-2]}\|_{\omega, k-2}^2}{\hat{P}_n^{[k-1]}(\xi) \hat{P}_n^{[k-2]}(\xi)} \\ &= \frac{(-1)^k (\hat{P}_{n+1}^{[k-2]}(x))'_{x=\xi}}{\hat{P}_n^{[k-1]}(\xi) \hat{P}_n^{[k-2]}(\xi)} \prod_{i=1}^{k-2} \frac{\hat{P}_{n+1}^{[i-1]}(\xi)}{\hat{P}_n^{[i-1]}(\xi)} \|\hat{P}_n\|_{\omega}^2. \quad \square \end{aligned}$$

Denote by $x_{r,n}^{[k]}$, $r = 1, 2, \dots, n$, the zeros of $\hat{P}_n^{[k]}(x)$ in increasing order.

Proposition 3. (i) *The zeros of $\hat{P}_n^{[k]}(x)$ interlace with both the zeros of $\hat{P}_{n+1}^{[k-1]}(x)$ and $\hat{P}_n^{[k-1]}(x)$, i.e.,*

$$x_{r,n}^{[k-1]} < x_{r,n}^{[k]} < x_{r+1,n+1}^{[k-1]}, \quad r = 1, 2, \dots, n.$$

(ii) *Between two consecutive zeros of $\hat{P}_{n+1}^{[k-2]}$, $k \geq 2$, there is exactly one zero of $\hat{P}_n^{[k]}$.*

(iii) $\text{sgn } \hat{P}_n^{[k-2]}(x_{r,n-1}^{[k]}) = (-1)^{n-r} = -\text{sgn } \hat{P}_{n-2}^{[k+2]}(x_{r,n-1}^{[k]})$ for $r = 1, 2, \dots, n-1$.

Proof. (i) Here we will use the same argument as in [Chihara 1978, page 65] (see also [Bracciali et al. 2002, Lemma 1]). It is well known that the zeros of $\hat{P}_{n+1}^{[k-1]}$ interlace with the zeros of $\hat{P}_n^{[k-1]}$, i.e.,

$$0 < x_{1,n+1}^{[k-1]} < x_{1,n}^{[k-1]} < x_{2,n+1}^{[k-1]} < \dots < x_{n,n}^{[k-1]} < x_{n+1,n+1}^{[k-1]} < \infty.$$

From (5) $\hat{P}_{n+1}^{[k-1]}(\xi) / \hat{P}_n^{[k-1]}(\xi) < 0$ and taking (8) into account we have

$$\begin{aligned} \text{sgn } \hat{P}_n^{[k]}(x_{r,n+1}^{[k-1]}) &= \text{sgn } \hat{P}_n^{[k-1]}(x_{r,n+1}^{[k-1]}) = (-1)^{n-r+1} \quad \text{for } r = 1, 2, \dots, n+1, \\ \text{sgn } \hat{P}_n^{[k]}(x_{r,n}^{[k-1]}) &= \text{sgn } \hat{P}_{n+1}^{[k-1]}(x_{r,n}^{[k-1]}) = (-1)^{n-r+1} \quad \text{for } r = 1, 2, \dots, n. \end{aligned}$$

Thus, there exist zeros $x_{r,n}^{[k]}$, $r = 2, 3, \dots, n$, of $\hat{P}_n^{[k]}(x)$ satisfying

$$x_{r,n}^{[k-1]} < x_{r,n}^{[k]} < x_{r+1,n+1}^{[k-1]}, \quad r = 1, 2, \dots, n.$$

(ii) By using (8) and the recurrence relation we obtain

$$(x - \xi)^2 \hat{P}_n^{[k]}(x) = (d_{1,n}x + d_{2,n}) \hat{P}_{n+1}^{[k-2]}(x) + d_{3,n} \hat{P}_n^{[k-2]}(x).$$

Since $\hat{P}_{n+1}^{[k-2]}(\xi) \neq 0$ we have $d_{3,n} \neq 0$. Now, the rest of the proof can be done in a similar way as in [Meijer 1993a, Lemma 6.1]; see also [Meijer 1993b, Lemma 4.1].

(iii) From (ii) we have $x_{r,n}^{[k-2]} < x_{r,n-1}^{[k]} < x_{r+1,n}^{[k-2]}$ for $r = 1, 2, \dots, n - 1$. Therefore,

$$\operatorname{sgn} \hat{P}_n^{[k-2]}(x_{r,n-1}^{[k]}) = (-1)^{n-r}.$$

Again, according to (ii), $x_{r-1,n-2}^{[k+2]} < x_{r,n-1}^{[k]} < x_{r,n-2}^{[k+2]}$ for $r = 1, 2, \dots, n - 2$, and $x_{n-2,n-2}^{[k+2]} < x_{n-1,n-1}^{[k]}$. Therefore,

$$\operatorname{sgn} \hat{P}_{n-2}^{[k+2]}(x_{r,n-1}^{[k]}) = (-1)^{n-r-1} \quad \text{and} \quad \operatorname{sgn} \hat{P}_{n-2}^{[k+2]}(x_{n-1,n-1}^{[k]}) = 1.$$

As a conclusion,

$$\operatorname{sgn} \hat{P}_n^{[k-2]}(x_{r,n-1}^{[k]}) = -\operatorname{sgn} \hat{P}_{n-2}^{[k+2]}(x_{r,n-1}^{[k]}), \quad r = 1, 2, \dots, n - 1. \quad \square$$

3. Discrete Sobolev orthogonal polynomials

Connection formula. We consider the inner product

$$(12) \quad \langle f, g \rangle_S = \int_0^\infty \omega(x) f(x) g(x) dx + M f(\xi) g(\xi) + N f'(\xi) g'(\xi),$$

where $\xi \leq 0$, and $M, N \geq 0$. Let $\{\hat{S}_n\}_{n \geq 0}$ denote the SMOP with respect to the discrete Sobolev inner product (12)).

Theorem 1. *Let $M \geq 0$ and $N \geq 0$. There are real constants $A_{n,1}$ and $A_{n,2}$ such that*

$$\hat{S}_n(x) = \hat{P}_n(x) + A_{n,1}(x - \xi) \hat{P}_{n-1}^{[2]}(x) + A_{n,2}(x - \xi)^2 \hat{P}_{n-2}^{[4]}(x),$$

where

$$A_{n,1} = \frac{N I_{2,n}(\xi) \hat{P}'_n(\xi) - M I_{3,n}(\xi) \hat{P}_n(\xi)}{I_{1,n}(\xi) I_{3,n}(\xi) - N I_{2,n}(\xi) \hat{P}_{n-1}^{[2]}(\xi)},$$

$$A_{n,2} = \frac{MN \hat{P}_n(\xi) \hat{P}_{n-1}^{[2]}(\xi) - N I_{1,n}(\xi) \hat{P}'_n(\xi)}{I_{1,n}(\xi) I_{3,n}(\xi) - N I_{2,n}(\xi) \hat{P}_{n-1}^{[2]}(\xi)},$$

$$I_{1,n}(\xi) = -\frac{\hat{P}_n(\xi)}{K_{n-1}(\xi, \xi)},$$

$$I_{2,n}(\xi) = \frac{\hat{P}_{n-1}(\xi) \hat{P}_{n-1}^{[1]}(\xi) \hat{P}_{n-1}^{[2]'}(\xi)}{\hat{P}_{n-2}(\xi) \hat{P}_{n-2}^{[1]}(\xi) \hat{P}_{n-2}^{[2]}(\xi) \hat{P}_{n-2}^{[3]}(\xi)} \|\hat{P}_{n-2}\|_\omega^2,$$

$$I_{3,n}(\xi) = -\frac{\hat{P}_{n-1}(\xi) \hat{P}_{n-1}^{[1]}(\xi) \hat{P}_{n-1}^{[2]}(\xi)}{\hat{P}_{n-2}(\xi) \hat{P}_{n-2}^{[1]}(\xi) \hat{P}_{n-2}^{[2]}(\xi) \hat{P}_{n-2}^{[3]}(\xi)} \|\hat{P}_{n-2}\|_\omega^2.$$

Proof. We will prove that

$$\langle \hat{S}_n, (\cdot - \xi)^k \rangle_S = 0 \quad \text{for } k = 0, 1, \dots, n - 1.$$

For $k \geq 2$ and $n > k$,

$$\begin{aligned} &\langle \hat{S}_n, (\cdot - \xi)^k \rangle_S \\ &= \int_0^\infty \omega(x) \hat{S}_n(x) (x - \xi)^k dx \\ &= \int_0^\infty \omega(x) \hat{P}_n(x) (x - \xi)^k dx + A_{n,1} \int_0^\infty (x - \xi)^2 \omega(x) \hat{P}_{n-1}^{[2]}(x) (x - \xi)^{k-1} dx \\ &\quad + A_{n,2} \int_0^\infty (x - \xi)^4 \omega(x) \hat{P}_{n-2}^{[4]}(x) (x - \xi)^{k-2} dx \\ &= 0, \end{aligned}$$

Now consider $k = 0$ and $n \geq 1$. We have

$$\begin{aligned} \langle \hat{S}_n, 1 \rangle_S &= \int_0^\infty \omega(x) \hat{S}_n(x) dx + M \hat{S}_n(\xi) \\ &= A_{n,1} \int_0^\infty (x - \xi) \omega(x) \hat{P}_{n-1}^{[2]}(x) dx + A_{n,2} \int_0^\infty (x - \xi)^2 \omega(x) \hat{P}_{n-2}^{[4]}(x) dx \\ &\quad + M \hat{P}_n(\xi). \end{aligned}$$

On the other hand, by using Proposition 2(i),

$$(13) \quad I_{1,n}(\xi) = \int_0^\infty (x - \xi) \omega(x) \hat{P}_{n-1}^{[2]}(x) dx = -\frac{\hat{P}_n(\xi)}{\hat{P}_{n-1}(\xi) \hat{P}_{n-1}^{[1]}(\xi)} \|\hat{P}_{n-1}\|_\omega^2,$$

and taking derivatives in (8) and then substituting $x = \xi$ we get

$$(14) \quad \hat{P}_{n-1}^{[k]}(\xi) = (\hat{P}_n^{[k-1]}(x))'_{x=\xi} - \frac{\hat{P}_n^{[k-1]}(\xi)}{\hat{P}_{n-1}^{[k-1]}(\xi)} (\hat{P}_{n-1}^{[k-1]}(x))'_{x=\xi}.$$

Combining (3), (13), and (14), we get

$$I_{1,n}(\xi) = -\frac{\hat{P}_n(\xi)}{K_{n-1}(\xi, \xi)}.$$

Using Proposition 2(ii),

$$\begin{aligned} (15) \quad I_{2,n}(\xi) &= \int_0^\infty (x - \xi)^2 \omega(x) \hat{P}_{n-2}^{[4]}(x) dx = \frac{(\hat{P}_{n-1}^{[2]}(x))'_{x=\xi} \|\hat{P}_{n-2}\|_{\omega,2}^2}{\hat{P}_{n-2}^{[2]}(\xi) \hat{P}_{n-2}^{[3]}(\xi)} \\ &= \frac{\hat{P}_{n-1}(\xi) \hat{P}_{n-1}^{[1]}(\xi) (\hat{P}_{n-1}^{[2]}(x))'_{x=\xi}}{\hat{P}_{n-2}(\xi) \hat{P}_{n-2}^{[1]}(\xi) \hat{P}_{n-2}^{[2]}(\xi) \hat{P}_{n-2}^{[3]}(\xi)} \|\hat{P}_{n-2}\|_\omega^2. \end{aligned}$$

Therefore,

$$\langle \hat{S}_n, 1 \rangle_S = A_{n,1}I_{1,n}(\xi) + A_{n,2}I_{2,n}(\xi) + M\hat{P}_n(\xi).$$

In the same way, for $k = 1$ and $n \geq 2$, we have

$$\begin{aligned} \langle \hat{S}_n, (\cdot - \xi) \rangle_S &= \int_0^\infty \omega(x)\hat{S}_n(x)(x - \xi) dx + N\hat{S}'_n(\xi) \\ &= A_{n,2}I_{3,n}(\xi) + NA_{n,1}\hat{P}_{n-1}^{[2]}(\xi) + N\hat{P}'_n(\xi), \end{aligned}$$

where

$$\begin{aligned} I_{3,n}(\xi) &= \int_0^\infty (x - \xi)^3 \omega(x)\hat{P}_{n-2}^{[4]}(x) dx = \frac{\|\hat{P}_{n-2}^{[3]}\|_{\omega,3}^2}{\hat{P}_n^{[3]}(\xi)} \\ &= -\frac{\hat{P}_{n-1}(\xi)\hat{P}_{n-1}^{[1]}(\xi)\hat{P}_{n-1}^{[2]}(\xi)}{\hat{P}_{n-2}(\xi)\hat{P}_{n-2}^{[1]}(\xi)\hat{P}_{n-2}^{[2]}(\xi)\hat{P}_{n-2}^{[3]}(\xi)}\|\hat{P}_{n-2}\|_{\omega}^2. \end{aligned}$$

Finally, using the expressions of $A_{n,1}$ and $A_{n,2}$, our statement follows. □

Next, we will study the behavior of the coefficients $A_{n,1}$ and $A_{n,2}$.

Proposition 4.

(i) $I_{1,n}(\xi)I_{3,n}(\xi) - NI_{2,n}(\xi)\hat{P}_{n-1}^{[2]}(\xi) = -I_{2,n}(\xi)\hat{P}_{n-1}^{[2]}(\xi)(N + \alpha_n\beta_n)$, where

$$0 < \alpha_n = \frac{I_{1,n}(\xi)}{\hat{P}_{n-1}^{[2]}(\xi)} < d_0^1 \quad \text{and} \quad \frac{d_0^3}{d_0^2} < -\frac{\hat{P}_{n-1}^{[2]'}(\xi)}{\hat{P}_{n-1}^{[2]}(\xi)} = \frac{I_{2,n}(\xi)}{I_{3,n}(\xi)} = \frac{1}{\beta_n} < -\frac{n}{\xi}.$$

(ii) $NI_{2,n}(\xi)\hat{P}'_n(\xi) - MI_{3,n}(\xi)\hat{P}_n(\xi) = I_{2,n}(\xi)\hat{P}'_n(\xi)(N + M\beta_n\gamma_n)$, where

$$\frac{d_0^1}{c_0} < -\frac{\hat{P}'_n(\xi)}{\hat{P}_n(\xi)} = \frac{1}{\gamma_n} < -\frac{n}{\xi}.$$

(iii) $MN\hat{P}_n(\xi)\hat{P}_{n-1}^{[2]}(\xi) - NI_{1,n}(\xi)\hat{P}'_n(\xi) = N\hat{P}_n(\xi)\hat{P}_{n-1}^{[2]}(\xi)\left(M + \frac{\alpha_n}{\gamma_n}\right)$.

Proof. (i) From the Christoffel–Darboux formula for polynomials $\{\hat{P}_n^{[2]}\}_{n \geq 0}$ we have

$$\begin{aligned} (16) \quad (x - \xi) \sum_{k=0}^n \frac{\hat{P}_k^{[2]}(x)\hat{P}_k^{[2]}(y)}{\|\hat{P}_k^{[2]}\|_{\omega,2}^2} - \sum_{k=0}^n \frac{\hat{P}_k^{[2]}(x)}{\|\hat{P}_k^{[2]}\|_{\omega,2}^2} (y - \xi)\hat{P}_k^{[2]}(y) \\ = \frac{1}{\|\hat{P}_n^{[2]}\|_{\omega,2}^2} (\hat{P}_{n+1}^{[2]}(x)\hat{P}_n^{[2]}(y) - \hat{P}_n^{[2]}(x)\hat{P}_{n+1}^{[2]}(y)). \end{aligned}$$

If we multiply (16) by $(y - \xi)\omega(y)$ and integrate over $(0, \infty)$, evaluation at $x = \xi$ yields

$$\begin{aligned}
 - \sum_{k=0}^n \frac{\hat{P}_k^{[2]}(\xi)}{\|\hat{P}_k^{[2]}\|_{\omega,2}^2} \int_0^\infty (y - \xi)^2 \omega(y) \hat{P}_k^{[2]}(y) dy \\
 = \frac{1}{\|\hat{P}_n^{[2]}\|_{\omega,2}^2} (\hat{P}_{n+1}^{[2]}(\xi) I_{1,n+1}(\xi) - \hat{P}_n^{[2]}(\xi) I_{1,n+2}(\xi)).
 \end{aligned}$$

Since

$$\int_0^\infty (y - \xi)^2 \omega(y) \hat{P}_k^{[2]}(y) dy = 0 \quad \text{for } k = 1, 2, \dots, n$$

and $\hat{P}_0^{[2]} = 1$, the left-hand side is negative. Therefore,

$$\hat{P}_{n+1}^{[2]}(\xi) I_{1,n+1}(\xi) - \hat{P}_n^{[2]}(\xi) I_{1,n+2}(\xi) < 0.$$

From (5) we have

$$\text{sgn } \hat{P}_{n+1}^{[2]}(\xi) = (-1)^{n+1} \quad \text{and} \quad \text{sgn } \hat{P}_n^{[2]}(\xi) = (-1)^n.$$

Thus, $\hat{P}_{n+1}^{[2]}(\xi) \hat{P}_n^{[2]}(\xi)$ is negative and, as a consequence,

$$\frac{I_{1,n+2}(\xi)}{\hat{P}_{n+1}^{[2]}(\xi)} < \frac{I_{1,n+1}(\xi)}{\hat{P}_n^{[2]}(\xi)}.$$

Using this relation recursively, we get

$$\frac{I_{1,n}(\xi)}{\hat{P}_{n-1}^{[2]}(\xi)} < I_{1,1}(\xi) = d_0^1.$$

On the other hand, (5) and (13) imply that $\text{sgn } I_{1,n}(\xi) = (-1)^{n+1}$; therefore,

$$0 < \frac{I_{1,n}(\xi)}{\hat{P}_{n-1}^{[2]}(\xi)} < d_0^1.$$

From (16)

$$0 < \sum_{k=0}^n \frac{(\hat{P}_k^{[2]}(\xi))^2}{\|\hat{P}_k^{[2]}\|_{\omega,2}^2} = \frac{1}{\|\hat{P}_n^{[2]}\|_{\omega,2}^2} (\hat{P}_{n+1}^{[2]'}(\xi) \hat{P}_n^{[2]}(\xi) - \hat{P}_n^{[2]'}(\xi) \hat{P}_{n+1}^{[2]}(\xi)).$$

Since $\hat{P}_{n+1}^{[2]}(\xi) \hat{P}_n^{[2]}(\xi)$ is negative this yields

$$\frac{\hat{P}_{n+1}^{[2]'}(\xi)}{\hat{P}_{n+1}^{[2]}(\xi)} < \frac{\hat{P}_n^{[2]'}(\xi)}{\hat{P}_n^{[2]}(\xi)}.$$

Using this relation recursively, we obtain

$$\frac{\hat{P}_{n+1}^{[2]'}(\xi)}{\hat{P}_{n+1}^{[2]}(\xi)} < \frac{\hat{P}_1^{[2]'}(\xi)}{\hat{P}_1^{[2]}(\xi)} = -\frac{d_0^3}{d_0^2}.$$

Let $0 < x_{1,n}^{[2]} < x_{2,n}^{[2]} < \dots < x_{n,n}^{[2]}$ denote the zeros of $\hat{P}_n^{[2]}$. Then

$$-\frac{\hat{P}_n^{[2]'}(\xi)}{\hat{P}_n^{[2]}(\xi)} = \frac{1}{x_{1,n}^{[2]} - \xi} + \frac{1}{x_{2,n}^{[2]} - \xi} + \dots + \frac{1}{x_{n,n}^{[2]} - \xi} < -\frac{n}{\xi}.$$

Statements (ii) and (iii) can be proved in a similar way as (i). \square

Proposition 5. *Let $M, N \geq 0$ and not both zero. Then*

$$\operatorname{sgn} A_{n,1} = -1 \quad \text{and} \quad \operatorname{sgn} A_{n,2} = -\operatorname{sgn} N.$$

Proof. From (5) and Proposition 4

$$\operatorname{sgn} A_{n,1} = -\operatorname{sgn} \frac{\hat{P}_n'(\xi)}{\hat{P}_{n-1}^{[2]}(\xi)} = \operatorname{sgn} \left(-\frac{\hat{P}_n'(\xi)}{\hat{P}_n(\xi)} \right) \operatorname{sgn} \frac{\hat{P}_n(\xi)}{\hat{P}_{n-1}^{[2]}(\xi)} = -1.$$

In a similar way,

$$\begin{aligned} \operatorname{sgn} A_{n,2} &= -\operatorname{sgn} N \operatorname{sgn} \frac{\hat{P}_n(\xi)}{I_{2,n}} \\ &= \operatorname{sgn} N \operatorname{sgn} \left(-\frac{\hat{P}_{n-1}^{[2]}(\xi)}{\hat{P}_{n-1}^{[2]'}(\xi)} \right) \operatorname{sgn} \frac{\hat{P}_n(\xi) \hat{P}_{n-2}(\xi) \hat{P}_{n-2}^{[1]}(\xi) \hat{P}_{n-2}^{[2]}(\xi) \hat{P}_{n-2}^{[3]}(\xi)}{\hat{P}_{n-1}(\xi) \hat{P}_{n-1}^{[1]}(\xi) \hat{P}_{n-1}^{[2]}(\xi)} \\ &= -\operatorname{sgn} N. \end{aligned} \quad \square$$

The zeros. We now analyze the zeros of the polynomials \hat{S}_n . The techniques are the same as those used by Meijer [1993a; 1993b].

Theorem 2. *The discrete Sobolev orthogonal polynomial \hat{S}_n has n real simple zeros and at most one of them is outside of $[\xi, \infty)$.*

Proof. Since for $N = 0$, \hat{S}_n is a standard orthogonal polynomial, in the sequel we will consider the cases when $N > 0$ and $M \geq 0$. Let $v_1 < v_2 < \dots < v_k$ be the zeros of $\hat{S}_n(x)$ on (ξ, ∞) with odd multiplicity. Let us introduce the polynomial

$$\phi(x) = (x - v_1)(x - v_2) \cdots (x - v_k).$$

Notice that $\phi(\xi)$ and $\phi'(\xi)$ have opposite signs and $\phi(x)\hat{S}_n(x)$ does not change sign on $[\xi, \infty)$. If $\deg \phi \leq n - 2$, then

$$0 = \langle \phi, \hat{S}_n \rangle_S = \int_0^\infty \omega(x) \phi(x) \hat{S}_n(x) dx + M \phi(\xi) \hat{S}_n(\xi) + N \phi'(\xi) \hat{S}_n'(\xi)$$

and

$$0 = \langle (\cdot - \xi)\phi, \hat{S}_n \rangle_S = \int_0^\infty \omega(x)(x - \xi)\phi(x)\hat{S}_n(x) dx + N\phi(\xi)\hat{S}'_n(\xi).$$

This means that $\phi'(\xi)\hat{S}'_n(\xi)$ and $\phi(\xi)\hat{S}'_n(\xi)$ have the same sign, and therefore $\phi'(\xi)$ and $\phi(\xi)$ have the same sign. This yields a contradiction.

As a conclusion, $\deg \phi = n - 1$ or $\deg \phi = n$, which proves our statement. \square

Next, we prove that the zeros of $\hat{S}_n(x)$ interlace with the zeros of $\hat{P}_{n-1}^{[2]}(x)$ if $\hat{S}_n(x)$ has a zero outside $[\xi, \infty)$. Notice that, by Theorem 1, $\hat{S}_n(\xi) \neq 0$.

Theorem 3. Denote by $v_{r,n}$, $r = 1, 2, \dots, n$, the zeros of $\hat{S}_n(x)$ in increasing order. Suppose that $v_{1,n} < \xi$. Then $2\xi - x_{1,n-1}^{[2]} < v_{1,n} < \xi$ and

$$\xi < v_{2,n} < x_{1,n-1}^{[2]} < \dots < v_{n,n} < x_{n-1,n-1}^{[2]}.$$

Proof. From Theorem 1 we have

$$\hat{S}_n(x_{r,n-1}^{[2]}) = \hat{P}_n(x_{r,n-1}^{[2]}) + A_{n,2}(x_{r,n-1}^{[2]} - \xi)^2 \hat{P}_{n-2}^{[4]}(x_{r,n-1}^{[2]}), \quad r = 1, 2, \dots, n - 1.$$

Then from Proposition 3(iii) and Proposition 5 we get

$$\operatorname{sgn} \hat{S}_n(x_{r,n-1}^{[2]}) = (-1)^{n-r}, \quad r = 1, 2, \dots, n - 1,$$

On the other hand, from (5) and Theorem 1,

$$\operatorname{sgn} \hat{S}_n(\xi) = \operatorname{sgn} \hat{P}_n(\xi) = (-1)^n.$$

Therefore, every interval $(\xi, x_{1,n-1}^{[2]})$ and $(x_{r,n-1}^{[2]}, x_{r+1,n-1}^{[2]})$, for $r = 1, \dots, n - 2$, contains an odd number of zeros of $\hat{S}_n(x)$. Since \hat{S}_n has n real zeros and at most one of them is outside of (ξ, ∞) , then

$$\xi < v_{2,n} < x_{1,n-1}^{[2]} < \dots < v_{n,n} < x_{n-1,n-1}^{[2]}.$$

Now, we will prove that $2\xi - x_{1,n-1}^{[2]} < v_{1,n} < \xi$. Let

$$\hat{S}_n(x) = (x - v_{1,n})(x - v_{2,n}) \cdots (x - v_{n,n}).$$

By Theorem 1 and Proposition 4,

$$\hat{S}'_n(\xi) = \hat{P}'_n(\xi) + A_{n,1} \hat{P}_{n-2}^{[2]}(\xi) = \frac{\beta_n \hat{P}_n(\xi)(M + \alpha_n/\gamma_n)}{N + \alpha_n \beta_n}.$$

Therefore,

$$\operatorname{sgn} \hat{S}'_n(\xi) = \operatorname{sgn} \hat{P}'_n(\xi) = \operatorname{sgn} \hat{S}_n(\xi)$$

and

$$0 < \frac{\hat{S}'_n(\xi)}{\hat{S}_n(\xi)} = \frac{1}{\xi - v_{1,n}} - \frac{1}{v_{2,n} - \xi} - \dots - \frac{1}{v_{n,n} - \xi}.$$

Hence $\frac{1}{\xi - \nu_{1,n}} > \frac{1}{\nu_{2,n} - \xi}$, which implies successively

$$x_{1,n-1}^{[2]} - \xi > \nu_{2,n} - \xi > \xi - \nu_{1,n} \quad \text{and} \quad 2\xi - x_{1,n-1}^{[2]} < \nu_{1,n}.$$

Our statement follows. □

4. Discrete Laguerre–Sobolev orthogonal polynomials: asymptotics

Laguerre polynomials. For $\alpha \in \mathbb{R}$, the Laguerre polynomials are defined by

$$L_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} \frac{(-x)^k}{k!}.$$

For $\alpha > -1$, the $\{L_n^{(\alpha)}(x)\}_{n \geq 0}$ are orthogonal on $[0, +\infty)$ with respect to the weight function $\omega(x) = x^\alpha e^{-x}$ [Szegő 1975, Chapter V]. Let $\{L_n^{(\alpha,k)}\}_{n=0}^\infty$, $k \in \mathbb{N}$, denote the sequence of polynomials orthogonal with respect to the modified Laguerre weight $(x - \xi)^k \omega(x)$, $\xi < 0$, normalized by the condition that $L_n^{(\alpha,k)}$ has the same leading coefficient as the classical Laguerre orthogonal polynomial $L_n^{(\alpha)} = L_n^{(\alpha,0)}$. That is, $k(L_n^{(\alpha,k)}) = (-1)^n/n!$.

We summarize some properties of the $L_n^{(\alpha,k)}(x)$, $k \in \mathbb{N} \cup \{0\}$, to be used later.

Proposition 6 [Fejzullahu 2011]. (i) For $\alpha > -1$,

$$\|L_n^{(\alpha)}\|_\alpha^2 = \int_0^\infty (L_n^{(\alpha)}(x))^2 x^\alpha e^{-x} dx = \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)}.$$

(ii) For every $n \in \mathbb{N}$,

$$(L_n^{(\alpha)}(x))' = -L_{n-1}^{(\alpha+1)}(x).$$

(iii) (Perron's formula) Let $\alpha \in \mathbb{R}$. Then

$$L_n^{(\alpha)}(x) = 2^{-1} \pi^{-1/2} e^{x/2} (-x)^{-\alpha/2-1/4} n^{\alpha/2-1/4} e^{2\sqrt{-nx}} (1 + O(n^{-1/2})).$$

This relation holds for x in the complex plane cut along the positive real semi-axis; both $(-x)^{-\alpha/2-1/4}$ and $\sqrt{-x}$ must be taken real and positive if $x < 0$. The bound of the remainder holds uniformly in every closed domain which does not overlap the positive real semi-axis.

Moreover, we get the outer ratio asymptotics

$$\lim_{n \rightarrow \infty} n^{(l-j)/2} \frac{L_{n+k}^{(\alpha+j)}(x)}{L_{n+h}^{(\alpha+l)}(x)} = (-x)^{(l-j)/2}, \quad j, l \in \mathbb{R}, \quad h, k \in \mathbb{Z},$$

$$\lim_{n \rightarrow \infty} \frac{L_n^{(\alpha,k)}(x)}{n^{k/2} L_n^{(\alpha)}(x)} = \frac{1}{(\sqrt{-x} + \sqrt{-\xi})^k},$$

uniformly on compact subsets of $\mathbb{C} \setminus [0, \infty)$.

(iv) (Mehler–Heine formula) *Uniformly on compact subsets of \mathbb{C} , we have*

$$\lim_{n \rightarrow \infty} \frac{L_n^{(\alpha)}(x/(n+j))}{n^\alpha} = x^{-\alpha/2} J_\alpha(2\sqrt{x})$$

and

$$\lim_{n \rightarrow \infty} \frac{L_n^{(\alpha,k)}(x/(n+j))}{n^{\alpha+k/2}} = \frac{1}{(\sqrt{-\xi})^k} x^{-\alpha/2} J_\alpha(2\sqrt{x})$$

where $j \in \mathbb{N} \cup 0$ and J_α is the Bessel function of the first kind.

(v) (Plancherel–Rotach type outer asymptotics for $L_n^{(\alpha,N)}$) *Uniformly on compact subsets of $\mathbb{C} \setminus [0, 4]$ and uniformly on $j \in \mathbb{N} \cup \{0\}$, we have*

$$\lim_{n \rightarrow \infty} \frac{L_{n-1}^{(\alpha)}((n+j)x)}{L_n^{(\alpha)}((n+j)x)} = -\frac{1}{\phi((x-2)/2)}$$

and

$$\lim_{n \rightarrow \infty} \frac{L_n^{(\alpha,N)}((n+j)x)}{L_n^{(\alpha)}((n+j)x)} = \left(\frac{\phi((x-2)/2) + 1}{x} \right)^N,$$

where ϕ is the conformal mapping of $\mathbb{C} \setminus [-1, 1]$ onto the exterior of the unit circle given by

$$\phi(x) = x + \sqrt{x^2 - 1}, \quad x \in \mathbb{C} \setminus [-1, 1],$$

with $\sqrt{x^2 - 1} > 0$ when $x > 1$.

Proposition 7. $L_n^{(\alpha,2)'}(\xi) \cong \frac{n}{4\xi} L_n^{(\alpha+1)}(\xi).$

Proof. Using integration by parts we have

$$\int_0^\infty (L_n^{(\alpha,2)}(x))' L_k^{(\alpha+1,3)}(x) (x-\xi)^3 x^{\alpha+1} e^{-x} dx = \begin{cases} 0 & \text{if } k \leq n-3, \\ n(n-1) \|\hat{L}_n^{(\alpha,2)}\|_{\alpha,2}^2 & \text{if } k = n-2. \end{cases}$$

Therefore,

$$(L_n^{(\alpha,2)}(x))' = -L_{n-1}^{(\alpha+1,3)}(x) + H_n L_{n-2}^{(\alpha+1,3)}(x),$$

where

$$H_n = \frac{n(n-1) \|\hat{L}_n^{(\alpha,2)}\|_{\alpha,2}^2}{\|\hat{L}_{n-2}^{(\alpha+1,3)}\|_{\alpha+1,3}^2}.$$

Using (8) and Proposition 6(iii),

$$\begin{aligned} H_n &= \frac{(n+1)^2(n+\alpha)}{(n-1)^3} \frac{L_{n-2}^{(\alpha+1,2)}(\xi)}{L_{n-1}^{(\alpha+1,2)}(\xi)} \prod_{i=1}^2 \frac{L_{n-2}^{(\alpha+1,i-1)}(\xi)}{L_{n-1}^{(\alpha+1,i-1)}(\xi)} \frac{L_{n+1}^{(\alpha,i-1)}(\xi)}{L_n^{(\alpha,i-1)}(\xi)} \\ &= \frac{L_{n-2}^{(\alpha+1,2)}(\xi)}{L_{n-1}^{(\alpha+1,2)}(\xi)} \prod_{i=1}^2 \frac{L_{n-2}^{(\alpha+1,i-1)}(\xi)}{L_{n-1}^{(\alpha+1,i-1)}(\xi)} \frac{L_{n+1}^{(\alpha,i-1)}(\xi)}{L_n^{(\alpha,i-1)}(\xi)} + O\left(\frac{1}{n}\right). \end{aligned}$$

On the other hand, [Fejzullahu 2011, Proposition 2.2] gives

$$(17) \quad (L_n^{(\alpha,2)}(x))' = -L_{n-1}^{(\alpha,3)}(x) + G_n L_{n-2}^{(\alpha+1,3)}(x),$$

where

$$\begin{aligned} G_n &= H_n - \frac{n^3}{(n-1)^3} \prod_{i=1}^3 \frac{L_{n-2}^{(\alpha+1,i-1)}(\xi)}{L_{n-1}^{(\alpha+1,i-1)}(\xi)} \frac{L_n^{(\alpha,i-1)}(\xi)}{L_{n-1}^{(\alpha,i-1)}(\xi)} \\ &= \prod_{i=1}^3 \frac{L_{n-2}^{(\alpha+1,i-1)}(\xi)}{L_{n-1}^{(\alpha+1,i-1)}(\xi)} \left(\frac{L_{n+1}^{(\alpha)}(\xi) L_{n+1}^{(\alpha,1)}(\xi)}{L_n^{(\alpha)}(\xi) L_n^{(\alpha,1)}(\xi)} - \frac{L_n^{(\alpha)}(\xi) L_n^{(\alpha,1)}(\xi) L_n^{(\alpha,2)}(\xi)}{L_{n-1}^{(\alpha)}(\xi) L_{n-1}^{(\alpha,1)}(\xi) L_{n-1}^{(\alpha,2)}(\xi)} \right) + O\left(\frac{1}{n}\right). \end{aligned}$$

Again, from [Fejzullahu 2011, Proposition 2.2],

$$\begin{aligned} \frac{L_{n+1}^{(\alpha)}(\xi) L_{n+1}^{(\alpha,1)}(\xi)}{L_n^{(\alpha)}(\xi) L_n^{(\alpha,1)}(\xi)} &= \frac{L_{n+1}^{(\alpha)}(\xi) L_{n+1}^{(\alpha-1,1)}(\xi)}{L_n^{(\alpha)}(\xi) L_n^{(\alpha,1)}(\xi)} + \frac{L_{n+2}^{(\alpha-1)}(\xi)}{L_{n+1}^{(\alpha-1)}(\xi)} + O\left(\frac{1}{n}\right), \\ \frac{L_n^{(\alpha)}(\xi) L_n^{(\alpha,1)}(\xi) L_n^{(\alpha,2)}(\xi)}{L_{n-1}^{(\alpha)}(\xi) L_{n-1}^{(\alpha,1)}(\xi) L_{n-1}^{(\alpha,2)}(\xi)} &= \frac{L_n^{(\alpha)}(\xi) L_n^{(\alpha,1)}(\xi) L_n^{(\alpha-1,2)}(\xi)}{L_{n-1}^{(\alpha)}(\xi) L_{n-1}^{(\alpha,1)}(\xi) L_{n-1}^{(\alpha,2)}(\xi)} \\ &\quad + \frac{L_{n+1}^{(\alpha-1)}(\xi) L_{n+1}^{(\alpha-1,1)}(\xi)}{L_n^{(\alpha-1)}(\xi) L_n^{(\alpha-1,1)}(\xi)} + O\left(\frac{1}{n}\right), \end{aligned}$$

and

$$\begin{aligned} \frac{L_{n+2}^{(\alpha-1)}(\xi)}{L_{n+1}^{(\alpha-1)}(\xi)} - \frac{L_{n+1}^{(\alpha-1)}(\xi) L_{n+1}^{(\alpha-1,1)}(\xi)}{L_n^{(\alpha-1)}(\xi) L_n^{(\alpha-1,1)}(\xi)} &= \frac{L_{n+2}^{(\alpha-2)}(\xi)}{L_{n+1}^{(\alpha-1)}(\xi)} + 1 \\ &\quad - \frac{L_{n+1}^{(\alpha-1)}(\xi) L_{n+1}^{(\alpha-2,1)}(\xi)}{L_n^{(\alpha-1)}(\xi) L_n^{(\alpha-1,1)}(\xi)} - \frac{L_{n+2}^{(\alpha-2)}(\xi)}{L_{n+1}^{(\alpha-2)}(\xi)} + O\left(\frac{1}{n}\right) \\ &= \frac{L_{n+2}^{(\alpha-2)}(\xi)}{L_{n+1}^{(\alpha-1)}(\xi)} - \frac{L_{n+1}^{(\alpha-1)}(\xi) L_{n+1}^{(\alpha-2,1)}(\xi)}{L_n^{(\alpha-1)}(\xi) L_n^{(\alpha-1,1)}(\xi)} - \frac{L_{n+2}^{(\alpha-3)}(\xi)}{L_{n+1}^{(\alpha-2)}(\xi)} + O\left(\frac{1}{n}\right). \end{aligned}$$

Therefore, by using Proposition 6(iii),

$$\sqrt{n}G_n \cong -\sqrt{-\xi}.$$

and taking into account (17) the result follows. □

Discrete Laguerre–Sobolev orthogonal polynomials. Let $\{S_n\}_{n \geq 0}$ be the sequence of polynomials orthogonal with respect to the discrete Sobolev inner product (12), where $\omega(x) = x^\alpha e^{-x}$ and $\xi < 0$, normalized by the condition that S_n has the same leading coefficient as the classical Laguerre orthogonal polynomial $L_n^{(\alpha)}$, i.e., $k(S_n) = (-1)^n/n!$.

Theorem 4. *Let $M \geq 0$ and $N \geq 0$. There are real constants $B_{n,0}$, $B_{n,1}$, and $B_{n,2}$ such that*

$$(18) \quad S_n(x) = B_{n,0}L_n^{(\alpha)}(x) + B_{n,1}(x - \xi)L_{n-1}^{(\alpha,2)}(x) + B_{n,2}(x - \xi)^2L_{n-2}^{(\alpha,4)}(x),$$

where $B_{n,0} = \frac{1}{1 + A_{n,1} + A_{n,2}}$, $B_{n,1} = -\frac{A_{n,1}}{n(1 + A_{n,1} + A_{n,2})}$, and

$$B_{n,2} = \frac{A_{n,2}}{n(n-1)(1 + A_{n,1} + A_{n,2})}.$$

Moreover:

(i) *If $M > 0$ and $N > 0$, then*

$$(19) \quad B_{n,0} \cong \frac{8\xi n^\alpha}{M(L_n^{(\alpha)}(\xi))^2}, \quad B_{n,1} \cong -\frac{32\xi\sqrt{-\xi}n^{\alpha-1/2}}{M(L_n^{(\alpha)}(\xi))^2}, \quad B_{n,2} \cong \frac{1}{n^2}.$$

(ii) *If $M = 0$ and $N > 0$, then*

$$B_{n,0} \cong \frac{1}{4\sqrt{-\xi}n}, \quad B_{n,1} \cong -\frac{1}{n}, \quad B_{n,2} \cong \frac{1}{4n^2\sqrt{-\xi}n}.$$

(iii) *If $M > 0$ and $N = 0$, then*

$$B_{n,0} \cong \frac{\sqrt{-\xi}}{Mn^{1/2-\alpha}(L_{n-1}^{(\alpha)}(\xi))^2}, \quad B_{n,1} \cong -\frac{1}{n}, \quad B_{n,2} = 0.$$

Proof. From Theorem 1,

$$S_n(x) = \frac{(-1)^n \hat{S}_n(x)}{n!(1 + A_{n,1} + A_{n,2})}$$

and, as a consequence,

$$S_n(x) = B_{n,0}L_n^{(\alpha)}(x) + B_{n,1}(x - \xi)L_{n-1}^{(\alpha,2)}(x) + B_{n,2}(x - \xi)^2L_{n-2}^{(\alpha,4)}(x),$$

where $B_{n,0}$, $B_{n,1}$, and $B_{n,2}$ are as in the statement of the theorem.

Now, from Proposition 4 we can obtain the behavior of the coefficients $B_{n,0}$, $B_{n,1}$ and $B_{n,2}$ for n large enough. In order to estimate $A_{n,1}$ and $A_{n,2}$, first we

compute $\alpha_n\beta_n$, α_n/γ_n , $\beta_n\gamma_n$ and $I_{2,n}(\xi)$. From (13) and Proposition 6, we can write

$$\begin{aligned} \alpha_n\beta_n &= -\frac{I_{1,n}(\xi)}{\hat{L}_{n-1}^{(\alpha,2)' }(\xi)} = \frac{\hat{L}_n^{(\alpha)}(\xi)}{\hat{L}_{n-1}^{(\alpha)}(\xi)\hat{L}_{n-1}^{(\alpha,1)}(\xi)\hat{L}_{n-1}^{(\alpha,2)' }(\xi)} \|\hat{L}_{n-1}^{(\alpha)}\|_\alpha^2 \\ &= -\frac{\Gamma(n+\alpha)}{\Gamma(n)} \frac{nL_n^{(\alpha)}(\xi)}{L_{n-1}^{(\alpha)}(\xi)L_{n-1}^{(\alpha,1)}(\xi)L_{n-1}^{(\alpha,2)' }(\xi)} \cong \frac{8(-\xi)^{3/2}n^{\alpha-1/2}}{L_n^{(\alpha)}(\xi)L_n^{(\alpha+1)}(\xi)}, \\ \frac{\alpha_n}{\gamma_n} &= -\frac{I_{1,n}(\xi)\hat{L}_n^{(\alpha)' }(\xi)}{\hat{L}_n^{(\alpha)}(\xi)\hat{L}_{n-1}^{(\alpha,2)}(\xi)} = \frac{\hat{L}_n^{(\alpha)' }(\xi)}{\hat{L}_{n-1}^{(\alpha)}(\xi)\hat{L}_{n-1}^{(\alpha,1)}(\xi)\hat{L}_{n-1}^{(\alpha,2)}(\xi)} \|\hat{L}_{n-1}^{(\alpha)}\|_\alpha^2 \\ &= \frac{\Gamma(n+\alpha)}{\Gamma(n)} \frac{nL_{n-1}^{(\alpha+1)}(\xi)}{L_{n-1}^{(\alpha)}(\xi)L_{n-1}^{(\alpha,1)}(\xi)L_{n-1}^{(\alpha,2)}(\xi)} \cong \frac{8(-\xi)^{3/2}n^{\alpha-1/2}L_n^{(\alpha+1)}(\xi)}{\left(L_n^{(\alpha)}(\xi)\right)^3}, \\ \beta_n\gamma_n &= \alpha_n\beta_n \frac{\gamma_n}{\alpha_n} \cong \left(\frac{L_n^{(\alpha)}(\xi)}{L_n^{(\alpha+1)}(\xi)}\right)^2 \cong -\frac{\xi}{n}, \\ I_{2,n}(\xi) &\cong (-1)^{n-1}(n-2)!n^{\alpha+3} \frac{L_{n-1}^{(\alpha)}(\xi)L_{n-1}^{(\alpha,1)}(\xi)L_{n-1}^{(\alpha,2)' }(\xi)}{L_{n-2}^{(\alpha)}(\xi)L_{n-2}^{(\alpha,1)}(\xi)L_{n-2}^{(\alpha,2)}(\xi)L_{n-3}^{(\alpha,3)}(\xi)} \\ &\cong \frac{8\xi(-1)^{n-1}(n-2)!n^{\alpha+2}}{L_n^{(\alpha)}(\xi)}. \end{aligned}$$

Next, we will analyze the following three situations.

(i) Let $M > 0$ and $N > 0$. Then,

$$A_{n,1} \cong -\frac{\hat{L}_n^{(\alpha)' }(\xi)}{\hat{L}_{n-1}^{(\alpha,2)}(\xi)} = \frac{nL_n^{(\alpha)' }(\xi)}{L_{n-1}^{(\alpha,2)}(\xi)} = -\frac{nL_{n-1}^{(\alpha+1)}(\xi)}{L_{n-1}^{(\alpha,2)}(\xi)} \cong -4\sqrt{-\xi n}$$

and

$$A_{n,2} \cong -\frac{M\hat{L}_n^{(\alpha)}(\xi)}{I_{n,2}(\xi)} \cong \frac{M(L_n^{(\alpha)}(\xi))^2}{8\xi n^\alpha}.$$

Therefore,

$$B_{n,0} \cong \frac{8\xi n^\alpha}{M(L_n^{(\alpha)}(\xi))^2}, \quad B_{n,1} \cong \frac{32\xi\sqrt{-\xi}n^{\alpha-1/2}}{M(L_n^{(\alpha)}(\xi))^2}, \quad B_{n,2} \cong \frac{1}{n^2}.$$

(ii) Let $M = 0$ and $N > 0$. Then,

$$A_{n,1} \cong -4\sqrt{-\xi n} \quad \text{and} \quad A_{n,2} = -\frac{\hat{L}_n^{(\alpha)}(\xi)}{I_{n,2}(\xi)} \frac{\alpha_n}{\gamma_n} \cong -1.$$

Therefore,

$$B_{n,0} \cong -\frac{1}{4\sqrt{-\xi n}}, \quad B_{n,1} \cong -\frac{1}{n}, \quad B_{n,2} \cong \frac{1}{4n^2\sqrt{-\xi n}}.$$

(iii) Let $M > 0$ and $N = 0$. Then,

$$A_{n,1} = \frac{M\hat{L}_n^{(\alpha)}(\xi)}{I_{n,1}(\xi)} = -\frac{M\hat{L}_{n-1}^{(\alpha)}(\xi)\hat{L}_{n-1}^{(\alpha,1)}(\xi)}{\|L_{n-1}^{(\alpha)}\|_\alpha^2} \cong -\frac{Mn^{1/2-\alpha}}{\sqrt{-\xi}}(L_{n-1}^{(\alpha)}(\xi))^2, \quad A_{n,2} = 0.$$

Therefore,

$$B_{n,0} \cong -\frac{\sqrt{-\xi}}{Mn^{1/2-\alpha}(L_{n-1}^{(\alpha)}(\xi))^2}, \quad B_{n,1} \cong -\frac{1}{n}, \quad B_{n,2} = 0. \quad \square$$

Next we deduce several asymptotic properties for discrete Laguerre–Sobolev polynomials when $M, N \geq 0$. (For $M > 0$ and $N = 0$, the same asymptotic results for corresponding Laguerre-type polynomials has been deduced in [Dueñas et al. 2011] and [Fejzullahu and Zejnullahu 2010].)

Theorem 5. (i) (Outer relative asymptotics) *Uniformly on compact subsets of $\mathbb{C} \setminus [0, \infty)$ we have:*

- If $M > 0$ and $N > 0$, then

$$\lim_{n \rightarrow \infty} \frac{S_n(x)}{L_n^{(\alpha)}(x)} = \left(\frac{\sqrt{-x} - \sqrt{-\xi}}{\sqrt{-x} + \sqrt{-\xi}} \right)^2.$$

Notice that, according to the Hurwitz’s Theorem, the point ξ attracts two negative zeros of $S_n(x)$ for n large enough.

- If $M = 0$ and $N > 0$ or $M > 0$ and $N = 0$, then

$$\lim_{n \rightarrow \infty} \frac{S_n(x)}{L_n^{(\alpha)}(x)} = \frac{\sqrt{-x} - \sqrt{-\xi}}{\sqrt{-x} + \sqrt{-\xi}}.$$

Notice that, according to the Hurwitz’s Theorem, the point ξ attracts one negative zero of $S_n(x)$ for n large enough.

(ii) (Mehler–Heine formula)

- If $M > 0$ and $N > 0$

$$\lim_{n \rightarrow \infty} \frac{S_n(x/n)}{n^\alpha} = x^{-\alpha/2} J_\alpha(2\sqrt{x}),$$

- If $M = 0$ and $N > 0$ or $M > 0$ and $N = 0$

$$\lim_{n \rightarrow \infty} \frac{S_n(x/n)}{n^\alpha} = -x^{-\alpha/2} J_\alpha(2\sqrt{x}),$$

uniformly on compact subsets of \mathbb{C} .

(iii) (Plancherel–Rotach type outer asymptotics for S_n)

• If $M \geq 0$ and $N \geq 0$, then

$$\lim_{n \rightarrow \infty} \frac{S_n(nx)}{L_n^{(\alpha)}(nx)} = 1,$$

uniformly on compact subsets of $\mathbb{C} \setminus [0, 4]$.

Proof. We will prove the theorem when $M > 0$ and $N > 0$. The proofs of the other cases can be done in a similar way.

(i) From (18)

$$\frac{S_n(x)}{L_n^{(\alpha)}(x)} = B_{n,0} + nB_{n,1}(x - \xi) \frac{L_{n-1}^{(\alpha,2)}(x)}{nL_n^{(\alpha)}(x)} + n^2B_{n,2}(x - \xi)^2 \frac{L_{n-2}^{(\alpha,4)}(x)}{n^2L_n^{(\alpha)}(x)}.$$

Now, Proposition 6(iii) and (19) yield

$$\lim_{n \rightarrow \infty} \frac{S_n(x)}{L_n^{(\alpha)}(x)} = (x - \xi)^2 \lim_{n \rightarrow \infty} \frac{L_{n-2}^{(\alpha,4)}(x)}{n^2L_n^{(\alpha)}(x)} = \left(\frac{\sqrt{-x} - \sqrt{-\xi}}{\sqrt{-x} + \sqrt{-\xi}} \right)^2.$$

(ii) Scaling the variable as $x \rightarrow x/n$ in (18) then dividing by n^α we get

$$\begin{aligned} & \frac{S_n(x/n)}{n^\alpha} \\ &= B_{n,0} \frac{L_n^{(\alpha)}(x/n)}{n^\alpha} + nB_{n,1}(x/n - \xi) \frac{L_{n-1}^{(\alpha,2)}(x/n)}{n^{\alpha+1}} + n^2B_{n,2}(x/n - \xi)^2 \frac{L_{n-2}^{(\alpha,4)}(x/n)}{n^{\alpha+2}}. \end{aligned}$$

Now, Proposition 6(iv) and (19) yield

$$\lim_{n \rightarrow \infty} \frac{S_n(x/n)}{n^\alpha} = (-\xi)^2 \lim_{n \rightarrow \infty} \frac{L_{n-2}^{(\alpha,4)}(x)}{n^{\alpha+2}} = x^{-\alpha/2} J_\alpha(2\sqrt{x}).$$

(iii) Dividing (18) by $L_n^{(\alpha)}(x)$ then scaling the variable as $x \rightarrow nx$ we get

$$\begin{aligned} \frac{S_n(nx)}{L_n^{(\alpha)}(nx)} &= B_{n,0} + nB_{n,1} \frac{nx - \xi}{n} \frac{L_{n-1}^{(\alpha,2)}(nx)}{L_{n-1}^{(\alpha)}(nx)} \frac{L_{n-1}^{(\alpha)}(nx)}{L_n^{(\alpha)}(nx)} \\ &\quad + n^2B_{n,2} \frac{(nx - \xi)^2}{n^2} \frac{L_{n-2}^{(\alpha,4)}(nx)}{L_{n-2}^{(\alpha)}(nx)} \frac{L_{n-2}^{(\alpha)}(nx)}{L_n^{(\alpha)}(nx)}. \end{aligned}$$

From Proposition 6(v) and (19)

$$\lim_{n \rightarrow \infty} \frac{S_n(nx)}{L_n^{(\alpha)}(nx)} = x^2 \left(\frac{\phi((x-2)/2) + 1}{x} \right)^4 \frac{1}{(\phi((x-2)/2))^2}.$$

Now, using the fact that $(\phi(z) + 1)^2 = 2(z+1)\phi(z)$ if $|z| > 1$, we get our result. \square

Acknowledgements

The third author (BF) acknowledges the kind reception of the Departamento de Matemáticas of Universidad Carlos III de Madrid during his visit from December 2010 to March 2011. In this period the content of manuscript was discussed in several seminars. The work of the first (FM) and fourth (EH) authors has been supported by Dirección General de Investigación, Ministerio de Ciencia e Innovación of Spain, grant MTM2009-12740-C03-01. The authors thank the valuable comments by the referees which have contributed to improve the presentation of the manuscript.

References

- [Alfaro et al. 1992] M. Alfaro, F. Marcellán, M. L. Rezola, and A. Ronveaux, “On orthogonal polynomials of Sobolev type: algebraic properties and zeros”, *SIAM J. Math. Anal.* **23**:3 (1992), 737–757. MR 93g:42015 Zbl 0764.33003
- [Álvarez-Nodarse and Moreno-Balcázar 2004] R. Álvarez-Nodarse and J. J. Moreno-Balcázar, “Asymptotic properties of generalized Laguerre orthogonal polynomials”, *Indag. Math. (N.S.)* **15**:2 (2004), 151–165. MR 2005e:33003 Zbl 1064.41022
- [Bracciali et al. 2002] C. F. Bracciali, D. K. Dimitrov, and A. Sri Ranga, “Chain sequences and symmetric generalized orthogonal polynomials”, *J. Comput. Appl. Math.* **143**:1 (2002), 95–106. MR 2003f:33007 Zbl 1006.33008
- [Chihara 1978] T. S. Chihara, *An introduction to orthogonal polynomials*, Mathematics and its Applications **13**, Gordon and Breach, New York, 1978. MR 58 #1979 Zbl 0389.33008
- [Dueñas et al. 2011] H. Dueñas, E. J. Huertas, and F. Marcellán, “Analytic properties of Laguerre-type orthogonal polynomials”, *Integral Transforms Spec. Funct.* **22**:2 (2011), 107–122. MR 2011k:33029 Zbl 1213.33017
- [Fejzullahu 2011] B. X. Fejzullahu, “Asymptotics for orthogonal polynomials with respect to the Laguerre measure modified by a rational factor”, *Acta Sci. Math. (Szeged)* **77**:1-2 (2011), 73–85. MR 2841145 Zbl 05990930
- [Fejzullahu and Marcellán 2009] B. X. Fejzullahu and F. Marcellán, “A Cohen type inequality for Laguerre–Sobolev expansions”, *J. Math. Anal. Appl.* **352**:2 (2009), 880–889. MR 2010a:42093 Zbl 1160.42312
- [Fejzullahu and Zejnullahu 2010] B. X. Fejzullahu and R. X. Zejnullahu, “Orthogonal polynomials with respect to the Laguerre measure perturbed by the canonical transformations”, *Integral Transforms Spec. Funct.* **21**:8 (2010), 569–580. MR 2011i:33022 Zbl 1215.33007
- [Koekoek and Meijer 1993] R. Koekoek and H. G. Meijer, “A generalization of Laguerre polynomials”, *SIAM J. Math. Anal.* **24**:3 (1993), 768–782. MR 94b:33007 Zbl 0780.33007
- [López et al. 1995] G. López, F. Marcellán, and W. Van Assche, “Relative asymptotics for polynomials orthogonal with respect to a discrete Sobolev inner product”, *Constr. Approx.* **11**:1 (1995), 107–137. MR 96c:42051 Zbl 0840.42017
- [Marcellán and Moreno-Balcázar 2006] F. Marcellán and J. J. Moreno Balcázar, “Asymptotics and zeros of Sobolev orthogonal polynomials on unbounded supports”, *Acta Appl. Math.* **94**:2 (2006), 163–192. MR 2007i:42002 Zbl 1137.42312

- [Marcellán and Ronveaux 1990] F. Marcellán and A. Ronveaux, “On a class of polynomials orthogonal with respect to a discrete Sobolev inner product”, *Indag. Math. (N.S.)* **1**:4 (1990), 451–464. MR 92f:42029 Zbl 0732.42016
- [Marcellán and Van Assche 1993] F. Marcellán and W. Van Assche, “Relative asymptotics for orthogonal polynomials with a Sobolev inner product”, *J. Approx. Theory* **72**:2 (1993), 193–209. MR 94h:42037 Zbl 0771.42014
- [Meijer 1993a] H. G. Meijer, “Laguerre polynomials generalized to a certain discrete Sobolev inner product space”, *J. Approx. Theory* **73**:1 (1993), 1–16. MR 94d:42027 Zbl 0771.42015
- [Meijer 1993b] H. G. Meijer, “Zero distribution of orthogonal polynomials in a certain discrete Sobolev space”, *J. Math. Anal. Appl.* **172**:2 (1993), 520–532. MR 94a:42027 Zbl 0780.42016
- [Szegő 1975] G. Szegő, *Orthogonal polynomials*, 4th ed., Amer. Math. Soc. Colloq. Publ. **23**, Amer. Math. Soc., Providence, RI, 1975. MR 51 #8724 Zbl 0305.42011

Received June 20, 2011. Revised January 23, 2012.

FRANCISCO MARCELLÁN
DEPARTAMENTO DE MATEMÁTICAS
ESCUELA POLITÉCNICA SUPERIOR
UNIVERSIDAD CARLOS III DE MADRID
AVENIDA DE LA UNIVERSIDAD 30
28911 LEGANÉS
SPAIN
pacomarc@ing.uc3m.es

RAMADAN ZEJNULLAHU
FACULTY OF MATHEMATICS AND SCIENCES
UNIVERSITY OF PRISHTINA
MOTHER TERESA 5
10000 PRISHTINA
KOSOVO
zejnullahu@yahoo.com

BUJAR FEJZULLAHU
FACULTY OF MATHEMATICS AND SCIENCES
UNIVERSITY OF PRISHTINA
MOTHER TERESA 5
1000 PRISHTINA
KOSOVO
bujar.feizullahu@uni-pr.edu

EDMUNDO HUERTAS
DEPARTAMENTO DE MATEMÁTICAS
ESCUELA POLITÉCNICA SUPERIOR
UNIVERSIDAD CARLOS III DE MADRID
AVENIDA DE LA UNIVERSIDAD 30
28911 LEGANÉS
SPAIN
ehuertas@math.uc3m.es

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 257 No. 1 May 2012

Energy and volume of vector fields on spherical domains	1
FABIANO G. B. BRITO, ANDRÉ O. GOMES and GIOVANNI S. NUNES	
Maps on 3-manifolds given by surgery	9
BOLDIZSÁR KALMÁR and ANDRÁS I. STIPSICZ	
Strong solutions to the compressible liquid crystal system	37
YU-MING CHU, XIAN-GAO LIU and XIAO LIU	
Presentations for the higher-dimensional Thompson groups nV	53
JOHANNA HENNIG and FRANCESCO MATUCCI	
Resonant solutions and turning points in an elliptic problem with oscillatory boundary conditions	75
ALFONSO CASTRO and ROSA PARDO	
Relative measure homology and continuous bounded cohomology of topological pairs	91
ROBERTO FRIGERIO and CRISTINA PAGLIANTINI	
Normal enveloping algebras	131
ALEXANDRE N. GRISHKOV, MARINA RASSKAZOVA and SALVATORE SICILIANO	
Bounded and unbounded capillary surfaces in a cusp domain	143
YASUNORI AOKI and DAVID SIEGEL	
On orthogonal polynomials with respect to certain discrete Sobolev inner product	167
FRANCISCO MARCELLÁN, RAMADAN ZEJNULLAHU, BUJAR FEJZULLAHU and EDMUNDO HUERTAS	
Green versus Lempert functions: A minimal example	189
PASCAL THOMAS	
Differential Harnack inequalities for nonlinear heat equations with potentials under the Ricci flow	199
JIA-YONG WU	
On overtwisted, right-veering open books	219
PAOLO LISCA	
Weakly Krull domains and the composite numerical semigroup ring $D + E[\Gamma^*]$	227
JUNG WOOK LIM	
Arithmeticity of complex hyperbolic triangle groups	243
MATTHEW STOVER	



0030-8730(201205)257:1;1-7