

*Pacific
Journal of
Mathematics*

**GREEN VERSUS LEMPERT FUNCTIONS:
A MINIMAL EXAMPLE**

PASCAL THOMAS

GREEN VERSUS LEMPERT FUNCTIONS: A MINIMAL EXAMPLE

PASCAL THOMAS

The Lempert function for a set of poles in a domain of \mathbb{C}^n at a point z is obtained by taking a certain infimum over all analytic disks going through the poles and the point z ; it majorizes the corresponding multipole pluricomplex Green function. Coman proved that both coincide in the case of sets of two poles in the unit ball. We give an example of a set of three poles in the unit ball where this equality fails.

1. Introduction

Let Ω be a domain in \mathbb{C}^n , and $a_j \in \Omega$, $j = 1, \dots, N$. The pluricomplex Green function with logarithmic singularities at $S := \{a_1, \dots, a_N\}$ is defined by

$$G_S(z) := \sup\{u \in \text{PSH}(\Omega, \mathbb{R}_-) : u(z) \leq \log|z - a_j| + C_j, j = 1, \dots, N\},$$

where $\text{PSH}(\Omega, \mathbb{R}_-)$ stands for the set of all negative plurisubharmonic functions in Ω . When Ω is hyperconvex, this solves the Monge–Ampère equation with right hand side equal to $\sum_{i=1}^N \delta_{a_j}$.

Pluricomplex Green functions have been studied by many authors at different levels of generality. See [Demailly 1987; Zahariuta 1984; Lempert 1981; Lelong 1989; Lárusson and Sigurdsson 1998].

A deep result due to Poletsky [1993], and see also [Lárusson and Sigurdsson 1998; Edigarian 1997], is that the Green function may be computed from analytic disks:

$$(1-1) \quad G_S(z) = \inf \left\{ \sum_{\alpha: \varphi(\alpha) \in S} \log|\alpha| : \text{such that there exists } \varphi \in \mathcal{O}(\mathbb{D}, \Omega) \text{ with } \varphi(0) = z \right\}.$$

MSC2010: 32U35, 32F45.

Keywords: pluricomplex Green function, Lempert function, analytic disks, Schwarz Lemma.

However, it is tempting to pick only one $\alpha_j \in \varphi^{-1}(a_j)$ in the range $1 \leq j \leq N$, which motivated the Coman's definition of the Lempert function [2000]:

$$(1-2) \quad \ell_S(z) := \inf \left\{ \sum_{j=1}^N \log |\zeta_j| : \varphi(0) = z, \varphi(\zeta_j) = a_j, j = 1, \dots, N \right. \\ \left. \text{for some } \varphi \in \mathbb{C}(\mathbb{D}, \Omega) \right\},$$

where \mathbb{D} is the unit disc in \mathbb{C} .

One easily sees that $\ell_S(z) \geq G_S(z)$ without recourse to (1-1); the fact that equality holds when $N = 1$ and Ω is convex is part of Lempert's celebrated theorem [1981], which was, in fact, the starting point for many of the notions defined above; see also [Edigarian 1995]. Coman [2000] proved that equality holds when $N = 2$ and $\Omega = \mathbb{B}^2$, the unit ball of \mathbb{C}^2 . The goal of this note is to present an example that shows that this is as far as it can go.

Theorem 1.1. *There exists a set of 3 points $S \subset \mathbb{B}^2$ such that $\ell_S(z) > G_S(z)$ for some $z \in \mathbb{B}^2$.*

Other examples in the same vein have been found in [Carlehed and Wiegerinck 2003; Thomas and Trao 2003; Nikolov and Zwonek 2005]. The interesting features of this one are that it involves no multiplicities and is minimal in the ball. Examples with an arbitrary number of points can be deduced from it. Let $z_0 \in \mathbb{B}^2$ satisfy $\ell_S(z_0) - G_S(z_0) =: \varepsilon_0 > 0$. Consider $S' := S \cup \{a_1, \dots, a_N\}$ with all the a_j close enough to the boundary so that $\ell_{S'}(z_0) \geq \ell_S(z_0) - \varepsilon_0/2$ (the Schwarz lemma shows that $|\zeta_j| \rightarrow 1$ when $\varphi(\zeta_j) = a_j$ and $|a_j| \rightarrow 1$). Then $\ell_{S'}(z_0) > G_S(z_0) \geq G_{S'}(z_0)$, as was to be shown. (I thank Nikolai Nikolov for sharing this observation with me).

Moreover, the corresponding Green function can be recovered, up to a bounded error, by using an analytic disk with just one more preimage than the number of points: One of the points has exactly two preimages and each of the other two points, only one; see [Magnússon et al. 2012, §6.8.2, Lemma 6.16].

More specifically, the theorem will follow from a precise calculation in the bidisk \mathbb{D}^2 . Let $S_\varepsilon = \{(0, 0), (\rho(\varepsilon), 0), (0, \varepsilon)\} \subset \mathbb{D}^2$, where $\lim_{\varepsilon \rightarrow 0} \rho(\varepsilon)/\varepsilon = 0$.

Proposition 1.2. *There exists $C_1 > 0$ such that for any $\delta \in (0, 1/4)$ there exists $\varepsilon_0 = \varepsilon_0(\delta) > 0$ and $r_0 = r_0(\delta) > 0$ such that*

$$(1-3) \quad G_{S_\varepsilon}(z) \leq 2 \log |z_2| + C_1,$$

$$(1-4) \quad \ell_{S_\varepsilon}(z) \geq (2 - \delta) \log |z_2|.$$

for any ε with $|\varepsilon| < \varepsilon_0$ and any $z = (z_1, z_2) \in \mathbb{D}^2$ such that

$$(1-5) \quad \frac{1}{2} |z_2|^{3/2} \leq |z_1| \leq |z_2|^{3/2} \quad \text{and} \quad \|z\| < r_0.$$

Proof of Theorem 1.1. If U and V are domains, and $S \subset U \subset V$, then the definitions of the Green and Lempert functions imply that $G_S^U(z) \geq G_S^V(z)$ and $\ell_S^U(z) \geq \ell_S^V(z)$.

For $|\varepsilon|$ small enough, $S_\varepsilon \subset \mathbb{B}^2$. When $|z_1| = |z_2|^{3/2}$, so that z verifies (1-5), the inclusion $\mathbb{B}^2 \subset \mathbb{D}^2$ implies

$$\ell_{S_\varepsilon}^{\mathbb{B}^2}(z) \geq \ell_{S_\varepsilon}^{\mathbb{D}^2}(z) \geq (2 - \delta) \log |z_2|.$$

Using the fact that $\mathbb{D}^2/\sqrt{2} \subset \mathbb{B}^2$ and the invariance of the Green function under biholomorphic mappings, we have

$$G_{S_\varepsilon}^{\mathbb{B}^2}(z) \leq G_{S_\varepsilon/\sqrt{2}}^{\mathbb{D}^2}(z) = G_{\sqrt{2}S_\varepsilon}^{\mathbb{D}^2}(\sqrt{2}z) \leq 2 \log |z_2| + \log 2 + C_1.$$

The last inequality follows from the fact that $\sqrt{2}z$ still verifies (1-5), and $\sqrt{2}S_\varepsilon$ has the same form as S_ε , so we can apply (1-3).

Comparing the last two estimates, we see that $G_{S_\varepsilon}^{\mathbb{B}^2}(z) < \ell_{S_\varepsilon}^{\mathbb{B}^2}(z)$ for $|z_2|$ small enough and $|\varepsilon| < \varepsilon_0$. □

Open questions

This example is minimal in the ball, in terms of number of poles; what is the situation for the bidisk? Are the Green and Lempert functions equal when one takes two poles, not lying on a line parallel to the coordinate axes? Do they at least have the same order of singularity as one pole tends to the other?

What is the precise order of the singularity of the limit as $\varepsilon \rightarrow 0$ of the Lempert function in this case? Looking at the available analytic disks that give the correct order of the singularity of the limit of the Green function, one finds $\frac{3}{2} \log |z_2|$, so one would hope that the proposition can still be proved at least for $\delta < 1/2$.

Do the analytic disks from [Magnússon et al. 2012] yield the Green function itself, without any bounded error term?

More generally, when one is given a finite number of points in a given bounded (hyperconvex) domain, is there a bound on the number of preimages required to attain the Green function in the Poletsky formula? For instance, is 4 the largest possible number of preimages required when looking at 3 points in the ball?

2. Upper estimate for the Green function

Proof of (1-3) of Proposition 1.2. The upper bound (1-3) follows from [Magnússon et al. 2012, §6.8.2, Lemma 6.16]. For the reader’s convenience, and since that paper is not generally available, we repeat the proof here in the case that concerns us.

We now construct an analytic disk passing twice through one of the poles. Our disk will be a perturbation of the Neil parabola $\zeta \mapsto (\zeta^3, \zeta^2)$.

We write $s(\varepsilon) = \rho(\varepsilon)/\varepsilon = o(1)$.

Choose complex numbers λ and μ such that

$$\lambda^2 := \frac{z_1}{z_2(z_2 - \varepsilon)} \left(\frac{z_1}{z_2 - \varepsilon} + s(\varepsilon) \right) \quad \text{and} \quad \mu^2 := \varepsilon + \left(\frac{s(\varepsilon)}{2\lambda} \right)^2.$$

Let

$$\Psi_{\lambda,\mu}(\zeta) := \left((\lambda\zeta - \frac{1}{2}s(\varepsilon))(\zeta^2 - \mu^2), \zeta^2 - \left(\frac{s(\varepsilon)}{2\lambda} \right)^2 \right).$$

Then by construction $\Psi_{\lambda,\mu}(\mu) = \Psi_{\lambda,\mu}(-\mu) = (0, \varepsilon)$,

$$\Psi_{\lambda,\mu} \left(\frac{s(\varepsilon)}{2\lambda} \right) = (0, 0) \quad \text{and} \quad \Psi_{\lambda,\mu} \left(-\frac{s(\varepsilon)}{2\lambda} \right) = (\varepsilon s(\varepsilon), 0),$$

so we have a disk passing through all three poles of G_ε . Furthermore, choosing

$$\zeta_z := \frac{1}{\lambda} \left(\frac{z_1}{z_2 - \varepsilon} + \frac{s(\varepsilon)}{2} \right),$$

we have $\Psi_{\lambda,\mu}(\zeta_z) = z$. Notice that

$$\zeta_z^2 = \frac{z_2(z_2 - \varepsilon)}{z_1} \left(\frac{z_1}{z_2 - \varepsilon} + \frac{s(\varepsilon)}{2} \right)^2 \left(\frac{z_1}{z_2 - \varepsilon} + s(\varepsilon) \right)^{-1},$$

so for any $\eta > 0$ there exists $\varepsilon_0(\delta, \eta) > 0$ such that for $|\varepsilon| < \varepsilon_0(\delta, \eta)$

$$(2-1) \quad \left| |\zeta_z| - |z_2|^{1/2} \right| \leq \eta$$

for any z such that $\delta \leq \frac{1}{2}|z_2|^{3/2} \leq |z_1| \leq |z_2|^{3/2} \leq 1$. In particular, by choosing η small enough we ensure that $\zeta_z \in \mathbb{D}$. We need a more general fact.

Claim. *Let $\eta > 0$, and $\delta > 0$. Then there exists $\varepsilon_1 = \varepsilon_1(\delta, \eta) > 0$ such that for any ε with $|\varepsilon| \leq \varepsilon_1$, we have $\Psi_{\lambda,\mu}(D(0, 1 - \eta)) \subset \mathbb{D}^2$ for any z such that $\delta \leq \frac{1}{2}|z_2|^{3/2} \leq |z_1| \leq |z_2|^{3/2} \leq 1$.*

Proof. For $|\varepsilon| \leq \delta^{2/3}/2$, we have $|z_2|/2 \leq |z_2 - \varepsilon| \leq 2|z_2|$, so

$$|\lambda|^2 \geq \left| \frac{z_1}{2z_2^2} \right| \left(\left| \frac{z_1}{2z_2} \right| - |s(\varepsilon)| \right) \geq \left| \frac{z_1^2}{8z_2^3} \right| \geq \frac{1}{32}$$

for ε small enough. So when $|\zeta| \leq 1 - \eta$,

$$|\Psi_{\lambda,\mu,2}(\zeta)| \leq (1 - \eta)^2 + 256|s(\varepsilon)|^2 < 1$$

for ε small enough.

In a similar way, given η' , for ε small enough depending on δ and η' , we have $|z_2| \leq (1 + \eta')|z_2 - \varepsilon|$, so

$$|\lambda|^2 \leq (1 + \eta')^2 \left| \frac{z_1}{z_2^2} \right| \left(\left| \frac{z_1}{z_2} \right| + \frac{|s(\varepsilon)|}{(1 + \eta')} \right) \leq (1 + \eta')^3 \left| \frac{z_1^2}{z_2^3} \right| \leq (1 + \eta')^3$$

for ε small enough. Choose η' so that $(1 + \eta')^3 = (1 + \eta)$. When $|\zeta| \leq 1 - \eta$,

$$|\Psi_{\lambda,\mu,1}(\zeta)| \leq \left((1 + \eta)(1 - \eta) + \frac{1}{2}|s(\varepsilon)| \right) \left((1 - \eta)^2 + |\varepsilon| + 64^2|s(\varepsilon)|^2 \right) < 1$$

for ε small enough. □

So now the function $v(\zeta) := G_\varepsilon(\Psi_{\lambda,\mu}((1 - \eta)\zeta))$ is negative and subharmonic on \mathbb{D} . Furthermore, it has logarithmic poles at the points

$$\pm \frac{\mu}{1 - \eta} \quad \text{and} \quad \pm \frac{s(\varepsilon)}{2\lambda(1 - \eta)};$$

in the cases when $\mu = 0$ or $s(\varepsilon) = 0$, we get a double logarithmic pole at the corresponding point.

Denote by $d_G(\zeta, \xi) := |(\zeta - \xi)/(1 - \zeta\bar{\xi})|$ the invariant (pseudohyperbolic) distance between points of the unit disk. Then

$$\begin{aligned} G_\varepsilon(z) = v(\zeta_z) &\leq \log d_G\left(\zeta_z, \frac{\mu}{1 - \eta}\right) + \log d_G\left(\zeta_z, -\frac{\mu}{1 - \eta}\right) \\ &\quad + \log d_G\left(\zeta_z, \frac{s(\varepsilon)}{2\lambda(1 - \eta)}\right) + \log d_G\left(\zeta_z, -\frac{s(\varepsilon)}{2\lambda(1 - \eta)}\right). \end{aligned}$$

By (2-1), choosing $m(\delta, \eta)$ accordingly, we have $G_\varepsilon(z) \leq 4 \log|z_2|^{1/2} + O(\eta)$ for $|\varepsilon| \leq m$. □

3. Lower estimate for the Lempert function

Proof of (1-4) of Proposition 1.2. The proof will follow the methods and notations of [Thomas 2007]. We will make repeated use of the involutive automorphisms of the unit disk given by $\phi_a(\zeta) := (a - \zeta)/(1 - \bar{a}\zeta)$ for $a \in \mathbb{D}$, which exchange 0 and a . Notice that the invariant (pseudohyperbolic) distance verifies

$$d_G(a, b) := |\phi_a(b)| = |\phi_b(a)|.$$

Write $\rho(\varepsilon) = \varepsilon s(\varepsilon)$ with $\lim_{\varepsilon \rightarrow 0} s(\varepsilon) = 0$.

We will assume that the conclusion fails. That is, for any $\delta \in (0, 1/4)$, there exist arbitrarily small values of $|z_2| = \max(|z_1|, |z_2|)$, and $|\varepsilon|$ such that

$$(3-1) \quad \ell_{S_\varepsilon}(z) < (2 - \delta) \log|z_2|.$$

After applying, for each analytic disk, an automorphism of the disk that exchanges the preimage of $(0, 0)$ and 0, the assumption implies that there exists a holomorphic map φ from \mathbb{D} to \mathbb{D}^2 and points $\zeta_j \in \mathbb{D}$, depending on z and ε , satisfying the conditions

$$(3-2) \quad \begin{aligned} \varphi(0) &= (0, 0), & \varphi(\zeta_1) &= (\varepsilon s(\varepsilon), 0), \\ \varphi(\zeta_0) &= (z_1, z_2), & \varphi(\zeta_2) &= (0, \varepsilon), \end{aligned}$$

with

$$(3-3) \quad \log|\zeta_0| + \log|\phi_{\zeta_0}(\zeta_1)| + \log|\phi_{\zeta_0}(\zeta_2)| \leq (2 - \delta) \log|z_2|.$$

The interpolation conditions in (3-2) are equivalent to the existence of holomorphic functions h_1 and h_2 from \mathbb{D} to itself such that

$$\varphi(\zeta) = (\zeta \phi_{\zeta_2}(\zeta) h_1(\zeta), \zeta \phi_{\zeta_1}(\zeta) h_2(\zeta)),$$

such that furthermore

$$(3-4) \quad h_1(\zeta_1) = \frac{\varepsilon s(\varepsilon)}{\zeta_1 \phi_{\zeta_2}(\zeta_1)} =: w_1,$$

$$(3-5) \quad h_1(\zeta_0) = \frac{z_1}{\zeta_0 \phi_{\zeta_2}(\zeta_0)} =: w_2,$$

$$(3-6) \quad h_2(\zeta_2) = \frac{\varepsilon}{\zeta_2 \phi_{\zeta_1}(\zeta_2)} =: w_4,$$

$$(3-7) \quad h_2(\zeta_0) = \frac{z_2}{\zeta_0 \phi_{\zeta_1}(\zeta_0)} =: w_3.$$

By the invariant Schwarz lemma, the existence of a holomorphic function h_1 mapping \mathbb{D} to itself and satisfying (3-4) and (3-5) is equivalent to

$$(3-8) \quad |w_1| < 1, \quad |w_2| < 1 \quad \text{and} \quad d_G(w_1, w_2) < d_G(\zeta_1, \zeta_0) = |\phi_{\zeta_1}(\zeta_0)|.$$

In the same way, the existence of h_2 is equivalent to

$$(3-9) \quad |w_3| < 1, \quad |w_4| < 1 \quad \text{and} \quad d_G(w_3, w_4) < d_G(\zeta_2, \zeta_0) = |\phi_{\zeta_2}(\zeta_0)|.$$

As in [Thomas 2007], we start by remarking that (3-3) can be rewritten as

$$(3-10) \quad -\log|w_2| - \log|w_3| = \log \left| \frac{\zeta_0 \phi_{\zeta_1}(\zeta_0)}{z_2} \right| + \log \left| \frac{\zeta_0 \phi_{\zeta_0}(\zeta_2)}{z_1} \right| \\ \leq \log|\zeta_0| + (2 - \delta) \log|z_2| - \log|z_1| - \log|z_2| \\ \leq \log|\zeta_0| - \left(\frac{1}{2} + \delta\right) \log|z_2| + \log 2,$$

by (1-5). We can rewrite this in a more symmetric fashion:

$$(3-11) \quad \log \frac{1}{|w_2|} + \log \frac{1}{|w_3|} + \log \frac{1}{|\zeta_0|} \leq \left(\frac{1}{2} + \delta\right) \log \frac{1}{|z_2|} + \log 2.$$

Since all terms are positive by (3-8) and (3-9), each of the terms on the left hand side is bounded by the right hand side.

We will proceed as follows: We have used the contradiction hypothesis (3-3) to prove that $|\zeta_0|$ and $|w_3|$ are relatively big. We will prove that $|\phi_{\zeta_2}(\zeta_0)|$ has to be relatively small, which by (3-9) forces $|w_4|$ to be roughly as large as $|w_3|$. This then allows us to bound $|\phi_{\zeta_1}(\zeta_2)|$ by a quantity that becomes as small as desired

when ε can be made small, and hence allows us to bound $|\phi_{\zeta_1}(\zeta_0)|$ by the triangle inequality.

The final contradiction will concern $w_2 = z_1/(\zeta_0\phi_{\zeta_2}(\zeta_0))$. On the one hand, (3-11) guarantees that it is not too small; but an explicit computation of the quotient w_1/w_4 shows that w_1 must be small, and by (3-8) and the estimate on $|\phi_{\zeta_1}(\zeta_0)|$, $|w_2|$ is small as well.

We provide the details. From (3-11),

$$(3-12) \quad \log|w_3| \geq \left(\frac{1}{2} + \delta\right) \log|z_2| - \log 2.$$

From (3-5) and (3-10),

$$(3-13) \quad \begin{aligned} \log|\phi_{\zeta_2}(\zeta_0)| &= \log|z_1/\zeta_0| - \log|w_2| \\ &\leq \log|z_1/\zeta_0| + \log|\zeta_0| - \left(\frac{1}{2} + \delta\right) \log|z_2| + \log 2 \\ &\leq (1 - \delta) \log|z_2| + \log 2. \end{aligned}$$

Since $\delta < 1/4$, (3-13) and (3-12) imply that $|\phi_{\zeta_2}(\zeta_0)| < \frac{1}{2}|w_3|$ for $|z_2| \leq r_1(\delta)$, so by (3-9) and the triangle inequality for d_G ,

$$(3-14) \quad |w_4| \geq \frac{1}{2}|w_3|.$$

We now prove that both ζ_1 and ζ_2 must be close to ζ_0 and even closer to each other. First, since (3-11) implies that $\log|\zeta_0| \geq \left(\frac{1}{2} + \delta\right) \log|z_2| - \log 2$, by (3-13), $|\phi_{\zeta_2}(\zeta_0)| \leq \frac{1}{2}|\zeta_0|$ for $|z_2| \leq r_2(\delta)$. By the triangle inequality for d_G ,

$$(3-15) \quad \frac{1}{2}|\zeta_0| \leq |\zeta_2| \leq \frac{3}{2}|\zeta_0|.$$

On the other hand, from (3-11),

$$\log|w_3| + \log|\zeta_0| \geq \left(\frac{1}{2} + \delta\right) \log|z_2| - \log 2, \quad \text{that is, } |w_3\zeta_0| \geq \frac{1}{2}|z_2|^{\delta+1/2}.$$

Therefore, applying (3-14) and (3-15),

$$(3-16) \quad |\phi_{\zeta_1}(\zeta_2)| = \left| \frac{\varepsilon}{\zeta_2 w_4} \right| \leq 4 \left| \frac{\varepsilon}{\zeta_0 w_3} \right| \leq 8|\varepsilon||z_2|^{-\delta-1/2}.$$

In particular, for

$$(3-17) \quad |\varepsilon| < \frac{1}{8}|z_2|^{3/2},$$

this implies $|\phi_{\zeta_1}(\zeta_2)| < |z_2|^{1-\delta}$, and by the triangle inequality,

$$(3-18) \quad |\phi_{\zeta_1}(\zeta_0)| < |\phi_{\zeta_2}(\zeta_0)| + |\phi_{\zeta_1}(\zeta_2)| < 3|z_2|^{1-\delta}.$$

We now establish the two (contradictory) estimates for w_2 . On the one hand, (3-11) implies that

$$(3-19) \quad \log|w_2| \geq \left(\frac{1}{2} + \delta\right) \log|z_2| - \log 2, \quad \text{that is, } |w_2| \geq \frac{1}{2}|z_2|^{\delta+1/2}.$$

On the other hand,

$$\left| \frac{w_1}{w_4} \right| = \left| \frac{\varepsilon s(\varepsilon)}{\zeta_1 \phi_{\zeta_2}(\zeta_1)} \frac{\zeta_2 \phi_{\zeta_1}(\zeta_2)}{\varepsilon} \right| = \left| s(\varepsilon) \frac{\zeta_2}{\zeta_1} \right|.$$

By the triangle inequality for d_G , when (3-17) holds, the lower bound in (3-15) and the corollary to (3-16) imply

$$|\zeta_1| \geq |\zeta_2| - |\phi_{\zeta_1}(\zeta_2)| \geq \frac{1}{2}|\zeta_0| - |z_2|^{1-\delta} \geq \frac{1}{4}|\zeta_0|$$

for $|z_2|$ small enough, because of (3-11) again. So finally, using the upper bound in (3-15), $|w_1/w_4| \leq 6|s(\varepsilon)|$. We choose $\varepsilon_0 < \frac{1}{8}|z_2|^{3/2}$ so that for any ε with $|\varepsilon| \leq \varepsilon_0$,

$$(3-20) \quad |s(\varepsilon)| < |z_2|^{1-\delta}.$$

The triangle inequality for d_G and (3-18) imply that when $|\varepsilon| \leq \varepsilon_0$,

$$|w_2| \leq |w_1| + |\phi_{\zeta_1}(\zeta_0)| \leq 6|s(\varepsilon)| + 3|z_2|^{1-\delta} \leq 9|z_2|^{1-\delta}.$$

Finally, if we choose $|z_2| \leq r_0(\delta)$, with

$$r_0(\delta) \leq \min(r_1(\delta), r_2(\delta)) \quad \text{and} \quad 9r_0(\delta)^{1-\delta} < \frac{1}{2}r_0(\delta)^{1/2+\delta},$$

we see that for any ε with $|\varepsilon| \leq \varepsilon_0$, this last bound contradicts (3-19). □

Acknowledgments

Part of this work was done when I was a guest of the Hanoi University of Education in March 2011. I wish to thank my colleagues there for their hospitality, in particular Do Duc Thai and Nguyen Van Trao. A detailed exposition of the results in a workshop organized in Bedlewo by colleagues from the Jagiellonian University in Cracow resulted in some streamlining of the proof. Finally, the Semester in Complex Analysis and Spectral Theory of the Centre de Recerca Matemàtica at the Universitat Autònoma de Barcelona provided the occasion to mention the result in a talk, and to put the finishing touches on the submitted manuscript.

I thank Nguyen Van Trao for useful discussions on this topic, and the referee for pointing out and correcting a mistake in the original exposition.

References

- [Carlehed and Wiegerinck 2003] M. Carlehed and J. Wiegerinck, “Le cône des fonctions plurisous-harmoniques négatives et une conjecture de Coman”, *Ann. Polon. Math.* **80** (2003), 93–108. MR 2004e:32034 Zbl 1026.32066
- [Coman 2000] D. Coman, “The pluricomplex Green function with two poles of the unit ball of \mathbb{C}^n ”, *Pacific J. Math.* **194**:2 (2000), 257–283. MR 2001g:32081 Zbl 1015.32029

- [Demailly 1987] J.-P. Demailly, “Mesures de Monge–Ampère et mesures pluriharmoniques”, *Math. Z.* **194**:4 (1987), 519–564. [MR 88g:32034](#) [Zbl 0595.32006](#)
- [Edigarian 1995] A. Edigarian, “A remark on the Lempert theorem”, *Univ. Iagel. Acta Math.* **32** (1995), 83–88. [MR 96h:32034](#) [Zbl 0833.32005](#)
- [Edigarian 1997] A. Edigarian, “On definitions of the pluricomplex Green function”, *Ann. Polon. Math.* **67**:3 (1997), 233–246. [MR 99f:32039](#) [Zbl 0909.31007](#)
- [Lárússon and Sigurdsson 1998] F. Lárússon and R. Sigurdsson, “Plurisubharmonic functions and analytic discs on manifolds”, *J. Reine Angew. Math.* **501** (1998), 1–39. [MR 99e:32020](#) [Zbl 0901.31004](#)
- [Lelong 1989] P. Lelong, “Fonction de Green pluricomplexe et lemmes de Schwarz dans les espaces de Banach”, *J. Math. Pures Appl.* (9) **68**:3 (1989), 319–347. [MR 91c:46065](#) [Zbl 0633.32019](#)
- [Lempert 1981] L. Lempert, “La métrique de Kobayashi et la représentation des domaines sur la boule”, *Bull. Soc. Math. France* **109** (1981), 427–474. [MR 84d:32036](#) [Zbl 0492.32025](#)
- [Magnússon et al. 2012] J. I. Magnússon, A. Rashkovskii, R. Sigurdsson, and P. J. Thomas, “Limits of multipole pluricomplex Green functions”, *Internat. J. Math* **23**:6 (2012), #1250065. [arXiv 1103.2296](#)
- [Nikolov and Zwonek 2005] N. Nikolov and W. Zwonek, “On the product property for the Lempert function”, *Complex Var. Theory Appl.* **50**:12 (2005), 939–952. [MR 2006g:32017](#) [Zbl 1085.32006](#)
- [Poletsky 1993] E. A. Poletsky, “Holomorphic currents”, *Indiana Univ. Math. J.* **42**:1 (1993), 85–144. [MR 94c:32007](#) [Zbl 0811.32010](#)
- [Thomas 2007] P. J. Thomas, “An example of limit of Lempert functions”, *Vietnam J. Math.* **35**:3 (2007), 317–330. [MR 2009h:32017](#) [Zbl 1158.32015](#) [arXiv math/0601642](#)
- [Thomas and Trao 2003] P. J. Thomas and N. V. Trao, “Pluricomplex Green and Lempert functions for equally weighted poles”, *Ark. Mat.* **41**:2 (2003), 381–400. [MR 2004m:32067](#) [Zbl 1038.32028](#)
- [Zahariuta 1984] V. P. Zahariuta, *Spaces of analytic functions and maximal plurisubharmonic functions*, thesis, Rostov State University, Rostov-on-Don, 1984.

Received June 20, 2011. Revised October 18, 2011.

PASCAL THOMAS
INSTITUT DE MATHÉMATIQUES DE TOULOUSE
UNIVERSITÉ DE TOULOUSE
UPS, INSA, UT1, UTM
31062 TOULOUSE
FRANCE

pthomas@math.univ-toulouse.fr
<http://www.math.univ-toulouse.fr/~thomas>

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 257 No. 1 May 2012

Energy and volume of vector fields on spherical domains	1
FABIANO G. B. BRITO, ANDRÉ O. GOMES and GIOVANNI S. NUNES	
Maps on 3-manifolds given by surgery	9
BOLDIZSÁR KALMÁR and ANDRÁS I. STIPSICZ	
Strong solutions to the compressible liquid crystal system	37
YU-MING CHU, XIAN-GAO LIU and XIAO LIU	
Presentations for the higher-dimensional Thompson groups nV	53
JOHANNA HENNIG and FRANCESCO MATUCCI	
Resonant solutions and turning points in an elliptic problem with oscillatory boundary conditions	75
ALFONSO CASTRO and ROSA PARDO	
Relative measure homology and continuous bounded cohomology of topological pairs	91
ROBERTO FRIGERIO and CRISTINA PAGLIANTINI	
Normal enveloping algebras	131
ALEXANDRE N. GRISHKOV, MARINA RASSKAZOVA and SALVATORE SICILIANO	
Bounded and unbounded capillary surfaces in a cusp domain	143
YASUNORI AOKI and DAVID SIEGEL	
On orthogonal polynomials with respect to certain discrete Sobolev inner product	167
FRANCISCO MARCELLÁN, RAMADAN ZEJNULLAHU, BUJAR FEJZULLAHU and EDMUNDO HUERTAS	
Green versus Lempert functions: A minimal example	189
PASCAL THOMAS	
Differential Harnack inequalities for nonlinear heat equations with potentials under the Ricci flow	199
JIA-YONG WU	
On overtwisted, right-veering open books	219
PAOLO LISCA	
Weakly Krull domains and the composite numerical semigroup ring $D + E[\Gamma^*]$	227
JUNG WOOK LIM	
Arithmeticity of complex hyperbolic triangle groups	243
MATTHEW STOVER	