

*Pacific
Journal of
Mathematics*

**ODD HAMILTONIAN SUPERALGEBRAS
AND SPECIAL ODD HAMILTONIAN SUPERALGEBRAS
OF FORMAL VECTOR FIELDS**

LI REN, QIANG MU AND YONGZHENG ZHANG

ODD HAMILTONIAN SUPERALGEBRAS AND SPECIAL ODD HAMILTONIAN SUPERALGEBRAS OF FORMAL VECTOR FIELDS

LI REN, QIANG MU AND YONGZHENG ZHANG

The natural filtrations of odd Hamiltonian superalgebras and special odd Hamiltonian superalgebras of formal vector fields are proved to be invariant under their automorphism group respectively, by determining the set of ad-quasi-nilpotent elements. Thereby, the automorphism groups of these Lie superalgebras are determined.

1. Introduction

As is well known, filtration structures and automorphism groups play an important role in the classification of modular Lie algebras (see [Jin 1992; Strade and Farnsteiner 1988; Wilson 1975]) and nonmodular Lie superalgebras (see [Kac 1977; Kac 1998; Scheunert 1979]), respectively. Cartan-type Lie algebras and Lie superalgebras possess natural filtration structures. The natural filtrations of the infinite-dimensional Lie algebras $L(m)$ and $\hat{L}(m)$ were proved to be invariant in [Rudakov 1986], where $L = W, S, H$ or K . The natural filtrations of the general Lie superalgebra and special Lie superalgebra of formal vector fields were proved to be invariant in [Zhang and Liu 2004]. The invariance of natural filtrations of Cartan-type Lie algebras or Lie superalgebras provides a useful method of determining intrinsic properties and automorphism groups (see [Wilson 1971; Zhang and Liu 2004]).

We consider the infinite-dimensional odd Hamiltonian superalgebra HO and special odd Hamiltonian superalgebra SHO of formal vector fields, which are involved in [Kac 1998]. The corresponding results of Cartan-type Lie algebras are generalized and Jin's methods are used (see [Jin 1992]). Denote by $\{X_i\}_{i \geq -1}$

Qiang Mu is the corresponding author.

Supported by the Tianyuan Fund for Mathematics (grant number 11126051), the Heilongjiang Provincial Natural Science Foundation of China (grant number A201008), the Scientific Research Fund of Heilongjiang Provincial Education Department (grant number 12511157), and the Doctoral Research Foundation of Harbin Normal University (grant number KJB201105).

MSC2010: 17B40, 17B65, 17B66.

Keywords: superalgebra of formal vector fields, ad-nilpotent element, filtration, continuous automorphism.

the natural filtration of X . By determining the ad-quasi-nilpotent elements in the even part and the subalgebras generated by certain ad-quasi-nilpotent elements, we prove that the natural filtration of X is invariant under automorphisms in the following sense: If φ is an automorphism of X , then $\varphi(X_i) \subseteq X_i$ for every $i \geq -1$. Besides, we prove that every automorphism of X is continuous and can be induced from an automorphism of $\Lambda(n, n)$. Finally, we prove that the automorphism group of X is isomorphic to the admissible automorphism group of the base superalgebra $\Lambda(n, n)$.

This paper is arranged as follows. In Section 2, we recall the necessary definitions concerning Lie superalgebras of Cartan type HO and SHO of formal vector fields. In Section 3, we characterize the ad-quasi-nilpotent elements with certain properties, and prove the invariance of their natural filtrations. In Section 4, we determine the automorphism groups of Lie superalgebras of Cartan type HO and SHO of formal vector fields.

2. Preliminaries

In this paper, \mathbb{F} denotes an algebraically closed field of characteristic zero, and n is a positive integer greater than 3. Let \mathbb{N} and \mathbb{N}_0 denote the sets of positive integers and nonnegative integers, respectively. Let $\mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ denote the ring of integers modulo 2. Let $P(n) = \mathbb{F}[[x_1, \dots, x_n]]$ denote the ring of formal power series in n variables over field \mathbb{F} . For $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}_0^n$, we abbreviate $x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}$ to $x^{(\alpha)}$, and put $|\alpha| = \sum_{i=1}^n \alpha_i$. Let $\Lambda(n)$ be the Grassmann algebra over \mathbb{F} in n variables $x_{n+1}, x_{n+2}, \dots, x_{2n}$. Denote by $\Lambda(n, n)$ the tensor product $P(n) \otimes_{\mathbb{F}} \Lambda(n)$. Then $\Lambda(n, n)$ is a noncommutative linearly compact topological superalgebra with a fundamental system $\{(\Lambda_1)^j\}_{j \geq 1}$ of neighborhoods of 0, where $(\Lambda_1)^j = \text{span}_{\mathbb{F}}\{x_{i_1} \cdots x_{i_k} \mid j \leq k\}$. In particular, $(\Lambda_1)^1$ is the ideal of $\Lambda(n, n)$ generated by $\{x_1, \dots, x_{2n}\}$ (see [Kac 1998]). For $g \in P(n)$ and $f \in \Lambda(n)$, we abbreviate $g \otimes f$ to gf .

Put $Y_0 = \{1, 2, \dots, n\}$, $Y_1 = \{n + 1, \dots, 2n\}$ and $Y = Y_0 \cup Y_1$. Let

$$\mathbb{B}_k = \{(i_1, i_2, \dots, i_k) \mid n + 1 \leq i_1 < i_2 < \cdots < i_k \leq 2n\}$$

and $\mathbb{B}(n) = \bigcup_{k=0}^n \mathbb{B}_k$, where $\mathbb{B}_0 = \emptyset$. Given $u = (i_1, i_2, \dots, i_k) \in \mathbb{B}_k$, set $|u| = k$, $\{u\} = \{i_1, i_2, \dots, i_k\}$ and $x^u = x_{i_1} x_{i_2} \cdots x_{i_k}$ (with the convention that $|\emptyset| = 0$ and $x^\emptyset = 1$). Then $\{x^{(\alpha)} x^u \mid \alpha \in \mathbb{N}_0^n, u \in \mathbb{B}(n)\}$ is an \mathbb{F} -basis of the infinite-dimensional superalgebra $\Lambda(n, n)$. Clearly, $\Lambda(n, n)$ has a \mathbb{Z} -grading structure $\Lambda(n, n) = \bigoplus_{i \geq 0} \Lambda(n, n)_{[i]}$, where

$$\Lambda(n, n)_{[i]} = \text{span}_{\mathbb{F}}\{x^{(\alpha)} x^u \mid |\alpha| + |u| = i, \alpha \in \mathbb{N}_0^n, u \in \mathbb{B}(n)\}.$$

An arbitrary element $f \in \Lambda(n, n)$ can be uniquely decomposed into $f = \sum_{i=0}^{\infty} f_i$, where $f_i \in \Lambda(n, n)_{[i]}$. The continuation of the addition of topological algebra $\Lambda(n, n)$ allows us to get the sum of infinite nonzero elements of $\Lambda(n, n)$. Set $\Lambda(n, n)_j = \bigoplus_{i \geq j} \Lambda(n, n)_{[i]}$. Then $\{\Lambda(n, n)_j\}_{j \geq 0}$ is a filtration of $\Lambda(n, n)$. Clearly, $\Lambda(n, n)_j = \{(\Lambda_1)^j\}$, where $j \in \mathbb{N}_0$.

Let D_1, D_2, \dots, D_{2n} be the linear transformations of $\Lambda(n, n)$ such that

$$D_i(x^{(\alpha)}x^u) = \begin{cases} x^{(\alpha-\varepsilon_i)}x^u & \text{if } i \in Y_0, \\ x^{(\alpha)} \cdot (\partial x^u / \partial x_i) & \text{if } i \in Y_1. \end{cases}$$

Then D_i is a derivation of superalgebra $\Lambda(n, n)$ for every $i \in Y$. Let $\text{Der } \Lambda(n, n)$ be the Lie superalgebra consisting of all continuous derivations of $\Lambda(n, n)$. Then $\text{Der } \Lambda(n, n) = W(n, n)$, where $W(n, n) = \{\sum_{i=1}^{2n} f_i D_i \mid f_i \in \Lambda(n, n)\}$, and we call $W(n, n)$ the general superalgebra of formal vector fields (see [Kac 1998]). Clearly, $W(n, n)$ has a \mathbb{Z} -grading structure $W(n, n) = \bigoplus_{i \geq -1} W(n, n)_{[i]}$, where $W(n, n)_{[i]} = \text{span}_{\mathbb{F}}\{f D_j \mid f \in \Lambda(n, n)_{[i+1]}, j \in Y\}$. Let

$$W(n, n)_j = \bigoplus_{i \geq j} W(n, n)_{[i]}.$$

Then $\{W(n, n)_j\}_{j \geq -1}$ is called the natural filtration of $W(n, n)$. Therefore, $W(n, n)$ is a linearly compact topological Lie superalgebra with $\{W(n, n)_j\}_{j \geq -1}$ as a fundamental system of neighborhoods of 0.

If $\text{deg } f$ appears in some expression in this paper, we always regard f as a \mathbb{Z}_2 -homogenous element and $\text{deg } f$ as the \mathbb{Z}_2 -degree of f . Then $\text{deg } D_i = \mu(i)$, where

$$\mu(i) = \begin{cases} \bar{0} & \text{if } i \in Y_0, \\ \bar{1} & \text{if } i \in Y_1. \end{cases}$$

The following formula holds in $W(n, n)$ (see [Zhang 1997]):

$$[f D_i, g D_j] = f D_i(g) D_j - (-1)^{\text{deg } f D_i \text{ deg } g D_j} g D_j(f) D_i,$$

where $f, g \in \Lambda(n, n)$ and $i, j \in Y$.

Put

$$i' = \begin{cases} i + n & \text{if } i \in Y_0, \\ i - n & \text{if } i \in Y_1. \end{cases}$$

Let $T_H : \Lambda(n, n) \rightarrow W(n, n)$ be the linear mapping such that

$$(1) \quad T_H(f) = \sum_{i=1}^{2n} (-1)^{\mu(i) \text{ deg } f} D_i(f) D_{i'}.$$

Put $HO(n) = \{T_H(f) \mid f \in \Lambda(n, n)\}$. Then $HO(n)$ is an infinite-dimensional Lie superalgebra (see [Kac 1998]), called the odd Hamiltonian superalgebra of formal

vector fields. For $f, g \in \Lambda(n, n)$ the equation

$$(2) \quad [T_H(f), T_H(g)] = T_H(T_H(f)(g))$$

holds (see [Kac 1998]). Clearly, the algebra $HO(n)$ has a \mathbb{Z} -grading structure $HO(n) = \bigoplus_{i \geq -1} HO(n)_{[i]}$, where $HO(n)_{[i]} = \{T_H(f) \mid f \in \Lambda(n, n)_{[i+2]}\}$. Set $HO(n)_i = HO(n) \cap W(n, n)_i$. Then $\{HO(n)_i\}_{i \geq -1}$ is called the natural filtration of $HO(n)$.

Let $HO(n, n)$ be the \mathbb{Z}_2 -graded space $\Lambda(n, n)/\mathbb{F} \cdot 1$ with reversed parity, that is, $HO(n, n) = HO(n, n)_{\bar{0}} + HO(n, n)_{\bar{1}}$, where

$$HO(n, n)_{\theta} = \text{span}_{\mathbb{F}}\{x^{(\alpha)} x^u \in \Lambda(n, n)_{\theta+\bar{1}} \mid |\alpha| + |u| \geq 1\}, \quad \theta \in \mathbb{Z}_2.$$

We denote by $p(y)$ the \mathbb{Z}_2 -degree of the element y of $HO(n, n)$ to distinguish it from the \mathbb{Z}_2 -degree in $\Lambda(n, n)$. By (2), we can define a Lie multiplication in $HO(n, n)$ by

$$(3) \quad [y, z] = \sum_{i=1}^{2n} (-1)^{\mu(i)p(y)+\mu(i)} D_i(y) D_{i'}(z).$$

Clearly, Lie superalgebra $HO(n, n)$ is isomorphic to $HO(n)$.

Let $\Delta = \sum_{i=1}^n D_i D_{i'}$ be a linear mapping on $\Lambda(n, n)$, let

$$\Lambda^\Delta(n, n) = \{f \in \Lambda(n, n) \mid \Delta f = 0\},$$

and let $\overline{SHO}(n, n) = \Lambda^\Delta(n, n)/\mathbb{F} \cdot 1$. Then $\overline{SHO}(n, n)$ is a \mathbb{Z}_2 -graded subspace of $HO(n, n)$. For $f, g \in \Lambda(n, n)$ we have

$$\Delta(T_H(f)(g)) = (-1)^{\deg f + \bar{1}} T_H(f)(\Delta g) - (-1)^{\deg f \deg g + \deg f} T_H(g)(\Delta f);$$

see [Kac 1998]. Therefore, with the multiplication defined in (3), $\overline{SHO}(n, n)$ is a subalgebra of $HO(n, n)$. Set

$$(4) \quad SHO(n, n) = \text{span}_{\mathbb{F}} \{[x^{(\alpha)}, x^u] \mid \alpha \in \mathbb{N}_0^n, u \in \mathbb{B}(n), |\alpha| + |u| \geq 3\}.$$

Then $SHO(n, n)$ is an infinite-dimensional subalgebra of $\overline{SHO}(n, n)$, called the special odd Hamiltonian superalgebra of formal vector fields (see [Kac 1998]). Clearly, $SHO(n, n)$ has a \mathbb{Z} -grading structure $SHO(n, n) = \bigoplus_{i \geq -1} SHO(n, n)_{[i]}$, where

$$SHO(n, n)_{[i]} = \text{span}_{\mathbb{F}} \{[x^{(\alpha)}, x^u] \mid \alpha \in \mathbb{N}_0^n, u \in \mathbb{B}(n), |\alpha| + |u| = i + 4\}.$$

Set $SHO(n, n)_i = SHO(n, n) \cap W(n, n)_i$. Then $\{SHO(n, n)_i\}_{i \geq -1}$ is called the natural filtration of $SHO(n, n)$.

Set $SHO(n) = T_H(SHO(n, n))$. Clearly, with the multiplication defined in (2), $SHO(n)$ is a Lie superalgebra that is isomorphic to $SHO(n, n)$. For the sake of simplicity, we always write SHO for $SHO(n, n)$ or $SHO(n)$.

In the following, we simply write HO for $HO(n)$, and let X denote the Lie superalgebra HO or SHO .

3. Invariant subalgebras and natural filtrations

Lemma 3.1. *Suppose that $y \in X_{[-1]} \cap X_{\bar{0}}$ and that $y \neq 0$. Then*

$$y(\Lambda(n, n)_{[r]}) = \Lambda(n, n)_{[r-1]}$$

for all $r \in \mathbb{N}$; hence $y(\Lambda(n, n)) = \Lambda(n, n)$.

Proof. We first prove that $\Lambda(n, n)_{[r-1]} \subseteq y(\Lambda(n, n)_{[r]})$. Write $y = \sum_{j=1}^n c_j D_j$, where $c_j \in \mathbb{F}$. Then there exists at least one nonzero element in $\{c_1, \dots, c_n\}$. Let $c_{1j} = c_j$, where $1 \leq j \leq n$. Then there exist $c_{lj} \in \mathbb{F}$, where $2 \leq l \leq n, 1 \leq j \leq n$, such that the matrix $(c_{ij})_{1 \leq i, j \leq n}$ is invertible. Let $(a_{ij})_{1 \leq i, j \leq n} = (c_{ij})_{1 \leq i, j \leq n}^{-1}$. Note that $(1, 0, \dots, 0) (c_{ij})_{1 \leq i, j \leq n} = (c_1, c_2, \dots, c_n)$. It follows that

$$(5) \quad (1, 0, \dots, 0) = (c_1, c_2, \dots, c_n) (a_{ij})_{1 \leq i, j \leq n}.$$

Let $h_j = \sum_{i=1}^n a_{ij} x_i$, where $j \in Y_0$, and let $h_k = x_k$, where $k \in Y_1$. Then the set $\{h_1, h_2, \dots, h_{2n}\}$ is an \mathbb{F} -basis of $\Lambda(n, n)_{[1]}$. Therefore, for every $r \in \mathbb{N}$, we have

$$\Lambda(n, n)_{[r-1]} = \text{span}_{\mathbb{F}}\{h_1^{\alpha_1} \cdots h_n^{\alpha_n} h_{i_1} \cdots h_{i_k}\},$$

where $(\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$, $(i_1, \dots, i_k) \in \mathbb{B}_k$ and $\sum_{i=1}^n \alpha_i + k = r - 1$. Noting that $\alpha_1 \in \mathbb{N}_0$, we see that $\alpha_1 + 1 \neq 0$, since $\text{char } \mathbb{F} = 0$. By (5), we have $y(h_j) = \delta_{j1}$, where $j \in Y_0$. Consequently, we have

$$y((\alpha_1 + 1)^{-1} h_1 h_1^{\alpha_1} \cdots h_n^{\alpha_n} h_{i_1} \cdots h_{i_k}) = h_1^{\alpha_1} \cdots h_n^{\alpha_n} h_{i_1} \cdots h_{i_k}.$$

Thus $\Lambda(n, n)_{[r-1]} \subseteq y(\Lambda(n, n)_{[r]})$. The reverse inclusion can be verified trivially.

Suppose that $f = \sum_{s \geq 0} f_s$ is an arbitrary element of $\Lambda(n, n)$, where f_s is in $\Lambda(n, n)_{[s]}$. According to the results above, for every $s \in \mathbb{N}_0$, there exists a g_{s+1} in $\Lambda(n, n)_{[s+1]}$ such that $f_s = y(g_{s+1})$. Since y is continuous, it follows that $f = \sum_{s \geq 0} y(g_{s+1}) = y(\sum_{s \geq 0} g_{s+1}) \in y(\Lambda(n, n))$. Thus $\Lambda(n, n) = y(\Lambda(n, n))$. \square

Lemma 3.2. *Suppose that $y \in X_{[-1]} \cap X_{\bar{0}}$ and that $y \neq 0$. Then $[y, X_{[r]}] = X_{[r-1]}$ for all $r \in \mathbb{N}_0$.*

Proof. It suffices to show that $X_{[r-1]} \subseteq [y, X_{[r]}]$. Consider the case of HO . Suppose that $T_H(f)$ is an element of $HO_{[r-1]}$, where $f \in \Lambda(n, n)_{[r+1]}$. Then by Lemma 3.1 there exists $g \in \Lambda(n, n)_{[r+2]}$ such that $y(g) = f$, which combined with (2) yields that $T_H(f) = [y, T_H(g)] \in [y, HO_{[r]}]$.

Consider the case of SHO . Suppose that $[x^{(\alpha)}, x^u]$ is a standard basis element of $SHO_{[r-1]}$, and assume that $y = x_{i'}$, where $i \in Y_0$. Then by (3), we have that $-[x^{(\alpha)}, x^u] = [x_{i'}, [x^{(\alpha+\varepsilon_i)}, x^u]]$ is in $[y, SHO_{[r]}]$. \square

Lemma 3.3. *Suppose that $y \in X_{-1} \cap X_{\bar{0}} \setminus X_0$ and that $y \neq 0$. Then $[y, X] = X$.*

Proof. It suffices to show that $X \subseteq [y, X]$. Suppose that $y = \sum_{i \geq -1} y_i$, where $y_i \in X_{[i]}$, and suppose that $z = \sum_{i \geq -1} z_i$ is an arbitrary element of X , where $z_i \in X_{[i]}$. Then by Lemma 3.2, we can inductively pick $w_j \in X_{[j]}$, such that $[y_{-1}, w_0] = z_{-1}$ when $j = 0$, and $[y_{-1}, w_j] = z_{j-1} - \sum_{i=0}^{j-1} [y_i, w_{j-1-i}]$ when $j > 0$. For arbitrary $k \in \mathbb{N}_0$, direct calculations show that

$$(6) \quad \left[y_{-1}, \sum_{j=0}^k w_j \right] = [y_{-1}, w_0] + \sum_{j=1}^k [y_{-1}, w_j] = \sum_{j=-1}^{k-1} z_j - \sum_{0 \leq i+j \leq k-1} [y_i, w_j],$$

and

$$(7) \quad \left[\sum_{i \geq 0} y_i, \sum_{j=0}^k w_j \right] = \left[\sum_{i=0}^{k-1} y_i, \sum_{j=0}^k w_j \right] + \left[\sum_{i \geq k} y_i, \sum_{j=0}^k w_j \right] \\ = \sum_{0 \leq i+j \leq k-1} [y_i, w_j] + \sum_{i+j \geq k} [y_i, w_j].$$

Combining (6) and (7), we have

$$\left[y, \sum_{j=0}^k w_j \right] = \left[y_{-1}, \sum_{j=0}^k w_j \right] + \left[\sum_{i \geq 0} y_i, \sum_{j=0}^k w_j \right] \\ = \sum_{j=-1}^{k-1} z_j + \sum_{i+j \geq k} [y_i, w_j] \in \sum_{j=-1}^{k-1} z_j + X_k.$$

Noting that $X_k = \bigcap_{i=0}^k X_i$, we see that $[y, \sum_{j=0}^k w_j] \equiv \sum_{j=-1}^{k-1} z_j \pmod{\bigcap_{i=0}^k X_i}$. Let $w = \sum_{j \geq 0} w_j$. Then $[y, w] = [y, \sum_{j \geq 0} w_j] \equiv \sum_{j \geq -1} z_j \pmod{\bigcap_{i \geq 0} X_i}$, whence $[y, w] = z$. Thus $X \subseteq [y, X]$. \square

For an element y of Lie superalgebra L , we call y ad-nilpotent if there exists a positive integer t such that $(\text{ad } y)^t(L) = 0$. We call y ad-quasi-nilpotent if $\bigcap_{t=1}^{\infty} (\text{ad } y)^t(L) = 0$ (see [Humphreys 1972; Jin 1992]). Obviously, ad-nilpotent elements are ad-quasi-nilpotent elements. In particular, D_i is an ad-nilpotent element of X for every $i \in Y_1$.

Let J be a subset of L . Put

$$\text{qn}_L(J) := \{y \in J \mid y \text{ is an ad-quasi-nilpotent element of } L\}.$$

In the following, we simply write $\text{qn}(J)$ for $\text{qn}_X(J)$, and denote by $\text{Qn}(J)$ the subalgebra of X generated by $\text{qn}(J)$. It is clear that $X_1 \subseteq \text{qn}(X)$. In the following, we will determine the ad-quasi-nilpotent elements of $X_{[0]}$, and prove the invariance of natural filtration of X .

We denote by $M_{2n}(\Lambda(n, n))$ the \mathbb{F} -algebra consisting of all $2n \times 2n$ matrices over $\Lambda(n, n)$, denote by $\text{pr}_{[0]}$ the projection of $\Lambda(n, n)$ on $\Lambda(n, n)_{[0]}$, and denote

by pr_1 the projection on $\Lambda(n, n)_1$. For $(a_{ij})_{i,j \in Y} \in \mathbf{M}_{2n}(\Lambda(n, n))$, we also denote

$$\text{pr}_{[0]} : (a_{ij})_{i,j \in Y} \mapsto (\text{pr}_{[0]}(a_{ij}))_{i,j \in Y} \quad \text{and} \quad \text{pr}_1 : (a_{ij})_{i,j \in Y} \mapsto (\text{pr}_1(a_{ij}))_{i,j \in Y}.$$

Lemma 3.4. *Suppose that $h_1, h_2, \dots, h_{2n} \in \Lambda(n, n)_1$ with $\deg(h_j) = \mu(j)$ such that the matrix $(\text{pr}_{[0]}(D_i h_j))_{i,j \in Y}$ is invertible. Then there exists an automorphism σ of $\Lambda(n, n)$ such that*

$$(8) \quad \sigma(x_i) = h_i \quad \text{for all } i \in Y.$$

Proof. Let $\sigma : \Lambda(n, n) \rightarrow \Lambda(n, n)$ be an even endomorphism such that (8) holds. Note that the natural filtration of $\Lambda(n, n)$ is invariant under σ . Then σ induces a linear transformation σ_i of $\Lambda(n, n)_i / \Lambda(n, n)_{i+1}$ for every $i \geq 0$. We first use induction on k to show that σ_k is bijective. Since the matrix of σ_1 with respect to \mathbb{F} -basis $\{x_1 + \Lambda(n, n)_2, \dots, x_{2n} + \Lambda(n, n)_2\}$ is just $(\text{pr}_{[0]}(D_i h_j))_{i,j \in Y}$, we see that σ_1 is bijective. Suppose that $k > 1$ and $x^{(\alpha)}x^u$ is an element of $\Lambda(n, n)_{[k]}$. Then we can write $x^{(\alpha)}x^u = f_j f_{k-j}$, where $f_j \in \Lambda(n, n)_{[j]}$ and $f_{k-j} \in \Lambda(n, n)_{[k-j]}$ with $1 \leq j < k$. By induction, there exist $f'_j \in \Lambda(n, n)_{[j]}$ and $f'_{k-j} \in \Lambda(n, n)_{[k-j]}$ such that $\sigma(f'_j) \equiv f_j \pmod{\Lambda(n, n)_{j+1}}$ and $\sigma(f'_{k-j}) \equiv f_{k-j} \pmod{\Lambda(n, n)_{k-j+1}}$, whence

$$\sigma(f'_j f'_{k-j}) = \sigma(f'_j)\sigma(f'_{k-j}) \equiv f_j f_{k-j} = x^{(\alpha)}x^u \pmod{\Lambda(n, n)_{k+1}}.$$

Thus σ_k is surjective. Note that since $\Lambda(n, n)_k / \Lambda(n, n)_{k+1}$ is finite-dimensional, it follows that σ_k is bijective.

We next prove that σ is bijective. Suppose that $f \in \ker(\sigma) \cap \Lambda(n, n)_i$ for any $i \geq 0$. Then $\sigma_i(f + \Lambda(n, n)_{i+1}) = 0$. It follows that $f \in \Lambda(n, n)_{i+1}$, since σ_i is injective. Thus $\ker(\sigma) \subseteq \bigcap_{j \geq i} \Lambda(n, n)_j = 0$, and σ is injective. Suppose that $g = g_0 + g_1 \in \Lambda(n, n)$, where $g_0 \in \mathbb{F}$, $g_1 \in \Lambda(n, n)_1$. Since σ_1 is surjective, there exists $g'_1 \in \Lambda(n, n)_1$ such that $\sigma_1(g'_1 + \Lambda(n, n)_2) = g_1 + \Lambda(n, n)_2$. It follows that $g_2 := g_1 - \sigma(g'_1) \in \Lambda(n, n)_2$. Note that σ_i is surjective for every $i \geq 0$. Then we can inductively pick $g'_i \in \Lambda(n, n)_i$, and define $g_{i+1} \in \Lambda(n, n)_{i+1}$ by

$$(9) \quad g_{i+1} := g_i - \sigma(g'_i).$$

Let $g' = g_0 + \sum_{i \geq 0} g'_i$. Since σ is continuous, it follows from (9) that

$$\sigma(g') = \sigma(g_0) + \sum_{i \geq 0} \sigma(g'_i) = g_0 + \sum_{i \geq 1} (g_i - g_{i+1}) = g_0 + g_1 = g,$$

whence σ is surjective. □

Let ρ be the corresponding representation with respect to $X_{[0]}$ -module $X_{[-1]}$, that is, $\rho(y) = \text{ad } y|_{X_{[-1]}}$ for all $y \in X_{[0]}$. It is easy to see that ρ is faithful. For $y \in X_{[0]}$, we also denote by $\rho(y)$ the matrix of $\rho(y)$ relative to the fixed ordered

\mathbb{F} -basis $\{D_1, D_2, \dots, D_{2n}\}$. Denote by $\mathfrak{gl}(n, n)$ the general linear Lie superalgebra of $2n \times 2n$ matrices over \mathbb{F} (see [Scheunert 1979]).

Lemma 3.5. *Suppose that A is an invertible matrix of $\mathfrak{gl}(n, n)$, and $y \in W(n, n)_{[0]}$. Then there exists an automorphism φ of $W(n, n)$ such that $\rho(\varphi(y)) = A\rho(y)A^{-1}$.*

Proof. Suppose that $A = (a_{ij})_{1 \leq i, j \leq 2n}$, and let $h_j = \sum_{i=1}^{2n} a_{ij}x_i$, where $1 \leq j \leq 2n$. Then $\{h_1, h_2, \dots, h_{2n}\}$ is an \mathbb{F} -basis of $\Lambda(n, n)_{[1]}$. Note that $D_i(h_j) = a_{ij}$ for all $i, j \in Y$. It follows from Lemma 3.4 that there exists $\sigma \in \text{Aut } \Lambda(n, n)$ such that $\sigma(x_i) = h_i$ for all $i \in Y$. Clearly, $\sigma \in \text{Aut}(\Lambda(n, n) : W(n, n))$.

Let $\varphi : W(n, n) \rightarrow W(n, n)$ be the linear mapping such that $z \mapsto \sigma z \sigma^{-1}$ for all z in $W(n, n)$. Then φ is an automorphism of $W(n, n)$. We claim that φ is the desired automorphism. Suppose that $A^{-1} = (c_{ij})_{1 \leq i, j \leq 2n}$, and let $y = \sum_{s,r \in Y} b_{sr}x_s D_r$ be an arbitrary element of $W(n, n)_{[0]}$, where $b_{sr} \in \mathbb{F}$. Then $\rho(y) = (b_{sr})_{1 \leq s, r \leq 2n}$. Noting that $(\varphi(y))(\sigma x_i) = \sigma(yx_i)$, we see that $\varphi(y) = \sum_{t,j \in Y} \sigma(yx_t)D_j$. Thus

$$\begin{aligned} \varphi(y) &= \sum_{t,j \in Y} c_{tj} \sigma \left(\sum_{s \in Y} b_{st}x_s \right) D_j = \sum_{t,j,s \in Y} b_{st}c_{tj} \sigma(x_s) D_j \\ &= \sum_{t,j,s \in Y} b_{st}c_{tj} h_s D_j = \sum_{t,j,s,k \in Y} a_{ks} b_{st}c_{tj} x_k D_j. \end{aligned}$$

It follows that $\rho(\varphi(y)) = A\rho(y)A^{-1}$. □

Lemma 3.6. *Suppose that $y \in W(n, n)_{[0]}$. Then $\text{ad } y$ is a nilpotent linear transformation of $W(n, n)_{[r]}$ for every $r \geq -1$ if and only if $\rho(y)$ is a nilpotent matrix.*

Proof. If $\text{ad } y|_{W(n, n)_{[-1]}}$ is nilpotent, then the definition of ρ shows that $\rho(y)$ is a nilpotent matrix. Conversely, suppose that $\rho(y)$ is nilpotent. By Lemma 3.5, it suffices to consider the case when $\rho(y)$ is a strictly upper triangular matrix. Suppose that $y = \sum_{i,j \in Y, i < j} a_{ij}x_i D_j$, where $a_{ij} \in \mathbb{F}$.

We first prove that $\text{ad } x_i D_j$ is nilpotent linear transformation of $W(n, n)_{[r]}$ for every $r \geq -1$ when $i < j$. For any standard basis element $x^{(\alpha)}x^u D_k$ of $W(n, n)_{[r]}$, where $\alpha \in \mathbb{N}_0^n$, $u \in \mathbb{B}(n)$ and $k \in Y$, two cases arise.

Case 1. $j \in Y_0$. If $k \neq i$, then

$$(\text{ad } x_i D_j)^t (x^{(\alpha)}x^u D_k) = x_i^t D_j^t (x^{(\alpha)}x^u D_k) = 0$$

when $t \geq r + 2$. If $k = i$, then

$$(\text{ad } x_i D_j)^t (x^{(\alpha)}x^u D_i) = x_i^t D_j^t (x^{(\alpha)}x^u D_i) - t x_i^{t-1} D_j^{t-1} (x^{(\alpha)}x^u D_j) = 0$$

when $t \geq r + 3$.

Case 2. $j \in Y_1$. If $k \neq i$, then

$$(\text{ad } x_i D_j)^t (x^{(\alpha)}x^u D_k) = (\text{ad } x_i D_j)^{t-2} (x_i^2 x^{(\alpha)} D_j^2 (x^u) D_k) = 0$$

when $t \geq 2$, and if $k = i$, then

$$(\text{ad } x_i D_j)^t (x^{(\alpha)} x^u D_i) = l(\text{ad } x_i D_j)^{t-3} (x_i^2 x^{(\alpha)} D_j^2 (x^u) D_j) = 0$$

when $t \geq 3$, where $l = 1$ or $l = -1$.

Therefore $(\text{ad } x_i D_j)^t (x^{(\alpha)} x^u D_k) = 0$ when $t \geq r + 4$. Let $f_k = \sum_{\alpha,u} c_{\alpha,u} x^{(\alpha)} x^u$ be an arbitrary element of $\Lambda(n, n)_{[r+1]}$, where $c_{\alpha,u} \in \mathbb{F}$, $k \in Y$. Then for any $t \geq r + 4$,

$$(\text{ad } x_i D_j)^t (f_k D_k) = \sum_{\alpha,u} (\text{ad } x_i D_j)^t (c_{\alpha,u} x^{(\alpha)} x^u D_k) = 0,$$

since $(\text{ad } x_i D_j)^t$ is continuous. Consequently, we see that $\text{ad } x_i D_j|_{W(n,n)_{[r]}}$ is nilpotent for every $r \geq -1$ when $i < j$.

Note that the set $\{\pm x_i D_j, 0 | i < j\}$ is closed under the multiplication of $W(n, n)$, and the Lie superalgebra $\text{span}_{\mathbb{F}}\{\pm x_i D_j, 0 | i < j\}$ is finite-dimensional. It follows from [Strade and Farnsteiner 1988, Theorem 1.3.1] that $\text{ad } y|_{W(n,n)_{[r]}}$ is nilpotent. □

Lemma 3.7. *Suppose that $y \in X_{[0]}$. Then $\text{ad } y$ is a nilpotent linear transformation of $X_{[r]}$ for every $r \geq -1$ if and only if $\rho(y)$ is a nilpotent matrix.*

Proof. Clearly, $\rho(y)$ is a nilpotent matrix when $\text{ad } y|_{X_{[-1]}}$ is nilpotent. Conversely, suppose that $\rho(y)$ is a nilpotent matrix. Then by Lemma 3.6, $\text{ad } y$ is a nilpotent linear transformation of $W(n, n)_{[r]}$ for every $r \geq -1$. Since X is a subalgebra of $W(n, n)$, it follows that $\text{ad } y|_{X_{[r]}}$ is nilpotent. □

Lemma 3.8. *Suppose that $y = y_0 + y_1 \in \text{qn}(X_0)$, where $y_0 \in X_{[0]}$, $y_1 \in X_1$. Then $\rho(y_0)$ is a nilpotent matrix, and hence, $y_0 \in \text{qn}(X_{[0]})$.*

Proof. Let $X_{(i)} = X/X_{i+1}$ for every $i \geq -1$. Then $X_{(i)} \cong \bigoplus_{j \leq i} X_{[j]}$. For every $i \geq -1$, let τ_i be the endomorphism on $X_{(i)}$ satisfying $\tau_i(z) \equiv [y, z] \pmod{X_{i+1}}$ for all $z \in X_{(i)}$. Assume that $\rho(y_0)$ is not a nilpotent matrix. Then τ_i is not nilpotent for every $i \geq -1$. Let $X_{(i)} = U_i \oplus V_i$ be the Fitting decomposition of $X_{(i)}$ with respect to τ_i , where $U_i \neq 0$, $\tau_i|_{U_i}$ is invertible, $\tau_i|_{V_i}$ is nilpotent. Since $X_{(i)} = X_{(i+1)}/X_{[i+1]}$, $\tau_i = \tau_{i+1} \pmod{X_{i+1}}$ and $\tau_{i+1}(X_{[i+1]}) \subseteq (X_{[i+1]})$, it follows that

$$X_{(i)} = (U_{i+1} + X_{[i+1]}/X_{[i+1]}) \oplus (V_{i+1} + X_{[i+1]}/X_{[i+1]}).$$

is also the Fitting decomposition of $X_{(i)}$ with respect to τ_i , and by the uniqueness of the Fitting decomposition, we get $U_i = U_{i+1} + X_{[i+1]}/X_{[i+1]}$. This implies that U_i is the projection of U_{i+1} on $X_{(i)}$. Set

$$\bar{U} = \{z \in X \mid \text{the projection of } z \text{ on } X_{(i)} \text{ belongs to } U_i \text{ for all } i \geq -1\}.$$

By the completeness of X , the set \bar{U} is nonempty, and its projection on $X_{(i)}$ is U_i for each $i \geq -1$. It follows that $[y, \bar{U}] = \bar{U}$. So $\bigcap_{i=0}^{\infty} (\text{ad } y)^i(X) \supseteq \bar{U} \neq 0$,

contradicting the hypothesis that y is ad-quasi-nilpotent. Therefore, $\rho(y_0)$ is a nilpotent matrix, which combined with Lemma 3.7 yields $y_0 \in \text{qn}(X_{[0]})$. \square

Proposition 3.9. *One has $\text{qn}(X_0) = AN_0 \cap X_{[0]} + X_1$, where*

$$AN_0 = \{y \mid y \in W(n, n)_{[0]} \text{ such that } \rho(y) \text{ is a nilpotent matrix} \}.$$

Proof. By Lemma 3.8, it suffices to show that $AN_0 \cap X_{[0]} + X_1 \subseteq \text{qn}(X_0)$. Suppose that $y_0 \in AN_0 \cap X_{[0]}$, and suppose that $y_1 \in X_1$. Let $y = y_0 + y_1$. Then $\rho(y_0)$ is a nilpotent matrix. According to Lemma 3.7, there exists a positive integer t_i such that $(\text{ad } y_0)^{t_i}(X_{[i]}) = 0$ for each $i \geq -1$. This implies that $(\text{ad } y_0)^{t_i}(X_i) \subseteq X_{i+1}$. Consequently, we have $(\text{ad } y)^{t_1 + \dots + t_k}(X_0) \subseteq X_{k+1}$ for any $k \geq -1$. It follows that $\bigcap_{l=1}^{\infty} (\text{ad } y)^l(X_0) \subseteq \bigcap_{k=1}^{\infty} X_k = 0$, whence $y \in \text{qn}(X_0)$. \square

Lemma 3.10. $\text{Qn}(X_{\bar{0}}) = \text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}}$, and then $\text{Qn}(X_{\bar{0}}) \subseteq X_0 \cap X_{\bar{0}}$.

Proof. Note that $X_1 \subseteq \text{Qn}(X)$. It follows that $X_1 \cap X_{\bar{0}} \subseteq \text{Qn}(X_{\bar{0}})$. Consequently, we have $\text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}} \subseteq \text{Qn}(X_{\bar{0}})$.

Conversely, suppose that $y = y_{-1} + y_0 \in \text{qn}(X_{\bar{0}})$, where $y_{-1} \in X_{[-1]} \cap X_{\bar{0}}$, and $y_0 \in X_0 \cap X_{\bar{0}}$. Assume that $y_{-1} \neq 0$. It follows from Lemma 3.3 that $[y, X] = X$, which implies that y is not an ad-quasi-nilpotent element of X , a contradiction. Thus $y \in X_0 \cap X_{\bar{0}}$.

Now we can write $y = y_0 + y_1$, where $y_0 \in X_{[0]} \cap X_{\bar{0}}$, and $y_1 \in X_1 \cap X_{\bar{0}}$. By Lemma 3.8, we have $y_0 \in \text{qn}(X_{[0]} \cap X_{\bar{0}}) \subseteq \text{Qn}(X_{[0]} \cap X_{\bar{0}})$. It follows that $y \in \text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}}$. Thus

$$(10) \quad \text{qn}(X_{\bar{0}}) \subset \text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}}.$$

The right-hand side of (10) is a subalgebra of $X_{\bar{0}}$. Then

$$\text{Qn}(X_{\bar{0}}) \subset \text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}}.$$

Thus the lemma holds. \square

Lemma 3.11. *Suppose that $i, j \in Y$ with $i \neq j'$. Then $T_H(x_i x_j) \in \text{qn}(X_{[0]})$.*

Proof. It suffices to show that $\text{ad } T_H(x_i x_j)$ is a nilpotent linear transformation of $W(n, n)_{[t]}$ for every $t \geq -1$. Suppose that $x^{(\alpha)} x^u D_k$ is a standard basis element of $W(m, n)_{[t]}$, where $t \geq -1$. To simplify our proof for the lemma, we only verify the case $i \in Y_0, j \in Y_1$, and $k \neq i, j$ as the proofs for the other cases are similar and hence omitted.

An induction on l shows that

$$\begin{aligned} (\text{ad } T_H(x_i x_j))^l(x^{(\alpha)} x^u D_k) \\ = (-1)^{l-1} l x_i^{l-1} D_j^{l-1}(x^{(\alpha)}) x_j D_{i'}(x^u) D_k + (-1)^l x_i^l D_j^l(x^{(\alpha)}) x^u D_k = 0. \end{aligned}$$

This yields that $(\text{ad } T_H(x_i x_j))^l(x^{(\alpha)} x^u D_k) = 0$ when $l \geq t + 3$. \square

Let I_n denote the identity matrix of size $2n \times 2n$, and let e_{ij} denote the $2n \times 2n$ matrix whose (i, j) -entry is 1 and whose other entries are 0. Set

$$\tilde{p}(n) = \left\{ \begin{pmatrix} A & B \\ C & -A^T \end{pmatrix} \in \mathfrak{gl}(n, n) \mid B = -B^T, C = C^T \right\},$$

$$p(n) = \left\{ \begin{pmatrix} A & B \\ C & -A^T \end{pmatrix} \in \tilde{p}(n) \mid \text{tr } A = 0 \right\},$$

where A^T is the transpose of A . Then $\tilde{p}(n)$, $p(n)$ are subalgebras of $\mathfrak{gl}(n, n)$ (see [Kac 1998]). Clearly,

$$\tilde{p}(n)_{\bar{0}} = \left\{ \begin{pmatrix} A & 0 \\ 0 & -A^T \end{pmatrix} \in \mathfrak{gl}(n, n) \right\},$$

$$p(n)_{\bar{0}} = \left\{ \begin{pmatrix} A & 0 \\ 0 & -A^T \end{pmatrix} \in \mathfrak{gl}(n, n) \mid \text{tr } A = 0 \right\}.$$

Lemma 3.12. (1) $\rho(T_H(x_i x_j)) = (-1)^{\mu(i')} e_{i'j} - (-1)^{\mu(i')\mu(j)} e_{j'i}$ for all $i, j \in Y$, hence $\rho(HO_{[0]}) = \tilde{p}(n)$ and $\rho(HO_{[0]} \cap HO_{\bar{0}}) = \tilde{p}(n)_{\bar{0}}$.

(2) $\rho(T_H(x_i x_j)) = (-1)^{\mu(i')} e_{i'j} - (-1)^{\mu(i')\mu(j)} e_{j'i}$ for all $i, j \in Y$ with $i \neq j'$, $\rho(T_H(x_k x_{k'} - x_l x_{l'})) = e_{kk} - e_{k'k'} - e_{ll} + e_{l'l'}$ for all $k, l \in Y_0$ with $k \neq l$, hence $\rho(SHO_{[0]}) = p(n)$ and $\rho(SHO_{[0]} \cap SHO_{\bar{0}}) = p(n)_{\bar{0}}$.

Proof. (1) Direct calculation shows that $\text{ad } T_H(x_i x_j)(D_j) = (-1)^{\mu(i')} D_{i'}$, and $\text{ad } T_H(x_i x_j)(D_i) = -(-1)^{\mu(i')\mu(j)} D_{j'}$. It follows that

$$\rho(T_H(x_i x_j)) = (-1)^{\mu(i')} e_{i'j} - (-1)^{\mu(i')\mu(j)} e_{j'i}.$$

Note that $HO_0 = \text{span}_{\mathbb{F}}\{T_H(x_i x_j) \mid i, j \in Y\}$. Consequently, (1) holds.

(2) The proof is similar to that of (1), hence omitted. □

Lemma 3.13. $\rho(\text{Qn}(X_{[0]} \cap X_{\bar{0}})) = p(n)_{\bar{0}}$.

Proof. Suppose that $y \in \text{qn}(X_{[0]} \cap X_{\bar{0}})$. Then $\rho(y)$ is a nilpotent matrix by Lemma 3.8. It follows from Lemma 3.12 that $\rho(y) = \text{diag}(A, -A^T)$, where A and $-A^T$ are $n \times n$ nilpotent matrices. This shows that $\text{tr } A = 0$, that is $\rho(y) \in p(n)_{\bar{0}}$. Thus $\rho(\text{qn}(X_{[0]} \cap X_{\bar{0}})) \subseteq p(n)_{\bar{0}}$.

Conversely, set

$$R = \{T_H(x_i x_j) \mid i \in Y_0, j \in Y_1, \text{ with } i \neq j'\}.$$

Then $R \subseteq \text{qn}(X_{[0]} \cap X_{\bar{0}})$, by Lemma 3.11, whence $\rho(R) \subseteq \rho(\text{qn}(X_{[0]} \cap X_{\bar{0}}))$. Noting that $\rho(T_H(x_i x_j)) = e_{j'i} - e_{i'j}$ for all $i \in Y_0, j \in Y_1$ with $i \neq j'$, by Lemma 3.12, we see that $\rho(R)$ generates $p(n)_{\bar{0}}$. Thus $p(n)_{\bar{0}} \subseteq \rho(\text{Qn}(X_{[0]} \cap X_{\bar{0}}))$. □

Note that $\text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}})) = \{y \in X_{\bar{0}} \mid [y, \text{Qn}(X_{\bar{0}})] \subseteq \text{Qn}(X_{\bar{0}})\}$. Clearly, the set $\text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}}))$ is invariant under automorphisms of X .

Proposition 3.14. $X_0 \cap X_{\bar{0}} = \text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}}))$. In particular, $X_0 \cap X_{\bar{0}}$ is an invariant subalgebra.

Proof. Note that $[\tilde{p}(n)_{\bar{0}}, p(n)_{\bar{0}}] = p(n)_{\bar{0}}$. It follows from Lemmas 3.12 and 3.13 that

$$[\rho(X_{[0]} \cap X_{\bar{0}}), \rho(\text{Qn}(X_{[0]} \cap X_{\bar{0}}))] = [\tilde{p}(n)_{\bar{0}}, p(n)_{\bar{0}}] = \rho(\text{Qn}(X_{[0]} \cap X_{\bar{0}})),$$

whence

$$(11) \quad [X_{[0]} \cap X_{\bar{0}}, \text{Qn}(X_{[0]} \cap X_{\bar{0}})] = \text{Qn}(X_{[0]} \cap X_{\bar{0}}),$$

since ρ is faithful. By Lemma 3.10 and (11), we have

$$\begin{aligned} [X_0 \cap X_{\bar{0}}, \text{Qn}(X_{\bar{0}})] &= [X_{[0]} \cap X_{\bar{0}} + X_1 \cap X_{\bar{0}}, \text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}}] \\ &\subseteq [X_{[0]} \cap X_{\bar{0}}, \text{Qn}(X_{[0]} \cap X_{\bar{0}})] + X_1 \cap X_{\bar{0}} \\ &\subseteq \text{Qn}(X_{[0]} \cap X_{\bar{0}}) + X_1 \cap X_{\bar{0}} = \text{Qn}(X_{\bar{0}}). \end{aligned}$$

Thus $X_0 \cap X_{\bar{0}} \subseteq \text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}}))$.

Conversely, suppose that $y = y_{-1} + y_0 \in \text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}}))$, where $y_{-1} \in X_{[-1]} \cap X_{\bar{0}}$, $y_0 \in X_0 \cap X_{\bar{0}}$. We want to show that $y_{-1} = 0$. Assume $y_{-1} = \sum_{t=1}^n a_t D_t \neq 0$, where $a_t \in \mathbb{F}$. Then we can pick $k \in Y_0$ such that $a_k \neq 0$, and then pick $j_k \in Y_1$ such that $j_k \neq k'$. From Lemma 3.11 we see that $T_H(x_k x_{j_k}) \in \text{Qn}(X_{\bar{0}})$, which combined with our hypothesis that $y \in \text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}}))$, yields $[y, T_H(x_k x_{j_k})] \in \text{Qn}(X_{\bar{0}})$, whence $[y, T_H(x_k x_{j_k})] \in X_0 \cap X_{\bar{0}}$ by Lemma 3.10. On the other hand, a direct calculation shows that

$$[y, T_H(x_k x_{j_k})] = [y_{-1}, T_H(x_k x_{j_k})] + [y_0, T_H(x_k x_{j_k})] = -a_k D_{j'_k} + [y_0, T_H(x_k x_{j_k})].$$

Since $[y_0, T_H(x_k x_{j_k})] \in X_0 \cap X_{\bar{0}}$, we see that $a_k = 0$, contradicting our assumption, thus $y_{-1} = 0$. So $y = y_0 \in X_0 \cap X_{\bar{0}}$, proving $\text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}})) \subseteq X_0 \cap X_{\bar{0}}$. Since $\text{Nor}_{X_{\bar{0}}}(\text{Qn}(X_{\bar{0}}))$ is invariant, we see that $X_0 \cap X_{\bar{0}}$ is invariant. \square

Set $\Omega = \{y \in \text{qn}(X_0 \cap X_{\bar{0}}) \mid [y, X_0 \cap X_{\bar{0}}] \subseteq \text{qn}(X_0 \cap X_{\bar{0}})\}$. Then Proposition 3.14 shows that Ω is invariant under automorphisms of X .

Proposition 3.15. $X_1 \cap X_{\bar{0}} = \Omega$. In particular, $X_1 \cap X_{\bar{0}}$ is an invariant subalgebra.

Proof. We only verify the case of SHO , as the proof for HO is similar and hence omitted. Suppose that $y = y_0 + y_1$ is an arbitrary element of Ω , where y_0 is in $SHO_{[0]} \cap SHO_{\bar{0}}$ and y_1 is in $SHO_1 \cap SHO_{\bar{0}}$. Suppose that

$$y_0 = \sum_{i,j \in Y_0, i \neq j} a_{ij} T_H(x_i x_{j'}) + \sum_{i=1}^{n-1} a_{ii} T_H(x_i x_{i'} - x_{i+1} x_{(i+1)'}),$$

where $a_{ij} \in \mathbb{F}$. Let $b_{ij} = a_{ij}$, where $i, j \in Y_0$ with $i \neq j$, and let $b_{ii} = a_{ii} - a_{i-1, i-1}$, where $i \in Y_0$, and $a_{00} = a_{nn} = 0$. Let

$$l = \min \{i \in Y_0 \mid b_{ij_0} \neq 0 \text{ for some } j_0 \in Y\},$$

$$t = \min \{j \in Y_0 \mid b_{i_0j} \neq 0 \text{ for some } i_0 \in Y\}.$$

We first consider the case $l \leq t$. Set $k = \max\{j \in Y_0 \mid b_{lj} \neq 0\}$. Then $l \leq t \leq k$, and $b_{lk} \neq 0$. If $l = k$, then

$$y_0 = \sum_{i=l+1}^n \sum_{j=l, i \neq j}^n a_{ij} T_H(x_i x_j) + \sum_{i=l}^{n-1} a_{ii} T_H(x_i x_{i'} - x_{i+1} x_{(i+1)'}),$$

and

$$\rho(y_0) = \sum_{i=l}^n a_{ii} (e_{ii} - e_{i+1i+1}) - \sum_{i=l}^n a_{ii} (e_{i'i'} - e_{(i+1)'(i+1)'}) + \sum_{i=l+1}^n \sum_{j=l, i \neq j}^n a_{ij} (e_{ji} - e_{i'j'}) = \begin{pmatrix} a_{ll} A_{ll} & * \\ 0 & * \end{pmatrix},$$

where A_{ll} is the $l \times l$ matrix whose (l, l) -entry is 1 and 0 elsewhere. Since $a_{ll} \neq 0$, we conclude that $\rho(y_{[0]})$ is not a nilpotent matrix. It follows from Lemma 3.8 that $y \notin \text{qn}(SHO_0)$, contradicting the hypothesis that $y \in \Omega$. Therefore $l < k$ and

$$y_0 = \sum_{j=t, j \neq l}^k a_{lj} T_H(x_l x_j) + \sum_{i=l+1}^n \sum_{j=t, i \neq j}^n a_{ij} T_H(x_i x_j) + \sum_{i=t}^n a_{ii} T_H(x_i x_{i'} - x_{i+1} x_{(i+1)'}).$$

A direct calculation shows that

$$\begin{aligned} \rho([T_H(x_k x_{l'}), y_0]) &= [e_{lk} - e_{k'l'}, \rho(y_0)] \\ &= a_{lk} (e_{ll} - e_{l'l'} - e_{kk} + e_{k'k'}) - \sum_{j=t, j \neq l}^{k-1} a_{lj} (e_{jk} - e_{k'j'}) \\ &\quad + \sum_{i=l+1, i \neq k}^n a_{ik} (e_{li} - e_{l'i'}) + (a_{kk} - a_{k-1, k-1} - \delta_{ll} a_{ll}) (e_{lk} - e_{k'l'}) \\ &= \begin{pmatrix} a_{lk} A_{ll} & * \\ 0 & * \end{pmatrix}, \end{aligned}$$

so $\rho([T_H(x_k x_{l'}), y_0])$ is not a nilpotent matrix. Since

$$[T_H(x_k x_{l'}), y] = [T_H(x_k x_{l'}), y_0] + [T_H(x_k x_{l'}), y_1],$$

it follows from Lemma 3.8 that $[T_H(x_k x_{l'}), y] \notin \text{qn}(SHO_0)$, contradicting our hypothesis that $y \in \Omega$.

We now consider the case $l > t$. Set $k = \max\{i \in Y_0 \mid b_{it} \neq 0\}$. Then $t < l \leq k$, $b_{kt} \neq 0$ and

$$y_0 = \sum_{i=l}^k a_{it} T_H(x_i x_{i'}) + \sum_{i=l}^n \sum_{j=t+1, i \neq j}^n a_{ij} T_H(x_i x_{j'}) + \sum_{i=l}^n a_{ii} T_H(x_i x_{i'} - x_{i+1} x_{(i+1)'})$$

A direct computation shows that

$$\begin{aligned} \rho([T_H(x_t x_{t'}), y_0]) &= [e_{kt} - e_{t'k'}, \rho(y_0)] \\ &= -a_{kt}(e_{tt} - e_{t't'} - e_{kk} + e_{k'k'}) + \sum_{i=l}^{k-1} a_{it}(e_{ki} - e_{i'k'}) \\ &\quad - \sum_{j=t+1, j \neq k}^n a_{kj}(e_{jt} - e_{t'j'}) + (a_{k-1, k-1} - a_{k, k})(e_{kt} - e_{t'k'}) \\ &= \begin{pmatrix} -a_{kt} B_{tt} & 0 \\ * & * \end{pmatrix}, \end{aligned}$$

where B_{tt} is the $t \times t$ matrix whose (t, t) -entry is 1 and 0 elsewhere. It follows that $\rho([T_H(x_t x_{t'}), y_0])$ is not a nilpotent matrix, whence $[T_H(x_t x_{t'}), y] \notin \text{qn}(SHO_0)$ by Lemma 3.8, a contradiction which yields $y_0 = 0$. Therefore $y = y_1 \in SHO_1 \cap SHO_{\bar{0}}$, proving $\Omega \subseteq SHO_1 \cap SHO_{\bar{0}}$.

Conversely, noting that $SHO_1 \subseteq \text{qn}(SHO)$, we see that

$$\begin{aligned} [SHO_1 \cap SHO_{\bar{0}}, SHO_0 \cap SHO_{\bar{0}}] \\ \subseteq SHO_1 \cap SHO_{\bar{0}} \subseteq SHO_0 \cap SHO_{\bar{0}} \cap \text{qn}(SHO) = \text{qn}(SHO_0 \cap SHO_{\bar{0}}). \end{aligned}$$

Thus $SHO_1 \cap SHO_{\bar{0}} \subseteq \Omega$. Since Ω is invariant, we see that $SHO_1 \cap SHO_{\bar{0}}$ is invariant. \square

Lemma 3.16. $[X_{\bar{1}}, X_1 \cap X_{\bar{0}}] = X_0 \cap X_{\bar{1}}$.

Proof. It suffices to show that $X_0 \cap X_{\bar{1}} \subseteq [X_{\bar{1}}, X_1 \cap X_{\bar{0}}]$. We first consider the case of HO . Suppose that $T_H(x^{(\alpha)} x^u) \in HO_0 \cap HO_{\bar{1}}$, where $\alpha \in \mathbb{N}_0^n$, $u \in \mathbb{B}(n)$. Note that if $|u| \neq n$, then there exists $k \in Y_1 \setminus \{u\}$ such that

$$T_H(x^{(\alpha)} x^u) = [D_k, T_H(x^{(\alpha)} x_k x^u)] \in [HO_{\bar{1}}, HO_1 \cap HO_{\bar{0}}],$$

and if $|u| = n$, then

$$-T_H(x^{(\alpha)} x^u) = [T_H(x_i x_j), T_H(x_i x^{(\alpha)} D_j(x^u))] \in [HO_{\bar{1}}, HO_1 \cap HO_{\bar{0}}],$$

for all $i \in Y_0, j \in Y_1$.

Next, we consider the case of SHO . Suppose that $[x^{(\alpha)}, x^u] \in SHO_0 \cap SHO_{\bar{1}}$, where $\alpha \in \mathbb{N}_0^n, u \in \mathbb{B}(n)$. If $|u| \neq n$, then there exists $k \in Y_1$ such that $k \notin \{u\}$. It

follows from (3) that

$$-[x^{(\alpha)}, x^u] = [x_{k'}, [x^{(\alpha)}, x_k x^u]] \in [SHO_{\bar{1}}, SHO_1 \cap SHO_{\bar{0}}].$$

If $|u| = n$, then by the hypothesis, we see that $|u|$ is even. It follows that

$$[x^{(\alpha)}, x^u] = [x_{3'} \cdots x_{n'}, [x^{(\alpha+\varepsilon_3)}, x_{3'} x_{1'} x_{2'}]] \in [SHO_{\bar{1}}, SHO_1 \cap SHO_{\bar{0}}]. \quad \square$$

Theorem 3.17. *The natural filtration of X is invariant under automorphisms of X .*

Proof. By Proposition 3.15, we see that $X_1 \cap X_{\bar{0}}$ is invariant under automorphisms of X . It follows from Lemma 3.16 that $X_0 \cap X_{\bar{1}}$ is invariant, which combined with Proposition 3.14 yields that X_0 is invariant. Noting that

$$X_{[i]} = \{y \in X \mid [y, X] \subseteq X_{[i-1]}\}$$

for every $i \geq 1$, we see that $X_{[i]}$ is invariant. Thus X is invariant. □

4. The automorphism group of X

Lemma 4.1. *Suppose that $\varphi_X \in \text{Aut } X$. Then:*

- (1) φ_X is a continuous automorphism.
- (2) There exists an \mathbb{F} -basis $\{e_1, e_2, \dots, e_{2n}\}$ of $X_{[-1]}$ such that $\varphi_X(D_i) \equiv e_i \pmod{X}$.

Proof. (1) By Theorem 3.17, we have $\varphi_X(X_i) \subseteq X_i$ for every $i \geq -1$. It follows that $X_i \subseteq \varphi_X^{-1}(X_i)$. On the other hand, noting that $\varphi_X^{-1} \in \text{Aut } X$, we see that $\varphi_X^{-1}(X_i) \subseteq X_i$. Consequently, φ_X is a continuous automorphism.

(2) By Theorem 3.17, φ_X can induce an \mathbb{F} -isomorphism $\bar{\varphi}_X$ of the quotient spaces

$$\bar{\varphi}_X : X/X_0 \rightarrow X/X_0,$$

such that $\bar{\varphi}_X(y + X_0) = \varphi_X(y) + X_0$ for all $y \in X$. Since $\{D_i + X_0 \mid i \in Y\}$ is an \mathbb{F} -basis of X/X_0 , it follows that $\{\varphi_X(D_i) + X_0 \mid i \in Y\}$ is an \mathbb{F} -basis of X/X_0 . Then for every $i \in Y$, there exists $e_i \in X_{[-1]}$ such that $\varphi_X(D_i) + X_0 = e_i + X_0$. Thus (2) holds. □

Proposition 4.2. *Suppose that $\varphi, \psi \in \text{Aut } X$. If $\varphi|_{X_{[-1]}} = \psi|_{X_{[-1]}}$, then $\varphi = \psi$.*

Proof. We first use induction on k to show that $\varphi|_{X_{[k]}} = \psi|_{X_{[k]}}$, where $k \geq -1$. The result is obvious for $k = -1$. We assume it for $k - 1$. Suppose that $y \in X_{[k]}$, and let $z = \varphi(y) - \psi(y)$. Then

$$[z, \psi(D_i)] = [\varphi(y) - \psi(y), \psi(D_i)] = \varphi([y, D_i]) - \psi([y, D_i]) = 0$$

for every $i \in Y$. By Lemma 4.1 (2), we can write $\psi(D_i) = e_i + w_i$, where $\{e_1, \dots, e_{2n}\}$ is an \mathbb{F} -basis of $X_{[-1]}$ and $w_i \in X_0$. It follows that $[z, e_i + w_i] = 0$.

Suppose that $e_i = \sum_{j \in Y} a_{ij} D_j$, where $a_{ij} \in \mathbb{F}$, and let $(c_{ij})_{1 \leq i, j \leq 2n} = (a_{ij})_{1 \leq i, j \leq 2n}^{-1}$. Then $[z, -w_i] = [z, e_i] = \sum_{j \in Y} a_{ij} [z, D_j]$, whence

$$(12) \quad [z, D_l] = \sum_{i=1}^{2n} c_{li} [z, -w_i] \quad \text{for all } l \in Y.$$

Noting that $y \in X_0$, we see that $z \in X_0$. Thus we can write $z = \sum_{j \geq 0} z_j$, where $z_j \in X_{[j]}$. Applying (12), we have $[z, D_l] \in X_0$, thus $[z_0, D_l] \in X_0 \cap X_{[-1]} = 0$ for all $l \in Y$. This yields $z_0 = 0$. Now we can write $z = \sum_{j \geq 1} z_j$. Repeating the argument above, we can see that $z_j = 0$ for each $j \geq 1$. It follows that $z = 0$, whence $\varphi|_{X_{[k]}} = \psi|_{X_{[k]}}$. Hence $\varphi = \psi$, by Lemma 4.1 (1). \square

Given $\sigma \in \text{Aut } \Lambda(n, n)$ and $D \in \text{Der } \Lambda(n, n)$, we set $D^\sigma = \sigma D \sigma^{-1}$. Then $\tilde{\sigma} : D \mapsto D^\sigma$ is an automorphism of $\text{Der } \Lambda(n, n)$. Let

$$\text{Aut}(\Lambda(n, n) : X) = \{ \sigma \in \text{Aut } \Lambda(n, n) \mid \tilde{\sigma}(X) \subseteq X \}.$$

Then $\text{Aut}(\Lambda(n, n) : X)$ is a subgroup of $\text{Aut } \Lambda(n, n)$, and it is called the admissible automorphism group of $\Lambda(n, n)$ relative to X . Obviously, the morphism $\Phi : \text{Aut}(\Lambda(n, n) : X) \rightarrow \text{Aut } X$ given by $\sigma \mapsto \tilde{\sigma}|_X$ is a homomorphism of groups.

Lemma 4.3. (1) *Suppose that $A \in \mathbf{M}_{2n}(\Lambda(n, n))$. Then $\text{pr}_{[0]}(A)$ is invertible if and only if A is invertible.*

(2) *Suppose that $\{e_1, \dots, e_{2n}\}$ is a $\Lambda(n, n)$ -basis of $W(n, n)$. Let $\text{pr}_{[-1]}$ be the projection of $W(n, n)$ onto $W(n, n)_{[-1]}$. Then $\{\text{pr}_{[-1]} e_1, \dots, \text{pr}_{[-1]} e_{2n}\}$ is an \mathbb{F} -basis of $W(n, n)_{[-1]}$.*

(3) *Suppose that φ is an automorphism of X , and suppose that $\{y_i \mid i \in Y\} \subset X$ is a $\Lambda(n, n)$ -basis of $W(n, n)$. Then $\{\varphi(y_i) \mid i \in Y\}$ is also a $\Lambda(n, n)$ -basis of $W(n, n)$.*

(4) *The natural filtration of $\Lambda(n, n)$ is invariant under automorphisms of $\Lambda(n, n)$.*

Proof. (1) We first prove that A is invertible when $\text{pr}_{[0]} A$ is invertible. Set

$$\begin{aligned} \mathbf{P}(n)_1 &= \{f \in \mathbf{P}(n) \mid \text{pr}_{[0]}(f) = 0\} \quad \text{and} \\ T &= \text{span}_{\mathbb{F}}\{x^{(\alpha)} x^u \mid D_i(x^u) \neq 0 \text{ for some } i \in Y\}. \end{aligned}$$

Then we can write $A = \text{pr}_{[0]}(A) + B + C$, where $B \in \mathbf{M}_{2n}(\mathbf{P}(n)_1)$, $C \in \mathbf{M}_{2n}(T)$. Let $D = \text{pr}_{[0]}(A) + B$. Since $\mathbf{P}(n)$ is commutative, we see that $\det D$ is well defined. Note that $\text{pr}_{[0]}(\det D) = \det(\text{pr}_{[0]} D) \neq 0$, we can write $\det D = a + f$, where $0 \neq a \in \mathbb{F}$, $f \in \mathbf{P}(n)_1$. Put $g = a^{-1}(\sum_{i=0}^{\infty} (-1)^i (a^{-1} f)^i)$. A direct calculation shows that $g \det D = 1$. It follows that $\det D$ is invertible, whence D is invertible.

Let E be the inverse of D . Since $C \in \mathbf{M}_{2n}(T)$, we have $CE \in \mathbf{M}_{2n}(T)$, which combined with the fact that the product of any $n + 1$ elements of T is 0 yields that

CE is nilpotent. Thus $I + CE$ is invertible. Consequently, we have

$$AE(I + CE)^{-1} = (C + D)E(I + CE)^{-1} = (CE + DE)(I + CE)^{-1} = I.$$

Therefore A is invertible.

Using the fact that $\text{pr}_{[0]}(AB) = \text{pr}_{[0]}(A) \text{pr}_{[0]}(B)$ for arbitrary matrices A and B in $M_{2n}(\Lambda(n, n))$, we can prove the converse implication.

(2) Suppose that $(D_1, \dots, D_{2n})^T = A(e_1, \dots, e_{2n})^T$, where $A \in M_{2n}(\Lambda(n, n))$. Then $(D_1, \dots, D_{2n})^T = \text{pr}_{[0]} A(\text{pr}_{[-1]}(e_1), \dots, \text{pr}_{[-1]}(e_{2n}))^T$. Since $\{D_1, \dots, D_{2n}\}$ is an \mathbb{F} -basis of $W(n, n)_{[-1]}$, it follows that $\{\text{pr}_{[-1]}(e_1), \dots, \text{pr}_{[-1]}(e_{2n})\}$ is an \mathbb{F} -basis of $W(n, n)_{[-1]}$.

(3) By Theorem 3.17, φ induces canonically $\bar{\varphi} \in \text{gl}(X/X_0)$. Denote by \bar{y}_i the image of y_i under the canonically map $X \rightarrow X/X_0$. Then $\{\bar{y}_i \mid i \in Y\}$ is an \mathbb{F} -basis of X/X_0 . Assume that

$$(\varphi(y_1), \dots, \varphi(y_{2n}))^T = A(D_1, \dots, D_{2n})^T,$$

where $A \in M_{2n}(\Lambda(n, n))$. Decomposing $A = \text{pr}_{[0]} A + \text{pr}_1 A$, we obtain

$$(\bar{\varphi}(\bar{y}_1), \dots, \bar{\varphi}(\bar{y}_{2n}))^T = (\overline{\varphi(y_1)}, \dots, \overline{\varphi(y_{2n})})^T = \text{pr}_{[0]} A(\bar{D}_1, \dots, \bar{D}_{2n})^T.$$

This implies that $\text{pr}_{[0]} A$ is invertible. It follows from (1) that A is invertible. Therefore $\{\varphi(y) \mid y \in Y\}$ is a $\Lambda(n, n)$ -basis of $W(n, n)$.

(4) Since $\text{Der } \Lambda(n, n) = W(n, n)$, we have $\text{Aut } \Lambda(n, n) = \text{Aut}(\Lambda(n, n) : W(n, n))$. By [Zhang and Liu 2004, Theorem 2.12], the natural filtration of $W(n, n)$ is invariant under $\text{Aut } W(n, n)$. Note that for every $i \in Y$, $\tilde{\sigma}(f D_i) = (\sigma f)(\tilde{\sigma} D_i)$, where $\sigma \in \text{Aut } \Lambda(n, n)$ and $f \in \Lambda(n, n)$, which implies the desired result. \square

Theorem 4.4. *The map $\Phi : \text{Aut}(\Lambda(n, n) : X) \rightarrow \text{Aut } X$ given by $\sigma \mapsto \tilde{\sigma}|_X$ is an isomorphism.*

Proof. It suffices to show that Φ is bijective. Assume that $\sigma \in \text{Aut}(\Lambda(n, n) : X)$ is an element such that $\tilde{\sigma}|_X = 1|_X$. We first use induction on $|\alpha| + |u|$ to show that $\sigma(x^{(\alpha)}x^u) = x^{(\alpha)}x^u$, where $x^{(\alpha)}x^u$ is a standard basis element of $\Lambda(n, n)$, $\alpha \in \mathbb{N}_0^n$, $u \in \mathbb{B}(n)$. If $|\alpha| + |u| = 1$, then $x^{(\alpha)}x^u = x_i$ for some $i \in Y$. Since for every $k \in Y$

$$D_k(\sigma(x_i)) = (\tilde{\sigma}(D_k))(\sigma(x_i)) = \sigma D_k \sigma^{-1} \sigma(x_i) = \sigma D_k(x_i) = \sigma(\delta_{ik}) = \delta_{ik} = D_k(\delta_{ik}),$$

it follows that $D_k(\sigma(x_i) - x_i) = 0$, which combined with Lemma 4.3(4) yields $\sigma(x_i) = x_i$. If $|\alpha| + |u| > 1$, then by induction

$$D_k(\sigma(x^{(\alpha)}x^u) - x^{(\alpha)}x^u) = (\tilde{\sigma} D_k)\sigma(x^{(\alpha)}x^u) - D_k(x^{(\alpha)}x^u) = 0,$$

for every $k \in Y$. Thus $\sigma(x^{(\alpha)}x^u) - x^{(\alpha)}x^u \in \mathbb{F} \cap \Lambda(n, n)_1 = 0$. Consequently, $\sigma = 1$ and Φ is injective.

We next prove that Φ is surjective. Suppose that φ is in $\text{Aut } X$. Then by Lemma 4.3 (3), $\{\varphi(D_1), \dots, \varphi(D_{2n})\}$ is a $\Lambda(n, n)$ -basis of $W(n, n)$. Therefore, we can suppose that $\varphi(T_H(x_i x_j)) = \sum_{t=1}^{2n} h_{ijt} \varphi(D_t)$, where $i, j \in Y$ with $i \neq j'$, and $h_{ijt} \in \Lambda(n, n)$. Applying Lemma 4.3 (2), we have $h_{ijt} \in \Lambda(n, n)_1$. Thus

$$(13) \quad \varphi([D_k, T_H(x_i x_j)]) = \left[\varphi(D_k), \sum_{t=1}^{2n} h_{ijt} \varphi(D_t) \right] = \sum_{t=1}^{2n} (\varphi(D_k)(h_{ijt})) \varphi(D_t).$$

On the other hand,

$$(14) \quad \begin{aligned} \varphi([D_k, T_H(x_i x_j)]) &= \varphi[D_k, (-1)^{\mu(i)\mu(j')} x_j D_{i'} + (-1)^{\mu(j)} x_i D_{j'}] \\ &= (-1)^{\mu(i)\mu(j')} \delta_{kj} \varphi(D_{i'}) + (-1)^{\mu(j)} \delta_{ki} \varphi(D_{j'}). \end{aligned}$$

In particular, by letting $i = 1$ and $j \in Y \setminus \{1'\}$ in equations (13) and (14), one sees that $\varphi(D_k)(h_{1j1'}) = \delta_{kj} + \delta_{k1} \delta_{j1}$ for all $k \in Y$. Similarly, by letting $i = 2'$ and $j = 1'$, we obtain $\varphi(D_k)(h_{2'1'2}) = \delta_{k1'}$. Let $h_1 = \frac{1}{2} h_{111'}$, $h_{1'} = h_{2'1'2}$ and $h_j = h_{1j1'}$ for $j \in Y \setminus \{1, 1'\}$. Then $h_j \in \Lambda(n, n)_1$ with $\deg(h_j) = \mu(j)$, and

$$(15) \quad \varphi(D_i)(h_j) = \delta_{ij} \quad \text{for all } i, j \in Y.$$

Suppose that $\varphi(D_i) = \sum_{t=1}^{2n} f_{it} D_t$, where $f_{it} \in \Lambda(n, n)$. It follows from (15) that

$$(\delta_{ij})_{i,j \in Y} = (\varphi(D_i)h_j)_{i,j \in Y} = (f_{ij})_{i,j \in Y} (D_i h_j)_{i,j \in Y}.$$

This implies that $(D_i h_j)_{i,j \in Y}$ is invertible, whence $(\text{pr}_{[0]}(D_i h_j))_{i,j \in Y}$ is invertible by Lemma 4.3(1). Consequently, there exists $\sigma \in \text{Aut } \Lambda(n, n)$ such that $\sigma(x_i) = h_i$ by Lemma 3.4, which combined with (15) yields

$$(\tilde{\sigma} D_i - \varphi D_i)(h_j) = \sigma(D_i x_j) - \delta_{ij} = 0$$

for all $i, j \in Y$. Since h_1, h_2, \dots, h_{2n} generate $\Lambda(n, n)$, we see that $\tilde{\sigma} D_i = \varphi D_i$, whence $\tilde{\sigma}|_X = \varphi$ by Proposition 4.2. □

Acknowledgments

This paper was completed when Q. Mu visited the College of William and Mary, and he thanks the university and Professor Junping Shi for their warm hospitality.

References

[Humphreys 1972] J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics **9**, Springer, New York, 1972. MR 48 #2197 Zbl 0254.17004

[Jin 1992] N. Jin, “ad-nilpotent elements, quasi-nilpotent elements and invariant filtrations of infinite-dimensional Lie algebras of Cartan type”, *Sci. China Ser. A* **35**:10 (1992), 1191–1200. MR 94j:17027 Zbl 0772.17007

- [Kac 1977] V. G. Kac, “Lie superalgebras”, *Advances in Math.* **26**:1 (1977), 8–96. MR 58 #5803 Zbl 0366.17012
- [Kac 1998] V. G. Kac, “Classification of infinite-dimensional simple linearly compact Lie superalgebras”, *Adv. Math.* **139**:1 (1998), 1–55. MR 99m:17006 Zbl 0929.17026
- [Rudakov 1986] A. N. Rudakov, “Subalgebras and automorphisms of Lie algebras of Cartan type”, *Funktional. Anal. i Prilozhen.* **20**:1 (1986), 83–84. In Russian; translated in *Funct. Anal. Appl.* **20** (1986), 72–73. MR 87m:17030 Zbl 0594.17015
- [Scheunert 1979] M. Scheunert, *The theory of Lie superalgebras*, Lecture Notes in Mathematics **716**, Springer, Berlin, 1979. MR 80i:17005 Zbl 0407.17001
- [Strade and Farnsteiner 1988] H. Strade and R. Farnsteiner, *Modular Lie algebras and their representations*, Monographs and Textbooks in Pure and Applied Mathematics **116**, Marcel Dekker, New York, 1988. MR 89h:17021 Zbl 0648.17003
- [Wilson 1971] R. L. Wilson, “Classification of generalized Witt algebras over algebraically closed fields”, *Trans. Amer. Math. Soc.* **153** (1971), 191–210. MR 47 #5070 Zbl 0218.17009
- [Wilson 1975] R. L. Wilson, “Automorphisms of graded Lie algebras of Cartan type”, *Comm. Algebra* **3**:7 (1975), 591–613. MR 52 #3259 Zbl 0318.17009
- [Zhang 1997] Y. Zhang, “Finite-dimensional Lie superalgebras of Cartan type over fields of prime characteristic”, *Chinese Sci. Bull.* **42**:9 (1997), 720–724. MR 98h:17023 Zbl 0886.17022
- [Zhang and Liu 2004] Y. Zhang and W. Liu, “The general and special superalgebras of formal vectorfields”, *Sci. China Ser. A* **47**:2 (2004), 272–283. MR 2005g:17040 Zbl 1124.17304

Received September 22, 2011. Revised October 27, 2011.

LI REN
SCHOOL OF MATHEMATICS AND STATISTICS
NORTHEAST NORMAL UNIVERSITY
CHANGCHUN, 130024
CHINA
ren1944@nenu.edu.cn

QIANG MU
SCHOOL OF MATHEMATICAL SCIENCES
HARBIN NORMAL UNIVERSITY
HARBIN, 150025
CHINA
qmu520@gmail.com

YONGZHENG ZHANG
SCHOOL OF MATHEMATICS AND STATISTICS
NORTHEAST NORMAL UNIVERSITY
CHANGCHUN, 130024
CHINA
zhyz@nenu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 257 No. 2 June 2012

Extending triangulations of the 2-sphere to the 3-disk preserving a 4-coloring	257
RUI PEDRO CARPENTIER	
Orthogonal quantum group invariants of links	267
LIN CHEN and QINGTAO CHEN	
Some properties of squeezing functions on bounded domains	319
FUSHENG DENG, QIAN GUAN and LIYOU ZHANG	
Representations of little q -Schur algebras	343
JIE DU, QIANG FU and JIAN-PAN WANG	
Renormalized weighted volume and conformal fractional Laplacians	379
MARÍA DEL MAR GONZÁLEZ	
The L_4 norm of Littlewood polynomials derived from the Jacobi symbol	395
JONATHAN JEDWAB and KAI-UWE SCHMIDT	
On a conjecture of Kaneko and Ohno	419
ZHONG-HUA LI	
Categories of unitary representations of Banach–Lie supergroups and restriction functors	431
STÉPHANE MERIGON, KARL-HERMANN NEEB and HADI SALMASIAN	
Odd Hamiltonian superalgebras and special odd Hamiltonian superalgebras of formal vector fields	471
LI REN, QIANG MU and YONGZHENG ZHANG	
Interior derivative estimates for the Kähler–Ricci flow	491
MORGAN SHERMAN and BEN WEINKOVE	
Two-dimensional disjoint minimal graphs	503
LINFENG ZHOU	



0030-8730(201206)257:2;1-5