

*Pacific  
Journal of  
Mathematics*

**REMARKS ON THE BEHAVIOR OF NONPARAMETRIC  
CAPILLARY SURFACES AT CORNERS**

KIRK E. LANCASTER

Volume 258    No. 2

August 2012

## REMARKS ON THE BEHAVIOR OF NONPARAMETRIC CAPILLARY SURFACES AT CORNERS

KIRK E. LANCASTER

**Consider a nonparametric capillary or prescribed mean curvature surface  $z = f(x)$  defined in a cylinder  $\Omega \times \mathbb{R}$  over a two-dimensional region  $\Omega$  whose boundary has a corner at  $\mathbb{O}$  with an opening angle of  $2\alpha$ . Suppose the contact angle approaches limiting values  $\gamma_1$  and  $\gamma_2$  in  $(0, \pi)$  as  $\mathbb{O}$  is approached along each side of the opening angle. We will prove the nonconvex Concus–Finn conjecture, determine the exact sizes of the radial limit fans of  $f$  at  $\mathbb{O}$  when  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$  and discuss the continuity of the Gauss map.**

### 1. Introduction

Let  $\Omega \subset \mathbb{R}^2$  be a connected, open set. Consider the classical capillary problem in a cylinder

$$(1) \quad Nf = \kappa f + \lambda \quad \text{in } \Omega,$$

$$(2) \quad Tf \cdot \nu = \cos \gamma \quad (\text{a.e.}) \quad \text{on } \partial\Omega,$$

and, more generally, the prescribed mean curvature problem in a cylinder

$$(3) \quad Nf = H(\cdot, f(\cdot)) \quad \text{in } \Omega,$$

$$(4) \quad Tf \cdot \nu = \cos \gamma \quad (\text{a.e.}) \quad \text{on } \partial\Omega,$$

where  $Tf = \nabla f / \sqrt{1 + |\nabla f|^2}$ ,  $Nf = \nabla \cdot Tf$ ,  $\nu$  is the exterior unit normal on  $\partial\Omega$ ,  $H(x, t)$  is a weakly increasing function of  $t$  for each  $x \in \Omega$  and  $\gamma = \gamma(x)$  is in  $[0, \pi]$ . We will let  $\mathcal{S}_f$  denote the closure in  $\mathbb{R}^3$  of the graph of  $f$  over  $\Omega$ . When  $H(x, t) = \kappa t + \lambda$  (i.e.,  $f$  satisfies (1)–(2)) with  $\kappa$  and  $\lambda$  constants such that  $\kappa \geq 0$ , then the surface  $\mathcal{S}_f \cap (\Omega \times \mathbb{R})$  represents the stationary liquid-gas interface formed by an incompressible fluid in a vertical cylindrical tube with cross-section  $\Omega$  in a microgravity environment or in a downward-oriented gravitational field, the subgraph  $U = \{(x, t) \in \Omega \times \mathbb{R} : t < f(x)\}$  represents the fluid filled portion of the cylinder and  $\gamma(x)$  is the angle (within the fluid) at which the liquid-gas interface meets the vertical cylinder at  $(x, f(x))$ ; Paul Concus and Robert Finn have made

---

*MSC2010:* primary 76B45, 35J93; secondary 53A10, 35J62.

*Keywords:* capillary graph, Concus–Finn conjecture, Gauss map, sizes of fans.

fundamental contributions to the mathematical theory of capillary surfaces and have discovered that capillary surfaces can behave in unexpected ways (cf. [Concus and Finn 1996; Finn 1986; 1999; 2002b; 2002a]). For a function  $f \in C^2(\Omega)$ , we let


$$\vec{n}(X) = \vec{n}_f(X) = \frac{(\nabla f(x), -1)}{\sqrt{1+|\nabla f(x)|^2}}, \quad X = (x, t) \in \Omega,$$

denote the downward unit normal to the graph of  $f$ ; when  $f$  is a solution of (1)–(2) and  $\kappa \geq 0$ ,  $\vec{n}$  represents the inward unit normal with respect to the fluid region. Of interest here is the behavior of capillary surfaces and prescribed mean curvature surfaces over domains  $\Omega \subset \mathbb{R}^2$  whose boundaries contain corners (e.g., [Concus and Finn 1996; Finn 1996]).


Let us suppose  $\mathbb{O} = (0, 0) \in \partial\Omega$  and  $\Omega$  is a connected, simply connected open set in  $\mathbb{R}^2$  such that  $\partial\Omega \setminus \{\mathbb{O}\}$  is a piecewise  $C^1$  curve,  $\Omega$  has a corner of size  $2\alpha$  at  $\mathbb{O}$  and the tangent cone to  $\partial\Omega$  at  $\mathbb{O}$  is  $L^+ \cup L^-$ , where polar coordinates relative to  $\mathbb{O}$  are denoted by  $r$  and  $\theta$ ,  $L^+ = \{\theta = \alpha\}$  and  $L^- = \{\theta = -\alpha\}$ . We will assume there exist  $\delta^* > 0$ ,  $\rho^* \in (0, 1]$  such that  $\partial^+\Omega = \partial\Omega \cap \overline{B(\mathbb{O}, \delta^*)} \cap T^+$  and  $\partial^-\Omega = \partial\Omega \cap \overline{B(\mathbb{O}, \delta^*)} \cap T^-$  are connected,  $C^{1,\rho^*}$  arcs, where  $T^+ = \{x \in \mathbb{R}^2 : x_2 \geq 0\}$ ,  $T^- = \{x \in \mathbb{R}^2 : x_2 \leq 0\}$  and  $B(\mathbb{O}, \epsilon) = \{x \in \mathbb{R}^2 : |x| < \epsilon\}$ ; hence the tangent rays to  $\partial^+\Omega$  and  $\partial^-\Omega$  at  $\mathbb{O}$  are  $L^+$  and  $L^-$  respectively. Set  $\Omega_0 = \Omega_0(\delta^*) = \Omega \cap B(\mathbb{O}, \delta^*)$ . Let  $\gamma^+(s)$  and  $\gamma^-(s)$  denote  $\gamma$  along the arcs  $\partial^+\Omega_0$  and  $\partial^-\Omega_0$ , respectively, where  $s = 0$  corresponds to the point  $\mathbb{O}$ ; here we have parametrized  $\partial^+\Omega_0$  and  $\partial^-\Omega_0$  by, for example, arclength  $s$  from  $\mathbb{O}$  and write these parametrizations as  $x^+$  and  $x^-$  respectively. We will assume there exist  $\gamma_1, \gamma_2 \in (0, \pi)$  such that

$$(5) \quad \lim_{\partial^+\Omega \ni x \rightarrow \mathbb{O}} \gamma(x) = \gamma_1 \quad \text{and} \quad \lim_{\partial^-\Omega \ni x \rightarrow \mathbb{O}} \gamma(x) = \gamma_2.$$

Suppose first that  $2\alpha \leq \pi$  (i.e., the corner is convex or  $\partial\Omega$  is  $C^1$  at  $\mathbb{O}$ ); such an  $\Omega$  is illustrated in Figure 1. Figure 2 can then be used to illustrate our knowledge



**Figure 1.**  $\Omega$  with  $2\alpha < \pi$ .



**Figure 2.** The Concus–Finn rectangle for convex corners.


of the behavior of a solution  $f$  of (3)–(4) at the corner  $\mathbb{O}$ ; here let  $R, D_1^\pm, D_2^\pm$  be the indicated open regions in the (open) square  $(0, \pi) \times (0, \pi)$ . If  $(\gamma_1, \gamma_2)$  is in  $\bar{R} \cap (0, \pi) \times (0, \pi)$ , then  $f$  is continuous at  $\mathbb{O}$  [Concus and Finn 1996, Theorem 1; Lancaster and Siegel 1996b; 1996a, Corollary 4; Tam 1986]. If  $(\gamma_1, \gamma_2) \in D_1^\pm$ , then  $f$  is unbounded in any neighborhood of  $\mathbb{O}$  and the capillary problem has no solution if  $\kappa = 0$  [Concus and Finn 1996; Finn 1996]. If  $(\gamma_1, \gamma_2) \in D_2^\pm$ , then  $f$  is bounded [Lancaster and Siegel 1996a, Proposition 1] but its continuity at  $\mathbb{O}$  was unknown until recently. Concus and Finn discovered bounded solutions of (1)–(2) in domains with corners whose unit normals (i.e., Gauss maps) cannot extend continuously as functions of  $x$  to a corner on the boundary of the domain [Finn 1988a, page 15; 1988b; 1996; Concus and Finn 1996, Example 2]. They formulated the conjecture that the solution  $f$  of (1)–(2) must be discontinuous at  $\mathbb{O}$  when  $(\gamma_1, \gamma_2) \in D_2^\pm$ . Writing the conditions required for a pair of angles to be in  $D_2^\pm$  yields the following formulation of their conjecture:

**Concus–Finn conjecture.** *Suppose  $0 < \alpha < \pi/2$ , that the limits (5) exist and  $0 < \gamma_1, \gamma_2 < \pi$ . If  $2\alpha + |\gamma_1 - \gamma_2| > \pi$ , then any solution of (3)–(4), with  $H(x, z) = \kappa z + \lambda$ ,  $\kappa \geq 0$ , has a jump discontinuity at  $\mathbb{O}$ .*

This conjecture was proven for solutions of (3)–(4) (i.e., without the restriction that  $H(x, z) = \kappa z + \lambda$ ) in [Lancaster 2010].<sup>1</sup>

Thus, when  $2\alpha \leq \pi$ ,  $(\gamma_1, \gamma_2) \in D_2^\pm$ , and  $f$  satisfies (3)–(4),  $f$  is discontinuous at  $\mathbb{O}$  and there is a countable set  $\mathcal{F} \subset (-\alpha, \alpha)$  such that the radial limit function of  $f$  at

<sup>1</sup>For convenience, we will abbreviate this reference as [L]. Similarly, [Lancaster and Siegel 1996a] and [Lancaster and Siegel 1996b] will be abbreviated [LS a] and [LS b], respectively.



**Figure 3.**  $\Omega$  with  $2\alpha > \pi$ .

$\mathbb{C}$ ,  $Rf$ , defined by  $Rf(\alpha) = \lim_{\partial^+\Omega \ni x \rightarrow \mathbb{C}} f(x)$ ,  $Rf(-\alpha) = \lim_{\partial^-\Omega \ni x \rightarrow \mathbb{C}} f(x)$  and

$$(6) \quad Rf(\theta) = \lim_{r \downarrow 0} f(r \cos \theta, r \sin \theta), \quad -\alpha < \theta < \alpha,$$

is well-defined and continuous on  $[-\alpha, \alpha] \setminus \mathcal{F}$  and behaves as in Proposition 1(i) of [LS b]; if  $H(x, z)$  is strictly increasing in  $z$  [LS a, §5] or real-analytic [LS b] for  $x$  in a neighborhood of  $\mathbb{C}$ , then  $\mathcal{F} = \emptyset$ . (See [LS a], Step 3 of the proof of Theorem 1 and §5, and [LS b] regarding the sets  $\mathcal{F}$  and cusp solutions.) We may assume for the moment that  $(\gamma_1, \gamma_2) \in D_2^+$  since the other case follows by interchanging  $x_1$  and  $x_2$ ; then Theorems 1 and 2 of [LS a] and Proposition 1 and Theorem 1 of [LS b] imply there is a countable set  $\mathcal{F} \subset [\alpha_1, \alpha_2]$  such that

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly increasing} & \text{on } [\alpha_1, \alpha_2] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, \alpha_1], \end{cases}$$

with  $\alpha_1 < \alpha_2$ ,  $\alpha - \alpha_2 \geq \gamma_1$  and  $\alpha_1 - (-\alpha) \geq \pi - \gamma_2$ . In fact, determining the exact sizes of these radial limit fans when  $f$  is discontinuous at  $\mathbb{C}$  follows easily from [L]. (Notice that  $D_1^\pm = \emptyset$  if  $2\alpha = \pi$ .)

**Proposition 1.1.** *Let  $\Omega$  be as above with  $2\alpha < \pi$  and  $f$  be a bounded solution to (3)–(4). Suppose that  $(\gamma_1, \gamma_2) \in D_2^\pm$  and that there exist constants  $\underline{\gamma}^\pm, \bar{\gamma}^\pm$ ,  $0 < \underline{\gamma}^\pm \leq \bar{\gamma}^\pm < \pi$ , satisfying*

$$\underline{\gamma}^+ + \underline{\gamma}^- > \pi - 2\alpha \quad \text{and} \quad \bar{\gamma}^+ + \bar{\gamma}^- < 2\alpha + \pi,$$

so that  $\underline{\gamma}^\pm \leq \gamma^\pm(s) \leq \bar{\gamma}^\pm$  for all  $s$ ,  $0 < s < s_0$ , for some  $s_0$ . Then  $Rf(\theta)$  exists for  $\theta \in [-\alpha, \alpha] \setminus \mathcal{F}$  and  $Rf(\theta)$  is a continuous function of  $\theta \in [-\alpha, \alpha] \setminus \mathcal{F}$ , where  $\mathcal{F}$  is a countable subset of  $(-\alpha, \alpha)$ .

**Case (I).** If  $(\gamma_1, \gamma_2) \in D_2^+$  (i.e.,  $\gamma_1 - \gamma_2 < 2\alpha - \pi$ ) then  $\alpha_1 = -\alpha + \pi - \gamma_2, \alpha_2 = \alpha - \gamma_1$  and

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha - \gamma_1, \alpha], \\ \text{strictly increasing} & \text{on } [-\alpha + (\pi - \gamma_2), \alpha - \gamma_1] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, -\alpha + (\pi - \gamma_2)], \end{cases}$$

where  $\mathcal{F}$  is a countable subset of  $[-\alpha + (\pi - \gamma_2), \alpha - \gamma_1]$ .

**Case (D).** If  $(\gamma_1, \gamma_2) \in D_2^-$  (i.e.,  $\gamma_1 - \gamma_2 > \pi - 2\alpha$ ) then  $\alpha_1 = -\alpha + \gamma_2, \alpha_2 = \alpha - \pi + \gamma_1$  and

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha - (\pi - \gamma_1), \alpha], \\ \text{strictly decreasing} & \text{on } [-\alpha + \gamma_2, \alpha - (\pi - \gamma_1)] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, -\alpha + \gamma_2], \end{cases}$$

where  $\mathcal{F}$  is a countable subset of  $[-\alpha + \gamma_2, \alpha - \pi + \gamma_1]$ .

*Proof.* Using the information from [LS b] and [LS a] given above and assuming  $(\gamma_1, \gamma_2) \in D_2^+$ , we will argue by contradiction. Suppose that  $\alpha_2 < \alpha - \gamma_1$ . Let

$$\Omega_0 \subset \{(r \cos \theta, r \sin \theta) \in \Omega : r > 0, \alpha_2 < \theta < \alpha - \gamma_1/2\}.$$

be an open set whose boundary  $\partial\Omega_0$  contains  $\{\theta = \alpha - \gamma_1/2\}$  and is tangent to  $\{\theta = \alpha_2\}$  at  $\mathbb{O}$  so that the appropriate analogue of [L, (43)] tends to zero. Then  $f$  is continuous on  $\overline{\Omega}$  and, from Theorem 2.1 of [L], we obtain

$$(7) \quad \lim_{r \downarrow 0} \vec{n}_f(r \cos(\alpha - \frac{1}{2}\gamma_1), r \sin(\alpha - \frac{1}{2}\gamma_1)) = (-\sin(\alpha - \gamma_1), \cos(\alpha - \gamma_1), 0),$$

$$(8) \quad \lim_{\substack{x \rightarrow \mathbb{O} \\ x \in \partial\Omega_0 \setminus \{\theta = \alpha - \frac{1}{2}\gamma_1\}}} \vec{n}_f(x) = (-\sin \alpha_2, \cos \alpha_2, 0).$$

Notice that the limiting contact angles at  $\mathbb{O}$  are  $\frac{1}{2}\gamma_1$  (on  $\theta = \alpha - \frac{1}{2}\gamma_1$ ) and  $\pi$  (on  $\theta = \alpha_2$ ). Now, using Theorem 2.1 of [L], we see that the arguments in §3 of [L] yield a contradiction to the assumption that  $\alpha_2 < \alpha - \gamma_1$ . (If  $\gamma_2 = \pi$  were allowed in Theorem 1.1 of [L], then a contradiction would follow immediately since  $2\alpha = \alpha - \alpha_2 - \frac{1}{2}\gamma_1, |\gamma_1 - \gamma_2| = \pi - \frac{1}{2}\gamma_1$  and  $2\alpha + |\gamma_1 - \gamma_2| = \pi + \alpha - \gamma_1 - \alpha_2 > \pi$ .) In the case that  $\alpha_1 > -\alpha + \pi - \gamma_2$  or  $(\gamma_1, \gamma_2) \in D_2^-$ , the proof follows in a similar manner. □

The focus of this note is to give a direct proof of the nonconvex Concus–Finn conjecture and, when  $(\gamma_1, \gamma_1) \in D_1^\pm \cup D_2^\pm$ , establish the exact sizes of radial limit fans at reentrant corners and discuss the continuity of the Gauss map. We note that Danzhu Shi assumes the (convex) Concus–Finn conjecture holds when  $\gamma_1 \in \{0, \pi\}$  or  $\gamma_2 \in \{0, \pi\}$  and then, in her extremely interesting paper [Shi 2006], gives an argument for the proof of the nonconvex Concus–Finn conjecture. Unfortunately,

these cases (e.g.,  $\gamma_j \in \{0, \pi\}$ ,  $j = 1, 2$ ) are not covered in [L]. Our interest in proving the nonconvex Concus–Finn conjecture arises from our need, when determining the exact sizes of fans at reentrant corners, for the information developed during its proof (e.g., analogs of Theorem 2.1 of [L]) and from a belief in the value of presenting a proof which directly uses the ideas and techniques in [L].

### 2. The nonconvex Concus–Finn conjecture

The following theorem implies that the nonconvex Concus–Finn conjecture (cf. [Shi 2006]) is true; the proof will be given in Section 2B.

**Theorem 2.1.** *Let  $\Omega$  and  $\gamma$  be as above with  $\alpha \in [\frac{\pi}{2}, \pi]$ . Let*

$$f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{O}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{O}\})$$

*be a bounded solution of (3)–(4) with*

$$|H|_\infty = \sup_{x \in \Omega} |H(x, f(x))| < \infty$$

*for some  $\delta > 0$  and  $\rho \in (0, 1)$ . Suppose (5) holds and  $\gamma_1, \gamma_2 \in (0, \pi)$ . Then  $f$  is discontinuous at  $\mathbb{O}$  whenever  $|\gamma_1 - \gamma_2| > 2\alpha - \pi$  or  $|\gamma_1 + \gamma_2 - \pi| > 2\pi - 2\alpha$  (i.e.,  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ ).*

Throughout this section, we will consider  $f$  to be a fixed solution of (3)–(4) that satisfies the hypotheses of this theorem. We may parametrize the graph of  $f$  as in [LS a], using the unit disk  $E = \{(u, v) : u^2 + v^2 < 1\}$  as our parameter domain. From Step 1 of the proof of Theorem 1 of [LS a] and §3 of [L], we see that there is a parametric description  $X : \overline{E} \rightarrow \mathbb{R}^3$  of the closure  $S$  of  $S_0 = \{(x, f(x)) : x \in \Omega\}$ ,


$$X(u, v) = (x(u, v), y(u, v), z(u, v)), \quad (u, v) \in \overline{E},$$

such that:

- (i)  $X \in C^2(E : \mathbb{R}^3) \cap W^{1,2}(E : \mathbb{R}^3)$ .
- (ii)  $X$  is a homeomorphism of  $E$  onto  $S_0$ .
- (iii)  $X$  maps  $\partial E$  onto  $\{(x, f(x)) : x \in \partial\Omega\} \cup (\{\mathbb{O}\} \times [z_1, z_2])$ , where

$$z_1 = \liminf_{\overline{\Omega} \ni x \rightarrow \mathbb{O}} f(x) \quad \text{and} \quad z_2 = \limsup_{\overline{\Omega} \ni x \rightarrow \mathbb{O}} f(x).$$

- (iv)  $X$  is conformal on  $E$ :  $X_u \cdot X_v = 0$ ,  $|X_u| = |X_v|$  on  $E$ .
- (v) Let  $\tilde{H}(u, v) = H(X(u, v))$  denote the prescribed mean curvature of  $\mathcal{S}_f$  at  $X(u, v)$ . Then  $\Delta X := X_{uu} + X_{vv} = \tilde{H} X_u \times X_v$ .
- (vi)  $X \in C^0(\overline{E})$ .



**Figure 4.** The Concus-Finn rectangle for nonconvex corners.

(vii) Writing  $G(u, v) = (x(u, v), y(u, v))$ ,  $G(\cos t, \sin t)$  moves clockwise about  $\partial\Omega$  as  $t$  increases,  $0 \leq t \leq 2\pi$ , and  $G$  is an orientation reversing homeomorphism from  $E$  onto  $\Omega$ ;  $G$  maps  $\bar{E}$  onto  $\bar{\Omega}$  and, if  $f$  is continuous at  $\mathbb{O}$ , then  $G$  is a homeomorphism from  $\bar{E}$  onto  $\bar{\Omega}$ .

(viii) Let  $\pi_S : S^2 \rightarrow \mathbb{C}$  denote the stereographic projection from the North Pole and define  $g(u + iv) = \pi_S(\vec{n}_f(G(u, v)))$ ,  $(u, v) \in E$ . Then

$$(9) \quad |g_{\bar{\zeta}}| = \frac{1}{2} |\tilde{H}| (1 + |g|^2) |X_u|,$$

where  $\zeta = u + iv$ ,  $\partial/\partial\zeta = \frac{1}{2}(\partial/\partial u - i\partial/\partial v)$  and  $\partial/\partial\bar{\zeta} = \frac{1}{2}(\partial/\partial u + i\partial/\partial v)$ . For convenience when working with complex variables, set  $E_1 = \{\zeta \in \mathbb{C} : |\zeta| < 1\}$ .

(ix) The parametric Gauss map  $N : E \rightarrow S^2$  is  $N = (X_u \times X_v)/|X_u \times X_v|$  and satisfies  $N(u, v) = \vec{n}_f(G(u, v))$ ,  $(u, v) \in E$ ; the domain of  $N$  is taken as the largest subset of  $\bar{E}$  on which  $N$  extends continuously.

It is convenient to introduce some notation. Suppose  $V \subset \mathbb{R}^2$  with  $\mathbb{O} \in \partial V$ . For  $t > 0$ , set  $V_t = \{(x, y) \in V : x^2 + y^2 < t^2\}$ . Let  $s(V)$  denote the set of sequences in  $V$  that converge to  $\mathbb{O}$ . If  $h \in C^1(V)$ , we define  $\Pi_h(V) = \bigcap_{t>0} \overline{\vec{n}_h(V_t)}$ ; then

$$\Pi_h(V) = \left\{ Y \in S^2 : \text{there exists } (x_j) \in s(V) \text{ such that } Y = \lim_{j \rightarrow \infty} \vec{n}_h(x_j) \right\}.$$

Without assuming that  $f$  is or is not continuous at  $\mathbb{O}$ , we have:

**Lemma 2.2.** *Let  $\Lambda$  be an open, connected, simply connected subset of  $\Omega$  with  $\mathbb{O}$  in  $\partial\Lambda$  and suppose that there is a rotation  $M$  of  $\mathbb{R}^2$  about  $\mathbb{O}$  such that*

$$\{(M(y_1, y_2), y_3) : Y \in \Pi_f(\Lambda)\}$$



is contained in a compact subset of  $\{Y \in S^2 : y_2 > 0, y_3 \leq 0\}$ . Let  $\phi$  be a conformal map from  $E$  to  $G^{-1}(\Lambda)$  and define

$$(10) \quad \tilde{g}(u + iv) = \pi_S(\vec{n}_f(G \circ \phi(u, v))), \quad (u, v) \in E.$$

Then there exists  $p > 2$  such that

$$(11) \quad \tilde{g}(\zeta) = \psi(\zeta) + h(\zeta), \quad \zeta \in E_1,$$

where  $\psi$  is a holomorphic function on  $E_1$  and  $h \in L^\infty(E_1)$  is a Hölder continuous function on  $\bar{E}_1$  with Hölder exponent  $\mu = (p - 2)/p$ .

*Proof.* In §3 of [L], the fact that the limits at  $\mathbb{O}$  of the Gauss map are contained in a compact subset of  $\{Y \in S^2 : y_2 > 0, y_3 \leq 0\}$  implies that  $(u, v) \mapsto (z(u, v), x(u, v))$  is quasiconformal and has a quasiconformal extension to  $\mathbb{R}^2$ ; Gehring’s lemma and the isothermal parametrization imply  $X \in W^{1,p}$  for some  $p > 2$  and the classical literature implies  $g = \psi + h$  with  $\psi$  and  $h$  as above. We can argue as in §3 of [L]; we find that  $X \in W^{1,p}(E : \mathbb{R}^3)$  for some  $p > 2$  and

$$(12) \quad \tilde{g}(\zeta) = \psi(\zeta) + h(\zeta),$$

where  $\psi$  is a holomorphic function and  $h \in L^\infty(E_1)$  is an uniformly Hölder continuous function on  $E_1$  with Hölder exponent  $\mu$ . □

**Remark 2.3.** Notice that  $\tilde{g} = g \circ \phi_1$ , where  $\phi_1$  is a conformal map from  $E_1$  onto  $\{u + iv : (u, v) \in G^{-1}(\Lambda)\}$ .

**2A. Image of the Gauss map.** The (nonparametric) Gauss map on  $\mathcal{S}_f$  is the (downward) unit normal map to  $\mathcal{S}_f$  when this is defined and equals  $\vec{n}_f$  on  $\mathcal{S}_f \cap (\Omega \times \mathbb{R})$ ; here we consider  $\vec{n}_f : \Omega \times \mathbb{R} \rightarrow S^2_-$  by letting  $(x, t) \mapsto \vec{n}_f(x)$ . In this section, we characterize in Theorems 2.4 and 2.5 the behavior of the limits at points of  $\{\mathbb{O}\} \times \mathbb{R}$  of the Gauss map for the graph of  $f$  when  $(\gamma_1, \gamma_2) \notin \bar{R}$ . Let  $S^2_- = \{\omega \in \mathbb{R}^3 : |\omega| = 1, \omega_3 \leq 0\}$  be the (closed) lower half of the unit sphere.

**Theorem 2.4.** Let  $2\alpha > \pi$  and  $\Omega$  and  $\gamma$  be as in Section 1 and suppose (5) holds with  $\gamma_1, \gamma_2 \in (0, \pi)$ . Let  $\beta \in (-\alpha, \alpha)$  and  $(x_j) \in s(\Omega)$  such that

$$(13) \quad \lim_{j \rightarrow \infty} \frac{x_j}{|x_j|} = (\cos \beta, \sin \beta).$$

Let us write  $\omega(\theta) = (\cos \theta, \sin \theta, 0)$  for  $\theta \in \mathbb{R}$ .

$(D_2^+)$  If  $(\gamma_1, \gamma_2) \in D_2^+$  (i.e.,  $\gamma_1 - \gamma_2 < \pi - 2\alpha$ ) then

$$\lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \begin{cases} \omega(\alpha - \gamma_1 + \frac{\pi}{2}) & \text{if } \beta \in [\alpha - \gamma_1, \alpha), \\ \omega(\beta + \frac{\pi}{2}) & \text{if } \beta \in [-\alpha + (\pi - \gamma_2), \alpha - \gamma_1], \\ \omega(-\alpha - \gamma_2 + \frac{3\pi}{2}) & \text{if } \beta \in (-\alpha, -\alpha + (\pi - \gamma_2)]. \end{cases}$$

$(D_2^-)$  If  $(\gamma_1, \gamma_2) \in D_2^-$  (i.e.,  $\gamma_1 - \gamma_2 > 2\alpha - \pi$ ) then

$$\lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \begin{cases} \omega(\alpha + \gamma_1 - \frac{3\pi}{2}) & \text{if } \beta \in [\alpha + \gamma_1 - \pi, \alpha), \\ \omega(\beta - \frac{\pi}{2}) & \text{if } \beta \in [-\alpha + \gamma_2, \alpha + \gamma_1 - \pi], \\ \omega(-\alpha + \gamma_2 - \frac{\pi}{2}) & \text{if } \beta \in (-\alpha, -\alpha + \gamma_2]. \end{cases}$$

*Proof.* Let us assume  $(\gamma_1, \gamma_2) \in D_2^\pm$ . Let  $\beta \in (-\alpha, \alpha)$  and  $(x_j)$  be an arbitrary sequence in  $\Omega$  converging to  $\mathbb{O}$  and satisfying (13). Since  $(\vec{n}_f(x_j) : j \in \mathbb{N})$  is a sequence in the compact set  $S_-^2$ , there is a subsequence of  $(x_j : j \in \mathbb{N})$ , still denoted  $(x_j : j \in \mathbb{N})$ , and  $\theta \in (-\frac{\pi}{2}, \frac{3\pi}{2}]$ ,  $\tau \in [0, 1]$  such that  $(\vec{n}_f(x_{j_k}) : k \in \mathbb{N})$  is convergent and

$$\lim_{k \rightarrow \infty} \vec{n}_f(x_{j_k}) = (\tau \cos \theta, \tau \sin \theta, -\sqrt{1 - \tau^2}).$$

Using [Jeffres and Lancaster 2008] and the techniques and arguments in §2 of [L], we see that  $\tau = 1$ ,  $\lim_{k \rightarrow \infty} \vec{n}_f(x_{j_k}) = \omega(\theta)$ , and  $\omega(\theta)$  is normal to  $\partial\mathcal{P}$  and points into  $\mathcal{P}$ , where  $\omega(\beta) \in \partial\mathcal{P}$  and  $\mathcal{P}$  is given in Theorem 2.2 of [Jeffres and Lancaster 2008]. (In §2 of [L], the function  $u(x) = f(x) - Rf(\beta)$  is blown up about  $(0, 0, 0)$ ; that is, the graphs of a subsequence of the sequence  $(u_j)$  in  $C^2(\Omega)$ , where  $u_j$  is defined by  $u_j(x) = (f(\epsilon_j x) - Rf(\beta))/\epsilon_j$  and  $\epsilon_j = |x_j|$  for  $j \in \mathbb{N}$ , are shown to converge to the intersection of  $\Omega \times \mathbb{R}$  with a vertical plane  $\pi_1$ . The (downward) unit normal to  $\pi_1$  is shown to be normal to the vertical plane  $\partial\mathcal{P}$  which contains  $(\cos \beta, \sin \beta, 0)$  and point into  $\mathcal{P}$ , where  $\mathcal{P}$  satisfies Theorem 2.1 of [Jeffres and Lancaster 2007].)

If  $(\gamma_1, \gamma_2) \in D_2^+$ , then the conclusions of Theorem 2.4 follow from Corollary 2.4 of [Jeffres and Lancaster 2008]; Figure 5 illustrates the graph of the argument of  $\vec{n}(\beta) = \lim_{r \downarrow 0} \vec{n}_f(r \cos \beta, r \sin \beta)$ . If  $(\gamma_1, \gamma_2) \in D_2^-$ , then the conclusions of Theorem 2.4 follow from Corollary 2.5 of [ibid.].  $\square$

Suppose that  $\alpha \in (\frac{\pi}{2}, \pi]$ ,  $\gamma_1, \gamma_2 \in (0, \pi)$  and  $\gamma_1 + \gamma_2 < 2\alpha - \pi$ . Let us define  $\mathcal{F} = \mathcal{F}(\alpha, \gamma_1, \gamma_2)$  as follows: Set

$$\mathcal{F}_1 = [-\alpha, -\alpha - \gamma_2 + \pi] \times \{-\alpha - \gamma_2 - \pi/2\},$$

$$\mathcal{F}_2 = [-\alpha, -\alpha + \gamma_2 + \pi] \times \{-\alpha + \gamma_2 - \pi/2\},$$

$$\mathcal{F}_3 = [\alpha - \gamma_1 - \pi, \alpha] \times \{\alpha - \gamma_1 - 3\pi/2\},$$


$$\mathcal{F}_4 = [\alpha + \gamma_1 - \pi, \alpha] \times \{\alpha + \gamma_1 - 3\pi/2\},$$

$$\mathcal{F}_5 = \{(\beta, \beta - \pi/2) : \beta \in [-\alpha + \gamma_2, \alpha + \gamma_1 - \pi]\},$$


$$\mathcal{F}_6 = \{(\beta, \beta - 3\pi/2) : \beta \in [-\alpha + \gamma_2 + \pi, \alpha - \gamma_1]\},$$

$$\mathcal{F}_7 = \left\{ (\beta + t, -\alpha + \gamma_2 - \pi/2 + t) : \begin{array}{l} \beta \in [-\alpha + \gamma_2, -\alpha + \gamma_2 + \pi] \\ t \in [0, 2\alpha - \pi - \gamma_1 - \gamma_2] \end{array} \right\},$$


and define  $\mathcal{F} = \bigcup_{j=1}^7 \mathcal{F}_j = \mathcal{F}_1 \cup \dots \cup \mathcal{F}_7$  (see Figures 6, 7 and 8 for illustrations).




**Figure 5.**  $\{(\beta, \arg(\bar{m}_f(\beta)))\}$  at a  $D_2^+$  corner;  $\alpha = \frac{3\pi}{4}$ ,  $\gamma_1 = \frac{\pi}{6}$ ,  $\gamma_2 = \frac{5\pi}{6}$ .



**Figure 6.**  $\mathcal{F}$  at a  $D_1^+$  corner;  $\alpha = \frac{5\pi}{6}$ ,  $\gamma_1 = \frac{\pi}{3}$ ,  $\gamma_2 = \frac{\pi}{6}$ .



**Figure 7.**  $\mathcal{F}$  at a  $D_1^+$  corner;  $\alpha = \frac{3\pi}{4}$ ,  $\gamma_1 = \frac{\pi}{6}$ ,  $\gamma_2 = \frac{\pi}{4}$ .



**Figure 8.**  $\mathcal{F}$  at a  $D_1^+$  corner;  $\alpha = \frac{11\pi}{12}$ ,  $\gamma_1 = \frac{7\pi}{12}$ ,  $\gamma_2 = \frac{\pi}{12}$ .

**Theorem 2.5.** *Let  $2\alpha > \pi$  and  $\Omega$  and  $\gamma$  be as in Section 1 and suppose (5) holds with  $\gamma_1, \gamma_2 \in (0, \pi)$ . Let  $\beta \in (-\alpha, \alpha)$  and  $(x_j) \in s(V)$  such that*

$$(14) \quad \lim_{j \rightarrow \infty} \frac{x_j}{|x_j|} = (\cos \beta, \sin \beta).$$

*Continuing to write  $\omega(\theta) = (\cos \theta, \sin \theta, 0)$  for  $\theta \in \mathbb{R}$ , we see:*

(i) *Suppose  $(\gamma_1, \gamma_2) \in D_1^+$  (i.e.,  $\gamma_1 + \gamma_2 < 2\alpha - \pi$ ),  $\lim_{j \rightarrow \infty} \vec{n}_f(x_j)$  exists and*

$$\lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \omega(\theta) \quad \text{for some } \theta \in \mathbb{R}.$$

*Then  $(\beta, \theta) \in \mathcal{F}$ .*

(ii) *Suppose  $(\gamma_1, \gamma_2) \in D_1^-$  (i.e.,  $\gamma_1 + \gamma_2 > 2\alpha + \pi$ ),  $\lim_{j \rightarrow \infty} \vec{n}_f(x_j)$  exists and*

$$\lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \omega(\theta) \quad \text{for some } \theta \in \mathbb{R}.$$

*Then  $(-\beta, \theta) \in \mathcal{F}$ .*

(iii) *Connectedness at  $\beta$ : Suppose  $(\gamma_1, \gamma_2) \in D_1^+$  and  $(x_j), (y_j) \in s(\Omega)$  such that*

$$\begin{aligned} \lim_{j \rightarrow \infty} \frac{x_j}{|x_j|} = \lim_{j \rightarrow \infty} \frac{y_j}{|y_j|} &= (\cos \beta, \sin \beta), \\ \lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \omega(\theta_1) \quad \text{and} \quad \lim_{j \rightarrow \infty} \vec{n}_f(y_j) &= \omega(\theta_2), \end{aligned}$$

*for some  $\theta_1 \leq \theta_2$  such that  $(\beta, \theta_1), (\beta, \theta_2) \in \mathcal{F}$ . Then the set  $\{\theta \in [\theta_1, \theta_2] : (\beta, \theta) \in \mathcal{F}\}$  must be connected.*

(iv) *Connectedness: Suppose  $(\gamma_1, \gamma_2) \in D_1^+$ . Let  $\beta_1, \beta_2 \in (-\alpha, \alpha)$  with  $\beta_1 \leq \beta_2$ . Suppose  $(x_j), (y_j) \in s(\Omega)$  such that*

$$\begin{aligned} \lim_{j \rightarrow \infty} \frac{x_j}{|x_j|} = (\cos \beta_1, \sin \beta_1), \quad \lim_{j \rightarrow \infty} \frac{y_j}{|y_j|} &= (\cos \beta_2, \sin \beta_2), \\ \lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \omega(\theta_1) \quad \text{and} \quad \lim_{j \rightarrow \infty} \vec{n}_f(y_j) &= \omega(\theta_2), \end{aligned}$$

*for some  $\theta_1, \theta_2$  such that  $(\beta_1, \theta_1), (\beta_2, \theta_2) \in \mathcal{F}$ . Set  $L = [\min\{\theta_1, \theta_2\}, \max\{\theta_1, \theta_2\}]$ . Then the set  $\mathcal{F} \cap ([\beta_1, \beta_2] \times L)$  must be connected.*

*Proof.* The proof of Theorem 2.5 (i) and (ii) is essentially the same as that of Theorem 2.4 with Corollaries 2.6 and 2.7 of [Jeffres and Lancaster 2008] replacing Corollaries 2.4 and 2.5 respectively. Conclusion (iii) follows from (i) by standard arguments (e.g., proof of Lemma 4.2). Conclusion (iv) follows from (i) by standard arguments which take into account the specific geometry of  $\mathcal{F}$ . □

**2B. Proof of Theorem 2.1.** Assume  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ ,  $f$  satisfies (3) in  $\Omega$  and (4) on  $B(\mathbb{C}, \delta^*) \cap \partial\Omega \setminus \{\mathbb{0}\}$  and  $f$  is continuous at  $\mathbb{0}$ ; then  $f$  is bounded in a neighborhood of  $\mathbb{0}$ . Since  $f$  is continuous at  $\mathbb{0}$ , we have the following modifications of (i)–(viii) in Section 2A:

- (iii)'  $X$  maps  $\partial E$  strictly monotonically onto  $\{(x, f(x)) : x \in \partial\Omega\}$ .
- (vi)'  $X \in C^0(\bar{E})$  and  $X(1, 0) = (0, 0, z_0)$ , where  $z_0 = f(0, 0)$ .
- (vii)' Continuing to write  $G(u, v) = (x(u, v), y(u, v))$ ,  $G(\cos t, \sin t)$  moves clockwise about  $\partial\Omega$  as  $t$  increases,  $0 \leq t \leq 2\pi$ , and  $G$  is an orientation reversing homeomorphism from  $\bar{E}$  onto  $\bar{\Omega}$ .

We will prove Theorem 2.1 in the cases  $(\gamma_1, \gamma_2) \in D_2^+$  and  $(\gamma_1, \gamma_2) \in D_1^+$ ; this will suffice to prove the lemma since the mapping

$$\mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad (x_1, x_2, x_3) \mapsto (x_1, -x_2, -x_3),$$

converts a  $D_2^-$  corner into a  $D_2^+$  corner and converts a  $D_1^-$  corner into a  $D_1^+$  corner.

Suppose  $(\gamma_1, \gamma_2) \in D_2^+$ . Set  $\theta_1 = (\pi - (\gamma_1 + \gamma_2))/2$  and let  $\theta_2 \in (\alpha - \gamma_1, \alpha)$ . By choosing  $\delta_0 > 0$  small, we may assume

$$\Omega^* = \{(r \cos \theta, r \sin \theta) : 0 < r < \delta_0, \theta_1 < \theta < \theta_2\} \subset \Omega.$$

Notice that Theorem 2.4 ( $D_2^+$ ) implies

$$\Pi_f(\Omega^*) = \{(\cos \theta, \sin \theta, 0) : \theta_1 + \frac{1}{2}\pi \leq \theta \leq \alpha - \gamma_1 + \frac{1}{2}\pi\}.$$

Since  $\alpha - \gamma_1 - \theta_1 = \frac{1}{2}(2\alpha - \pi - \gamma_1 + \gamma_2) \in (2\alpha - \pi, \alpha) \subset (0, \pi)$ , the hypotheses of Lemma 2.2 are satisfied (with  $M$  a rotation through an angle of  $\pi/2 - \alpha$ ). If  $\phi$  is a conformal map from  $E$  onto  $G^{-1}(\Omega^*)$  which maps  $(1, 0)$  to  $(1, 0)$  and  $\tilde{g}$  is defined by (10), then Lemma 2.2 implies there exists  $p > 2$  such that

$$(15) \quad \tilde{g}(\zeta) = \psi(\zeta) + h(\zeta),$$


where  $\psi$  is a holomorphic function and  $h \in L^\infty(E_1)$  is a Hölder continuous function on  $\bar{E}_1$  with Hölder exponent  $\mu = (p - 2)/p$ . The assumption that  $f$  is continuous at  $\mathbb{0}$  yields a contradiction as in §3 of [L] (i.e., the Phragmén–Lindelöf theorem is violated).

Now suppose  $(\gamma_1, \gamma_2) \in D_1^+$ . Let  $\theta_1 \in (-\alpha, -\alpha + \gamma_2)$  and  $\theta_2 \in (\alpha - \gamma_1, \alpha)$  and choose  $\delta_0 > 0$  small enough that


$$\Omega^* = \{(r \cos \theta, r \sin \theta) : 0 < r < \delta_0, \theta_1 < \theta < \theta_2\} \subset \Omega.$$

Using Theorem 2.5, we see that

$$\Pi_f(\Omega^*) \subset \{(\cos \theta, \sin \theta, 0) : \beta \in (-\alpha, \alpha), (\beta, \theta) \in \mathcal{F}_L\},$$




**Figure 9.**  $\mathcal{F}_A$  at a  $D_1^+$  corner;  $\alpha = \frac{5\pi}{6}$ ,  $\gamma_1 = \frac{\pi}{3}$ ,  $\gamma_2 = \frac{\pi}{6}$ .



**Figure 10.**  $\mathcal{F}_B$  at a  $D_1^+$  corner;  $\alpha = \frac{5\pi}{6}$ ,  $\gamma_1 = \frac{\pi}{3}$ ,  $\gamma_2 = \frac{\pi}{6}$ .

where  $\mathcal{F}_L$  is one of the sets  $\mathcal{F}_A$ ,  $\mathcal{F}_B$  or  $\mathcal{F}_C$  illustrated in Figures 9, 10 and 11 respectively. When  $\mathcal{F}_L$  is  $\mathcal{F}_A$  or  $\mathcal{F}_C$ , the proof is essentially that same as that above for the case in which  $(\gamma_1, \gamma_2) \in D_2^+$ . When  $\mathcal{F}_L$  is  $\mathcal{F}_B$ , the proof is essentially that same as that in §3 of [L]. □



**Figure 11.**  $\mathcal{F}_C$  at a  $D_1^+$  corner;  $\alpha = \frac{5\pi}{6}$ ,  $\gamma_1 = \frac{\pi}{3}$ ,  $\gamma_2 = \frac{\pi}{6}$ .

### 3. The exact sizes of fans

We recall that a solution  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{O}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{O}\})$  of (3)–(4) is unbounded if  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2) \in D_1^\pm$ , for some  $\delta > 0$  and  $\rho \in (0, 1)$ . The following lemma justifies the definition of  $\vec{m}_f : (-\alpha, \alpha) \rightarrow S_-^2$ , given by


$$\vec{m}_f(\beta) = \lim_{j \rightarrow \infty} \vec{n}_f(x_j) \text{ whenever } (x_j) \in s(\Omega) \text{ with } \lim_{j \rightarrow \infty} \frac{x_j}{|x_j|} = (\cos \beta, \sin \beta),$$

when  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ .

**Lemma 3.1.** *Let  $\Omega$  and  $\gamma$  be as in Section 1, with  $\alpha \in [0, \pi]$ . For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{O}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{O}\})$  is a bounded solution of (3)–(4) with  $|H|_\infty = \sup_{\mathbf{x} \in \Omega} |H(\mathbf{x}, f(\mathbf{x}))| < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ ; that is, either  $\alpha \in [0, \frac{\pi}{2}]$  and  $|\gamma_1 - \gamma_2| > \pi - 2\alpha$  holds or  $\alpha \in [\frac{\pi}{2}, \pi]$  and one of  $|\gamma_1 - \gamma_2| > 2\alpha - \pi$  or  $|\gamma_1 + \gamma_2 - \pi| > 2\pi - 2\alpha$  holds. Then the Gauss map from  $\mathcal{S}_f$  to  $S_-^2$  is continuous on  $\mathcal{S}_f \cap (\overline{\Omega}(\epsilon) \times \mathbb{R})$  for each  $\epsilon > 0$ , where  $\Omega(\epsilon) = \{(r \cos \theta, r \sin \theta) \in \Omega : r > 0, |\theta| < \alpha - \epsilon\}$ . In particular,  $\vec{m}_f(\beta)$  exists for all  $\beta \in (-\alpha, \alpha)$  and  $\vec{m}_f \in C^0((-\alpha, \alpha) : S_-^2)$ .*

*Proof.* Using Theorem 2.1 of [LSa] when  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2) \in D_2^\pm$  and Theorems 2.4 and 2.5 and the proof of Theorem 2.1 when  $\alpha \geq \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_1^\pm \cup D_2^\pm$ , we see that the hypotheses of Lemma 2.2 are satisfied when  $\epsilon > 0$  and  $\Lambda = \Omega(\epsilon)$ . Therefore the restriction of the Gauss map to  $\mathcal{S}_f \cap (\overline{\Omega}(\epsilon) \times \mathbb{R})$  is continuous.  $\square$





**Figure 12.** Radial limits: Side fans at a convex corner.

Now we wish to determine the exact sizes of the side fans (illustrated in Figure 12) when  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ . From [LS a], Theorems 1 and 2, we know that if  $f$  is discontinuous at  $\mathbb{O}$ , then  $Rf$  and the limits at  $\mathbb{O}$  of the Gauss map behave in the following ways; here  $\mathcal{I}$  denotes a countable subset of the appropriate interval(s) and  $\omega(\theta) = (\cos \theta, \sin \theta, 0)$  for  $\theta \in \mathbb{R}$ .

**Case (I)**  $Rf$  is  $\begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly increasing} & \text{on } [\alpha_1, \alpha_2] \setminus \mathcal{I}, \\ \text{constant} & \text{on } [-\alpha, \alpha_1]. \end{cases}$

This case can only occur when  $(\gamma_1, \gamma_2) \in R \cup D_2^+ \cup D_1^\pm$ . Theorem 2 of [LS a] implies  $\alpha_2 \leq \alpha - \gamma_1$  and  $\alpha_1 \geq -\alpha + \pi - \gamma_2$ . If  $\beta \in (\alpha_1, \alpha_2)$  then  $\vec{m}_f(\beta) = \omega(\beta + \frac{\pi}{2})$ .

**Case (D)**  $Rf$  is  $\begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly decreasing} & \text{on } [\alpha_1, \alpha_2] \setminus \mathcal{I}, \\ \text{constant} & \text{on } [-\alpha, \alpha_1]. \end{cases}$

This case can only occur when  $(\gamma_1, \gamma_2) \in R \cup D_2^- \cup D_1^\pm$ . Theorem 2 of [LS a] implies  $\alpha_2 \leq \alpha - \pi + \gamma_1$  and  $\alpha_1 \geq -\alpha + \gamma_2$ . If  $\beta \in (\alpha_1, \alpha_2)$ , then  $\vec{m}_f(\beta) = \omega(\beta - \frac{\pi}{2})$ .

**Case (DI)** There exists  $\theta_0 \in (-\alpha + \gamma_2, \alpha - \gamma_1 - \pi)$  such that

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly increasing} & \text{on } [\theta_0 + \pi, \alpha_2] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [\theta_0, \theta_0 + \pi], \\ \text{strictly decreasing} & \text{on } [\alpha_1, \theta_0] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, \alpha_2]. \end{cases}$$

This case can only occur when  $(\gamma_1, \gamma_2) \in D_1^+$ . Theorem 2 of [LS a] implies  $\alpha_2 \leq \alpha - \gamma_1$  and  $\alpha_1 \geq -\alpha + \gamma_2$ . If  $\beta \in (-\alpha, \alpha)$ , then

$$\vec{m}_f(\beta) = \begin{cases} \omega(\beta + \frac{\pi}{2}) & \text{if } \beta \in (\theta_0 + \pi, \alpha_2), \\ \omega(\theta_0 - \frac{\pi}{2}) & \text{if } \beta \in [\theta_0, \theta_0 + \pi], \\ \omega(\beta - \frac{\pi}{2}) & \text{if } \beta \in (\alpha_1, \theta_0). \end{cases}$$

**Case (ID)** There exists  $\theta_0 \in (-\alpha + \pi - \gamma_2, \alpha + \gamma_1 - 2\pi)$  such that

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly decreasing} & \text{on } [\theta_0 + \pi, \alpha_2] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [\theta_0, \theta_0 + \pi], \\ \text{strictly increasing} & \text{on } [\alpha_1, \theta_0] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, \alpha_1]. \end{cases}$$

This case can only occur when  $(\gamma_1, \gamma_2) \in D_1^-$ . Theorem 2 of [LS a] implies  $\alpha_2 \leq \alpha - \pi + \gamma_1$  and  $\alpha_1 \geq -\alpha + \pi - \gamma_2$ . If  $\beta \in (-\alpha, \alpha)$ , then

$$\vec{m}_f(\beta) = \begin{cases} \omega(\beta - \frac{\pi}{2}) & \text{if } \beta \in (\theta_0 + \pi, \alpha_2), \\ \omega(\theta_0 + \frac{\pi}{2}) & \text{if } \beta \in [\theta_0, \theta_0 + \pi], \\ \omega(\beta + \frac{\pi}{2}) & \text{if } \beta \in (\alpha_1, \theta_0). \end{cases}$$

**Theorem 3.2.** *Let  $\Omega$  and  $\gamma$  be as in Section 1, with  $\alpha \in [\frac{\pi}{2}, \pi]$ . For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{C}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{C}\})$  is a bounded solution of (3)–(4) with  $|H|_\infty < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ . Then:*

- (i) In Case (I),  $\alpha_1 = -\alpha + \pi - \gamma_2$  and  $\alpha_2 = \alpha - \gamma_1$ .
- (ii) In Case (D),  $\alpha_1 = -\alpha + \gamma_2$  and  $\alpha_2 = \alpha - \pi + \gamma_1$ .
- (iii) In Case (DI),  $\alpha_1 = -\alpha + \gamma_2$  and  $\alpha_2 = \alpha - \gamma_1$ .
- (iv) In Case (ID),  $\alpha_1 = -\alpha + \pi - \gamma_2$  and  $\alpha_2 = \alpha - \pi + \gamma_1$ .

*Proof.* Suppose  $(\gamma_1, \gamma_2) \in D_2^\pm$ ; the argument is the same when  $\alpha < \pi/2$  and when  $\alpha \geq \pi/2$ . Let us assume  $(\gamma_1, \gamma_2) \in D_2^+$ ; hence Case (I) holds. Then Figure 5 illustrates the conclusions of Theorem 2.1 of [L] and Theorem 2.4. Suppose there exists  $\alpha_2 < \alpha - \gamma_1$  (and  $\alpha_1 \geq -\alpha + \pi - \gamma_2$ ) such that

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly increasing} & \text{on } [\alpha_1, \alpha_2] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, \alpha_1]. \end{cases}$$

If we define

$$\Omega' = \{(r \cos \beta, r \sin \beta) \in \Omega : 0 < r < \delta, \alpha_2 < \beta < \pi\}$$

for  $\delta > 0$  sufficiently small, then  $f \in C^0(\overline{\Omega'})$  and we may apply the arguments in the proof of Theorem 2.1, using  $\Omega'$  as our domain, to obtain a contradiction. If  $\alpha_1 > -\alpha + \pi - \gamma_2$ , a similar argument yields a contradiction.

Now suppose  $(\gamma_1, \gamma_2) \in D_1^+$ , Case (I) holds and there exist  $\alpha_2 < \alpha - \gamma_1$  and  $\alpha_1 \geq -\alpha + \pi - \gamma_2$  such that

$$Rf \text{ is } \begin{cases} \text{constant} & \text{on } [\alpha_2, \alpha], \\ \text{strictly increasing} & \text{on } [\alpha_1, \alpha_2] \setminus \mathcal{F}, \\ \text{constant} & \text{on } [-\alpha, \alpha_1]. \end{cases}$$

Let  $\theta_1 \in (-\alpha, -\alpha_1 + \pi - \gamma_2)$  and  $\theta_2 \in (\alpha - \gamma_1, \alpha)$ . By choosing  $\delta_0 > 0$  small, we may assume  $\Omega^* = \{(r \cos \theta, r \sin \theta) : 0 < r < \delta_0, \theta_1 < \theta < \theta_2\}$  is a subset of  $\Omega$ . Set  $\Omega' = \{(r \cos \beta, r \sin \beta) : 0 < r < \delta_0, \alpha_2 < \theta < \theta_2\}$  and notice that  $f \in C^0(\overline{\Omega'})$ . Now Theorem 2.5, Lemma 3.1 and the fact that

$$\lim_{j \rightarrow \infty} \vec{n}_f(x_j) = \omega(\beta + \frac{\pi}{2})$$

when  $\beta \in (\alpha_1, \alpha_2)$  and  $(x_j) \in s(\Omega)$  such that  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos \beta, \sin \beta)$  implies that

$$\Pi_f(\Omega^*) \subset \{\omega(\theta) : \beta \in [\theta_1, \theta_2], (\beta, \theta) \in \mathcal{F}_C\}$$

and  $\vec{m}_f(\cdot) \in C^0((-\alpha, \alpha))$ . If  $\vec{m}_f(\alpha_2) \neq \omega(\alpha - \gamma_1 + \frac{\pi}{2})$ , then we may apply the arguments in the proof of Theorem 2.1, using  $\Omega'$  as our domain, to get a contradiction. If  $\vec{m}_f(\alpha_2) = \omega(\alpha - \gamma_1 + \frac{\pi}{2})$ , then  $\vec{m}_f$  is discontinuous at  $\alpha_2$ , which is a contradiction. Therefore  $\alpha_2 = \alpha - \gamma_1$ . The argument that  $\alpha_1 = -\alpha + \pi - \gamma_2$  is similar.

The proof of the theorem when  $(\gamma_1, \gamma_2) \in D_1^+$  and one of Cases (D), (DI) or (ID) occurs follows in a similar manner. The situation where  $(\gamma_1, \gamma_2) \in D_1^-$  follows from this. □

### 4. Continuity of the Gauss map

Notice that Lemma 3.1 and the proof of Theorem 3.2 imply that the (nonparametric) Gauss map is continuous on  $\mathcal{S}_f \cap (\overline{\Omega_\epsilon} \times \mathbb{R})$  for each  $\epsilon > 0$  and, in each case, we have:

- (I):  $\lim_{\beta \rightarrow \alpha} \vec{m}_f(\beta) = \omega(\alpha_2 + \pi/2)$  and  $\lim_{\beta \rightarrow -\alpha} \vec{m}_f(\beta) = \omega(\alpha_1 + \pi/2)$ .
- (D):  $\lim_{\beta \rightarrow \alpha} \vec{m}_f(\beta) = \omega(\alpha_2 - \pi/2)$  and  $\lim_{\beta \rightarrow -\alpha} \vec{m}_f(\beta) = \omega(\alpha_1 - \pi/2)$ .
- (DI):  $\lim_{\beta \rightarrow \alpha} \vec{m}_f(\beta) = \omega(\alpha_2 + \pi/2)$  and  $\lim_{\beta \rightarrow -\alpha} \vec{m}_f(\beta) = \omega(\alpha_1 - \pi/2)$ .
- (ID):  $\lim_{\beta \rightarrow \alpha} \vec{m}_f(\beta) = \omega(\alpha_2 - \pi/2)$  and  $\lim_{\beta \rightarrow -\alpha} \vec{m}_f(\beta) = \omega(\alpha_1 + \pi/2)$ .

In order to conclude that the Gauss map is in  $C^0(\mathcal{S}_f \cap (B(\mathbb{C}, \delta) \times \mathbb{R}) : S^2_-)$ , it would be sufficient to blow up the graph of  $u(x) = f(x) - Rf(\alpha)$  (or  $u(x) = f(x) - Rf(-\alpha)$ ) about  $(0, 0, 0)$  tangent to  $\partial^+ \Omega$  (or  $\partial^- \Omega$  respectively) and know that a subsequence converges to an appropriate cone. If one is willing to accept this hypothesis, then the claim that the Gauss map is in  $C^0(\mathcal{S}_f \cap (B(\mathbb{C}, \delta) \times \mathbb{R}) : S^2_-)$  can be proven.

**Hypothesis (B±).** For all  $(x_j) \in s(\Omega)$  with  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos(\pm\alpha), \sin(\pm\alpha))$ , there is a subsequence  $(x_{j_k})$  and a function  $u_\infty : \Omega_\infty \rightarrow [-\infty, \infty]$  such that the subgraph  $U_\infty = \{(x, t) \in \Omega_\infty \times \mathbb{R} : t < u_\infty(x)\}$  of  $u_\infty$  is a cone with respect to  $(0, 0, 0)$ , there exists  $\vec{\xi} \in S^2_-$  such that  $\lim_{k \rightarrow \infty} \vec{n}(x_{j_k}) = \vec{\xi} = (\xi_1, \xi_2, \xi_3)$ ,

$$\lim_{k \rightarrow \infty} \text{dist}(\{(x, u_{j_k}(x)) \in \Omega_{j_k}(\delta, b)\}, \partial U_\infty \cap \Omega_{j_k}(\delta, b)) = 0$$

for each  $\delta > 0$  and  $b > 0$ , where  $\epsilon_j = |x_j|$ ,  $u_j(x) = (f(\epsilon_j x) - Rf(\pm\alpha))/\epsilon_j$  and  $\Omega_j(\delta, b) = \{(x, t) \in \mathbb{R}^3 : x \in B(\mathbb{C}, \delta), \epsilon_j x \in \Omega, t \in (-b, b)\}$  for  $j \in \mathbb{N}$ , and

(a) if  $\xi_3 < 0$ , then  $\partial U_\infty = \pi_1 \cap (\overline{\Omega_\infty} \times \mathbb{R})$ ,  $\pi_1$  is a nonvertical plane with downward unit normal  $\vec{\xi} \in S^2_-$ ,  $\vec{\xi}$  makes an angle of  $\gamma_1$  with the exterior unit normal to  $\partial^+ \Omega_\infty \times \mathbb{R}$  and an angle of  $\gamma_2$  with the exterior unit normal to  $\partial^- \Omega_\infty \times \mathbb{R}$  and  $\vec{n}_{u_{j_k}} \rightarrow \vec{\xi}$  uniformly on compacta in  $\Omega \times \mathbb{R}$  as  $k \rightarrow \infty$ ,

(b) if  $\xi_3 = 0$ , then  $\partial U_\infty = \partial \mathcal{P} \cap (\overline{\Omega_\infty} \times \mathbb{R})$ ,  $\mathcal{P} = \{(r \cos \theta, r \sin \theta) : r > 0, \theta \in \Theta\}$  and, for each  $x \in \partial \mathcal{P} \cap \Omega_\infty$ ,  $\vec{n}_{u_{j_k}}(x) \rightarrow \vec{\xi}(x) \times \{0\}$ , where  $\mathcal{P} = \{x \in \Omega_\infty : u_\infty(x) = \infty\}$ ,  $\vec{\xi}(x)$  is the interior (with respect to  $\mathcal{P}$ ) unit normal vector to  $\partial \mathcal{P}$  at  $x$  and  $\Theta$  is one of the following sets:  $(\alpha - \gamma_1, \alpha)$ ,  $(-\alpha, -\alpha + \gamma_2)$ ,  $(-\alpha + \pi - \gamma_2, \alpha - \pi + \gamma_1)$  (provided  $(\alpha - \pi + \gamma_1) - (-\alpha + \pi - \gamma_2) \geq \pi$ ) or  $(-\alpha, -\alpha + \gamma_2) \cup (\alpha - \gamma_1, \alpha)$  (provided  $(\alpha - \gamma_1) - (-\alpha + \gamma_2) \geq \pi$ ).

**Theorem 4.1.** *Let  $\Omega$  and  $\gamma$  be as in Section 1. For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{C}, \delta) \cap \overline{\Omega} \setminus \{0\})$  is a bounded solution of (3)–(4) with  $|H|_\infty < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and either  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_2^\pm$  or  $\alpha \geq \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_1^\pm \cup D_2^\pm$ . Suppose that Hypotheses (B±) are true. Then  $\vec{n}_f : \mathcal{S}_f \times (\Omega \times \mathbb{R})$  extends to be continuous on  $\mathcal{S}_f \cap (B(\mathbb{C}, \delta) \times \mathbb{R})$  and  $N \in C^0(E \cup \{(u, v) \in \partial E : G(u, v) \in \partial \Omega \cap B(\mathbb{C}, \delta)\})$ .*

The proof of this theorem will follow from the information at the beginning of this section about the behavior of  $\vec{m}_f$  and from Lemmas 4.2–4.5. Set

$$\begin{aligned}
 C(x) &= \{X \in S_-^2 : X \cdot v(x) = \cos \gamma(x)\}, \\
 \Gamma_1 &= \{X \in S_-^2 : X \cdot (\cos(\alpha + \frac{\pi}{2}), \sin(\alpha + \frac{\pi}{2}), 0) = \cos \gamma_1\}, \\
 \Gamma_2 &= \{X \in S_-^2 : X \cdot (\cos(-\alpha - \frac{\pi}{2}), \sin(-\alpha - \frac{\pi}{2}), 0) = \cos \gamma_2\}, \\
 \vec{\xi}_A &= \omega(\alpha - \gamma_1 + \frac{\pi}{2}) \in \Gamma_1, \quad \vec{\xi}_B = \omega(\alpha + \gamma_1 - \frac{3\pi}{2}) \in \Gamma_1, \\
 \vec{\xi}_C &= \omega(-\alpha - \gamma_2 + \frac{3\pi}{2}) \in \Gamma_2, \quad \vec{\xi}_D = \omega(-\alpha + \gamma_2 - \frac{\pi}{2}) \in \Gamma_2, \\
 \Omega_\infty &= \{(r \cos(\theta), r \sin(\theta)) : r > 0, -\alpha < \theta < \alpha\}, \\
 \Sigma_\infty^1 &= \{(r \cos(\alpha), r \sin(\alpha)) : r > 0\}, \quad \Sigma_\infty^2 = \{(r \cos(\alpha), -r \sin(\alpha)) : r > 0\}, \\
 v_\infty^+ &= (\cos(\alpha + \frac{\pi}{2}), \sin(\alpha + \frac{\pi}{2}), 0) \quad \text{and} \quad v_\infty^- = (\cos(-\alpha - \frac{\pi}{2}), \sin(-\alpha - \frac{\pi}{2}), 0).
 \end{aligned}$$

**Lemma 4.2.** *Let  $\Omega$  and  $\gamma$  be as in Section 1. For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{O}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{O}\})$  is a bounded solution of (3)–(4) with  $|H|_\infty < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and either  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_2^+$ , so that  $Rf$  behaves as in Case (I), or  $\alpha \geq \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_1^\pm \cup D_2^+$  and  $Rf$  behaves as in Case (I) or Case (DI). Assume Hypothesis (B+) is true. Then*

$$(16) \quad \lim_{j \rightarrow \infty} \vec{n}(x_j) = (\cos(\alpha - \gamma_1 + \pi/2), \sin(\alpha - \gamma_1 + \pi/2), 0) = \vec{\xi}_A$$

for every  $(x_j) \in s(\Omega)$  with  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos \alpha, \sin \alpha)$ .

*Proof.* Since  $\gamma(x) \rightarrow \gamma_1$  and  $v(x) \rightarrow v_\infty^+$  as  $x \in \partial^+ \Omega$  converges to  $\mathbb{O}$ , we see that  $\text{dist}(C(x), \Gamma_1) \rightarrow 0$  as  $x \in \partial^+ \Omega$  converges to  $\mathbb{O}$ . Similarly,  $\gamma(x) \rightarrow \gamma_2$ ,  $v(x) \rightarrow v_\infty^-$  and  $\text{dist}(C(x), \Gamma_2) \rightarrow 0$  as  $x \in \partial^- \Omega$  converges to  $\mathbb{O}$ . Thus

$$(17) \quad \text{dist}(\vec{n}(x^+(s)), \Gamma_1) \rightarrow 0 \quad \text{and} \quad \text{dist}(\vec{n}(x^-(s)), \Gamma_2) \rightarrow 0$$

as  $s \rightarrow 0+$ .

Suppose that  $(x_j) \in s(\Omega)$  with  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos \alpha, \sin \alpha)$ ; then there is a subsequence, still denoted  $(x_j)$ , and  $\vec{\xi} \in S_-^2$  such that  $\lim_{j \rightarrow \infty} \vec{n}(x_j, y_j) \rightarrow \vec{\xi}$ . Notice that  $\vec{\xi} \in \Gamma_1$  since  $f \in C^{1,\rho}(B(\mathbb{O}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{O}\})$  and (17) holds.

Assume first that  $\vec{\xi} = (\xi_1, \xi_2, \xi_3)$  with  $\xi_3 < 0$ . For each  $j \in \mathbb{N}$ , define  $\epsilon_j = |x_j|$ ,  $\Omega_j = \{x \in \mathbb{R}^2 : \epsilon_j x \in \Omega\}$  and  $u_j \in C^\infty(\Omega_j) \cap C^1(\overline{\Omega_j} \setminus \{\mathbb{O}\})$  by

$$(18) \quad u_j(x) = \frac{1}{\epsilon_j} (f(\epsilon_j x) - Rf(\alpha)).$$

Let  $\gamma_j$  be defined on  $\partial\Omega_j \setminus \{\mathbb{O}\}$  by  $\gamma_j(x) = \gamma(\epsilon_j x)$  and let  $\nu_j$  denote the outward unit normal to  $\partial\Omega_j$ . Then  $u_j$  satisfies the prescribed mean curvature problem

$$(19) \quad Nu_j(x) = \epsilon_j H(\epsilon_j x, f(\epsilon_j x)), \quad x \in \Omega_j,$$

$$(20) \quad Tf_j \cdot \nu_j = \cos \gamma_j \quad \text{on } \partial\Omega_j \setminus \{\mathbb{O}\}.$$

Hypothesis (B+) implies that there is a nonvertical plane  $\pi_1$  with downward unit normal  $\vec{\xi}$  which meets  $\partial^+\Omega_\infty$  in an angle of  $\gamma_1$  and  $\partial^-\Omega_\infty$  in an angle of  $\gamma_2$  in the sense described in (a); however this is impossible since  $(\gamma_1, \gamma_2) \in D_1^\pm \cup D_2^\pm$ . Thus  $\xi_3 = 0$  and so either  $\vec{\xi} = \vec{\xi}_A$  or  $\vec{\xi} = \vec{\xi}_B$ . The intermediate value theorem implies that

$$(21) \quad \text{if } \vec{\xi} = \vec{\xi}_A, \quad \text{then } \vec{n}(x) \rightarrow \vec{\xi}_A \text{ as } x \in \partial^+\Omega \text{ converges to } \mathbb{O},$$

$$(22) \quad \text{if } \vec{\xi} = \vec{\xi}_B, \quad \text{then } \vec{n}(x) \rightarrow \vec{\xi}_B \text{ as } x \in \partial^+\Omega \text{ converges to } \mathbb{O}.$$

Suppose (22) holds. Notice then that

$$(23) \quad \lim_{s \rightarrow 0^+} \frac{d}{ds} f^+(s) = -\infty,$$

since  $(\cos \alpha, \sin \alpha, 0) \cdot \vec{\xi}_B = -\sin(\gamma_1) < 0$ , and so  $f^+(s) = f(x^+(s))$  is a strictly decreasing function of  $s$  for  $0 \leq s \leq s_0$ , where  $s_0 > 0$  is sufficiently small. Since  $\vec{m}_f(\beta) = \omega(\alpha - \gamma_1 + \frac{\pi}{2})$  when  $\beta \in [\alpha - \gamma_1, \alpha)$ , we have

$$(24) \quad \lim_{r \rightarrow 0^+} \nabla f(r \cos \beta, r \sin \beta) \cdot (\cos \beta, \sin \beta) = +\infty \quad \text{for } \beta \in (\alpha - \gamma_1, \alpha).$$

Since  $Rf$  behaves as in Case (I) or (DI), we have

$$(25) \quad Rf(\beta) < Rf(\alpha) = f^+(0) \quad \text{if } \beta \in [\alpha - \pi, \alpha - \gamma_1).$$

Let  $\Omega_H$  be the connected component of

$$\{(r \cos \beta, r \sin \beta) \in \Omega : r > 0, \alpha - \pi < \beta < 5\pi/4\}$$

that contains  $\{(r \cos \beta, r \sin \beta) : 0 < r < \delta, \alpha - \gamma_1 < \beta < \alpha - \epsilon\}$  for sufficiently small  $\epsilon, \delta > 0$ . Consider the  $k = f^+(0) (= Rf(\alpha))$  level set of  $f$  in  $\Omega_H$ . From (23), we see that there is a component  $C$  of  $\{x \in \Omega_H : f(x) < k\}$  whose boundary contains  $\partial^+\Omega \cap B(\mathbb{O}, \tau)$  for  $\tau > 0$  sufficiently small; let  $c_\alpha$  be the component of  $\Omega_H \cap \partial C$  whose closure contains  $\mathbb{O}$ . Then (23) and (24) imply that for every  $\beta_1 < \alpha$  and  $\beta_2 > \alpha$ , there exists  $\epsilon > 0$  such that

$$c_\alpha \cap B(\mathbb{O}, \epsilon) \subset \{(r \cos \theta, r \sin \theta) : 0 < r < \epsilon, \beta_1 < \theta < \beta_2\};$$

in this sense,  $c_\alpha$  is tangent to  $\theta = \alpha$  at  $\mathbb{O}$ . Similarly, using (24) and (25), we see that there is a  $k$ -level curve of  $f$ , denoted  $c_{\alpha-\gamma_1}$ , which is tangent to  $\theta = \alpha - \gamma_1$  at  $\mathbb{O}$  in the sense that for every  $\beta_1 < \alpha - \gamma_1$  and  $\beta_2 > \alpha - \gamma_1$ , there exists  $\epsilon > 0$  such that

$$c_{\alpha-\gamma_1} \cap B(\mathbb{O}, \epsilon) \subset \{(r \cos \theta, r \sin \theta) : 0 < r < \epsilon, \beta_1 < \theta < \beta_2\}.$$

Now pick  $\tau > 0$  small enough that the region bounded by  $c_\alpha, c_{\alpha-\gamma_1}$  and  $\{r = \tau\}$  is well-defined, connected and simply connected; let us rotate this region about  $\mathbb{O}$  through an angle  $(\pi + \gamma_1)/2 - \alpha$  and denote this open set as  $\Omega^\tau$ , so that  $\partial\Omega^\tau$  is tangent to  $\theta = (\pi \pm \gamma_1)/2$  at  $\mathbb{O}$ . Notice that  $\tilde{f} = f \circ R^{-1} \in C^0(\overline{\Omega^\tau})$  if  $R$  denotes the rotation above. We will let a particular portion of a suitable nodoid be the graph of a comparison function  $h$  over a domain  $U^\tau \subset \Omega^\tau$  with  $B(\mathbb{O}, \epsilon) \cap U^\tau = B(\mathbb{O}, \epsilon) \cap \Omega^\tau$  for some  $\epsilon > 0$ . Now  $\partial U^\tau$  will consist of two disjoint, connected curves,  $\partial_1 U^\tau \subset \partial\Omega^\tau \setminus \{r = \tau\}$  and  $\partial_2 U^\tau$ , with  $\mathbb{O} \in \partial_1 U^\tau$  and  $\mathbb{O} \notin \partial_2 U^\tau$ . The comparison function  $h \in C^0(\overline{U^\tau}) \cap C^1(U^\tau \cup \partial_1 U^\tau)$  will have the properties  $h(\mathbb{O}) = k, \frac{\partial h}{\partial x_2}(\mathbb{O}) < \infty, h \geq k = \tilde{f}$  on  $\partial_1 U^\tau, Nh \leq \inf_{x \in \Omega} Nf(x)$  on  $U^\tau$  and  $\frac{\partial h}{\partial \eta} = \nabla f \cdot \eta = +\infty$  on  $\partial_2 U^\tau$ , where  $\eta$  is the exterior unit normal to  $\partial_2 U^\tau$ . The comparison principle then implies  $\tilde{f} \leq h$  on  $\overline{U^\tau}$ . This yields a contradiction of (22) since (24) implies

$$\lim_{x_2 \downarrow 0} \frac{\partial \tilde{f}}{\partial x_2}(0, x_2) = +\infty,$$

and the facts that  $\tilde{f}(\mathbb{O}) = h(\mathbb{O}), \frac{\partial h}{\partial x_2}(\mathbb{O}) < \infty$  and  $\tilde{f} \leq h$  imply

$$\liminf_{x_2 \downarrow 0} \frac{\partial \tilde{f}}{\partial x_2}(0, x_2) < \infty.$$

This implies (21) holds and completes the proof of Lemma 4.2 except for the construction of the comparison function  $h$ .

Let  $\mathcal{C}$  be a nodary in  $\{x \in \mathbb{R}^2 : x_2 > 0\}$  which, when rotated about the  $x_1$ -axis, generates a nodoid in  $\mathbb{R}^3$  with constant mean curvature  $H_D = |H|_\infty$ , which we assume is positive; if not, set  $H_D = 1$ . (See, for example, [Eells 1987; Mladenov 2002; Rossman 2005] for discussions of Delaunay surfaces and nodoids.) Let the minimal and maximal radii of the nodary be  $r_0$  and  $R_0$  respectively, so that  $r_0 \leq x_2 \leq R_0$  whenever  $(x_1, x_2) \in \mathcal{C}$ ; we will assume  $(0, r_0) \in \mathcal{C}$ . Now let  $\mathcal{D} \subset \mathcal{C}$  be the particular open inner loop of the nodary which contains  $(0, r_0)$  (i.e.,  $(0, r_0) \in \mathcal{D}$  and  $\mathcal{D}$  does not contain endpoints); notice that the unit normal to the nodary at the endpoints of  $\overline{\mathcal{D}}$  are parallel to the axis of rotation of the nodoid and the surface

$$\mathcal{S}_{\mathcal{D}} = \{(x_1, x_2 \cos \theta, x_2 \sin \theta) : (x_1, x_2) \in \mathcal{D}, -\pi \leq \theta \leq 0\}$$

obtained by partially rotating  $\mathcal{D}$  about the  $x_1$ -axis has constant mean curvature  $-H_D$  with respect to its upward unit normal.

Now fix  $t, 0 < t < r_0$ , large enough that  $\Phi_1 = \{(x_1, x_2 + t) : x \in \partial\Omega^\tau \cap R(c_\alpha)\}$  and  $\Phi_2 = \{(x_1, x_2 + t) : x \in \partial\Omega^\tau \cap R(c_{\alpha-\gamma_1})\}$  both intersect  $\mathcal{D}$ . Let  $\Sigma$  denote the component of  $\overline{\Phi_1} \cup \overline{\Phi_2} \setminus \mathcal{D}$  that contains  $(0, t)$  and let  $W$  be the region bounded by  $\Sigma$  and  $\mathcal{D}$ . Set  $W^\tau = \{(x_1, x_2) : (x_1, x_2 + t) \in W\}, \partial_1 W^\tau = \{(x_1, x_2) : (x_1, x_2 + t) \in \Sigma\}$  and  $\partial_2 W^\tau = \partial W^\tau \setminus \partial_1 W^\tau$ . Notice that  $\partial_2 W^\tau \subset \{(x_1, x_2) : (x_1, x_2 + t) \in \mathcal{D}\}$ . Now

define  $h \in C^0(\overline{W^\tau}) \cap C^1(W^\tau \cup \partial_1 W^\tau)$  by

$$h(x_1, x_2) = w(x_1, x_2 + t) - w(0, t) + k$$

for  $x \in \overline{W^\tau}$ , where  $w : \mathcal{D} \rightarrow \mathbb{R}$  such that  $S_{\mathcal{D}}$  is the graph of  $w$ . It follows that  $h$  has the properties mentioned previously.  $\square$

In a similar manner, we can prove each of the following lemmas.

**Lemma 4.3.** *Let  $\Omega$  and  $\gamma$  be as in Section 1. For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{C}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{C}\})$  is a bounded solution of (3)–(4) with  $|H|_\infty < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and either  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_2^-$ , so that  $Rf$  behaves as in Case (D), or  $\alpha \geq \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_1^\pm \cup D_2^-$  and  $Rf$  behaves as in Case (D) or Case (ID). Assume Hypothesis (B+) is true. Then*

$$(26) \quad \lim_{j \rightarrow \infty} \vec{n}(x_j) = (\cos(\alpha + \gamma_1 - 3\pi/2), \sin(\alpha + \gamma_1 - 3\pi/2), 0)$$

for every  $(x_j) \in s(\Omega)$  with  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos \alpha, \sin \alpha)$ .

**Lemma 4.4.** *Let  $\Omega$  and  $\gamma$  be as in Section 1. For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{C}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{C}\})$  is a bounded solution of (3)–(4) with  $|H|_\infty < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and either  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_2^+$ , so that  $Rf$  behaves as in Case (I), or  $\alpha \geq \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_1^\pm \cup D_2^+$  and  $Rf$  behaves as in Case (I) or Case (ID). Assume Hypothesis (B-) is true. Then*

$$(27) \quad \lim_{j \rightarrow \infty} \vec{n}(x_j) = (\cos(-\alpha - \gamma_1 + 3\pi/2), \sin(-\alpha - \gamma_1 + 3\pi/2), 0)$$

for every  $(x_j) \in s(\Omega)$  with  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos(-\alpha), \sin(-\alpha))$ .

**Lemma 4.5.** *Let  $\Omega$  and  $\gamma$  be as in Section 1. For some  $\rho \in (0, 1)$  and  $\delta > 0$ , suppose  $f \in C^2(\Omega) \cap C^{1,\rho}(B(\mathbb{C}, \delta) \cap \overline{\Omega} \setminus \{\mathbb{C}\})$  is a bounded solution of (3)–(4) with  $|H|_\infty < \infty$ . Suppose (5) holds,  $\gamma_1, \gamma_2 \in (0, \pi)$  and either  $\alpha < \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_2^-$ , so that  $Rf$  behaves as in Case (D), or  $\alpha \geq \pi/2$  and  $(\gamma_1, \gamma_2)$  is in  $D_1^\pm \cup D_2^-$  and  $Rf$  behaves as in Case (D) or Case (DI). Assume Hypothesis (B-) is true. Then*

$$(28) \quad \lim_{j \rightarrow \infty} \vec{n}(x_j) = (\cos(-\alpha + \gamma_1 - \pi/2), \sin(-\alpha + \gamma_1 - \pi/2), 0)$$

for every  $(x_j) \in s(\Omega)$  with  $\lim_{j \rightarrow \infty} x_j/|x_j| = (\cos(-\alpha), \sin(-\alpha))$ .

### References

[Concus and Finn 1996] P. Concus and R. Finn, “Capillary wedges revisited”, *SIAM J. Math. Anal.* **27**:1 (1996), 56–69. MR 96m:76006 Zbl 0843.76012

[Eells 1987] J. Eells, “The surfaces of Delaunay”, *Math. Intelligencer* **9**:1 (1987), 53–57. MR 88h:53011 Zbl 0605.53002

[Finn 1986] R. Finn, *Equilibrium capillary surfaces*, Grundlehren der Mathematischen Wissenschaften **284**, Springer, New York, 1986. MR 88f:49001 Zbl 0583.35002



- [Finn 1988a] R. Finn, “Comparison principles in capillarity”, pp. 156–197 in *Partial differential equations and calculus of variations*, edited by S. Hildebrandt and R. Leis, Lecture Notes in Math. **1357**, Springer, Berlin, 1988. MR 90d:53010 Zbl 0692.35006
- [Finn 1988b] R. Finn, “Moon surfaces, and boundary behaviour of capillary surfaces for perfect wetting and nonwetting”, *Proc. London Math. Soc.* (3) **57**:3 (1988), 542–576. MR 89m:49076 Zbl 0668.76019
- [Finn 1996] R. Finn, “Local and global existence criteria for capillary surfaces in wedges”, *Calc. Var. Partial Differential Equations* **4**:4 (1996), 305–322. MR 97f:53005 Zbl 0872.76017
- [Finn 1999] R. Finn, “Capillary surface interfaces”, *Notices Amer. Math. Soc.* **46**:7 (1999), 770–781. MR 2000g:76033 Zbl 1194.76049
- [Finn 2002a] R. Finn, “Eight remarkable properties of capillary surfaces”, *Math. Intelligencer* **24**:3 (2002), 21–33. MR 2003f:76041
- [Finn 2002b] R. Finn, “Some properties of capillary surfaces”, *Milan J. Math.* **70** (2002), 1–23. MR 2003m:53012 Zbl 1053.76009
- [Jeffres and Lancaster 2007] T. Jeffres and K. Lancaster, “Vertical blow ups of capillary surfaces in  $\mathbb{R}^3$ . I. Convex corners”, *Electron. J. Differential Equations* **152** (2007), 24 pp. MR 2009b:49103 Zbl 1139.49041
- [Jeffres and Lancaster 2008] T. Jeffres and K. Lancaster, “Vertical blow ups of capillary surfaces in  $\mathbb{R}^3$ . II. Nonconvex corners”, *Electron. J. Differential Equations* **160** (2008), 25 pp. MR 2009j:49098 Zbl 1163.49044
- [Lancaster 2010] = [L] K. E. Lancaster, “A proof of the Concus–Finn conjecture”, *Pacific J. Math.* **247**:1 (2010), 75–108. MR 2011m:53011 Zbl 05809032
- [Lancaster and Siegel 1996a] = [LS a] K. E. Lancaster and D. Siegel, “Existence and behavior of the radial limits of a bounded capillary surface at a corner”, *Pacific J. Math.* **176**:1 (1996), 165–194. MR 98g:58030a Zbl 0866.76018
- [Lancaster and Siegel 1996b] = [LS b] K. E. Lancaster and D. Siegel, “Behavior of a bounded non-parametric  $H$ -surface near a reentrant corner”, *Z. Anal. Anwendungen* **15**:4 (1996), 819–850. MR 97m:53011 Zbl 0866.35046
- [Mladenov 2002] I. M. Mladenov, “Delaunay surfaces revisited”, *C. R. Acad. Bulgare Sci.* **55**:5 (2002), 19–24. MR 2003g:53005 Zbl 1017.53010
- [Rossman 2005] W. Rossman, “The first bifurcation point for Delaunay nodoids”, *Experiment. Math.* **14**:3 (2005), 331–342. MR 2006j:53011 Zbl 1092.53010
- [Shi 2006] D. Shi, “Capillary surfaces at a reentrant corner”, *Pacific J. Math.* **224**:2 (2006), 321–353. MR 2008a:58013 Zbl 1125.76013
- [Tam 1986] L.-F. Tam, “Regularity of capillary surfaces over domains with corners: borderline case”, *Pacific J. Math.* **124**:2 (1986), 469–482. MR 87k:49049 Zbl 0604.49029

Received July 12, 2010. Revised July 8, 2012.

KIRK E. LANCASTER  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
WICHITA STATE UNIVERSITY  
WICHITA, KS 67260-0033  
UNITED STATES  
lancaster@math.wichita.edu

# PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by  
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [pacificmath.org](http://pacificmath.org) for submission instructions.

---

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2012 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 258    No. 2    August 2012

---

Uniqueness theorems for CR and conformal mappings YOUNG-JUN CHOI and JAE-CHEON JOO	257
Some finite properties for vertex operator superalgebras CHONGYING DONG and JIANZHI HAN	269
On the geometric flows solving Kählerian inverse $\sigma_k$ equations HAO FANG and MIJIA LAI	291
An optimal anisotropic Poincaré inequality for convex domains GUOFANG WANG and CHAO XIA	305
Einstein metrics and exotic smooth structures MASASHI ISHIDA	327
Noether's problem for $\hat{S}_4$ and $\hat{S}_5$ MING-CHANG KANG and JIAN ZHOU	349
Remarks on the behavior of nonparametric capillary surfaces at corners KIRK E. LANCASTER	369
Generalized normal rulings and invariants of Legendrian solid torus links MIKHAIL LAVROV and DAN RUTHERFORD	393
Classification of singular $\mathbb{Q}$ -homology planes II: $\mathbb{C}^1$ - and $\mathbb{C}^*$ -rulings. KAROL PALKA	421
A dynamical interpretation of the profile curve of CMC twizzler surfaces OSCAR M. PERDOMO	459
Classification of Ising vectors in the vertex operator algebra $V_L^+$ HIROKI SHIMAKURA	487
Highest-weight vectors for the adjoint action of $GL_n$ on polynomials RUDOLF TANGE	497



0030-8730(201208)258:2;1-2