

*Pacific
Journal of
Mathematics*

**ON A GALOIS CONNECTION
BETWEEN THE SUBFIELD LATTICE
AND THE MULTIPLICATIVE SUBGROUP LATTICE**

JOHN K. MCVEY

Volume 264 No. 1

July 2013

ON A GALOIS CONNECTION BETWEEN THE SUBFIELD LATTICE AND THE MULTIPLICATIVE SUBGROUP LATTICE

JOHN K. MCVEY

Given finite fields $F < E$, we present a collection of subgroups $C \leq E^\times$ and establish, to each C , a Galois connection between the intermediate field lattice $\mathcal{C} = \{L \mid F \leq L \leq E\}$ and C 's subgroup lattice. Our main result is that, in all but an extremely limited and completely determined family, the closed subset of \mathcal{C} is \mathcal{C} itself, establishing a natural bijection between \mathcal{C} and the lattice $\{L \cap C \mid L \in \mathcal{C}\}$. As an application, we use this bijection to calculate the set of degrees for the complex-valued irreducible representations of the split extension $C \rtimes \text{Gal}(E/F)$.

1. Introduction

In §3 of [McVey 2004], generalizing results in §5 of [Riedl 1999], we worked towards (among other things) a better understanding of the groups $C \rtimes \text{Gal}(E/F)$ for finite fields $F < E$, where $C < E^\times$ is the subgroup of order $|E^\times : F^\times|$. While working to generalize those results further, we discovered a Galois connection which itself is worthy of further study. This paper's intent is to record the Galois connection as well as the research that motivated its initial study. The primary assertion of the Main Theorem is that, but for a completely determined and rather limited family, the intermediate field lattice $\mathcal{C} = \{L \mid F \leq L \leq E\}$ is itself one of the two closed subsets in the Galois connection, thereby determining a canonical bijection between \mathcal{C} and the other closed set $\{L \cap C \mid L \in \mathcal{C}\}$. As to the motivating research, we use this bijection to calculate the degrees of the irreducible complex representations of the aforementioned split extension $C \rtimes \text{Gal}(E/F)$, showing every integer allowed by Itô's theorem is a degree.

2. Towards the Galois connection

Our focus in this paper is on monotone Galois connections. To avoid confusion between monotone and antitone connections, we define the term and present the basic relevant results. Two monotone nondecreasing functions $f : \mathcal{A} \rightarrow \mathcal{B}$ and $g : \mathcal{B} \rightarrow \mathcal{A}$ on partially ordered sets (\mathcal{A}, \leq) and (\mathcal{B}, \leq) form a *monotone Galois*

MSC2010: primary 06A15; secondary 20C15.

Keywords: Galois correspondence, lattice, character degree, finite field.

connection if

$$f(a) \leq b \iff a \leq g(b)$$

over all $a \in \mathcal{A}$ and $b \in \mathcal{B}$. The function f is the *lower adjoint* and g is the *upper adjoint*. The *closed sets* \mathcal{A}_0 and \mathcal{B}_0 of \mathcal{A} and \mathcal{B} respectively are defined by $\mathcal{A}_0 = g(\mathcal{B})$ and $\mathcal{B}_0 = f(\mathcal{A})$, and satisfy

$$\begin{aligned} \mathcal{A}_0 &= \{a \in \mathcal{A} \mid g \circ f(a) = a\} = g \circ f(\mathcal{A}), \\ \mathcal{B}_0 &= \{b \in \mathcal{B} \mid f \circ g(b) = b\} = f \circ g(\mathcal{B}). \end{aligned}$$

The functions f and g are inverse bijections between the sets \mathcal{A}_0 and \mathcal{B}_0 .

Turning now specifically to our setting of finite fields $F < E$, label by π the set of primes which divide $|F^\times| = |F| - 1$. The collection of groups to which the Galois connection applies consists of all subgroups $C \leq E^\times$ for which the index $|E^\times : C|$ is a π -number (thus naturally generalizing results in [Riedl 1999] where hypotheses guaranteed $|E^\times : C| = |F^\times|$). Fixing a group C , the upper adjoint is very easy to describe; it is the function “intersect with C ”.

As to the lower adjoint, define the F -closure \widehat{X} of a subset $X \subseteq E$ to be the smallest subfield of E which contains $X \cup F$. In other words, \widehat{X} is the intersection of all fields L satisfying $X \cup F \subseteq L \leq E$. It should be obvious that F -closure actually is a closure operator (i.e.,

$$\widehat{X} \supseteq X \quad \text{and} \quad \widehat{\widehat{X}} = \widehat{X}$$

over all subsets $X \subseteq E$), and that a Galois automorphism $\sigma \in \text{Gal}(E/F)$ centralizes X if and only if it centralizes \widehat{X} . The partially ordered sets in our Galois connection are the lattices

$$(1) \quad \mathcal{L} = \{L \mid F \leq L \leq E \text{ is a field}\} \quad \text{and} \quad \mathcal{C} = \{D \mid D \leq C \text{ is a group}\},$$

ordered by inclusion. The functions $X \mapsto X \cap C$ and $X \mapsto \widehat{X}$ are obviously monotone. Given $D \in \mathcal{C}$ and $L \in \mathcal{L}$, and noting that $L \cap C = L^\times \cap C \in \mathcal{C}$, we have

$$\widehat{D} \subseteq L \iff D \subseteq L \iff D \subseteq L \cap C,$$

showing that $\widehat{\cdot}$ is a lower adjoint while $(\cdot) \cap C$ is an upper adjoint. Therefore, as $\mathcal{A}_0 = g(\mathcal{B})$, the closed subset of \mathcal{L} is $\mathcal{L}_0 = \{L \cap C \mid L \in \mathcal{L}\}$.

We are now ready to state the Main Theorem. All but the last two sentences were proven in the above discussion. Those last two sentences are the true content of the theorem, and their proof is at the end of this section.

Main Theorem. *Let $F < E$ be finite fields and label by π the set of primes dividing $|F| - 1$. Let C be a subgroup of E^\times whose index $|E^\times : C|$ is a π -number. Given the partially ordered sets defined by (1), the functions $\widehat{\cdot} : \mathcal{C} \rightarrow \mathcal{L}$ and $(\cdot) \cap C : \mathcal{L} \rightarrow \mathcal{C}$*

are respectively the lower and upper adjoints of a monotone Galois connection, and thus provide inverse bijections between the closed subsets $\mathcal{C}_0 \subseteq \mathcal{C}$ and $\mathcal{E}_0 \subseteq \mathcal{E}$. The closed subset \mathcal{C}_0 of \mathcal{C} is the lattice $\mathcal{C}_0 = \{L \cap C \mid L \in \mathcal{E}\}$. If $|F|$ is a Mersenne prime, $|E : F|$ is even, and 4 does not divide $|C|$, then the closed subset \mathcal{E}_0 of \mathcal{E} is the set $\mathcal{E}_0 = \mathcal{E} \setminus \{K\}$ where $|K : F| = 2$. Otherwise, $\mathcal{E} = \mathcal{E}_0$.

Our argument for the as yet unproven portion of the Main Theorem relies fundamentally on number theory. We ask the reader to recall Zsigmondy’s prime theorem, as it is the foundation for what follows.

Theorem 1 [Zsigmondy 1892]. *Let a, b, n be positive integers and assume a, b are coprime and not both 1. Then, $a^n - b^n$ has a prime divisor which does not divide $a^k - b^k$ for integers $0 < k < n$, except when either*

$$n = 6 \text{ and } \{a, b\} = \{1, 2\} \quad \text{or} \quad n = 2 \text{ and } a + b \text{ is a 2-power.}$$

Aside from specifying $\{a, b\}$ as $\{q, 1\}$ with q a prime-power, the main point behind Corollary 2 is that the order of the quantifiers changed (from ‘ \exists prime $\forall k$ ’ in Zsigmondy’s theorem to ‘ $\forall k \exists$ prime’ in the corollary).

Corollary 2. *Let $n > 1$ be an integer and q a power of a prime. For each integer k with $0 < k < n$, there is a prime which divides $q^n - 1$ and not $q^k - 1$, except when q is a Mersenne prime and $n = 2$. Conversely, when q is a Mersenne prime, every prime dividing $q^2 - 1$ divides $q - 1$.*

Proof. As stated previously, Zsigmondy’s theorem provides a universal prime (over all k) unless we are in one of the exceptional cases. First, assume $n = 6$ and $q = 2$, in which case $q^n - 1 = 2^6 - 1 = 63 = 3^2 \cdot 7$. It suffices to check that 3 divides none of $1 = 2^1 - 1$, $7 = 2^3 - 1$, and $31 = 2^5 - 1$, and that 7 divides neither $3 = 2^2 - 1$ nor $15 = 2^4 - 1$.


In the other exceptional case, $n = 2$ and $q + 1$ is a 2-power. However, Catalan’s conjecture (proven in [Mihăilescu 2004]) says that the integer equation $x^a - y^b = 1$ with $a, b > 1$ only has the solution $3^2 - 2^3$. Because $q + 1$ is a 2-power already, q itself must be prime, hence a Mersenne prime.

As to the converse, when q is a Mersenne prime, the only prime dividing $q + 1$ is 2, which necessarily divides $(q + 1) - 2 = q - 1$. As $q^2 - 1 = (q + 1)(q - 1)$, the result follows. □

We now leave number theory and move to algebra proper. Our first algebraic goal is a lemma which shows how the number theory embedded in the previous corollary can be applied to finite fields.

Lemma 3. *Let $F \leq K \leq L \leq E$ be finite fields. For the set π of prime divisors of $|F^\times|$, let C be a subgroup of E^\times whose index is a π -number. If the prime p divides $|L^\times|$ and not $|K^\times|$, then p divides $|L \cap C : K \cap C|$.*

Proof. The following picture provides insight into this proof.



That p does not divide $|K^\times|$ implies p does not divide $|F^\times| = q - 1$. Since

$$|L^\times : L \cap C| = |L^\times C : C| \text{ divides } |E^\times : C|,$$

which is a π -number and thus coprime to p , necessarily p divides $|L \cap C|$. As p does not divide $|K^\times|$, it also does not divide $|K \cap C|$. \square

Theorem 4. *Let q be a prime-power, $e > 1$ an integer, and π the set of primes dividing $q - 1$. Label $F = \mathbb{F}_q$ and $E = \mathbb{F}_{q^e}$, and let C be a subgroup of E^\times whose index $|E^\times : C|$ is a π -number. Then, for all fields $F \leq L \leq E$, the equality $L = \widehat{L \cap C}$ holds, except when the following conditions are all satisfied.*

- (1) q is a Mersenne prime.
- (2) e is even.
- (3) $L = \mathbb{F}_{q^2}$.
- (4) 4 does not divide $|C|$.

When these simultaneously hold, $L \cap C = F \cap C$, so $\widehat{L \cap C} = F < L$.

Proof. Fix the field L . Obviously, the set $L \cap C$ is a subset both of $\widehat{L \cap C}$ and of C . Therefore,


$$L \cap C \subseteq \widehat{L \cap C} \cap C \subseteq \widehat{L \cap C} = L \cap C,$$

and we have equality throughout. Applying (the contrapositive of) Lemma 3 to $K = \widehat{L \cap C}$, the equality $K \cap C = L \cap C$ shows that every prime dividing $|L^\times|$ divides $|K^\times|$. Labelling $|L^\times| = q^n - 1$ and $|K^\times| = q^k - 1$, either $k = n$ (and we are done) or we are in the exceptional case of Corollary 2.

Henceforth, assume $n = 2$ and q is a Mersenne prime. As e is a multiple of n , it is even. Write $q = 2^p - 1$. Consequently,

- $|F^\times| = q - 1 = 2(2^{p-1} - 1)$, which has 2-part exactly 2, while
- $|L^\times| = q^2 - 1 = 2^p |F^\times|$, which has 2-part 2^{p+1} .

In particular, the $2'$ -part of $|L^\times|$ is exactly the $2'$ -part of $|F^\times|$.



We now split the argument as to whether or not 4 divides $|C|$. If 4 divides $|C|$, then because 4 also divides $|L^\times|$, it divides $|L \cap C|$. However, 4 does not divide $|F^\times|$, so $L \cap C \not\leq F^\times$. Accordingly, $F < \widehat{L \cap C} \leq L$, and $F = \mathbb{F}_q$ being a maximal subfield of $L = \mathbb{F}_{q^2}$ shows $L = \widehat{L \cap C}$.

When 4 does not divide $|C|$, the 2-part of $|L \cap C|$ divides $|F^\times|$. Generally, the $2'$ -part of $|L \cap C|$ divides $|L^\times|_{2'} = |F^\times|_{2'}$. It follows that $|L \cap C|$ divides $|F^\times|$. Since E^\times is cyclic, this shows $L \cap C \leq F^\times$, so $L \cap C = F \cap C$. \square

With the above result in place, we use that $\mathcal{B}_0 = f \circ g(\mathcal{B})$ to conclude

$$\mathcal{E}_0 = \{\widehat{L \cap C} \mid L \in \mathcal{E}\}.$$

Meanwhile, the F -closure $\widehat{L \cap C}$ equals L but for the one exception \mathbb{F}_{q^2} when q is Mersenne, e is even, and 4 fails to divide $|C|$. This finishes the proof of the Main Theorem.

3. Application to degrees

Our concluding section presents the computations for the character degree set of the split extension $C \rtimes \text{Gal}(E/F)$ when $\mathcal{E} = \mathcal{E}_0$. We emphasize once more that this result was the principal impetus for our study of this Galois connection. All standard notations and conventions regarding character theory are taken from [Isaacs 1976]. The following generalizes Theorem 3.2 in [McVey 2004], and the proof here is fundamentally the same as is presented there, the main modification being the use of Theorem 4.

Theorem 5. *Fix a prime-power q and an exponent $1 < e \in \mathbb{Z}$, and label by F the field \mathbb{F}_q , by E the field \mathbb{F}_{q^e} , and by π the set of primes dividing $q - 1$. Let $\Gamma = \text{Gal}(E/F)$, and fix $C \leq E^\times$ under the assumption $|E^\times : C|$ is a π -number. If q is Mersenne and e is even, assume 4 divides $|C|$. Then, Γ normalizes C and*

$$\text{cd}(C\Gamma) = \{n \mid n \text{ divides } e\}.$$

Proof. Because E^\times is cyclic, every subgroup is characteristic. In particular, C is fixed (setwise) under every field automorphism of E , so Γ normalizes C . As C is cyclic, $\text{Irr}(C)$ contains only linear characters and forms a cyclic group under multiplication. Let $\lambda \in \text{Irr}(C)$ be a generator, noting λ is both faithful and a homomorphism. In summary, $\lambda(d_1) = \lambda(d_2)$ implies $d_1 = d_2$, $\lambda^m(d) = \lambda(d^m)$,

and $\lambda^\tau(d^\tau) = \lambda(d)$ for all $d, d_1, d_2 \in C$, $m \in \mathbb{Z}$, and $\tau \in \Gamma$. Recalling $C \triangleleft C\Gamma$ is abelian, Itô's theorem says every degree in $\text{cd}(C\Gamma)$ divides $|C\Gamma : C| = |\Gamma| = e$.

Conversely, fix a divisor n of e , and we will demonstrate an irreducible character of $C\Gamma$ whose degree is n . Let σ be a generator of Γ , and label $\Phi = \langle \sigma^n \rangle$, observing that $|\Gamma : \Phi| = n$. Let L be the fixed field for σ^n in the (usual) Galois correspondence for E over F . Hence, $\Phi = \text{Gal}(E/L)$ and σ^n fixes the subgroup $L \cap C$ of C . For some generator $c \in C$, let $L \cap C = \langle c^m \rangle$.

We claim the stabilizer of λ^m in Γ is Φ . Given the claim, the stabilizer of λ^m in $C\Gamma$ is $C\Phi$, and λ^m extends to a character $\varphi \in \text{Irr}(C\Phi)$ through for example Corollary 11.22 in [Isaacs 1976]. Also, φ induces irreducibly to $C\Gamma$ by Clifford correspondence. Therefore,

$$n = |\Gamma : \Phi| = |C\Gamma : C\Phi| = |C\Gamma : C\Phi|\varphi(1) = \varphi^{C\Gamma}(1) \in \text{cd}(C\Gamma).$$

As n was an arbitrary divisor of e , we will have shown the result.

Given $\tau \in \Gamma$ and recalling λ is faithful, the equalities

$$(\lambda^m)^\tau(d) = \lambda^m(d^{\tau^{-1}}) = \lambda((d^{\tau^{-1}})^m) = \lambda((d^m)^{\tau^{-1}})$$

and

$$\lambda^m(d) = \lambda(d^m)$$

imply that τ centralizes λ^m (the left ends are equal) if and only if τ^{-1} centralizes d^m for every $d \in C$ (the right ends are equal). The latter happens exactly when τ^{-1} centralizes $\langle c^m \rangle = L \cap C$, which occurs if and only if τ^{-1} centralizes $\widehat{L \cap C}$. As $L = \widehat{L \cap C}$ (Theorem 4), this is equivalent to $\tau^{-1} \in \text{Gal}(E/L) = \Phi$. \square

In closing, we would be remiss in not mentioning an application of Theorem 5 to a remark made in [Lewis 2001]. For the subsequent, we use the notation of [Lewis 2001]. In the paragraph preceding Lemma 3.4, Dr. Lewis made the comment that "... every divisor of m occurs in $\text{cd}(G/V)$ ", but that particular conclusion was superfluous to Lemma 3.4, so it went unproven. Reading through the first two and a half paragraphs of that proof, V can be viewed as the additive group of the field \mathbb{F}_{q^m} , K/Z acts on \mathbb{F}_{q^m} by multiplication as if it were a subgroup of $\mathbb{F}_{q^m}^\times$, and the quotient H/K behaves as a Galois group. Lastly, the hypotheses to Example 2.4 imply m is coprime to $(q^m - 1)/(q - 1)$. Hence, our result applies to the group H/Z , and Lewis' claim about the degrees is an immediate corollary of Theorem 5 and the relations $G/V \cong H$ and $\text{cd}(H) = \text{cd}(H/Z)$.

References

- [Isaacs 1976] I. M. Isaacs, *Character theory of finite groups*, Pure and Applied Mathematics **69**, Academic Press, New York, 1976. MR 57 #417 Zbl 0337.20005
- [Lewis 2001] M. L. Lewis, "Solvable groups whose degree graphs have two connected components", *J. Group Theory* **4:3** (2001), 255–275. MR 2002g:20015 Zbl 0998.20009

- [McVey 2004] J. K. McVey, “Prime divisibility among degrees of solvable groups”, *Comm. Algebra* **32**:9 (2004), 3391–3402. MR 2005i:20012 Zbl 1077.20009
- [Mihăilescu 2004] P. Mihăilescu, “Primary cyclotomic units and a proof of Catalan’s conjecture”, *J. Reine Angew. Math.* **572** (2004), 167–195. MR 2005f:11051 Zbl 1067.11017
- [Riedl 1999] J. M. Riedl, “Character degrees, class sizes, and normal subgroups of a certain class of p -groups”, *J. Algebra* **218**:1 (1999), 190–215. MR 2000f:20025 Zbl 0932.20015
- [Zsigmondy 1892] K. Zsigmondy, “Zur Theorie der Potenzreste”, *Monatsh. Math. Phys.* **3**:1 (1892), 265–284. MR 1546236 JFM 24.0176.02

Received June 2, 2012. Revised December 10, 2012.

JOHN K. MCVEY
DEPARTMENT OF MATHEMATICS
KENT STATE UNIVERSITY
MATHEMATICS AND COMPUTER SCIENCE BUILDING 233
SUMMIT STREET
KENT, OH 44242
UNITED STATES
jmcvey@math.kent.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 264 No. 1 July 2013

On the center of fusion categories	1
ALAIN BRUGUIÈRES and ALEXIS VIRELIZIER	
Connected quandles associated with pointed abelian groups	31
W. EDWIN CLARK, MOHAMED ELHAMDADI, XIANG-DONG HOU, MASAHICO SAITO and TIMOTHY YEATMAN	
Entropy and lowest eigenvalue on evolving manifolds	61
HONGXIN GUO, ROBERT PHILIPOWSKI and ANTON THALMAIER	
Poles of certain residual Eisenstein series of classical groups	83
DIHUA JIANG, BAIYING LIU and LEI ZHANG	
Harmonic maps on domains with piecewise Lipschitz continuous metrics	125
HAIGANG LI and CHANGYOU WANG	
q -hypergeometric double sums as mock theta functions	151
JEREMY LOVEJOY and ROBERT OSBURN	
Monic representations and Gorenstein-projective modules	163
XIU-HUA LUO and PU ZHANG	
Helicoidal flat surfaces in hyperbolic 3-space	195
ANTONIO MARTÍNEZ, JOÃO PAULO DOS SANTOS and KETI TENENBLAT	
On a Galois connection between the subfield lattice and the multiplicative subgroup lattice	213
JOHN K. MCV EY	
Some characterizations of Campanato spaces via commutators on Morrey spaces	221
SHAOGUANG SHI and SHANZHEN LU	
The Siegel–Weil formula for unitary groups	235
SHUNSUKE YAMANA	



0030-8730(201307)264:1;1-4