

*Pacific  
Journal of  
Mathematics*

**TWISTED K-THEORY FOR THE ORBIFOLD  $[*/G]$**

MARIO VELÁSQUEZ, EDWARD BECERRA AND HERMES MARTINEZ

## TWISTED K-THEORY FOR THE ORBIFOLD $[*/G]$

MARIO VELÁSQUEZ, EDWARD BECERRA AND HERMES MARTINEZ

*The second author dedicates this paper to Hesel María*

The main result of this paper establishes an explicit ring isomorphism between the twisted orbifold K-theory  ${}^\omega K_{\text{orb}}([*/G])$  and  $R(D^\omega(G))$  for any element  $\omega \in Z^3(G; S^1)$ . We also study the relation between the twisted orbifold K-theories  ${}^\alpha K_{\text{orb}}(\mathcal{X})$  and  ${}^{\alpha'} K_{\text{orb}}(\mathcal{Y})$  of the orbifolds  $\mathcal{X} = [*/G]$  and  $\mathcal{Y} = [*/G']$ , where  $G$  and  $G'$  are different finite groups, and  $\alpha \in Z^3(G; S^1)$  and  $\alpha' \in Z^3(G'; S^1)$  are different twistings. We prove that if  $G'$  is an extraspecial group with prime number  $p$  as an index and order  $p^n$  (for some fixed  $n \in \mathbb{N}$ ), under a suitable hypothesis over the twisting  $\alpha'$  we can obtain a twisting  $\alpha$  on the group  $(\mathbb{Z}_p)^n$  such that there exists an isomorphism between the twisted K-theories  ${}^{\alpha'} K_{\text{orb}}([*/G'])$  and  ${}^\alpha K_{\text{orb}}([*/(\mathbb{Z}_p)^n])$ .

### 1. Introduction

The twisted K-theory is a successful example of the increasing flow of physical ideas into mathematics. Brought from the physical setup, the twisted orbifold K-theory has been, for the last twenty-five years, a fruitful field of ideas and development in K-theory and algebraic topology. It emerged from two sources: the consideration of the D-brane charge on a smooth manifold by Witten [1998], and the concept of discrete torsion on an orbifold by Vafa [2001]. Although for any element  $\alpha \in H^3(\mathcal{X}; \mathbb{Z})$  one can associate the twisted K-theory  ${}^\alpha K(\mathcal{X})$ , its structure is simpler if the element  $\alpha$  lies in the image of the pullback associated to the map  $X \rightarrow *$ . In such a case, we call this element a *discrete torsion* since we can see it as an element in the cohomology  $H^3(G; \mathbb{Z})$ .

On the other hand, an orbifold is a type of generalization of a smooth manifold. It is a topological space locally modeled as a quotient of a manifold by an action of a finite group. When  $X$  represents an orbifold, the twisted K-theory is more

---

Velásquez was partially supported by Colciencias through the grant Becas Generación del Bicentenario #494, Fundación Mazda para el Arte y la Ciencia, and Prof. Wolfgang Lück through his Leibniz Prize. We want to express our gratitude to Professor Bernardo Uribe for his important suggestions and ideas for our work.

MSC2010: primary 19L47, 19L50; secondary 20J06, 20C25.

Keywords: inverse transgression map, twisted double Drinfeld, twisted K-theory.

interesting because it is naturally related to equivariant theories if we specialize in the orbifold  $\mathcal{X} = [X/G]$ , where  $X$  is a smooth manifold and  $G$  is a compact Lie group acting almost freely on  $X$ . In the case of orbifolds, we have another important advantage to work with; it is the cohomological counterpart given by the Chen–Ruan cohomology on orbifolds  $H_{CR}^*(\mathcal{X}; \mathbb{C})$  related to K-theory by the Chern character. The Chen–Ruan cohomology of orbifolds has an interesting nontrivial internal product which makes it an algebra. This product can be presented in the setup of the K-theory to obtain a stringy product on the K-theory of the orbifold  $K_{\text{orb}}(\mathcal{X})$  (see [Becerra and Uribe 2009; Adem and Ruan 2003]). If the orbifold considered has the form  $[X/G]$ , then the twisted orbifold K-theory can be related to the equivariant K-theory of the spaces of fixed points by the  $G$ -action on  $X$ .

On the other hand, the tensor product defines a product

$${}^{\alpha}K_{\text{orb}}(\mathcal{X}) \otimes {}^{\beta}K_{\text{orb}}(\mathcal{X}) \rightarrow {}^{\alpha+\beta}K_{\text{orb}}(\mathcal{X})$$

for any pair of elements  $\alpha$  and  $\beta$  in  $H^3(\mathcal{X}; \mathbb{Z})$ . In fact, one can obtain a stringy product for the twisted K-theory on orbifolds by using the stringy product defined on each space of fixed points to define an explicit stringy product in each  ${}^{\alpha}K(\mathcal{X})$  for any  $\alpha \in H^3(\mathcal{X}; \mathbb{Z})$ . Nevertheless, the crucial information to define the stringy product on the twisted K-theory of orbifolds does not lie in  $H^3(\mathcal{X}; \mathbb{Z})$ ; instead it lies in  $H^4(\mathcal{X}; \mathbb{Z})$ . Given an element  $\phi$  in  $H^4(\mathcal{X}; \mathbb{Z})$ , it defines an element  $\theta(\phi)$  in  $H^3(\wedge X; \mathbb{Z})$ , where  $\wedge \mathcal{X}$  is the inertia orbifold associated to  $\mathcal{X}$ . Hence, we can define a stringy product over the twisted K-theory orbifold  ${}^{\theta(\phi)}K_{\text{orb}}(\wedge \mathcal{X})$  by using a suitable structure of the inertia orbifold  $\wedge X$ . One such product structure is based on the map  $\theta(\phi)$  called the *inverse transgression map*, which is considered to be the inverse of the classical transgression map.

For this paper, the stringy product in  $K_{\text{orb}}(\mathcal{X})$  has a trivial expression as we will restrict our observations to the case in which  $\mathcal{X} = [*/G]$ , where  $G$  is a finite group.

The main result in this paper is to present an explicit relation between the twisted Drinfeld double  $D^{\omega}(G)$  and the twisted orbifold K-theory  ${}^{\omega}K_{\text{orb}}([*/G])$  for an element  $\omega$  of discrete torsion (see Section 4). This allows us to relate the twisted orbifold K-theories  ${}^{\omega}K_{\text{orb}}([*/G])$  and  ${}^{\omega'}K_{\text{orb}}([*/G'])$  for two orbifolds  $[*/G]$  and  $[*/G']$  with the twistings  $\omega \in H^4(G; \mathbb{Z})$  and  $\omega' \in H^4(G'; \mathbb{Z})$ . To obtain such a relation, we modify the stringy product defined in [Adem et al. 2007] by an element in  $R_{\alpha_g}(C(g) \cap C(h))$ .

## 2. Pushforward map in the twisted representation ring

In this section, we introduce the pushforward map. Although this map can be defined for almost complex manifolds, we will focus only on the case of homogeneous

spaces  $G/H$ . To define this map, let us recall the Thom isomorphism theorem in equivariant K-theory.

**Fact [Segal 1968, Proposition 3.2].** *Let  $X$  be a compact  $G$ -manifold and  $p : E \rightarrow X$  be a complex  $G$ -vector bundle over  $X$ . There exists an isomorphism*

$$\phi : K_G^*(X) \rightarrow K_G^*(E, E \setminus E_0), \quad \phi([F]) := p^*(F) \otimes \lambda_{-1}(E),$$

where  $E_0$  is the zero section and the class  $\lambda_{-1}(E)$  is the Thom class associated to  $[E]$ .

**Remark 2.1.** We need to recall how to define the *normal bundle*. If  $M$  and  $N$  are  $G$ -manifolds (that means a manifold with a smooth  $G$ -action) and  $f : M \rightarrow N$  is a  $G$ -embedding, we can define a (real) vector bundle  $\tau$  such that  $df(TM) \oplus \tau \cong TN$  (for details in this construction, consult [tom Dieck 1987]). If the map  $f$  is not a  $G$ -embedding, we can consider  $f : M \rightarrow N \times D^j$  ( $D^j$  is the unitary disk in  $\mathbb{R}^j$  with the trivial  $G$ -action) for sufficiently large  $j$  and by Corollary 1.10 [Wasserman 1969] we can approximate  $f$  by an immersion  $g_f$ , then we define the normal bundle for  $f$  as the normal bundle of  $g_f$ .

Now, we proceed to define the pushforward map  $f_* : K_G^*(X) \rightarrow K_G^*(Y)$  for a differentiable map  $f : X \rightarrow Y$  between almost complex  $G$ -manifolds by letting  $\tau$  be the normal bundle associated to the map  $f : X \rightarrow Y$ . We define the pushforward, which will be denoted by  $f_*$ , as the composition

$$K_G^*(X) \xrightarrow{\phi} K_G^*(\tau, \tau \setminus \tau_0) \xrightarrow{j} K_G^*(Y \times D^j, (Y \times D^j) \setminus g_f(X)) \xrightarrow{i_{\sharp}} K_G^*(Y \times D^j) \cong K_G^*(Y),$$

where  $\phi$  is the Thom isomorphism. The map  $j$  is given by excision, the map  $i_{\sharp}$  is the pullback map induced by the inclusion, and the last isomorphism is induced by the natural inclusion. The pushforward map can be defined also in the twisted case (see [Carey and Wang 2008]). Consider the following diagram of inclusions:

$$\begin{array}{ccc} G & \longleftarrow & K \\ \uparrow & & \uparrow \\ H & \longleftarrow & H \cap K \end{array}$$

from which we get a diagram of surjections:

$$\begin{array}{ccc} G/G & \xleftarrow{i_1} & G/K \\ \uparrow j_1 & & \uparrow j_2 \\ G/H & \xleftarrow{i_2} & G/(H \cap K) \end{array}$$

Using this diagram we obtain the map

$$j_{2*} \circ i_2^* : K_G^*(G/H) \rightarrow K_G^*(G/K), \quad [E] \mapsto [\lambda_{-1}(\tau_{j_2}) \otimes i_2^*(p^*(E))],$$

where  $\tau_{j_2}$  is the normal bundle of  $j_2$ , and the map

$$i_1^* \circ j_{1*} : K_G^*(G/H) \rightarrow K_G^*(G/K), \quad [E] \mapsto [i_1^*(\lambda_{-1}(\tau_{j_1})) \otimes i_1^*(p^*(E))].$$

Afterwards, we compare the two maps and we conclude that the *obstruction bundle* is  $\lambda_{-1}(i_1^*(\tau_{j_1})/\tau_{j_2})$ . This means that

$$(2-1) \quad i_1^* \circ j_{1*}([E]) = j_{2*} \circ i_2^*([E]) \otimes \lambda_{-1}(i_1^*(\tau_{j_1})/\tau_{j_2}).$$

We consider the particular case of the groups  $H = C_G(x)$  and  $K = C_G(y)$ , where  $x$  and  $y$  are elements in the group  $G$  and  $C_G(x)$  and  $C_G(y)$  denote their centralizers in  $G$ . Then by (2-1) we get an obstruction bundle which is denoted as  $\gamma_{x,y}$ .

### 3. Twisted orbifold K-theory for the orbifold $[*/G]$

The goal of this section is to consider a K-theory structure on an orbifold structure defined by the trivial action of a finite group over the space  $\{*\}$ . This is a particular case of a more general kind of spaces that are obtained by almost free actions of a compact Lie group over compact manifolds. These spaces naturally have an orbifold structure that sets a basis for all developments in this paper. When the manifold is one point and the group is finite, all the hypotheses in the already defined theory hold.

Let us consider the inertia orbifold  $\wedge[*/G]$  for a finite group  $G$ . We define the orbifold K-theory for the orbifold  $[*/G]$  as the module

$$K_{\text{orb}}([*/G]) := K(\wedge[*/G]) \cong \bigoplus_{(g)} K([*/C_G(g)]) \cong \bigoplus_{(g)} K_{C_G(g)}(*),$$

where  $(g)$  denotes the class of conjugation of the element  $g \in G$  and  $C_G(g)$  denotes the centralizer of the element  $g \in G$ . In this case, the orbifold K-theory introduced in [Adem et al. 2007] turns out to be simply  $K_G(*)$ , which is additively isomorphic to the group  $\bigoplus_{(g)} R(C_G(g))$  (see [Adem and Ruan 2003]), where  $R(C_G(g))$  denotes the Grothendieck ring associated to the semigroup of isomorphism classes of linear representations of the group  $C_G(g)$ , and the sum is taken over conjugacy classes. The product structure in  $K_G(*)$  is defined as follows: consider the maps

$$\begin{aligned} e_1 : C_G(g) \cap C_G(h) \times C_G(g) \cap C_G(h) &\rightarrow C_G(g), & e_1(a, b) &= a, \\ e_2 : C_G(g) \cap C_G(h) \times C_G(g) \cap C_G(h) &\rightarrow C_G(h), & e_2(a, b) &= b, \\ e_{12} : C_G(g) \cap C_G(h) \times C_G(g) \cap C_G(h) &\rightarrow C_G(gh), & e_{12}(a, b) &= ab. \end{aligned}$$

Note that for any element  $\tau \in G$ , the map  $\phi_\tau : G \rightarrow G$  defined by  $\phi_\tau(g) = \tau g \tau^{-1}$  implies that  $\phi_\tau \circ e_i = e_i \circ (\phi_\tau, \phi_\tau)$ . Thus, the maps  $e_i$  are  $\phi_\tau$ -equivariant for any element  $\tau \in C_G(g)$ . Given  $E$  in  $R(C_G(g))$  and  $F$  in  $R(C_G(h))$ , we define the product

$$(3-1) \quad E \star F := e_{12*}(e_1^*(E) \otimes e_2^*(F) \otimes \gamma_{g,h}) \in R(C_G(gh)).$$

Since the action is trivial, it follows from Theorem 2.2 in [Segal 1968] that  $R(C_G(g)) = K_{C_G(g)}(*)$ . Thus, the product can be seen as

$$\star : K_{C_G(g)}(*) \times K_{C_G(h)}(*) \rightarrow K_{C_G(gh)}(*)$$

in the setup of equivariant K-theory. We note that the product defined in (3-1) is analogous to the stringy and twisted stringy product defined in [Becerra and Uribe 2009] in the case in which  $G$  is an abelian group. Let  $\alpha$  be a cocycle in  $Z^3(G; S^1)$ , i.e.,  $\alpha : G \times G \times G \rightarrow S^1$  satisfies  $\alpha(a, b, c)\alpha(a, bc, d)\alpha(d, c, d) = \alpha(ab, c, d)\alpha(a, b, cd)$  for all  $a, b, c, d \in G$ . We proceed to define the twisted orbifold K-theory  ${}^\alpha K_{\text{orb}}([*/G])$ .

For the global quotient  $[X/G]$  and the element  $\alpha \in Z^3(G; S^1)$ , the twisted orbifold K-theory is defined as the sum

$$(3-2) \quad {}^\alpha K_{\text{orb}}([X/G]) := \bigoplus_{g \in C} {}^{\alpha_g} K_{C_G(g)}(X^g),$$

where  $C$  is a set of representatives of the conjugacy classes in  $G$  and  $\alpha_g$  is the inverse transgression map (see below for details). In particular, if  $G$  is an abelian group, the set  $C$  is the group  $G$ . For every group  $H$  and  $\beta \in Z^2(H; S^1)$  we take its associated group  $H_\beta$  given by the central extension

$$1 \rightarrow S^1 \rightarrow H_\beta \rightarrow H \rightarrow 1.$$

Recall that the group  $H_\beta$  is the set  $S^1 \times H$ , with the group operation defined by

$$(s_1, h_1) * (s_2, h_2) := (s_1 s_2 \beta(h_1, h_2), h_1 h_2).$$

The twisted equivariant K-theory  ${}^\beta K_H(X)$  is defined as the class of  $H_\beta$ -equivariant vector bundles such that the action of the center  $S^1$  restricts to multiplication on the fibers. In the case of the space  $X = \{*\}$ , the twisted equivariant K-theory  ${}^\beta K_H(*)$  coincides with  $R_\beta(H)$ , the Grothendieck ring of classes of projective representations for the group  $H$  (see [Karpilovsky 1993] for a precise definition of  $R_\beta(H)$ ).

Returning to the case of the orbifold  $[*/G]$  for a finite group  $G$ , the twisted orbifold K-theory defined in (3-2) takes the form

$${}^\alpha K_{\text{orb}}([*/G]) := \bigoplus_{g \in C} {}^{\alpha_g} K_{C_G(g)}(*) \cong \bigoplus_{g \in C} R_{\alpha_g}(C_G(g)).$$

**Inverse transgression map.** We review the inverse transgression map for finite groups to describe the multiplicative structure in the module  ${}^{\alpha}K_{\text{orb}}([*/G])$ . Throughout this subsection we follow the development presented in Section 3.2 in [Becerra and Uribe 2009]. Let us recall the definition of the inverse transgression map for a global quotient  $[M/G]$ . For  $g \in G$ , consider the action of  $C_G(g) \times \mathbb{Z}$  on  $M^g = \{x \in M \mid gx = x\}$  given by  $(h, m) \cdot x := hg^m x$  and the homomorphism

$$\psi_g : C_G(g) \times \mathbb{Z} \rightarrow G, \quad (h, m) \mapsto hg^m.$$

Thus, the inclusion  $i_g : M^g \rightarrow M$  becomes a  $\psi_g$  equivariant map and induces a homomorphism

$$i_g^* : H_G^*(M; \mathbb{Z}) \rightarrow H_{C_G(g) \times \mathbb{Z}}^*(M^g; \mathbb{Z}).$$

From the isomorphisms

$$\begin{aligned} H_{C_G(g) \times \mathbb{Z}}^*(M^g; \mathbb{Z}) &\cong H^*(M^g \times_{C_G(g)} \times EC_G(g) \times B\mathbb{Z}; \mathbb{Z}) \\ &\cong H_{C_G(g)}^*(M^g; \mathbb{Z}) \otimes_{\mathbb{Z}} H^*(S^1; \mathbb{Z}), \end{aligned}$$

we have, for each  $k$ ,

$$i_g^* : H_G^k(M; \mathbb{Z}) \rightarrow H_{C_G(g)}^k(M^g; \mathbb{Z}) \oplus H_{C_G(g)}^{k-1}(M^g; \mathbb{Z}).$$

Hence, we define the inverse transgression map as the map induced by projecting on the second factor

$$\tau_g : H_G^k(M; \mathbb{Z}) \rightarrow H_{C_G(g)}^{k-1}(M^g; \mathbb{Z}).$$

In the particular case that  $[M/G] = [*/G]$  the definition above turns into:

**Definition 3.1.** For any element  $\alpha \in Z^3(G; S^1)$ , the inverse transgression map is defined as the map

$$\tau_g : H_G^k(*; \mathbb{Z}) \rightarrow H_{C_G(g)}^{k-1}(*; \mathbb{Z})$$

induced by  $\tau_g$  on each  $k$ .

**Product in the twisted case.** Take  $\alpha \in Z^3(G; \mathbb{Z})$ . Let us consider the orbifold  $[*/G]$  where  $G$  is a finite group. Now, consider the module

$$(3-3) \quad {}^{\alpha}K_{\text{orb}}([*/G]) := \bigoplus_{g \in C} R_{\alpha_g}(C_G(g)),$$

where  $\alpha_g \in H^2(C(g); \mathbb{Z})$  denotes the inverse transgression map of  $\alpha$ . The goal of this subsection is to define an associative product for this module; specifically, we show that it's possible to endow the module  ${}^{\alpha}K_{\text{orb}}([*/G])$  with a ring structure. For simplicity, we denote  $C_G(g)$  as  $C(g)$ . Consider the inclusion maps of groups

$$i_g : C(g) \cap C(h) \rightarrow C(g), \quad i_h : C(g) \cap C(h) \rightarrow C(h)$$

and

$$i_{gh} : C(g) \cap C(h) \rightarrow C(gh)$$

for  $g, h \in G$ . These maps induce the restriction maps

$$\begin{aligned} i_g^* : H^2(C(g); S^1) &\rightarrow H^2(C(g) \cap C(h); S^1), \\ i_h^* : H^2(C(h); S^1) &\rightarrow H^2(C(g) \cap C(h); S^1), \end{aligned}$$

and the morphism  $i_{gh}$  induces a map

$$(i_{gh})_* : H^2(C(g) \cap C(h); S^1) \rightarrow H^2(C(gh); S^1),$$

which is the induction morphism in group cohomology (see for example [tom Dieck 1987]).

Given  $E \in R_{\alpha_g}(C(g))$ , we consider it as a  $C(g)_{\alpha_g}$ -module that restricts to multiplication on the fibers over  $S^1$ . Therefore, we get the following commutative diagram for the inclusion  $i_g$  and the identity map  $s$  on  $S^1$ :

$$(3-4) \quad \begin{array}{ccccccc} 1 & \longrightarrow & S^1 & \longrightarrow & C(g)_{\alpha_g} & \longrightarrow & C(g) \longrightarrow 1 \\ & & \uparrow s & & \uparrow (s, i_g) & & \uparrow i_g \\ 1 & \longrightarrow & S^1 & \longrightarrow & (C(g) \cap C(h))_{i_g^*(\alpha_g)} & \longrightarrow & C(g) \cap C(h) \longrightarrow 1 \end{array}$$

This implies that any  $C(g)_{\alpha_g}$ -module restricts to a  $(C(g) \cap C(h))_{i_g^*(\alpha_g)}$ -module, denoted by  $i_g^*(E)$ . In particular, for any  $(E, F) \in R_{\alpha_g}(C(g)) \times R_{\alpha_h}(C(h))$ , we get the map

$$\begin{aligned} R_{\alpha_g}(C(g)) \times R_{\alpha_h}(C(h)) &\rightarrow R_{i_g^*(\alpha_g)}(C(g) \cap C(h)) \times R_{i_h^*(\alpha_h)}(C(g) \cap C(h)), \\ (E, F) &\mapsto (i_g^*(E), i_h^*(F)). \end{aligned}$$

Now, from the central extensions

$$\begin{aligned} 1 \rightarrow S^1 &\rightarrow (C(g) \cap C(h))_{i_g^*(\alpha_g)} \rightarrow C(g) \cap C(h) \rightarrow 1, \\ 1 \rightarrow S^1 &\rightarrow (C(g) \cap C(h))_{i_h^*(\alpha_h)} \rightarrow C(g) \cap C(h) \rightarrow 1 \end{aligned}$$

induced by  $i_g^*(\alpha)$  and  $i_h^*(\alpha) \in H^2(C(g) \cap C(h); S^1)$ , we get

$$\begin{aligned} 1 \rightarrow S^1 \times S^1 &\rightarrow (C(g) \cap C(h))_{i_g^*(\alpha_g)} \times (C(g) \cap C(h))_{i_h^*(\alpha_h)} \\ &\rightarrow C(g) \cap C(h) \times C(g) \cap C(h) \rightarrow 1 \end{aligned}$$

For  $E \in R_{i_g^*(\alpha_g)}(C(g))$  and  $F \in R_{i_h^*(\alpha_h)}(C(g))$ , the tensor product  $E \otimes F$  is naturally a  $(C(g) \cap C(h))_{i_g^*(\alpha_g)} \times (C(g) \cap C(h))_{i_h^*(\alpha_h)}$ -module that restricts to multiplication



on the fibers by elements of  $S^1$ . By considering the action restricted to the diagonal

$$\Delta(C(g) \cap C(h)) \subset C(g) \cap C(h) \times C(g) \cap C(h),$$

we get the central extension

$$1 \rightarrow S^1 \rightarrow (C(g) \cap C(h))_{i_g^*(\alpha_g) i_h^*(\alpha_h)} \rightarrow \Delta(C(g) \cap C(h)) \rightarrow 1$$

corresponding to the element

$$i_g^*(\alpha_g) i_h^*(\alpha_h) \in H^2(C(g) \cap C(h); S^1).$$

Thus, we have

$$\begin{aligned} R_{i_g^*(\alpha_g)}(C(g) \cap C(h)) \times R_{i_h^*(\alpha_h)}(C(g) \cap C(h)) &\rightarrow R_{i_g^*(\alpha_g) i_h^*(\alpha_h)}(C(g) \cap C(h)), \\ (E, F) &\mapsto E \otimes F. \end{aligned}$$

Now, since  $i_g^*(\alpha_g) i_h^*(\alpha_h) = i_{gh}^*(\alpha_{gh})$  in  $H^2(C(g) \cap C(h); S^1)$  since the cocycles are cohomologous (see [Adem et al. 2007, Proposition 4.3]), it follows that

$$R_{i_g^*(\alpha_g) i_h^*(\alpha_h)}(C(g) \cap C(h)) \cong R_{i_{gh}^*(\alpha_{gh})}(C(g) \cap C(h)).$$

Therefore, the induction map can be defined as

$$R_{i_{gh}^*(\alpha_{gh})}(C(g) \cap C(h)) \rightarrow R_{(i_{gh})^*(\alpha_{gh})}(C(gh)), \quad A \mapsto \text{Ind}_{C(g) \cap C(h)}^{C(gh)}(A).$$

Thus, a product on the module (3-3) can be obtained from the previously described morphisms to get

$$(3-5) \quad R_{\alpha_g}(C(g)) \times R_{\alpha_h}(C(h)) \rightarrow R_{\alpha_{gh}}(C(gh))$$

defined by

$$(E, F) \mapsto E \star_{\alpha} F := \text{Ind}_{C(g) \cap C(h)}^{C(gh)}(i_g^*(E) \otimes i_h^*(F) \otimes \gamma_{g,h}),$$

where  $\gamma_{g,h}$  is defined as the excess bundle as in Section 2.

**Definition 3.2.** By using the restriction notation, we define the twisted stringy product in the module  ${}^{\alpha}K_{\text{orb}}([*/G])$  as the map

$$\begin{aligned} R_{\alpha_g}(C(g)) \times R_{\alpha_h}(C(h)) &\rightarrow R_{\alpha_g \alpha_h}(C(gh)) \\ (E, F) &\mapsto I_{C(g) \cap C(h)}^{C(gh)}(\text{Res}_{C(g) \cap C(h)}^{C(g)}(E) \otimes \text{Res}_{C(g) \cap C(h)}^{C(h)}(F) \otimes \gamma_{g,h}), \end{aligned}$$

where  $\text{Res}_{C(g) \cap C(h)}^{C(g)}$  denotes the restriction of  $\alpha$ -twisted representations of  $C(g)$  to  $i_g(\alpha)$ -representations of  $C(g) \cap C(h)$  (and likewise for  $h$ ).

### 4. Twisted orbifold K-theory and the algebra $D^\omega(G)$

The goal of this section is to give an introduction of the *twisted Drinfeld double*  $D^\omega(G)$  and to show an explicit relation with the twisted orbifold K-theory. Our main reference is [Witherspoon 1996]. Let us recall the definition and the main properties of the twisted Drinfeld double to clarify the nature of this structure and its representations. From a different point of view, we can also obtain some properties of the stringy product defined on the sections above, using the properties of the representations of the twisted Drinfeld double. Namely, the Grothendieck ring of these representations is isomorphic to the twisted orbifold K-theory with the structure induced by the stringy product, which will be proven on page 481. Because of the associativity of the tensor product of the  $D^\omega(G)$ -modules, this yields a proof of the associativity of the stringy product defined above; see Corollary 4.3.

Let  $G$  be a finite group and  $k$  an algebraically closed field. Let  $\omega$  be an element in  $Z^3(G, k^*)$ , that is, a function  $\omega : G \times G \times G \rightarrow k^*$  such that

$$\omega(a, b, c)\omega(a, bc, d)\omega(d, c, d) = \omega(ab, c, d)\omega(a, b, cd)$$

for all  $a, b, c, d \in G$ . We define the quasitriangular quasi-Hopf algebra  $D^\omega(G)$  as the vector space  $(kG)^* \otimes (kG)$ , where  $(kG)^*$  denotes the dual of the algebra  $kG$  (see [Drinfel'd 1987]) and the algebra structure in  $D^\omega(G)$  is given as follows: consider the canonical basis  $\{\delta_g \otimes \bar{x}\}_{g,x \in G}$  of  $D^\omega(G)$ , where  $\delta_g$  is the function such that  $\delta_g(h) = 1$  if  $h = g$  and 0 otherwise. We denote  $\delta_g \otimes \bar{x}$  by  $\delta_g \bar{x}$ . Now, we define the product of elements in the basis by

$$(4-1) \quad (\delta_g \bar{x})(\delta_h \bar{y}) = \omega_g(x, y)\delta_g \delta_{xhx^{-1}} \bar{xy},$$

where  $\omega_g$  is the image of  $\omega$  via the inverse transgression map of the element  $g \in G$  as in Definition 3.1. The multiplicative identity for this product is the element  $1_{D^\omega(G)} = \bigoplus_{g \in G} \delta_g \bar{1}$ . Now, we use the notation  $\delta_g$  for the element  $\delta_g \bar{1}$ . The coproduct  $\Delta : D^\omega(G) \rightarrow D^\omega(G) \otimes D^\omega(G)$  in the algebra  $D^\omega(G)$  is defined by the map

$$(4-2) \quad \Delta(\delta_g \bar{x}) = \bigoplus_{h \in G} \gamma_x(h, h^{-1}g)(\delta_h \bar{x}) \otimes (\delta_{h^{-1}g} \bar{x}),$$

where

$$\gamma_x(h, l) = \frac{\omega(h, l, x)\omega(x, x^{-1}hx, x^{-1}lx)}{\omega(h, x, x^{-1}lx)}.$$

The algebra  $D^\omega(G)$  endowed with these operations is usually called the twisted Drinfeld double.

**Representations of  $D^\omega(G)$ .** Let  $U, V$  be modules over the algebra  $D^\omega(G)$ . Consider the tensor product  $U \otimes V$  as a  $D^\omega(G)$ -module endowed with the action from  $D^\omega(G)$ , induced by the coproduct  $\Delta$ . Note that the field  $k$  can be considered as a

trivial  $D^\omega(G)$ -module, which is the multiplicative identity for the tensor product of  $D^\omega(G)$ -modules. In particular, for  $k = \mathbb{C}$ , we define the ring of representations  $R(D^\omega(G))$  of  $D^\omega(G)$  as the  $\mathbb{C}$ -algebra generated by the set of isomorphism classes of  $D^\omega(G)$ -modules with the direct sum of modules as the sum operation and the tensor as the product operation. We define the ideal  $R_0(D^\omega(G))$  generated by all combinations  $[U] - [U'] - [U'']$  (brackets denoting the isomorphism class) where  $0 \rightarrow U' \rightarrow U \rightarrow U'' \rightarrow 0$  is a short exact sequence of  $D^\omega(G)$ -modules. Now, we define the Grothendieck ring  $R(D^\omega(G))$  as the quotient between  $\text{Rep}(D^\omega(G))$  and the ideal  $R_0(D^\omega(G))$ .

The algebra  $D^\omega(G)$  is quasitriangular with

$$A = \bigoplus_{g,h \in G} \delta_g \bar{1} \otimes \delta_h \bar{g} \quad \text{and} \quad A^{-1} = \bigoplus_{g,h \in G} \omega_{ghg^{-1}}(g, g^{-1})^{-1} \delta_g \bar{1} \otimes \delta_h \overline{g^{-1}}.$$

Thus,  $A\Delta(a)A^{-1} = \sigma(\Delta(a))$  for all  $a \in D^\omega(G)$ , where  $\sigma$  is the automorphism that exchanges the images in the coproduct. Therefore, if  $U$  and  $V$  are  $D^\omega(G)$ -modules, this equation implies that  $U \otimes V$  and  $V \otimes U$  are isomorphic as  $D^\omega(G)$ -modules; that is, the algebra  $R(D^\omega(G))$  is commutative. Now, assume that  $\beta : G \times G \rightarrow \mathbb{C}^*$  is a cochain with coboundary

$$\delta\beta(a, b, c) = \beta(b, c)\beta(a, bc)\beta(ab, c)^{-1}\beta(a, b)^{-1}.$$

Then, the algebra  $D^{\omega\delta\beta}(G)$  is isomorphic to  $D^\omega(G)$  given through the map

$$\nu(\delta_g \bar{x}) = \frac{\beta(g, x)}{\beta(x, xgx^{-1})} \delta_g \bar{x}.$$

In particular, we get the isomorphism

$$\nu^* : R(D^{\omega\delta\beta}(G)) \xrightarrow{\cong} R(D^\omega(G)).$$

Next, we consider the following theorem (compare [Willerton 2008, Theorem 19]):

**Theorem 4.1.** *The ring  $R(D^\omega(G))$  is additively isomorphic to the ring*

$$\bigoplus_{(g) \subset G} R_{\omega_g}(C(g)),$$

where  $(g)$  denotes the conjugacy class of  $g \in G$ .

*Proof.* For all  $x \in G$ , we take the subspaces

$$S^\omega(x) := \bigoplus_{g \in C(x)} \mathbb{C} \delta_x \bar{g} \quad \text{and} \quad D^\omega(x) := \bigoplus_{g \in G} \mathbb{C} \delta_x \bar{g}$$

of  $D^\omega(G)$ . Then  $S^\omega(x)$  is a subalgebra of  $D^\omega(G)$  with identity element  $\delta_x \bar{1}$  such that, from the product defined in  $D^\omega(G)$ , it follows that  $S^\omega(x) \cong R_{\omega_x} C(x)$  where

$R_{\omega_x}C(x)$  is defined in [Karpilovsky 1993]. Given  $(g) \subset G$ , consider

$$D^\omega((g)) := \bigoplus_{h \in (g)} D^\omega(h).$$

Note that  $D^\omega(G) \cong \bigoplus_{(g) \subset G} D^\omega((g))$  (additively). For an element  $h$  in a fixed conjugacy class  $(g)$ , take a  $S^\omega(h)$ -module (i.e., a  $R_{\omega_h}C(h)$ -module)  $U$ , and define the map

$$U \mapsto U \otimes_{S^\omega(h)} D^\omega(h),$$

whose image is a  $D^\omega((g))$ -module if we take the action of  $D^\omega((g))$  on it as right multiplication in the second factor. On the other hand, for a  $D^\omega((g))$ -module  $V$ , we define the map

$$V \mapsto V \delta_h \bar{1},$$

whose image is a  $R_{\omega_h}C(h)$ -module. Thus, there is an equivalence between  $R_{\omega_h}C(h)$ -modules and  $D^\omega((g))$ -modules. Therefore, from [Karpilovsky 1993, Theorem I.3.2], we have  $R_{\omega_h}(C(h)) \cong R(D^\omega((g)))$  for any  $h \in (g)$ , and the theorem follows.  $\square$

From [Dijkgraaf et al. 1991], we get that it is possible to explicitly describe the morphism using the *induction DPR* which is defined on each  $R_\alpha(C(g))$  for  $g \in G$ . Namely, let  $(\rho, V)$  be a twisted representation of the group  $C(g)$  and define the representation  $\psi((\rho, V)) := (\pi_\rho, A)$  of  $D^\omega(G)$  as given by

$$(4-3) \quad A := \text{Ind}_{C(g)}^G(V), \quad \pi_\rho := \pi_\rho(\delta_k \bar{x})x_j \otimes v = \delta_k \delta_{x_s g x_s^{-1}} \frac{\omega_k(x, x_j)}{\omega_k(x_s, r)} x_s \otimes \rho(r)v,$$

where  $x_j$  is a representative of a class in  $G/C(g)$ ,  $r \in C(g)$  and the element  $x_s$  is a representative of a class in  $G/C(g)$ , such that  $xx_j = x_s r$ .

**Relation between  $R(D^\omega(G))$  and the twisted K-theory of the orbifold  $[*/G]$ .** Let us consider an element  $\omega \in Z^3(G; S^1)$ . By (3-3) the twisted orbifold K-theory of the orbifold  $[*/G]$  is the ring

$${}^\omega K_{\text{orb}}([*/G]) = \bigoplus_{(g) \subset G} {}^\omega K_{C(g)}(*) \cong \bigoplus_{(g) \subset G} R_{\omega_g}(C(g)).$$

By Theorem 4.1, there exists an additive isomorphism between  $R(D^\omega(G))$  and the twisted orbifold K-theory  ${}^\omega K_{\text{orb}}([*/G])$ . We will show that if we endow this ring with the twisted product  $\star_\alpha$ , then the additive isomorphism is in fact a ring isomorphism. The DPR induction is defined as  $(I_{C(g)}^G(E), \rho_\pi)$  where  $(E, \pi)$  is an element in  $R_{\omega_g}(C(g))$ . Let us consider two elements  $E$  and  $F$  in  $R_{\omega_g}(C(g))$  and  $R_{\omega_h}(C(h))$  respectively. The tensor product of the DPR-induction of these elements

can be related to the twisted product  $\star$  via the Frobenius reciprocity as follows:

$$\begin{aligned}
 I_{C(g)}^G(E) \otimes I_{C(h)}^G(F) &\cong I_{C(g)}^G(E \otimes R_{C(g)}^G(I_{C(h)}^G(F))) \\
 &\cong I_{C(g)}^G(E \otimes I_{C(g)}^G(R_{C(g) \cap C(h)}^{C(h)}(F) \otimes \gamma_{g,h})) \\
 &\cong I_{C(g)}^G(I_{C(g) \cap C(h)}^{C(g)}(R_{C(g) \cap C(h)}^{C(g)}(E) \otimes R_{C(g) \cap C(h)}^{C(h)}(F) \otimes \gamma_{g,h})) \\
 &\cong I_{C(g) \cap C(h)}^G(R_{C(g) \cap C(h)}^{C(g)}(E) \otimes R_{C(g) \cap C(h)}^{C(h)}(F) \otimes \gamma_{g,h}) \\
 &\cong I_{C(g)}^G(I_{C(g) \cap C(h)}^{C(gh)}(R_{C(g) \cap C(h)}^{C(g)}(E) \otimes R_{C(g) \cap C(h)}^{C(h)}(F) \otimes \gamma_{g,h})) \\
 &\cong I_{C(gh)}^G(E \star_\omega F).
 \end{aligned}$$

**Proposition 4.2.** *There exists a ring isomorphism*

$$(\omega K_{\text{orb}}([*/G]), \star_\omega) \cong (R(D^\omega(G)), \otimes).$$

*Proof.* DPR induction defines a morphism  $\phi : \bigoplus_{(g) \subset G} R_\omega(C(g)) \rightarrow R(D^\omega(G))$ . Moreover, we proved above that for  $E \in R_{\omega_g}(C(g))$  and  $F \in R_{\omega_h}(C(h))$ , we have

$$\phi(E) \otimes \phi(F) = \phi(E \star_\omega F);$$

that is, it is a ring homomorphism. By [Theorem 4.1](#), the result follows. □

**Corollary 4.3.** *The stringy product  $\star_\omega$  is associative.*

### 5. Twisted K-theory for an extraspecial $p$ -group

The goal of this section is to establish a relation between the twisted orbifold K-theories for the orbifolds  $[*/H]$  and  $[*/G]$ , where  $H$  is an extraspecial group with exponent  $p$ , order  $p^{2n+1}$  and  $G = (\mathbb{Z}_p)^{2n+1}$ . For an odd prime number  $p$ , a  $p$ -group  $H$  is called *extraspecial* if its center  $Z(H)$  is a cyclic group of order  $p$ , that is  $Z(H) \cong \mathbb{Z}_p$ , and  $H/Z(H)$  is an elementary abelian group. Any extraspecial  $p$ -group has order  $p^{2n+1}$  for some  $n \in \mathbb{N}$ . On the other hand, for any  $n$  there exist two extraspecial groups of order  $p^{2n+1}$  such that a group has exponent  $p$  and the other group has exponent  $p^2$ . The motivation for these kinds of relations comes from works such as [\[Goff et al. 2007\]](#), where these relations are studied for  $p = 2$ , and to some extent results due to A. Duman [\[2009\]](#). However, there exists a deeper interest to study these kinds of relations by establishing correspondences with the twisted Drinfeld algebras. In particular, the following result is of utmost importance for obtaining the results of this section:

**Theorem 5.1** [\[Naidu and Nikshych 2008, Corollary 4.20\]](#). *Let  $H$  be a finite group  $\omega' \in Z^3(H; S^1)$  such that*

- $H$  contains an abelian normal subgroup  $K$ ,
- $\omega'|_{K \times K \times K}$  is trivial in cohomology (in  $H^3(K; S^1)$ ),

- there exists an  $H$ -invariant 2-cochain  $\mu$  over  $H$  such that  $\delta(\mu)|_{K \times K \times K} = \omega'|_{K \times K \times K}$ .

Then, there exists a group  $G$  and an element  $\omega \in Z^2(G; S^1)$  such that  $R(D^\omega(G)) \cong R(D^{\omega'}(H))$ .

From the relation established in the previous section between the twisted Drinfeld’s algebras and the twisted orbifold K-theory, we get the following corollary under the same assumptions as in the last theorem.

**Corollary 5.2.** *There exists a ring isomorphism*

$${}^\omega K_{\text{orb}}([*/G]) \cong {}^{\omega'} K_{\text{orb}}([*/H]).$$

Now, we follow with a nice application of this result.

**Proposition 5.3.** *Let  $H$  be an extraspecial group with order  $p^{2n+1}$  and exponent  $p$ . Then*

$$K_{\text{orb}}([*/H]) \cong {}^\omega K_{\text{orb}}([*/(\mathbb{Z}_p)^{2n+1}])$$

for some nontrivial twisting  $\omega$ .

*Proof.* Let  $H$  be an extraspecial group. From definition we may assume  $K = Z(H) \cong \mathbb{Z}_p$ . Now, suppose there exists  $\mu \in C^2(H; S^1)$  such that  $\delta(\mu)|_{K \times K \times K} = \omega'|_{K \times K \times K}$ , which is  $H$ -invariant; that is, if we take the action of  $H$  on the 2-cochains  $C^2(H; S^1)$  defined by  ${}^y\mu := \mu(yx_1y^{-1}, yx_2y^{-1})$ , then  ${}^y\mu = \mu$  in  $C^2(H; S^1)$  for all  $y \in H$ . Now, since  $K = Z(H)$ , it follows that  ${}^y\mu|_K = \mu|_K$  for all  $y \in H$ . Thus, for all  $y \in H$  there exists a 1-chain  $\eta_y$  on  $H$  such that  $\delta\eta_y = {}^y\mu/\mu = 1$ . Since  $K$  is abelian, we can define the map

$$v : H/K \times H/K \rightarrow C^1(H; S^1), \quad (y_1, y_1) \mapsto \frac{{}^{y_2}\eta_{y_1}\eta_{y_2}}{\eta_{y_1y_2}}.$$

**Lemma 5.4** [Naidu 2007, Lemma 4.2, Corollary 4.3]. *The function  $v$  defines an element in  $H^2(H/K; \hat{K})$ .*

However, this element represents a short exact sequence

$$1 \rightarrow \hat{K} \rightarrow \hat{K} \times_v H/K \rightarrow H/K \rightarrow 1,$$

where the product in  $\hat{K} \times_v H/K$  is defined by the formula

$$(\rho_1, x_1)(\rho_2, x_2) := (v(x_1, x_2)\rho_1\rho_2, x_1x_2).$$

Now, the element  $\omega \in Z^3(G; S^1)$ , with  $G := \hat{K} \times_v H/K$ , is defined for all  $(\rho_1, x_1), (\rho_2, x_2), (\rho_3, x_3)$  in  $\hat{K} \times_v H/K$  by the formula

$$\omega((\rho_1, x_1)(\rho_2, x_2)(\rho_3, x_3)) := (v(x_1, x_2)(u(x_3)))(1)\rho_1(k_{x_2, x_3}),$$

where  $u : H/K \rightarrow H$  is a function such that, when composed with the projection

$p : H \rightarrow H/K$ , yields  $p(u(x)) = x$ , and  $k_{x_2, x_3} \in H$  is an element that satisfies  $u(x_1)u(x_2) = k_{x_1, x_2}u(p(u(x_1)u(x_2)))$ .

Clearly, when  $\omega'$  is the trivial 3-cocycle, we can choose  $\mu$  to be trivial and so  $\nu$  is also trivial. By definition of an extraspecial group,  $H/K$  is an elementary abelian group and if  $\nu$  is trivial, it follows easily that  $\hat{K} \times_\nu H/K \cong (\mathbb{Z}_p)^{2n+1}$ . It remains to show that  $\omega$  is nontrivial in  $H^3(G; S^1)$ . Take  $H = \{h_1, \dots, h_{p^{2n+1}}\}$ ,  $K = Z(H) = \{z_1, \dots, z_p\}$  and  $\hat{K} = \{\rho_1, \dots, \rho_p\}$ , with  $\rho_i$  nontrivial for  $i \neq 1$ . Denote the quotient group  $H/K = \{x_1K, \dots, x_{p^{2n}}K\}$ , with  $x_1 = 1_{H/K}$ . Now, we define the function  $u : H/K \rightarrow H$  such that  $u(x_iK) = x_i(z_i^{-1})$  for  $x_iK \in H/K$ ,  $z_j \in K$ . Consider the element  $((\rho, x_iK), (\rho, x_iK), (\rho, x_iK))$  with  $\rho \in \hat{K}$  fixed and nontrivial. Since  $\nu$  is trivial, the element  $\omega$  is reduced to  $\rho(k_{x_iK, x_iK}) = \rho(z_i) \neq 1$ , which implies that  $\omega$  is nontrivial.  $\square$

**Twisted orbifold K-theory for the orbifold  $[*/(\mathbb{Z}_p)^n]$ .** With the above result, to calculate the orbifold K-theory structure for  $[*/H]$ , with  $H$  an extraspecial  $p$ -group, we only have to calculate the twisted orbifold K-theory for  $[*/(\mathbb{Z}_p)^n]$  and a twist element in  $H^3((\mathbb{Z}_p)^n; S^1)$ , following the constructions presented in Section 3. Because all those constructions are based on the inverse transgression map, we proceed to give an explicit way of calculating it. Later we present an example with a particular twist element, having no trivial inverse transgression map.

\*. Inverse transgression map for the group  $(\mathbb{Z}_p)^n$  Let us consider the following commutative diagram given by two natural short exact sequences:

$$(5-1) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{\times p} & \mathbb{Z} & \xrightarrow{\pi} & \mathbb{Z}_p & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathbb{Z}_p & \xrightarrow{\times p} & \mathbb{Z}_{p^2} & \xrightarrow{\tau} & \mathbb{Z}_p & \longrightarrow & 0 \end{array}$$

where  $\pi$  and  $\tau$  are the natural projections, and likewise for the downarrow maps. These two exact sequences in the diagram above induce long exact sequences

$$(5-2) \quad \dots \longrightarrow H^{k-1}(BG; \mathbb{Z}_p) \xrightarrow{\partial} H^k(BG; \mathbb{Z}) \xrightarrow{(\times p)_*} H^k(BG; \mathbb{Z}) \xrightarrow{(\pi)_*} H^k(BG; \mathbb{Z}_p) \xrightarrow{\partial} H^{k+1}(BG; \mathbb{Z}) \longrightarrow \dots,$$

$$(5-3) \quad \dots \longrightarrow H^{k-1}(BG; \mathbb{Z}_p) \xrightarrow{\beta} H^k(BG; \mathbb{Z}_p) \xrightarrow{(\times p)_*} H^k(BG; \mathbb{Z}_{p^2}) \xrightarrow{(\tau)_*} H^k(BG; \mathbb{Z}_p) \xrightarrow{\beta} H^{k+1}(BG; \mathbb{Z}_p) \longrightarrow \dots$$

**Remark 5.5.** The connection morphism  $\beta$  of the long exact sequence (5-3) is known as the Bockstein map. It induces a map  $\beta : H^*(BG; \mathbb{Z}_p) \rightarrow H^*(BG; \mathbb{Z}_p)$

which has the multiplicative property

$$\beta(xy) = \beta(x)y + (-1)^{\deg(x)}x\beta(y).$$

Since  $G$  is a  $p$ -group,  $H^k(BG; -)$  is also a  $p$ -group and this implies that the morphism  $(\times p)_*$  in the long exact sequences (5-2) and (5-3) is the zero map. Thus,  $\pi_*$  and  $\tau_*$  are injective maps and  $H^k(BG; \mathbb{Z}) \cong H^k(BG; \mathbb{Z}_{p^2})$ . On the other hand, by the exactness of the sequence (5-3), we have  $H^k(BG; \mathbb{Z}_{p^2}) \cong \text{Ker}(\beta : H^k(BG; \mathbb{Z}_p) \rightarrow H^{k+1}(BG; \mathbb{Z}_p))$  and then

$$H^k(BG; \mathbb{Z}) \cong \text{Ker}(\beta : H^k(BG; \mathbb{Z}_p) \rightarrow H^{k+1}(BG; \mathbb{Z}_p)).$$

*Relation to the inverse transgression map.* By definition, the inverse transgression map  $\tau_g$  is a map defined between the groups  $H^k(BG; \mathbb{Z})$  and  $H^{k-1}(BC_G(g); \mathbb{Z})$ . Since  $G = (\mathbb{Z}_p)^n$  is an abelian group, the inverse transgression map can be factorized as

$$\begin{aligned} \tilde{\tau}_g : \text{Ker}(\beta : H^k(BG; \mathbb{Z}_p) \rightarrow H^{k+1}(BG; \mathbb{Z}_p)) \\ \rightarrow \text{Ker}(\beta : H^{k-1}(BG; \mathbb{Z}_p) \rightarrow H^k(BG; \mathbb{Z}_p)). \end{aligned}$$

Consider the cohomology ring  $H^*(BG; \mathbb{Z}_p) \cong \mathbb{F}_p[x_1, \dots, x_n] \otimes \Lambda[y_1, \dots, y_n]$  with  $|x_i| = 2$  and  $|y_i| = 1$  for  $i = 1, \dots, n$ . By the calculations above we need to find a polynomial  $p(x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{F}_p[x_1, \dots, x_n] \otimes \Lambda[y_1, \dots, y_n]$  of degree  $k$ , such that  $\beta(p) = 0$  and  $\tilde{\tau}_g(p) \neq 0$  for some  $g \in G$ .

To obtain the desired polynomial, first we do the calculation of the inverse transgression map. Take an element  $g = (a_1, \dots, a_n) \in G$  and consider the map

$$G \times \mathbb{Z} \rightarrow G \times \langle g \rangle \rightarrow G$$

defined by

$$(h, m) \mapsto (h, g^m) \mapsto hg^m.$$

At the level of cohomology we get

$$\begin{aligned} (5-4) \quad H^*(BG; \mathbb{F}_p) &\rightarrow H^*(BG \times B(\mathbb{Z}_p); \mathbb{F}_p) \rightarrow H^*(BG \times B\mathbb{Z}; \mathbb{F}_p), \\ x_i &\mapsto x_i + a_i w && \mapsto x_i, \\ y_i &\mapsto y_i + a_i z && \mapsto y_i + a_i z, \end{aligned}$$

where

$$\begin{aligned} H^*(BG; \mathbb{F}_p) &= \mathbb{F}_p[x_1, \dots, x_n] \otimes \Lambda[y_1, \dots, y_n], \\ H^*(BG \times B(\mathbb{Z}_p); \mathbb{F}_p) &= \mathbb{F}_p[x_1, \dots, x_n, w] \otimes \Lambda[y_1, \dots, y_n, z], \\ H^*(BG \times B\mathbb{Z}; \mathbb{F}_p) &= \mathbb{F}_p[x_1, \dots, x_n] \otimes \Lambda[y_1, \dots, y_n, z]. \end{aligned}$$

Now, for the products  $x_i y_j, x_i x_j, y_i y_j \in H^*(BG; \mathbb{F}_p)$  we can obtain the calculation



of the inverse transgression maps. For the first product  $x_i y_j$  we get

$$(5-5) \quad (x_i y_j) \mapsto (x_i + a_i w)(y_j + a_i z) = x_i y_j + x_i a_j z + a_i w y_j + a_i a_j w z$$

in  $H^*(BG \times B(\mathbb{Z}_p); \mathbb{F}_p)$  and  $(x_i y_j) \mapsto x_i y_j + x_i a_j z$  in  $H^*(BG \times B\mathbb{Z}; \mathbb{F}_p)$ , from [Definition 3.1](#) it follows that  $\tilde{\tau}_g(x_i y_j) = x_i a_j$ . For the second product  $x_i x_j$  we get

$$(x_i x_j) \mapsto (x_i + a_i w)(x_j + a_i w) = x_i x_j + x_i a_j w + a_i w x_j$$

in  $H^*(BG \times B(\mathbb{Z}_p); \mathbb{F}_p)$  and  $(x_i x_j) \mapsto x_i x_j$  in  $H^*(BG \times B\mathbb{Z}; \mathbb{F}_p)$ ; hence

$$(5-6) \quad \tilde{\tau}_g(x_i x_j) = 0.$$

Finally, for the product  $y_i y_j$  we get

$$(5-7) \quad (y_i y_j) \mapsto (y_i + a_i z)(y_j + a_i z) = y_i y_j + y_i a_j z + a_i z y_j$$

in  $H^*(BG \times B(\mathbb{Z}_p); \mathbb{F}_p)$  and  $(y_i y_j) \mapsto y_i y_j + (a_j y_i - a_i y_j)z$  in  $H^*(BG \times B\mathbb{Z}; \mathbb{F}_p)$ ; therefore  $\tilde{\tau}_g(y_i y_j) = (a_j y_i - a_i y_j)$ .

Since we are interested in calculating the inverse transgression map for elements  $\alpha \in H^4(G; \mathbb{Z})$ , we consider only polynomials of degree 4 in

$$H^*(BG; \mathbb{F}_p) = \mathbb{F}_p[x_1, \dots, x_n] \otimes \Lambda[y_1, \dots, y_n].$$

Now, we present some examples of the inverse transgression map. It is easiest to consider the cases  $n = 2$  and  $n = 3$ . In the first case the inverse transgression map is a trivial map. In the latter the inverse transgression map is more interesting.

**Example 5.6.**  $n = 2$ . For  $p \neq 2$  we have  $H^*(BG; \mathbb{F}_p) = \mathbb{F}_p[x_1, x_2] \otimes \Lambda[y_1, y_2]$  with  $|y_i| = 1$  and  $|\beta y_i| = |x_i| = 2$ . Thus, we can just consider linear combinations of the polynomials  $p_1(x_1, x_2, y_1, y_2) := x_1 x_2$ ,  $p_2(x_1, x_2, y_1, y_2) := x_1 y_1 y_2$  and  $p_3(x_1, x_2, y_1, y_2) := x_2 y_1 y_2$ . For  $p_1$  the calculations leading up to [\(5-6\)](#) show that  $\tilde{\tau}_g(p_1) = 0$ . Thus we need to find a  $(\mathbb{Z}_p)$ -linear combination  $p$  of the polynomials  $p_2$  and  $p_3$  such that  $\beta(p) = 0$ . But we have

$$\beta(p_3) = x_2(\beta(y_1)y_2 - y_1\beta(y_2)) = x_2(x_1 y_2 - y_1 x_2),$$

$$\beta(p_2) = x_1(\beta(y_1)y_2 - y_1\beta(y_2)) = x_1(x_1 y_2 - y_1 x_2).$$

Therefore, there does not exist such a  $(\mathbb{Z}_p)$ -linear combination.

$n = 3$ . By analyzing the degree of the polynomials, we obtain the element

$$(5-8) \quad p(x_1, x_2, x_3, y_1, y_2, y_3) = x_1 y_2 y_3 - x_2 y_1 y_3 + x_3 y_1 y_2,$$

which satisfies the condition  $\beta(p) = 0$ . To check this, we use the property of  $\beta$

noted in [Remark 5.5](#):

$$\begin{aligned} \beta(p) &= \beta(x_1y_2y_3) - \beta(x_2y_1y_3) + \beta(x_3y_1y_2) \\ &= \beta(x_1)y_2y_3 + x_1\beta(y_2y_3) - \beta(x_2)y_1y_3 - x_2\beta(y_1y_3) + \beta(x_3)y_1y_2 + x_3\beta(y_1y_2) \\ &= x_1\beta(y_2)y_3 - x_1y_2\beta(y_3) - x_2\beta(y_1)y_3 + x_2y_1\beta(y_3) + x_3\beta(y_1)y_2 - x_3y_1\beta(y_2) \\ &= x_1x_2y_3 - x_1y_2x_3 - x_2x_1y_3 + x_2y_1x_3 + x_3x_1y_2 - x_3y_1x_2 \\ &= 0. \end{aligned}$$

The inverse transgression map for an element  $g = (a_1, a_2, a_3) \in (\mathbb{Z}_p)^3$  evaluated in the polynomial  $p$  gives

$$\begin{aligned} (5-9) \quad \tau_g(p) &= \tau_g(x_1y_2y_3) - \tau_g(x_2y_1y_3) + \tau_g(x_3y_1y_2) \\ &= x_1(a_3y_2 - a_2y_3) - x_2(a_3y_1 - a_1y_3) + x_3(a_2y_1 - a_1y_2) \\ &= a_1(x_2y_3 - x_3y_2) + a_2(x_3y_1 - x_1y_3) + a_3(x_1y_2 - x_2y_1). \end{aligned}$$

**Lemma 5.7.** *Let  $g = (a_1, a_2, a_3)$  and  $h = (b_1, b_2, b_3)$  be elements in  $G = (\mathbb{Z}_p)^3$ . The double inverse transgression map of  $p$  is equal to*

$$(5-10) \quad \tau_h\tau_g(p) = [(a_1, a_2, a_3) \times (b_1, b_2, b_3)] \cdot (x_1, x_2, x_3).$$

**Remark 5.8.** With  $n = 3$ , this example shows that for  $n \geq 3$  the inverse transgression map is nontrivial. We can always consider the element  $p(x_1, \dots, x_n, y_1, \dots, y_n) = x_iy_jy_k - x_jy_iy_k + x_ky_iy_j$  as being in  $H^4((\mathbb{Z}_p)^n; \mathbb{Z})$ . By calculations similar to those leading to (5-9), we can prove that  $\beta(p) = 0$  while  $\tau_g(p) \neq 0$  for  $g \in (\mathbb{Z}_p)^n$ .

By using the inverse transgression map for the group  $(\mathbb{Z}_p)^n$  presented above and by the decomposition formula presented in Theorem 3.6 in [\[Becerra and Uribe 2009\]](#), we can calculate the explicit structure of the twisted orbifold K-theory for the orbifold  $[*/(\mathbb{Z}_p)^3]$  and the twist element  $\alpha$  as the element in  $H^3((\mathbb{Z}_p)^3; S^1)$  associated to the polynomial defined in (5-8) via the isomorphism  $H^3((\mathbb{Z}_p)^3; S^1) \cong H^4((\mathbb{Z}_p)^3; \mathbb{Z})$ . Note that in this case

$${}^{\alpha}K_{\text{orb}}([*/(\mathbb{Z}_p)^3]) = \bigoplus_{g \in (\mathbb{Z}_p)^3} {}^{\alpha_g}K_{(\mathbb{Z}_p)^3}(\ast) \cong \bigoplus_{g \in (\mathbb{Z}_p)^3} R_{\alpha_g}((\mathbb{Z}_p)^3).$$

Now, for each  $g \in \mathbb{Z}_p$ , the decomposition formula implies

$$R_{\alpha_g}((\mathbb{Z}_p)^3) \otimes \mathbb{Q} \cong \prod_{g, h \in (\mathbb{Z}_p)^3} (\mathbb{Q}(\zeta_p)_{h, \alpha_g})^{(\mathbb{Z}_p)^3},$$

where  $\zeta_p$  is a  $p$ -root of the unity. Note that the action of  $(\mathbb{Z}_p)^3$  on  $\mathbb{Q}(\zeta_p)_{h, \alpha_g}$  is to multiply by the double inverse transgression map evaluated on  $k \in (\mathbb{Z}_p)^3$ ,  $\tau_h(\alpha_g)(k)$ .

By [Lemma 5.7](#), we get

$$(5-11) \quad (\mathbb{Q}(\zeta_p)_{h, \alpha_g})^{(\mathbb{Z}_p)^3} = \begin{cases} \mathbb{Q}(\zeta_p) & \text{if } g = \lambda h, \lambda \in \mathbb{Z}_p, \\ 0 & \text{else.} \end{cases}$$

So, for  $h \neq 0$ , we have

$$R_{\alpha_g}((\mathbb{Z}_p)^3) \otimes \mathbb{Q} = \prod_{\lambda \in \mathbb{Z}_p} \mathbb{Q}(\zeta_p),$$

while for  $g = 0$ , we get

$$R_{\alpha_1}((\mathbb{Z}_p)^3) \otimes \mathbb{Q} = \prod_{\lambda \in (\mathbb{Z}_p)^3} \mathbb{Q}(\zeta_p).$$

Then, the twisted orbifold K-theory module for the orbifold  $[*/(\mathbb{Z}_p)^3]$  turns out to be

$${}^\alpha K_{\text{orb}}([*/(\mathbb{Z}_p)^3]) \otimes \mathbb{Q} = \prod_{\lambda \in \mathbb{Z}_p} \mathbb{Q}(\zeta_p) \oplus \prod_{\lambda \in (\mathbb{Z}_p)^3} \mathbb{Q}(\zeta_p)$$

and the product structure is defined via the product of the elements in  $\mathbb{Q}(\zeta_p)$ .

## 6. Final remarks

With the result presented in [Section 4](#) about the Grothendieck ring associated to the semigroup of representations of the twisted Drinfeld double  $D^\omega(G)$  and the twisted orbifold K-theory, we found a nice relation between two structures coming from different sources. As we already said, the orbifold  $[*/G]$  is a particular case of a more general kind of orbifolds obtained by the almost free action of a compact Lie group  $G$  on a compact manifold  $M$ . With a little more structure, the stringy product introduced in [Section 3](#) can be extended to a stringy product on the module  ${}^\alpha K_{\text{orb}}([M/G])$  (in the same way as in [\[Becerra and Uribe 2009\]](#)), where  $[M/G]$  denotes the orbifold structure obtained by the almost free action (see [\[Adem and Ruan 2003\]](#) for the details of this structure). Therefore, under suitable hypotheses we can think about the twisted orbifold K-theory  ${}^\omega K_{\text{orb}}([M/G])$  as a more general object which coincides with the Grothendieck ring  $R(D^\omega(G))$  if  $G$  is a finite group and  $M = \{*\}$ . Nevertheless, we shall explore the interpretation and consequences of this more general object. Next, we focus our attention on the results obtained in [Section 5](#), where we establish an explicit relation between the twisted orbifold K-theories of the orbifolds  $[*/H]$  and  $[*/(\mathbb{Z}_p)^n]$ , where  $H$  is a particular extraspecial  $p$ -group. In the same spirit, we look for some general relation between the twisted orbifold K-theories  ${}^\alpha K_{\text{orb}}([M/G])$  and  ${}^\beta K_{\text{orb}}([M/K])$  of the orbifolds  $[M/G]$  and  $[M/K]$ , for suitable twistings  $\alpha \in H^3(G; S^1)$  and  $\beta \in H^3(H; S^1)$ , and appropriate actions of the finite groups  $G$  and  $K$  on a compact manifold  $M$ . In the same way,

we hope that some analogous results may be obtained if  $G$  and  $K$  are compact Lie groups acting almost freely on a compact manifold  $M$ . By our preliminary observations, in order to obtain such results, some hypothesis on the almost free actions of the compact Lie groups  $G$  and  $K$  must be added.

## References

- [Adem and Ruan 2003] A. Adem and Y. Ruan, “Twisted orbifold  $K$ -theory”, *Comm. Math. Phys.* **237**:3 (2003), 533–556. [MR 2004e:19004](#) [Zbl 1051.57022](#)
- [Adem et al. 2007] A. Adem, Y. Ruan, and B. Zhang, “A stringy product on twisted orbifold  $K$ -theory”, *Morfismos* **11**:2 (2007), 33–64. [arXiv math/0605534](#)
- [Becerra and Uribe 2009] E. Becerra and B. Uribe, “Stringy product on twisted orbifold  $K$ -theory for abelian quotients”, *Trans. Amer. Math. Soc.* **361**:11 (2009), 5781–5803. [MR 2011b:19008](#) [Zbl 1191.14068](#)
- [Carey and Wang 2008] A. L. Carey and B.-L. Wang, “Thom isomorphism and push-forward map in twisted  $K$ -theory”, *J. K-Theory* **1**:2 (2008), 357–393. [MR 2009g:55005](#) [Zbl 1204.55006](#)
- [tom Dieck 1987] T. tom Dieck, *Transformation groups*, de Gruyter Studies in Mathematics **8**, de Gruyter, Berlin, 1987. [MR 89c:57048](#) [Zbl 0611.57002](#)
- [Dijkgraaf et al. 1991] R. Dijkgraaf, V. Pasquier, and P. Roche, “Quasi Hopf algebras, group cohomology and orbifold models”, *Nuclear Phys. B Proc. Suppl.* **18**:2 (1991), 60–72. [MR 92m:81238](#) [Zbl 0957.81670](#)
- [Drinfel’d 1987] V. G. Drinfel’d, “Quantum groups”, pp. 798–820 in *Proceedings of the International Congress of Mathematicians* (Berkeley, CA, 1986), vol. 1, edited by A. M. Gleason, Amer. Math. Soc., Providence, RI, 1987. [MR 89f:17017](#) [Zbl 0667.16003](#)
- [Goff et al. 2007] C. Goff, G. Mason, and S.-H. Ng, “On the gauge equivalence of twisted quantum doubles of elementary abelian and extra-special 2-groups”, *J. Algebra* **312**:2 (2007), 849–875. [MR 2008d:16057](#) [Zbl 1171.16021](#)
- [Karpilovsky 1993] G. Karpilovsky, *Group representations*, vol. 2, North-Holland Mathematics Studies **177**, North-Holland, Amsterdam, 1993. [MR 94f:20001](#) [Zbl 0778.20001](#)
- [Nabi Duman 2009] A. Nabi Duman, “The fusion algebra of an extraspecial  $p$ -group”, *J. Math. Phys.* **50**:8 (2009), Article ID #082301. [MR 2010h:20023](#) [Zbl 1203.81077](#)
- [Naidu 2007] D. Naidu, “Categorical Morita equivalence for group-theoretical categories”, *Comm. Algebra* **35**:11 (2007), 3544–3565. [MR 2008j:18007](#) [Zbl 1143.18009](#)
- [Naidu and Nikshych 2008] D. Naidu and D. Nikshych, “Lagrangian subcategories and braided tensor equivalences of twisted quantum doubles of finite groups”, *Comm. Math. Phys.* **279**:3 (2008), 845–872. [MR 2009b:16092](#) [Zbl 1139.16028](#)
- [Segal 1968] G. Segal, “Equivariant  $K$ -theory”, *Inst. Hautes Études Sci. Publ. Math.* **34** (1968), 129–151. [MR 38 #2769](#) [Zbl 0199.26202](#)
- [Vafa 2001] C. Vafa, “Superstrings and topological strings at large  $N$ ”, *J. Math. Phys.* **42**:7 (2001), 2798–2817. [MR 2002h:81227](#) [Zbl 1060.81594](#)
- [Wasserman 1969] A. G. Wasserman, “Equivariant differential topology”, *Topology* **8** (1969), 127–150. [MR 40 #3563](#) [Zbl 0215.24702](#)
- [Willerton 2008] S. Willerton, “The twisted Drinfeld double of a finite group via gerbes and finite groupoids”, *Algebr. Geom. Topol.* **8**:3 (2008), 1419–1457. [MR 2009g:57050](#) [Zbl 1154.57029](#)

[Witherspoon 1996] S. J. Witherspoon, “The representation ring of the twisted quantum double of a finite group”, *Canad. J. Math.* **48**:6 (1996), 1324–1338. [MR 97m:16078](#) [Zbl 0876.16027](#)

[Witten 1998] E. Witten, “D-branes and K-theory”, *J. High Energy Phys.* **12** (1998), Article ID #19. [MR 2000e:81151](#) [Zbl 0959.81070](#)

Received May 16, 2012. Revised November 28, 2012.

MARIO VELÁSQUEZ  
UNIVERSIDAD DE LOS ANDES  
DEPARTAMENTO DE MATEMÁTICAS  
CARRERA 1 No 18A–12 EDIFICIO H  
BOGOTÁ  
COLOMBIA

[ma.velasquez109@uniandes.edu.co](mailto:ma.velasquez109@uniandes.edu.co)

EDWARD BECERRA  
DEPARTMENT OF MATHEMATICS  
UNIVERSIDAD NACIONAL DE COLOMBIA  
CARRERA 30 CALLE 45, CIUDAD UNIVERSITARIA  
BOGOTÁ  
COLOMBIA

[esbecerrar@unal.edu.co](mailto:esbecerrar@unal.edu.co)

HERMES MARTINEZ  
ESCUELA DE MATEMÁTICAS  
UNIVERSIDAD SERGIO ARBOLEDA  
CALLE 74 No. 14–14  
BOGOTÁ  
COLOMBIA

[hermes.martinez@ima.usergioarboleda.edu.co](mailto:hermes.martinez@ima.usergioarboleda.edu.co)

# PACIFIC JOURNAL OF MATHEMATICS

[msp.org/pjm](http://msp.org/pjm)

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Don Blasius  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Vijayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published ten to twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 264    No. 2    August 2013

---

On 4-manifolds, folds and cusps	257
STEFAN BEHRENS	
Thin $r$ -neighborhoods of embedded geodesics with finite length and negative Jacobi operator are strongly convex	307
PHILIPPE DELANOË	
Eigenvalues of perturbed Laplace operators on compact manifolds	333
ASMA HASSANNEZHAD	
Four equivalent versions of nonabelian gerbes	355
THOMAS NIKOLAUS and KONRAD WALDORF	
On nonlinear nonhomogeneous resonant Dirichlet equations	421
NIKOLAOS S. PAPAGEORGIOU and GEORGE SMYRLIS	
A geometric model of an arbitrary real closed field	455
STANISŁAW SPODZIEJA	
Twisted K-theory for the orbifold $[*/G]$	471
MARIO VELÁSQUEZ, EDWARD BECERRA and HERMES MARTINEZ	
Linear restriction estimates for the wave equation with an inverse square potential	491
JUNYONG ZHANG and JIQIANG ZHENG	