

*Pacific
Journal of
Mathematics*

RANK GRADIENT OF SMALL COVERS

DARLAN GIRÃO

Volume 266 No. 1

November 2013

RANK GRADIENT OF SMALL COVERS

DARLAN GIRÃO

We prove that if $M \rightarrow P$ is a small cover of a compact right-angled hyperbolic polyhedron then M admits a cofinal tower of finite sheeted covers with positive rank gradient. As a corollary, if $\pi_1(M)$ is commensurable with the reflection group of P , then M admits a cofinal tower of finite sheeted covers with positive rank gradient.

1. Introduction

Let P^n be an n -dimensional simple convex polytope. Here P^n is *simple* if the number of codimension-one faces meeting at each vertex is n . Equivalently, the dual K_P of its boundary complex ∂P^n is an $(n-1)$ -dimensional simplicial sphere. A *small cover* of P^n is an n -dimensional manifold endowed with an action of the group \mathbb{Z}_2^n whose orbit space is P^n . The notion of small cover was introduced and studied by Davis and Januszkiewicz [1991]. We will be dealing mostly with 3-dimensional polytopes. In the case P is a compact right-angled polyhedron in \mathbb{H}^3 then Andreev's theorem [1970] implies that all vertices have valence three and in particular P is a simple convex polytope.

Let G be a finitely generated group. The *rank of G* is the minimal number of elements needed to generate G , and is denoted by $\text{rk}(G)$. If G_j is a finite index subgroup of G , the Reidemeister–Schreier process [Lyndon and Schupp 1977] gives an upper bound on the rank of G_j .

$$\text{rk}(G_j) - 1 \leq [G : G_j](\text{rk}(G) - 1)$$

Lackenby [2006] introduced the notion of *rank gradient*. Given a finitely generated group G and a collection $\{G_j\}$ of finite index subgroups, the rank gradient of the pair $(G, \{G_j\})$ is defined by

$$\text{rgr}(G, \{G_j\}) = \lim_{j \rightarrow \infty} \frac{\text{rk}(G_j) - 1}{[G : G_j]}.$$

We say that the collection of finite index subgroups $\{G_j\}$ is *cofinal* if $\bigcap_j G_j = \{1\}$, and we call it a *tower* if $G_{j+1} < G_j$.

MSC2010: 57M05.

Keywords: rank gradient, hyperbolic 3-manifolds.

In general it is very hard to construct cofinal families $(G, \{G_j\})$ with positive rank gradient. For instance, it seems that only recently the first examples of torsion-free finite covolume Kleinian groups with this property were given in [Girão 2011]. Before stating the main result we need some terminology.

If M is a finite volume hyperbolic 3-manifold, we call the family of covers $\{M_j \rightarrow M\}$ *cofinal* (resp. a *tower*) if $\{\pi_1(M_j)\}$ is cofinal (resp. a tower). By the rank gradient of the pair $(M, \{M_j\})$, $\text{rgr}(M, \{M_j\})$, we mean the rank gradient of $(\pi_1(M), \{\pi_1(M_j)\})$.

Theorem 1.1. *Let $M \rightarrow P$ be a small cover of a compact, right-angled hyperbolic polyhedron of dimension 3. Then M admits a cofinal tower of finite sheeted covers $\{M_j \rightarrow M\}$ with positive rank gradient.*

We remark here that this is not true for 3-dimensional polytopes in general. Let $T^3 \rightarrow C$ be the covering of a cube in 3-dimensional Euclidean space by the 3-torus T^3 . It is easy to see that any subgroup of $\pi_1(T^3) = \mathbb{Z}^3$ has bounded rank and therefore the rank gradient with respect to any tower of covers is zero.

This theorem has the following consequence:

Corollary 1.2. *Let M be a finite volume hyperbolic 3-manifold such that $\pi_1(M)$ is commensurable with the group generated by reflections along the faces of a compact, right-angled hyperbolic polyhedron $P \subset \mathbb{H}^3$. Then M admits a cofinal tower of finite sheeted covers $\{M_j \rightarrow M\}$ with positive rank gradient.*

We note that this corollary is complementary to the results of [Girão 2011], where ideal right-angled polyhedra were considered. The key idea there was to estimate the rank of the fundamental group of the manifolds by estimating their number of cusps. Here the estimates on the rank of the fundamental groups are given in terms of the rank of the mod 2 homology.

The study of the rank of the fundamental group of (finite volume hyperbolic) 3-manifolds has always been a central theme in low-dimensional topology. In recent years the study of the rank gradient for this class of groups has received special attention. Motivated by the seminal paper [Lackenby 2006], for instance, Long, Lubotzky, Reid proved in [Long et al. 2008] that every finite volume hyperbolic 3-manifold has a cofinal tower of covers in which the *Heegaard genus* grows linearly with the degree of the covers. Whether or not the same happens to the rank of their fundamental groups is a major open problem. Another important recent work using these notions is [Abért and Nikolov 2012]. There they connect the problem related to the growth of the rank of π_1 and the growth of the Heegaard genus in a cofinal tower of hyperbolic 3-manifolds to a problem in topological dynamics, the *fixed price problem* (see [Farber 1998; Gaboriau 2000]). These papers have all been motivation for the current work.

2. Small covers

Recall that an n -dimensional convex polytope P^n is simple if the number of codimension-one faces meeting at each vertex is n . Equivalently, the dual K_P of its boundary complex ∂P is an $(n-1)$ -dimensional simplicial complex. A small cover of P is an n -dimensional manifold endowed with an action of the group \mathbb{Z}_2^n whose orbit space is P .

Let K be a finite simplicial complex of dimension $n-1$. For $0 \leq i \leq n-1$, let f_i be the number of i -simplices of K . Define a polynomial $\Phi_K(t)$ of degree n by

$$\Phi_K(t) = (t-1)^n + \sum_{i=0}^{n-1} f_i (t-1)^{n-1-i}$$

and let h_i be the coefficient of t^{n-i} in this polynomial, that is,

$$\Phi_K(t) = \sum_{i=0}^n h_i t^{n-i}.$$

If we restrict to the case where K is the dual K_P of the boundary complex of a convex simple polytope P^n , then one can see that f_i is the number of faces of P^n of codimension $i+1$. Let $h_i(P^n)$ denote the coefficient of t^{n-i} in $\Phi_{K_P}(t)$.

We state one of the main results of [Davis and Januszkiewicz 1991] in our particular setting:

Theorem 2.1. *If $\pi : M^n \rightarrow P^n$ is a small cover of a simple convex polytope P^n and $b_i(M^n, \mathbb{Z}_2)$ is the i -th mod 2 Betti number of M^n , then $b_i(M^n, \mathbb{Z}_2) = h_i(P^n)$.*

As Davis and Januszkiewicz observe, it is somewhat surprising that all mod 2 Betti numbers of a small cover M^n depend on P^n only. They showed that this theorem does not hold for homology groups in general. They proved that small covers of a square Q by tori and Klein bottles are such that the rational Betti numbers are not determined by Q .

When P is a right-angled dodecahedron in \mathbb{H}^3 then [Garrison and Scott 2003] shows that up to homeomorphism there exist exactly 25 small covers of P . Choi [2010] estimates the number of orientable small covers of the n -dimensional cube. Also, if P is a 3-dimensional convex polytope, [Nakayama and Nishimura 2005] proves that P admits an orientable small cover. They also prove that unless P is a 3-simplex, then it admits a nonorientable small cover.

3. Proof of theorem

Theorem 1.1. *Let $M \rightarrow P$ be a small cover of a compact, right-angled hyperbolic polyhedron of dimension 3. Then M admits a cofinal tower of finite sheeted covers $\{M_j \rightarrow M\}$ with positive rank gradient.*

Proof. As observed above, when P is a compact right-angled polyhedron in \mathbb{H}^3 then Andreev's theorem [1970] implies that all vertices have valence three and in particular P is a simple convex polytope. Let V , E and F denote the number of vertices, edges and faces, respectively, of a 3-dimensional simple polyhedron P . Straightforward computations show that

$$\Phi_{K_P}(t) = t^3 + (F - 3)t^2 + (3 - 2F + E)t + (V - E + F - 1)$$

and thus $h_0(P) = 1$, $h_1(P) = F - 3$, $h_2(P) = 3 - 2F + E$ and $h_3(P) = V - E + F - 1$. Since P is simple we also have $E = \frac{3}{2}V$. And since $V - E + F = 2$ (∂P is topologically a sphere) this gives $F = \frac{1}{2}V + 2$ and therefore $h_1(P) = \frac{1}{2}V - 1$.

The strategy involved in the proof is similar to the proof of the main theorem in [Girão 2011]. Given $P \in \mathbb{H}^3$, construct a family of polyhedra

$$P = P_0, P_1, \dots, P_j, \dots$$

such that P_{j+1} is obtained from P_j by reflecting P_j along one of its faces. This must be done in a way such that the following holds: If $x \in \mathbb{H}^3$, then there exists j sufficiently large so that x lies in the interior of P_j . This means that the family $\{P_j\}$ is an exhaustion of \mathbb{H}^3 . Denote by G_j the group generated by reflections along the faces of P_j . If the family $\{P_j\}$ is constructed as above, then it is easy to see that $G_{j+1} < G_j$ (with index 2) and it can be shown that the tower $\{G_j\}$ is cofinal (see [Agol 2008]). We refer the reader to [Girão 2011] for a detailed proof of this fact.

Now let $M \rightarrow P$ be a small cover of P , and let $M_j \rightarrow M$ be the cover corresponding to the group $\pi_1(M) \cap G_j$. Recall that the degree of the cover $M \rightarrow P$ is 2^3 .

Lemma 3.2. $[\pi_1(M_j) : \pi_1(M_{j+1})] = 2$.

Proof of lemma. First observe that $[G_j : G_{j+1}] = 2$. Since $\pi_1(M_1) = G_1 \cap \pi_1(M)$, we must have $[\pi_1(M) : \pi_1(M_1)] \leq 2$. If this index were 1, then it would mean that $\pi_1(M_1) = \pi_1(M) < G_1$ from which would follow that M_1 is a manifold cover of the simple polyhedron P_1 of degree 2^2 . But this is not possible, since any manifold cover of a 3-dimensional simple polyhedron must have degree at least 2^3 (see [Davis and Januszkiewicz 1991; Garrison and Scott 2003]). The remaining cases follow by induction. \square

Since $[G_j : G_{j+1}] = 2$, from the above lemma and an inductive argument we see that $M_j \rightarrow P_j$ is a cover of degree 2^3 . In particular this implies that M_j is a small cover of P_j . From Theorem 2.1 we have

$$b_1(M_j, \mathbb{Z}_2) = h_1(P_j).$$

Denote by V_j the number of vertices of P_j . From the computations of h_1 ,

$$b_1(M_j, \mathbb{Z}_2) = h_1(P_j) = \frac{1}{2}V_j - 1.$$

Also note that a lower bound for $\text{rk}(\pi_1(M_j))$ is $b_1(M_j, \mathbb{Z}_2)$ and thus

$$\text{rk}(\pi_1(M_j)) \geq \frac{1}{2}V_j - 1.$$

We also have $[\pi_1(M) : \pi_1(M_j)] = 2^j$. Therefore

$$\text{rgr}(\pi_1(M), \{\pi_1(M_j)\}) = \lim_{j \rightarrow \infty} \frac{\text{rk}(\pi_1(M_j)) - 1}{[\pi_1(M) : \pi_1(M_j)]} \geq \lim_{j \rightarrow \infty} \frac{V_j - 3}{2^{j+1}}.$$

We thus need to show that V_j is of magnitude 2^j . This follows from the next result:

Theorem 3.3 [Atkinson 2009]. *There exist constants $C, D > 0$ such that if P is a compact right-angled polyhedron in \mathbb{H}^3 with V vertices then*

$$C(V - 8) \leq \text{vol}(P) \leq D(V - 10).$$

We now observe that, in our setting, $\text{vol}(P_j) = 2^j \text{vol}(P)$ and thus

$$D(V_j - 10) \geq 2^j \text{vol}(P) \geq 2^j C(V - 8)$$

which gives

$$V_j \geq 2^j \frac{C}{D}(V - 8) + 10,$$

where V is the number of vertices in P . Also, the second inequality in Atkinson's theorem provides $V > 8$. \square

4. Extending the examples

Theorem 1.1 has an interesting corollary, which complements the family of manifolds provided in [Girão 2011].

Corollary 1.2. *Let N be a closed hyperbolic 3-manifold such that $\pi_1(N)$ is commensurable with the group generated by reflections along the faces of a compact, right-angled hyperbolic polyhedron $P \subset \mathbb{H}^3$. Then N admits a cofinal tower of finite sheeted covers $\{N_j \rightarrow N\}$ with positive rank gradient.*

Proof. First we note that, by passing to a finite cover, we may assume N is orientable. Note also that [Nakayama and Nishimura 2005] implies orientable small covers of P exist and therefore N is commensurable with a small cover $M \rightarrow P$. Let N' be the manifold cover of both M and N corresponding to the group $\pi_1(M) \cap \pi_1(N)$. Consider now $N_j \rightarrow N$ corresponding to the group $\pi_1(N') \cap G_j$, where the family $\{G_j\}$ is given as in the proof of Theorem 1.1. Consider also $\{M_j\}$, the tower where M_j is a small cover of P_j , as in the proof of Theorem 1.1.

Note that $\pi_1(N_j) = \pi_1(N') \cap G_j = \pi_1(N') \cap \pi_1(M_j) < \pi_1(M_j)$ and therefore we have the following diagram of covers, where the labels in the arrows indicate

the degree of the cover:

$$\begin{array}{ccccccc}
 P & \xleftarrow{2} & P_1 & \xleftarrow{2} & \cdots & \xleftarrow{2} & P_j & \xleftarrow{2} & \cdots \\
 \uparrow & & \uparrow & & & & \uparrow & & \\
 & & 2^3 & & & & 2^3 & & \\
 & & \uparrow & & & & \uparrow & & \\
 M & \xleftarrow{2} & M_1 & \xleftarrow{2} & \cdots & \xleftarrow{2} & M_j & \xleftarrow{2} & \cdots \\
 \uparrow & & \uparrow & & & & \uparrow & & \\
 N & \xleftarrow{\quad} & N' & \xleftarrow{\quad} & N_1 & \xleftarrow{\quad} & \cdots & \xleftarrow{\quad} & N_j & \xleftarrow{\quad} & \cdots
 \end{array}$$

Agol, Culler and Shalen proved:

Theorem 4.2 ([Agol et al. 2006]; see also [Shalen 2007]). *Let M be a closed, orientable hyperbolic 3-manifold such that $b_1(M, \mathbb{Z}_p) = r$ for a given prime p . Then for any finite sheeted covering space M' of M , $b_1(M', \mathbb{Z}_p) \geq r - 1$.*

We thus have

$$\text{rk}(\pi_1(N_j)) \geq b_1(N_j, \mathbb{Z}_2) \geq b_1(M_j, \mathbb{Z}_2) - 1 = \frac{1}{2}V_j - 2$$

and therefore all we need to do is show that $[\pi_1(N) : \pi_1(N_j)]$ grows at most as fast as 2^j . But from the above diagram we see that $[\pi_1(N_j) : \pi_1(N_{j+1})] \leq 2$ and we are done. \square

Acknowledgements

The author thanks Alan Reid for suggesting he look at [Davis and Januszkiewicz 1991]. He also thanks the Erwin Schrödinger International Institute for Mathematical Physics for the hospitality and financial support during his visit in the occasion of the workshop “Golod–Shafarevich Groups and Algebras and Rank Gradient”. Finally, he would like to thank the organizers of this event.

References

- [Abért and Nikolov 2012] M. Abért and N. Nikolov, “Rank gradient, cost of groups and the rank versus Heegaard genus problem”, *J. Eur. Math. Soc.* **14**:5 (2012), 1657–1677. MR 2966663 Zbl 06095894
- [Agol 2008] I. Agol, “Criteria for virtual fibering”, *J. Topol.* **1**:2 (2008), 269–284. MR 2009b:57033 Zbl 1148.57023
- [Agol et al. 2006] I. Agol, M. Culler, and P. B. Shalen, “Dehn surgery, homology and hyperbolic volume”, *Algebr. Geom. Topol.* **6** (2006), 2297–2312. MR 2008f:57024 Zbl 1129.57019
- [Andreev 1970] E. M. Andreev, “Convex polyhedra in Lobachevski spaces”, *Mat. Sb. (N.S.)* **81 (123)** (1970), 445–478. In Russian; translated in *Math. USSR Sbornik* **10**:3 (1970), 413–440. MR 41 #4367 Zbl 0217.46801
- [Atkinson 2009] C. K. Atkinson, “Volume estimates for equiangular hyperbolic Coxeter polyhedra”, *Algebr. Geom. Topol.* **9**:2 (2009), 1225–1254. MR 2010k:57035 Zbl 1170.57012

- [Choi 2010] S. Choi, “The number of orientable small covers over cubes”, *Proc. Japan Acad. Ser. A Math. Sci.* **86**:6 (2010), 97–100. MR 2011i:57046 Zbl 1198.37074
- [Davis and Januszkiewicz 1991] M. W. Davis and T. Januszkiewicz, “Convex polytopes, Coxeter orbifolds and torus actions”, *Duke Math. J.* **62**:2 (1991), 417–451. MR 92i:52012 Zbl 0733.52006
- [Farber 1998] M. Farber, “Geometry of growth: Approximation theorems for L^2 invariants”, *Math. Ann.* **311**:2 (1998), 335–375. MR 2000b:58042 Zbl 0911.53026
- [Gaboriau 2000] D. Gaboriau, “Coût des relations d’équivalence et des groupes”, *Invent. Math.* **139**:1 (2000), 41–98. MR 2001f:28030 Zbl 0939.28012
- [Garrison and Scott 2003] A. Garrison and R. Scott, “Small covers of the dodecahedron and the 120-cell”, *Proc. Amer. Math. Soc.* **131**:3 (2003), 963–971. MR 2003h:57018 Zbl 1009.57019
- [Girão 2011] D. Girão, “Rank gradient in cofinal towers of certain Kleinian groups”, preprint, 2011. To appear in *Groups, Geometry, and Dynamics*. arXiv 1102.4281
- [Lackenby 2006] M. Lackenby, “Heegaard splittings, the virtually Haken conjecture and property (τ) ”, *Invent. Math.* **164**:2 (2006), 317–359. MR 2007c:57030 Zbl 1110.57015
- [Long et al. 2008] D. D. Long, A. Lubotzky, and A. W. Reid, “Heegaard genus and property τ for hyperbolic 3-manifolds”, *J. Topol.* **1**:1 (2008), 152–158. MR 2008j:57036 Zbl 1158.57018
- [Lyndon and Schupp 1977] R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete **89**, Springer, Berlin, 1977. MR 58 #28182 Zbl 0368.20023
- [Nakayama and Nishimura 2005] H. Nakayama and Y. Nishimura, “The orientability of small covers and coloring simple polytopes”, *Osaka J. Math.* **42**:1 (2005), 243–256. MR 2006a:57023 Zbl 1065.05041
- [Shalen 2007] P. B. Shalen, “Hyperbolic volume, Heegaard genus and ranks of groups”, pp. 335–349 in *Workshop on Heegaard splittings* (Haifa, 2005), edited by C. Gordon and Y. Moriah, Geom. Topol. Monogr. **12**, Geometry & Topology Publications, Coventry, 2007. MR 2009k:57029 Zbl 1140.57009

Received September 3, 2012. Revised January 2, 2013.

DARLAN GIRÃO
DEPARTMENT OF MATHEMATICS
UNIVERSIDADE FEDERAL DO CEARÁ
AV. HUMBERTO MONTE S/N
CAMPUS DO PICI - BLOCO 914
60455-760 FORTALEZA, CE
BRAZIL
dgirao@mat.ufc.br

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 266 No. 1 November 2013

Multiplicity of solutions to the Yamabe problem on collapsing Riemannian submersions	1
RENATO G. BETTIOL and PAOLO PICCIONE	
Rank gradient of small covers	23
DARLAN GIRÃO	
Nonrationality of nodal quartic threefolds	31
KYUSIK HONG	
Supertropical linear algebra	43
ZUR IZHAKIAN, MANFRED KNEBUSCH and LOUIS ROWEN	
Isometry groups among topological groups	77
PIOTR NIEMIEC	
Singularities and Liouville theorems for some special conformal Hessian equations	117
QIANZHONG OU	
Attaching handles to Delaunay nodoids	129
FRANK PACARD and HAROLD ROSENBERG	
Some new canonical forms for polynomials	185
BRUCE REZNICK	
Applications of the deformation formula of holomorphic one-forms	221
QUANTING ZHAO and SHENG RAO	



0030-8730(201311)266:1;1-5