

Pacific Journal of Mathematics

**EIGENVALUES AND ENTROPIES UNDER THE
HARMONIC-RICCI FLOW**

YI LI

Volume 267 No. 1

January 2014

EIGENVALUES AND ENTROPIES UNDER THE HARMONIC-RICCI FLOW

YI LI

In this paper, the author discusses the eigenvalues and entropies under the harmonic-Ricci flow, which is the Ricci flow coupled with the harmonic map flow. We give an alternative proof of results for compact steady and expanding harmonic-Ricci breathers. In the second part, we derive some monotonicity formulas for eigenvalues of the Laplacian under the harmonic-Ricci flow. Finally, we obtain the first variation of the shrinker and expanding entropies of the harmonic-Ricci flow.

1. Introduction	141
2. Notation and commuting identities	148
3. Harmonic-Ricci flow and the evolution equations	150
4. Entropies for harmonic-Ricci flow	150
5. Compact steady harmonic-Ricci breathers	153
6. Compact expanding harmonic-Ricci breathers	154
7. Eigenvalues of the Laplacian under the harmonic-Ricci flow	159
8. Eigenvalues of the Laplacian-type under the harmonic-Ricci flow	166
9. Another formula for $d\lambda(f(t))/dt$	171
10. The first variation of expander and shrinker entropies	176
Acknowledgements	183
References	183

1. Introduction

Since the successful application of the Ricci flow to topological and geometric problems, several analogous flows have been studied, including the harmonic-Ricci flow [List 2006; Müller 2012], connection Ricci flow [Streets 2008], Ricci–Yang–Mills flow [Streets 2007; 2010; Young 2008], and renormalization group flows [He et al. 2008; Li 2012; Oliynyk et al. 2006; Streets 2009]. In this article, we study the

MSC2010: 53C44, 35K55.

Keywords: Eigenvalue, entropies, harmonic-Ricci flow, harmonic-Ricci breathers.

eigenvalue problems of the harmonic-Ricci flow, which is the following coupled system:

$$(1-1) \quad \frac{\partial}{\partial t} g(t) = -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t),$$

$$(1-2) \quad \frac{\partial}{\partial t} u(t) = \Delta_{g(t)} u(t).$$

For convenience, we introduce a new symmetric 2-tensor $\mathcal{S}_{g(t), u(t)}$ whose components S_{ij} are defined by

$$S_{ij} := R_{ij} - 2\partial_i u \partial_j u.$$

Its trace is $S_{g(t), u(t)} := g^{ij} S_{ij} = R_{g(t)} - 2|\nabla_{g(t)} u(t)|_{g(t)}^2$.

Suppose that M is a compact Riemannian manifold. For any Riemannian metric g and any smooth functions u, f , we have a number of functionals:

$$\mathcal{F}(g, u, f) = \int_M (R_g + |\nabla_g f|_g^2 - 2|\nabla_g u|_g^2) e^{-f} dV_g,$$

$$\mathcal{E}(g, u, f) = \int_M (R_g - 2|\nabla_g u|_g^2) e^{-f} dV_g,$$

$$\mathcal{F}_k(g, u, f) = \int_M (kR_g + |\nabla_g f|_g^2 - 2k|\nabla_g u|_g^2) e^{-f} dV_g.$$

List [2006] and Müller [2012] showed that, as in the case of Perelman's $\mathcal{F}$ -functional, under the evolution equation

$$\frac{\partial}{\partial t} g(t) = -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t),$$

$$(1-3) \quad \frac{\partial}{\partial t} u(t) = \Delta_{g(t)} u(t),$$

$$\frac{\partial}{\partial t} f(t) = -\Delta_{g(t)} f(t) - R_{g(t)} + |\nabla_{g(t)} f(t)|_{g(t)}^2 + 2|\nabla_{g(t)} u(t)|_{g(t)}^2,$$

the evolution equation for the $\mathcal{F}$ -functional is

$$(1-4) \quad \begin{aligned} \frac{d}{dt} \mathcal{F}(g(t), u(t), f(t)) &= 2 \int_M |\mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 4 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)}, \end{aligned}$$

which is nonnegative. Based on (1-4), we derive the following.

Theorem 1.1. *Under the evolution equation (1-3), one has*

$$(1-5) \quad \begin{aligned} \frac{d}{dt} \mathcal{E}(g(t), u(t), f(t)) \\ = 2 \int_M |\mathcal{S}_{g(t), u(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 4 \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)}, \end{aligned}$$

and

$$\begin{aligned}
 (1-6) \quad & \frac{d}{dt} \mathcal{F}_k(g(t), u(t), f(t)) \\
 &= 2(k-1) \int_M |\mathcal{S}_{g(t), u(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 2 \int_M |\mathcal{S}_{g(t), u(t)} \\
 &\quad + \nabla_{g(t)}^2 f(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 4(k-1) \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\
 &\quad + 4 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)}.
 \end{aligned}$$

As a corollary we give a new proof of the following result.

Corollary 1.2. *There is no compact steady harmonic-Ricci breather unless the manifold $(M, g(t))$ is Ricci-flat and $u(t)$ is a constant.*

To deal with the expanding harmonic-Ricci breather, we need the functionals

$$\begin{aligned}
 \mathcal{L}_+(g, u, \tau, f) &= \tau^2 \int_M \left(R_g + \frac{n}{2\tau} + \Delta_g f - 2|\nabla_g u|_g^2 \right) e^{-f} dV_g, \\
 \mathcal{L}_{+,k}(g, u, \tau, f) &= \tau^2 \int_M \left(k \left(R_g + \frac{n}{2\tau} \right) + \Delta_g f - 2k|\nabla_g u|_g^2 \right) e^{-f} dV_g.
 \end{aligned}$$

Under the evolution equation

$$\begin{aligned}
 \frac{\partial}{\partial t} g(t) &= -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t), \\
 \frac{\partial}{\partial t} u(t) &= \Delta_{g(t)} u(t), \\
 \frac{\partial}{\partial t} f(t) &= -\Delta_{g(t)} f(t) + |\nabla_{g(t)} f(t)|_{g(t)}^2 - R_{g(t)} + 2|\nabla_{g(t)} u(t)|_{g(t)}^2, \\
 \frac{d}{dt} \tau(t) &= 1,
 \end{aligned}$$

we have:

Theorem 1.3. *Under the evolution equation, one has*

$$\begin{aligned}
 (1-7) \quad & \frac{d}{dt} \mathcal{L}_+(g(t), u(t), \tau(t), f(t)) \\
 &= 2\tau(t)^2 \int_M \left| \mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t) + \frac{1}{2\tau(t)} g(t) \right|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\
 &\quad + 4\tau(t)^2 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)}
 \end{aligned}$$

and

$$\begin{aligned}
 (1-8) \quad & \frac{d}{dt} \mathcal{L}_{+,k}(g(t), u(t), \tau(t), f(t)) \\
 &= 2\tau(t)^2 \int_M |\mathcal{G}_{g(t),u(t)} + \nabla_{g(t)}^2 f(t) + \frac{1}{2\tau(t)} g(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\
 &\quad + 2(k-1)\tau(t)^2 \int_M |\mathcal{G}_{g(t),u(t)} + \frac{1}{2\tau(t)} g(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\
 &\quad + 4\tau(t)^2 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\
 &\quad + 4(k-1)\tau(t)^2 \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)}.
 \end{aligned}$$

As a corollary, we obtain a new proof of the following.

Corollary 1.4. *There is no expanding harmonic-Ricci breather on compact Riemannian manifolds unless the manifold M is an Einstein manifold and $u(t)$ a constant.*

The second part of this paper focuses on the eigenvalue of the Laplacian operator under the harmonic-Ricci flow.

Theorem 1.5. *If $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $\lambda(t)$ denotes the eigenvalue of the Laplacian $\Delta_{g(t)}$ with eigenfunction $f(t)$,*

$$\begin{aligned}
 (1-9) \quad & \frac{d}{dt} \lambda(t) \cdot \int_M f(t)^2 dV_{g(t)} \\
 &= \lambda(t) \int_M S_{g(t),u(t)} f(t)^2 dV_{g(t)} - \int_M S_{g(t),u(t)} |\nabla_{g(t)} f|_{g(t)}^2 dV_{g(t)} \\
 &\quad + 2 \int_M \langle \mathcal{G}_{g(t),u(t)}, df(t) \otimes df(t) \rangle_{g(t)} dV_{g(t)}.
 \end{aligned}$$

Equation (1-9) is a general formula to describe the evolution of $\lambda(t)$ under the harmonic-Ricci flow. Under a curvature assumption, we can derive some monotonicity formulas for the eigenvalue $\lambda(t)$. Set

$$(1-10) \quad S_{\min}(0) := \min_{x \in M} S_{g(0),u(0)}(x),$$

the minimum of $S_{g(t),u(t)}$ over M at the time 0.

Theorem 1.6. *Let $(g(t), u(t))_{t \in [0, T]}$ be a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$. Suppose that $\mathcal{G}_{g(t),u(t)} - \alpha S_{g(t),u(t)} g(t) \geq 0$ along the harmonic-Ricci flow for some $\alpha \geq \frac{1}{2}$.*

- (1) If $S_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.
- (2) If $S_{\min}(0) > 0$, the quantity

$$\left(1 - \frac{2}{n} S_{\min}(0) t\right)^{n\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for $T \leq n/(2S_{\min}(0))$.

- (3) If $S_{\min}(0) < 0$, the quantity

$$\left(1 - \frac{2}{n} S_{\min}(0) t\right)^{n\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.

Corollary 1.7. Let $(g(t), u(t))_{t \in [0, T]}$ be a solution of the harmonic-Ricci flow on a compact Riemannian surface Σ and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$.

- (1) Suppose that $\text{Ric}_{g(t)} \leq \epsilon du(t) \otimes du(t)$ where

$$\epsilon \leq 4 \frac{1 - \alpha}{1 - 2\alpha}, \quad \alpha > \frac{1}{2}.$$

- (i) If $S_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.
- (ii) If $S_{\min}(0) > 0$, the quantity

$$(1 - S_{\min}(0)t)^{2\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for $T \leq 1/S_{\min}(0)$.

- (iii) If $S_{\min}(0) < 0$, the quantity

$$(1 - S_{\min}(0)t)^{2\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.

- (2) Suppose that

$$|\nabla_{g(t)} u(t)|_{g(t)}^2 g(t) \geq 2 du(t) \otimes du(t).$$

- (i) If $S_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.
- (ii) If $S_{\min}(0) > 0$, the quantity

$$(1 - S_{\min}(0)t) \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for $T \leq 1/S_{\min}(0)$.

(iii) If $S_{\min}(0) < 0$, the quantity

$$(1 - S_{\min}(0)t)\lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.

When we restrict to the Ricci flow, we obtain:

Corollary 1.8. *Let $(g(t))_{t \in [0, T]}$ be a solution of the Ricci flow on a compact Riemannian surface Σ and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$.*

- (1) *If $R_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the Ricci flow for any $t \in [0, T]$.*
- (2) *If $R_{\min}(0) > 0$, the quantity $(1 - R_{\min}(0)t)\lambda(t)$ is nondecreasing along the Ricci flow for $T \leq 1/R_{\min}(0)$.*
- (3) *If $R_{\min}(0) < 0$, the quantity $(1 - R_{\min}(0)t)\lambda(t)$ is nondecreasing along the Ricci flow for any $t \in [0, T]$.*

Remark 1.9. Let $(g(t))_{t \in [0, T]}$ be a solution of the Ricci flow on a compact Riemannian surface Σ with nonnegative scalar curvature and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$. Then $\lambda(t)$ is nondecreasing along the Ricci flow for any $t \in [0, T]$.

Since

$$(1-11) \quad \mu(g, u) := \inf \left\{ \mathcal{F}(g, u, f) \mid f \in C^\infty(M), \int_M e^{-f} dV_g = 1 \right\}$$

is the smallest eigenvalue of the operator $\Delta_{g,u} := -4\Delta_g + R_g - 2|\nabla_g u|_g^2$, we can consider the evolution equation for this eigenvalue under the harmonic-Ricci flow. To the operator $\Delta_{g,u}$ we associate a functional

$$(1-12) \quad \lambda_{g,u}(f) := \int_M f \Delta_{g,u} f dV_g.$$

When f is an eigenfunction of the operator $\Delta_{g,u}$ with the eigenvalue λ and normalized by $\int_X f^2 dV_g = 1$, we obtain $\lambda_{g,u}(f) = \lambda$. Hence it suffices to study the evolution equation for $(d/dt)\lambda_{g,u}(f)$ under the harmonic-Ricci flow.

Theorem 1.10. *Suppose that $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $f(t)$ is an eigenfunction of $\Delta_{g(t), u(t)}$, that is, $\Delta_{g(t), u(t)} f(t) = \lambda(t) f(t)$ (where $\lambda(t)$ is only a function of time t), with the normalized condition $\int_M f(t)^2 dV_{g(t)} = 1$. Then we have*

$$(1-13) \quad \begin{aligned} \frac{d}{dt} \lambda(t) = \frac{d}{dt} \lambda_{g,u}(f(t)) &= \int_M 2 \langle \mathcal{I}_{g(t), u(t)}, df(t) \otimes df(t) \rangle_{g(t)} dV_{g(t)} \\ &\quad + \int_M f(t)^2 (|\mathcal{I}_{g(t), u(t)}|_{g(t)}^2 + 2|\Delta_{g(t)} u(t)|_{g(t)}^2) dV_{g(t)}. \end{aligned}$$

List [2006] proved the nonnegativity of the operator $\mathcal{P}_{g(t),u(t)}$ is preserved by the harmonic-Ricci flow. Hence we get the following.

Corollary 1.11. *If $\text{Ric}_{g(0)} - 2du(0) \otimes du(0) \geq 0$, the eigenvalues of the operator $\Delta_{g(t),u(t)}$ are nondecreasing under the harmonic-Ricci flow.*

Remark 1.12. If we choose $u(t) \equiv 0$, we obtain X. Cao's result [2007].

There is another expression for $d\lambda(t)/dt$.

Theorem 1.13. *Suppose that $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $f(t)$ is an eigenfunction of $\Delta_{g(t),u(t)}$, that is, $\Delta_{g(t),u(t)} f(t) = \lambda(t) f(t)$ (where $\lambda(t)$ is only a function of time t), with the normalized condition $\int_M f(t)^2 dV_{g(t)} = 1$. Then we have*

$$(1-14) \quad \begin{aligned} \frac{d}{dt} \lambda(t) &= \frac{d}{dt} \lambda_{g,u}(f(t)) = \frac{1}{2} \int_M |\mathcal{P}_{g(t),u(t)} + \nabla_{g(t)}^2 \varphi(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} \\ &\quad + \frac{1}{4} \int_M |\mathcal{P}_{g(t),u(t)}|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} + \int_M |\langle du(t), d\varphi(t) \rangle_{g(t)}|^2 e^{-\varphi(t)} dV_{g(t)} \\ &\quad + 2 \int_M |\nabla_{g(t)}^2 u(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} - \int_M \Delta_{g(t)} (|\nabla_{g(t)} u(t)|_{g(t)}^2) e^{-\varphi(t)} dV_{g(t)} \\ &\quad + \frac{1}{4} \int_M |\mathcal{P}_{g(t),u(t)} + 4du(t) \otimes du(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)}, \end{aligned}$$

where $f(t)^2 = e^{-\varphi(t)}$.

Remark 1.14. When $u \equiv 0$, (1-14) reduces to J. Li's formula [2007].

Suppose that M is a compact manifold of dimension n . For any Riemannian metric g , any smooth functions u , f , and any positive number τ , we define

$$(1-15) \quad \mathcal{W}_{\pm}(g, u, f, \tau) := \int_M [\tau(S_g + |\nabla_g f|_g^2) \mp f \pm n] \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g.$$

Set

$$\begin{aligned} \mu_{\pm}(g, u, \tau) &:= \inf \left\{ \mathcal{W}_{\pm}(g, u, f, \tau) \mid f \in C^{\infty}(M), \int_M \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g = 1 \right\}, \\ \nu_{-}(g, u) &:= \inf \{ \mu_{-}(g, u, \tau) \mid \tau > 0 \}, \quad \nu_{+}(g, u) := \sup \{ \mu_{+}(g, u, \tau) \mid \tau > 0 \}. \end{aligned}$$

The first variation of $\nu_{\pm}(g(s), u(s))$ is the following.

Theorem 1.15. *Suppose that (M, g) is a compact Riemannian manifold and u a smooth function on M . Let h be any symmetric covariant 2-tensor on M and set $g(s) := g + sh$. Let v be any smooth function on M and $u(s) := u + sv$. If*

$v_{\pm}(g(s), u(s)) = \mathcal{W}_{\pm}(g(s), u(s), f_{\pm}(s), \tau_{\pm}(s))$ for some smooth functions $f_{\pm}(s)$ with $\int_M e^{-f_{\pm}(s)} dV_g / (4\pi \tau_{\pm}(s))^{n/2} = 1$ and constants $\tau_{\pm}(s) > 0$,

$$(1-16) \quad \frac{d}{ds} \Big|_{s=0} v_{\pm}(g(s), u(s)) = 4\tau_{\pm} \int_M v(\Delta_g u - \langle du, df_{\pm} \rangle_g) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g \\ - \tau_{\pm} \int_M \left(\langle h, \mathcal{P}_{g,u} \rangle_g + \langle h, \nabla_g^2 f \rangle_g \pm \frac{\text{tr}_g h}{2\tau_{\pm}} \right) \frac{e^{-f_{\pm}} dV_g}{(4\pi \tau_{\pm})^{n/2}},$$

where $f_{\pm} := f_{\pm}(0)$ and $\tau_{\pm} := \tau_{\pm}(0)$. In particular, the critical points of $v_{\pm}(\cdot, \cdot)$ satisfy

$$\mathcal{P}_{g,u} + \nabla_g^2 f \pm \frac{1}{2\tau_{\pm}} g = 0, \quad \Delta_g u = \langle du, df_{\pm} \rangle_g.$$

Consequently, if $\mathcal{W}_{\pm}(g, u, f, \tau)$ and $v_{\pm}(g, u)$ achieve their extremum, (M, g) is a gradient expanding and shrinker harmonic-Ricci soliton according to the sign.

Corollary 1.16. Suppose that (M, g) is a compact Riemannian manifold and u a smooth function on M . Let h be any symmetric covariant 2-tensor on M and set $g(s) := g + sh$. Let v be any smooth function on M and $u(s) := u + sv$. If $v_{\pm}(g(s), u(s)) = \mathcal{W}_{\pm}(g(s), u(s), f_{\pm}(s), \tau_{\pm}(s))$ for some smooth function $f_{\pm}(s)$ with $\int_M e^{-f_{\pm}(s)} dV / (4\pi \tau_{\pm}(s))^{n/2} = 1$ and a constant $\tau_{\pm}(s) > 0$, and (g, u) is a critical point of $v_{\pm}(\cdot, \cdot)$, then

$$\text{Ric}_g = \mp \frac{1}{2\tau_{\pm}} g, \quad f_{\pm} \equiv \text{constant}, \quad u \equiv \text{constant}.$$

Thus, if $\mathcal{W}_{\pm}(g, u, \cdot, \cdot)$ achieve their minimum and (g, u) is a critical point of $v_{\pm}(\cdot, \cdot)$, (M, g) is an Einstein manifold and u is a constant function.

Remark 1.17. In the situation of Corollary 1.16, by normalization, we may choose $f_{\pm} = n/2$ and $u = 0$.

2. Notation and commuting identities

Let M be a compact Riemannian manifold of dimension n . For any vector bundle E over M , we denote by $\Gamma(M, E)$ the space of smooth sections of E . Set

$$\odot^2(M) := \{v = (v_{ij}) \in \Gamma(M, T^*M \otimes T^*M) \mid v_{ij} = v_{ji}\}, \\ \odot_+^2(M) := \{g = (g_{ij}) \in \odot^2(M) \mid g_{ij} > 0\}.$$

Thus $\odot^2(M)$ is the space of all symmetric covariant 2-tensors on M , while $\odot_+^2(M)$ is the space of all Riemannian metrics on M . The space of all smooth functions on M is denoted by $C^{\infty}(M)$.

For a given Riemannian metric $g \in \odot_+^2(M)$, the corresponding Levi-Civita connection $\Gamma_g = (\Gamma_{ij}^k)$ is given by

$$(2-1) \quad \Gamma_{ij}^k = \frac{1}{2} g^{k\ell} (\partial_i g_{j\ell} + \partial_j g_{i\ell} - \partial_\ell g_{ij})$$

where $\partial_i := \partial/\partial x^i$ for a local coordinate system $\{x^1, \dots, x^n\}$. The Riemann tensor $\text{Rm}_g = (R_{ijl}^k)$ is determined by

$$(2-2) \quad R_{ijl}^k = \partial_i \Gamma_{jl}^k - \partial_j \Gamma_{il}^k + \Gamma_{ip}^k \Gamma_{jl}^p - \Gamma_{jp}^k \Gamma_{il}^p.$$

The Ricci curvature $\text{Ric}_g = (R_{ij})$ is

$$(2-3) \quad R_{ij} = g^{k\ell} R_{kij}^\ell.$$

The scalar curvature R_g of the metric g now is given by

$$(2-4) \quad R_g = g^{ij} R_{ij}.$$

For any tensor $A = (A_{j_1 \dots j_p}^{k_1 \dots k_q})$ the covariant derivative of A is

$$\nabla_i A_{j_1 \dots j_p}^{k_1 \dots k_q} = \partial_i A_{j_1 \dots j_p}^{k_1 \dots k_q} - \sum_{r=1}^p \Gamma_{ij_r}^m A_{j_1 \dots m \dots j_p}^{k_1 \dots k_q} + \sum_{s=1}^q \Gamma_{im}^{k_s} A_{j_1 \dots j_p}^{k_1 \dots m \dots k_q}.$$

Next we recall the Ricci identity:

$$\nabla_i \nabla_j A_{k_1 \dots k_p}^{\ell_1 \dots \ell_q} - \nabla_j \nabla_i A_{k_1 \dots k_p}^{\ell_1 \dots \ell_q} = \sum_{r=1}^q R_{ijm}^{l_r} A_{k_1 \dots m \dots k_p}^{\ell_1 \dots m \dots \ell_q} - \sum_{s=1}^p R_{ijk_s}^m A_{k_1 \dots m \dots k_p}^{\ell_1 \dots \ell_q}.$$

In particular, for any smooth function $f \in C^\infty(M)$, we have

$$\nabla_i \nabla_j f = \nabla_j \nabla_i f.$$

The Bianchi identities are

$$(2-5) \quad 0 = R_{ijk\ell} + R_{iklj} + R_{iljk},$$

$$(2-6) \quad 0 = \nabla_q R_{ijk\ell} + \nabla_i R_{jqk\ell} + \nabla_j R_{qik\ell},$$

and the contracted Bianchi identities are

$$(2-7) \quad 0 = 2\nabla^j R_{ij} - \nabla_i R_g,$$

$$(2-8) \quad 0 = \nabla_i R_{jk} - \nabla_j R_{ik} + \nabla^\ell R_{\ell kij}.$$

3. Harmonic-Ricci flow and the evolution equations

Motivated by the static Einstein vacuum equation, List [2006] introduced the harmonic-Ricci flow (originally called the Ricci flow coupled with the harmonic map flow). This flow is similar to the Ricci flow and is given by the coupled system

$$(3-1) \quad \frac{\partial}{\partial t} g(t) = -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t),$$

$$(3-2) \quad \frac{\partial}{\partial t} u(t) = \Delta_{g(t)} u(t)$$

for a family of Riemannian metrics $g(t)$ and a family of smooth functions $u(t)$. Locally, we have

$$(3-3) \quad \frac{\partial}{\partial t} g_{ij} = -2R_{ij} + 4\partial_i u \cdot \partial_j u, \quad \frac{\partial}{\partial t} u = \Delta_{g(t)} u(t).$$

Introduce a new symmetric tensor field $\mathcal{S}_{g(t), u(t)} = (S_{ij}) \in \odot^2(M)$,

$$(3-4) \quad S_{ij} := R_{ij} - 2\partial_i u \cdot \partial_j u.$$

Then its trace $S_{g(t), u(t)}$ is equal to

$$(3-5) \quad S_{g(t), u(t)} = g^{ij} S_{ij} = R_{g(t)} - 2|\nabla_{g(t)} u(t)|_{g(t)}^2.$$

The evolution equation for $R_{g(t)}$ is

$$(3-6) \quad \begin{aligned} \frac{\partial}{\partial t} R_{g(t)} &= \Delta_{g(t)} R_{g(t)} + 2|\operatorname{Ric}_{g(t)}|_{g(t)}^2 + 4|\Delta_{g(t)} u(t)|_{g(t)}^2 \\ &\quad - 4|\nabla_{g(t)}^2 u(t)|_{g(t)}^2 - 8\langle \operatorname{Ric}_{g(t)}, du(t) \otimes du(t) \rangle_{g(t)}. \end{aligned}$$

Also, we have the evolution equation for $|\nabla_{g(t)} u|_{g(t)}^2$,

$$(3-7) \quad \frac{\partial}{\partial t} |\nabla_{g(t)} u(t)|_{g(t)}^2 = \Delta_{g(t)} |\nabla_{g(t)} u(t)|_{g(t)}^2 - 2|\nabla_{g(t)}^2 u(t)|_{g(t)}^2 - 4|\nabla_{g(t)} u(t)|_{g(t)}^4,$$

and the evolution equation for $S_{g(t), u(t)}$,

$$(3-8) \quad \frac{\partial}{\partial t} S_{g(t), u(t)} = \Delta_{g(t)} S_{g(t), u(t)} + 2|\mathcal{S}_{g(t), u(t)}|_{g(t)}^2 + 4|\Delta_{g(t)} u(t)|_{g(t)}^2.$$

4. Entropies for harmonic-Ricci flow

Motivated by Perelman's entropy, List [2006] introduced a similar functional for the harmonic-Ricci flow:

$$\odot_+^2(M) \times C^\infty(M) \times C^\infty(M) \rightarrow \mathbb{R}, \quad (g, u, f) \mapsto \mathcal{F}(g, u, f)$$

where

$$(4-1) \quad \mathcal{F}(g, u, f) := \int_M (R_g + |\nabla_g f|_g^2 - 2|\nabla_g u|_g^2) e^{-f} dV_g.$$

He also showed that if $(g(t), u(t), f(t))$ satisfies the system

$$(4-2) \quad \begin{aligned} \frac{\partial}{\partial t} g(t) &= -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t) - 2 \nabla_{g(t)}^2 f(t), \\ \frac{\partial}{\partial t} u(t) &= \Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}, \\ \frac{\partial}{\partial t} f(t) &= -\Delta_{g(t)} f(t) - R_{g(t)} + 2|\nabla_{g(t)} u(t)|_{g(t)}^2, \end{aligned}$$

the evolution of the entropy is given by

$$(4-3) \quad \begin{aligned} \frac{d}{dt} \mathcal{F}(g(t), u(t), f(t)) \\ = 2 \int_M \left(|\mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t)|_{g(t)}^2 \right. \\ \left. + 2|\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 \right) e^{-f(t)} dV_{g(t)} \\ \geq 0. \end{aligned}$$

Remark 4.1. The system (4-2) is equivalent to

$$(4-4) \quad \begin{aligned} \frac{\partial}{\partial t} g(t) &= -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t), \\ \frac{\partial}{\partial t} u(t) &= \Delta_{g(t)} u(t), \\ \frac{\partial}{\partial t} f(t) &= -\Delta_{g(t)} f(t) - R_{g(t)} + |\nabla_{g(t)} f(t)|_{g(t)}^2 + 2|\nabla_{g(t)} u(t)|_{g(t)}^2. \end{aligned}$$

The same evolution of the entropy holds for system (4-4).

In particular, the entropy is nondecreasing and the equality holds if and only if $(g(t), u(t), f(t))$ satisfies

$$(4-5) \quad \begin{aligned} \mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t) &= 0, \\ \Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)} &= 0. \end{aligned}$$

Definition 4.2. The $\mathcal{E}$ -functional is defined as

$$\odot_+^2(M) \times C^\infty(M) \times C^\infty(M) \rightarrow \mathbb{R}, \quad (g, u, f) \mapsto \mathcal{E}(g, u, f),$$

where

$$(4-6) \quad \mathcal{E}(g, u, f) := \int_M (R_g - 2|\nabla_g u|_g^2) e^{-f} dV_g.$$

Proposition 4.3. *Under the evolution equation (4-4), one has*

$$(4-7) \quad \begin{aligned} & \frac{d}{dt} \mathcal{E}(g(t), u(t), f(t)) \\ &= 2 \int_M |\mathcal{J}_{g(t), u(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 4 \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)}. \end{aligned}$$

Proof. Since $S_{g(t), u(t)} = R_{g(t)} - 2|\nabla_{g(t)} u(t)|_{g(t)}^2$ and

$$\begin{aligned} \frac{\partial}{\partial t} S_{g(t), u(t)} &= \Delta_{g(t)} S_{g(t), u(t)} + 2|\mathcal{J}_{g(t), u(t)}|_{g(t)}^2 + 4|\Delta_{g(t)} u(t)|_{g(t)}^2, \\ \frac{\partial}{\partial t} dV_{g(t)} &= -S_{g(t), u(t)} dV_{g(t)}, \end{aligned}$$

we have

$$\begin{aligned} & \frac{d}{dt} \mathcal{E}(g(t), u(t), f(t)) \\ &= \int_M \left(\frac{\partial}{\partial t} S_{g(t), u(t)} \right) e^{-f(t)} dV_{g(t)} + \int_M S_{g(t), u(t)} \frac{\partial}{\partial t} (e^{-f(t)} dV_{g(t)}) \\ &= \int_M (\Delta_{g(t)} S_{g(t), u(t)} + 2|\mathcal{J}_{g(t), u(t)}|_{g(t)}^2 + 4|\Delta_{g(t)} u(t)|_{g(t)}^2) e^{-f(t)} dV_{g(t)} \\ & \quad + \int_M S_{g(t), u(t)} \left(-\frac{\partial}{\partial t} f(t) - S_{g(t), u(t)} \right) e^{-f(t)} dV_{g(t)} \\ &= 2 \int_M |\mathcal{J}_{g(t), u(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 4 \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ & \quad - \int_M S_{g(t), u(t)} \left(\Delta_{g(t)} f(t) - |\nabla_{g(t)} f(t)|_{g(t)}^2 + \frac{\partial}{\partial t} f(t) + S_{g(t), u(t)} \right) e^{-f(t)} dV_{g(t)}, \end{aligned}$$

which implies (4-7). □

Definition 4.4. For any $k \geq 1$ we define

$$(4-8) \quad \mathcal{F}_k(g, u, f) := \int_M (kR_g + |\nabla_g f|_g^2 - 2k|\nabla_g u|_g^2) e^{-f} dV_g.$$

Using the definition, it is easy to show that

$$(4-9) \quad \mathcal{F}_k(g, u, f) = (k-1)\mathcal{E}(g, u, f) + \mathcal{F}(g, u, f).$$

When $k = 1$, this is the $\mathcal{F}$ -functional.

Theorem 4.5. *Under the evolution equation (4-4), one has*

$$\begin{aligned}
 (4-10) \quad & \frac{d}{dt} \mathcal{F}_k(g(t), u(t), f(t)) \\
 &= 2(k-1) \int_M |\mathcal{S}_{g(t), u(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 2 \int_M |\mathcal{S}_{g(t), u(t)} \\
 &\quad + \nabla_{g(t)}^2 f(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} + 4(k-1) \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\
 &\quad + 4 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)}.
 \end{aligned}$$

Furthermore, the monotonicity is strict unless $g(t)$ is Ricci-flat, $u(t)$ is constant, and $f(t)$ is constant.

Proof. It immediately follows from (4-3) and (4-7). $\square$

Set

$$(4-11) \quad \mu_k(g, u) := \inf \left\{ \mathcal{F}_k(g, u, f) \mid f \in C^\infty(M), \int_M e^{-f} dV_g = 1 \right\}.$$

Then $\mu_k(g, u)$ is the lowest eigenvalue of $-4\Delta_g + k(R_g - 2|\nabla_g u|^2)$.

5. Compact steady harmonic-Ricci breathers

In this section we give an alternative proof on some results on compact steady harmonic-Ricci breathers that were proved in [List 2006; Müller 2012].

Definition 5.1. A solution $(g(t), u(t))$ of the harmonic-Ricci flow (1-1)–(1-2) is called a *harmonic-Ricci breather* if there exist $t_1 < t_2$, a diffeomorphism $\psi : M \rightarrow M$, and a constant $\alpha > 0$ such that

$$g(t_2) = \alpha \psi^* g(t_1), \quad u(t_2) = \psi^* u(t_1).$$

The cases $\alpha < 1$, $\alpha = 1$, and $\alpha > 1$, correspond to *shrinking*, *steady*, and *expanding harmonic-Ricci breathers*.

Theorem 5.2. *If $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M , the lowest eigenvalue $\mu_k(g(t), u(t))$ of the operator $-4\Delta_{g(t)} + k(R_{g(t)} - 2|\nabla_{g(t)} u(t)|_{g(t)}^2)$ is nondecreasing under the harmonic-Ricci flow. The monotonicity is strict unless $g(t)$ is Ricci-flat and $u(t)$ is constant.*

Proof. The proof is similar to that given in [Li 2007]. For any $t_1 < t_2$, suppose that

$$\mu_k(g(t_2), u(t_2)) = \mathcal{F}_k(g(t_2), u(t_2), f_k(t_2))$$

for some smooth function $f_k(x)$. Being an initial value, $f_k(x) = f_k(x, t_2)$ for some smooth function $f_k(x, t)$ satisfying the evolution equation (4-4). The monotonicity

formula (4-10) implies $\mu_k(g(t_2), u(t_2)) \geq \mathcal{F}_k(g(t_1), u(t_1), f_k(t_1)) \geq \mu_k(g(t_1), u(t_1))$. This completes the proof. $\square$

Corollary 5.3. *On a compact Riemannian manifold, the lowest eigenvalues of $-\Delta_{g(t)} + (1/2)(R_{g(t)} - 2|\nabla_{g(t)}u(t)|_{g(t)}^2)$ are nondecreasing under the harmonic-Ricci flow.*

Proof. Since $\mu_2(g(t), u(t))/4$ is the lowest eigenvalue of this operator, the result immediately follows from Theorem 5.2. $\square$

Corollary 5.4. *There is no compact steady harmonic-Ricci breather unless the manifold $(M, g(t))$ is Ricci-flat and u is a constant.*

Proof. If $(g(t), u(t))$ is a steady harmonic-Ricci breather, then, for $t_1 < t_2$ given in the definition, we have

$$\mu_k(g(t_1), u(t_1)) = \mu_k(g(t_2), u(t_2)).$$

Hence, using Theorem 5.2, for any $t \in [t_1, t_2]$, we must have

$$\frac{d}{dt}\mu_k(g(t), u(t)) \equiv 0.$$

Thus $(M, g(t))$ is Ricci-flat and $u(t)$ is constant. $\square$

6. Compact expanding harmonic-Ricci breathers

Inspired by [Li 2007], we define a new functional

$$\odot_+^2(M) \times C^\infty(M) \times C^\infty(\mathbb{R}) \times C^\infty(M) \rightarrow \mathbb{R}, \quad (g, u, \tau, f) \mapsto \mathcal{W}_+(g, u, \tau, f),$$

where $(\tau = \tau(t), t \in \mathbb{R})$.

$$(6-1) \quad \mathcal{W}_+(g, u, \tau, f) := \tau^2 \int_M \left(R_g + \frac{n}{2\tau} + \Delta_g f - 2|\nabla_g u|_g^2 \right) e^{-f} dV_g.$$

Similarly, we define a family of functionals

$$(6-2) \quad \mathcal{W}_{+,k}(g, u, \tau, f) := \tau^2 \int_M \left(k \left(R_g + \frac{n}{2\tau} \right) + \Delta_g f - 2k|\nabla_g u|_g^2 \right) e^{-f} dV_g.$$

It's clear that $\mathcal{W}_{+,1}(g, u, \tau, f) = \mathcal{W}_+(g, u, \tau, f)$.

Lemma 6.1. *One has*

$$\mathcal{W}_+(g, u, \tau, f) = \tau^2 \mathcal{F}(g, u, f) + \frac{n}{2} \tau \int_M e^{-f} dV_g,$$

$$\mathcal{W}_{+,k}(g, u, \tau, f) = \tau^2 \mathcal{F}_k(g, u, f) + \frac{kn}{2} \tau \int_M e^{-f} dV_g,$$

$$\mathcal{W}_{+,k}(g, u, \tau, f) = \mathcal{W}_+(g, u, \tau, f) + (k-1) \left(\tau^2 \mathcal{E}(g, u, f) + \frac{n}{2} \tau \int_M e^{-f} dV_g \right).$$

Proof. Since $\Delta(e^{-f}) = (-\Delta f + |\nabla f|^2)e^{-f}$, it follows that

$$\begin{aligned} \mathcal{W}_+(g, u, \tau, f) - \tau^2 \mathcal{F}(g, u, f) &= \frac{n}{2} \tau \int_M e^{-f} dV_g + \tau^2 \int_M (\Delta_g f - |\nabla_g f|_g^2) e^{-f} dV_g \\ &= \frac{n}{2} \tau \int_M e^{-f} dV_g + \tau^2 \int_M \Delta_g(e^{-f}) dV_g = \frac{n}{2} \tau \int_M e^{-f} dV_g. \end{aligned}$$

We can similarly prove the remaining two relations. $\square$

Theorem 6.2. *Under the coupled system*

$$\begin{aligned} \frac{\partial}{\partial t} g(t) &= -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t) - 2 \nabla_{g(t)}^2 f(t), \\ \frac{\partial}{\partial t} u(t) &= \Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}, \\ \frac{\partial}{\partial t} f(t) &= -\Delta_{g(t)} f(t) - R_{g(t)} + 2 |\nabla_{g(t)} u(t)|_{g(t)}^2, \\ \frac{d}{dt} \tau(t) &= 1, \end{aligned}$$

the first variation formula for $\mathcal{W}_+(g(t), u(t), \tau(t), f(t))$ is

$$\begin{aligned} (6-3) \quad \frac{d}{dt} \mathcal{W}_+(g(t), u(t), \tau(t), f(t)) &= 2\tau(t)^2 \int_M |\mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t) + \frac{1}{2\tau(t)} g(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 4\tau(t)^2 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)}, \end{aligned}$$

and the first variation formula for $\mathcal{W}_{+,k}(g(t), u(t), \tau(t), f(t))$ is

$$\begin{aligned} (6-4) \quad \frac{d}{dt} \mathcal{W}_{+,k}(g(t), u(t), \tau(t), f(t)) &= 2\tau(t)^2 \int_M |\mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t) + \frac{1}{2\tau(t)} g(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 2(k-1)\tau(t)^2 \int_M |\mathcal{S}_{g(t), u(t)} + \frac{1}{2\tau(t)} g(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 4\tau(t)^2 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 4(k-1)\tau(t)^2 \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)}. \end{aligned}$$

Proof. Under this coupled system, we first observe that

$$(6-5) \quad \frac{d}{dt} \left(\int_M e^{-f(t)} dV_{g(t)} \right) = 0.$$

In fact, from $\frac{\partial}{\partial t} dV_{g(t)} = -S_{g(t),u(t)} - \Delta_{g(t)} f(t) dV_{g(t)}$ we obtain

$$\begin{aligned} \frac{d}{dt} \left(\int_M e^{-f(t)} dV_{g(t)} \right) &= \int_M \left(-\frac{\partial}{\partial t} f(t) \cdot dV_{g(t)} + \frac{\partial}{\partial t} dV_{g(t)} \right) e^{-f(t)} \\ &= \int_M [\Delta_{g(t)} f(t) + S_{g(t),u(t)} - S_{g(t),u(t)} - \Delta_{g(t)} f(t)] e^{-f(t)} dV_{g(t)} \\ &= 0. \end{aligned}$$

Lemma 6.1 and the identity (6-5) imply

$$\begin{aligned} \frac{d}{dt} \mathcal{W}_+(g(t), u(t), \tau(t), f(t)) &= \tau(t)^2 \frac{d}{dt} \mathcal{F}(g(t), u(t), f(t)) + 2\tau(t) \mathcal{F}(g(t), u(t), f(t)) + \frac{n}{2} \int_M e^{-f(t)} dV_{g(t)} \\ &= 2\tau(t)^2 \int_M |\mathcal{G}_{g(t),u(t)} + \nabla_{g(t)}^2 f(t)|_{g(t)}^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 4\tau(t)^2 \int_M |\Delta_{g(t)} u(t) - \langle du(t), df(t) \rangle_{g(t)}|^2 e^{-f(t)} dV_{g(t)} \\ &\quad + 2\tau(t) \int_M (S_{g(t),u(t)} + |\nabla_{g(t)} f(t)|_{g(t)}^2) e^{-f(t)} dV_{g(t)} + \frac{n}{2} \int_M e^{-f(t)} dV_{g(t)}, \end{aligned}$$

which is (6-3). Using Lemma 6.1 and the same method, we can prove (6-4). $\square$

Remark 6.3. Under the coupled system

$$\begin{aligned} \frac{\partial}{\partial t} g(t) &= -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t), \\ \frac{\partial}{\partial t} u(t) &= \Delta_{g(t)} u(t), \\ \frac{\partial}{\partial t} f(t) &= -\Delta_{g(t)} f(t) + |\nabla_{g(t)} f(t)|_{g(t)}^2 - R_{g(t)} + 2|\nabla_{g(t)} u(t)|_{g(t)}^2, \\ \frac{d}{dt} \tau(t) &= 1, \end{aligned}$$

the same formulas (6-3) and (6-4) hold for $\mathcal{W}_+$ and $\mathcal{W}_{+,k}$.

Define

$$(6-6) \quad \mu_+(g, u, \tau) := \inf \left\{ \mathcal{W}_+(g, u, \tau, f) \mid f \in C^\infty(M), \int_M e^{-f} dV_g = 1 \right\}.$$

Lemma 6.4. For any $\alpha > 0$, one has

$$(6-7) \quad \mu_+(\alpha g, u, \alpha \tau) = \alpha \mu_+(g, u, \tau).$$

Proof. Set $\bar{g} := \alpha g$; then $R_{\bar{g}} = \alpha^{-1} R_g$, $\Delta_{\bar{g}} f = \alpha^{-1} \Delta_g f$, and $|\nabla_{\bar{g}} u|_{\bar{g}}^2 = \alpha^{-1} |\nabla_g u|_g^2$. Hence

$$\begin{aligned}
\mathcal{W}_+(\bar{g}, u, \alpha\tau, f) &= \alpha^2\tau^2 \int_M \left(R_{\bar{g}} + \frac{n}{2\alpha\tau} + \Delta_{\bar{g}}f - 2|\nabla_{\bar{g}}u|_{\bar{g}}^2 \right) e^{-f} dV_{\bar{g}} \\
&= \alpha\tau^2 \int_M \left(R_g + \frac{n}{2\tau} + \Delta_g f - 2|\nabla_{g(t)}u|_g^2 \right) \alpha^{n/2} e^{-f} dV_g.
\end{aligned}$$

Since $f \mapsto f - (n/2) \ln \alpha$ is one-to-one and onto, by taking the infimum, we derive $\mu_+(\alpha g, u, \alpha\tau) = \alpha \mu_+(g, u, \tau)$. $\square$

Definition 6.5. A solution $(g(t), u(t))$ of the harmonic-Ricci flow is called a *harmonic-Ricci soliton* if there exists a one-parameter family of diffeomorphisms $\psi_t : M \rightarrow M$, satisfying $\psi_0 = \text{id}_M$, and a positive scaling function $\alpha(t)$ such that

$$g(t) = \alpha(t) \psi_t^* g(0), \quad u(t) = \psi_t^* u(0).$$

The cases $(\partial/\partial t)\alpha(t) = \dot{\alpha} < 0$, $\dot{\alpha} = 0$, and $\dot{\alpha} > 0$ correspond to *shrinking*, *steady*, and *expanding harmonic-Ricci solitons*, respectively. If the diffeomorphisms ψ_t are generated by a (possibly time-dependent) vector field $X(t)$ that is the gradient of some function $f(t)$ on M , the soliton is called a *gradient harmonic-Ricci soliton* and f is called the *potential of the harmonic-Ricci soliton*.

Müller [2012] showed that if $(g(t), u(t))$ is a gradient harmonic-Ricci soliton with potential f ,

$$\begin{aligned}
0 &= \text{Ric}_{g(t)} - 2du(t) \otimes du(t) + \nabla_{g(t)}^2 f(t) + cg(t), \\
0 &= \Delta_{g(t)} u(t) - \langle \nabla_{g(t)} u(t), \nabla_{g(t)} f(t) \rangle_{g(t)}
\end{aligned}$$

for some constant c .

Corollary 6.6. *There is no expanding breather on compact Riemannian manifolds other than expanding gradient harmonic-Ricci solitons.*

Proof. The proof is similar to that given in [Li 2007]. Suppose there is an expanding breather on a compact Riemannian manifold M . Then, by definition, we have

$$g(t_2) = \alpha \Phi^* g(t_1), \quad u(t_2) = \Phi^* u(t_1)$$

for some $t_1 < t_2$, where Φ be a diffeomorphism and the constant $\alpha > 1$. Let $f_+(x)$ be a smooth function where $\mathcal{W}_+(g(t_2), u(t_2), \tau(t_2), f(t_2))$ attains its minimum. Then there exists a smooth function $f_+(x, t) : M \times [t_1, t_2] \rightarrow \mathbb{R}$ with initial value $f_+(x, t_2) = f_+(x)$ that satisfies the coupled system in Remark 6.3. Define a linear function

$$\tau : [t_1, t_2] \rightarrow (0, +\infty), \quad \tau(t_2) = T + t_2$$

where T is a constant. By the monotonicity formula, we have

$$\begin{aligned}\mu_+(g(t_2), u(t_2), \tau(t_2)) &= \mathcal{W}_+(g(t_2), u(t_2), \tau(t_2), f_+(t_2)) \\ &\geq \mathcal{W}_+(g(t_1), u(t_1), \tau(t_1), f_+(t_1)) \\ &\geq \mu_+(g(t_1), u(t_1), \tau(t_1)).\end{aligned}$$

Lemma 6.4 and the diffeomorphic invariant property of the functionals shows

$$\mu_+(g(t_1), u(t_1), \tau(t_1)) \leq \alpha \mu_+(g(t_1), u(t_1), \tau(t_1)),$$

which yields

$$\mu_+(g(t_1), u(t_1), \tau(t_1)) \geq 0,$$

since $\alpha > 1$.

If we impose an additional condition $\tau(t_2) = \alpha \tau(t_1)$ and $\tau(t_1) = T + t_1$, we have

$$\tau(t) = \frac{\alpha(t - t_1) - (t - t_2)}{\alpha - 1}, \quad T = \frac{t_2 - \alpha t_1}{\alpha - 1}.$$

Then

$$\frac{\tau(t_2)^{n/2}}{V_{g(t_2)}} = \frac{[\alpha(t_2 - t_1)/(\alpha - 1)]^{n/2}}{\alpha^{n/2} V_{g(t_1)}} = \frac{\tau(t_1)^{n/2}}{V_{g(t_1)}}.$$

The mean value theorem tells us that there exists a time $\bar{t} \in [t_1, t_2]$ with

$$\begin{aligned}0 &= \frac{d}{dt} \Big|_{t=\bar{t}} \log \frac{\tau(t)^{n/2}}{V_{g(t)}} \\ &= \frac{V_{g(\bar{t})}}{\tau(\bar{t})^{n/2}} \cdot \frac{(n/2)\tau(\bar{t})^{n/2-1} V_{g(\bar{t})} - \tau(\bar{t})^{n/2} (d/dt)|_{t=\bar{t}} V_{g(t)}}{V_{g(\bar{t})}^2} \\ &= \frac{n}{2\tau(\bar{t})} - \frac{1}{V_{g(\bar{t})}} \frac{\partial}{\partial t} \Big|_{t=\bar{t}} V_{g(\bar{t})}.\end{aligned}$$

From the evolution equation for the volume element $dV_{g(t)}$, we have

$$\frac{d}{dt} V_{g(t)} = \int_M \frac{\partial}{\partial t} dV_{g(t)} = \int_M (-S_{g(t), u(t)} - \Delta_{g(t)} f(t)) dV_{g(t)} = - \int_M S_{g(t), u(t)} dV_{g(t)}.$$

Putting these together yields

$$0 = \frac{n}{2\tau(\bar{t})} + \frac{1}{V_{g(\bar{t})}} \int_M S_{g(\bar{t}), u(\bar{t})} dV_{g(\bar{t})} = \frac{1}{V_{g(\bar{t})}} \int_M \left(S_{g(\bar{t}), u(\bar{t})} + \frac{n}{2\tau(\bar{t})} \right) dV_{g(\bar{t})}.$$

If we set $\bar{f} = \log V_{g(\bar{t})}$,

$$0 = \mathcal{W}_+(g(\bar{t}), u(\bar{t}), \tau(\bar{t}), \bar{f}) \geq \mu_+(g(\bar{t}), u(\bar{t}), \tau(\bar{t})).$$

By the monotonicity of μ_+ we obtain

$$0 \leq \mu_+(g(t_1), u(t_1), \tau(t_1)) \leq \mu_+(g(\bar{t}), u(\bar{t}), \tau(\bar{t})) \leq 0$$

Hence $\mu_+(g(t_1), u(t_1), \tau(t_1)) = \mu_+(g(t_2), u(t_2), \tau(t_2)) = 0$ and $\mathcal{W}_+ = 0$ on the interval $[t_1, t_2]$. This indicates that the first variation of $\mathcal{W}_+$ must vanish. So the expanding breather is a gradient soliton, that is,

$$\mathcal{J}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t) + \frac{1}{2\tau(t)} g(t) = 0.$$

Moreover, in this case $\Delta_{g(t)} u(t) = \langle du(t), df(t) \rangle_{g(t)}$. □

Because of (6-7), we define

$$(6-8) \quad \mu_{+,k}(g, u, \tau) := \inf \left\{ \mathcal{W}_{+,k}(g, u, \tau, f) \mid f \in C^{+\infty}(M), \int_M e^{-f} dV_g = 1 \right\}.$$

Due to Lemma 6.4, we still have

$$(6-9) \quad \mu_{+,k}(\alpha g, u, \alpha \tau) = \alpha \mu_{+,k}(g, u, \tau).$$

Corollary 6.7. *If $(g(t), u(t))$ is an expanding harmonic-Ricci breather on compact Riemannian manifolds, M is an Einstein manifold and $u(t)$ is constant.*

Proof. Using the same method as in Corollary 6.6 and $\mu_{+,k}$, we can show that the first variation of $\mathcal{W}_{+,k}$ must vanish. Hence, from (6-4), one has

$$\begin{aligned} \mathcal{J}_{g(t), u(t)} + \nabla_{g(t)}^2 f(t) + \frac{1}{2\tau(t)} g(t) &= 0, \\ \mathcal{J}_{g(t), u(t)} + \frac{1}{2\tau(t)} g(t) &= 0, \\ \Delta_{g(t)} u(t) &= \langle du(t), df(t) \rangle_{g(t)}, \\ \Delta_{g(t)} u(t) &= 0. \end{aligned}$$

The above four equations can be reduced to the coupled equation

$$\mathcal{J}_{g(t), u(t)} + \frac{1}{2\tau(t)} g(t) = 0 = \Delta_{g(t)} u(t),$$

which indicates that $u(t)$ is a constant and $\text{Ric}_{g(t)} = -(1/(2\tau(t)))g(t)$. □

7. Eigenvalues of the Laplacian under the harmonic-Ricci flow

In this section we consider the eigenvalues of the Laplacian $\Delta_{g(t)}$ under the harmonic-Ricci flow

$$(7-1) \quad \frac{\partial}{\partial t} g(t) = -2 \operatorname{Ric}_{g(t)} + 4 du(t) \otimes du(t),$$

$$(7-2) \quad \frac{\partial}{\partial t} u(t) = \Delta_{g(t)} u(t).$$

Suppose that $\lambda(t)$, which is a function of time t only, is an eigenvalue of the Laplacian $\Delta_{g(t)}$ with an eigenfunction $f(t) = f(x, t)$, that is,

$$(7-3) \quad -\Delta_{g(t)} f(t) = \lambda(t) f(t).$$

Taking the derivative with respect to t , we get

$$-\left(\frac{\partial}{\partial t} \Delta_{g(t)}\right) f(t) - \Delta_{g(t)} \left(\frac{\partial}{\partial t} f(t)\right) = \left(\frac{d}{dt} \lambda(t)\right) f(t) + \lambda(t) \frac{\partial}{\partial t} f(t).$$

Integrating the above equation with f yields

$$\begin{aligned} -\int_M f(t) \left(\frac{\partial}{\partial t} \Delta_{g(t)}\right) f(t) dV_{g(t)} - \int_M f(t) \Delta_{g(t)} \left(\frac{\partial}{\partial t} f(t)\right) dV_{g(t)} \\ = \frac{d}{dt} \lambda(t) \cdot \int_M f(t)^2 dV_{g(t)} + \lambda(t) \int_M f(t) \frac{\partial}{\partial t} f(t) dV_{g(t)}. \end{aligned}$$

Since

$$\begin{aligned} -\int_M f(t) \Delta \left(\frac{\partial}{\partial t} f(t)\right) dV_{g(t)} &= -\int_M \Delta_{g(t)} f(t) \cdot \frac{\partial}{\partial t} f(t) dV_{g(t)} \\ &= \lambda(t) \int_M f(t) \frac{\partial}{\partial t} f(t) dV_{g(t)}, \end{aligned}$$

it follows that

$$(7-4) \quad \frac{d}{dt} \lambda(t) \cdot \int_M f(t)^2 dV_{g(t)} = -\int_M f(t) \left(\frac{\partial}{\partial t} \Delta_{g(t)}\right) f(t) dV_{g(t)}.$$

If we set $v_{ij} = -2R_{ij} + 4\partial_i u \partial_j u$,

$$\frac{\partial}{\partial t} \Gamma_{ij}^k = \frac{1}{2} g^{k\ell} (\partial_i v_{\ell j} + \partial_j v_{i\ell} - \partial_\ell v_{ij}).$$

We temporarily omit all subscripts t . Multiplying with g^{ij} on both sides, we obtain

$$\begin{aligned} g^{ij} \frac{\partial}{\partial t} \Gamma_{ij}^k &= \frac{1}{2} g^{kl} (2\nabla^i v_{li} - \nabla_l (g^{ij} v_{ij})) = g^{kl} \nabla^i v_{il} + \nabla^k S \\ &= g^{kl} \nabla^i (-2R_{il} + 4\nabla_i u \nabla_l u) + \nabla^k (R - 2|\nabla u|^2) \end{aligned}$$

$$\begin{aligned}
&= -\nabla^k R + 4\Delta u \cdot \nabla^k u + 4\nabla_i u \cdot \nabla^i \nabla^k u + \nabla^k R - 4\nabla^k \nabla^i u \cdot \nabla_i u \\
&= 4\Delta u \cdot \nabla^k u.
\end{aligned}$$

Therefore,

$$\begin{aligned}
\frac{\partial}{\partial t}(\Delta f) &= \frac{\partial}{\partial t}(g^{ij} \nabla_i \nabla_j f) \\
&= \left(\frac{\partial}{\partial t} g^{ij} \right) \nabla_i \nabla_j f + g^{ij} \left[\partial_i \partial_j \frac{\partial f}{\partial t} - \left(\frac{\partial}{\partial t} \Gamma_{ij}^k \right) \partial_k f - \Gamma_{ij}^k \partial_k \frac{\partial f}{\partial t} \right] \\
&= \left(\frac{\partial}{\partial t} g^{ij} \right) \nabla_i \nabla_j f + \Delta_{g(t)} \left(\frac{\partial}{\partial t} f \right) - g^{ij} \left(\frac{\partial}{\partial t} \Gamma_{ij}^k \right) \nabla_k f \\
&= (2R_{ij} - 4\nabla_i u \nabla_j u) \nabla^i \nabla^j f - 4\Delta u \cdot \nabla^k u \nabla_k f + \Delta_{g(t)} \left(\frac{\partial}{\partial t} f \right).
\end{aligned}$$

Plugging this into (7-4), we derive

$$\begin{aligned}
&\frac{d}{dt} \lambda(t) \cdot \int_M f(t)^2 dV_{g(t)} \\
&= -2 \int_M R_{ij} \nabla^i \nabla^j f dV + 4 \int_M f \nabla^i u \nabla^j u \nabla_i \nabla_j f dV + 4 \int_M f \Delta u \cdot \nabla^k u \nabla_k f dV.
\end{aligned}$$

The first term can be rewritten as

$$\begin{aligned}
-2 \int_M f R_{ij} \nabla^i \nabla^j f dV &= \int_M \nabla^i (2f R_{ij}) \nabla^j f dV \\
&= 2 \int_M (\nabla^i f \cdot R_{ij} + f \cdot \nabla^i R_{ij}) \nabla^j f dV \\
&= 2 \int_M R_{ij} \nabla^i f \nabla^j f dV + \int_M f \nabla_j R \nabla^j f dV \\
&= 2 \int_M R_{ij} \nabla^i f \nabla^j f dV - \int_M R \nabla_j (f \nabla^j f) dV \\
&= \lambda \int_R f^2 dV - \int_M R |\nabla f|^2 dV + 2 \int_M R_{ij} \nabla^i f \nabla^j f dV.
\end{aligned}$$

Hence

$$\begin{aligned}
&\left(\frac{d}{dt} \lambda(t) \right) \int_M f(t)^2 dV_{g(t)} \\
&= \lambda(t) \int_M R_{g(t)} f(t)^2 dV_{g(t)} + 2 \int_M R_{ij} \nabla^i f \nabla^j f dV - \int_M R_{g(t)} |\nabla_{g(t)} f(t)|_{g(t)}^2 dV_{g(t)} \\
&\quad + 4 \int_M f (\nabla^i u \nabla^j u \nabla_i \nabla_j f + \Delta u \nabla^k u \nabla_k f) dV.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \int_M f \nabla^i u \nabla^j u \nabla_i \nabla_j f \, dV \\
&= - \int_M \nabla_i (f \nabla^i u \nabla^j u) \nabla_j f \, dV \\
&= - \int_M (\nabla_i f \nabla^i u \nabla^j u + f \Delta u \nabla^j u + f \nabla^i u \nabla_i \nabla^j u) \nabla_j f \, dV \\
&= - \int_M f \Delta u \langle \nabla u, \nabla f \rangle \, dV - \int_M \nabla^i u \nabla^j u \nabla_i f \nabla_j f \, dV - \int_M f \nabla^i u \nabla^j f \nabla_i \nabla_j u \, dV
\end{aligned}$$

and therefore

$$\begin{aligned}
\frac{d}{dt} \lambda(t) \int_M f(t)^2 \, dV_{g(t)} &= \lambda(t) \int_M R_{g(t)} f(t)^2 \, dV_{g(t)} - 4 \int_M f \nabla^i u \nabla^j f \nabla_i \nabla_j u \, dV \\
&\quad + 2 \int_M S_{ij} \nabla^i f \nabla_j f \, dV - \int_M R_{g(t)} |\nabla_{g(t)} f(t)|_{g(t)}^2 \, dV_{g(t)}.
\end{aligned}$$

The last term here can be simplified as follows:

$$\begin{aligned}
& - \int_M f \nabla^i u \nabla^j f \nabla_i \nabla_j u \, dV \\
&= \int_M \nabla^j (f \nabla_i u \nabla_j f) \nabla^i u \, dV \\
&= \int_M (\nabla^j f \nabla_i u \nabla_j f + f \nabla^j \nabla_i u \nabla_j f + f \nabla_i u \Delta f) \nabla^i u \, dV \\
&= \int_M |\nabla u|^2 |\nabla f|^2 \, dV + \int_M f \Delta f |\nabla u|^2 \, dV + \int_M f \nabla^i u \nabla^j f \nabla_i \nabla_j u \, dV.
\end{aligned}$$

Consequently,

$$-2 \int_M f \nabla^i u \nabla^j f \nabla_i \nabla_j u \, dV = \int_M |\nabla u|^2 |\nabla f|^2 \, dV - \lambda \int_M f^2 |\nabla u|^2 \, dV.$$

Therefore we derive the following.

Theorem 7.1. *If $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $\lambda(t)$ denotes the eigenvalue of the Laplacian $\Delta_{g(t)}$, then*

$$\begin{aligned}
(7-5) \quad & \frac{d}{dt} \lambda(t) \cdot \int_M f(t)^2 \, dV_{g(t)} \\
&= \lambda(t) \int_M S_{g(t), u(t)} f(t)^2 \, dV_{g(t)} - \int_M S_{g(t), u(t)} |\nabla_{g(t)} f(t)|_{g(t)}^2 \, dV_{g(t)} \\
&\quad + 2 \int_M \langle \mathcal{G}_{g(t), u(t)}, df(t) \otimes df(t) \rangle \, dV_{g(t)}.
\end{aligned}$$

We set

$$(7-6) \quad S_{\min}(0) := \min_{x \in M} S(x, 0).$$

Theorem 7.2. *Let $(g(t), u(t))_{t \in [0, T]}$ be a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$. Suppose that $\mathcal{S}_{g(t), u(t)} - \alpha S_{g(t), u(t)} g(t) \geq 0$ along the harmonic-Ricci flow for some $\alpha \geq \frac{1}{2}$.*

- (1) *If $S_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.*
- (2) *If $S_{\min}(0) > 0$, the quantity*

$$\left(1 - \frac{2}{n} S_{\min}(0) t\right)^{n\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for $T \leq n/(2S_{\min}(0))$.

- (3) *If $S_{\min}(0) < 0$, the quantity*

$$\left(1 - \frac{2}{n} S_{\min}(0) t\right)^{n\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.

Proof. By Theorem 7.1, we have

$$\frac{d}{dt} \lambda(t) \geq \frac{\int_M S_{g(t), u(t)} f(t)^2 dV_{g(t)}}{\int_M f(t)^2 dV_{g(t)}} \lambda(t) + (2\alpha - 1) \frac{\int_M S_{g(t), u(t)} |\nabla_{g(t)} f(t)|_{g(t)}^2}{\int_M f(t)^2 dV_{g(t)}}.$$

By definition we have $-f(t) \Delta_{g(t)} = \lambda(t) f(t)$. Integrating both sides yields that $\lambda(t) \geq 0$. Since

$$\frac{\partial}{\partial t} S_{g(t), u(t)} = \Delta_{g(t)} S_{g(t), u(t)} + 2|\mathcal{S}_{g(t), u(t)}|_{g(t)}^2 + 4|\Delta_{g(t)} u(t)|_{g(t)}^2$$

and $|\mathcal{S}_{g(t), u(t)}|^2 \geq (1/n) S_{g(t), u(t)}^2$, it follows that

$$\frac{\partial}{\partial t} S_{g(t), u(t)} \geq \Delta_{g(t)} S_{g(t), u(t)} + \frac{2}{n} S_{g(t), u(t)}^2.$$

The corresponding ODE

$$\frac{d}{dt} a(t) = \frac{2}{n} a(t)^2, \quad a(t) = S_{\min}(0)$$

has the solution

$$a(t) = \frac{S_{\min}(0)}{1 - (2/n) S_{\min}(0) t}.$$

Then the maximum principle implies $S_{g(t),u(t)} \geq a(t)$ and hence, using the assumption that $2\alpha - 1 \geq 0$,

$$\frac{d}{dt}\lambda(t) \geq a(t)\lambda(t) + (2\alpha - 1)a(t) \frac{\int_M |\nabla_{g(t)} f(t)|_{g(t)}^2 dV_{g(t)}}{\int_M f(t)^2 dV_{g(t)}}.$$

By integration by parts, we note that

$$\int_M |\nabla f|^2 dV = - \int_M f \cdot \Delta f dV = \lambda \int_M f^2 dV,$$

which shows that

$$\frac{d}{dt}\lambda(t) \geq a(t)\lambda(t) + (2\alpha - 1)a(t)\lambda = 2\alpha a(t)\lambda(t)$$

and

$$\frac{d}{dt} \left(\lambda(t) \cdot \exp \left(-2\alpha \int_0^t a(\tau) d\tau \right) \right) \geq 0.$$

This inequality clearly implies the desired result. If $S_{\min}(0) \geq 0$, by the nonnegativity of $\mathcal{F}_{g(t)}$ preserved along the harmonic-Ricci flow, we conclude that $d\lambda(t)/dt \geq 0$. $\square$

Corollary 7.3. *Let $(g(t), u(t))_{t \in [0, T]}$ be a solution of the harmonic-Ricci flow on a compact Riemannian surface Σ and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$.*

(1) *Suppose that $\text{Ric}_{g(t)} \leq \epsilon du(t) \otimes du(t)$ where*

$$(7-7) \quad \epsilon \leq 4 \frac{1 - \alpha}{1 - 2\alpha}, \quad \alpha > \frac{1}{2}.$$

(i) *If $S_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.*

(ii) *If $S_{\min}(0) > 0$, the quantity*

$$(1 - S_{\min}(0)t)^{2\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for $T \leq 1/S_{\min}(0)$.

(iii) *If $S_{\min}(0) < 0$, the quantity*

$$(1 - S_{\min}(0)t)^{2\alpha} \lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.

(2) *Suppose that*

$$(7-8) \quad |\nabla_{g(t)} u(t)|_{g(t)}^2 g(t) \geq 2du(t) \otimes du(t).$$

(i) *If $S_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.*

(ii) If $S_{\min}(0) > 0$, the quantity

$$(1 - S_{\min}(0)t)\lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for $T \leq 1/S_{\min}(0)$.

(iii) If $S_{\min}(0) < 0$, the quantity

$$(1 - S_{\min}(0)t)\lambda(t)$$

is nondecreasing along the harmonic-Ricci flow for any $t \in [0, T]$.

Proof. As above, we always omit subscripts t . In the surface case, we have $R_{ij} = \frac{1}{2}Rg_{ij}$. Then

$$\begin{aligned} T_{ij} &:= S_{ij} - \alpha Sg_{ij} = \frac{R}{2}g_{ij} - 2\nabla_i u \nabla_j u - \alpha(R - 2|\nabla u|^2)g_{ij} \\ &= \left(\frac{1}{2} - \alpha\right)Rg_{ij} - 2\nabla_i u \nabla_j u + 2\alpha|\nabla u|^2g_{ij}. \end{aligned}$$

For any vector $V = (V^i)$, we calculate

$$\begin{aligned} T_{ij}V^iV^j &= \left(\frac{1}{2} - \alpha\right)R|V|^2 - 2(\nabla_i u V^i)^2 + 2\alpha|\nabla u|^2|V|^2 \\ &\geq \left(\frac{1}{2} - \alpha\right)R|V|^2 - 2|\nabla u|^2|V|^2 + 2\alpha|\nabla u|^2|V|^2. \end{aligned}$$

If $R_{ij} \leq \epsilon \nabla_i u \nabla_j u$, then $T_{ij}V^iV^j = [(\frac{1}{2} - \alpha)\epsilon - 2 + 2\alpha]|\nabla u|^2|V|^2 \geq 0$.

For the second case, we note that

$$\begin{aligned} T_{ij}V^iV^j &= R_{ij}V^iV^j - 2\nabla_i u V^i \nabla_j u V^j - \frac{R}{2}|V|^2 + |\nabla u|^2|V|^2 \\ &\geq R_{ij}V^iV^j - |\nabla u|^2|V|^2 - \frac{R}{2}|V|^2 + |\nabla u|^2|V|^2 = 0. \end{aligned}$$

Hence the corresponding results follow by Theorem 7.2. $\square$

When we consider the Ricci flow, we have the following two results derived from Corollary 7.3.

Corollary 7.4. *Let $(g(t))_{t \in [0, T]}$ be a solution of the Ricci flow on a compact Riemannian surface Σ and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$.*

- (1) *If $R_{\min}(0) \geq 0$, $\lambda(t)$ is nondecreasing along the Ricci flow for any $t \in [0, T]$.*
- (2) *If $R_{\min}(0) > 0$, the quantity $(1 - R_{\min}(0)t)\lambda(t)$ is nondecreasing along the Ricci flow for $T \leq 1/R_{\min}(0)$.*
- (3) *If $R_{\min}(0) < 0$, the quantity $(1 - R_{\min}(0)t)\lambda(t)$ is nondecreasing along the Ricci flow for any $t \in [0, T]$.*

Remark 7.5. Let $(g(t))_{t \in [0, T]}$ be a solution of the Ricci flow on a compact Riemannian surface Σ with nonnegative scalar curvature and let $\lambda(t)$ denote the eigenvalue of the Laplacian $\Delta_{g(t)}$. Then $\lambda(t)$ is nondecreasing along the Ricci flow for $t \in [0, T]$.

8. Eigenvalues of the Laplacian-type under the harmonic-Ricci flow

Recall that

$$(8-1) \quad \mu(g, u) = \mu_1(g, u) = \inf \left\{ \mathcal{F}(g, u, f) \mid \int_M e^{-f} dV_g = 1 \right\}.$$

We showed that $\mu(g, u)$ is the smallest eigenvalue of the operator

$$-4\Delta_g + R_g - 2|\nabla_g u|_g^2.$$

Inspired by [Cao 2007; 2008], we define a Laplacian-type operators associated with quantities g, u, c :

$$(8-2) \quad \Delta_{g, u, c} := -\Delta_g + c(R_g - 2|\nabla_g u|_g^2),$$

$$(8-3) \quad \Delta_{g, u} := \Delta_{g, u, \frac{1}{2}} = -\Delta_g + \frac{1}{2}(R_g - 2|\nabla_g u|_g^2).$$

Then $\mu(g, u)$ is the smallest eigenvalue of the operator $4\Delta_{g, u, 1/4}$.

To the operator $\Delta_{g, u}$ we associate the functional

$$(8-4) \quad C^\infty(M) \rightarrow \mathbb{R}, \quad f \mapsto \lambda_{g, u}(f) := \int_M f \Delta_{g, u} f dV_g.$$

When f is an eigenfunction of the operator $\Delta_{g, u}$ with the eigenvalue λ , that is, $\Delta_{g, u} f = \lambda f$ and is normalized by $\int_M f^2 dV_g = 1$, we obtain $\lambda_{g, u}(f) = \lambda$. The next lemma will deal with the evolution equation for $\lambda(f(t))$, where $f(t)$ is an eigenfunction of $\Delta_{g(t), u(t)}$ and the couple $(g(t), u(t))$ satisfies the harmonic-Ricci flow. Set

$$(8-5) \quad v_{ij} := -2S_{ij} = -2R_{ij} + 4\partial_i u \cdot \partial_j u, \quad v := g^{ij} v_{ij}.$$

The symmetric tensor field thus obtained is denoted by $\mathcal{V}_{g(t), u(t)} = (v_{ij})$.

Lemma 8.1. *Suppose that $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $f(t)$ is an eigenfunction of $\Delta_{g(t), u(t)}$, that is, $\Delta_{g(t), u(t)} f(t) = \lambda(t) f(t)$ (where $\lambda(t)$ is only a function of time t only), with the normalized condition*

$$\int_M f(t)^2 dV_{g(t)} = 1.$$

Then we have

$$\begin{aligned}
(8-6) \quad & \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) \\
&= \int_M f(t) (\nabla^i v_{ik} - \frac{1}{2} \nabla_k v) \nabla^k f(t) dV_{g(t)} - \int_M f^2(t) \frac{\partial}{\partial t} |\nabla_{g(t)} u(t)|_{g(t)}^2 dV_{g(t)} \\
&\quad + \int_M \left(\langle \nabla_{g(t), u(t)}, \nabla_{g(t)}^2 f(t) \rangle_{g(t)} + \frac{1}{2} \left(\frac{\partial}{\partial t} R_{g(t)} \right) f(t) \right) f(t) dV_{g(t)}.
\end{aligned}$$

Before proving the lemma, we recall a formula that is an immediate consequence of the evolution equation:

$$\begin{aligned}
(8-7) \quad & \frac{\partial}{\partial t} (\Delta_{g(t)} f) \\
&= -g^{ip} g^{jq} v_{pq} \nabla_i \nabla_j f - g^{ij} g^{kl} \nabla_i v_{jl} \nabla_k f + \frac{1}{2} \langle \nabla_{g(t)} v_{g(t)}, \nabla_{g(t)} f(t) \rangle_{g(t)}
\end{aligned}$$

where the metric $g(t)$ evolves by $\partial g_{ij} / \partial t = v_{ij}$.

Proof. Using (8-7) and integration by parts, we get

$$\begin{aligned}
& \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) \\
&= \frac{\partial}{\partial t} \int_M \left(-\Delta_{g(t)} f(t) + \left(\frac{R_{g(t)}}{2} - |\nabla_{g(t)} u(t)|_{g(t)}^2 \right) f(t) \right) f(t) dV_{g(t)} \\
&= \int_M \left(g^{ip} g^{jq} v_{pq} \nabla_i \nabla_j f + g^{ij} g^{kl} \nabla_i v_{jl} \nabla_k f - \frac{1}{2} \langle \nabla_{g(t)} v_{g(t)}, \nabla_{g(t)} f(t) \rangle_{g(t)} \right) f(t) dV_{g(t)} \\
&\quad + \int_M \left(-\Delta_{g(t)} \left(\frac{\partial}{\partial t} f(t) \right) + \left(\frac{R_{g(t)}}{2} - |\nabla_{g(t)} u(t)|_{g(t)}^2 \right) \frac{\partial}{\partial t} f(t) \right. \\
&\quad \left. + \left(\frac{\partial}{\partial t} \left(\frac{1}{2} R_{g(t)} \right) - \frac{\partial}{\partial t} (|\nabla_{g(t)} u(t)|_{g(t)}^2) \right) f(t) \right) f(t) dV_{g(t)} \\
&\quad + \int_M \left(-\Delta_{g(t)} f(t) + \left(\frac{R_{g(t)}}{2} - |\nabla_{g(t)} u(t)|_{g(t)}^2 \right) f(t) \right) \frac{\partial}{\partial t} (f(t) dV_{g(t)}) \\
&= \int_M \left(g^{ip} g^{jq} v_{pq} \nabla_i \nabla_j f + \frac{1}{2} \left(\frac{\partial}{\partial t} R_{g(t)} \right) f(t) \right) f(t) dV_{g(t)} \\
&\quad + \int_M (g^{ij} g^{kl} \nabla_i v_{jl} \nabla_k f - \frac{1}{2} g^{kl} \nabla_l v \nabla_k f) f(t) dV_{g(t)} \\
&\quad + \int_M \Delta_{g(t), u(t)} f(t) \left(\frac{\partial}{\partial t} f(t) dV_{g(t)} + \frac{\partial}{\partial t} (f(t) dV_{g(t)}) \right) \\
&\quad \quad \quad - \int_M \frac{\partial}{\partial t} (|\nabla_{g(t)} u(t)|_{g(t)}^2) f(t)^2 dV_{g(t)}.
\end{aligned}$$

Since $f(t)$ is an eigenfunction of $\Delta_{g(t), u(t)}$, it follows that

$$\int_M \Delta_{g(t), u(t)} f(t) \left(\frac{\partial}{\partial t} f(t) dV_{g(t)} + \frac{\partial}{\partial t} (f(t) dV_{g(t)}) \right) = \lambda(t) \frac{\partial}{\partial t} \int_M f(t)^2 dV_{g(t)} = 0$$

by the normalization condition. This completes the proof. $\square$

Using (3-6), we find that the first term in the right side of (8-6) can be written as

$$\begin{aligned}
& \int_M \left(v_{ij} \nabla^i \nabla^j f + \frac{1}{2} \left(\frac{\partial}{\partial t} R_{g(t)} \right) f(t) \right) f(t) dV_{g(t)} \\
&= \int_M \left(-2f(t) \langle \text{Ric}_{g(t)}, \nabla_{g(t)}^2 f(t) \rangle_{g(t)} + 4f(t) \langle du(t) \otimes du(t), \nabla_{g(t)}^2 f(t) \rangle_{g(t)} \right) dV_{g(t)} \\
&\quad + \int_M \left(\left(\frac{1}{2} \Delta_{g(t)} R_{g(t)} + |\text{Ric}_{g(t)}|_{g(t)}^2 \right) f(t)^2 + 2f(t)^2 |\Delta_{g(t)} u(t)|_{g(t)}^2 \right. \\
&\quad \left. - 2f(t)^2 |\nabla_{g(t)}^2 u(t)|_{g(t)}^2 - 4f(t)^2 \langle \text{Ric}_{g(t)}, du(t) \otimes du(t) \rangle_{g(t)} \right) dV_{g(t)} \\
&= \int_M \left(-2f(t) \langle \text{Ric}_{g(t)}, \nabla_{g(t)}^2 f(t) \rangle_{g(t)} + \left(\frac{1}{2} \Delta_{g(t)} R_{g(t)} + |\text{Ric}_{g(t)}|_{g(t)}^2 \right) f(t)^2 \right) dV_{g(t)} \\
&\quad + \int_M \left(4f(t) \langle du \otimes du, \nabla_{g(t)}^2 f(t) \rangle_{g(t)} - 4f^2 \langle du(t) \otimes du(t), \text{Ric}_{g(t)} \rangle_{g(t)} \right. \\
&\quad \left. + 2f(t)^2 |\Delta_{g(t)} u(t)|_{g(t)}^2 - 2f(t)^2 |\nabla_{g(t)}^2 u(t)|_{g(t)}^2 \right) dV_{g(t)}
\end{aligned}$$

For the second term in (8-6), using the contracted Bianchi identities, one has

$$\begin{aligned}
& \int_M \left(g^{ij} \nabla_i v_{jk} - \frac{1}{2} \nabla_k v \right) \nabla^k f \cdot f(t) dV_{g(t)} \\
&= \int_M \left(g^{ij} \nabla_i (-2R_{jk} + 4\partial_j u \partial_k u) \right. \\
&\quad \left. - \frac{1}{2} \nabla_k (-2R_{g(t)} + 4|\nabla_{g(t)} u(t)|_{g(t)}^2) \right) \nabla^k f \cdot f(t) dV_{g(t)} \\
&= \int_M 4f(t) \Delta_{g(t)} u(t) \langle \nabla_{g(t)} u(t), \nabla_{g(t)} f(t) \rangle_{g(t)} dV_{g(t)} \\
&\quad + \int_M (4g^{ij} \nabla_j u \cdot \nabla_i \nabla_k u - 2\nabla_k |\nabla_{g(t)} u(t)|_{g(t)}^2) \nabla^k f \cdot f(t) dV_{g(t)} \\
&= \int_M 4f(t) \Delta_{g(t)} u(t) \langle \nabla_{g(t)} u(t), \nabla_{g(t)} f(t) \rangle_{g(t)} dV_{g(t)}
\end{aligned}$$

where in the last step we use the identity $\nabla_k |\nabla u|^2 = 2g^{pq} \nabla_k \nabla_p u \cdot \nabla_q u$. Therefore

$$\begin{aligned}
(8-8) \quad & \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) \\
&= \int_M \left(-2f(t) \langle \text{Ric}_{g(t)}, \nabla_{g(t)}^2 f(t) \rangle_{g(t)} + \left(\frac{1}{2} \Delta_{g(t)} R_{g(t)} + |\text{Ric}_{g(t)}|_{g(t)}^2 \right) f(t)^2 \right) dV_{g(t)} \\
&\quad + 4f(t) \int_M \left(\langle du(t) \otimes du(t), \nabla_{g(t)}^2 f(t) \rangle_{g(t)} - f(t) \langle du(t) \otimes du(t), \text{Ric}_{g(t)} \rangle_{g(t)} \right. \\
&\quad \left. + 2f(t)^2 |\Delta_{g(t)} u(t)|_{g(t)}^2 - 2f(t)^2 |\nabla_{g(t)}^2 u(t)|_{g(t)}^2 \right. \\
&\quad \left. + 4f(t) \Delta_{g(t)} u(t) \langle \nabla_{g(t)} u(t), \nabla_{g(t)} f(t) \rangle_{g(t)} \right) dV_{g(t)} \\
&\quad - \int_M (\Delta_{g(t)} |\nabla_{g(t)} u(t)|_{g(t)}^2 - 2|\nabla_{g(t)}^2 u(t)|_{g(t)}^2 - 4|\nabla_{g(t)} u(t)|_{g(t)}^4) f(t)^2 dV_{g(t)}.
\end{aligned}$$

The above evolution equation can be simplified as follows.

Theorem 8.2. *Suppose $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $f(t)$ is an eigenfunction of $\Delta_{g(t), u(t)}$, that is, $\Delta_{g(t), u(t)} f(t) = \lambda(t) f(t)$ (where $\lambda(t)$ is only a function of time t only), with the normalized condition $\int_M f(t)^2 dV_{g(t)} = 1$. Then we have*

$$(8-9) \quad \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) = \int_M 2 \langle \mathcal{P}_{g(t)}, df(t) \otimes df(t) \rangle_{g(t)} dV_{g(t)} \\ + \int_M f(t)^2 (|\mathcal{P}_{g(t)}|_{g(t)}^2 + 2|\Delta_{g(t)} u(t)|_{g(t)}^2) dV_{g(t)}.$$

Proof. Calculate

$$\int_M 4f(t) \Delta_{g(t)} u(t) \langle \nabla_{g(t)} u(t), \nabla_{g(t)} f(t) \rangle_{g(t)} dV_{g(t)} \\ = -4 \int_M \nabla_i u [\nabla^i f \cdot \langle \nabla u, \nabla f \rangle + f(\nabla^i \langle \nabla u, \nabla f \rangle)] dV \\ = -4 \int_M |\langle \nabla u, \nabla f \rangle|^2 dV_g - 4 \int_M f \nabla_i u (\langle \nabla^i \nabla u, \nabla f \rangle + \langle \nabla u, \nabla^i \nabla f \rangle) dV.$$

By the same method, we have

$$\int_M -\Delta_{g(t)} |\nabla_{g(t)} u(t)|_{g(t)}^2 f(t)^2 dV_{g(t)} \\ = - \int_M |\nabla u|^2 (2f \Delta f + 2|\nabla f|^2) dV \\ = -2 \int_M |\nabla f|^2 |\nabla u|^2 dV - 2 \int_M f \Delta f |\nabla u|^2 dV.$$

However,

$$\int_M f \Delta f |\nabla u|^2 dV = \int_M -\nabla_i f \cdot \nabla^i (f |\nabla u|^2) dV \\ = - \int_M \nabla_i f (\nabla^i f |\nabla u|^2 + f \nabla^i |\nabla u|^2) dV \\ = - \int_M |\nabla u|^2 |\nabla f|^2 dV - \int_M f \nabla_i f \cdot \nabla^i |\nabla u|^2 dV.$$

Therefore we arrive at

$$\int_M -\Delta_{g(t)} |\nabla_{g(t)} u(t)|_{g(t)}^2 f(t)^2 dV_{g(t)} \\ = 2 \int_M f \nabla_i f \cdot \nabla^i |\nabla u|^2 dV \\ = 4 \int_M f(t) \langle du(t) \otimes df(t), \nabla_{g(t)}^2 u(t) \rangle_{g(t)} dV_{g(t)}.$$

Using the contracted Bianchi identities, we may simplify the term $\int_M \frac{1}{2} f^2 \Delta R dV$ as follows:

$$\begin{aligned}
& \int_N \frac{f(t)^2}{2} \Delta_{g(t)} R_{g(t)} dV_{g(t)} \\
&= -\frac{1}{2} \int_M \nabla_i R \cdot \nabla^i (f^2) dV \\
&= -\int_M \nabla_i R \cdot f \nabla^i f dV = -2 \int_M \nabla^k R_{ki} \cdot f \nabla^i f dV \\
&= 2 \int_M R_{ki} \nabla^k (f \nabla^j f) dV = 2 \int_M R_{ki} (\nabla^k f \cdot \nabla^j f + f \nabla^k \nabla^j f) dV \\
&= 2 \int_M \langle \text{Ric}_{g(t)}, df(t) \otimes df(t) \rangle_{g(t)} dV_{g(t)} + 2 \int_M f(t) \langle \text{Ric}_{g(t)}, \nabla_{g(t)}^2 f(t) \rangle_{g(t)} dV_{g(t)}.
\end{aligned}$$

Hence (8-8) becomes

$$\begin{aligned}
& \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) \\
&= \int_M (2 \langle \text{Ric}_{g(t)}, df(t) \otimes df(t) \rangle_{g(t)} + |\text{Ric}_{g(t)}|_{g(t)}^2 f(t)^2) dV_{g(t)} \\
&\quad + \int_M (2 |\Delta_{g(t)} u(t)|_{g(t)}^2 + 4 |\nabla_{g(t)} u(t)|_{g(t)}^4) f(t)^2 dV_{g(t)} \\
&\quad - \int_M 4 f(t)^2 \langle du(t) \otimes du(t), \text{Ric}_{g(t)} \rangle_{g(t)} dV_{g(t)} \\
&\quad - \int_M 4 |\langle \nabla_{g(t)} u(t), \nabla_{g(t)} f(t) \rangle_{g(t)}|^2 dV_{g(t)} \\
&= \int_M 2 \langle \mathcal{S}_{g(t)}, df(t) \otimes df(t) \rangle_{g(t)} dV_{g(t)} \\
&\quad + \int_M f(t)^2 (|\text{Ric}_{g(t)} - 2 du(t) \otimes du(t)|_{g(t)}^2 + 2 |\Delta_{g(t)} u(t)|_{g(t)}^2) dV_{g(t)}
\end{aligned}$$

where, by definition, $S_{ij} = R_{ij} - 2\partial_i u \partial_j u$. □

List [2006] proved that the nonnegativity of the operator $\mathcal{S}_{g(t)}$ is preserved by the harmonic-Ricci flow. Hence we get the following.

Corollary 8.3. *If $\text{Ric}_{g(0)} - 2du(0) \otimes du(0) \geq 0$, the eigenvalues of the operator $\Delta_{g(t), u(t)}$ are nondecreasing under the harmonic-Ricci flow.*

Remark 8.4. If we choose $u(t) \equiv 0$, we obtain Cao's result [2007].

9. Another formula for $\frac{d}{dt}\lambda(f(t))$

In this section we give another formula for $\frac{d}{dt}\lambda(f(t))$ using a method similar to that in [Li 2007]. Recall the formula

$$\begin{aligned} \frac{d}{dt}\lambda_{g(t),u(t)}(f(t)) &= \int_M 2\langle \mathcal{G}_{g(t),u(t)}, df(t) \otimes df(t) \rangle_{g(t)} dV_{g(t)} \\ &\quad + \int_M f(t)^2 (|\mathcal{G}_{g(t),u(t)}|_{g(t)}^2 + 2|\Delta_{g(t)}u(t)|_{g(t)}^2) dV_{g(t)}. \end{aligned}$$

Consider the function φ determined by $f^2(t) = e^{-\varphi(t)}$. Then we have

$$df = \frac{-e^\varphi d\varphi}{2f}, \quad \frac{\nabla f}{f} = -\frac{\nabla \varphi}{2}, \quad \frac{\Delta f}{f} = -\frac{1}{2}\Delta \varphi + \frac{1}{4}|\nabla \varphi|^2.$$

Hence

$$\begin{aligned} 2\frac{d}{dt}\lambda_{g(t),u(t)}(f(t)) &= \int_M \langle \mathcal{G}_{g(t),u(t)}, d\varphi(t) \otimes d\varphi(t) \rangle_{g(t)} e^{-\varphi(t)} dV_{g(t)} \\ &\quad + 2 \int_M (|\mathcal{G}_{g(t),u(t)}|_{g(t)}^2 + 2|\Delta_{g(t)}u(t)|_{g(t)}^2) e^{-\varphi} dV_{g(t)}. \end{aligned}$$

Using integration by parts and contracted Bianchi identities yields

$$\begin{aligned} &\int_M \langle \mathcal{G}_{g(t),u(t)}, d\varphi(t) \otimes d\varphi(t) \rangle_{g(t)} e^{-\varphi(t)} dV_{g(t)} \\ &= \int_M S_{ij} \nabla^i \varphi \nabla^j \varphi e^{-\varphi} dV = - \int_M S_{ij} \nabla^j \varphi \nabla^i (e^{-\varphi}) dV \\ &= \int_M e^{-\varphi} \nabla^i (S_{ij} \nabla^j \varphi) dV \\ &= \int_M \nabla^i S_{ij} \cdot \nabla^j \varphi \cdot e^{-\varphi} dV + \int_M S_{ij} \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV \\ &= \int_M \nabla^i R_{ij} \cdot \nabla^j \varphi \cdot e^{-\varphi} dV_g + \int_M S_{ij} \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV \\ &\quad + \int_M \nabla^i (-2\nabla_i u \nabla_j u) \nabla^j \varphi \cdot e^{-\varphi} dV_g \\ &= \frac{1}{2} \int_M R \Delta(e^{-\varphi}) dV + \int_M S_{ij} \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV - 2 \int_M (\nabla^i u \nabla_j u) \nabla^i \nabla^j (e^{-\varphi}) dV. \end{aligned}$$

Thus

$$\begin{aligned} &\int_M S_{ij} \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV \\ &= \int_M S_{ij} \nabla^i \varphi \nabla^j \varphi e^{-\varphi} dV - \frac{1}{2} \int_M R \Delta(e^{-\varphi}) dV + 2 \int_M (\nabla^i u \nabla_j u) \nabla^i \nabla^j (e^{-\varphi}). \end{aligned}$$

On the other hand, one gets

$$\begin{aligned}
 \int_M |\nabla_{g(t)}^2 \varphi(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} &= \int_M \nabla_i \nabla_j \varphi \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV \\
 &= - \int_M \nabla_j \varphi \cdot \nabla_i \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV - \int_M \nabla_j \varphi \cdot \nabla^i \nabla^j \varphi \cdot \nabla_i (e^{-\varphi}) dV \\
 &= - \int_M \nabla_j \varphi \cdot \nabla_i \nabla^j \nabla^i \varphi \cdot e^{-\varphi} dV - \int_M \nabla_j \varphi \cdot \nabla^i \nabla^j \varphi \cdot \nabla_i (e^{-\varphi}) dV.
 \end{aligned}$$

Since

$$\begin{aligned}
 \int_M \nabla_j \varphi \cdot \nabla^i \nabla^j \varphi \cdot \nabla_i (e^{-\varphi}) dV &= - \int_M \nabla^i (\nabla_j \varphi \cdot \nabla_i (e^{-\varphi})) \nabla^j \varphi dV \\
 &= - \int_M \nabla^j \varphi \cdot \nabla^i \nabla_j \varphi \cdot \nabla_i (e^{-\varphi}) dV - \int_M |\nabla \varphi|^2 \Delta (e^{-\varphi}) dV,
 \end{aligned}$$

which implies

$$\int_M \nabla_j \varphi \cdot \nabla^i \nabla^j \varphi \cdot \nabla_i (e^{-\varphi}) dV = -\frac{1}{2} \int_M |\nabla \varphi|^2 \Delta (e^{-\varphi}) dV,$$

it follows that

$$\int_M |\nabla^2 \varphi|^2 e^{-\varphi} dV = - \int_M \nabla_j \varphi \cdot \nabla_i \nabla^j \nabla^i \varphi \cdot e^{-\varphi} dV + \frac{1}{2} \int_M |\nabla \varphi|^2 \Delta (e^{-\varphi}) dV.$$

By the Ricci identity the term $\nabla^i \nabla^j \nabla^i \varphi$ equals

$$\begin{aligned}
 \nabla_i \nabla^j \nabla^i \varphi &= g^{jk} g^{il} \nabla_i \nabla_k \nabla_l \varphi = g^{jk} g^{il} (\nabla_k \nabla_i \nabla_l \varphi - R_{ikl}^p \nabla_p \varphi) \\
 &= \nabla^j \nabla_i \nabla^i \varphi - g^{jk} g^{il} R_{iklp} \nabla^p \varphi \\
 &= \nabla^j \Delta \varphi + g^{jk} g^{il} R_{ikpl} \nabla^p \varphi = \nabla^j \Delta \varphi + g^{jk} R_{kp} \nabla^p \varphi.
 \end{aligned}$$

Hence

$$\begin{aligned}
 &- \int_M \nabla_j \varphi \cdot \nabla_i \nabla^j \nabla^i \varphi \cdot e^{-\varphi} dV \\
 &= - \int_M \nabla_i \varphi \cdot \nabla^j \Delta \varphi \cdot e^{-\varphi} dV - \int_M R_{kp} \nabla^k \varphi \cdot \nabla^p \varphi e^{-\varphi} dV \\
 &= \int_M \nabla^j \Delta \varphi \cdot \nabla_j (e^{-\varphi}) + \int_M R_{kp} \nabla^k \varphi \cdot \nabla^p (e^{-\varphi}) dV \\
 &= - \int_M \Delta \varphi \cdot \Delta (e^{-\varphi}) - \int_M e^{-\varphi} (\nabla^p R_{kp} \cdot \nabla^k \varphi + R_{kp} \nabla^p \nabla^k \varphi) \\
 &= - \int_M \Delta (e^{-\varphi}) \cdot \Delta \varphi dV + \frac{1}{2} \int_M \nabla_k R \cdot \nabla^k (e^{-\varphi}) dV - \int_M e^{-\varphi} R_{kp} \nabla^k \nabla^p \varphi dV \\
 &= - \int_M \Delta (e^{-\varphi}) (\Delta \varphi + \frac{1}{2} R) - \int_M R_{kp} \nabla^k \nabla^p \varphi \cdot e^{-\varphi} dV.
 \end{aligned}$$

Putting those formulas together, we obtain

$$\begin{aligned}
& \int_M 2S_{ij} \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV + \int_M |\nabla^2 \varphi|^2 e^{-\varphi} dV \\
&= \int_M S_{ij} \nabla^i \nabla_j \varphi \cdot e^{-\varphi} dV + \int_M (-2\nabla_i u \nabla_j u) \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV \\
&\quad - \int_M \Delta(e^{-\varphi}) \left(\Delta \varphi + \frac{R}{2} - \frac{1}{2} |\nabla \varphi|^2 \right) dV \\
&= \int_M S_{ij} \nabla^i \varphi \nabla^j \varphi \cdot e^{-\varphi} dV - \int_M \Delta(e^{-\varphi}) (\Delta \varphi + R - \frac{1}{2} |\nabla \varphi|^2) dV \\
&\quad + 2 \int_M (\nabla_i u \nabla_j u \cdot \nabla^i \nabla^j (e^{-\varphi}) - \nabla_i u \nabla_j u \cdot \nabla^i \nabla^j \varphi \cdot e^{-\varphi}) dV.
\end{aligned}$$

Since f is an eigenfunction of λ , it induces

$$\lambda = -\frac{\Delta f}{f} + \frac{R}{2} - |\nabla u|^2 = \frac{1}{2} \Delta \varphi - \frac{1}{4} |\nabla \varphi|^2 + \frac{R}{2} - |\nabla u|^2,$$

and therefore

$$\begin{aligned}
& \int_M 2S_{ij} \nabla^i \nabla^j \varphi \cdot e^{-\varphi} dV + \int_M |\nabla^2 \varphi|^2 e^{-\varphi} dV \\
&= \int_M S_{ij} \nabla^i \varphi \nabla^j \varphi \cdot e^{-\varphi} dV - 2 \int_M \Delta(|\nabla u|^2) \cdot e^{-\varphi} dV \\
&\quad + 2 \int_M \nabla_i u \nabla_j u (\nabla^i \nabla^j (e^{-\varphi}) - \nabla^i \nabla^j \varphi \cdot e^{-\varphi}) dV.
\end{aligned}$$

Plugging this into the expression of $\frac{d}{dt} \lambda(f(t))$ yields

$$\begin{aligned}
& 2 \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) \\
&= \int_M S_{ij} \nabla^i \varphi \nabla^j \varphi \cdot e^{-\varphi} dV + \int_M |\mathcal{P}|^2 e^{-\varphi} dV + \int_M |\mathcal{S}|^2 e^{-\varphi} dV + 4 \int_M |\Delta u|^2 e^{-\varphi} dV \\
&= \int_M |\mathcal{P}_{g(t), u(t)} + \nabla_{g(t)}^2 \varphi(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} + \int_M |\mathcal{P}_{g(t), u(t)}|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} \\
&\quad + 4 \int_M |\Delta_{g(t)} u(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} + 2 \int_M \Delta_{g(t)} |\nabla_{g(t)} u(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} \\
&\quad + 2 \int_M \nabla_i u \nabla_j u (-\nabla^i \nabla^j (e^{-\varphi}) + \nabla^i \nabla^j \varphi \cdot e^{-\varphi}) dV
\end{aligned}$$

Now define

$$\begin{aligned}
I &:= \int_M (\nabla_i u \nabla_j u \cdot \nabla^i \nabla^j \varphi) e^{-\varphi} dV = - \int_M \nabla^i (\nabla_i u \nabla_j u \cdot e^{-\varphi}) \nabla^j \varphi dV \\
&= - \int_M \nabla^j \varphi (\Delta u \cdot \nabla_j u \cdot e^{-\varphi} + \nabla_i u \nabla^i \nabla_j u \cdot e^{-\varphi} - \nabla_i u \nabla_j u \nabla^i \varphi \cdot e^{-\varphi}) dV \\
&= - \int_M \nabla_j u \nabla^j \varphi \Delta u \cdot e^{-\varphi} dV - \int_M \nabla_i u \nabla^j \varphi \nabla^i \nabla_j u \cdot e^{-\varphi} dV + \int_M |\langle du, d\varphi \rangle|^2 e^{-\varphi} dV,
\end{aligned}$$

$$\begin{aligned}
II &:= \int_M \nabla_i u \nabla_j u \nabla^i \nabla^j (e^{-\varphi}) dV = \int_M \nabla^i \nabla^j (\nabla_i u \nabla_j u) e^{-\varphi} dV \\
&= \int_M \nabla^i (\nabla^j \nabla_i u \cdot \nabla_j u + \nabla_i u \Delta u) e^{-\varphi} dV \\
&= \int_M (\Delta \nabla^i u \cdot \nabla_i u + \nabla^i \Delta u \cdot \nabla_i u + |\nabla^2 u|^2 + |\Delta u|^2) e^{-\varphi} dV, \\
III &:= \int_M \Delta (|\nabla u|^2) e^{-\varphi} dV = 2 \int_M \nabla^i (\nabla_i \nabla_j u \cdot \nabla^j u) e^{-\varphi} dV \\
&= 2 \int_M (\Delta \nabla_j u \cdot \nabla^j u + |\nabla^2 u|^2) e^{-\varphi} dV.
\end{aligned}$$

If we set

$$B := 2(III + I - II),$$

then

$$\begin{aligned}
\frac{B}{2} &= \int_M (\Delta \nabla_i u \cdot \nabla^i u - \nabla_i \Delta u \cdot \nabla^i u + |\nabla^2 u|^2 - |\Delta u|^2 + |\langle du, d\varphi \rangle|^2 \\
&\quad - \nabla_i u \cdot \nabla^i \varphi \cdot \Delta u - \nabla_i u \cdot \nabla^j \varphi \cdot \nabla^i \nabla_j u) e^{-\varphi} dV \\
&= \int_M (R_{ij} \nabla^i u \nabla^j u + |\nabla^2 u|^2 - |\Delta u|^2 + |\langle du, d\varphi \rangle|^2 \\
&\quad - \nabla_i u \cdot \nabla^i \varphi \cdot \Delta u - \nabla_i u \cdot \nabla^j \varphi \cdot \nabla^i \nabla_j u) e^{-\varphi} dV.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
- \int_M \nabla_i u \cdot \nabla^i \varphi \cdot \Delta u \cdot e^{-\varphi} dV &= \int_M (\nabla_i u \cdot \Delta u) \nabla^i (e^{-\varphi}) dV \\
&= - \int_M \nabla^i (\nabla_i u \cdot \Delta u) e^{-\varphi} dV \\
&= \int_M (-|\Delta u|^2 - \nabla_i u \cdot \nabla^i \Delta u) e^{-\varphi} dV
\end{aligned}$$

and

$$\begin{aligned}
- \int_M \nabla_i u \nabla^j \varphi \nabla^i \nabla_j u \cdot e^{-\varphi} dV &= \int_M \nabla_i u \nabla^i \nabla_j u \nabla^j (e^{-\varphi}) dV \\
&= - \int_M \nabla^j (\nabla_i u \nabla^i \nabla_j u) e^{-\varphi} dV \\
&= \int_M (-|\nabla^2 u|^2 - \nabla_i u \Delta \nabla^i u) e^{-\varphi} dV.
\end{aligned}$$

Therefore

$$(9-1) \quad \frac{B}{2} = \int_M (-2|\Delta u|^2 + |\langle du, d\varphi \rangle|^2 - 2\langle \nabla u, \nabla \Delta u \rangle) e^{-\varphi} dV.$$

By definition,

$$\Delta(|\nabla u|^2) = \Delta(\nabla^i u \cdot \nabla_i u) = 2\nabla^i u \cdot \Delta \nabla_i u + 2|\nabla^2 u|^2.$$

So

$$\begin{aligned} \Delta|\nabla u|^2 &= 2|\nabla^2 u|^2 + 2(\nabla_i \Delta u + R_{ij} \nabla^j u) \nabla^i u \\ &= 2|\nabla^2 u|^2 + 2R_{ij} \nabla^i u \cdot \nabla^j u + 2\langle \nabla u, \nabla \Delta u \rangle. \end{aligned}$$

Plugging this into (9-1) yields

$$\frac{B}{2} = \int_M (-2|\Delta u|^2 + |\langle du, d\varphi \rangle|^2 + 2|\nabla^2 u|^2 - \Delta|\nabla u|^2 + 2R_{ij} \nabla^i u \nabla^j u) e^{-\varphi} dV.$$

Since

$$\begin{aligned} 2R_{ij} \nabla^i u \nabla^j u &= 2(S_{ij} + 2\nabla_i u \nabla_j u) \nabla^i u \nabla^j u \\ &= 2S_{ij} \nabla^i u \nabla^j u + 4|\nabla u|^4 \\ &= \frac{1}{4}|\mathcal{S} + 4du \otimes du|^2 - \frac{1}{4}|\mathcal{S}|^2, \end{aligned}$$

it follows that

$$\begin{aligned} \frac{B}{2} &= III + I - II \\ &= \int_M (|\langle du, d\varphi \rangle|^2 - 2|\Delta u|^2 - \frac{1}{4}|\mathcal{S}|^2 + 2|\nabla^2 u|^2 + \frac{1}{4}|\mathcal{S} + 4du \otimes du|^2) e^{-\varphi} dV - III. \end{aligned}$$

Hence

$$B = \int_M (-4|\Delta u|^2 + 2|\langle du, d\varphi \rangle|^2 - \frac{1}{2}|\mathcal{S}|^2 + 4|\nabla^2 u|^2 + \frac{1}{2}|\mathcal{S} + 4du \otimes du|^2) e^{-\varphi} dV - 2III.$$

Theorem 9.1. *Suppose that $(g(t), u(t))$ is a solution of the harmonic-Ricci flow on a compact Riemannian manifold M and $f(t)$ is an eigenfunction of $\Delta_{g(t), u(t)}$, that is, $\Delta_{g(t), u(t)} f(t) = \lambda(t) f(t)$ (where $\lambda(t)$ is only a function of time t), with the normalized condition $\int_M f(t)^2 dV_{g(t)} = 1$. Then we have*

$$\begin{aligned} \frac{d}{dt} \lambda(t) &= \frac{d}{dt} \lambda_{g(t), u(t)}(f(t)) \\ &= \frac{1}{2} \int_M |\mathcal{S}_{g(t), u(t)} + \nabla_{g(t)}^2 \varphi(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} + \frac{1}{4} \int_M |\mathcal{S}_{g(t), u(t)}|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} \\ &\quad + \int_M |\langle du(t), d\varphi(t) \rangle_{g(t)}|^2 e^{-\varphi(t)} dV_{g(t)} + 2 \int_M |\nabla_{g(t)}^2 u(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} \\ &\quad + \frac{1}{4} \int_M |\mathcal{S}_{g(t), u(t)} + 4du(t) \otimes du(t)|_{g(t)}^2 e^{-\varphi(t)} dV_{g(t)} \\ &\quad - \int_M \Delta_{g(t)} (|\nabla_{g(t)} u(t)|_{g(t)}^2) e^{-\varphi(t)} dV_{g(t)}. \end{aligned}$$

Remark 9.2. When $u \equiv 0$, this equation reduces to Li's formula [2007].

10. The first variation of expander and shrinker entropies

Suppose that M is a closed manifold of dimension n . We define

$$\mathcal{W}_{\pm} : \odot_+^2(M) \times C^\infty(M) \times C^\infty(M) \times \mathbb{R}^+ \rightarrow \mathbb{R}, \quad (g, u, f, \tau) \mapsto \mathcal{W}_{\pm}(g, u, f, \tau)$$

where

$$(10-1) \quad \mathcal{W}_{\pm}(g, u, f, \tau) := \int_M (\tau(S_{g,u} + |\nabla_g f|_g^2) \mp f \pm n) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g.$$

Set

$$\mu_{\pm}(g, u, \tau) := \inf \left\{ \mathcal{W}_{\pm}(g, u, f, \tau) \mid f \in C^\infty(M), \int_M \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g = 1 \right\},$$

$$v_{\pm}(g, u) := \sup \{ \mu_{\pm}(g, u, \tau) \mid \tau > 0 \}.$$

Lemma 10.1. Suppose $v_{\pm}(g, u) = \mathcal{W}_{\pm}(g, u, f_{\pm}, \tau_{\pm})$ for some functions $f_{\pm}$ and constants $\tau_{\pm}$ satisfying

$$\int_M \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV_g = 1, \quad \tau_{\pm} > 0.$$

Then we must have

$$\tau_{\pm}(-2\Delta_g f_{\pm} + |\nabla_g f_{\pm}|_g^2 - S_{g,u}) \pm f_{\pm} \mp n + v_{\pm}(g, u) = 0,$$

$$\int_M \frac{f_{\pm} e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV_g = \frac{n}{2} \mp v_{\pm}(g, u).$$

Proof. Since g and u are fixed, we consider the corresponding Lagrangian multiplier function

$$\mathfrak{L}_{\pm}(f, \tau; \lambda) := \mathcal{W}_{\pm}(g, u, f, \tau) - \lambda \left(\int_M \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g - 1 \right).$$

Then the variation of $\mathfrak{L}_{\pm}$ in f direction is

$$\delta_f \mathfrak{L}_{\pm}(f, \tau; \lambda) = \int_M (2\tau \nabla^i f \nabla_i (\delta f) \mp \delta f + \lambda \delta f) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g$$

$$- \int_M (\tau(S_{g,u} + |\nabla_g f|_g^2) \mp f \pm n) \delta f \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g.$$

By the divergence theorem, we calculate

$$\int_M \nabla^i f \cdot \nabla_i (\delta f) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g = - \int_M \nabla_i (\nabla^i f \frac{e^{-f}}{(4\pi\tau)^{n/2}}) \delta f dV_g$$

$$= - \int_M (\Delta_g f - |\nabla_g f|_g^2) \delta f \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g.$$

Hence

$$\delta_f \mathfrak{L}_\pm(f, \tau; \lambda) = \int_M (\tau(-2\Delta_g f + |\nabla_g f|_g^2 - S_{g,u}) \pm f \mp n \mp 1 + \lambda) \delta f \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV.$$

This implies that

$$\tau_\pm(-2\Delta_g f_\pm + |\nabla_g f_\pm|_g^2 - S_{g,u}) \pm f_\pm \mp n \mp 1 + \lambda_\pm = 0.$$

Since $f_\pm$ satisfies the normalized condition, it follows that

$$0 = \lambda_\pm \mp 1 + \int_M (\tau_\pm(-2\Delta_g f_\pm + |\nabla_g f_\pm|_g^2 - S_{g,u}) \pm f_\pm \mp n) \frac{e^{-f_\pm}}{(4\pi\tau_\pm)^{n/2}} dV_g.$$

From the identity

$$\int_M \Delta_g f \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g = \int_M |\nabla_g f|_g^2 \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g$$

and the definition (10-1), we obtain

$$v_\pm(g, u) = \mathcal{W}_\pm(g, u, f_\pm, \tau_\pm) = \lambda_\pm \mp 1,$$

and, consequently,

$$\tau_\pm(-2\Delta_g f_\pm + |\nabla_g f_\pm|_g^2 - S_{g,u}) \pm f_\pm \mp n + v_\pm(g, u) = 0.$$

The variation of $\mathfrak{L}_\pm$ with respect to τ indicates that

$$\begin{aligned} \delta_\tau \mathfrak{L}_\pm(f, \tau; \lambda) &= \int_M \delta\tau (S_{g,u} + |\nabla_g f|_g^2) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g - \lambda \int_M \left(-\frac{n}{2} \frac{\delta\tau}{\tau}\right) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g \\ &\quad + \int_M \left(-\frac{n}{2} \frac{\delta\tau}{\tau}\right) (\tau(S_{g,u} + |\nabla_g f|_g^2) \mp f \pm n) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g \\ &= \int_M \delta\tau \left(\left(1 - \frac{n}{2}\right) (S_{g,u} + |\nabla_g f|_g^2) + \frac{n}{2\tau} (\lambda \pm f \mp n) \right) \frac{e^{-f}}{(4\pi\tau)^{n/2}} dV_g. \end{aligned}$$

Using the first proved equation, we have

$$\begin{aligned} 0 &= \int_M ((v_\pm(g, u) \pm f_\pm \mp n) \left(1 - \frac{n}{2}\right) + \frac{n}{2} (v_\pm(g, u) \pm f_\pm \mp n \pm 1)) \frac{e^{-f_\pm}}{(4\pi\tau_\pm)^{n/2}} dV_g \\ &= \int_M \left(v_\pm \pm f_\pm \mp \frac{n}{2}\right) \frac{e^{-f_\pm}}{(4\pi\tau_\pm)^{n/2}} dV_g \end{aligned}$$

and therefore we obtain the second one. $\square$

For a symmetric 2-tensor $h = (h_{ij}) \in \odot^2(M)$, we set

$$g(s) := g + sh$$

Then the variation of $g(s)$ is

$$(10-2) \quad \frac{\partial}{\partial s} \Big|_{s=0} R_{g(s)} = -h^{ij} R_{ij} + \nabla^i \nabla^j h_{ij} - \Delta_g(\text{tr}_g h).$$

Theorem 10.2. *Suppose that (M, g) is a compact Riemannian manifold and u a smooth function on M . Let h be any symmetric covariant 2-tensor on M and set $g(s) := g + sh$. Let v be any smooth function on M and $u(s) := u + sv$. If*

$$v_{\pm}(g(s), u(s)) = {}^{\circ}W_{\pm}(g(s), u(s), f_{\pm}(s), \tau_{\pm}(s))$$

for some smooth functions $f_{\pm}(s)$ with

$$\int_M e^{-f_{\pm}(s)} dV / (4\pi \tau_{\pm}(s))^{n/2} = 1$$

and constants $\tau_{\pm}(s) > 0$,

$$\begin{aligned} \frac{d}{ds} \Big|_{s=0} v_{\pm}(g(s), u(s)) = & -\tau_{\pm} \int_M \left(\langle h, \mathcal{S}_{g,u} \rangle_g + \langle h, \nabla_g^2 f \rangle_g \pm \frac{1}{2\tau_{\pm}} \text{tr}_g h \right) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g \\ & + 4\tau_{\pm} \int_M v(\Delta_g u - \langle du, df_{\pm} \rangle_g) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g, \end{aligned}$$

where $f_{\pm} := f_{\pm}(0)$ and $\tau_{\pm} := \tau_{\pm}(0)$. In particular, the critical points of $v_{\pm}(\cdot, \cdot)$ satisfy

$$\mathcal{S}_{g,u} + \nabla_g^2 f \pm \frac{1}{2\tau_{\pm}} g = 0, \quad \Delta_g u = \langle du, df_{\pm} \rangle_g.$$

Consequently, if ${}^{\circ}W_{\pm}(g, u, f, \tau)$ and $v_{\pm}(g, u)$ achieve their minimums, (M, g) is a gradient expanding and shrinker harmonic-Ricci soliton according to the sign.

Proof. By definition, one has

$$\begin{aligned} \frac{d}{ds} v_{\pm}(g(s), u(s)) &= \frac{d}{ds} {}^{\circ}W_{\pm}(g(s), u(s), f_{\pm}(s), \tau_{\pm}(s)) \\ &= \int_M \left(\frac{\partial}{\partial s} \tau_{\pm}(s) (S_{g(s), u(s)} + |\nabla_{g(s)} f_{\pm}(s)|_{g(s)}^2) \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} \\ &\quad + \int_M \left(\tau_{\pm}(s) \frac{\partial}{\partial s} (S_{g(s), u(s)} + |\nabla_{g(s)} f_{\pm}(s)|_{g(s)}^2) \mp \frac{\partial}{\partial s} f_{\pm}(s) \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} \\ &\quad + \int_M (\tau_{\pm}(s) (S_{g(s), u(s)} + |\nabla_{g(s)} f_{\pm}(s)|_{g(s)}^2) \mp f_{\pm}(s) \pm n) \\ &\quad \cdot \frac{\partial}{\partial s} \left(\frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} \right). \end{aligned}$$

Since

$$\begin{aligned}
\frac{\partial}{\partial s} S_{g(s), u(s)} &= \frac{\partial}{\partial s} R_{g(s)} - 2 \frac{\partial}{\partial s} |\nabla_{g(s)} u(s)|_{g(s)}^2 \\
&= \frac{\partial}{\partial s} R_{g(s)} - 2 \left(\frac{\partial}{\partial s} g^{ij} \right) \nabla_i u \nabla_j u - 4 g^{ij} \frac{\partial}{\partial s} \nabla_i u \cdot \nabla_j u \\
&= \frac{\partial}{\partial s} R_{g(s)} - 2 (-g^{ip} g^{jq} h_{pq}) \nabla_i u \nabla_j u - 4 g^{ij} \nabla_i \left(\frac{\partial}{\partial s} u \right) \nabla_j u \\
&= \frac{\partial}{\partial s} R_{g(s)} + 2 h_{pq} \nabla^p u \nabla^q u - 4 \nabla_i \left(\frac{\partial}{\partial s} u \right) \nabla^i u
\end{aligned}$$

and

$$\begin{aligned}
&\frac{\partial}{\partial s} \left(\frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} \right) \\
&= \left(-\frac{\partial}{\partial s} f_{\pm}(s) - \frac{n}{2\tau_{\pm}(s)} \frac{\partial}{\partial s} \tau_{\pm}(s) \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} + \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} \frac{\partial}{\partial s} dV_{g(s)} \\
&= \left(-\frac{\partial}{\partial s} f_{\pm}(s) - \frac{n}{2\tau_{\pm}(s)} \frac{\partial}{\partial s} \tau_{\pm}(s) + \frac{1}{2} \operatorname{tr}_g h \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)},
\end{aligned}$$

it follows that

$$\begin{aligned}
&\frac{d}{ds} v_{\pm}(g(s), u(s)) \\
&= \int_M \frac{\partial}{\partial s} \tau_{\pm}(s) (S_{g(s), u(s)} + |\nabla_{g(s)} f_{\pm}(s)|_{g(s)}^2) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} \\
&\quad + \int_M \left(\tau_{\pm}(s) \left(\frac{\partial}{\partial s} R_{g(s)} + 2 h_{pq} \nabla^p u \nabla^q u - 4 \nabla_i \left(\frac{\partial}{\partial s} u \right) \nabla^i u \right. \right. \\
&\quad \left. \left. - h_{pq} \nabla^p f \nabla^q f + 2 \nabla_i \left(\frac{\partial}{\partial s} f \right) \nabla^i f \right) \mp \frac{\partial}{\partial s} f_{\pm}(s) \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)} \\
&\quad + \int_M \left(-\frac{\partial}{\partial s} f_{\pm}(s) - \frac{n}{2\tau_{\pm}(s)} \frac{\partial}{\partial s} \tau_{\pm}(s) + \frac{1}{2} \operatorname{tr}_g h \right) \\
&\quad \cdot \left(\tau_{\pm}(s) (S_{g(s), u(s)} + |\nabla_{g(s)} f_{\pm}(s)|_{g(s)}^2) \mp f_{\pm}(s) \pm n \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_{g(s)}.
\end{aligned}$$

From the equalities

$$\begin{aligned}
\int_M \Delta_g \operatorname{tr}_g h \cdot e^{-f} dV_g &= \int_M \operatorname{tr}_g h \cdot \Delta_g (e^{-f}) dV_g \\
&= \int_M \operatorname{tr}_g h (-\Delta_g f + |\nabla_g f|_g^2) e^{-f} dV_g,
\end{aligned}$$

$$\begin{aligned}
\int_M \nabla^i \nabla^j h_{ij} \cdot e^{-f} dV_g &= \int_M h_{ij} \nabla^i \nabla^j (e^{-f}) dV \\
&= \int_M h_{ij} (-\nabla^i \nabla^j f + \nabla^i f \nabla^j f) e^{-f} dV_g, \\
\int_M \nabla_i \left(\frac{\partial}{\partial s} f \right) \nabla^i f e^{-f} dV_g &= \int_M -\frac{\partial}{\partial s} f (\Delta_g f - |\nabla_g f|_g^2) e^{-f} dV_g, \\
\int_M \Delta_g (e^{-f}) dV_g &= \int_M (-\Delta_g f + |\nabla_g f|_g^2) e^{-f} dV_g,
\end{aligned}$$

and Lemma 10.1, we obtain

$$\begin{aligned}
&\frac{d}{ds} \Big|_{s=0} v_{\pm}(g(s), u(s)) \\
&= \int_M \frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) (S_{g,u} + |\nabla_g f|_g^2) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g \\
&\quad + \int_M \left(\tau_{\pm} \left(-h^{ij} R_{ij} + \nabla^i \nabla_j h_{ij} - \Delta_g (\text{tr}_g h) + 2h_{pq} \nabla^p u \nabla^q u \right. \right. \\
&\quad \left. \left. - 4\nabla_i v \nabla^i u - h_{pq} \nabla^p f \nabla^q f + 2\nabla_i \left(\frac{\partial}{\partial s} \Big|_{s=0} f(s) \right) \nabla^i f \right) \mp \frac{\partial}{\partial s} \Big|_{s=0} f(s) \right) \\
&\quad \cdot \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g + \int_M \left(-\frac{\partial}{\partial s} \Big|_{s=0} f_{\pm}(s) - \frac{n}{2\tau_{\pm}}(s) \frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) + \frac{1}{2} \text{tr}_g h \right) \\
&\quad \cdot (\tau_{\pm}(S_{g,u} + |\nabla_g f_{\pm}|_g^2) \mp f_{\pm} \pm n) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g.
\end{aligned}$$

If we denote by B the last term and by A the remaining terms,

$$\begin{aligned}
A &= \int_M \left(\frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) (|\nabla_g f_{\pm}|_g^2 + S_{g,u}) \right. \\
&\quad \left. - \tau_{\pm} (h^{ij} \nabla_i \nabla_j f_{\pm} + h^{ij} S_{ij} + 4\nabla_i v \cdot \nabla^i u) \mp \frac{\partial}{\partial s} f_{\pm} \right) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g \\
&\quad + \int_M \tau_{\pm} (\Delta_g f_{\pm} - |\nabla_g f_{\pm}|_g^2) \left(\text{tr}_g h - 2 \frac{\partial}{\partial s} \Big|_{s=0} f(s) \right) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g.
\end{aligned}$$

The normalized condition

$$1 = \int_M \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_g$$

implies

$$0 = \int_M \left(-\frac{\partial}{\partial s} \Big|_{s=0} f_{\pm}(s) - \frac{n}{2\tau_{\pm}} \frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) + \frac{1}{2} \text{tr}_g h \right) \frac{e^{-f_{\pm}(s)}}{(4\pi \tau_{\pm}(s))^{n/2}} dV_g.$$

From Lemma 10.1, we conclude that

$$\tau_{\pm} S_{g,u} - \tau_{\pm} (|\nabla_g f_{\pm}|_g^2 - 2\Delta_g f_{\pm}) = \pm f_{\pm} \mp n + v_{\pm}(g, u).$$

Therefore

$$\tau_{\pm} (S_{g,u} + |\nabla_g f_{\pm}|_g^2) \mp f_{\pm} \pm n = 2\tau_{\pm} (|\nabla_g f_{\pm}|_g^2 - \Delta_g f_{\pm}) + v_{\pm}(g, u).$$

Plugging this into the definition of B yields

$$\begin{aligned} B &= \int_M \left(-\frac{\partial}{\partial s} \Big|_{s=0} f_{\pm}(s) - \frac{n}{2\tau_{\pm}} \frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) + \frac{1}{2} \operatorname{tr}_g h \right) \\ &\quad \cdot (2\tau_{\pm} (|\nabla_g f_{\pm}|_g^2 - \Delta_g f_{\pm}) + v_{\pm}(g, u)) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g \\ &= \int_M \left(-\frac{\partial}{\partial s} \Big|_{s=0} f_{\pm}(s) - \frac{n}{2\tau_{\pm}} \frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) + \frac{1}{2} \operatorname{tr}_g h \right) \\ &\quad \cdot (2\tau_{\pm} (|\nabla_g f_{\pm}|_g^2 - \Delta_g f_{\pm})) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g \\ &= \int_M \left(-\frac{\partial}{\partial s} \Big|_{s=0} f_{\pm}(s) + \frac{1}{2} \operatorname{tr}_g h \right) 2\tau_{\pm} (|\nabla_g f_{\pm}|_g^2 - \Delta_g f_{\pm}) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g, \end{aligned}$$

where we use the fact that

$$\int_M \Delta_g (e^{-f}) dV_g = 0.$$

Hence B cancels with the last term in A . Therefore the above variation equals

$$\begin{aligned} &\frac{d}{ds} \Big|_{s=0} v_{\pm}(g(s), u(s)) \\ &= \int_M \left(\frac{\partial}{\partial s} \Big|_{s=0} \tau_{\pm}(s) \left(|\nabla_g f_{\pm}|_g^2 + S_{g,u} \pm \frac{n}{2\tau_{\pm}} \right) - \tau_{\pm} \left(h^{ij} \nabla_i \nabla_j f + h^{ij} S_{ij} \right. \right. \\ &\quad \left. \left. \pm \frac{1}{2\tau_{\pm}} \operatorname{tr}_g h + 4v(\langle du, df \rangle - \Delta_g u) \right) \right) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV_g. \end{aligned}$$

To prove the theorem, it is sufficient to show that

$$\int_M \left(|\nabla_g f_{\pm}|_g^2 + S_{g,u} \pm \frac{n}{2\tau_{\pm}} \right) \frac{e^{-f_{\pm}}}{(4\pi \tau_{\pm})^{n/2}} dV = 0.$$

Since M is compact, we have

$$0 = \int_M \Delta_g (e^{-f_{\pm}}) = \int_M (-\Delta_g f_{\pm} + |\nabla_g f_{\pm}|_g^2) e^{-f_{\pm}} dV.$$

Hence

$$\begin{aligned} \int_M \left(|\nabla_g f_{\pm}|^2 + S_{g,u} \pm \frac{n}{2\tau_{\pm}} \right) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV \\ = \int_M \left(2\Delta_g f_{\pm} - |\nabla_g f|_g^2 + S_{g,u} \pm \frac{n}{2\sigma_{\pm}} \right) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV. \end{aligned}$$

Lemma 10.1 now indicates

$$\begin{aligned} \int_M \left(|\nabla_g f_{\pm}|^2 + S_{g,u} \pm \frac{n}{2\tau_{\pm}} \right) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV \\ = \int_M \left(\frac{\pm f_{\pm} \mp n + v_{\pm}(g, u)}{\tau_{\pm}} \pm \frac{n}{2} \right) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV \\ = \int_M \frac{1}{\tau_{\pm}} \left(\pm f_{\pm} \mp \frac{n}{2} + v_{\pm}(g, u) \right) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV \\ = \frac{1}{\tau_{\pm}} \left(\pm \frac{n}{2} - v_{\pm}(g, u) \mp \frac{n}{2} + v_{\pm}(g, u) \right) = 0. \end{aligned}$$

The sign + corresponds to the gradient expanding soliton and the sign - to the gradient shrinker soliton. $\square$

Corollary 10.3. *Suppose that (M, g) is a compact Riemannian manifold and u is a smooth function on M . Let h be any symmetric covariant 2-tensor on M and set $g(s) := g + sh$. Let v be any smooth function on M and $u(s) := u + sv$. If $v_{\pm}(g(s), u(s)) = \mathcal{W}_{\pm}(g(s), u(s), f_{\pm}(s), \tau_{\pm}(s))$ for some smooth function $f_{\pm}(s)$ with $\int_M e^{-f_{\pm}(s)} dV / (4\pi\tau_{\pm}(s))^{n/2} = 1$ and a constant $\tau_{\pm}(s) > 0$, and (g, u) is a critical point of $v_{\pm}(\cdot, \cdot)$, then*

$$\mathcal{S}_{g,u} = \mp \frac{1}{2\tau_{\pm}} g, \quad f_{\pm} \equiv \text{constant}.$$

Thus, if $\mathcal{W}_{\pm}(g, u, \cdot, \cdot)$ achieve their minimum and (g, u) is a critical point of $v_{\pm}(\cdot, \cdot)$, (M, g, u) satisfies the static Einstein vacuum equation.

Proof. According to Lemma 10.1 and Theorem 10.2, we have

$$\begin{aligned} \tau_{\pm}(-2\Delta_g f_{\pm} + |\nabla_g f_{\pm}|_g^2 - S_{g,u}) \pm f_{\pm} \mp n \\ = -v_{\pm} = - \int_M \left(\tau_{\pm}(S_{g,u} + |\nabla_g f|_g^2) \mp f_{\pm} \pm n \right) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV_g, \end{aligned}$$

and hence

$$\begin{aligned}
 2\Delta_g f_{\pm} - |\nabla_g f_{\pm}|_g^2 + S_{g,u} &= \int_M (S_{g,u} + |\nabla_g f_{\pm}|_g^2) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV_g \\
 &= \int_M (S_{g,u} + \Delta_g f_{\pm}) \frac{e^{-f_{\pm}}}{(4\pi\tau_{\pm})^{n/2}} dV_g \\
 &= \mp \frac{n}{2\tau_{\pm}} = S_{g,u} + \Delta_g f_{\pm}.
 \end{aligned}$$

From this we get $\Delta_g f_{\pm} = |\nabla_g f_{\pm}|_g^2$. After integrating on both sides, the functions $f_{\pm}$ must be constant, which implies $\mathcal{P}_g \pm (1/(2\tau_{\pm}))g = 0$. $\square$

Remark 10.4. In the situation of Corollary 10.3, by normalization, we may choose $f_{\pm} = n/2$.

Acknowledgements

The author thanks Professor Kefeng Liu, who teaches the author mathematics, Professor Hongwei Xu, and other staff members at the Center of Mathematical Science, where part of this work was done.

References

- [Cao 2007] X. Cao, “Eigenvalues of $(-\Delta + R/2)$ on manifolds with nonnegative curvature operator”, *Math. Ann.* **337**:2 (2007), 435–441. MR 2007g:53071 Zbl 1105.53051
- [Cao 2008] X. Cao, “First eigenvalues of geometric operators under the Ricci flow”, *Proc. Amer. Math. Soc.* **136**:11 (2008), 4075–4078. MR 2009f:53098 Zbl 1166.58007
- [He et al. 2008] C.-L. He, S. Hu, D.-X. Kong, and K. Liu, “Generalized Ricci flow. I. Local existence and uniqueness”, pp. 151–171 in *Topology and physics*, Nankai Tracts Math. **12**, World Sci. Publ., Hackensack, NJ, 2008. MR 2010k:53098
- [Li 2007] J.-F. Li, “Eigenvalues and energy functionals with monotonicity formulae under Ricci flow”, *Math. Ann.* **338**:4 (2007), 927–946. MR 2008c:53068 Zbl 1127.53059
- [Li 2012] Y. Li, “Generalized Ricci flow I: higher derivative estimates for compact manifolds”, *Analysis & PDE* **5**:4 (2012), 747–775.
- [List 2006] B. List, *Evolution of an extended Ricci flow system*, Ph.D. thesis, Fachbereich Mathematik und Informatik der Freie Universität Berlin, 2006, Available at <http://www.diss.fu-berlin.de/2006/180/index.html>.
- [Müller 2012] R. Müller, “Ricci flow coupled with harmonic map flow”, *Ann. Sci. Éc. Norm. Supér.* (4) **45**:1 (2012), 101–142. MR 2961788
- [Oliynyk et al. 2006] T. Oliynyk, V. Suneeta, and E. Woolgar, “A gradient flow for worldsheet nonlinear sigma models”, *Nuclear Phys. B* **739**:3 (2006), 441–458. MR 2006m:81185 Zbl 1109.81058
- [Streets 2007] J. D. Streets, *Ricci Yang–Mills flow*, Ph.D. thesis, Duke University, 2007, Available at <http://www.math.uci.edu/~jstreets/papers/StreetsThesis.pdf>.
- [Streets 2008] J. Streets, “Regularity and expanding entropy for connection Ricci flow”, *J. Geom. Phys.* **58**:7 (2008), 900–912. MR 2009f:53105 Zbl 1144.53326

- [Streets 2009] J. Streets, “Singularities of renormalization group flows”, *J. Geom. Phys.* **59**:1 (2009), 8–16. MR 2010a:53143 Zbl 1153.53329
- [Streets 2010] J. Streets, “Ricci Yang–Mills flow on surfaces”, *Adv. Math.* **223**:2 (2010), 454–475. MR 2011c:53164 Zbl 1190.53069
- [Young 2008] A. N. Young, *Modified Ricci flow on a principal bundle*, Ph.D. thesis, The University of Texas at Austin, 2008, Available at <http://search.proquest.com/docview/193674070>.

Received August 27, 2012. Revised January 6, 2013.

YI LI

DEPARTMENT OF MATHEMATICS

JOHNS HOPKINS UNIVERSITY

3400 N. CHARLES STREET

BALTIMORE, MD 21218

UNITED STATES

yli@math.jhu.edu

Current address:

DEPARTMENT OF MATHEMATICS

SHANGHAI JIAO TONG UNIVERSITY

800 DONG CHUAN ROAD, MIN HANG DISTRICT

SHANGHAI, 200240

CHINA

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic.

Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 267 No. 1 January 2014

Numerical study of unbounded capillary surfaces	1
YASUNORI AOKI and HANS DE STERCK	
Dual R -groups of the inner forms of $SL(N)$	35
KUOK FAI CHAO and WEN-WEI LI	
Automorphisms and quotients of quaternionic fake quadrics	91
AMIR DŽAMBIĆ and XAVIER ROULLEAU	
Distance of bridge surfaces for links with essential meridional spheres	121
YEONHEE JANG	
Normal states of type III factors	131
YASUYUKI KAWAHIGASHI, YOSHIKO OGATA and ERLING STØRMER	
Eigenvalues and entropies under the harmonic-Ricci flow	141
YI LI	
Quantum extremal loop weight modules and monomial crystals	185
MATHIEU MANSUY	
Lefschetz fibrations with small slope	243
NAOYUKI MONDEN	



0030-8730(2014)267:1;1-7