

*Pacific
Journal of
Mathematics*

LEFSCHETZ FIBRATIONS WITH SMALL SLOPE

NAOYUKI MONDEN

LEFSCHETZ FIBRATIONS WITH SMALL SLOPE

NAOYUKI MONDEN

We construct Lefschetz fibrations over S^2 which do not satisfy the slope inequality. This disproves a conjecture of Hain.

1. Introduction

Lefschetz fibrations have been an active area of research ever since the remarkable work in [Donaldson 1999] and [Gompf and Stipsicz 1999] revealed a close connection between them and symplectic 4-manifolds. In this paper, we consider the geography problem of Lefschetz fibrations over S^2 , which derives from that of complex surfaces fibred over curves.

We are interested in two kinds of geography problems. Let σ and e be the signature and the Euler characteristic of a closed oriented smooth 4-manifold X , respectively. For an almost complex closed 4-manifold X , we set $K^2 := 3\sigma + 2e$ and $\chi_h := (\sigma + e)/4$ (the *holomorphic Euler characteristic*).

One is the geography problem for complex surfaces: the characterization of pairs (K^2, χ_h) corresponding to minimal complex surfaces. It is well known that any minimal complex surface of general type satisfies $K^2 > 0$, $\chi_h > 0$, the *Noether inequality* $2\chi_h - 6 \leq K^2$ and the *Bogomolov–Miyaoka–Yau inequality* $K^2 \leq 9\chi_h$ (see [Barth et al. 1984], for example). The above geography problem can be extended to the symplectic 4-manifolds. However, Fintushel and Stern [1998] constructed Lefschetz fibration which does not satisfy the Noether inequality. In particular, for most pairs (p, q) satisfying $p < 2q - 6$, there exists a minimal symplectic 4-manifold with $p = K^2$ and $q = \chi_h$ (see [Gompf and Stipsicz 1999]). On the other hand, no examples of a minimal symplectic 4-manifold with $K^2 > 9\chi_h$ have been found yet.

The other is the geography problem for complex surfaces fibred over curves. Hereafter, we assume $g \geq 2$. Let $f : S \rightarrow C$ be a relatively minimal holomorphic genus- g fibration, where S is a complex surface and C is a complex curve of genus k . We define relative numerical invariants $\chi_f := \chi_h - (g - 1)(k - 1)$ and $K_f^2 := K^2 - 8(g - 1)(k - 1)$ for $f : S \rightarrow C$. Then, we have two inequalities $\chi_f \geq 0$

MSC2010: primary 57R20; secondary 57N13, 14D06, 20F38.

Keywords: Lefschetz fibration, mapping class group, slope.

and $K_f^2 \geq 0$, known as *Beauville's inequality* (see [Beauville 1979]) and *Arakelov's inequality* (see [Arakelov 1971]), respectively. For $\chi_f \neq 0$, which is equivalent to the fact that f is not a holomorphic bundle, we define λ_f to be the quotient K_f^2/χ_f , called the slope of f . Xiao [1987] proved that $4 - 4/g \leq \lambda_f \leq 12$ (that is, $(4 - 4/g)\chi_f \leq K_f^2 \leq 12\chi_f$). The former inequality is called the *slope inequality*. For a relatively minimal genus- g Lefschetz fibration, χ_f , K_f^2 and the slope λ_f are defined in the same way as for complex surfaces fibred over curves. To the author's knowledge, the slope of all known Lefschetz fibrations over S^2 is greater than or equal to $4 - 4/g$.

Conjecture 1.1 (Hain; see [Amorós et al. 2000, Question 5.10; Endo and Nagami 2005, Conjecture 4.12]). *For every relatively minimal genus- g Lefschetz fibration $f : X \rightarrow S^2$, the slope inequality $\lambda_f \geq 4 - 4/g$ holds.*

In this paper, we give a negative answer to [Conjecture 1.1](#).

Theorem 3.1. *For each $g \geq 3$, there exists a genus- g Lefschetz fibration over S^2 with slope $\lambda_f = 4 - 4/g - 1/3g$ whose total space is simply connected.*

Moreover, by fiber sum operations, we have the following results:

Corollary 3.6. *For each $g \geq 3$, $m \geq 0$ and $l \geq 0$, there exists a genus- g Lefschetz fibration $f_{m,l} : X_{m,l} \rightarrow S^2$ with slope $\lambda_{f_{m,l}} = 4 - 4/g - 1/(m+3)g$ such that $\pi_1(X_{m,l}) = 1$. Moreover, if $(m, l) \neq (0, 0)$, then $X_{m,l}$ is a minimal symplectic 4-manifold.*

Corollary 3.7. *For each $g \geq 3$, $m \geq 1$ and $l \geq 0$, there exists a genus- g Lefschetz fibration $f'_{m,l} : Y_{m,l} \rightarrow S^2$ with slope $\lambda_{f'_{m,l}} = 4 - 4/g - 1/2g + 1/(2 \cdot 3^{m-1}g)$ such that $\pi_1(Y_{m,l}) = 1$. Moreover, if $l \geq 1$, then $Y_{m,l}$ is a minimal symplectic 4-manifold.*

As a consequence, we have the following results.

Corollary 4.2. *The Lefschetz fibrations $f_{m,l}$ ($m \geq 0$) and $f'_{m,l}$ ($m \geq 2$) are non-holomorphic.*

Let $f : X \rightarrow S^2$ be a relatively minimal genus- g Lefschetz fibration with $n > 0$ singular fibers. From $e(X) = -4(g-1) + n$ and results from [Smith 1999; Stipsicz 1999; Ozbagci 2002], we have $\chi_f > 0$, $K_f^2 \geq 4g - 4$ and $\lambda_f \leq 10$. Moreover, it is well known that any hyperelliptic Lefschetz fibration satisfies the slope inequality. This fact follows from the signature formula for genus- g hyperelliptic Lefschetz fibrations obtained by Matsumoto [1983; 1996] for $g = 1, 2$ and Endo [2000] for $g \geq 3$. Therefore, genus-2 Lefschetz fibrations satisfy the slope inequality. In particular, if f is a hyperelliptic Lefschetz fibration with only nonseparating vanishing cycles, then λ_f is equal to $4 - 4/g$. For Lefschetz fibrations with $b_2^+ = 1$, we prove the following result.

Theorem 5.1. *Let $g \geq 2$ and let $f : X \rightarrow S^2$ be a genus- g Lefschetz fibration with $b_2^+(X) = 1$.*

(1) *If X is not diffeomorphic to the blow-up of a ruled surface, then*

- (i) $4 - 4/g \leq \lambda_f \leq 8$ for $b_1(X) = 0$,
- (ii) $4 \leq \lambda_f \leq 8$ for $b_1(X) = 2$.

(2) *If X is diffeomorphic to the blow-up of an S^2 -bundle over Σ_k , then*

$$4 + 4(k - 1)/(g - k) \leq \lambda_f \leq 8,$$

and the lower bound is sharp.

The study of the slope of holomorphic fibrations was mainly motivated by Severi's inequality, which states that if S is a minimal surface of general type of maximal Albanese dimension, then $K^2 \geq 4\chi_h$. Equivalently, if $K^2 < 4\chi_h$, then S is a surface fibred over C of genus $b_1(S)/2$. Severi [1932] claimed it, but his proof was not correct (see [Catanese 1983]). The inequality was independently posed as a conjecture by Reid [1979] and by Catanese [1983]. Xiao [1987] proved the conjecture when S is a surface fibred over a curve of positive genus. He showed that if S admits a holomorphic genus- g fibration f over C of positive genus k with $K^2 < 4\chi_h + 4(g - 1)(k - 1)$ (that is, $\lambda_f < 4$), then $k = b_1(S)/2$. Konno [1996] proved it in the case of *even* surfaces. The conjecture was solved by Manetti [2003] when S has ample canonical bundle. Pardini [2005] proved the conjecture completely by using the slope inequality for holomorphic fibrations over $\mathbb{C}\mathbb{P}^1$.

In Section 2, we review some standard facts on Lefschetz fibrations. Our main results are proved in Section 3. We give Lefschetz fibrations which violate the slope inequality. Consequently, we obtain examples of nonholomorphic Lefschetz fibrations in Section 4. In the last section, we investigate the slopes of Lefschetz fibrations with $b_2^+ = 1$.

Remark 1.2. The slope inequality of Conjecture 1.1 can be reformulated in terms of the Deligne–Mumford compactified moduli space of stable curves of genus g , denoted by $\overline{\mathcal{M}}_g$, as follows. For a relatively minimal genus- g Lefschetz fibration $f : X \rightarrow S^2$ with n singular fibers, we obtain a symplectic structure on X such that for all $x \in S^2$, $f^{-1}(x)$ is a pseudoholomorphic curve. Since a 2-dimensional almost-complex structure is integrable, $f^{-1}(x)$ determines a point in $\overline{\mathcal{M}}_g$. Thus, we obtain the moduli map $\phi_f : S^2 \rightarrow \overline{\mathcal{M}}_g$ which is defined by $\phi_f(x) = [f^{-1}(x)] \in \overline{\mathcal{M}}_g$ for $x \in S^2$. We denote by \mathcal{H}_g the Hodge bundle on $\overline{\mathcal{M}}_g$ with fiber the determinant line $\bigwedge^g H^0(C; K_C)$, where C is the set of critical points of f . By Smith's signature formula [1999] and the slope inequality, we have the following inequality:

$$(8g + 4)\langle c_1(\mathcal{H}_g), [\phi_f(S^2)] \rangle - g \cdot n \geq 0.$$

2. Preliminaries

In this section, we first recall the definition and basic properties of Lefschetz fibrations. More details can be found in [Gompf and Stipsicz 1999].

Let Σ_g be a closed oriented surface of genus $g \geq 2$ and let Γ_g be the *mapping class group* of Σ_g , which is the group of isotopy classes of orientation-preserving diffeomorphisms of Σ_g . We denote by t_c the right-handed *Dehn twist* about a simple closed curve c on an oriented surface. The notation $t_c t_d$ means that we first apply t_d then t_c .

Definition 2.1. Let X be a closed, oriented smooth 4-manifold. A smooth map $f : X \rightarrow S^2$ is a *genus- g Lefschetz fibration* if it satisfies the following conditions:

- (i) f has finitely many critical values $b_1, \dots, b_n \in S^2$, and f is a smooth Σ_g -bundle over $S^2 - \{b_1, \dots, b_n\}$.
- (ii) For each i ($i = 1, \dots, n$), there exists a unique critical point p_i in the *singular fiber* $f^{-1}(b_i)$ such that about each p_i and b_i there are local complex coordinate charts agreeing with the orientations of X and S^2 on which f is of the form $f(z_1, z_2) = z_1^2 + z_2^2$.
- (iii) f is relatively minimal (no fiber contains a (-1) -sphere).

Each singular fiber is obtained by collapsing a simple closed curve (the *vanishing cycle*) in the regular fiber. The monodromy of the fibration around a singular fiber is given by a right-handed Dehn twist along the corresponding vanishing cycle. A Lefschetz fibration $f : X \rightarrow S^2$ is *holomorphic* if there are complex structures on both X and S^2 with holomorphic projection f .

Once we fix an identification of Σ_g with the fiber over a base point of S^2 , we can characterize the Lefschetz fibration $f : X \rightarrow S^2$ by its *monodromy representation* $\pi_1(S^2 - \{b_1, \dots, b_n\}) \rightarrow \Gamma_g$. This map is really an antihomomorphism, since elements of $\pi_1(S^2 - \{b_1, \dots, b_n\})$ are written left-to-right and elements of Γ_g are written right-to-left. Let $\gamma_1, \dots, \gamma_n$ be an ordered system of generating loops for $\pi_1(S^2 - \{b_1, \dots, b_n\})$, such that each γ_i encircles only b_i and $\prod \gamma_i$ is homotopically trivial. Thus, the monodromy of f comprises a factorization

$$t_{v_n} \dots t_{v_2} t_{v_1} = 1 \in \Gamma_g,$$

where v_i are vanishing cycles of the singular fibers. This factorization is called the *positive relator*.

According to theorems of Kas [1980] and Matsumoto [1996], if $g \geq 2$, then the isomorphism class of a Lefschetz fibration is determined by a positive relator modulo simultaneous conjugations

$$t_{v_n} \dots t_{v_2} t_{v_1} \sim t_{\phi(v_n)} \dots t_{\phi(v_2)} t_{\phi(v_1)} \quad \text{for all } \phi \in \Gamma_g$$




Figure 1. The curves $d_1, d_2, d_3, d_4, x_1, x_2, x_3$.

and elementary transformations

$$t_{v_n} \cdots t_{v_{i+2}} t_{v_{i+1}} t_{v_i} t_{v_{i-1}} t_{v_{i-2}} \cdots t_{v_1} \sim t_{v_n} \cdots t_{v_{i+2}} t_{v_i} t_{v_{i-1}}^{-1} t_{v_{i+1}} t_{v_{i-1}} t_{v_{i-2}} \cdots t_{v_1},$$

$$t_{v_n} \cdots t_{v_{i+2}} t_{v_{i+1}} t_{v_i} t_{v_{i-1}} t_{v_{i-2}} \cdots t_{v_1} \sim t_{v_n} \cdots t_{v_{i+2}} t_{v_{i+1}} t_{v_i} t_{v_{i-1}} t_{v_i} t_{v_{i-2}} \cdots t_{v_1}.$$

Note that $\phi t_{v_i} \phi^{-1} = t_{\phi(v_i)}$. For all $\phi \in \Gamma_g$, let $\phi(\varrho)$ be the positive relator which is obtained by applying simultaneous conjugations by ϕ to a positive relator ϱ . We denote a Lefschetz fibration associated to a positive relator $\varrho \in \Gamma_g$ by $f_\varrho : X_\varrho \rightarrow S^2$. Clearly, if $\varrho_1 \sim \varrho_2$ in Γ_g (that is, ϱ_2 is obtained by applying elementary transformations or simultaneous conjugations to ϱ_1), then

$$\chi_{f_{\varrho_1}} = \chi_{f_{\varrho_2}} \quad \text{and} \quad K_{f_{\varrho_1}}^2 = K_{f_{\varrho_2}}^2.$$

For positive relators ϱ_1 and ϱ_2 in Γ_g , the genus- g Lefschetz fibration

$$f_{\varrho_1 \varrho_2} : X_{\varrho_1 \varrho_2} \rightarrow S^2$$

is the (trivial) fiber sum of f_{ϱ_1} and f_{ϱ_2} . Since $\sigma(X_{\varrho_1 \varrho_2}) = \sigma(X_{\varrho_1}) + \sigma(X_{\varrho_2})$ and $e(X_{\varrho_1 \varrho_2}) = e(X_{\varrho_1}) + e(X_{\varrho_2}) + 4(g-1)$, we see that $\chi_{f_{\varrho_1 \varrho_2}} = \chi_{f_{\varrho_1}} + \chi_{f_{\varrho_2}}$ and $K_{f_{\varrho_1 \varrho_2}}^2 = K_{f_{\varrho_1}}^2 + K_{f_{\varrho_2}}^2$. In particular, if $\varrho_1 \sim \varrho_2$, then

$$\chi_{f_{\varrho_1 \varrho_2}} = 2\chi_{f_{\varrho_1}} = 2\chi_{f_{\varrho_2}} \quad \text{and} \quad K_{f_{\varrho_1 \varrho_2}}^2 = 2K_{f_{\varrho_1}}^2 = 2K_{f_{\varrho_2}}^2.$$

We next begin with a definition of the lantern relation (see [Dehn 1938; Johnson 1979]).

Definition 2.2. Let Σ_0^4 denote a sphere with 4 boundary components. Let d_1, d_2, d_3, d_4 be the 4 boundary curves of Σ_0^4 and let x_1, x_2, x_3 be the interior curves as shown in Figure 1. Then, we have the *lantern relation*

$$t_{d_1} t_{d_2} t_{d_3} t_{d_4} = t_{x_1} t_{x_2} t_{x_3}.$$

Let ϱ be a positive relator of Γ_g . Let $d_1, d_2, d_3, d_4, x_1, x_2, x_3$ be curves as in Definition 2.2. Suppose that ϱ includes $t_{d_1} t_{d_2} t_{d_3} t_{d_4}$ as a subword:

$$\varrho = U \cdot t_{d_1} t_{d_2} t_{d_3} t_{d_4} \cdot V,$$

where U and V are products of right-handed Dehn twists. Then, by the lantern relation, the product of right-handed Dehn twists

$$\varrho' = U \cdot t_{x_1} t_{x_2} t_{x_3} \cdot V$$

is also a positive relator of Γ_g .

This operation is one of substitution techniques introduced by Fuller.

Definition 2.3. We say that ϱ' is obtained by applying an L -substitution to ϱ . Conversely, ϱ is said to be obtained by applying an L^{-1} -substitution to ϱ' . We also call these two kinds of operations *lantern substitutions*.

Proposition 2.4 [Endo and Nagami 2005, Theorem 4.3 and Proposition 3.12]. *Let ϱ, ϱ' be positive relators of Γ_g and let $X_\varrho, X_{\varrho'}$ be the corresponding Lefschetz fibrations over S^2 , respectively. Suppose that ϱ is obtained by applying an L^{-1} -substitution to ϱ' . Then, $\sigma(X_\varrho) = \sigma(X_{\varrho'}) - 1$ and $e(X_\varrho) = e(X_{\varrho'}) + 1$. Therefore,*

$$\chi_{f_\varrho} = \chi_{f_{\varrho'}} \quad \text{and} \quad K_{f_\varrho}^2 = K_{f_{\varrho'}}^2 - 1.$$

Remark 2.5. Endo and Gurtas [2010] showed that $X_{\varrho'}$ is a rational blowdown of X_ϱ introduced by Fintushel and Stern [1997]. Such relations were also generalized by Endo, Mark, and Van Horn-Morris [Endo et al. 2011].

3. Main results

In this section, we give a negative answer to [Conjecture 1.1](#).

Theorem 3.1. *For each $g \geq 3$, there exists a genus- g Lefschetz fibration over S^2 with slope $\lambda_f = 4 - 4/g - 1/3g$ whose total space is simply connected.*

In order to prove [Theorem 3.1](#), we recall some standard facts on hyperelliptic Lefschetz fibrations. Let Δ_g be the *hyperelliptic mapping class group* of genus g , that is, the subgroup of Γ_g which consists of all isotopy classes of orientation-preserving diffeomorphisms of Σ_g commuting with the isotopy class of ι , called the hyperelliptic involution. Note that $\Delta_g = \Gamma_g$ for $g = 1, 2$.

Definition 3.2. Let $\varrho = t_{a_1} \dots t_{a_n}$ be a positive relator in Γ_g . A genus- g Lefschetz fibration $f_\varrho : X_\varrho \rightarrow S^2$ is called *hyperelliptic* if for each $k \in \{1, \dots, n\}$, t_{a_k} is in Δ_g . Equivalently, $\iota(a_k) = a_k$ for each k .

The following theorem was established in [Matsumoto 1983; 1996] for $g = 1, 2$ and in [Endo 2000] for $g \geq 3$.




Figure 2. The curves $c_1, \dots, c_{2g+1}, x_1, x_2, x_3, y$.

Theorem 3.3 (Matsumoto, Endo). *Let $f_\varrho : X_\varrho \rightarrow S^2$ be a genus- g hyperelliptic Lefschetz fibration with m nonseparating and*

$$s = \sum_{h=1}^{\lfloor g/2 \rfloor} s_h$$

separating vanishing cycles, where s_h denotes the number of separating vanishing cycles that separate Σ_g into two surfaces, one of which has genus h . Then, we have

$$\sigma(X_\varrho) = -\frac{g+1}{2g+1}m + \sum_{h=1}^{\lfloor g/2 \rfloor} \left(\frac{4h(g-h)}{2g+1} - 1 \right) s_h.$$

We need the following positive relator to prove [Theorem 3.1](#). As shown in [Figure 2](#), let $c_1, c_2, \dots, c_{2g+1}$ be the curves in Σ_g . We denote by $h_g (\in \Gamma_g)$ the product of $8g + 4$ right-handed Dehn twists

$$h_g := (t_{c_1} t_{c_2} \dots t_{c_{2g+1}}^2 \dots t_{c_2} t_{c_1})^2.$$

It is well known that h_g is a positive relator in Δ_g and that $\sigma(X_{h_g}) = -4(g+1)$, by [Theorem 3.3](#) and $e(X_{h_g}) = 4(g+2)$. This gives $\chi_{f_{h_g}} = g$, $K_{f_{h_g}}^2 = 4g - 4$ and $\lambda_{f_{h_g}} = 4 - 4/g$ (that is, f_{h_g} is lying on the slope line).

Proof of [Theorem 3.1](#). Suppose $g \geq 3$. Let x_1, x_2, x_3, y be the curves as shown in [Figure 2](#). Since c_1, x_i are nonseparating curves, there exists a diffeomorphism ϕ_i such that $\phi_i(c_1) = x_i$. Hence, we have the following positive relator r_i ($i = 1, 2, 3$):

$$\begin{aligned} r_i &= \phi_i h_g \phi_i^{-1} = \phi_i (t_{c_1} t_{c_2} \dots t_{c_{2g+1}}^2 \dots t_{c_2} t_{c_1})^2 \phi_i^{-1} \\ &= (t_{\phi_i(c_1)} t_{\phi_i(c_2)} \dots t_{\phi_i(c_{2g+1})}^2 \dots t_{\phi_i(c_2)} t_{\phi_i(c_1)})^2 \\ &= (t_{x_i} t_{\phi_i(c_2)} \dots t_{\phi_i(c_{2g+1})}^2 \dots t_{\phi_i(c_2)} t_{\phi_i(c_1)})^2 \\ &= 1 \in \Gamma_g. \end{aligned}$$

Let $r'_g = r_1 r_2 r_3 = (t_{x_1} \dots t_{\phi_1(c_1)})^2 (t_{x_2} \dots)^2 (t_{x_3} \dots)^2$. Since $f_{r'_g}$ is the fiber sum of f_{r_1}, f_{r_2} and f_{r_3} which are obtained by applying simultaneous conjugations to h_g , we have

$$\chi_{f_{r'_g}} = 3\chi_{f_{h_g}} = 3g \quad \text{and} \quad K_{f_{r'_g}}^2 = 3K_{f_{h_g}}^2 = 3(4g - 4).$$

We apply elementary transformations to r'_g as follows:

$$\begin{aligned}
 r'_g &= r_1 r_2 r_3 \\
 &= t_{x_1} t_{\phi_1(c_2)} \cdots t_{\phi_1(c_2)} t_{\phi_1(c_1)} \cdot t_{x_2} t_{\phi_2(c_2)} \cdots t_{\phi_2(c_1)} \cdot t_{x_3} t_{\phi_3(c_2)} \cdots t_{\phi_3(c_1)} \\
 &\sim t_{x_1} t_{\phi_1(c_2)} \cdots t_{\phi_1(c_2)} t_{x_2} t_{x_2^{-1}(\phi_1(c_1))} t_{\phi_2(c_2)} \cdots t_{\phi_2(c_1)} \cdot t_{x_3} t_{\phi_3(c_2)} \cdots t_{\phi_3(c_1)} \\
 &\quad \vdots \qquad \qquad \qquad \vdots \\
 &\sim t_{x_1} t_{x_2} t_{x_2^{-1}(\phi_1(c_2))} \cdots t_{x_2^{-1}(\phi_1(c_2))} t_{x_2^{-1}(\phi_1(c_1))} t_{\phi_2(c_2)} \cdots t_{\phi_2(c_1)} \cdot t_{x_3} t_{\phi_3(c_2)} \cdots t_{\phi_3(c_1)} \\
 &\sim t_{x_1} t_{x_2} t_{x_2^{-1}(\phi_1(c_2))} \cdots t_{x_2^{-1}(\phi_1(c_2))} t_{x_2^{-1}(\phi_1(c_1))} t_{\phi_2(c_2)} \cdots t_{x_3} t_{x_3^{-1}(\phi_2(c_1))} t_{\phi_3(c_2)} \cdots t_{\phi_3(c_1)} \\
 &\quad \vdots \qquad \qquad \qquad \vdots \\
 &\sim (t_{x_1} t_{x_2} t_{x_3}) W,
 \end{aligned}$$

where W is a product of $24g + 9$ right-handed Dehn twists. By the lantern relation, we get the following positive relator r_g :

$$r_g := (t_{c_1} t_{c_3} t_{c_5} t_y) W.$$

Since r_g is obtained by applying an L^{-1} -substitution to r'_g , by [Proposition 2.4](#)

$$\chi_{f_{r_g}} = 3g \quad \text{and} \quad K_{f_{r_g}}^2 = 3(4g - 4) - 1.$$

Then, the slope of f_{r_g} is equal to $4 - 4/g - 1/3g$.

Since it is easy to check that r_g includes the Dehn twist about a curve $\phi_3(c_i)$ for $1 \leq i \leq 2g + 1$, $\pi_1(X_{r_g}) = 1$. This follows from [\[Gompf and Stipsicz 1999\]](#) and the fact that f_{r_g} has a section. This completes the proof of [Theorem 3.1](#). □

Remark 3.4. Since r_g is obtained by applying an L^{-1} -substitution to r'_g , X_{r_g} is a rational blow-up of $X_{r'_g}$. By applying elementary transformations to a relator corresponding to a Lefschetz fibration which is obtained by taking a twisted fiber sum with sufficiently many Lefschetz fibrations, we obtain a positive relator such that we can apply a monodromy substitution, which corresponds to the operation of rational blowdown (resp. rational blow-up) in [\[Endo et al. 2011\]](#), to it.

Remark 3.5. Miyachi and Shiga [\[2011\]](#) produced genus- g Lefschetz fibrations over Σ_{2m} which do not satisfy the slope inequality.

Moreover, by fiber sum operations, we have the following results:

Corollary 3.6. *For each $g \geq 3$, $m \geq 0$ and $l \geq 0$, there exists a genus- g Lefschetz fibration $f_{m,l} : X_{m,l} \rightarrow S^2$ with slope $\lambda_{f_{m,l}} = 4 - 4/g - 1/(m + 3)g$ such that $\pi_1(X_{m,l}) = 1$. Moreover, if $(m, l) \neq (0, 0)$, then $X_{m,l}$ is a minimal symplectic 4-manifold.*

Proof of Corollary 3.6. For any $m \geq 0$, we consider the Lefschetz fibration $f_{r_g h_g^m} : X_{r_g h_g^m} \rightarrow S^2$ which is the fiber sum of f_{r_g} and m copies of f_{h_g} . Then,

$$\begin{aligned}\chi_{f_{r_g h_g^m}} &= \chi_{f_{r_g}} + m \chi_{f_{h_g}} = (3 + m)g, \\ K_{f_{r_g h_g^m}}^2 &= K_{f_{r_g}}^2 + m K_{f_{h_g}}^2 = (3 + m)(4g - 4) - 1.\end{aligned}$$

Therefore, we obtain

$$(1) \quad \lambda_{f_{r_g h_g^m}} = 4 - 4/g - 1/(m + 3)g.$$

Let $f_{m,l} : X_{m,l} \rightarrow S^2$ be the fiber sum of l copies of $f_{r_g h_g^m}$ (that is, $f_{m,l} = f_{(r_g h_g^m)^l}$). By Using (1) and an argument similar to the proof of Theorem 3.1, we have $\lambda_{f_{m,l}} = 4 - 4/g - 1/(m + 3)g$ and $\pi_1(X_{m,l}) = 1$. By the result of Usher [2006], $X_{m,l}$ is minimal for $(m, l) \neq (0, 0)$. This completes the proof. \square

Corollary 3.7. *For each $g \geq 3$, $m \geq 1$ and $l \geq 0$, there exists a genus- g Lefschetz fibration $f'_{m,l} : Y_{m,l} \rightarrow S^2$ with slope $\lambda_{f'_{m,l}} = 4 - 4/g - 1/2g + 1/(2 \cdot 3^{m-1}g)$ such that $\pi_1(Y_{m,l}) = 1$. Moreover, if $l \geq 1$, then $Y_{m,l}$ is a minimal symplectic 4-manifold.*

Proof of Corollary 3.7. When we apply the argument of Theorem 3.1 again, with $\varrho_1 = h_g$ replaced by $\varrho_2 = r_g$, we obtain a genus- g Lefschetz fibration $f_{\varrho_3} : X_{\varrho_3} \rightarrow S^2$ with

$$\begin{aligned}\chi_{f_{\varrho_3}} &= 3\chi_{f_{\varrho_2}} = 3 \cdot 3\chi_{f_{e_1}} \\ K_{f_{\varrho_3}}^2 &= 3K_{f_{\varrho_2}}^2 - 1 = 3(3K_{f_{e_1}}^2 - 1) - 1.\end{aligned}$$

By repeating this argument, we get a genus- g Lefschetz fibration $f_{\varrho_m} (m \geq 1)$ with

$$\begin{aligned}\chi_{f_{\varrho_m}} &= 3^{m-1}\chi_{f_{e_1}} = 3^{m-1}g, \\ K_{f_{\varrho_m}}^2 &= 3(\dots(3(3K_{f_{e_1}}^2 - 1) - 1)\dots) - 1 = 3^{m-1}K_{f_{e_1}}^2 - 3^{m-2} - \dots - 3 - 1 \\ &= 3^{m-1}(4g - 4) - (3^{m-1} - 1)/2.\end{aligned}$$

Therefore, $\lambda_{f_{\varrho_m}} = 4 - 4/g - 1/2g + 1/(2 \cdot 3^{m-1}g)$.

Let $f'_{m,l} : Y_{m,l} \rightarrow S^2$ be the fiber sum of l copies of f_{ϱ_m} , and so $\lambda_{f'_{m,l}} = 4 - 4/g - 1/2g + 1/(2 \cdot 3^{m-1}g)$. Similar to the proof of Corollary 3.6, we see that $\pi_1(Y_{m,l}) = 1$ and that $Y_{m,l}$ is minimal for $l \geq 1$. \square

4. Nonholomorphic Lefschetz fibrations

There are various kinds of nonholomorphic Lefschetz fibrations. By fiber summing two copies of genus-2 Lefschetz fibration due to Matsumoto [1996], Ozbagci and Stipsicz [2000] constructed nonholomorphic genus-2 Lefschetz fibrations whose total space does not admit a complex structure. Korkmaz [2001] generalized their examples to $g \geq 3$. The above mentioned examples of Fintushel and Stern are

also nonholomorphic Lefschetz fibrations. From the study of divisors in moduli space, Smith [2001] showed that a genus-3 Lefschetz fibration over S^2 which was produced by Fuller is nonholomorphic. Endo and Nagami [2005] constructed some examples of nonholomorphic Lefschetz fibrations which violate lower bounds of the slope for nonhyperelliptic fibrations of genus 3, 4 and 5 from the results of Konno [1991; 1993] and Chen [1993]. Hirose [2010] also gave some examples of $g = 3, 4$. In this section, we give new examples of nonholomorphic Lefschetz fibrations.

From the slope inequality for holomorphic fibrations, we have the following necessary condition for a Lefschetz fibration to be holomorphic:

Proposition 4.1 [Xiao 1987]. *If a Lefschetz fibration f is holomorphic, then the slope inequality $\lambda_f \geq 4 - 4/g$ holds.*

As a consequence, we have the following results.

Corollary 4.2. *The Lefschetz fibrations $f_{m,l}$ ($m \geq 0$) and $f'_{m,l}$ ($m \geq 2$) are non-holomorphic.*

Remark 4.3. The above mentioned examples of Fintushel and Stern satisfy the slope inequality but violate the Noether inequality. On the other hand, $f_{m,l}$ and $f'_{m,l}$ satisfy the Noether inequality but violate the slope inequality. Therefore, these two necessary conditions for a Lefschetz fibration to be holomorphic are independent in the sense that neither one implies the other.

5. The slopes of Lefschetz fibrations with $b_2^+ = 1$

We have the following natural question: Which Lefschetz fibrations satisfy the slope inequality? By Proposition 4.1, holomorphic Lefschetz fibrations satisfy the slope inequality. If a Lefschetz fibration is hyperelliptic, then $\lambda_f \geq 4 - 4/g$. This fact can be proved as follows. In the notation of Theorem 3.3, we have $e(X_\rho) = -4(g-1) + (m+s)$. Then, since $h \in \{1, \dots, [g/2]\}$ and $g \geq 2$, we have

$$K_{f_e}^2 - (4 - 4/g)\chi_{f_e} = \sum_{h=1}^{[g/2]} \frac{4h(g-h)-g}{g} s_h \geq 0.$$

In particular, this means that for any hyperelliptic Lefschetz fibrations with only nonseparating vanishing cycles, $\lambda_f = 4 - 4/g$.

In this section, we investigate the slopes of Lefschetz fibrations with $b_2^+ = 1$. By combining the results of [Stipsicz 1999; 2002] and [Li 2000], we can show that Lefschetz fibrations with $b_2^+ = 1$ satisfy the slope inequality. Stipsicz showed that if $X \rightarrow S^2$ is a genus- g Lefschetz fibration over S^2 with $b_2^+(X) = 1$ and X is not diffeomorphic to the blow-up of a ruled surface (that is, diffeomorphic to an S^2 -bundle over Σ_k), then $b_1(X) \in \{0, 2\}$ and $e \geq 0$ (see [Stipsicz 1999, Corollary 3.3

and 3.5]). In particular, if X is the blow-up of an S^2 -bundle over Σ_k , then $k \leq g/2$ (see [Li 2000, Proposition 4.4]). Then, we obtain the following result.

Theorem 5.1. *Let $g \geq 2$ and let $f : X \rightarrow S^2$ be a genus- g Lefschetz fibration with $b_2^+(X) = 1$.*

(1) *If X is not diffeomorphic to the blow-up of a ruled surface, then*

$$(i) \quad 4 - 4/g \leq \lambda_f \leq 8 \quad \text{for } b_1(X) = 0,$$

$$(ii) \quad 4 \leq \lambda_f \leq 8 \quad \text{for } b_1(X) = 2.$$

(2) *If X is diffeomorphic to the blow-up of an S^2 -bundle over Σ_k , then*

$$4 + 4(k - 1)/(g - k) \leq \lambda_f \leq 8,$$

and the lower bound is sharp.

An improvement of the previous result was suggested by the referee.

Proof of Theorem 5.1. For a genus- g Lefschetz fibration, a regular fiber has zero self-intersection. Since the intersection form is nondegenerate, it follows that $b_2^\pm \geq 1$. Let $f : X \rightarrow S^2$ be a nontrivial genus- g Lefschetz fibration with $b_2^+(X) = 1$. Note that $-4(g - 1) \leq K^2$, and so $4(g - 1) \leq K_f^2$ (see [Stipsicz 1999, Lemma 3.2]). Suppose that X is not diffeomorphic to the blow-up of a ruled surface.

First, suppose that $b_1 = 0$. Since $b_2^+ = 1$ and $\chi_f = (\sigma + e)/4 + (g - 1) = (b_2^+ - b_1 + 1)/2 + (g - 1) = g$, we have $4(g - 1)/g \leq K_f^2/\chi_f = \lambda_f$. On the other hand, since $K^2 = 3\sigma + 2e = 5b_2^+ - b_2^- + 4 - 4b_1 = 9 - b_2^-$, by $b_2^- \geq 1$, we have $\lambda_f = K_f^2/\chi_f = \{9 - b_2^- + 8(g - 1)\}/g \leq 8$.

Next, suppose $b_1 = 2$. Then, $\chi_f = g - 1$. Therefore, by $4(g - 1) \leq K_f^2$, we have $4 \leq \lambda_f$. Since $0 \leq e = 2 - 2b_1 + b_2^+ + b_2^- = 2 - 4 + 1 + b_2^- = -1 + b_2^-$, we obtain $\lambda_f = \{1 - b_2^- + 8(g - 1)\}/(g - 1) \leq 8$.

Finally, suppose that X is the m -fold blow-up of an S^2 -bundle over Σ_k . Let Y be the S^2 -bundle over Σ_k . Then, since $b_1(Y) = 2k$, $b_2^\pm(Y) = 1$ and $X = Y \# m\mathbb{C}\mathbb{P}^2$, we have $b_1(X) = 2k$, $b_2^+(X) = 1$, $b_2^-(X) = m + 1$, $e(X) = 4 - 4k + m$ and $\sigma(X) = -m$. Hence, we have $\lambda_f = 8 - m/(g - k)$. From $m \geq 0$, $\lambda_f \leq 8$. We will give lower bounds for λ_f . By Lemma 3.2 in [Stipsicz 2002], $4(2k - g) + m \leq 4$. We have $\lambda_f \geq 4 + 4(k - 1)/(g - k)$ from $\lambda_f = 8 - m/(g - k)$, $4(2k - g) + m \leq 4$ and $0 \leq k \leq g/2$ [Li 2000, Proposition 4.4]. Fintushel and Stern [2004] showed that $(\Sigma_k \times S^2) \# 4m\mathbb{C}\mathbb{P}^2$ admits a genus- $(2k + m - 1)$ Lefschetz fibration f_{FS} over S^2 . When $m = g - 2k + 1$, we find $b_2^+ = 1$ and that $\lambda_{f_{FS}} = 4 + 4(k - 1)/(g - k)$. \square

Remark 5.2. If two Lefschetz fibrations f_1 and f_2 satisfy $\lambda_{f_1}, \lambda_{f_2} \geq 4 - 4/g$, then the (twisted) fiber sum f_3 of f_1 and f_2 satisfies $\lambda_{f_3} \geq 4 - 4/g$.

Acknowledgments

The author would like to thank Hisaaki Endo for pointing out that the author's idea is effective in constructing a Lefschetz fibration which violates the slope inequality and for his encouragement. The author would like to express his gratitude to Kazuhiro Konno for his kind explanation of the history of the slope, and Mustafa Korkmaz and Kouichi Yasui for their comments on this paper. Finally, the referee's comments were especially helpful for improving the exposition.

References

- [Amorós et al. 2000] J. Amorós, F. Bogomolov, L. Katzarkov, and T. Pantev, “Symplectic Lefschetz fibrations with arbitrary fundamental groups”, *J. Differential Geom.* **54**:3 (2000), 489–545. [MR 2002g:57051](#) [Zbl 1031.57021](#)
- [Arakelov 1971] S. J. Arakelov, “Families of algebraic curves with fixed degeneracies”, *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 1269–1293. In Russian; translated in *Math. USSR Izv.* **5** (1971), 1277–1302. [MR 48 #298](#) [Zbl 0238.14012](#)
- [Barth et al. 1984] W. Barth, C. Peters, and A. Van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **4**, Springer, Berlin, 1984. [MR 86c:32026](#) [Zbl 0718.14023](#)
- [Beauville 1979] A. Beauville, “L’application canonique pour les surfaces de type général”, *Invent. Math.* **55**:2 (1979), 121–140. [MR 81m:14025](#) [Zbl 0403.14006](#)
- [Catanese 1983] F. Catanese, “Moduli of surfaces of general type”, pp. 90–112 in *Algebraic geometry: open problems* (Ravello, 1982), edited by C. Ciliberto et al., Lecture Notes in Math. **997**, Springer, Berlin, 1983. [MR 84k:14026](#) [Zbl 0517.14011](#)
- [Chen 1993] Z. J. Chen, “On the lower bound of the slope of a nonhyperelliptic fibration of genus 4”, *Internat. J. Math.* **4**:3 (1993), 367–378. [MR 94g:14012](#) [Zbl 0816.14006](#)
- [Dehn 1938] M. Dehn, “Die Gruppe der Abbildungsklassen”, *Acta Math.* **69**:1 (1938), 135–206. [MR 1555438](#) [Zbl 0019.25301](#)
- [Donaldson 1999] S. K. Donaldson, “Lefschetz pencils on symplectic manifolds”, *J. Differential Geom.* **53**:2 (1999), 205–236. [MR 2002g:53154](#) [Zbl 1040.53094](#)
- [Endo 2000] H. Endo, “Meyer’s signature cocycle and hyperelliptic fibrations”, *Math. Ann.* **316**:2 (2000), 237–257. [MR 2001b:57047](#) [Zbl 0948.57013](#)
- [Endo and Gurtas 2010] H. Endo and Y. Z. Gurtas, “Lantern relations and rational blowdowns”, *Proc. Amer. Math. Soc.* **138**:3 (2010), 1131–1142. [MR 2011a:57047](#) [Zbl 1198.57017](#)
- [Endo and Nagami 2005] H. Endo and S. Nagami, “Signature of relations in mapping class groups and non-holomorphic Lefschetz fibrations”, *Trans. Amer. Math. Soc.* **357**:8 (2005), 3179–3199. [MR 2006g:57051](#) [Zbl 1081.57023](#)
- [Endo et al. 2011] H. Endo, T. E. Mark, and J. Van Horn-Morris, “Monodromy substitutions and rational blowdowns”, *J. Topol.* **4**:1 (2011), 227–253. [MR 2012b:57051](#) [Zbl 1234.57037](#)
- [Fintushel and Stern 1997] R. Fintushel and R. J. Stern, “Rational blowdowns of smooth 4-manifolds”, *J. Differential Geom.* **46**:2 (1997), 181–235. [MR 98j:57047](#) [Zbl 0896.57022](#)
- [Fintushel and Stern 1998] R. Fintushel and R. J. Stern, “Constructions of smooth 4-manifolds”, pp. 443–452 in *Proceedings of the International Congress of Mathematicians, II* (Berlin, 1998), edited by G. Fischer and U. Rehmann, 1998. [MR 99g:57033](#) [Zbl 0899.57012](#)

- [Fintushel and Stern 2004] R. Fintushel and R. J. Stern, “Families of simply connected 4-manifolds with the same Seiberg–Witten invariants”, *Topology* **43**:6 (2004), 1449–1467. MR 2005d:57044 Zbl 1064.57036
- [Gompf and Stipsicz 1999] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics **20**, Amer. Math. Soc., Providence, RI, 1999. MR 2000h:57038 Zbl 0933.57020
- [Hirose 2010] S. Hirose, “Presentations of periodic maps on oriented closed surfaces of genera up to 4”, *Osaka J. Math.* **47**:2 (2010), 385–421. MR 2011h:57025 Zbl 1202.57022
- [Johnson 1979] D. L. Johnson, “Homeomorphisms of a surface which act trivially on homology”, *Proc. Amer. Math. Soc.* **75**:1 (1979), 119–125. MR 80h:57008 Zbl 0407.57003
- [Kas 1980] A. Kas, “On the handlebody decomposition associated to a Lefschetz fibration”, *Pacific J. Math.* **89**:1 (1980), 89–104. MR 82f:57012 Zbl 0457.14011
- [Konno 1991] K. Konno, “A note on surfaces with pencils of nonhyperelliptic curves of genus 3”, *Osaka J. Math.* **28**:3 (1991), 737–745. MR 93c:14035 Zbl 0766.14037
- [Konno 1993] K. Konno, “Nonhyperelliptic fibrations of small genus and certain irregular canonical surfaces”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **20**:4 (1993), 575–595. MR 95b:14026 Zbl 0822.14009
- [Konno 1996] K. Konno, “Even canonical surfaces with small K^2 , III”, *Nagoya Math. J.* **143** (1996), 1–11. MR 98a:14051 Zbl 0872.14033
- [Korkmaz 2001] M. Korkmaz, “Noncomplex smooth 4-manifolds with Lefschetz fibrations”, *Int. Math. Res. Not.* **2001**:3 (2001), 115–128. MR 2001m:57036 Zbl 0977.57020
- [Li 2000] T.-J. Li, “Symplectic Parshin–Arakelov inequality”, *Int. Math. Res. Not.* **2000**:18 (2000), 941–954. MR 2001i:57038 Zbl 0961.57022
- [Manetti 2003] M. Manetti, “Surfaces of Albanese general type and the Severi conjecture”, *Math. Nachr.* **261–262**:1 (2003), 105–122. MR 2004m:14081 Zbl 1044.14017
- [Matsumoto 1983] Y. Matsumoto, “On 4-manifolds fibered by tori, II”, *Proc. Japan Acad. Ser. A Math. Sci.* **59**:3 (1983), 100–103. MR 84j:57010b Zbl 0532.55020
- [Matsumoto 1996] Y. Matsumoto, “Lefschetz fibrations of genus two: a topological approach”, pp. 123–148 in *Topology and Teichmüller spaces* (Katinkulta, 1995), edited by S. Kojima et al., World Scientific, 1996. MR 2000h:14038 Zbl 0921.57006
- [Miyachi and Shiga 2011] H. Miyachi and H. Shiga, “Holonomies and the slope inequality of Lefschetz fibrations”, *Proc. Amer. Math. Soc.* **139**:4 (2011), 1299–1307. MR 2012a:57036 Zbl 1226.14015
- [Ozbagci 2002] B. Ozbagci, “Signatures of Lefschetz fibrations”, *Pacific J. Math.* **202**:1 (2002), 99–118. MR 2002k:57066 Zbl 1053.57013
- [Ozbagci and Stipsicz 2000] B. Ozbagci and A. I. Stipsicz, “Noncomplex smooth 4-manifolds with genus-2 Lefschetz fibrations”, *Proc. Amer. Math. Soc.* **128**:10 (2000), 3125–3128. MR 2000m:57036 Zbl 0951.57015
- [Pardini 2005] R. Pardini, “The Severi inequality $K^2 \geq 4\chi$ for surfaces of maximal Albanese dimension”, *Invent. Math.* **159**:3 (2005), 669–672. MR 2006a:14065 Zbl 1082.14041
- [Reid 1979] M. Reid, “ π_1 for surfaces with small K^2 ”, pp. 534–544 in *Algebraic geometry* (Copenhagen, 1978), edited by K. Lønsted, Lecture Notes in Math. **732**, Springer, Berlin, 1979. MR 81j:14018 Zbl 0423.14021
- [Severi 1932] F. Severi, “La serie canonica e la teoria delle serie principali di gruppi di punti sopra una superficie algebrica”, *Comment. Math. Helv.* **4**:1 (1932), 268–326. MR 1509461 Zbl 0005.17602

- [Smith 1999] I. Smith, “Lefschetz fibrations and the Hodge bundle”, *Geom. Topol.* **3** (1999), 211–233. [MR 2000j:57059](#) [Zbl 0929.53047](#)
- [Smith 2001] I. Smith, “Lefschetz pencils and divisors in moduli space”, *Geom. Topol.* **5** (2001), 579–608. [MR 2002f:57056](#) [Zbl 1066.57030](#)
- [Stipsicz 1999] A. I. Stipsicz, “On the number of vanishing cycles in Lefschetz fibrations”, *Math. Res. Lett.* **6**:3–4 (1999), 449–456. [MR 2000g:57046](#) [Zbl 0955.57026](#)
- [Stipsicz 2002] A. I. Stipsicz, “Singular fibers in Lefschetz fibrations on manifolds with $b_2^+ = 1$ ”, *Topology Appl.* **117**:1 (2002), 9–21. [MR 2002j:57048](#) [Zbl 1007.53059](#)
- [Usher 2006] M. Usher, “Minimality and symplectic sums”, *Int. Math. Res. Not.* **2006**:16 (2006), Art. ID 49857. [MR 2007h:53139](#) [Zbl 1110.57017](#)
- [Xiao 1987] G. Xiao, “Fibered algebraic surfaces with low slope”, *Math. Ann.* **276**:3 (1987), 449–466. [MR 88a:14046](#) [Zbl 0596.14028](#)

Received November 1, 2012. Revised May 19, 2013.

NAOYUKI MONDEN
DEPARTMENT OF MATHEMATICS
KYOTO UNIVERSITY
GRADUATE SCHOOL OF SCIENCE
KYOTO
JAPAN
n-monden@math.kyoto-u.ac.jp

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 267 No. 1 January 2014

Numerical study of unbounded capillary surfaces	1
YASUNORI AOKI and HANS DE STERCK	
Dual R -groups of the inner forms of $SL(N)$	35
KUOK FAI CHAO and WEN-WEI LI	
Automorphisms and quotients of quaternionic fake quadrics	91
AMIR DŽAMBIĆ and XAVIER ROULLEAU	
Distance of bridge surfaces for links with essential meridional spheres	121
YEONHEE JANG	
Normal states of type III factors	131
YASUYUKI KAWAHIGASHI, YOSHIKO OGATA and ERLING STØRMER	
Eigenvalues and entropies under the harmonic-Ricci flow	141
YI LI	
Quantum extremal loop weight modules and monomial crystals	185
MATHIEU MANSUY	
Lefschetz fibrations with small slope	243
NAOYUKI MONDEN	