

*Pacific
Journal of
Mathematics*

**A NEW MONOTONE QUANTITY
ALONG THE INVERSE MEAN CURVATURE FLOW IN \mathbb{R}^n**

KWOK-KUN KWONG AND PENGZI MIAO

A NEW MONOTONE QUANTITY ALONG THE INVERSE MEAN CURVATURE FLOW IN \mathbb{R}^n

KWOK-KUN KWONG AND PENGZI MIAO

We find a new monotone increasing quantity along smooth solutions to the inverse mean curvature flow in \mathbb{R}^n . As an application, we derive a sharp geometric inequality for mean convex, star-shaped hypersurfaces which relates the volume enclosed by a hypersurface to a weighted total mean curvature of the hypersurface.

1. Statement of the result

Monotone quantities along hypersurfaces evolving under the inverse mean flow have many applications in geometry and relativity. Huisken and Ilmanen [2001] applied the monotone increasing property of Hawking mass to give a proof of the Riemannian Penrose inequality. Brendle, Hung and Wang [Brendle et al. 2012] discovered a monotone decreasing quantity along the inverse mean curvature flow in anti-de Sitter–Schwarzschild manifolds and used it to establish a Minkowski-type inequality for star-shaped hypersurfaces.

In this note, we provide a new monotone increasing quantity along smooth solutions to the inverse mean curvature flow in \mathbb{R}^n :

Theorem 1. *Let Σ be a smooth, closed, embedded hypersurface with positive mean curvature in \mathbb{R}^n . Let I be an open interval and $X : \Sigma \times I \rightarrow \mathbb{R}^n$ be a smooth map satisfying*

$$(1-1) \quad \frac{\partial X}{\partial t} = \frac{1}{H} \nu,$$

where H is the mean curvature of the surface $\Sigma_t = X(\Sigma, t)$ and ν is the outward unit normal vector to Σ_t . Let Ω_t be the bounded region enclosed by Σ_t and $r = r(x)$ be the distance from x to a fixed point O . Then the function

$$(1-2) \quad Q(t) = e^{-\frac{n-2}{n-1}t} \left(n \operatorname{Vol}(\Omega_t) - \frac{1}{n-1} \int_{\Sigma_t} r^2 H \, d\mu \right)$$

MSC2010: primary 53C44; secondary 53A07.

Keywords: inverse mean curvature flow.

is monotone increasing and $Q(t)$ is a constant function if and only if Σ_t is a round sphere for each t . Here $\text{Vol}(\Omega)$ denotes the volume of a bounded region Ω and $d\mu$ denotes the volume form on a hypersurface.

As an application, we derive a sharp inequality for star-shaped hypersurfaces in \mathbb{R}^n which relates the volume enclosed by a hypersurface to an r^2 -weighted total mean curvature of the hypersurface.

Theorem 2. *Let Σ be a smooth, star-shaped hypersurface with positive mean curvature in \mathbb{R}^n . Then*

$$(1-3) \quad n \text{Vol}(\Omega) \leq \frac{1}{n-1} \int_{\Sigma} r^2 H \, d\mu$$

where $\text{Vol}(\Omega)$ is the volume of the region Ω enclosed by Σ , r is the distance to a fixed point O and H is the mean curvature of Σ . Furthermore, equality in (1-3) holds if and only if Σ is a sphere centered at O .

We give some remarks about [Theorem 1](#) and [Theorem 2](#). The discovery of the monotonicity of $Q(t)$ in [Theorem 1](#) is motivated by [\[Brendle et al. 2012, Section 5\]](#). To prove [Theorem 1](#), we also need a result of Ros, proved using Reilly’s formula. Once we know that $Q(t)$ is monotone increasing, to prove [Theorem 2](#), it may be tempting to ask whether $\lim_{t \rightarrow \infty} Q(t) = 0$? We do not know if this is true because both $\text{Vol}(\Omega_t)$ and $\int_{\Sigma_t} r^2 H \, d\mu$ grow like $\exp(\frac{n}{n-1} t)$ when $\{\Sigma_t\}$ are spheres, while there is only a factor of $\exp(-\frac{n-2}{n-1} t)$ in (1-2). Instead, we take an alternate approach by first proving [Theorem 2](#) for a convex hypersurface Σ . The proof in that case again makes use of Reilly’s formula. When Σ is merely assumed to be mean convex and star-shaped, we prove [Theorem 2](#) by reducing it to the convex case using solutions to the inverse mean curvature flow provided by [\[Gerhardt 1990\]](#) and [\[Urbas 1990\]](#).

2. Proof of the theorems

Given a compact Riemannian manifold (Ω, g) with boundary Σ , Reilly’s formula [\[1977\]](#) asserts that

$$(2-1) \quad \int_{\Omega} |\nabla^2 u|^2 + \langle \nabla(\Delta u), \nabla u \rangle + \text{Ric}(\nabla u, \nabla u) \, dV \\ = \int_{\Sigma} (\Delta u) \frac{\partial u}{\partial \nu} - II(\nabla^{\Sigma} u, \nabla^{\Sigma} u) - 2(\Delta_{\Sigma} u) \frac{\partial u}{\partial \nu} - H \left(\frac{\partial u}{\partial \nu} \right)^2 \, d\mu.$$

Here u is a smooth function on Ω ; ∇^2 , Δ and ∇ denote the Hessian, the Laplacian and the gradient on Ω ; Δ_{Σ} and ∇^{Σ} denote the Laplacian and the gradient on Σ ; ν is the unit outward normal vector to Σ ; II and H are the second fundamental

form and the mean curvature of Σ with respect to ν ; and Ric is the Ricci curvature of g .

To prove [Theorem 1](#), we need a result of [\[Ros 1987\]](#), which was proved by choosing $\Delta u = 1$ on Ω and $u = 0$ at Σ in the above Reilly’s formula.

Theorem 3 [\[Ros 1987\]](#). *Let (Ω, g) be an n -dimensional compact Riemannian manifold with nonnegative Ricci curvature with boundary Σ . Suppose Σ has positive mean curvature H ; then*

$$(2-2) \quad n \operatorname{Vol}(\Omega) \leq (n - 1) \int_{\Sigma} \frac{1}{H} d\mu$$

and equality holds if and only if (Ω, g) is isometric to a round ball in \mathbb{R}^n .

Proof of Theorem 1. We use $'$ to denote differentiation with respect to t . Some basic formulas along the inverse mean curvature flow (1-1) in \mathbb{R}^n are

$$(2-3) \quad H' = -\Delta_{\Sigma_t} \left(\frac{1}{H} \right) - \frac{|II|^2}{H}, \quad d\mu' = d\mu, \quad \operatorname{Vol}(\Omega_t)' = \int_{\Sigma_t} \frac{1}{H} d\mu.$$

Let $u = r^2$. Then u satisfies

$$(2-4) \quad \nabla^2 u = 2g \quad \text{and} \quad \Delta u = 2n,$$

where g is the Euclidean metric. Now

$$(2-5) \quad \left(\int_{\Sigma_t} uH d\mu \right)' = \int_{\Sigma_t} (u'H + uH' + uH) d\mu.$$

Let $\langle \cdot, \cdot \rangle$ be the Euclidean inner product. By (2-3), (2-4) and the divergence theorem, we have

$$(2-6) \quad \int_{\Sigma_t} u'H d\mu = \int_{\Sigma_t} \left\langle \nabla u, \frac{1}{H} \nu \right\rangle H d\mu = \int_{\Omega_t} \Delta u dV = 2n \operatorname{Vol}(\Omega_t).$$

By (2-4), we also have

$$\Delta_{\Sigma_t} u = \Delta u - H \frac{\partial u}{\partial \nu} - \nabla^2 u(\nu, \nu) = 2(n - 1) - H \frac{\partial u}{\partial \nu},$$

which together with (2-3) and (2-4) implies

$$(2-7) \quad \begin{aligned} \int_{\Sigma_t} uH' d\mu &= \int_{\Sigma_t} \left(-\frac{\Delta_{\Sigma_t} u}{H} - \frac{u|II|^2}{H} \right) d\mu \\ &= \int_{\Sigma_t} \left(-\frac{2(n-1)}{H} + \frac{\partial u}{\partial \nu} - \frac{u|II|^2}{H} \right) d\mu \\ &= -\int_{\Sigma_t} \frac{2(n-1)}{H} d\mu + 2n \operatorname{Vol}(\Omega_t) - \int_{\Sigma_t} \frac{u|II|^2}{H} d\mu. \end{aligned}$$

Substituting (2-6) and (2-7) into (2-5) yields

$$\begin{aligned}
 (2-8) \quad \left(\int_{\Sigma_t} uH \, d\mu \right)' &= 4n \operatorname{Vol}(\Omega_t) + \int_{\Sigma_t} \left(-\frac{2(n-1)}{H} - \frac{u|II|^2}{H} + uH \right) d\mu \\
 &\leq 4n \operatorname{Vol}(\Omega_t) + \int_{\Sigma_t} \left(-\frac{2(n-1)}{H} - \frac{uH}{n-1} + uH \right) d\mu \\
 &= 4n \operatorname{Vol}(\Omega_t) + \int_{\Sigma_t} \left(-\frac{2(n-1)}{H} + \frac{n-2}{n-1} uH \right) d\mu \\
 &\leq 4n \operatorname{Vol}(\Omega_t) - 2n \operatorname{Vol}(\Omega_t) + \frac{n-2}{n-1} \int_{\Sigma_t} uH \, d\mu \\
 &= 2n \operatorname{Vol}(\Omega_t) + \frac{n-2}{n-1} \int_{\Sigma_t} uH \, d\mu,
 \end{aligned}$$

where we have used $|II|^2 \geq \frac{1}{n-1} H^2$ in the second line and [Theorem 3](#) in the fourth. On the other hand, by [Theorem 3](#) again, we have

$$(2-9) \quad \operatorname{Vol}(\Omega_t)' = \int_{\Sigma_t} \frac{1}{H} \, d\mu \geq \frac{n}{n-1} \operatorname{Vol}(\Omega_t).$$

It follows from (2-8) and (2-9) that

$$\left(n(n-1) \operatorname{Vol}(\Omega_t) - \int_{\Sigma_t} uH \, d\mu \right)' \geq \frac{n-2}{n-1} \left(n(n-1) \operatorname{Vol}(\Omega_t) - \int_{\Sigma_t} uH \, d\mu \right)$$

or equivalently

$$(2-10) \quad \left[e^{-\frac{n-2}{n-1}t} \left(n \operatorname{Vol}(\Omega_t) - \frac{1}{n-1} \int_{\Sigma_t} r^2 H \, d\mu \right) \right]' \geq 0.$$

We conclude that $Q(t)$ is monotone increasing, moreover $Q(t)$ is a constant function if and only if equalities in (2-8) and (2-9) hold. By [Theorem 3](#), we know these equalities hold if and only if Σ_t is a round sphere for all t . This completes the proof of [Theorem 1](#). \square

Next, we prove [Theorem 2](#) in the case that Σ is a convex hypersurface.

Proposition 1. *Let Σ be a smooth, closed, convex hypersurface in \mathbb{R}^n . Then*

$$(2-11) \quad n \operatorname{Vol}(\Omega) \leq \frac{1}{n-1} \int_{\Sigma} r^2 H \, d\mu,$$

where $\operatorname{Vol}(\Omega)$ is the volume of the region Ω enclosed by Σ , r is the distance to a fixed point O and H is the mean curvature of Σ . Moreover, equality in (2-11) holds if and only if Σ is a sphere centered at O .

Remark. [Proposition 1](#) generalizes the first inequality in [Theorem 3.2\(1\)](#) of [[Kwong 2012](#)].

Proof. Apply Reilly’s formula (2-1) to the Euclidean region Ω and choose $u = r^2$; we have

$$4n(n-1) \text{Vol}(\Omega) = \int_{\Sigma} II(\nabla^{\Sigma}u, \nabla^{\Sigma}u) + 2(\Delta_{\Sigma}u) \frac{\partial u}{\partial \nu} + H \left(\frac{\partial u}{\partial \nu} \right)^2 d\mu$$

where $\Delta_{\Sigma}u = \Delta u - H \frac{\partial u}{\partial \nu} - \nabla^2 u(v, v) = 2(n-1) - H \frac{\partial u}{\partial \nu}$. Therefore,

$$(2-12) \quad \int_{\Sigma} H \left(\frac{\partial u}{\partial \nu} \right)^2 d\mu = \int_{\Sigma} II(\nabla^{\Sigma}u, \nabla^{\Sigma}u) d\mu + 4n(n-1) \text{Vol}(\Omega).$$

Since Σ is convex, $II(\cdot, \cdot)$ is positive definite. Hence, (2-12) implies

$$(2-13) \quad n(n-1) \text{Vol}(\Omega) \leq \frac{1}{4} \int_{\Sigma} H \langle \nabla(r^2), \nu \rangle^2 d\mu \leq \int_{\Sigma} H r^2 d\mu.$$

When $n(n-1) \text{Vol}(\Omega) = \int_{\Sigma} H r^2 d\mu$, we must have $II(\nabla^{\Sigma}u, \nabla^{\Sigma}u) = 0$, hence $\nabla^{\Sigma}u = 0$. This implies that $u = r^2$ is a constant on Σ , which shows that Σ is a sphere centered at O . \square

To deform a star-shaped hypersurface to a convex hypersurface through the inverse mean curvature flow, we make use of a special case of a general result of Gerhardt and Urbas:

Theorem 4 [Gerhardt 1990; Urbas 1990]. *Let Σ be a smooth, closed hypersurface in \mathbb{R}^n with positive mean curvature, given by a smooth embedding $X_0 : \mathbb{S}^{n-1} \rightarrow \mathbb{R}^n$. Suppose Σ is star-shaped with respect to a point P . Then the initial value problem*

$$(2-14) \quad \begin{cases} \frac{\partial X}{\partial t} = \frac{1}{H} \nu, \\ X(\cdot, 0) = X_0(\cdot), \end{cases}$$

has a unique smooth solution $X : \mathbb{S}^{n-1} \times [0, \infty) \rightarrow \mathbb{R}^n$, where ν is the unit outer normal vector to $\Sigma_t = X(\mathbb{S}^{n-1}, t)$ and H is the mean curvature of Σ_t . Moreover, Σ_t is star-shaped with respect to P and the rescaled hypersurface $\tilde{\Sigma}_t$, parametrized by $\tilde{X}(\cdot, t) = e^{-t/(n-1)} X(\cdot, t)$, converges to a sphere centered at P in the \mathcal{C}^{∞} topology as $t \rightarrow \infty$.

Proof of Theorem 2. By Theorem 4, there exists a smooth solution $\{\Sigma_t\}$ to the inverse mean curvature flow with initial condition Σ . Moreover, the rescaled hypersurface $\tilde{\Sigma}_t = \{e^{-t/(n-1)}x \mid x \in \Sigma_t\}$ converges exponentially fast in the C^{∞} topology to a sphere. In particular, $\tilde{\Sigma}_t$ and hence Σ_t , must be convex for large t .

Let T be a time when Σ_T becomes convex. By Proposition 1, we have

$$n \text{Vol}(\Omega_T) \leq \frac{1}{n-1} \int_{\Sigma_T} r^2 H d\mu;$$

thus $Q(T) \leq 0$. By [Theorem 1](#), we know that $Q(t)$ is monotone increasing. Hence $Q(0) \leq Q(T) \leq 0$, which proves [\(1-3\)](#).

If the equality in [\(1-3\)](#) holds, then $Q(0) = 0$. It follows from the monotonicity of $Q(t)$ and the fact $Q(t) \leq 0$ for large t that $Q(t) = 0$ for all t . By [Theorem 1](#), this implies that Σ_t is a sphere for each t . By [Theorem 1](#), Σ_t is a sphere centered at O for large t . Therefore, we conclude that the initial hypersurface Σ is a sphere centered at O . \square

Remark. It can be shown that [Theorem 2](#) is still true if the mean curvature is only assumed to be nonnegative. Please refer to the arXiv version of this paper ([1212.1906](#)) for details.

References

- [Brendle et al. 2012] S. Brendle, P.-K. Hung, and M.-T. Wang, “A Minkowski-type inequality for hypersurfaces in the Anti-deSitter–Schwarzschild manifold”, preprint, 2012. [arXiv 1209.0669](#)
- [Gerhardt 1990] C. Gerhardt, “Flow of nonconvex hypersurfaces into spheres”, *J. Differential Geom.* **32**:1 (1990), 299–314. [MR 91k:53016](#) [Zbl 0708.53045](#)
- [Huisken and Ilmanen 2001] G. Huisken and T. Ilmanen, “The inverse mean curvature flow and the Riemannian Penrose inequality”, *J. Differential Geom.* **59**:3 (2001), 353–437. [MR 2003h:53091](#) [Zbl 1055.53052](#)
- [Kwong 2012] K.-K. Kwong, “On convex hypersurfaces in space forms and eigenvalues estimates for differential forms”, preprint, 2012. [arXiv 1207.3999](#)
- [Reilly 1977] R. C. Reilly, “Applications of the Hessian operator in a Riemannian manifold”, *Indiana Univ. Math. J.* **26**:3 (1977), 459–472. [MR 57 #13799](#) [Zbl 0391.53019](#)
- [Ros 1987] A. Ros, “Compact hypersurfaces with constant higher order mean curvatures”, *Rev. Mat. Iberoamericana* **3**:3-4 (1987), 447–453. [MR 90c:53160](#) [Zbl 0673.53003](#)
- [Urbas 1990] J. I. E. Urbas, “On the expansion of starshaped hypersurfaces by symmetric functions of their principal curvatures”, *Math. Z.* **205**:3 (1990), 355–372. [MR 92c:53037](#) [Zbl 0691.35048](#)

Received November 12, 2012.

KWOK-KUN KWONG
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MIAMI
CORAL GABLES, FL 33146
UNITED STATES

kwong@math.miami.edu

PENGZI MIAO
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MIAMI
CORAL GABLES, FL 33146
UNITED STATES

pengzim@math.miami.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 267 No. 2 February 2014

Sums of squares in algebraic function fields over a complete discretely valued field	257
KARIM JOHANNES BECHER, DAVID GRIMM and JAN VAN GEEL	
On the equivalence problem for toric contact structures on S^3 -bundles over S^2	277
CHARLES P. BOYER and JUSTIN PATI	
An almost-Schur type lemma for symmetric $(2,0)$ tensors and applications	325
XU CHENG	
Algebraic invariants, mutation, and commensurability of link complements	341
ERIC CHESEBRO and JASON DEBLOIS	
Taut foliations and the action of the fundamental group on leaf spaces and universal circles	399
YOSUKE KANO	
A new monotone quantity along the inverse mean curvature flow in \mathbb{R}^n	417
KWOK-KUN KWONG and PENGZI MIAO	
Nonfibered L-space knots	423
TYE LIDMAN and LIAM WATSON	
Families and Springer's correspondence	431
GEORGE LUSZTIG	
Reflexive operator algebras on Banach spaces	451
FLORENCE MERLEVÈDE, COSTEL PELIGRAD and MAGDA PELIGRAD	
Harer stability and orbifold cohomology	465
NICOLA PAGANI	
Spectra of product graphs and permanents of matrices over finite rings	479
LE ANH VINH	
The concavity of the Gaussian curvature of the convex level sets of minimal surfaces with respect to the height	489
PEI-HE WANG	