

*Pacific  
Journal of  
Mathematics*

**NONFIBERED L-SPACE KNOTS**

TYE LIDMAN AND LIAM WATSON

Volume 267 No. 2

February 2014



## NONFIBERED L-SPACE KNOTS

TYE LIDMAN AND LIAM WATSON

**We construct an infinite family of knots in rational homology spheres with irreducible, nonfibered complements, for which every nonlongitudinal filling is an L-space.**

The Heegaard Floer homology of a rational homology three-sphere  $Y$  is an abelian group  $\widehat{HF}(Y)$  satisfying  $\text{rk } \widehat{HF}(Y) \geq |H_1(Y; \mathbb{Z})|$  [Ozsváth and Szabó 2004]. When equality is realized in this bound  $Y$  is called an L-space and any knot in  $Y$  admitting a nontrivial L-space surgery is called an L-space knot [Ozsváth and Szabó 2005]. A result of Ghiggini [2008] and Ni [2007] shows that L-space knots in the three-sphere must be fibered. Since manifolds with finite fundamental group provide examples of L-spaces,<sup>1</sup> this result implies that a knot  $K$  in  $S^3$  admitting a finite filling must be fibered. This observation should be compared with other restrictions related to finite fillings such as the Cyclic Surgery Theorem [Culler et al. 1987] and its extensions [Boyer and Zhang 2001].

The restriction to knots in  $S^3$  is not necessary. It is shown in [Boileau et al. 2012] that a primitive knot<sup>2</sup> in an irreducible L-space admitting a nontrivial L-space surgery must be fibered. Irreducibility of the complement is required: removing an unknot from an embedded three-ball in any L-space produces a nonfibered manifold with nontrivial L-space fillings. Even in the general setting of knots in rational homology spheres with irreducible complements fibered is not a necessary condition:

**Theorem 1.** *There exist infinitely many irreducible, nonfibered knot complements such that all nonlongitudinal Dehn fillings are L-spaces. Moreover, these examples arise as knots in manifolds with finite fundamental group.*

---

Lidman was supported by a UCLA Dissertation Year Fellowship. Watson was partially supported by an NSERC Postdoctoral Fellowship.

MSC2010: 57M27.

*Keywords:* Heegaard Floer homology, L-space, fibration.

<sup>1</sup>Ozsváth and Szabó [2005] have shown that manifolds admitting elliptic geometry are L-spaces; Perelman's Geometrization Theorem (see [Kleiner and Lott 2008], for example) implies that three-manifolds with finite fundamental group admit elliptic geometry.

<sup>2</sup>Recall that a knot  $K$  is primitive in  $Y$  if  $[K] \in H_1(Y; \mathbb{Z})$  is a generator.

In particular, our examples are nonprimitive knots in L-spaces.

Before turning to the construction, we fix some terminology. Fibrations will always be locally trivial surface bundles over a circle and we say the total space fibers. To avoid confusion, we will refer to Seifert fibrations as Seifert structures; these are foliations of a manifold by circles. The base orbifold is the leaf space of such a foliation, where the (possibly empty) collection of cone points records the multiplicities of the exceptional fibers in the Seifert structure. A circle bundle is a Seifert structure for which there are no exceptional fibers.

Given a three-manifold  $M$  with torus boundary, a slope  $\alpha$  is a primitive class in  $H_1(\partial M; \mathbb{Z})/\{\pm 1\}$ . We use  $M(\alpha)$  to denote Dehn filling along  $\alpha$ . If  $\partial M = T_1 \cup T_2$ , for tori  $T_i$ , then we denote  $\alpha$ -filling on  $T_1$  (respectively  $T_2$ ) by  $M(\alpha, -)$  (respectively  $M(-, \alpha)$ ). When  $M$  admits a Seifert structure, the slope given by a regular fiber in the boundary is called the fiber slope. For background on Seifert structures and Dehn filling we refer the reader to [Boyer 2002]. A key fact is that Dehn filling a Seifert manifold with torus boundary along any slope  $\alpha$  other than the fiber slope results in a Seifert manifold with a possible additional singular fiber. The multiplicity of this new fiber is  $\Delta(\alpha, \phi)$ , the distance between the slopes  $\alpha$  and  $\phi$  [Heil 1974].

Finally, for knots in rational homology three-spheres recall that there is a preferred slope given by the rational longitude. This slope is characterized by the property that some number of like-oriented parallel copies in the boundary of the knot complement bounds a properly embedded surface. We will refer to this slope as the longitude. Note that an oriented three-manifold  $M$  with torus boundary for which  $H_1(M; \mathbb{Q}) \cong \mathbb{Q}$  always arises (nonuniquely) as the complement of a knot in a rational homology three-sphere.

### 1. The twisted $I$ -bundle over the Klein bottle

Let  $N$  denote the twisted  $I$ -bundle over the Klein bottle. As this orientable three-manifold with torus boundary plays a central role in our construction we will study its construction in depth.


First consider the group  $G$  generated by  $f, g: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , where

$$\begin{aligned} f(x, y, z) &= (x + 1, y, z), \\ g(x, y, z) &= (-x, y + 1, -z), \end{aligned}$$

and consider the noncompact, orientable three-manifold  $N^\circ = \mathbb{R}^3/G$ . Note that the  $z$ -component of  $\mathbb{R}^3$  gives  $N^\circ$  the structure of a line bundle, the zero-section of which is a Klein bottle; this is the unique line bundle over the Klein bottle with orientable total space. By restricting the action of  $G$  to  $\tilde{N} = \mathbb{R}^2 \times [-\frac{1}{2}, \frac{1}{2}] \subset \mathbb{R}^3$  we obtain the twisted  $I$ -bundle over the Klein bottle  $N = \tilde{N}/G$ .

From this description two Seifert structures on  $N$  become apparent: the  $x$ - and  $y$ -components of  $\tilde{N}$  both determine foliations of  $N$  by circles. (This is essentially the observation that the Klein bottle is foliated by circles in two ways.) The leaf space of the foliation described by the  $x$ -components is a Möbius strip without cone points. Denote a regular fiber in this Seifert structure by  $\phi_0$ . The base orbifold of the foliation determined by  $y$ -components is  $D^2(2, 2)$ , with regular fiber denoted  $\phi_1$ ; this follows readily from a natural Heegaard decomposition which we now describe.

From the preceding construction, a fundamental domain for  $N$  is obtained by taking  $[-\frac{1}{2}, \frac{1}{2}]^2 \times [-\frac{1}{2}, \frac{1}{2}] \subset \mathbb{R}^3$ . Then given a disk  $D^2$  of radius less than  $\frac{1}{2}$  (and centered at the origin of the  $xy$ -plane), the result of removing  $D^2 \times [-\frac{1}{2}, \frac{1}{2}]$  is a genus-two handlebody. This gives rise to a Heegaard decomposition for  $N$ ; a Heegaard diagram corresponding to this decomposition is described in Figure 1, from which the fundamental group  $\pi_1(N) = \langle a, b \mid a^2b^2 \rangle$  may be calculated. Note that since  $fgfg^{-1}$  is trivial in the group  $G$ , the homomorphism determined by  $a \mapsto fgfg^{-1}$  and  $b \mapsto fg^{-1}$  is well-defined and gives an isomorphism  $G \cong \langle a, b \mid a^2b^2 \rangle$ . Further, by considering a separating disk decomposing the handlebody into solid



**Figure 1.** Two views of the Heegaard diagram for the twisted  $I$ -bundle over the Klein bottle  $N$ . With  $a$  and  $b$  generating the fundamental group of the genus-two handlebody,  $N$  is obtained by attaching a handle along a curve in the boundary representing  $a^2b^2$  so that  $\phi_0 \simeq ab$  and  $\phi_1 \simeq b^2$ . On the left, an annulus in the boundary with core representing the element  $\phi_0 \simeq ab$  may be used to find the fundamental group of  $M$ , the complement of a regular fiber in the interior of  $N$ , via HNN extension. On the right, the axis of rotational symmetry shows that the hyperelliptic involution on the handlebody induces a strong inversion on the pair  $(N, K_0)$  where  $K_0$  is a knot in  $N$  isotopic to a regular fiber  $\phi_0$  in the interior of  $N$ .

tori, it is immediate that  $N$  is the union of two solid tori along essential annuli in the boundary. By fixing Seifert structures on each of these solid tori with base orbifolds  $D^2(2)$ , these annuli are foliated by regular fibers. The identification along these essential annuli therefore extends to a Seifert structure on  $N$  with base orbifold  $D^2(2, 2)$  as claimed.

Both Seifert structures induce foliations on the torus  $\partial N$ . Let  $\phi_0$  and  $\phi_1$  be regular fibers in  $\partial N$ , and notice that  $\Delta(\phi_0, \phi_1) = 1$ . (These conventions are consistent with [Boyer et al. 2013, Section 3].) The longitude of  $N$  is homotopic to the element  $ab$  (this element has order two in the abelianization of  $\pi_1(N)$ ). That is,  $\phi_0$  represents the longitude of  $N$ . Any filling  $N(\alpha)$  for which  $\alpha \neq \phi_0, \phi_1$  admits a pair of Seifert structures with base orbifolds  $\mathbb{R}P^2(\Delta(\alpha, \phi_0))$  and  $S^2(2, 2, \Delta(\alpha, \phi_1))$ . We point out that these manifolds always admit elliptic geometry [Scott 1983].


Now consider a knot  $K_0$  in  $N$  that is isotopic to a regular fiber  $\phi_0$  in the interior of  $N$ . Define  $M$  by removing a neighborhood of  $K_0$  from  $N$ ; by construction  $M$  inherits a Seifert structure (the base orbifold is a punctured Möbius band). Now  $\partial M = T_1 \cup T_2$  where  $T_2$  denotes the boundary of a regular neighborhood of  $K_0$ .

The fundamental group of  $M$  is presented by

$$\pi_1(M) = \langle a, b, t \mid a^2b^2, [t, ab] \rangle.$$

To see this, consult Figure 1 and notice that  $M$  may be constructed by identifying (disjoint neighborhoods of) each boundary component of the annulus with core  $ab$  in  $\partial N$ . This gives rise to the HNN extension presented above. Notice that  $M(-, \mu) \cong N$  for any slope on  $T_2$  satisfying  $\Delta(\mu, \phi_0) = 1$ . A preferred choice for  $\mu$  is given by a representative of the homotopy class of  $t$  in the above presentation.

A final observation pertains to a natural strong inversion on  $(N, K_0)$  that descends to an involution on  $M$  with one-dimensional fixed point set. Recall that a strong inversion on  $(N, K_0)$  is an orientation preserving involution on  $N$  that reverses orientation on  $K_0$ ; such a symmetry is illustrated in Figure 1. The involution on  $N$  is induced by the hyperelliptic involution on the genus-two handlebody since the attaching curve is fixed (as a set) by this involution. A fundamental domain for this involution is a three-ball, with one dimensional fixed point set. That is,  $N$  is



**Figure 2.** The branch set for the manifold  $M = M(-, -)$  with branch sets for the fillings  $M(\phi_1, -) = N$  and  $M(\phi_0, -)$ . Notice that  $M(\phi_0, -)$  is reducible, containing an  $S^2 \times S^1$  summand.

the twofold branched cover of a two-tangle; this is the leftmost tangle in Figure 2. We leave the following step to the reader: the genus-two handlebody is the twofold branched cover of a three-tangle, and attaching the handle closes one of the arcs (the arc meeting the attaching curve) to an unknotted curve in the branch set. The same construction may be applied to the complement of  $K_0$  in  $N$ , to see that  $M$  is the twofold branched cover of a tangle in  $S^2 \times I$ . This tangle is shown in Figure 2.

Towards a proof of Theorem 1, our interest is in the family of manifolds

$$\{M(-, \alpha) \mid \text{for any slope } \alpha \text{ with } \Delta(\alpha, \phi_0) > 1\}.$$

Notice that each manifold in this set admits a Seifert structure with base orbifold a Möbius band with a single cone point of order  $\Delta(\alpha, \phi_0)$ . Since  $M(\phi_1, \alpha)$  admits a Seifert structure with base orbifold  $S^2(2, 2, n)$  it follows that  $M(-, \alpha)$  is the complement of a knot in an elliptic manifold for all  $\alpha$ .

### 2. The proof of Theorem 1

Let  $M$  be the complement of  $K_0$  in the twisted  $I$ -bundle over the Klein bottle  $N$ . We assume all of the notation introduced in the previous section.

**Lemma 2.** *Fix a slope  $\alpha$  on  $T_2$  with  $\Delta(\alpha, \phi_0) = p$ . Then*

$$M(\phi_0, \alpha) = \begin{cases} S^2 \times S^1 \# S^2 \times S^1 & \text{if } p = 0, \\ S^2 \times S^1 \# L(p, q) & \text{if } p > 1, \\ S^2 \times S^1 & \text{if } p = 1. \end{cases}$$

*Proof.* Since  $\pi_1(M) \cong \langle a, b, t \mid a^2b^2, [t, ab] \rangle$  and  $\phi_0 \simeq ab$ , we have that

$$\pi_1(M(\phi_0, -)) \cong \langle a, b, t \mid a^2b^2, [t, ab] \rangle / \langle \langle ab \rangle \rangle \cong \langle a, b, t \mid ab \rangle.$$

In other words,  $\pi_1(M(\phi_0, -)) \cong \mathbb{Z} * \mathbb{Z}$ . If  $\alpha = p\mu + q\phi_0$ , then

$$\pi_1(M(\phi_0, \alpha)) \cong \langle a, b, t \mid ab \rangle / \langle \langle t^p(ab)^q \rangle \rangle \cong \mathbb{Z} * \mathbb{Z} / p.$$

By Whitehead’s proof of Kneser’s conjecture [Whitehead 1958],  $M(\phi_0, \alpha)$  is a connect-sum of closed manifolds  $Y_1$  and  $Y_2$  with  $\pi_1(Y_1) \cong \mathbb{Z}$  and  $\pi_1(Y_2) \cong \mathbb{Z}/p$ . Geometrization now establishes the lemma. □

**Remark 3.** Alternatively, Lemma 2 follows from considering  $M(\phi_0, -)$  as the double branched cover of a tangle as in Figure 2. The unknotted component gives rise to the  $S^2 \times S^1$  summand. Dehn filling corresponds to attaching a rational tangle, which (ignoring the unknotted component) produces a two-bridge link and exhibits the lens space connect-summand.

**Proposition 4.** *For any  $\alpha$  on  $T_2$  with  $\Delta(\alpha, \phi_0) > 1$ , the manifold  $M(-, \alpha)$  does not fiber over the circle.*

*Proof.* Suppose that  $M(-, \alpha)$  fibers. Since  $\phi_0$  is the longitude, this is the only filling that extends the fibration on  $M(-, \alpha)$  as any other filling of  $M(-, \alpha)$  results in a rational homology sphere. By Lemma 2,  $M(\phi_0, \alpha) \cong S^2 \times S^1 \# L(p, q)$  for  $p = \Delta(\phi_0, \alpha) \geq 2$ . Since  $M(\phi_0, \alpha)$  is fibered and  $\pi_2(M(\phi_0, \alpha)) \neq 0$ , the fiber surface  $F$  must also have  $\pi_2(F) \neq 0$  by the long exact sequence for a fibration. Hence  $F$  must be  $S^2$  or  $\mathbb{R}P^2$ . However,  $\pi_1(M(\phi_0, \alpha))$  is not the fundamental group of such a fibration, since it does not admit a surjective homomorphism onto  $\mathbb{Z}$  with finite kernel.  $\square$

*Proof of Theorem 1.* Fix  $\alpha$  with  $\Delta(\alpha, \phi_0) \geq 2$ . As the fiber slope of the Seifert structure on  $M(-, \alpha)$  is the longitude, all nonlongitudinal fillings will extend the Seifert structure, yielding a base orbifold  $\mathbb{R}P^2$  with two cone points. By [Boyer et al. 2013, Proposition 5], such manifolds are always L-spaces. Proposition 4 shows that  $M(-, \alpha)$  is not fibered. Furthermore,  $M(-, \alpha)$  is irreducible, since the only orientable, reducible Seifert manifolds are  $S^2 \times S^1$  and  $\mathbb{R}P^3 \# \mathbb{R}P^3$  (and in particular, are closed). Finally,  $M(-, \alpha)$  is the complement of a knot in an elliptic manifold as observed in Section 1.  $\square$

**Remark 5.** Further examples may be constructed in an analogous way by removing a regular fiber from any manifold which has a Seifert structure with base orbifold  $\mathbb{R}P^2$  with any positive number of singular fibers. It is also possible to construct examples, in a similar manner, admitting Sol geometry. The main observation is that every Sol rational homology sphere is an L-space [Boyer et al. 2013, Theorem 2]. Since every such L-space arises by identifying two twisted  $I$ -bundles along the boundary tori, one may consider the complement of the knot  $K_0$  in one of the twisted  $I$ -bundles. In this setting, our construction goes through almost verbatim, having noticed that the obvious essential torus must be horizontal to the purported fibration of the exterior of  $K_0$ .

**Question 6.** All of our examples relied on the presence of an essential annulus, and have nonhyperbolic exterior. Do there exist examples of hyperbolic, nonfibered knots for which every nonlongitudinal surgery is an L-space?

### Acknowledgments

The authors thank Ciprian Manolescu and Yi Ni for their comments on and interest in this problem. This paper owes its existence to the Workshop on Topics in Dehn Surgery, held at UT Austin in April 2012. The authors thank the organizers for putting together a great conference.

### References

- [Boileau et al. 2012] M. Boileau, S. Boyer, R. Cebanu, and G. S. Walsh, “Knot commensurability and the Berge conjecture”, *Geom. Topol.* **16**:2 (2012), 625–664. MR 2928979 Zbl 1258.57001



- [Boyer 2002] S. Boyer, “Dehn surgery on knots”, pp. 165–218 in *Handbook of geometric topology*, edited by R. J. Daverman and R. B. Sher, North-Holland, Amsterdam, 2002. MR 2003f:57030 Zbl 1058.57004
- [Boyer and Zhang 2001] S. Boyer and X. Zhang, “A proof of the finite filling conjecture”, *J. Differential Geom.* **59**:1 (2001), 87–176. MR 2003k:57007 Zbl 1030.57024
- [Boyer et al. 2013] S. Boyer, C. M. Gordon, and L. Watson, “On L-spaces and left-orderable fundamental groups”, *Math. Ann.* **356**:4 (2013), 1213–1245. MR 3072799 Zbl 06194415
- [Culler et al. 1987] M. Culler, C. M. Gordon, J. Luecke, and P. B. Shalen, “Dehn surgery on knots”, *Ann. of Math. (2)* **125**:2 (1987), 237–300. MR 88a:57026 Zbl 0633.57006
- [Ghiggini 2008] P. Ghiggini, “Knot Floer homology detects genus-one fibred knots”, *Amer. J. Math.* **130**:5 (2008), 1151–1169. MR 2010f:57013 Zbl 1149.57019
- [Heil 1974] W. Heil, “Elementary surgery on Seifert fiber spaces”, *Yokohama Math. J.* **22** (1974), 135–139. MR 51 #11515 Zbl 0297.57006
- [Kleiner and Lott 2008] B. Kleiner and J. Lott, “Notes on Perelman’s papers”, *Geom. Topol.* **12**:5 (2008), 2587–2855. MR 2010h:53098 Zbl 1204.53033
- [Ni 2007] Y. Ni, “Knot Floer homology detects fibred knots”, *Invent. Math.* **170**:3 (2007), 577–608. MR 2008j:57053 Zbl 1138.57031
- [Ozsváth and Szabó 2004] P. Ozsváth and Z. Szabó, “Holomorphic disks and topological invariants for closed three-manifolds”, *Ann. of Math. (2)* **159**:3 (2004), 1027–1158. MR 2006b:57016 Zbl 1073.57009
- [Ozsváth and Szabó 2005] P. Ozsváth and Z. Szabó, “On knot Floer homology and lens space surgeries”, *Topology* **44**:6 (2005), 1281–1300. MR 2006f:57034 Zbl 1077.57012
- [Scott 1983] P. Scott, “The geometries of 3-manifolds”, *Bull. London Math. Soc.* **15**:5 (1983), 401–487. MR 84m:57009 Zbl 0561.57001
- [Whitehead 1958] J. H. C. Whitehead, “On finite cocycles and the sphere theorem”, *Colloq. Math.* **6** (1958), 271–281. MR 20 #7271 Zbl 0119.38605

Received September 30, 2012. Revised September 16, 2013.

TYE LIDMAN  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF TEXAS AT AUSTIN  
1 UNIVERSITY STATION  
AUSTIN, TX 78712  
UNITED STATES  
tlid@math.utexas.edu

LIAM WATSON  
SCHOOL OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF GLASGOW  
15 UNIVERSITY GARDENS  
GLASGOW, G12 8QW  
UNITED KINGDOM  
liam.watson@glasgow.ac.uk



# PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
balmer@math.ucla.edu

Don Blasius  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
blasius@math.ucla.edu

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
chari@math.ucr.edu

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
cooper@math.ucsb.edu

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
finn@math.stanford.edu

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
liu@math.ucla.edu

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
jhlu@maths.hku.hk

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
popa@math.ucla.edu

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
qing@cats.ucsc.edu

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
yang@math.princeton.edu

## PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 267    No. 2    February 2014

---

Sums of squares in algebraic function fields over a complete discretely valued field	257
KARIM JOHANNES BECHER, DAVID GRIMM and JAN VAN GEEL	
On the equivalence problem for toric contact structures on $S^3$ -bundles over $S^2$	277
CHARLES P. BOYER and JUSTIN PATI	
An almost-Schur type lemma for symmetric $(2,0)$ tensors and applications	325
XU CHENG	
Algebraic invariants, mutation, and commensurability of link complements	341
ERIC CHESEBRO and JASON DEBLOIS	
Taut foliations and the action of the fundamental group on leaf spaces and universal circles	399
YOSUKE KANO	
A new monotone quantity along the inverse mean curvature flow in $\mathbb{R}^n$	417
KWOK-KUN KWONG and PENGZI MIAO	
Nonfibered L-space knots	423
TYE LIDMAN and LIAM WATSON	
Families and Springer's correspondence	431
GEORGE LUSZTIG	
Reflexive operator algebras on Banach spaces	451
FLORENCE MERLEVÈDE, COSTEL PELIGRAD and MAGDA PELIGRAD	
Harer stability and orbifold cohomology	465
NICOLA PAGANI	
Spectra of product graphs and permanents of matrices over finite rings	479
LE ANH VINH	
The concavity of the Gaussian curvature of the convex level sets of minimal surfaces with respect to the height	489
PEI-HE WANG	



0030-8730(2014)267:2;1-6