

*Pacific
Journal of
Mathematics*

**NEW INVARIANTS FOR COMPLEX MANIFOLDS
AND RATIONAL SINGULARITIES**

RONG DU AND YUN GAO

Volume 269 No. 1

May 2014

NEW INVARIANTS FOR COMPLEX MANIFOLDS AND RATIONAL SINGULARITIES

RONG DU AND YUN GAO

Dedicated to Professor Stephen S.-T. Yau on the occasion of his sixtieth birthday.

Two new invariants $f^{(1,1)}$ and $g^{(1,1)}$ were introduced by Du and Yau for solving the complex Plateau problem. These invariants measure in some sense how far away the complex manifolds are from having global complex coordinates. In this paper, we study these two invariants further for rational surface singularities. We prove that these two invariants never vanish for rational surface singularities, which confirms Yau's conjecture for strict positivity of these two invariants. As an application, we solve regularity problem of the Harvey–Lawson solution to the complex Plateau problem for a strongly pseudoconvex compact rational CR manifold of dimension 3. We also construct resolution manifolds for rational triple points by means of local coordinates and show that $f^{(1,1)} = g^{(1,1)} = 1$ for rational triple points.

1. Introduction

Let M be a complex manifold of dimension n . It is a natural question to ask how far away this complex manifold is from having global complex coordinates. Together with Hing Sun Luk and Stephen Yau, the first author introduced in [Du et al. 2011] new biholomorphic invariants that give some measurements for this purpose.

It is well-known that if M is a complex submanifold in \mathbb{C}^N , then, given any global holomorphic p -form α on M , there exists a holomorphic p -form $\tilde{\alpha}$ on \mathbb{C}^N such that the restriction of $\tilde{\alpha}$ to M is α . Obviously, $\tilde{\alpha}$ is a p -th wedge product of holomorphic 1-forms. However, for non-Stein complex manifolds, such as the resolution manifolds of singularities, the situation is totally different. In particular,

Du's research was sponsored by the National Natural Science Foundation of China, the Shanghai Pujiang Program, and the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry. Gao's research was supported by the National Natural Science Foundation of China, the Innovation Program of Shanghai Municipal Education Commission, and the Research Fund for the Doctoral Program of Higher Education of China grant 20100073110052.

MSC2010: primary 14B05, 32V15; secondary 32S25, 57P05.

Keywords: complex Plateau problem, strongly pseudoconvex, CR manifold, rational triple points.

for surface singularities, we considered in [Du et al. 2011; Du and Yau 2012] two invariants $f^{(1,1)}$ and $g^{(1,1)}$ and showed that these two invariants are strictly positive for some special singularities. So the resolution manifolds of these special singularities do not have the above properties.

A direct application of the positivity of $g^{(1,1)}$ is for solving one of the most fundamental questions in complex geometry: the complex Plateau problem [Du and Yau 2012]. Given a strongly pseudoconvex CR manifold X in \mathbb{C}^N , the problem asks when X is the boundary of a complex manifold V in \mathbb{C}^N . By the beautiful work of Harvey and Lawson [1975], as well as [Yau 1981b; Luk and Yau 1998a], X is a boundary of a complex variety V with only isolated singularities if X is contained in the boundary of a strictly pseudoconvex domain in \mathbb{C}^N . Thus from the complex Plateau problem point of view, it is very desirable to introduce a numerical invariant for isolated singularities which never vanishes. However many numerical invariants such as the geometric genus p_g , the arithmetic genus p_a and the irregularity q vanish on rational singularities. In [Du et al. 2011; Du and Yau 2012] this idea was used to introduce two invariants $f^{(1,1)}$ and $g^{(1,1)}$ for isolated surface singularities. The invariant $g^{(1,1)}$ was used in the latter article to solve the regularity problem of the Harvey–Lawson solution to the complex Plateau problem.

Those two articles provided a detailed study of $f^{(1,1)}$ and $g^{(1,1)}$. Yau has the following conjecture:

Conjecture. For all normal surface singularities, the invariants $f^{(1,1)}$ and $g^{(1,1)}$ are strictly positive.

Du and Yau showed that these two numerical invariants are strictly positive when the surface singularities have a \mathbb{C}^* -action. They also gave explicit calculations for $f^{(1,1)}$ and $g^{(1,1)}$ for rational double points and cyclic quotient singularities and proved that they are exactly 1. In this paper, we shall prove that for rational surface singularities, $f^{(1,1)}$ and $g^{(1,1)}$ also never vanish. So, our results in this paper confirm the conjecture.

Theorem 2.8. *For rational surface singularities, $f^{(1,1)} = g^{(1,1)} \geq 1$.*

As an application, we solve the regularity problem of the Harvey–Lawson solution to the complex Plateau problem for a strongly pseudoconvex compact rational CR manifold of dimension 3.

Theorem 3.9. *Let X be a strongly pseudoconvex compact rational CR manifold of dimension 3. Suppose that X is contained in the boundary of a strongly pseudoconvex bounded domain D in \mathbb{C}^N . Then X is a boundary of the complex submanifold $V \subset D - X$ with boundary regularity if and only if $g^{(1,1)}(X) = 0$.*

We also construct resolution manifolds for rational triple points by local coordinates. By using these local coordinates, we give explicit calculations of $f^{(1,1)}$ and

$g^{(1,1)}$ for rational triple points and prove that they are also 1.

Theorem 4.4. *For 2-dimensional rational triple points, $f^{(1,1)} = g^{(1,1)} = 1$.*

2. Invariants of singularities

Let V be a n -dimensional complex analytic subvariety in \mathbb{C}^N with only isolated singularities. Yau [1982] considered four kinds of sheaves of germs of holomorphic p -forms:

- (1) $\bar{\Omega}_V^p := \pi_* \Omega_M^p$, where $\pi : M \rightarrow V$ is a resolution of singularities of V .
- (2) $\bar{\bar{\Omega}}_V^p := \theta_* \Omega_{V \setminus V_{\text{sing}}}^p$ where $\theta : V \setminus V_{\text{sing}} \rightarrow V$ is the inclusion map and V_{sing} is the singular set of V .
- (3) $\Omega_V^p := \Omega_{\mathbb{C}^N}^p / \mathcal{I}^p$, where $\mathcal{I}^p = \{f\alpha + dg \wedge \beta : \alpha \in \Omega_{\mathbb{C}^N}^p, \beta \in \Omega_{\mathbb{C}^N}^{p-1}, f, g \in \mathcal{I}\}$ and \mathcal{I} is the ideal sheaf of V in \mathbb{C}^N .
- (4) $\tilde{\Omega}_V^p := \Omega_{\mathbb{C}^N}^p / \mathcal{H}^p$, where $\mathcal{H}^p = \{\omega \in \Omega_{\mathbb{C}^N}^p : \omega|_{V \setminus V_{\text{sing}}} = 0\}$.

Ω_V^p is the Grauert–Grothendieck sheaf of germs of holomorphic p -form on V . In case V is a normal variety, the dualizing sheaf ω_V of Grothendieck is actually the sheaf $\bar{\bar{\Omega}}_V^n$. Clearly $\Omega_V^p, \tilde{\Omega}_V^p$ are coherent. $\bar{\Omega}_V^p$ is a coherent sheaf because π is a proper map. $\bar{\bar{\Omega}}_V^p$ is also a coherent sheaf by Theorem A of [Siu 1970].

In [Du et al. 2011] and [Du and Yau 2012], another two sheaves $\bar{\bar{\Omega}}_V^{1,1}$ and $\bar{\bar{\Omega}}_V^{1,1}$ were considered:

Definition 2.1. Let $(V, 0)$ be a 2-dimensional Stein analytic space with an isolated singularity at 0. Let $\pi : (M, A) \rightarrow (V, 0)$ be a resolution of the singularity with A as its exceptional set. Define a sheaf of germs $\bar{\bar{\Omega}}_V^{1,1}$ as the sheaf associated to the presheaf

$$U \mapsto \langle \Gamma(\pi^{-1}(U), \Omega_M^1) \wedge \Gamma(\pi^{-1}(U), \Omega_M^1) \rangle,$$

where U is an open set of V and $\langle \Gamma(\pi^{-1}(U), \Omega_M^1) \wedge \Gamma(\pi^{-1}(U), \Omega_M^1) \rangle$ represents the module generated by elements in $\Gamma(\pi^{-1}(U), \Omega_M^1) \wedge \Gamma(\pi^{-1}(U), \Omega_M^1)$ over the ring $\Gamma(\pi^{-1}(U), \mathbb{C}_M)$.

Definition 2.2. Let $(V, 0)$ be a Stein germ of a 2-dimensional analytic space with an isolated singularity at 0. Define a sheaf of germs $\bar{\bar{\bar{\Omega}}}_V^{1,1}$ by the sheaf associated to the presheaf

$$U \mapsto \langle \Gamma(U, \bar{\bar{\bar{\Omega}}}_V^1) \wedge \Gamma(U, \bar{\bar{\bar{\Omega}}}_V^1) \rangle,$$

where U is an open set of V .

Du and Yau showed that these two new sheaves are coherent and found the relation between $\bar{\bar{\bar{\Omega}}}_V^{1,1}$ (respectively, $\bar{\bar{\bar{\Omega}}}_V^{1,1}$) and $\bar{\bar{\Omega}}_V^2$ (respectively, $\bar{\bar{\bar{\Omega}}}_V^2$) by short exact sequence as follows:

Lemma 2.3 [Du and Yau 2012]. *Let $(V, 0)$ be a 2-dimensional Stein space with an isolated singularity at 0. Let $\pi : (M, A) \rightarrow (V, 0)$ be a resolution of the singularity with A as its exceptional set. Then $\overline{\Omega}_V^{1,1}$ is coherent and there is a short exact sequence*

$$(2-1) \quad 0 \longrightarrow \overline{\Omega}_V^{1,1} \longrightarrow \overline{\Omega}_V^2 \longrightarrow \mathcal{F}^{(1,1)} \longrightarrow 0,$$

where $\mathcal{F}^{(1,1)}$ is a sheaf supported on the singular point of V . Let

$$(2-2) \quad F^{(1,1)}(M) := \Gamma(M, \Omega_M^2) / \langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle;$$

then, $\dim \mathcal{F}_0^{(1,1)} = \dim F^{(1,1)}(M)$.

Lemma 2.4 [Du and Yau 2012]. *Let V be a 2-dimensional Stein space with 0 as its only singular point. Let $\pi : (M, A) \rightarrow (V, 0)$ be a resolution of the singularity with A as its exceptional set. Then $\overline{\overline{\Omega}}_V^{1,1}$ is coherent and there is a short exact sequence*

$$(2-3) \quad 0 \longrightarrow \overline{\overline{\Omega}}_V^{1,1} \longrightarrow \overline{\overline{\Omega}}_V^2 \longrightarrow \mathcal{G}^{(1,1)} \longrightarrow 0$$

where $\mathcal{G}^{(1,1)}$ is a sheaf supported on the singular point of V . Let

$$(2-4) \quad G^{(1,1)}(M \setminus A) := \Gamma(M \setminus A, \Omega_M^2) / \langle \Gamma(M \setminus A, \Omega_M^1) \wedge \Gamma(M \setminus A, \Omega_M^1) \rangle;$$

then, $\dim \mathcal{G}_0^{(1,1)} = \dim G^{(1,1)}(M \setminus A)$.

They defined local invariants of singularities which are independent of resolution:

Definition 2.5. Let V be a 2-dimensional Stein space with 0 as its only singular point. Let $\pi : (M, A) \rightarrow (V, 0)$ be a resolution of the singularity with A as its exceptional set. Let

$$(2-5) \quad f^{(1,1)}(0) := \dim \mathcal{F}_0^{(1,1)} = \dim F^{(1,1)}(M),$$

$$(2-6) \quad g^{(1,1)}(0) := \dim \mathcal{G}_0^{(1,1)} = \dim G^{(1,1)}(M \setminus A).$$

We will omit 0 in $f^{(1,1)}(0)$ and $g^{(1,1)}(0)$ if the context is clear.

The first author and Yau conjectured that for all normal surface singularities, the invariants $f^{(1,1)}$ and $g^{(1,1)}$ are strictly positive. This conjecture was confirmed when the singularities are with \mathbb{C}^* -action.

Theorem 2.6 [Du and Yau 2012]. *Let V be a 2-dimensional Stein space with 0 as its only normal singular point with \mathbb{C}^* -action. Then $f^{(1,1)} \geq 1$.*

Theorem 2.7 [Du et al. 2011]. *Let V be a 2-dimensional Stein space with 0 as its only normal singular point with \mathbb{C}^* -action. Then $g^{(1,1)} \geq 1$.*

Theorem 2.7 is the crucial part for the solution of the classical complex Plateau problem.

We will show that these two invariants are strictly positive for rational surface singularities. So these two invariants tell the difference between smoothness and singularity more precisely than the geometry genus does in some sense.

Theorem 2.8. *For rational surface singularities, $f^{(1,1)} = g^{(1,1)} \geq 1$.*

Proof. Let $(V, 0)$ be a 2-dimensional Stein space with 0 as its only rational singularity. So, the geometry genus p_g and the irregularity q are both 0. Let $\pi : (M, A) \rightarrow (V, 0)$ be a resolution of the singularity with A as its exceptional set. Then

$$\frac{\Gamma(M \setminus A, \Omega_M^2)}{\langle \Gamma(M \setminus A, \Omega_M^1) \wedge \Gamma(M \setminus A, \Omega_M^1) \rangle} = \frac{\Gamma(M, \Omega_M^2)}{\langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle},$$

and

$$g^{(1,1)} = f^{(1,1)} = \dim \frac{\Gamma(M, \Omega_M^2)}{\langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle}.$$

From [Yau 1981a], the canonical bundle K_M is generated by its global sections in a neighborhood of the exceptional set for rational surface singularities. So, there exists $\omega \in \Gamma(M, \Omega_M^2)$ such that ω does not vanish along some irreducible component A_k of A . As the singularity is rational, A_k is a smooth rational curve. Take a tubular neighborhood U of A_k such that $U \subset M$. By the proof of Proposition 3.9 in [Du et al. 2011], we know that the elements in $\langle \Gamma(U, \Omega_U^1) \wedge \Gamma(U, \Omega_U^1) \rangle$ vanish along A_k . Since $\Gamma(M, \Omega_M^1) \subset \Gamma(U, \Omega_U^1)$, the elements in $\langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle$ also vanish along A_k . Therefore $\omega \in \Gamma(M, \Omega_M^2) \setminus \langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle$, i.e.,

$$g^{(1,1)} = \dim \frac{\Gamma(M, \Omega_M^2)}{\langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle} \geq 1. \quad \square$$

3. The complex Plateau problem for 3-dimensional rational CR manifolds

Yau [1981b] solved the classical complex Plateau problem for the case $n \geq 3$.

Theorem 3.1 [ibid.]. *Let X be a compact connected strongly pseudoconvex CR manifold of real dimension $2n - 1$ for $n \geq 3$ in the boundary of a bounded strongly pseudoconvex domain D in \mathbb{C}^{n+1} . Then X is the boundary of the complex submanifold $V \subset D - X$ if and only if the Kohn–Rossi cohomology groups $H_{KR}^{p,q}(X)$ are zero for $1 \leq q \leq n - 2$.*

Luk and Yau [2012] introduced the so-called s -invariant in order to solve the complex Plateau problem as $n = 2$. But they could not give even a sufficient condition on the boundary such that it decides the smoothness in the interior.

Theorem 3.2 [Luk and Yau 2007]. *Let X be a strongly pseudoconvex compact Calabi–Yau CR manifold of dimension 3. Suppose that X is contained in the boundary of a strongly pseudoconvex bounded domain D in \mathbb{C}^N . If the holomorphic*

de Rham cohomology $H_h^2(X)$ is 0, then X is a boundary of a complex variety V in D with boundary regularity, and V has only isolated singularities in the interior and the normalizations of these singularities are Gorenstein surface singularities with vanishing s -invariant.

Corollary 3.3 [ibid.]. *Let X be a strongly pseudoconvex compact CR manifold of dimension 3. Suppose that X is contained in the boundary of a strongly pseudoconvex bounded domain D in \mathbb{C}^3 . If the holomorphic De Rham cohomology $H_h^2(X)$ is 0, then X is a boundary of a complex variety V in D with boundary regularity, and V has only isolated quasihomogeneous singularities such that the dual graphs of the exceptional sets in the resolution are star-shaped and all the curves are rational.*

Du and Yau [2012] used a new invariant $g^{(1,1)}$ for singularities to generate a new CR invariant $g^{(1,1)}(X)$.

Definition 3.4. Suppose X is a compact connected strongly pseudoconvex CR manifold of real dimension 3. Put

$$(3-1) \quad G^{(1,1)}(X) := \mathcal{G}^2(X) / \langle \mathcal{G}^1(X) \wedge \mathcal{G}^1(X) \rangle,$$

where \mathcal{G}^p denotes the holomorphic sections of $\wedge^p(\hat{T}(X)^*)$ and $\hat{T}(X)^*$ is the holomorphic cotangent bundle of X . Then we set

$$(3-2) \quad g^{(1,1)}(X) := \dim G^{(1,1)}(X).$$

Lemma 3.5 [Du and Yau 2012]. *Let X be a compact connected strongly pseudoconvex CR manifold of real dimension 3, which bounds a bounded strongly pseudoconvex variety V with only isolated singularities $\{0_1, \dots, 0_k\}$ in \mathbb{C}^N . Then $g^{(1,1)}(X) = \sum_i g^{(1,1)}(0_i)$.*

Note that this invariant $g^{(1,1)}(X)$ can be calculated on X directly. In [ibid.], we use this CR invariant to give the sufficient and necessary condition for the variety bounded by X to be smooth if $H_h^2(X) = 0$:

Theorem 3.6 [ibid.]. *Let X be a strongly pseudoconvex compact Calabi–Yau CR manifold of dimension 3. Suppose that X is contained in the boundary of a strongly pseudoconvex bounded domain D in \mathbb{C}^N with $H_h^2(X) = 0$. Then X is the boundary of the complex submanifold up to normalization $V \subset D - X$ with boundary regularity if and only if $g^{(1,1)}(X) = 0$.*

Theorem 3.7 [ibid.]. *Let X be a strongly pseudoconvex compact CR manifold of dimension 3. Suppose that X is contained in the boundary of a strongly pseudoconvex bounded domain D in \mathbb{C}^3 with $H_h^2(X) = 0$. Then X is the boundary of the complex submanifold $V \subset D - X$ if and only if $g^{(1,1)}(X) = 0$.*

In this paper, we solve the complex Plateau problem for a strongly pseudoconvex compact rational CR manifold of dimension 3, as a corollary of Theorem 2.8.

Recall the definition of a 3-dimensional rational CR manifold [Luk and Yau 1998b]:

Definition 3.8. Let X be a connected compact strongly pseudoconvex CR manifold of real dimension 3. Let V be the normal variety such that the boundary of V is X and V has isolated singularities at $\{0_1, \dots, 0_m\}$. Let $\pi : M \rightarrow V$ be a resolution of the singularities of V . Let U_i be a strongly pseudoconvex neighborhood of 0_i , for $1 \leq i \leq m$, such that the U_i are pairwise disjoint. Then

$$p_g(X) = \sum_{i=1}^m \dim \Gamma(U_i - \{0_i\}, \Omega^2) / L^2(U_i - \{0_i\}, \Omega^2).$$

If $p_g(X) = 0$, we call the CR manifold rational.

From the similar proof of Lemma 3.9 in [Du and Yau 2012], we know that the invariant $p_g(X)$ is also decided by the holomorphic sections of holomorphic cotangent bundle of X . So it is also a CR invariant.

Theorem 3.9. *Let X be a strongly pseudoconvex compact rational CR manifold of dimension 3. Suppose that X is contained in the boundary of a strongly pseudoconvex bounded domain D in \mathbb{C}^N . Then X is a boundary of the complex submanifold $V \subset D - X$ if and only if $g^{(1,1)}(X) = 0$.*


Proof. It is clear that rational CR manifolds can bound varieties with only rational singularities. Then from Theorem 2.8 and Lemma 3.5, we obtain our conclusion. \square

4. Explicit calculation of new invariants for special rational triple points

Du et al. [2011] calculated $f^{(1,1)}$ and $g^{(1,1)}$ for rational double points and quotient singularities. In this section we will calculate these two invariants for rational triple points. We suppose that V is a 2-dimensional Stein space with 0 as its only normal singularity and that V is contractible to 0.

Artin [1966] classified the dual graphs of rational triple points of dimension 2 into nine classes, and proved that each rational triple point can be embedded into \mathbb{C}^4 . Tyurina [1968a] gave explicitly three defining equations for each singularity. Tyurina [1968b] also proved that a rational triple point is determined uniquely by its dual graph ([Laufer 1973] totally gave all the dual graphs of singularities with such property). So, isomorphically, there are nine rational triple points, for which we use the notations defined in [Chen et al. 2007]:





where \circ is a (-2) -curve and \bullet is a (-3) -curve.

In the following computation, we shall use explicit resolutions $\pi : M \rightarrow V$ of $A_{m,n,k}$, $B_{m,n}$, $C_{m,n}$, $D_{n,5}$, $E_{6,0}$, $E_{7,0}$, $E_{0,7}$, $F_{n,6}$ and $G_{n,0}$ to compute our new invariants. Note that for these rational singularities of dimension 2, the irregularity q is 0, so $f^{(1,1)} = g^{(1,1)}$. In order to calculate our new invariants for these rational singularities, we must know all the holomorphic 1-forms and holomorphic 2-forms on M . In general, the difficulty for calculating these two invariants is that the holomorphic 1-forms on the resolution manifolds are hard to express. But for rational singularities, we can use the following proposition to simplify the calculation. Campana and Flenner [2002] gave a proof of the following proposition by using mixed Hodge structure theory, which is of independent interest. We give a short proof here:

Proposition 4.1. *If $(V, 0)$ is a rational isolated singularity of dimension 2 and M is a resolution of the singularity, then $H_h^1(M) = H_h^2(M) = 0$.*

Proof. We recall the similar proof in [Du and Yau 2010]. Let $\pi : M \rightarrow V$ be a good resolution of the singularity. Let $\pi^{-1}(0) = A = \bigcup A_i$, $1 \leq i \leq n$, be the irreducible decomposition of the exceptional set A .

We have the spectral sequence

$$(4-1) \quad E_1^{p,q} = H^q(M, \Omega_M^p) \Rightarrow H^{p+q}(M, \Omega_M^\bullet) \cong H^{p+q}(M, \mathbb{C}).$$

The spectral sequence induces an exact sequence of small-order terms

$$(4-2) \quad 0 \rightarrow H_h^1(M) \rightarrow H^1(M, \mathbb{C}) \rightarrow E_2^{0,1} \rightarrow H_h^2(M) \rightarrow H^2(M, \mathbb{C}) \rightarrow E_2^{1,1} \rightarrow 0,$$

where

$$(4-3) \quad E_2^{0,1} = \ker(H^1(M, \mathbb{C}_M) \rightarrow H^1(M, \Omega_M^1)),$$

$$(4-4) \quad E_2^{1,1} = \operatorname{coker}(H^1(M, \mathbb{C}_M) \rightarrow H^1(M, \Omega_M^1)).$$

So

$$(4-5) \quad h_h^1(M) - h^1(M) + \dim E_2^{0,1} - h_h^2(M) + h^2(M) - \dim E_2^{1,1} = 0.$$

Since

$$(4-6) \quad \dim E_2^{0,1} - \dim E_2^{1,1} = h^1(M, \mathbb{C}_M) - h^1(M, \Omega_M^1),$$

we have

$$(4-7) \quad h_h^1(M) - h^1(M) - h_h^2(M) + h^2(M) + h^1(M, \mathbb{C}_M) - h^1(M, \Omega_M^1) = 0.$$

From [Wahl 1985], we know that

$$(4-8) \quad h^1(M, \Omega_M^1) = \gamma + q + n = p_g - g - b - \alpha - \beta + n$$

and

$$h^1(M) = \dim H^1(A, \mathbb{C}) = 2g + b, \quad h^2(M) = n, \quad h^1(M, \mathbb{C}_M) = p_g.$$

So

$$(4-9) \quad h_h^2(M) - h_h^1(M) = \alpha + \beta - g.$$

From [van Straten and Steenbrink 1985], we know that

$$p_g = q + g + b + \alpha + \beta + \gamma.$$

As $p_g = 0$, $q = g = b = \alpha = \beta = \gamma$. So $h^1(M) = 2g + b = 0$. From (4-2) and (4-9), we get that $h_h^2(M) = h_h^1(M) = 0$. \square

Remark 4.2. In fact, from the above proof, we can get $h_h^2(M) = h_h^1(M) = 0$ if

$$E_2^{0,1} = \ker(H^1(M, \mathbb{C}_M) \rightarrow H^1(M, \Omega_M^1)) = 0.$$

So, rational singularity is a special case.

Now we can use holomorphic functions and holomorphic 2-forms to express holomorphic 1-forms on the resolution manifold from the following lemma:

Lemma 4.3. *If $(V, 0)$ is rational isolated singularity of dimension 2 and $\pi : M \rightarrow V$ is a resolution, then for any $\xi \in \Gamma(M, \Omega_M^1)$ and $\zeta \in d^{-1}(d\xi)$, there exists an $f \in \Gamma(M, \mathbb{C}_M)$ such that $\xi = \zeta + d(f)$, where d is the exterior differential operator.*

Proof. From Proposition 4.1 above, we have the exact sequence

$$0 \longrightarrow \Gamma(M, \mathbb{C}_M) \xrightarrow{d} \Gamma(M, \Omega_M^1) \xrightarrow{d} \Gamma(M, \Omega_M^2) \longrightarrow 0.$$

For $\xi \in \Gamma(M, \Omega_M^1)$ and any $\zeta \in d^{-1}(d\xi)$, $d(\xi - \zeta) = 0$. So there exists $f \in \Gamma(M, \mathbb{C}_M)$ such that $\xi = \zeta + d(f)$. \square

From the lemma above, we see that in order to get holomorphic 1-forms on M , we only need to calculate holomorphic functions and holomorphic 2-forms on M .

Laufer [1971] constructed local coordinates for the resolution manifolds of cyclic quotient singularities such that one can calculate everything explicitly on the manifolds. Now we are going to construct local coordinates for the resolution

manifolds of rational triple points to calculate our new invariants because of the tautness of rational triple points (“tautness” means that the singularity is determined by its dual graph).

An explicit resolution $\pi : M \rightarrow V$ can be given in terms of coordinates and transition functions on M for each type as follows:

Type $E_{6,0}$:

Coordinate charts:

$$\begin{aligned} W_t &= \{(u_t, v_t)\}, & t &= 0, 1, \dots, 4, \\ W_t &= \{(u_t, v_t) : u_t^2 v_t \neq -1\}, & t &= 5, 6, \\ W_7 &= \{(u_7, v_7)\} : u_7^5 v_7^2 \neq -1\}. \end{aligned}$$

Transition functions:

$$\begin{aligned} \begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} & 0 \leq t \leq 3, & \begin{cases} u_5 = 1/(u_4 v_4), \\ v_5 = u_4 v_4^2 (1 - u_4), \end{cases} \\ \begin{cases} u_6 = 1/(v_4(1 - u_4)), \\ v_6 = u_4 v_4^2 (1 - u_4), \end{cases} & & \begin{cases} u_7 = 1/v_6, \\ v_7 = u_6 v_6^3. \end{cases} \end{aligned}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \dots \cup C_7$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, \quad 1 \leq t \leq 4, \\ C_5 &= \{v_3 = 1\} \cup \{u_4 = 1\} \cup \{v_5 = 0\}, \\ C_6 &= \{u_4 = 0\} \cup \{v_6 = 0\}, \\ C_7 &= \{u_6 = 0\} \cup \{v_7 = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned} u_4^a v_4^b (1 - u_4)^c &= u_3^b v_3^{-a+2b-c} (v_3 - 1)^c \\ &= u_2^{-a+2b-c} v_2^{-2a+3b-2c} (u_2 v_2^2 - 1)^c \\ &= u_1^{-2a+3b-2c} v_1^{-3a+4b-3c} (u_1^2 v_1^3 - 1)^c \\ &= u_0^{-3a+4b-3c} v_0^{-4a+5b-4c} (u_0^3 v_0^4 - 1)^c \\ &= u_5^{-b+2c} v_5^c (1 + u_5^2 v_5)^{-a+b-c} \\ &= u_6^{2a-b} v_6^a (1 + u_6^2 v_6)^{-a+b-c} \\ &= u_7^{5a-3b} v_7^{2a-b} (1 + u_7^5 v_7^2)^{-a+b-c}, \end{aligned}$$

such that

$$(4-10) \quad \begin{cases} a, b, c \geq 0, \\ 5b \geq 4(a + c), \\ 2c \geq b, \\ 5a \geq 3b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned} \varphi_0 &= du_0 \wedge dv_0 = du_1 \wedge dv_1 = \cdots = du_4 \wedge dv_4 \\ &= -\frac{du_5 \wedge dv_5}{1 + u_5^2 v_5} = \frac{du_6 \wedge dv_6}{1 + u_6^2 v_6} = \frac{u_7 du_7 \wedge dv_7}{1 + u_7^5 v_7^2}, \end{aligned}$$

such that

$$(4-11) \quad \begin{cases} a, b, c \geq 0, \\ 5b \geq 4(a + c), \\ 2c \geq b, \\ 5a + 1 \geq 3b. \end{cases}$$

Type $E_{7,0}$:

Coordinate charts:

$$\begin{aligned} W_t &= \{(u_t, v_t)\}, & t = 0, 1, \dots, 5, \\ W_t &= \{(u_t, v_t) : u_t^2 v_t \neq -1\}, & t = 6, 7, \\ W_8 &= \{(u_8, v_8)\} : u_8^5 v_8^2 \neq -1\}. \end{aligned}$$

Transition functions:

$$\begin{aligned} \begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} & 0 \leq t \leq 4, & \begin{cases} u_6 = 1/(u_5 v_5), \\ v_6 = u_5 v_5^2 (1 - u_5), \end{cases} \\ \begin{cases} u_7 = 1/(v_5 (1 - u_5)), \\ v_7 = u_5 v_5^2 (1 - u_5), \end{cases} & & \begin{cases} u_8 = 1/v_7, \\ v_8 = u_7 v_7^3. \end{cases} \end{aligned}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \cdots \cup C_8$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, & 1 \leq t \leq 5, \\ C_6 &= \{v_4 = 1\} \cup \{u_5 = 1\} \cup \{v_6 = 0\}, \\ C_7 &= \{u_5 = 0\} \cup \{v_7 = 0\}, \\ C_8 &= \{u_7 = 0\} \cup \{v_8 = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned}
u_5^a v_5^b (1 - u_5)^c &= u_4^b v_4^{-a+2b-c} (v_4 - 1)^c \\
&= u_3^{-a+2b-c} v_3^{-2a+3b-2c} (u_3 v_3^2 - 1)^c \\
&= u_2^{-2a+3b-2c} v_2^{-3a+4b-3c} (u_2^2 v_2^3 - 1)^c \\
&= u_1^{-3a+4b-3c} v_1^{-4a+5b-4c} (u_1^3 v_1^4 - 1)^c \\
&= u_0^{-4a+5b-4c} v_0^{-5a+6b-5c} (u_0^4 v_0^5 - 1)^c \\
&= u_6^{-b+2c} v_6^c (1 + u_6^2 v_6)^{-a+b-c} \\
&= u_7^{2a-b} v_7^a (1 + u_7^2 v_7)^{-a+b-c} \\
&= u_8^{5a-3b} v_8^{2a-b} (1 + u_8^5 v_8^2)^{-a+b-c},
\end{aligned}$$

such that

$$(4-12) \quad \begin{cases} a, b, c \geq 0, \\ 6b \geq 5(a + c), \\ 2c \geq b, \\ 5a \geq 3b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned}
\varphi_0 &= du_0 \wedge dv_0 = du_1 \wedge dv_1 = \cdots = du_5 \wedge dv_5 \\
&= -\frac{du_6 \wedge dv_6}{1 + u_6^2 v_6} = \frac{du_7 \wedge dv_7}{1 + u_7^2 v_7} = \frac{u_8 du_8 \wedge dv_8}{1 + u_8^5 v_8^2},
\end{aligned}$$

such that

$$(4-13) \quad \begin{cases} a, b, c \geq 0, \\ 6b \geq 5(a + c), \\ 2c \geq b, \\ 5a + 1 \geq 3b. \end{cases}$$

Type $E_{0,7}$:

Coordinate charts:

$$\begin{aligned}
W_t &= \{(u_t, v_t)\}, & t &= 0, 1, \dots, 5, \\
W_t &= \{(u_t, v_t) : u_t^2 v_t \neq -1\}, & t &= 6, 7, \\
W_8 &= \{(u_8, v_8)\} : u_8^3 v_8^2 \neq -1\}.
\end{aligned}$$

Transition functions:

$$\begin{cases} u_1 = 1/v_0, \\ v_1 = u_0 v_0^3, \end{cases} \quad \begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} \quad 1 \leq t \leq 4,$$

$$\begin{cases} u_6 = 1/(u_5 v_5), \\ v_6 = u_5 v_5^2 (1 - u_5), \end{cases} \quad \begin{cases} u_7 = 1/(v_5 (1 - u_5)), \\ v_7 = u_5 v_5^2 (1 - u_5), \end{cases} \quad \begin{cases} u_8 = 1/v_7, \\ v_8 = u_7 v_7^2. \end{cases}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \dots \cup C_8$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, \quad 1 \leq t \leq 5, \\ C_6 &= \{v_4 = 1\} \cup \{u_5 = 1\} \cup \{v_6 = 0\}, \\ C_7 &= \{u_5 = 0\} \cup \{v_7 = 0\}, \\ C_8 &= \{u_7 = 0\} \cup \{v_8 = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned} u_5^a v_5^b (1 - u_5)^c &= u_4^b v_4^{-a+2b-c} (v_4 - 1)^c \\ &= u_3^{-a+2b-c} v_3^{-2a+3b-2c} (u_3 v_3^2 - 1)^c \\ &= u_2^{-2a+3b-2c} v_2^{-3a+4b-3c} (u_2^2 v_2^3 - 1)^c \\ &= u_1^{-3a+4b-3c} v_1^{-4a+5b-4c} (u_1^3 v_1^4 - 1)^c \\ &= u_0^{-4a+5b-4c} v_0^{-9a+11b-9c} (u_0^4 v_0^9 - 1)^c \\ &= u_6^{-b+2c} v_6^c (1 + u_6^2 v_6)^{-a+b-c} \\ &= u_7^{2a-b} v_7^a (1 + u_7^2 v_7)^{-a+b-c} \\ &= u_8^{3a-2b} v_8^{2a-b} (1 + u_8^3 v_8^2)^{-a+b-c}, \end{aligned}$$

such that

$$(4-14) \quad \begin{cases} a, b, c \geq 0, \\ 11b \geq 9(a + c), \\ 2c \geq b, \\ 3a \geq 2b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned} \varphi_0 &= v_0 du_0 \wedge dv_0 = du_1 \wedge dv_1 = \dots = du_5 \wedge dv_5 \\ &= -\frac{du_6 \wedge dv_6}{1 + u_6^2 v_6} = \frac{du_7 \wedge dv_7}{1 + u_7^2 v_7} = \frac{du_8 \wedge dv_8}{1 + u_8^5 v_8^2}, \end{aligned}$$

such that

$$(4-15) \quad \begin{cases} a, b, c \geq 0, \\ 11b + 1 \geq 9(a + c), \\ 2c \geq b, \\ 3a \geq 2b. \end{cases}$$

Type $G_{n,0}$:

Coordinate charts:

$$\begin{aligned} W_t &= \{(u_t, v_t)\}, & t = 0, 1, \dots, n+1, \\ W_t &= \{(u_t, v_t) : u_t^3 v_t^2 \neq -1\}, & t = n+2, n+4, \\ W_{n+3} &= \{(u_{n+3}, v_{n+3})\} : u_{n+3}^2 v_{n+3} \neq -1. \end{aligned}$$

Transition functions:

$$\begin{aligned} \begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} & 0 \leq t \leq n, & \begin{cases} u_{n+2} = 1/(u_{n+1} v_{n+1}), \\ v_{n+2} = u_{n+1}^2 v_{n+1}^3 (1 - u_{n+1}), \end{cases} \\ \begin{cases} u_{n+3} = 1/(v_{n+1}(1 - u_{n+1})), \\ v_{n+3} = u_{n+1} v_{n+1}^2 (1 - u_{n+1}), \end{cases} & & \begin{cases} u_{n+4} = 1/v_{n+3}, \\ v_{n+4} = u_{n+3} v_{n+3}^2. \end{cases} \end{aligned}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \dots \cup C_{n+4}$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, \quad 1 \leq t \leq n+1, \\ C_{n+2} &= \{v_n = 1\} \cup \{u_{n+1} = 1\} \cup \{v_{n+2} = 0\}, \\ C_{n+3} &= \{u_{n+1} = 0\} \cup \{v_{n+3} = 0\}, \\ C_{n+4} &= \{u_{n+3} = 0\} \cup \{v_{n+4} = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned} u_{n+1}^a v_{n+1}^b (1 - u_{n+1})^c &= u_t^{(n-t+1)b - (n-t)(a+c)} v_t^{(n-t+2)b - (n-t+1)(a+c)} \\ &\quad \cdot (u_t^{n-t} v_t^{n-t+1} - 1)^c \\ &= u_{n+2}^{-b+3c} v_{n+2}^c (1 + u_{n+2}^3 v_{n+2})^{-a+b-c} \\ &= u_{n+3}^{2a-b} v_{n+3}^a (1 + u_{n+3}^2 v_{n+3})^{-a+b-c} \\ &= u_{n+4}^{3a-2b} v_{n+4}^{2a-b} (1 + u_{n+4}^3 v_{n+4}^2)^{-a+b-c}, \end{aligned}$$

where $0 \leq t \leq n$, such that

$$(4-16) \quad \begin{cases} a, b, c \geq 0, \\ (n+2)b \geq (n+1)(a+c), \\ 3c \geq b, \\ 3a \geq 2b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned} \varphi_0 &= du_0 \wedge dv_0 = du_1 \wedge dv_1 = \cdots = du_{n+1} \wedge dv_{n+1} \\ &= -\frac{u_{n+2} du_{n+2} \wedge dv_{n+2}}{1 + u_{n+2}^3 v_{n+2}} = \frac{du_{n+3} \wedge dv_{n+3}}{1 + u_{n+3}^2 v_{n+3}} = \frac{du_{n+4} \wedge dv_{n+4}}{1 + u_{n+4}^3 v_{n+4}^2}, \end{aligned}$$

such that

$$(4-17) \quad \begin{cases} a, b, c \geq 0, \\ (n+2)b \geq (n+1)(a+c), \\ 3c + 1 \geq b, \\ 3a \geq 2b. \end{cases}$$

Type $D_{n,5}$:

Coordinate charts:

$$\begin{aligned} W_t &= \{(u_t, v_t)\}, & t &= 0, 1, \dots, n+3, \\ W_t &= \{(u_t, v_t) : u_t^2 v_t \neq -1\}, & t &= n+4, n+5, n+7, \\ W_{n+6} &= \{(u_{n+6}, v_{n+6})\} : u_{n+6}^3 v_{n+6}^2 \neq -1. \end{aligned}$$

Transition functions:

$$\begin{aligned} &\begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} & 0 \leq t \leq n-1 \text{ and } t = n+1, n+2, n+5, n+6, \\ &\begin{cases} u_{n+1} = 1/v_n, \\ v_{n+1} = u_n v_n^3, \end{cases} & \begin{cases} u_{n+4} = 1/(u_{n+3} v_{n+3}), \\ v_{n+4} = u_{n+3} v_{n+3}^2 (1 - u_{n+3}), \end{cases} \\ &\begin{cases} u_{n+5} = 1/(v_{n+3} (1 - u_{n+3})), \\ v_{n+5} = u_{n+3} v_{n+3}^2 (1 - u_{n+3}). \end{cases} \end{aligned}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \cdots \cup C_{n+7}$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, & 1 \leq t \leq n+3 \text{ and } t = n+6, n+7, \\ C_{n+4} &= \{v_{n+2} = 1\} \cup \{u_{n+3} = 1\} \cup \{v_{n+4} = 0\}, \\ C_{n+5} &= \{u_{n+3} = 0\} \cup \{v_{n+5} = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned}
u_{n+3}^a v_{n+3}^b (1 - u_{n+3})^c &= u_{n+2}^b v_{n+2}^{-a+2b-c} (v_{n+2} - 1)^c \\
&= u_{n+1}^{-a+2b-c} v_{n+1}^{-2a+3b-2c} (u_{n+1} v_{n+1}^2 - 1)^c \\
&= u_t^{(4n-4t+3)b - (3n-3t+2)(a+c)} v_t^{(4n-4t+7)b - (3n-3t+5)(a+c)} \\
&\quad \cdot (u_t^{3n-3t+2} v_t^{3n-3t+5} - 1)^c \\
&= u_{n+4}^{-b+2c} v_{n+4}^c (1 + u_{n+4}^2 v_{n+4})^{-a+b-c} \\
&= u_{n+5}^{2a-b} v_{n+5}^a (1 + u_{n+5}^2 v_{n+5})^{-a+b-c} \\
&= u_{n+6}^{3a-2b} v_{n+6}^{2a-b} (1 + u_{n+6}^3 v_{n+6}^2)^{-a+b-c} \\
&= u_{n+7}^{4a-3b} v_{n+7}^{3a-2b} (1 + u_{n+7}^4 v_{n+7}^3)^{-a+b-c},
\end{aligned}$$

where $0 \leq t \leq n$, such that

$$(4-18) \quad \begin{cases} a, b, c \geq 0, \\ (4n+7)b \geq (3n+5)(a+c), \\ 2c \geq b, \\ 4a \geq 3b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned}
\varphi_0 &= u_0^n v_0^{n+1} du_0 \wedge dv_0 = u_1^{n-1} v_1^n du_1 \wedge dv_1 \\
&= \cdots = u_{n-1} v_{n-1}^2 du_{n-1} \wedge dv_{n-1} = u_n du_n \wedge dv_n = du_{n+1} \wedge dv_{n+1} \\
&= du_{n+2} \wedge dv_{n+2} = du_{n+3} \wedge dv_{n+3} = -\frac{du_{n+4} \wedge dv_{n+4}}{1 + u_{n+4}^3 v_{n+4}} \\
&= \frac{du_{n+5} \wedge dv_{n+5}}{1 + u_{n+5}^2 v_{n+5}} = \frac{du_{n+6} \wedge dv_{n+6}}{1 + u_{n+6}^3 v_{n+6}^2} = \frac{du_{n+7} \wedge dv_{n+7}}{1 + u_{n+7}^4 v_{n+7}^3},
\end{aligned}$$

such that

$$(4-19) \quad \begin{cases} a, b, c \geq 0, \\ (4n+7)b + n + 1 \geq (3n+5)(a+c), \\ 2c \geq b, \\ 4a \geq 3b. \end{cases}$$

Type $F_{n,6}$:

Coordinate charts:

$$\begin{aligned}
 W_t &= \{(u_t, v_t)\}, & t &= 0, 1, \dots, n+4, \\
 W_t &= \{(u_t, v_t) : u_t^2 v_t \neq -1\}, & t &= n+5, n+6, \\
 W_{n+7} &= \{(u_{n+7}, v_{n+7})\} : u_{n+7}^3 v_{n+7}^2 \neq -1\}, \\
 W_{n+8} &= \{(u_{n+8}, v_{n+8})\} : u_{n+8}^4 v_{n+8}^3 \neq -1\}.
 \end{aligned}$$

Transition functions:

$$\begin{aligned}
 &\begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} & 0 \leq t \leq n-1 \text{ and } t = n+1, n+2, n+3, n+6, n+7, \\
 &\begin{cases} u_{n+1} = 1/v_n, \\ v_{n+1} = u_n v_n^3, \end{cases} & \begin{cases} u_{n+5} = 1/(u_{n+4} v_{n+4}), \\ v_{n+5} = u_{n+4} v_{n+4}^2 (1 - u_{n+4}), \end{cases} \\
 &\begin{cases} u_{n+6} = 1/(v_{n+4} (1 - u_{n+4})), \\ v_{n+6} = u_{n+4} v_{n+4}^2 (1 - u_{n+4}). \end{cases}
 \end{aligned}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \dots \cup C_{n+8}$, where

$$\begin{aligned}
 C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\} \quad 1 \leq t \leq n+4 \text{ and } t = n+7, n+8, \\
 C_{n+5} &= \{v_{n+3} = 1\} \cup \{u_{n+4} = 1\} \cup \{v_{n+5} = 0\}, \\
 C_{n+6} &= \{u_{n+4} = 0\} \cup \{v_{n+6} = 0\}.
 \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned}
 &u_{n+4}^a v_{n+4}^b (1 - u_{n+4})^c \\
 &= u_{n+3}^b v_{n+3}^{-a+2b-c} (v_{n+3} - 1)^c \\
 &= u_{n+2}^{-a+2b-c} v_{n+2}^{-2a+3b-2c} (u_{n+2} v_{n+2}^2 - 1)^c \\
 &= u_{n+1}^{-2a+3b-2c} v_{n+1}^{-3a+4b-3c} (u_{n+1}^2 v_{n+1}^3 - 1)^c \\
 &= u_t^{(5n-5t+4)b - (4n-4t+3)(a+c)} v_t^{(5n-5t+9)b - (4n-4t+7)(a+c)} \\
 &\quad \cdot (u_t^{4n-4t+3} v_t^{4n-4t+7} - 1)^c \\
 &= u_{n+5}^{-b+2c} v_{n+5}^c (1 + u_{n+5}^2 v_{n+5})^{-a+b-c} \\
 &= u_{n+6}^{2a-b} v_{n+6}^a (1 + u_{n+6}^2 v_{n+6})^{-a+b-c} \\
 &= u_{n+7}^{3a-2b} v_{n+7}^{2a-b} (1 + u_{n+7}^3 v_{n+7}^2)^{-a+b-c} \\
 &= u_{n+8}^{4a-3b} v_{n+8}^{3a-2b} (1 + u_{n+8}^4 v_{n+8}^3)^{-a+b-c},
 \end{aligned}$$

where $0 \leq t \leq n$, such that

$$(4-20) \quad \begin{cases} a, b, c \geq 0, \\ (5n+9)b \geq (4n+7)(a+c), \\ 2c \geq b, \\ 4a \geq 3b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned} \varphi_0 &= u_0^n v_0^{n+1} du_0 \wedge dv_0 = u_1^{n-1} v_1^n du_1 \wedge dv_1 \\ &= \cdots = u_{n-1} v_{n-1}^2 du_{n-1} \wedge dv_{n-1} = u_n du_n \wedge dv_n \\ &= du_{n+1} \wedge dv_{n+1} = \cdots = du_{n+4} \wedge dv_{n+4} \\ &= \frac{du_{n+5} \wedge dv_{n+5}}{1 + u_{n+5}^3 v_{n+5}} = \frac{du_{n+6} \wedge dv_{n+6}}{1 + u_{n+6}^2 v_{n+6}} \\ &= \frac{du_{n+7} \wedge dv_{n+7}}{1 + u_{n+7}^3 v_{n+7}^2} = \frac{du_{n+8} \wedge dv_{n+8}}{1 + u_{n+8}^4 v_{n+8}^3}, \end{aligned}$$

such that

$$(4-21) \quad \begin{cases} a, b, c \geq 0, \\ (5n+9)b + n + 1 \geq (4n+7)(a+c), \\ 2c \geq b, \\ 4a \geq 3b. \end{cases}$$

Type $C_{m,n}$:

Coordinate charts:

$$\begin{aligned} W_t &= \{(u_t, v_t)\}, & t &= 0, 1, \dots, m+n+2, \\ W_t &= \{(u_t, v_t) : u_t^2 v_t \neq -1\}, & t &= m+n+3, m+n+4. \end{aligned}$$

Transition functions:

$$\begin{aligned} &\begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} & 0 \leq t \leq n-1 \text{ and } n+1 \leq t \leq m+n+1, \\ &\begin{cases} u_{n+1} = 1/v_n, \\ v_{n+1} = u_n v_n^3, \end{cases} & \begin{cases} u_{m+n+3} = 1/(u_{m+n+2} v_{m+n+2}), \\ v_{m+n+3} = u_{m+n+2} v_{m+n+2}^2 (1 - u_{m+n+2}), \end{cases} \\ &\begin{cases} u_{m+n+4} = 1/(v_{m+n+2} (1 - u_{m+n+2})), \\ v_{m+n+4} = u_{m+n+2} v_{m+n+2}^2 (1 - u_{m+n+2}). \end{cases} \end{aligned}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \dots \cup C_{m+n+4}$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, \quad 1 \leq t \leq m+n+2, \\ C_{m+n+3} &= \{v_{m+n+1} = 1\} \cup \{u_{m+n+2} = 1\} \cup \{v_{m+n+3} = 0\}, \\ C_{m+n+4} &= \{u_{m+n+2} = 0\} \cup \{v_{m+n+4} = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned} u_{m+n+2}^a v_{m+n+2}^b (1 - u_{m+n+2})^c &= u_t^{(m+n+2-t)b - (m+n+1-t)(a+c)} v_t^{(m+n+3-t)b - (m+n+2-t)(a+c)} \\ &\quad \cdot (u_t^{m+n+1-t} v_t^{m+n+2-t} - 1)^c \\ &= u_s^{[(m+3)(n-s)+m+2]b - [(m+3)(n-s)+m+1](a+c)} \\ &\quad \cdot v_s^{[(m+3)(n-s)+m+2]b - [(m+3)(n-s)+m+1](a+c)} \\ &\quad \cdot (u_s^{(m+3)(n-s)+m+2} v_s^{(m+3)(n-s)+2m+3} - 1)^c \\ &= u_{m+n+3}^{-b+2c} v_{m+n+3}^c (1 + u_{m+n+3}^2 v_{m+n+3})^{-a+b-c} \\ &= u_{m+n+4}^{2a-b} v_{m+n+4}^a (1 + u_{m+n+4}^2 v_{m+n+4})^{-a+b-c}, \end{aligned}$$

where $n+1 \leq t \leq m+n+1$, $0 \leq s \leq n$, such that

$$(4-22) \quad \begin{cases} a, b, c \geq 0, \\ (mn + 3m + 2n + 5)b \geq (mn + 2m + 2n + 3)(a + c), \\ 2c \geq b, \\ 2a \geq b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned} \varphi_0 &= u_0^n v_0^{n+1} du_0 \wedge dv_0 = u_1^{n-1} v_1^n du_1 \wedge dv_1 \\ &= \dots = u_{n-1} v_{n-1}^2 du_{n-1} \wedge dv_{n-1} = u_n du_n \wedge dv_n = du_{n+1} \wedge dv_{n+1} \\ &= \dots = du_{m+n+2} \wedge dv_{m+n+2} = -\frac{du_{m+n+3} \wedge dv_{m+n+3}}{1 + u_{m+n+3}^2 v_{m+n+3}} \\ &= \frac{du_{m+n+4} \wedge dv_{m+n+4}}{1 + u_{m+n+4}^2 v_{m+n+4}}, \end{aligned}$$

such that

$$(4-23) \quad \begin{cases} a, b, c \geq 0, \\ (mn + 3m + 2n + 5)b + n + 1 \geq (mn + 2m + 2n + 3)(a + c), \\ 2c \geq b, \\ 2a \geq b. \end{cases}$$

Type $B_{m,n}$:

Coordinate charts:

$$\begin{aligned} W_t &= \{(u_t, v_t)\}, & t = 0, 1, \dots, m+2, \\ W_{m+3} &= \{(u_{m+3}, v_{m+3}) : u_{m+3}^2 v_{m+3} \neq -1\}, \\ W_t &= \{(u_t, v_t) : u_t^{t-m-2} v_t^{t-m-3} \neq -1\}, & m+4 \leq t \leq m+n+4. \end{aligned}$$

Transition functions:

$$\begin{cases} u_{t+1} = 1/v_t, \\ v_{t+1} = u_t v_t^2, \end{cases} \quad 0 \leq t \leq m-1, t = m+1, \text{ and } m+4 \leq t \leq m+n+3,$$

$$\begin{cases} u_{m+1} = 1/v_m, \\ v_{m+1} = u_m v_m^3, \end{cases} \quad \begin{cases} u_{m+3} = 1/(u_{m+2} v_{m+2}), \\ v_{m+3} = u_{m+2} v_{m+2}^2 (1 - u_{m+2}), \end{cases}$$

$$\begin{cases} u_{m+4} = 1/(v_{m+2} (1 - u_{m+2})), \\ v_{m+4} = u_{m+2} v_{m+2}^2 (1 - u_{m+2}). \end{cases}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \dots \cup C_{m+n+4}$, where

$$\begin{aligned} C_t &= \{u_{t-1} = 0\} \cup \{v_t = 0\}, \quad 1 \leq t \leq m+2 \text{ and } m+5 \leq t \leq m+n+4, \\ C_{m+3} &= \{v_{m+1} = 1\} \cup \{u_{m+2} = 1\} \cup \{v_{m+3} = 0\}, \\ C_{m+4} &= \{u_{m+2} = 0\} \cup \{v_{m+4} = 0\}. \end{aligned}$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned} &u_{m+2}^a v_{m+2}^b (1 - u_{m+2})^c \\ &= u_{m+1}^b v_{m+1}^{-a+2b-c} (v_{m+1} 1)^c \\ &= u_t^{(3m-3t+2)b-(2m-2t+1)(a+c)} v_t^{(3m-3t+5)b-(2m-2t+3)(a+c)} \\ &\quad \cdot (u_t^{2m-2t+1} v_t^{2m-2t+3} - 1)^c \\ &= u_{m+3}^{-b+2c} v_{m+3}^c (1 + u_{m+3}^2 v_{m+3})^{-a+b-c} \\ &= u_s^{(t-m-2)a-(t-m-3)b} v_s^{(t-m-3)a-(t-m-4)b} (u_s^{t-m-2} v_s^{t-m-3} - 1)^c, \end{aligned}$$

where $0 \leq t \leq m$, $m+4 \leq s \leq m+n+4$, such that

$$(4-24) \quad \begin{cases} a, b, c \geq 0, \\ (3m+5)b \geq (2m+3)(a+c), \\ 2c \geq b, \\ (n+2)a \geq (n+1)b. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned}
 \varphi_0 &= u_0^m v_0^{m+1} du_0 \wedge dv_0 = u_1^{m-1} v_1^m du_1 \wedge dv_1 \\
 &= \cdots = u_{m-1} v_{m-1}^2 du_{m-1} \wedge dv_{m-1} = u_m du_m \wedge dv_m \\
 &= du_{m+1} \wedge dv_{m+1} = du_{m+2} \wedge dv_{m+2} \\
 &= -\frac{du_{m+3} \wedge dv_{m+3}}{1 + u_{m+3}^2 v_{m+3}} \\
 &= \frac{du_t \wedge dv_t}{1 + u_t^{t-m-2} v_t^{t-m-3}},
 \end{aligned}$$

where $m + 4 \leq t \leq m + n + 4$, such that

$$(4-25) \quad \begin{cases} a, b, c \geq 0, \\ (3m + 5)b + m + 1 \geq (2m + 3)(a + c), \\ 2c \geq b, \\ (n + 2)a \geq (n + 1)b. \end{cases}$$

Type $A_{m,n,k}$:

Coordinate charts:

$$\begin{aligned}
 W_t &= \{(u_t, v_t)\}, & t &= 0, 1, \dots, m + 1, \\
 W_t &= \{(u_t, v_t) : u_t^{t-m} v_t^{t-m-1} \neq -1\}, & m + 2 &\leq t \leq m + n + 1, \\
 W_t &= \{(u_t, v_t) : u_t^{t-m-n} v_t^{t-m-n-1} \neq -1\}, & m + n + 2 &\leq t \leq m + n + k + 1.
 \end{aligned}$$

Transition functions:

$$\begin{cases} u_{t+1} = 1/v_t, & 0 \leq t \leq m - 1, m + 2 \leq t \leq m + n, \text{ or} \\ v_{t+1} = u_t v_t^2, & m + n + 2 \leq t \leq m + n + k, \end{cases}$$

$$\begin{cases} u_{m+1} = 1/v_m, \\ v_{m+1} = u_m v_m^3, \end{cases}$$

$$\begin{cases} u_{m+2} = 1/(v_{m+1}(1 - u_{m+1})), \\ v_{m+2} = u_{m+1} v_{m+1}^2 (1 - u_{m+1}), \end{cases}$$

$$\begin{cases} u_{m+n+2} = 1/(u_{m+1} v_{m+1}), \\ v_{m+n+2} = u_{m+1} v_{m+1}^2 (1 - u_{m+1}). \end{cases}$$

Exceptional set: $A = \pi^{-1}(0) = C_1 \cup \cdots \cup C_{m+n+k+1}$, where

$$C_t = \{u_{t-1} = 0\} \cup \{v_t = 0\}, \quad t \neq m+n+2,$$

$$C_{m+n+2} = \{v_m = 1\} \cup \{u_{m+1} = 1\} \cup \{v_{m+n+2} = 0\}.$$

A holomorphic function on M can be generated by the following forms:

$$\begin{aligned} & u_{m+1}^a v_{m+1}^b (1 - u_{m+1})^c \\ &= u_t^{(2m-2t+1)b - (m-t)(a+c)} v_t^{(2m-2t+3)b - (m-t+1)(a+c)} (u_t^{m-t} v_t^{m-t+1} - 1)^c \\ &= u_s^{(s-m)a - (s-m-1)b} v_s^{(s-m-1)a - (s-m-2)b} (u_s^{s-m} v_s^{s-m-1} - 1)^c, \\ &= u_r^{(r-m-n)c - (r-m-n-1)b} v_r^{(r-m-n-1)c - (r-m-n-2)b} \\ &\quad \cdot (u_r^{r-m-n} v_r^{r-m-n-1} - 1)^c, \end{aligned}$$

where $0 \leq t \leq m$, $m+2 \leq s \leq m+n+1$, $m+n+2 \leq r \leq m+n+k+1$, such that

$$(4-26) \quad \begin{cases} a, b, c \geq 0, \\ (2m+3)b \geq (m+1)(a+c), \\ (n+1)a \geq nb, \\ (k+1)c \geq kb. \end{cases}$$

A holomorphic 2-form can be written as $f\varphi_0$, where f is a holomorphic function on M and

$$\begin{aligned} \varphi_0 &= u_0^m v_0^{m+1} du_0 \wedge dv_0 = u_1^{m-1} v_1^m du_1 \wedge dv_1 \\ &= \cdots = u_{m-1} v_{m-1}^2 du_{m-1} \wedge dv_{m-1} = u_m du_m \wedge dv_m = du_{m+1} \wedge dv_{m+1} \\ &= \frac{du_t \wedge dv_t}{1 + u_t^{t-m} v_t^{t-m-1}} \\ &= -\frac{du_s \wedge dv_s}{1 + u_s^{s-m-n} v_s^{s-m-n-1}}, \end{aligned}$$

where $m+2 \leq t \leq m+n+1$, $m+n+2 \leq s \leq m+n+k+1$, such that

$$(4-27) \quad \begin{cases} a, b, c \geq 0, \\ (2m+3)b + m + 1 \geq (m+1)(a+c), \\ (n+1)a \geq nb, \\ (k+1)c \geq kb. \end{cases}$$

Theorem 4.4. *For 2-dimensional rational triple points, $f^{(1,1)} = g^{(1,1)} = 1$.*

Proof. We consider only type $E_{6,0}$ as an example; the other singularities are similar. From the above calculation, we know that the holomorphic functions on M are generated by a base $\{u_4^a v_4^b (1 - u_4)^c\}$ satisfying (4-10), and holomorphic 2-forms

are generated by a base $\{u_4^a v_4^b (1-u_4)^c du_4 \wedge dv_4\}$ satisfying (4-11). For every holomorphic 2-form $\omega = u_4^a v_4^b (1-u_4)^c du_4 \wedge dv_4$ on M , we consider

$$\xi = -\frac{u_4^a v_4^{b+1} (1-u_4)^c}{b+1} du_4.$$

So, ξ defines a holomorphic 1-form on W_0 and $d\xi = \omega$. In fact, we need to check that under all changes of coordinate charts, ξ transforms to define a holomorphic 1-form in each coordinate chart:

$$\begin{aligned} u_4^a v_4^{b+1} (1-u_4)^c du_4 &= -u_3^{b+1} v_3^{-a+2b-c} (v_3-1)^c dv_3 \\ &= -u_2^{-a+2b-c} v_2^{-2a+3b-2c+1} (u_2 v_2^2 - 1)^c du_2 \\ &\quad - 2u_2^{-a+2b-c+1} v_2^{-2a+3b-2c} (u_2 v_2^2 - 1)^c dv_2 \\ &= -2u_1^{-2a+3b-2c} v_1^{-3a+4b-3c+1} (u_1^2 v_1^3 - 1)^c du_1 \\ &\quad - 3u_1^{-2a+3b-2c+1} v_1^{-3a+4b-3c} (u_1^2 v_1^3 - 1)^c dv_1 \\ &= -3u_0^{-3a+4b-3c} v_0^{-4a+5b-4c+1} (u_0^3 v_0^4 - 1)^c du_0 \\ &\quad - 4u_0^{-3a+4b-3c+1} v_0^{-4a+5b-4c} (u_0^3 v_0^4 - 1)^c dv_0 \\ &= -2u_5^{-b+2c} v_5^{c+1} (1+u_5^2 v_5)^{-a+b-c-1} du_5 \\ &\quad - u_5^{-b+2c+1} v_5^c (1+u_5^2 v_5)^{-a+b-c-1} dv_5 \\ &= 2u_6^{2a-b} v_6^{a+1} (1+u_6^2 v_6)^{-a+b-c-1} du_6 \\ &\quad + u_6^{2a-b+1} v_6^a (1+u_6^2 v_6)^{-a+b-c-1} dv_6 \\ &= 5u_7^{5a-3b+1} v_7^{2a-b+1} (1+u_7^5 v_7^2)^{-a+b-c-1} du_7 \\ &\quad + 2u_7^{5a-3b+2} v_7^{2a-b} (1+u_7^5 v_7^2)^{-a+b-c-1} dv_7. \end{aligned}$$

By Lemma 4.3, we can get general expressions for elements in $\Gamma(M, \Omega_M^1)$. Therefore

$$\frac{\Gamma(M, \Omega_M^2)}{\langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle} = \langle u_4 \wedge v_4 \rangle,$$

and

$$f^{(1,1)} = \dim \frac{\Gamma(M, \Omega_M^2)}{\langle \Gamma(M, \Omega_M^1) \wedge \Gamma(M, \Omega_M^1) \rangle} = 1. \quad \square$$

References

- [Artin 1966] M. Artin, "On isolated rational singularities of surfaces", *Amer. J. Math.* **88** (1966), 129–136. MR 33 #7340 Zbl 0142.18602
- [Campana and Flenner 2002] F. Campana and H. Flenner, "Contact singularities", *Manuscripta Math.* **108**:4 (2002), 529–541. MR 2003k:32041 Zbl 1026.32057

- [Chen et al. 2007] Z. Chen, R. Du, S.-L. Tan, and F. Yu, “Cubic equations of rational triple points of dimension two”, pp. 63–76 in *Proceedings of the International Conference on Complex Geometry and Related Fields* (Shanghai, 2004), edited by Z. Chen et al., AMS/IP Stud. Adv. Math. **39**, Amer. Math. Soc., Providence, RI, 2007. MR 2008g:14056 Zbl 1127.14003
- [Du and Yau 2010] R. Du and S. Yau, “Local holomorphic de Rham cohomology”, *Comm. Anal. Geom.* **18**:2 (2010), 365–374. MR 2011g:32039 Zbl 1216.32018
- [Du and Yau 2012] R. Du and S. Yau, “Kohn–Rossi cohomology and its application to the complex Plateau problem, III”, *J. Differential Geom.* **90**:2 (2012), 251–266. MR 2899875 Zbl 1254.32051
- [Du et al. 2011] R. Du, H. S. Luk, and S. Yau, “New invariants for complex manifolds and isolated singularities”, *Comm. Anal. Geom.* **19**:5 (2011), 991–1021. MR 2886715 Zbl 1243.32018
- [Harvey and Lawson 1975] F. R. Harvey and H. B. Lawson, Jr., “On boundaries of complex analytic varieties, I”, *Ann. of Math. (2)* **102**:2 (1975), 223–290. MR 54 #13130 Zbl 0317.32017
- [Laufer 1971] H. B. Laufer, *Normal two-dimensional singularities*, Annals of Mathematics Studies **71**, Princeton University Press, 1971. MR 47 #8904 Zbl 0245.32005
- [Laufer 1973] H. B. Laufer, “Taut two-dimensional singularities”, *Math. Ann.* **205** (1973), 131–164. MR 48 #11563 Zbl 0281.32010
- [Luk and Yau 1998a] H. S. Luk and S. S.-T. Yau, “Counterexample to boundary regularity of a strongly pseudoconvex CR submanifold: An addendum to the paper of Harvey–Lawson”, *Ann. of Math. (2)* **148**:3 (1998), 1153–1154. MR 99j:32006 Zbl 0940.32014
- [Luk and Yau 1998b] H. S. Luk and S. S. T. Yau, “Explicit construction of graph invariant for strongly pseudoconvex compact 3-dimensional rational CR manifolds”, *Compositio Math.* **114**:1 (1998), 77–111. MR 2000d:32054 Zbl 0917.32008
- [Luk and Yau 2007] H. S. Luk and S. S.-T. Yau, “Kohn–Rossi cohomology and its application to the complex Plateau problem, II”, *J. Differential Geom.* **77**:1 (2007), 135–148. MR 2008k:32091 Zbl 1123.32020
- [Siu 1970] Y.-t. Siu, “Analytic sheaves of local cohomology”, *Trans. Amer. Math. Soc.* **148** (1970), 347–366. MR 41 #2054 Zbl 0195.36802
- [van Straten and Steenbrink 1985] D. van Straten and J. Steenbrink, “Extendability of holomorphic differential forms near isolated hypersurface singularities”, *Abh. Math. Sem. Univ. Hamburg* **55** (1985), 97–110. MR 87j:32025 Zbl 0584.32018
- [Tjurina 1968a] G. N. Tjurina, “Absolute isolatedness of rational singularities, and triple rational points”, *Funkcional. Anal. i Priložen.* **2**:4 (1968), 70–81. In Russian; translated in *Func. Anal. Appl.* **2**:4 (1968), 324–333. MR 39 #7136 Zbl 0176.50804
- [Tjurina 1968b] G. N. Tjurina, “The rigidity of rationally contractible curves on a surface”, *Izv. Akad. Nauk SSSR Ser. Mat.* **32** (1968), 943–970. In Russian. MR 40 #149 Zbl 0186.26301
- [Wahl 1985] J. M. Wahl, “A characterization of quasihomogeneous Gorenstein surface singularities”, *Compositio Math.* **55**:3 (1985), 269–288. MR 87e:32013 Zbl 0587.14024
- [Yau 1981a] S. S. T. Yau, “Existence of L^2 -integrable holomorphic forms and lower estimates of T_V^1 ”, *Duke Math. J.* **48**:3 (1981), 537–547. MR 83b:32009 Zbl 0474.14020
- [Yau 1981b] S. S. T. Yau, “Kohn–Rossi cohomology and its application to the complex Plateau problem, I”, *Ann. of Math. (2)* **113**:1 (1981), 67–110. MR 82k:32038 Zbl 0464.32012
- [Yau 1982] S. S. T. Yau, “Various numerical invariants for isolated singularities”, *Amer. J. Math.* **104**:5 (1982), 1063–1100. MR 84b:32012 Zbl 0523.14002

Received October 7, 2011.

RONG DU
DEPARTMENT OF MATHEMATICS
EAST CHINA NORMAL UNIVERSITY
DONGCHUAN ROAD 500
SHANGHAI 200241
CHINA
rd@math.ecnu.edu.cn

YUN GAO
DEPARTMENT OF MATHEMATICS
SHANGHAI JIAO TONG UNIVERSITY
DONGCHUAN ROAD 800
SHANGHAI 200240
CHINA
gaoyunmath@sjtu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 269 No. 1 May 2014

The asymptotic behavior of Palais–Smale sequences on manifolds with boundary	1
SÉRGIO ALMARAZ	
The cup subalgebra of a II_1 factor given by a subfactor planar algebra is maximal amenable	19
ARNAUD BROTHIER	
Representation theory of type B and C standard Levi W -algebras	31
JONATHAN BROWN and SIMON M. GOODWIN	
New invariants for complex manifolds and rational singularities	73
RONG DU and YUN GAO	
Homogeneity groups of ends of open 3-manifolds	99
DENNIS J. GARITY and DUŠAN REPOVŠ	
On the concircular curvature of a (κ, μ, ν) -manifold	113
FLORENCE GOULI-ANDREOU and EVAGGELIA MOUTAFI	
Genuses of cluster quivers of finite mutation type	133
FANG LI, JICHUN LIU and YICHAO YANG	
Taut foliations in knot complements	149
TAO LI and RACHEL ROBERTS	
On the set of maximal nilpotent supports of supercuspidal representations	169
QIN YUJUN	
The natural filtrations of finite-dimensional modular Lie superalgebras of Witt and Hamiltonian type	199
KELI ZHENG, YONGZHENG ZHANG and WEI SONG	
Free Brownian motion and free convolution semigroups: multiplicative case	219
PING ZHONG	



0030-8730(201405)269:1;1-0