

*Pacific
Journal of
Mathematics*

**ON THE CLASSIFICATION OF COMPLETE
AREA-STATIONARY AND STABLE SURFACES IN THE
SUBRIEMANNIAN SOL MANIFOLD**

MATTEO GALLI

Volume 271 No. 1

September 2014

ON THE CLASSIFICATION OF COMPLETE AREA-STATIONARY AND STABLE SURFACES IN THE SUBRIEMANNIAN SOL MANIFOLD

MATTEO GALLI

We study the classification of area-stationary and stable C^2 regular surfaces in the space of the rigid motions of the Minkowski plane $E(1, 1)$, equipped with its subriemannian structure. We construct examples of area-stationary surfaces that are not foliated by subriemannian geodesics. We also prove that there exist an infinite number of C^2 area-stationary surfaces with a singular curve. Finally we show the stability of C^2 area-stationary surfaces foliated by subriemannian geodesics.

1. Introduction

The study of the subriemannian area functional in three-dimensional pseudohermitian manifolds and in other subriemannian spaces has been largely investigated in the last years, see [Ambrosio et al. 2006; Barbieri and Citti 2011; Barone Adesi et al. 2007; Bigolin and Cassano 2010; Capogna et al. 2009; Cheng and Hwang 2010; Cheng et al. 2012; 2005; 2007; Danielli et al. 2007; 2008; 2009; Galli 2013; Galli and Ritoré 2013; Garofalo and Nhieu 1996; Hladky and Pauls 2008; Hurtado et al. 2010; Hurtado and Rosales 2008; \geq 2014; Ritoré 2009; Ritoré and Rosales 2008; Rosales 2012; Shcherbakova 2009], among others.

One of the more interesting questions concerning the subriemannian area functional is this:

Problem 1. Which are the area-minimizing surfaces in a given three-dimensional contact subriemannian manifold?

A surface Σ is *area-minimizing* if $A(\Sigma) \leq A(\tilde{\Sigma})$, for any compact deformation $\tilde{\Sigma}$ of Σ . To answer the previous question, a natural preliminary step is to study the *area-stationary* surfaces, the critical points of the area functional.

Research supported by MCyT-Feder grant MTM2010-21206-C02-01 and J. A. grant P09-FMQ-5088.
MSC2010: 49Q05, 49Q20, 53C17.

Keywords: subriemannian geometry, area-stationary surfaces, stable surfaces, pseudohermitian manifolds, Sol geometry.

Problem 2. Which are the area-stationary surfaces in a given three-dimensional contact subriemannian manifold?

We will consider these questions in the class of C^2 regular surfaces. For a general introduction about the study of the area functional in subriemannian spaces, we refer the interested reader to [Capogna et al. 2007] and [Galli 2012], which treat the case of \mathbb{H}^n and the contact subriemannian manifolds respectively.

In Sasakian space forms, the classification of C^2 area stationary surfaces was given in [Hurtado et al. 2010] in the case of the Heisenberg group \mathbb{H}^1 and in [Rosales 2012] for the Sasakian structures of S^3 and $\widetilde{\text{SL}}_2(\mathbb{R})$. In the case of pseudohermitian three-manifolds that are not Sasakian, the only known results concerning Problem 1 and Problem 2 are given in [Galli 2013], where the group of the rigid motions of the Euclidean plane $E(2)$ is studied.

Concerning the three-dimensional pseudohermitian manifolds, we have the following classification result, [Perrone 1998, Theorem 3.1], in terms of the Webster scalar curvature W and of the pseudohermitian torsion τ :

Proposition 1.1. *Let M be a simply connected contact 3-manifold that is homogeneous (in the sense of [Boothby and Wang 1958]). The following possibilities arise:*

1. *If M is unimodular, it is*
 - (i) *the first Heisenberg group \mathbb{H}^1 when $W = |\tau| = 0$;*
 - (ii) *the three-sphere group $\text{SU}(2)$ when $W > 2|\tau|$;*
 - (iii) *the group $\widetilde{\text{SL}}_2(\mathbb{R})$ when $-2|\tau| \neq W < 2|\tau|$;*
 - (iv) *the group $\widetilde{E}(2)$, universal cover of the group of rigid motions of the Euclidean plane, when $W = 2|\tau| > 0$;*
 - (v) *the group $E(1, 1)$ of rigid motions of Minkowski 2-space, when $W = -2|\tau| < 0$;*
2. *If M is not unimodular, the Lie algebra is given by*

$$[X, Y] = \alpha Y + 2T, \quad [X, T] = \gamma Y, \quad [Y, T] = 0, \quad \alpha \neq 0,$$

where $\{X, Y\}$ is an orthonormal basis of \mathcal{H} , $J(X) = Y$ and T is the Reeb vector field. In this case $W < 2|\tau|$ and when $\gamma = 0$ the structure is Sasakian and $W = -\alpha^2$.

The only case for which Problems 1 and 2 have not been investigated is that of Sol geometry, modeled by the space $E(1, 1)$. Its study is the aim of the present work.

After some preliminaries, the paper is organized as follows.

In Section 3, we compute explicitly the coordinates of the characteristic curves with given initial conditions. These curves play an important role in the study of area-stationary surfaces, since the regular part $\Sigma - \Sigma_0$ of a surface Σ is foliated by characteristic curves, that are not in general subriemannian geodesics, since $E(1, 1)$ is characterized by a nonvanishing pseudohermitian torsion.

Section 4 is the core of the paper. We first characterize the C^2 complete, area-stationary surfaces immersed in $E(1, 1)$ with singular points or singular curves that are subriemannian geodesics. On the other hand, for the first time in the three-dimensional pseudohermitian setting, we also find examples of area-stationary surfaces that are not foliated by subriemannian geodesics. We stress that these examples form an infinite family; that is, given an horizontal curve Γ , we can construct an area-stationary surface having Γ as singular set Σ_0 .

Finally in Section 5 we prove that complete area-stationary surfaces with non-empty singular set, whose characteristic curves are subriemannian geodesics, are stable. We also find three families of nonsingular planes that are area-minimizing, using a calibration argument.

We remark that Section 5 opens two interesting questions. Is a stable complete area-stationary surface in $E(1, 1)$ with a singular curve always foliated by subriemannian geodesics in $\Sigma - \Sigma_0$? Do some other complete stable area-stationary surfaces in $E(1, 1)$ with empty singular set exist?

2. Preliminaries

The group $E(1, 1)$ of rigid motions of the Minkowski plane. We consider the group of rigid motions of the Minkowski plane $E(1, 1)$, a unimodular Lie group with a natural subriemannian structure. As a model of $E(1, 1)$ we choose the underlying manifold \mathbb{R}^3 with the following orthonormal basis of left-invariant vector fields:

$$(2-1) \quad X = \frac{\partial}{\partial z}, \quad Y = \frac{1}{\sqrt{2}} \left(-e^z \frac{\partial}{\partial x} + e^{-z} \frac{\partial}{\partial y} \right), \quad T = \frac{1}{\sqrt{2}} \left(e^z \frac{\partial}{\partial x} + e^{-z} \frac{\partial}{\partial y} \right).$$

We have that $\{X, Y\}$ is an orthonormal basis of the horizontal distribution \mathcal{H} and T is the Reeb vector field. The scalar product of two vector fields W and V with respect to the metric induced by the basis $\{X, Y, T\}$ will be often denoted by $\langle W, V \rangle$. This structure of $E(1, 1)$ is characterized by the following Lie brackets, [Milnor 1976],

$$(2-2) \quad [X, Y] = -T, \quad [X, T] = -Y, \quad [Y, T] = 0.$$

In fact, applying [Galli 2013, (9.1) and (9.3)] we obtain that the Webster scalar curvature is $W = -\frac{1}{2}$ and the matrix of the pseudohermitian torsion τ in the X, Y, T

basis is

$$\begin{pmatrix} 0 & -\frac{1}{2} & 0 \\ -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The following derivatives can be easily computed:

$$(2-3) \quad \begin{aligned} \nabla_X X &= 0, & \nabla_Y X &= 0, & \nabla_T X &= \frac{1}{2}Y, \\ \nabla_X Y &= 0, & \nabla_Y Y &= 0, & \nabla_T Y &= -\frac{1}{2}X, \end{aligned}$$

where ∇ denotes the pseudohermitian connection; see [Dragomir and Tomassini 2006]. Moreover we have $-2|\tau|^2 = W < 0$, which characterizes $E(1, 1)$; see [Perrone 1998]. We also define the involution J on \mathcal{H} , called the complex structure, by $J(X) = Y$ and $J(Y) = -X$.

The geometry of regular surfaces in $E(1, 1)$. Consider a C^1 surface Σ immersed in $E(1, 1)$. We define the *subriemannian area* of Σ as

$$A(\Sigma) = \int_{\Sigma} |N_h| d\Sigma,$$

where N_h denotes the projection of the Riemannian unit normal N to \mathcal{H} and $d\Sigma$ denotes the Riemannian area element on Σ . In the sequel we always denote by N the inner unit normal. The singular set Σ_0 is composed of the points in which $T\Sigma$ coincides with \mathcal{H} . Outside Σ_0 , we can define the *horizontal unit normal* as

$$\nu_h := \frac{N_h}{|N_h|}$$

and the *characteristic vector field* as $Z := J(\nu_h)$. It is straightforward to verify that $\{Z, S\}$ is an orthonormal basis of $T\Sigma$ outside Σ_0 , where

$$S := \langle N, T \rangle \nu_h - |N_h| T.$$

Finally, outside Σ_0 , we define the *mean curvature* of Σ by

$$(2-4) \quad H := -\langle \nabla_Z \nu_h, Z \rangle.$$

Given a surface Σ as the zero level set of a function $u : \Omega \subset E(1, 1) \rightarrow \mathbb{R}$, we can express

$$(2-5) \quad \nu_h = -\frac{u_z X + \frac{1}{\sqrt{2}}(-e^z u_x + e^{-z} u_y)Y}{\sqrt{u_z^2 + \frac{1}{2}(-e^z u_x + e^{-z} u_y)^2}}$$

and

$$(2-6) \quad Z = \frac{\frac{1}{\sqrt{2}}(-e^z u_x + e^{-z} u_y)X - u_z Y}{\sqrt{u_z^2 + \frac{1}{2}(-e^z u_x + e^{-z} u_y)^2}}.$$

We define a *minimal surface* as a surface with vanishing mean curvature H .

Proposition 2.1. *Let Σ be a minimal surface defined as the zero level set of a C^2 function $u : \Omega \subset E(1, 1) \rightarrow \mathbb{R}$. Then u satisfies the equation*

$$(2-7) \quad u_{zz}(-e^z u_x + e^{-z} u_y)^2 + u_z^2(-e^{2z} u_{xx} - 2u_{xy} + e^{-2z} u_{yy}) \\ - u_z(-e^z u_x + e^{-z} u_y)(-2e^z u_{xz} - e^z u_x + 2e^{-z} u_{yz} - e^{-z} u_y) = 0$$

on Ω .

Proof. From (2-4), (2-5) and (2-6) we find that u has to satisfy

$$(2-8) \quad Y(u)^2 X(X(u)) - Y(u) X(u) Y(X(u)) - Y(u) X(u) X(Y(u)) + X(u)^2 Y(Y(u)) = 0$$

on Ω . Now, using (2-1), we can transform (2-8) into (2-7). \square

We will call (2-7) the *minimal surface equation*.

Remark 2.2. From (2-8), we immediately note that a surface Σ satisfying $u_z \equiv 0$ or $-e^z u_x + e^{-z} u_y \equiv 0$ is always minimal.

In the following lemma, we compute some important quantities related to the torsion and the geometry of a surface. The lemma follows from [Galli 2013, (9.8)].

Lemma 2.3. *Let Σ be a C^1 surface in $E(1, 1)$. Then we have*

$$\langle \tau(Z), Z \rangle = -\langle Z, X \rangle \langle Z, Y \rangle = \langle v_h, X \rangle \langle v_h, Y \rangle = -\langle \tau(v_h), v_h \rangle, \\ \langle \tau(Z), v_h \rangle = \frac{1}{2}(\langle Z, Y \rangle^2 - \langle Z, X \rangle^2).$$

3. Characteristic curves in $E(1, 1)$

In this section we will study the equation of the integral curves of Z on Σ , known as *characteristic curves*. It is well-known that a surface with constant mean curvature H is foliated by characteristic curves in $\Sigma - \Sigma_0$. In general, a *characteristic curve* is an arc-length parametrized horizontal curve γ in $E(1, 1)$ that satisfies the equation

$$(3-1) \quad \nabla_{\dot{\gamma}} \dot{\gamma} + HJ(\dot{\gamma}) = 0,$$

where $\dot{\gamma}$ denotes the tangent vector along γ and H is the (constant) curvature of γ . We stress that a curve γ satisfying (3-1) is not a subriemannian geodesic. In fact a characteristic curve γ is a subriemannian geodesic if and only if $H = 0$ and $\dot{\gamma}$ satisfies the additional equation

$$(3-2) \quad \langle \tau(\dot{\gamma}), \dot{\gamma} \rangle = 0,$$

see [Rumin 1994, Proposition 15], which forces γ to be an integral curve of X or Y , by Lemma 2.3.

Proposition 3.1. *Let γ be a characteristic curve in $E(1, 1)$ with curvature $H = 0$. Then γ belongs to the family of curves*

$$(3-3) \quad \gamma(t) = (x_0 + \dot{x}_0 t, y_0 + \dot{y}_0 t, z_0)$$

or to the family

$$(3-4) \quad \gamma(t) = \left(x_0 + \frac{\dot{x}_0}{\dot{z}_0} (e^{\dot{z}_0 t} - 1), y_0 - \frac{\dot{y}_0}{\dot{z}_0} (e^{-\dot{z}_0 t} - 1), z_0 + \dot{z}_0 t \right),$$

where $\gamma(0) = (x_0, y_0, z_0)$ and $\dot{\gamma}(0) = (\dot{x}_0, \dot{y}_0, \dot{z}_0)$.

Proof. We consider the curve $\gamma : I \rightarrow \Sigma$, where I denotes an interval. We express $\gamma(t) = (x(t), y(t), z(t))$ and we get

$$(3-5) \quad \begin{aligned} \dot{\gamma}(t) &= \dot{x} \frac{\partial}{\partial x} + \dot{y} \frac{\partial}{\partial y} + \dot{z} \frac{\partial}{\partial z} \\ &= \dot{z} X + \frac{1}{\sqrt{2}} (\dot{y} e^z - \dot{x} e^{-z}) Y + \frac{1}{\sqrt{2}} (\dot{y} e^z + \dot{x} e^{-z}) T, \end{aligned}$$

since

$$\frac{\partial}{\partial x} = \frac{1}{\sqrt{2}} e^{-z} (T - Y), \quad \frac{\partial}{\partial y} = \frac{1}{\sqrt{2}} e^z (Y + T).$$

From (3-5) and the fact that γ is horizontal, we have

$$(3-6) \quad \dot{y} e^z + \dot{x} e^{-z} = 0.$$

Now $\nabla_{\dot{\gamma}} \dot{\gamma} = 0$ is equivalent to the system

$$(3-7) \quad \begin{cases} \dot{z} = \dot{z}_0, \\ \dot{y} e^z - \dot{x} e^{-z} = c_0, \end{cases}$$

where \dot{z}_0 and c_0 are constants. We distinguish two cases. The first one corresponds to $\dot{z}_0 = 0$. This means which $z = z_0$, with $z_0 \in \mathbb{R}$, and so (3-6) and (3-7) reduce to

$$(3-8) \quad \begin{cases} 2\dot{y} = e^{-z_0} c_0, \\ 2\dot{x} = -e^{z_0} c_0, \end{cases}$$

which implies $\gamma(t) = (x_0 - e^{z_0} (c_0/2)t, y_0 + e^{-z_0} (c_0/2)t, z_0)$, where $c_0 \neq 0$ and $x_0, y_0 \in \mathbb{R}$.

The second possibility is $\dot{z}_0 \neq 0$, which implies $z(t) = z_0 + \dot{z}_0 t$, with $z_0 \in \mathbb{R}$. In this case integrating (3-8) we obtain

$$\gamma(t) = \left(x_0 + \frac{c_0 e^{z_0}}{2\dot{z}_0} - \frac{c_0}{2\dot{z}_0} e^{z_0 + \dot{z}_0 t}, y_0 + \frac{c_0 e^{-z_0}}{2\dot{z}_0} - \frac{c_0}{2\dot{z}_0} e^{-(z_0 + \dot{z}_0)t}, z_0 + \dot{z}_0 t \right),$$

where $\gamma(0) = (x_0, y_0, z_0)$. Finally, to conclude the result, we note that

$$\frac{c_0}{2} = \dot{y}_0 e^{z_0} = -\dot{x}_0 e^{-z_0}. \quad \square$$

4. Complete area-stationary surfaces with nonempty singular set in $E(1, 1)$

Complete area-stationary surfaces containing isolated singular points. The local structure of a C^1 surface Σ with prescribed mean curvature $H \in C$, in a neighborhood of an isolated singular point, is well understood [Cheng et al. 2012, Theorem D and Corollary E]. In the less general case of a bounded mean curvature surface of class C^2 , applying [Cheng et al. 2005, Theorem B and Section 7], we have:

Lemma 4.1. *Let Σ be a C^2 oriented immersed surface with constant mean curvature H in $E(1, 1)$. If $p \in \Sigma_0$ is an isolated singular point, then there exists $r > 0$ and $\lambda \in \mathbb{R}$ such that the set*

$$D_r(p) = \{ \gamma_{p,v}^H(s) : v \in T_p \Sigma, |v| = 1, s \in [0, r) \},$$

is an open neighborhood of p in Σ , where $\gamma_{p,v}^H$ denotes the characteristic curve starting from p in the direction v with curvature H .

First we construct the unique example, up to contact isometries, of a minimal surface with isolated singular points.

Proposition 4.2. *Let Σ be a C^2 complete, area-stationary surface immersed in $E(1, 1)$ with $H = 0$ and with an isolated singular point $p_0 = (x_0, y_0, z_0)$. Then $\Sigma = \{(x, y, z) \in E(1, 1) : e^{z-z_0}(y - y_0) + x - x_0 = 0\}$.*

Proof. By Lemma 4.1, the only possible way to construct a complete area-stationary surface, with a singular point p_0 , is to consider the union of all characteristic curves γ of curvature 0 with initial conditions $\gamma(0) = p_0$ and $\dot{\gamma}(0) \in T_{p_0} \Sigma = \mathcal{H}_{p_0}$, $|\dot{\gamma}(0)| = 1$. We can suppose $p_0 = 0$, since $E(1, 1)$ is homogeneous.

We consider the initial velocities

$$\begin{aligned} \dot{\gamma}_\alpha(0) &= \cos \alpha X(0) + \sin \alpha Y(0) \\ &= \cos \alpha \frac{\partial}{\partial z}(0) + \frac{\sin \alpha}{\sqrt{2}} \left(-\frac{\partial}{\partial x}(0) + \frac{\partial}{\partial y}(0) \right), \end{aligned}$$

for $\alpha \in [0, 2\pi[$. In this way we obtain as characteristic curves

$$(4-1) \quad \gamma_\alpha(t) = \left(-\frac{\sin \alpha}{\sqrt{2} \cos \alpha} (e^{\cos(\alpha)t} - 1), -\frac{\sin \alpha}{\sqrt{2} \cos \alpha} (e^{-\cos(\alpha)t} - 1), \cos(\alpha)t \right),$$

for $\alpha \in]0, 2\pi[$ and $\gamma_0(t) = (0, 0, t)$ when $\alpha = 0$. At this point it is easy show that Σ is the zero level set of the function $e^z y + x$ (or equivalently $e^{-z} x + y$), which satisfies (2-7). □

Complete area-stationary surfaces containing singular curves.

Lemma 4.3 [Galli 2013, Corollary 5.4]. *Let Σ be a C^2 minimal surface with nonempty singular set Σ_0 immersed in $E(1, 1)$. Then Σ is area stationary if and only if the characteristic curves meet the singular curves orthogonally with respect to the metric $\langle \cdot, \cdot \rangle$, induced by the orthonormal basis (2-1).*

A minimal area-stationary surface cannot contain more than one singular curve:

Lemma 4.4. *Let Σ be a C^2 complete, minimal, area-stationary surface, containing a singular curve Γ immersed in $E(1, 1)$. Then Σ cannot contain more singular curves.*

Proof. We consider a singular curve

$$\Gamma(\varepsilon) = (x(\varepsilon), y(\varepsilon), z(\varepsilon))$$

in Σ . Since Σ is foliated by characteristic curves, we can parametrize it by the map

$$F(\varepsilon, t) = \gamma_\varepsilon(t) = (x(\varepsilon, t), y(\varepsilon, t), z(\varepsilon, t)),$$

where $\gamma_\varepsilon(t)$ is the characteristic curve with initial data $\gamma_\varepsilon(0) = \Gamma(\varepsilon)$ and

$$(4-2) \quad \begin{aligned} \dot{\gamma}_\varepsilon(0) &= J(\dot{\Gamma}(\varepsilon)) = \dot{z}(\varepsilon)J(X) + \frac{1}{\sqrt{2}}(\dot{y}(\varepsilon)e^{z(\varepsilon)} - \dot{x}(\varepsilon)e^{-z(\varepsilon)})J(Y) \\ &= \frac{1}{\sqrt{2}}(-\dot{z}(\varepsilon)e^{z(\varepsilon)}, \dot{z}(\varepsilon)e^{-z(\varepsilon)}, \dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)}). \end{aligned}$$

We define

$$V_\varepsilon(t) := \frac{\partial F}{\partial \varepsilon}(t, \varepsilon),$$

which is a smooth Jacobi-like vector field along $\gamma_\varepsilon(t)$; see [Galli 2013, Section 4]. At a singular point (ε, t) , the vertical component of V_ε vanishes:

$$\langle V_\varepsilon, T \rangle(\varepsilon, t) = \frac{\partial x}{\partial \varepsilon}(\varepsilon, t)e^{-z(\varepsilon, t)} + \frac{\partial y}{\partial \varepsilon}(\varepsilon, t)e^{z(\varepsilon, t)} = 0.$$

We suppose that Γ is not an integral curve of X or Y . Then from the expressions of the component of $F(\varepsilon, t)$, which are

$$(4-3) \quad \begin{aligned} x(\varepsilon, t) &= x(\varepsilon) + \frac{\dot{z}(\varepsilon)e^{z(\varepsilon)}}{\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)}} (e^{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})t/\sqrt{2}} - 1), \\ y(\varepsilon, t) &= y(\varepsilon) - \frac{\dot{z}(\varepsilon)e^{-z(\varepsilon)}}{\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)}} (e^{-(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})t/\sqrt{2}} - 1), \\ z(\varepsilon, t) &= z(\varepsilon) + \frac{\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)}}{\sqrt{2}}t, \end{aligned}$$

we have

$$\begin{aligned} & \langle V_\varepsilon, T \rangle(\varepsilon, t) \\ &= \left(\dot{x}(\varepsilon)e^{-z(\varepsilon)} + \frac{\ddot{z}(\varepsilon)}{\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)}} - \frac{\dot{z}(\varepsilon) \frac{\partial}{\partial \varepsilon} (\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})}{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})^2} \right) \\ & \quad \cdot \left(e^{-\frac{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})t}{\sqrt{2}}} - e^{\frac{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})t}{\sqrt{2}}} \right) \\ & \quad + \frac{\dot{z}(\varepsilon)^2}{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})^2} \left(e^{-\frac{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})t}{\sqrt{2}}} + e^{\frac{(\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)})t}{\sqrt{2}}} - 2 \right), \end{aligned}$$

which, when t is positive, vanishes only for the values $(\varepsilon, 0)$. On the other hand, if Γ is an integral curve of Y we get

$$(4-4) \quad x(\varepsilon, t) = x(\varepsilon), \quad y(\varepsilon, t) = y(\varepsilon), \quad z(\varepsilon, t) = z(\varepsilon) + \frac{\dot{x}(\varepsilon)e^{-z(\varepsilon)} - \dot{y}(\varepsilon)e^{z(\varepsilon)}}{\sqrt{2}}t,$$

and if Γ is an integral curve of X we have

$$(4-5) \quad x(\varepsilon, t) = x(\varepsilon) - \frac{\dot{z}(\varepsilon)e^{z(\varepsilon)}}{\sqrt{2}}t, \quad y(\varepsilon, t) = y(\varepsilon) + \frac{\dot{z}(\varepsilon)e^{-z(\varepsilon)}}{\sqrt{2}}t, \quad z(\varepsilon, t) = z(\varepsilon).$$

In both cases, the singular set is only the curve $\Gamma(\varepsilon)$. \square

The vertical component of V_ε can be computed more directly using [Galli 2013, Proposition 4.3], since $H = 0$. On the other hand, the explicit computation of the components of the parametrization $F(\varepsilon, t)$ allows us to characterize all C^2 area-stationary complete surfaces with a singular curve that is a characteristic curve of curvature 0. We stress that, when the characteristic curves are subriemannian geodesics, these examples can also be constructed from Remark 2.2.

Proposition 4.5. *Let Σ be an area-stationary surface with $H = 0$, with a singular curve Γ that is a characteristic curve of curvature 0. Then, if Γ is a subriemannian geodesic, Σ belongs to one of the following families:*

- (i) $\{ax + by + c = 0 : (x, y, z) \in E(1, 1), a, b, c \in \mathbb{R}\}$;
- (ii) $\{e^{z-z_0}(y - y_0) + e^{z_0-z}(x - x_0) = 0 : (x, y, z) \in E(1, 1), x_0, y_0, z_0 \in \mathbb{R}\}$.

Otherwise, we suppose that Γ is a characteristic curve passing through (x_0, y_0, z_0) with velocity $(\dot{x}_0, \dot{y}_0, \dot{z}_0)$, $\dot{x}_0, \dot{y}_0, \dot{z}_0 \neq 0$. We can parametrize Σ by $F : \mathbb{R}^2 \rightarrow E(1, 1)$, with $F(\varepsilon, t) = (x(\varepsilon, t), y(\varepsilon, t), z(\varepsilon, t))$ and

$$\begin{aligned}
 x(\varepsilon, t) &= x_0 + \frac{\dot{x}_0}{\dot{z}_0}(e^{\dot{z}_0\varepsilon} - 1) + \frac{\dot{z}_0 e^{\dot{z}_0\varepsilon} + \dot{z}_0}{\dot{x}_0 e^{-\dot{z}_0\varepsilon} - \dot{y}_0 e^{\dot{z}_0\varepsilon}}(e^{(\dot{x}_0 e^{-\dot{z}_0\varepsilon} - \dot{y}_0 e^{\dot{z}_0\varepsilon})t/\sqrt{2}} - 1), \\
 (4-6) \quad y(\varepsilon, t) &= y_0 - \frac{\dot{y}_0}{\dot{z}_0}(e^{-\dot{z}_0\varepsilon} - 1) - \frac{\dot{z}_0 e^{-\dot{z}_0\varepsilon} - \dot{z}_0}{\dot{x}_0 e^{-\dot{z}_0\varepsilon} - \dot{y}_0 e^{\dot{z}_0\varepsilon}}(e^{-(\dot{x}_0 e^{-\dot{z}_0\varepsilon} - \dot{y}_0 e^{\dot{z}_0\varepsilon})t/\sqrt{2}} - 1), \\
 z(\varepsilon, t) &= z_0 + \dot{z}_0\varepsilon + \frac{\dot{x}_0 e^{-\dot{z}_0\varepsilon} - \dot{y}_0 e^{\dot{z}_0\varepsilon}}{\sqrt{2}}t.
 \end{aligned}$$

Remark 4.6. The surfaces parametrized by (4-6) are the first examples of area-stationary surfaces that are not foliated by subriemannian geodesics in three-dimensional contact subriemannian manifolds, up to our knowledge. In fact this phenomenon does not appear in the group of rigid motions [Galli 2013, Lemma 10.4], even if its pseudohermitian torsion is nonvanishing. In that case, the presence of two singular curves forces the surface to be foliated by subriemannian geodesics or to be not area-stationary. On the other hand, it is well-known that a minimal surface is foliated by subriemannian geodesics in any three-dimensional Sasakian manifold.

Remark 4.7. Given any horizontal curve $\Gamma = (x(\varepsilon), y(\varepsilon), z(\varepsilon))$ in $E(1, 1)$, we stress that (4-3) provides a parametrization $F(\varepsilon, t) : \mathbb{R}^2 \rightarrow \Sigma \subset E(1, 1)$ of a complete area-stationary surface Σ with $\Sigma_0 = \Gamma$.

5. Complete area-minimizing surfaces in $E(1, 1)$

Complete area-minimizing surfaces with empty singular set. Proposition 9.8 of [Galli 2013] gave a general necessary condition for the stability of a nonsingular surface in pseudohermitian Lie groups. This condition states that the quantity

$$W - \langle \tau(Z), v_h \rangle = \langle v_h, Y \rangle^2 - 1 = \langle Z, X \rangle^2 - 1$$

must be always nonpositive. This condition is trivial in $E(1, 1)$ due to the negativity of the Webster scalar curvature. On the other hand it has been used crucially in the classification of the stable, area-stationary surfaces without singular points in the manifolds \mathbb{H}^1 , $SU(2)$ and $\tilde{E}(2)$, see [Galli 2013; Hurtado et al. 2010; Rosales 2012]. In any case, we can prove:

Proposition 5.1. *The families of planes*

- (i) $\{x + c = 0 : (x, y, z) \in E(1, 1), c \in \mathbb{R}\}$,
- (ii) $\{y + c = 0 : (x, y, z) \in E(1, 1), c \in \mathbb{R}\}$,
- (iii) $\{z + c = 0 : (x, y, z) \in E(1, 1), c \in \mathbb{R}\}$,

are area-stationary, foliated by subriemannian geodesics, and area-minimizing.

Proof. We prove the result for $\Sigma = \{x = 0 : (x, y, z) \in E(1, 1)\}$, since all the cases are similar. In this case, from (2-5) and (2-6) we have

$$v_h = Y, \quad Z = -X.$$

So the integral curves of Z are subriemannian geodesics and $\Sigma_0 = \emptyset$. Now Remark 2.2 implies that Σ is area-stationary. Finally we can foliate a neighborhood of Σ in $E(1, 1)$ by translating Σ . We obtain a foliation by area-stationary surfaces, and a standard calibration argument implies that Σ is area-minimizing; see, for example, [Barone Adesi et al. 2007; Ritoré 2009; Ritoré and Rosales 2008, § 5]. \square

Remark 5.2. The planes in the family

$$\{ax + by + cz + d = 0 : (x, y, z) \in E(1, 1), a, b, c, d \in \mathbb{R}\}$$

are not minimal, since they do not satisfy (2-7).

A very natural question is: are the planes in Proposition 5.1 the unique complete area-minimizing surfaces with empty singular set in $E(1, 1)$? We have only been able to find the following sufficient condition:

Lemma 5.3. *Let Σ be a C^2 complete oriented minimal surface immersed in $E(1, 1)$, with empty singular set Σ_0 . If $\langle N, T \rangle \leq 0$ holds on Σ , then Σ is stable.*

Proof. Taking into account the expression of the stability operator for nonsingular surfaces in [Galli 2013, Lemma 8.3], we only need to show that

$$2Z(G) + G^2 \leq 0 \quad \text{on } \Sigma, \quad \text{where } G := \frac{\langle N, T \rangle}{|N_h|}.$$

Given a point p in Σ , let I be an open interval containing the origin and let $\alpha : I \rightarrow \Sigma$ be a piece of the integral curve of S passing through p . Consider the characteristic curve $\gamma_\varepsilon(s)$ of Σ with $\gamma_\varepsilon(0) = \alpha(\varepsilon)$. We define the map $F : I \times \mathbb{R} \rightarrow \Sigma$ by $F(\varepsilon, s) = \gamma_\varepsilon(s)$ and set $V(s) := (\partial F / \partial \varepsilon)(0, s)$, which is a Jacobi-like vector field along γ_0 ; see [ibid., Proposition 4.3]. Let $'$ represent differentiation with respect to s . Using [ibid., Lemma 3.1, (4.4) and (4.5)] we get

$$(5-1) \quad \langle V, T \rangle(0) = -|N_h|,$$

$$(5-2) \quad \langle V, T \rangle'(0) = -\langle N, T \rangle,$$

$$(5-3) \quad \langle V, T \rangle''(0) = -|N_h|(Z(G) + G^2).$$

It is easy to show that $g(V, T)$ never vanishes along γ_0 since Σ_0 is empty; see [ibid., proof of Proposition 9.5]. On the other hand, by [ibid., Proposition 4.3] and Lemma 2.3, we have that $\langle V, T \rangle$ satisfies the ordinary differential equation

$$\langle V, T \rangle'''(s) - \langle Z, X \rangle^2 \langle V, T \rangle'(s) = 0$$

along γ_0 . We suppose that $\langle Z, X \rangle \neq 0$. Taking into account the initial conditions (5-1), (5-2) and (5-3), we obtain

$$\langle V, T \rangle(s) = a \cosh(|\langle Z, X \rangle|s) + b \sinh(|\langle Z, X \rangle|s) + c,$$

where

$$a = \frac{|N_h|(Z(G) + G^2)}{\langle X, Z \rangle^2}, \quad b = -\frac{\langle N, T \rangle}{|\langle Z, X \rangle|}, \quad c = -|N_h| - a.$$

We have that $\langle V, T \rangle(s) \neq 0$ implies

$$a + b = \frac{|N_h|(Z(G) + G^2)}{\langle X, Z \rangle^2} - \frac{\langle N, T \rangle}{|\langle Z, X \rangle|} \leq 0.$$

Then we can conclude that

$$2Z(G) + G^2 \leq 2(Z(G) + G^2) \leq 2|\langle Z, X \rangle| \frac{\langle N, T \rangle}{|N_h|} \leq 0$$

on γ_0 . Now since the choice of p is arbitrary, we get the statement.

If $\langle Z, X \rangle = 0$, we conclude that Σ is stable if and only if $\langle N, T \rangle = 0$, by [Galli 2013, Proposition 9.8]. \square

Remark 5.4. The surfaces described in the points (i), (ii) and (iii) of Proposition 5.1 are characterized by $\langle N, T \rangle = -e^z/\sqrt{2}$, $\langle N, T \rangle = -e^z/\sqrt{2}$ and $\langle N, T \rangle \equiv 0$, respectively, where N denotes the inward unit normal on Σ . In the third family the planes are vertical surfaces and they satisfy $W - \langle \tau(Z), \nu_h \rangle \equiv 0$.

Taking into account the geometric invariants of $E(1, 1)$, we expect the existence of other examples of complete oriented minimal surface with empty singular set.

Complete area-minimizing surfaces with nonempty singular set. We consider the stability operator constructed in [Galli 2013, Theorem 8.6].

Lemma 5.5. *Let Σ be a C^2 oriented minimal surface immersed in $E(1, 1)$, with singular set Σ_0 and $\partial\Sigma = \emptyset$. If Σ is stable then, for any function $u \in C_0^1(\Sigma)$ such that $Z(u) = 0$ in a tubular neighborhood of a singular curve and constant in a tubular neighborhood of an isolated singular point, we have $Q(u) \geq 0$, where*

$$Q(u) := \int_{\Sigma} \{ |N_h|^{-1} Z(u)^2 + |N_h| ((1 + \langle Z, Y \rangle^2) - (|N_h|(\frac{1}{2} - \langle Z, Y \rangle^2) - \langle \nabla_S \nu_h, Z \rangle)^2) u^2 \} d\Sigma + 4 \int_{(\Sigma_0)_c} \langle N, T \rangle \langle Z, Y \rangle^2 \langle Z, \nu \rangle u^2 d(\Sigma_0)_c + \int_{(\Sigma_0)_c} S(u)^2 d(\Sigma_0)_c.$$

Here $d(\Sigma_0)_c$ is the Riemannian length measure on $(\Sigma_0)_c$ and ν is the external unit normal to $(\Sigma_0)_c$.

Corollary 5.6. *Let Σ be a plane in the family*

$$\{ax + by + c = 0 : (x, y, z) \in E(1, 1), a, b, c \in \mathbb{R}\}.$$

Then Σ is stable.

Proof. We know that Σ is area-stationary with a singular line, obtained intersecting Σ with the plane $z = \log \sqrt{b/a}$. From (2-6) we get

$$Z = \frac{-be^z + ae^{-z}}{|-be^z + ae^{-z}|} X,$$

which is orthogonal to the singular line. Since $\langle \nabla_S v_h, Z \rangle = \langle \nabla_S Y, X \rangle = \frac{|N_h|}{2}$, the stability operator

$$Q(u) = \int_{\Sigma} \{|N_h|^{-1} Z(u)^2 + |N_h| \langle N, T \rangle^2 u^2\} d\Sigma + \int_{\Sigma_0} S(u)^2 d\Sigma_0$$

is always nonnegative for any admissible test function u . □

Remark 5.7. The planes $\{ax + by + c = 0 : (x, y, z) \in E(1, 1), a, b, c \in \mathbb{R}\}$ are also area-minimizing, by calibration arguments.

Corollary 5.8. *The surface $\Sigma = \{e^z y + e^{-z} x = 0 : (x, y, z) \in E(1, 1)\}$ is stable.*

Proof. From (2-6) we get

$$Z = -\frac{(e^z y - e^{-z} x)Y}{|e^z y - e^{-z} x|}$$

and $\Sigma_0 = \{(0, 0, z) : (x, y, z) \in E(1, 1)\}$. From (2-3) we have

$$\langle \nabla_S v_h, Z \rangle = \langle \nabla_S Y, X \rangle = -\frac{|N_h|}{2},$$

which implies

$$Q(u) = \int_{\Sigma} \{|N_h|^{-1} Z(u)^2 + 2|N_h|^2 u^2\} d\Sigma + \int_{\Sigma_0} S(u)^2 d\Sigma_0 + 4 \int_{\Sigma_0} u^2 d\Sigma_0 \geq 0,$$

for all admissible u . □

Corollary 5.9. *The surfaces defined in Proposition 4.2 are stable.*

Proof. For simplicity we will prove the statement in the case of $x_0 = y_0 = z_0 = 0$. We note that, since $\Sigma_0 = (0, 0, 0)$, the argument in the proof of Lemma 5.3 works and the condition $\langle N, T \rangle = -(1 + e^z)/\sqrt{2} \leq 0$ is sufficient for the stability in the complement of any tubular neighborhood of Σ_0 . Finally we observe that the stability operator in Lemma 5.5 makes no contribution to the singular set in the case of isolated singular points. □

References

[Ambrosio et al. 2006] L. Ambrosio, F. Serra Cassano, and D. Vittone, “Intrinsic regular hypersurfaces in Heisenberg groups”, *J. Geom. Anal.* **16**:2 (2006), 187–232. MR 2007g:49072 Zbl 1085.49045

[Barbieri and Citti 2011] D. Barbieri and G. Citti, “Regularity of minimal intrinsic graphs in 3-dimensional sub-Riemannian structures of step 2”, *J. Math. Pures Appl.* (9) **96**:3 (2011), 279–306. MR 2012g:53050 Zbl 1231.53030

- [Barone Adesi et al. 2007] V. Barone Adesi, F. Serra Cassano, and D. Vittono, “The Bernstein problem for intrinsic graphs in Heisenberg groups and calibrations”, *Calc. Var. Partial Differential Equations* **30**:1 (2007), 17–49. MR 2009c:35044 Zbl 1206.35240
- [Bigolin and Cassano 2010] F. Bigolin and F. S. Cassano, “Distributional solutions of Burgers’ equation and intrinsic regular graphs in Heisenberg groups”, *J. Math. Anal. Appl.* **366**:2 (2010), 561–568. MR 2012a:35335 Zbl 1186.35030
- [Boothby and Wang 1958] W. M. Boothby and H. C. Wang, “On contact manifolds”, *Ann. of Math.* (2) **68** (1958), 721–734. MR 22 #3015 Zbl 0084.39204
- [Capogna et al. 2007] L. Capogna, D. Danielli, S. D. Pauls, and J. T. Tyson, *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*, Progress in Mathematics **259**, Birkhäuser, Basel, 2007. MR 2009a:53053 Zbl 1138.53003
- [Capogna et al. 2009] L. Capogna, G. Citti, and M. Manfredini, “Regularity of non-characteristic minimal graphs in the Heisenberg group \mathbb{H}^1 ”, *Indiana Univ. Math. J.* **58**:5 (2009), 2115–2160. MR 2010j:58032 Zbl 1182.35087
- [Cheng and Hwang 2010] J.-H. Cheng and J.-F. Hwang, “Variations of generalized area functionals and p -area minimizers of bounded variation in the Heisenberg group”, *Bull. Inst. Math. Acad. Sin. (N.S.)* **5**:4 (2010), 369–412. MR 2012g:49085 Zbl 1232.35178
- [Cheng et al. 2005] J.-H. Cheng, J.-F. Hwang, A. Malchiodi, and P. Yang, “Minimal surfaces in pseudo-hermitian geometry”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **4**:1 (2005), 129–177. MR 2006f:53008 Zbl 1158.53306
- [Cheng et al. 2007] J.-H. Cheng, J.-F. Hwang, and P. Yang, “Existence and uniqueness for p -area minimizers in the Heisenberg group”, *Math. Ann.* **337**:2 (2007), 253–293. MR 2009h:35120 Zbl 1109.35009
- [Cheng et al. 2012] J.-H. Cheng, J.-F. Hwang, A. Malchiodi, and P. Yang, “A Codazzi-like equation and the singular set for C^1 smooth surfaces in the Heisenberg group”, *J. Reine Angew. Math.* **671** (2012), 131–198. MR 2983199 Zbl 1255.53026
- [Danielli et al. 2007] D. Danielli, N. Garofalo, and D. M. Nhieu, “Sub-Riemannian calculus on hyper-surfaces in Carnot groups”, *Adv. Math.* **215**:1 (2007), 292–378. MR 2009h:53061 Zbl 1129.53017
- [Danielli et al. 2008] D. Danielli, N. Garofalo, and D. M. Nhieu, “A notable family of entire intrinsic minimal graphs in the Heisenberg group which are not perimeter minimizing”, *Amer. J. Math.* **130**:2 (2008), 317–339. MR 2009b:49102 Zbl 1158.53334
- [Danielli et al. 2009] D. Danielli, N. Garofalo, D. M. Nhieu, and S. D. Pauls, “Instability of graphical strips and a positive answer to the Bernstein problem in the Heisenberg group \mathbb{H}^1 ”, *J. Differential Geom.* **81**:2 (2009), 251–295. MR 2010e:53007 Zbl 1161.53024
- [Dragomir and Tomassini 2006] S. Dragomir and G. Tomassini, *Differential geometry and analysis on CR manifolds*, Progress in Mathematics **246**, Birkhäuser, Boston, MA, 2006. MR 2007b:32056 Zbl 1099.32008
- [Galli 2012] M. Galli, *Area-stationary surfaces in contact sub-Riemannian manifolds*, Ph.D. thesis, Universidad de Granada, 2012, Available at <http://hera.ugr.es/tesisugr/21013020.pdf>.
- [Galli 2013] M. Galli, “First and second variation formulae for the sub-Riemannian area in three-dimensional pseudo-Hermitian manifolds”, *Calc. Var. Partial Differential Equations* **47**:1-2 (2013), 117–157. MR 3044134 Zbl 1268.53039
- [Galli and Ritoré 2013] M. Galli and M. Ritoré, “Existence of isoperimetric regions in contact sub-Riemannian manifolds”, *J. Math. Anal. Appl.* **397**:2 (2013), 697–714. MR 2979606 Zbl 06109993

- [Garofalo and Nhieu 1996] N. Garofalo and D.-M. Nhieu, “Isoperimetric and Sobolev inequalities for Carnot–Carathéodory spaces and the existence of minimal surfaces”, *Comm. Pure Appl. Math.* **49**:10 (1996), 1081–1144. MR 97i:58032 Zbl 0880.35032
- [Hladky and Pauls 2008] R. K. Hladky and S. D. Pauls, “Constant mean curvature surfaces in sub-Riemannian geometry”, *J. Differential Geom.* **79**:1 (2008), 111–139. MR 2009m:53070 Zbl 1156.53038
- [Hurtado and Rosales 2008] A. Hurtado and C. Rosales, “Area-stationary surfaces inside the sub-Riemannian three-sphere”, *Math. Ann.* **340**:3 (2008), 675–708. MR 2008i:53038 Zbl 1132.53015
- [Hurtado and Rosales \geq 2014] A. Hurtado and C. Rosales, “Stable surfaces inside the sub-Riemannian three-sphere”. In preparation.
- [Hurtado et al. 2010] A. Hurtado, M. Ritoré, and C. Rosales, “The classification of complete stable area-stationary surfaces in the Heisenberg group \mathbb{H}^1 ”, *Adv. Math.* **224**:2 (2010), 561–600. MR 2011d:53054 Zbl 1192.53038
- [Milnor 1976] J. Milnor, “Curvatures of left invariant metrics on Lie groups”, *Advances in Math.* **21**:3 (1976), 293–329. MR 54 #12970 Zbl 0341.53030
- [Perrone 1998] D. Perrone, “Homogeneous contact Riemannian three-manifolds”, *Illinois J. Math.* **42**:2 (1998), 243–256. MR 99a:53067 Zbl 0906.53031
- [Ritoré 2009] M. Ritoré, “Examples of area-minimizing surfaces in the sub-Riemannian Heisenberg group \mathbb{H}^1 with low regularity”, *Calc. Var. Partial Differential Equations* **34**:2 (2009), 179–192. MR 2009h:53062 Zbl 1165.53023
- [Ritoré and Rosales 2008] M. Ritoré and C. Rosales, “Area-stationary surfaces in the Heisenberg group \mathbb{H}^1 ”, *Adv. Math.* **219**:2 (2008), 633–671. MR 2009h:49075 Zbl 1158.53022
- [Rosales 2012] C. Rosales, “Complete stable CMC surfaces with empty singular set in Sasakian sub-Riemannian 3-manifolds”, *Calc. Var. Partial Differential Equations* **43**:3-4 (2012), 311–345. MR 2875642 Zbl 1235.53036
- [Rumin 1994] M. Rumin, “Formes différentielles sur les variétés de contact”, *J. Differential Geom.* **39**:2 (1994), 281–330. MR 95g:58221 Zbl 0973.53524
- [Shcherbakova 2009] N. Shcherbakova, “Minimal surfaces in sub-Riemannian manifolds and structure of their singular sets in the (2, 3) case”, *ESAIM Control Optim. Calc. Var.* **15**:4 (2009), 839–862. MR 2011c:53055 Zbl 1190.53027

Received May 27, 2013. Revised August 9, 2013.

MATTEO GALLI
DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA
UNIVERSIDAD DE GRANADA
18071 GRANADA
SPAIN
galli@ugr.es

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2014 is US \$410/year for the electronic version, and \$535/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 271 No. 1 September 2014

Proper holomorphic maps between bounded symmetric domains revisited	1
GAUTAM BHARALI and JAIKRISHNAN JANARDHANAN	
An explicit Majorana representation of the group $3^2:2$ of $3C$ -pure type	25
HSIAN-YANG CHEN and CHING HUNG LAM	
Sofic groups: graph products and graphs of groups	53
LAURA CIOBANU, DEREK F. HOLT and SARAH REES	
Perturbations of a critical fractional equation	65
EDUARDO COLORADO, ARTURO DE PABLO and URKO SÁNCHEZ	
A density theorem in parametrized differential Galois theory	87
THOMAS DREYFUS	
On the classification of complete area-stationary and stable surfaces in the subriemannian Sol manifold	143
MATTEO GALLI	
Periodic orbits of Hamiltonian systems linear and hyperbolic at infinity	159
BAŞAK Z. GÜREL	
Nonsplittability of the rational homology cobordism group of 3-manifolds	183
SE-GOO KIM and CHARLES LIVINGSTON	
Biharmonic surfaces of constant mean curvature	213
ERIC LOUBEAU and CEZAR ONICIUC	
Foliations of a smooth metric measure space by hypersurfaces with constant f -mean curvature	231
JUNCHEOL PYO	
On the existence of large degree Galois representations for fields of small discriminant	243
JEREMY ROUSE and FRANK THORNE	



0030-8730(201409)271:1;1-1