

*Pacific
Journal of
Mathematics*

**SPANNING TREES AND RANDOM WALKS
ON WEIGHTED GRAPHS**

XIAO CHANG, HAO XU AND SHING-TUNG YAU

Volume 273 No. 1

January 2015

SPANNING TREES AND RANDOM WALKS ON WEIGHTED GRAPHS

XIAO CHANG, HAO XU AND SHING-TUNG YAU

Using two graph invariants arising from Chung and Yau's discrete Green's function, we derive explicit formulas and new estimates of hitting times of random walks on weighted graphs through the enumeration of spanning trees.

1. Introduction

Every reversible Markov chain can be viewed as a random walk on a weighted undirected graph $G = (V, E)$ with edge weights w_{xy} . We may assume that G has no multiedges but may have a loop of weight w_{xx} at each vertex x . The weighted degree d_x of x is the sum of all w_{xy} , $y \in V$. The *volume* of a graph is $\text{vol}(G) = \sum_{v \in V} d_v$.

The *Laplacian* of G is the matrix $L = D - A$, where D is the diagonal matrix whose entries are d_x , $x \in V$ and A is the adjacency matrix of G . *Chung's normalized Laplacian*, $\mathcal{L} = D^{-1/2} L D^{-1/2}$, is

$$\mathcal{L}(x, y) = \begin{cases} 1 - w_{xx}/d_x & \text{if } x = y, \\ -w_{xy}/\sqrt{d_x d_y} & \text{if } x \sim y, \\ 0 & \text{otherwise,} \end{cases}$$

where $x \sim y$ denotes that x and y are adjacent. Here we assume that $d_x \neq 0$ for all $x \in V$, since for a random walk it is natural to impose that G is connected and $w_{xy} > 0$ for all $xy \in E$.

If G is connected, denote by $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{n-1}$ the eigenvalues of \mathcal{L} with the corresponding orthonormal basis of eigenvectors $\phi_0, \phi_1, \dots, \phi_{n-1}$, as $n \times 1$ column vectors. Obviously $\phi_0(x) = \sqrt{d_x / \text{vol}(G)}$.

Chung and Yau [2000] defined the *discrete Green's function* \mathcal{G} by

$$\mathcal{G} = \sum_{j=1}^{n-1} \frac{1}{\lambda_j} \phi_j \phi_j^*,$$

MSC2010: 05C81.

Keywords: hitting time, random walk, spanning tree.

which is uniquely determined by the relations $\mathcal{G}\mathcal{L} = \mathcal{L}\mathcal{G} = I - P_0$ and $\mathcal{G}P_0 = 0$, where $P_0 = \phi_0\phi_0^t$ is an $n \times n$ matrix.

A random walk on G is a Markov chain on V with transition probability matrix $(p_{xy})_{x,y \in V}$, where

$$p_{xy} = \begin{cases} w_{xy}/d_x & \text{if } x \sim y, \\ 0 & \text{otherwise.} \end{cases}$$

The *hitting time* $H(x, y)$ is the expected number of steps to reach vertex y when started from vertex x . By using a result in [Aldous and Fill 2014], Chung and Yau proved an expression for $H(x, y)$ in terms of the discrete Green’s function:

Theorem 1.1 [Chung and Yau 2000]. *On a connected graph G , the hitting time $H(x, y)$ and Green’s function $\mathcal{G}(x, y)$ satisfy*

$$(1) \quad H(x, y) = \text{vol}(G) \left(\frac{\mathcal{G}(y, y)}{d_y} - \frac{\mathcal{G}(x, y)}{\sqrt{d_x d_y}} \right).$$

For a weighted graph G , we denote by $\Omega(G)$ the set of spanning trees of G . For $T \in \Omega(G)$, define the weight $w(T)$ of T to be $\prod_{e \in T} w_e$. Let $\tau(G)$ be the weighted counting of spanning trees:

$$(2) \quad \tau(G) = \sum_{T \in \Omega(G)} w(T).$$

Below is a typical identity expressing hitting times in terms of spanning trees (see Theorem 2.11) arising from our study of Chung and Yau’s discrete Green’s function.

Theorem 1.2. *Let G be a connected weighted graph and $x, y \in V(G)$. Then*

$$(3) \quad H(x, y) = \frac{1}{\tau(G)} \sum_{u \in V(G)} d_u \sum_{\substack{P \in \mathcal{P}_G(x,u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}).$$

The paper is organized as follows. In Section 2, we introduce two graph invariants and use them to derive explicit formulas for Chung and Yau’s discrete Green’s functions and hitting times of random walks on weighted graphs. In Section 3, we apply our formulas to obtain various estimates of hitting times on weighted graphs. In Section 4, we prove an explicit formula for hitting times of random walks on infinite trees. In Section 5, we apply our work to improve estimates of hitting times under different weight schemes on a given simple finite graph.

2. The hitting time of random walks on weighted graphs

Kirchhoff discovered his matrix-tree theorem in 1847 in his work on electrical networks, and this theorem gives an efficient way to calculate $\tau(G)$ using linear algebra.

Theorem 2.1 (Kirchhoff’s matrix-tree theorem). *Let G be a connected weighted graph.*

(i) *If the Laplacian L of G has eigenvalues $0 = \mu_0 < \mu_1 \leq \dots \leq \mu_{n-1}$, then*

$$(4) \quad \prod_{k=1}^{n-1} \mu_k = n\tau(G).$$

(ii) *Let L_{ij} be the matrix obtained from L by deleting the i -th row and j -th column. Then all cofactors $(-1)^{i+j} \det(L_{ij})$ of L are equal and*

$$(5) \quad \det L_{ii} = n\tau(G) \quad \text{for all } 1 \leq i \leq n.$$

We also need the following version of Kirchhoff’s matrix-tree theorem for weighted graphs. A proof, with slight changes, can be found in the cited reference.

Theorem 2.2 [Chung 2011, Theorem 1]. *For a connected weighted graph $G = (V, E)$, we have*

$$(6) \quad \prod_{k=1}^{n-1} \lambda_k = \frac{\text{vol}(G)\tau(G)}{\prod_{v \in V} d_v},$$

where $0 = \lambda_0 < \lambda_1 \leq \dots \leq \lambda_{n-1}$ are eigenvalues of \mathcal{L} and $\tau(G)$ is defined in (2).

A weighted graph is a graph G equipped with a function $w : E(G) \rightarrow \mathbb{R}_+$ that assigns a positive number to each edge. In the following, we fix a weighted graph (G, w) and introduce two invariants for any induced subgraph S of G . Consider the matrix

$$(7) \quad B(x, y) = \begin{cases} d_x^2 s + d_x - w_{xx} & \text{if } x = y, \\ d_x d_y s - w_{xy} & \text{if } x \sim y, \\ d_x d_y s & \text{otherwise,} \end{cases}$$

where $d_x = w_{xx} + \sum_{y \sim x} w_{xy}$. Denote by B_S the principle submatrix of B on indices corresponding to the vertices of S ; we define $R(S)$ and $Z(S)$ by

$$\det B_S = R(S) + Z(S) \cdot s.$$

In particular, $R(\emptyset) = 1$, $Z(\emptyset) = 0$ for the empty subgraph \emptyset , and $R(\{x\}) = d_x - w_{xx}$, $Z(\{x\}) = d_x^2$ for any $x \in V(G)$.

In the following four lemmas, S is an arbitrary induced subgraph of a fixed weighted graph (G, w) . The proofs of these lemmas are similar to those in [Xu and Yau 2013a, Section 2], where we considered unweighted simple graphs with $w_e \equiv 1$ for all $e \in E(G)$.

Lemma 2.3. *If S has k connected components S_1, \dots, S_k , then*

$$R(S) = \prod_{i=1}^k R(S_i), \quad Z(S) = \sum_{i=1}^k Z(S_i) \prod_{\substack{j=1 \\ j \neq i}}^k R(S_j).$$

Lemma 2.4. *For any fixed vertex $x \in V(S)$, we have*

$$R(S) = (d_x - w_{xx})R(S - \{x\}) - \sum_{\substack{y \in V(S) \\ y \sim x}} w_{xy} \sum_{P \in \mathcal{P}_S(x,y)} \prod_{e \in E(P)} w_e R(S - \{P\}),$$

and

$$\begin{aligned} Z(S) &= (d_x - w_{xx})Z(S - \{x\}) \\ &\quad - \sum_{\substack{y \in V(S) \\ y \sim x}} w_{xy} \sum_{P \in \mathcal{P}_S(x,y)} \prod_{e \in E(P)} w_e Z(S - \{P\}) + (d_x - w_{xx})^2 R(S - \{x\}) \\ &\quad + \sum_{\substack{u,v \in V(S) \\ u \neq v}} d_u d_v \sum_{\substack{P_1 \in \mathcal{P}_S(x,u) \\ P_2 \in \mathcal{P}_S(x,v) \\ P_1 \cap P_2 = x}} \prod_{e \in E(P_1 \cup P_2)} w_e R(S - \{P_1, P_2\}), \end{aligned}$$

where $\mathcal{P}_S(x, y)$ is the set of all simple paths (with no repeated vertices) connecting x and y in S . We assume that $\mathcal{P}_S(x, x)$ consists of the trivial path $\{x\}$ only. Here $S - \{P\}$ means the graph obtained by removing P together with incident edges.

Lemma 2.5. *We have*

$$(8) \quad Z(S) = \sum_{x,y \in V(S)} d_x d_y \sum_{P \in \mathcal{P}_S(x,y)} \prod_{e \in E(P)} w_e R(S - \{P\}).$$

Lemma 2.6. *Regarding G as a subgraph of itself, we have $R(G) = 0$ and $Z(G) = \text{vol}(G)^2 \tau(G)$. For any $x, y \in V(G)$, we have*

$$(9) \quad R(G - \{x\}) = \sum_{P \in \mathcal{P}_G(x,y)} \prod_{e \in E(P)} w_e R(G - \{P\}) = \tau(G).$$

Now we come to an explicit formula for the Green's function expressed in terms of the above two invariants.

Theorem 2.7. For a connected graph G and $x, y \in V(G)$, the value of the Green's function $\mathcal{G}(x, y)$ is equal to

$$\frac{\sqrt{d_x d_y}}{\text{vol}(G)^2 \tau(G)} \left(\sum_{P \in \mathcal{P}_G(x, y)} \prod_{e \in E(P)} w_e (R(G - \{P\}) + Z(G - \{P\})) - \sum_{\substack{u, v \in V(G) \\ u \neq v}} d_u d_v \sum_{\substack{P_1 \in \mathcal{P}_G(x, u) \\ P_2 \in \mathcal{P}_G(y, v) \\ P_1 \cap P_2 = \emptyset}} \prod_{e \in E(P_1 \cup P_2)} w_e R(G - \{P_1, P_2\}) \right) - \frac{\sqrt{d_x d_y}}{\text{vol}(G)^2}.$$

In particular, when $x = y$,

$$\mathcal{G}(y, y) = \frac{d_y}{\text{vol}(G)^2 \tau(G)} (R(G - \{y\}) + Z(G - \{y\})) - \frac{d_y}{\text{vol}(G)^2}.$$

Proof. The proof is almost the same as that of [Xu and Yau 2013a, Theorem 2.9]. \square

Theorem 2.8. Given a connected graph G and $x, y \in V(G)$, the expected hitting time $H(x, y)$ satisfies

$$(10) \quad H(x, y) = \frac{1}{\text{vol}(G) \tau(G)} \left(Z(G - \{y\}) - \sum_{P \in \mathcal{P}_G(x, y)} \prod_{e \in E(P)} w_e Z(G - \{P\}) + \sum_{\substack{u, v \in V(G) \\ u \neq v}} d_u d_v \sum_{\substack{P_1 \in \mathcal{P}_G(x, u) \\ P_2 \in \mathcal{P}_G(y, v) \\ P_1 \cap P_2 = \emptyset}} \prod_{e \in E(P_1 \cup P_2)} w_e R(G - \{P_1, P_2\}) \right).$$

Proof. This follows from Theorems 1.1 and 2.7 and Lemma 2.6. \square

Corollary 2.9. On a connected weighted graph G , $H(x, y) = H(y, x)$ for any $x, y \in V(G)$ if and only if $Z(G - \{x\})$ is independent of $x \in V(G)$.

Proof. By (10), we have

$$H(x, y) - H(y, x) = \frac{1}{\text{vol}(G) \tau(G)} (Z(G - \{y\}) - Z(G - \{x\})),$$

which implies the corollary. \square

Corollary 2.10. On a connected weighted graph G , $H(x, y) = H(y, x)$ for any $x, y \in V(G)$ if and only if $\det B_{G-\{x\}}|_{s=1}$ is independent of $x \in V(G)$.

Proof. From $\det B_{G-\{x\}}|_{s=1} = R(G - \{x\}) + Z(G - \{x\})$, the conclusion follows from Lemma 2.6 and the previous corollary. \square

Theorem 2.11. *Let G be a connected weighted graph and $x, y \in V(G)$. Then*

$$(11) \quad H(x, y) = \frac{1}{\tau(G)} \sum_{u \in V(G)} d_u \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e R(G - \{P, y\}).$$

In fact, $R(G - \{P, y\}) = \tau(G/\{P, y\})$, where $G/\{P, y\}$ is obtained from G by contracting $\{P, y\}$ to a point.

Proof. The proof is almost identical to that of [Xu and Yau 2013b, Theorem 2.7]. $R(G - \{P, y\}) = \tau(G/\{P, y\})$ follows from Theorem 2.1(ii). □

Corollary 2.12. *For any connected weighted graph G , we have*

$$(12) \quad Z(G - \{x\}) = \tau(G) \sum_{y \in V(G)} d_y H(y, x).$$

Proof. By (8) and (11), we have

$$\begin{aligned} Z(G - \{x\}, d_G) &= \sum_{u, y \in V(G - \{x\})} d_u d_y \sum_{P \in \mathcal{P}_{G - \{x\}}(u, y)} \prod_{e \in E(P)} w_e R(G - \{P, x\}) \\ &= \sum_{y, u \in V(G)} d_y d_u \sum_{\substack{P \in \mathcal{P}_G(y, u) \\ x \notin P}} \prod_{e \in E(P)} w_e R(G - \{P, x\}) \\ &= \tau(G) \sum_{y \in V(G)} d_y H(y, x). \end{aligned} \quad \square$$

3. Identities and estimates of hitting times

It is natural to regard a weighted graph as an electrical network, where an edge xy has conductance w_{xy} and hence resistance $1/w_{xy}$. Chandra et al. [1989] proved that the commute time $\kappa(x, y) := H(x, y) + H(y, x)$ can be expressed in terms of the effective resistance R_{xy} between x, y ,

$$(13) \quad \kappa(x, y) = \text{vol}(G) R_{xy}.$$

The effective resistance R_{xy} can be expressed in terms of spanning trees (cf. Theorem 3.2):

$$(14) \quad R_{xy} = \frac{\tau(G/\{x, y\})}{\tau(G)}.$$

Tetali’s formula [1991] expresses $H(x, y)$ in terms of effective resistances:

$$(15) \quad H(x, y) = \frac{1}{2} \sum_{z \in V(G)} d_z (R_{xy} + R_{yz} - R_{xz}).$$

We have the following well-known upper bound of R_{xy} .

Theorem 3.1. *Given a connected graph G and $x, y \in V(G)$, we have $R_{xy} \leq d(x, y)$, where the distance $d(x, y)$ between x, y is defined by*

$$d(x, y) = \min \left\{ \sum_{e \in E(P)} \frac{1}{w_e} \mid P \in \mathcal{P}_G(x, y) \right\}.$$

As remarked in [Lovász 1996, Corollary 4.2], the following formula could be proved by the method of electric networks. Here we give a proof by using (11).

Theorem 3.2. *Let $x, y \in V(G)$ be two distinct vertices of a connected weighted graph G . Then we have*

$$(16) \quad H(x, y) + H(y, x) = \text{vol}(G) \frac{\tau(G/\{x, y\})}{\tau(G)},$$

where $G/\{x, y\}$ is obtained from G by contracting $\{x, y\}$ to a point.

Proof. Define a graph G' by

$$G' = \begin{cases} G & \text{if } x \sim y, \\ G \cup \{xy\} & \text{otherwise.} \end{cases}$$

Namely, we modify G by adding an edge xy if x, y are not adjacent.

If $u \neq x, y$, define

$$\Omega_1 = \{T \in \Omega(G') \mid T \text{ contains } xy \text{ and a path from } u \text{ to } x \text{ not containing } y\},$$

$$\Omega_2 = \{T \in \Omega(G') \mid T \text{ contains } xy \text{ and a path from } u \text{ to } y \text{ not containing } x\},$$

$$\Omega_3 = \{T \in \Omega(G') \mid T \text{ contains } xy\}.$$

It is not difficult to see that $\Omega_1 \cup \Omega_2 = \Omega_3 = \Omega(G/\{x, y\})$. (More precisely, Ω_3 is in one-to-one correspondence with $\Omega(G/\{x, y\})$.) Then, by (11),

$$\begin{aligned} & H(x, y) + H(y, x) \\ &= \frac{1}{\tau(G)} \sum_{u \in V(G)} d_u \left(\sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) \right. \\ & \quad \left. + \sum_{\substack{P \in \mathcal{P}_G(y, u) \\ x \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, x\}) \right) \\ &= \frac{1}{\tau(G)} \sum_{u \in V(G)} d_u \left(\sum_{T \in \Omega_1} \prod_{e \in T} w_e + \sum_{T \in \Omega_2} \prod_{e \in T} w_e \right) \\ &= \frac{1}{\tau(G)} \sum_{u \in V(G)} d_u \tau(G/\{x, y\}). \end{aligned}$$

The term in parenthesis on the third line is equal to $\tau(G/\{x, y\})$, independently of $u \in V(G)$. □

In the rest of this section, we apply Theorem 2.11 to prove some estimates for the hitting time on weighted graphs. It is interesting to see the role of edge weights in these estimates.

Corollary 3.3. *Let G be a connected weighted graph with n vertices and $x, y \in V(G)$. Then*

$$(17) \quad H(x, y) \leq (n - 1)^2 \frac{d_{\max}}{w_{\min}},$$

where $d_{\max} = \max\{d_v \mid v \in V(G)\}$ and $w_{\min} = \min\{w_e \mid e \in E(G)\}$.

Proof. Fix $x, y, u \in V(G)$ with $y \neq u$. Given a spanning tree $T \in \Omega(G)$ and an edge $e \in E(T)$, denote by $T(e)$ the subgraph of G' obtained from T by removing e and adding an edge uy if $uy \notin E(T)$, namely,

$$T(e) = \begin{cases} T & \text{if } uy \in T, \\ T \cup \{uy\} - \{e\} & \text{if } uy \notin T. \end{cases}$$

Define a subset S of $\Omega(G) \times E(G)$ by

$$S = \{(T, e) \mid T \in \Omega(G), e \in E(T), T(e) \in \Omega(G')\}$$

and let $S' = \{T \in \Omega(G') \mid T \text{ contains } uy\}$. Then the map $(T, e) \rightarrow T(e)$ is a surjective map from S to S' . Since

$$\begin{aligned} & \bigcup_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \Omega(G/\{P, y\}) \\ &= \{T \in \Omega(G') \mid T \text{ contains } uy \text{ and a path from } u \text{ to } x \text{ not passing through } y\} \end{aligned}$$

is a subset of S' and the left-hand side is a disjoint union over $P \in \mathcal{P}_G(x, u), y \notin P$, we have

$$\sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) \leq (n - 1) \tau(G) \frac{1}{w_{\min}}.$$

Then (17) follows from (11). □

Corollary 3.4. *Let G be a connected weighted graph and $xy \in E(G)$. Then*

$$H(x, y) \leq \frac{\text{vol}(G) - d_y}{w_{xy}}.$$

Proof. Fix $x, y, u \in V(G)$ with $y \neq u$. It is not difficult to see that

$$\begin{aligned} & \bigcup_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \Omega(G/\{P, y\}) \\ &= \{T \in \Omega(G) \mid T \text{ contains } xy \text{ and a path from } u \text{ to } x \text{ not passing through } y\} \end{aligned}$$

is a subset of $\Omega(G)$ and the left-hand side is a disjoint union over $P \in \mathcal{P}_G(x, u)$, $y \notin P$. Thus (11) implies that

$$H(x, y) \leq \frac{1}{\tau(G)} \sum_{\substack{u \in V(G) \\ u \neq y}} d_u \frac{\tau(G)}{w_{xy}} = \frac{\text{vol}(G) - d_y}{w_{xy}},$$

as claimed. □

Corollary 3.5. *Let G be a connected weighted graph. Then for any two distinct vertices $x, y \in V(G)$, we have*

$$(18) \quad H(x, y) \leq \max \left\{ \frac{d_u}{w_{yu}} \mid u \in \mathcal{S} \right\},$$

where $\mathcal{S} = \{u \in V(G) \mid \text{there is a path from } x \text{ to } u \text{ not passing through } y\}$.

Proof. Fix two distinct vertices $x, y \in V(G)$. We may assume that $w_{yu} > 0$ for all $u \in \mathcal{S}$, i.e., there is an edge connecting y and u . Otherwise the right-hand side of (18) is infinite. Define $\Omega_{xy} = \{T \in \Omega(G) \mid xy \in T\}$ and

$$V_T = \{u \in V(G) \mid T \text{ contains a path from } x \text{ to } u \text{ not passing through } y\}.$$

Let $S = \{(T, u) \mid T \in \Omega_{xy}, u \in V_T\}$. Define a map $f : S \rightarrow \Omega(G)$ by

$$f(T, u) = \begin{cases} T & \text{if } u = x, \\ \{T - xy\} \cup \{uy\} & \text{if } u \neq x, \end{cases}$$

where we used the fact that $d_y = n - 1$. It is not difficult to see that f is injective. Thus, we have

$$\begin{aligned} \sum_{u \in V(G)} d_u \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) &= \sum_{u \in V(G)} \frac{d_u}{w_{yu}} \sum_{(T, u) \in S} w(f(T, u)) \\ &\leq \max \left\{ \frac{d_u}{w_{yu}} \mid u \in \mathcal{S} \right\} \tau(G). \end{aligned}$$

Therefore (11) implies (19). □

There is a direct probabilistic proof of Corollary 3.5 (see [Xu and Yau 2013b, Remark 2.14]). In fact, all the above three corollaries may be obtained from the following more refined estimates (see also [Chang and Xu \geq 2015]):

Theorem 3.6. *Let G be a connected weighted graph. Then*

$$(19) \quad H(x, y) \leq \max\{d_u \mid u \in \Gamma(y), u \neq y\} + \sum_{\substack{u \in V(G), u \neq y \\ u \notin \Gamma(y)}} d_u \min\{d(x, y), d(u, y)\},$$

where $\Gamma(y)$ is the set of vertices adjacent to y and $d(x, y)$ is defined in Theorem 3.1.

Proof. We split the right-hand side of (11) into two terms:

$$(20) \quad H(x, y) = \frac{1}{\tau(G)} \sum_{u \in \Gamma(y)} d_u \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) \\ + \frac{1}{\tau(G)} \sum_{\substack{u \in V(G) \\ u \notin \Gamma(y)}} d_u \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}).$$

If $u \in \Gamma(y)$, then define

$$\Omega_{uy} = \{T \in \Omega(G) \mid T \text{ contains } uy \text{ and a path from } x \text{ to } u \text{ not passing through } y\}.$$

Since $\bigcup_{u \in \Gamma(y)} \Omega_{uy}$ is a disjoint union in $\Omega(G)$ and

$$\sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) = \sum_{T \in \Omega_{uy}} w(T) \quad \text{for all } u \in \Gamma(G),$$

the first summand in the right-hand side of (20) satisfies

$$(21) \quad \frac{1}{\tau(G)} \sum_{u \in \Gamma(y)} d_u \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) \leq \max\{d_u \mid u \in \Gamma(y), u \neq y\}.$$

By using (14) and the inequality

$$(22) \quad \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) \leq \min\{\tau(G/\{x, y\}), \tau(G/\{u, y\})\},$$

the second summand in the right-hand of (20) satisfies

$$(23) \quad \frac{1}{\tau(G)} \sum_{\substack{u \in V(G) \\ u \notin \Gamma(y)}} d_u \sum_{\substack{P \in \mathcal{P}_G(x, u) \\ y \notin P}} \prod_{e \in E(P)} w_e \tau(G/\{P, y\}) \\ \leq \sum_{\substack{u \in V(G) \\ u \notin \Gamma(y)}} d_u \min\{R_{xy}, R_{uy}\} \leq \sum_{\substack{u \in V(G) \\ u \notin \Gamma(y)}} d_u \min\{d(x, y), d(u, y)\}.$$

The last inequality follows from Theorem 3.1. So (19) follows from (21) and (23). \square

Corollary 3.7. *Let G be a connected weighted graph. Then*

$$(24) \quad H(x, y) \leq \sum_{\substack{u \in V(G) \\ u \neq y}} d_u \min\{R_{xy}, R_{uy}\} \leq \sum_{\substack{u \in V(G) \\ u \neq y}} d_u \min\{d(x, y), d(u, y)\}.$$

Proof. The first inequality follows from (14), (22) and Theorem 2.11. The second inequality follows from Theorem 3.1. \square

4. Some examples

First we consider infinite but locally finite connected graphs. Since an infinite (locally finite) graph can be considered as a limit of a sequence of finite graphs, the hitting time formula (11) is still valid as long as the limit exists. A weighted tree T is a locally finite tree (possibly with loops) whose edges are assigned positive weights.

Theorem 4.1. *Let x, y be two distinct vertices of a weighted tree T , and denote by P_{xy} the path $[x = v_0, v_1, \dots, v_{k-1}, v_k = y]$ connecting x to y . For any $v_i \in V(P_{xy})$, we denote by T_i the component of $T - E(P_{xy})$ that contains v_i , and denote by $w_{i-1,i}$ the weight of the edge $v_{i-1}v_i$. Then the hitting time $H(x, y)$ is given by*

$$(25) \quad H(x, y) = \sum_{j=0}^{k-1} \left(\sum_{u \in T_j} d_u \right) \left(\sum_{i=j+1}^k \frac{1}{w_{i-1,i}} \right).$$

Proof. First we define induced subtrees $T(N)$ of T for $N \in \mathbb{N}$. The vertices of $T(N)$ are those vertices whose distances from x are within N . We may apply (11) to get $H(x, y)$ on $T(N)$, which increases as N increases. Then (25) follows easily. We omit the details. □

Corollary 4.2. *Let x, y be two distinct vertices of a weighted tree T . Then $H(x, y) < \infty$ if and only if*

$$\sum_{u \in \mathcal{S}} d_u < \infty,$$

where $\mathcal{S} = \{u \in V(T) \mid \text{there is a path from } x \text{ to } u \text{ not passing through } y\}$.

Corollary 4.3. *On the weighted one-dimensional lattice \mathbb{Z} ,*

$$(26) \quad H(j, j + 1) = \frac{\sum_{i \leq j} d_i}{w_{j,j+1}}.$$

Both corollaries follow easily from (25). For unweighted trees, formula (25) was obtained in [Haiyan and Fuji 2004] (see also [Moon 1973]). Formula (26) can be found in [Palacios and Tetali 1996], where it was used to study hitting times for birth and death chains.

Now let G be a locally finite connected weighted graph and $xy \in E(G)$. Then the inequality of Corollary 3.4 still holds: $H(x, y) \leq (\text{vol}(G) - d_y)/w_{xy}$. Next we show that the equality essentially holds when xy is a cut edge of G .

Let $\mathcal{S} = \{u \in V(G) \mid \text{there is a path from } x \text{ to } u \text{ not passing through } y\}$. Let G' be the subgraph obtained by removing all vertices in $V(G)/\{\mathcal{S} \cup y\}$ from G . If $xy \in E(G)$ is a cut edge of G , note that $H(x, y)$ is the same for random walks on either G or G' . Moreover, for each spanning tree T of G' and $u \in \mathcal{S}$, there exists a

path from x to u . Therefore,

$$H(x, y) = \frac{1}{\tau(G')} \sum_{u \in \mathcal{F}} d_u \frac{1}{w_{xy}} \tau(G') = \frac{1}{w_{xy}} \sum_{u \in \mathcal{F}} d_u = \frac{\text{vol}(G') - d'_y}{w_{xy}},$$

where d'_y is the degree of y in G' .

Following [Georgakopoulos 2012], we call a weighted graph G *reversible* if $H(x, y) = H(y, x)$ holds for any $x, y \in V(G)$. For simplicity, we assume that G has no loops, i.e., $w_{xx} = 0$ for all $x \in V(G)$, and all edge weights of G are positive. It is interesting to study restrictions on edge weights for a reversible graph G .

Conjecture 4.4. *Let G be a weighted cycle on n vertices. Assume all edge weights of G are positive. Denote $w_{n,n+1} = w_{n,1}$.*

- (i) *If n is odd, then G is reversible if and only if there exists some $a > 0$ such that $w_{i,i+1} = a$ for all $1 \leq i \leq n$.*
- (ii) *If n is even, then G is reversible if and only if there exist $a, b > 0$ such that $w_{1,2} = w_{3,4} = \dots = w_{n-1,n} = a$ and $w_{2,3} = w_{4,5} = \dots = w_{n,n} = b$.*

The sufficiency in (ii) follows from Corollary 2.10.

5. Weight schemes on graphs

Given a simple, connected, undirected graph G with n vertices, we obtain a weighted graph by assigning a positive number w_e to each edge $e \in E(G)$. The hitting and cover times of a simple random walk on G (i.e., $w_e = 1$, for all $e \in E(G)$) have order $O(n^3)$. The work of [Ikeda et al. 2009; Abdullah 2011] showed that if a token knows not only the degree of the current vertex that it is on, but also the degrees of neighboring vertices, we can guarantee $O(n^2)$ hitting times.

In this section, we will denote by $d(u)$ the number of edges adjacent to a vertex u in G and assume that G has no loops.

Lemma 5.1. *Let G be connected graph with n vertices and $u_0 = x, u_1, \dots, u_l = y$ a shortest path (achieving minimum l) connecting any two distinct vertices x and y . Then $\sum_{i=0}^l d(u_i) \leq 3n - 4$. More precisely,*

$$\sum_{i=0}^l d(u_i) \leq \begin{cases} 2n - 2 & \text{if } l = 1, \\ 3n - l - 3 & \text{if } l \geq 2. \end{cases}$$

Proof. The proof is due to [Ikeda et al. 2009, Theorem 2]. Each vertex of $V(G)$ not lying on the path can be connected to at most 3 vertices of the path, due to its minimality, which also implies that u_i, u_j are adjacent if and only if $|i - j| = 1$. The asserted inequalities follow easily. \square

Next we will apply Theorem 3.6 to estimate hitting times under three different weight schemes: $w_{uv} = 1/\sqrt{d(u)d(v)}$, $1/\min\{d(u), d(v)\}$ or $1/\max\{d(u), d(v)\}$. The leading terms of the bounds in Theorems 5.2 and 5.3 were obtained in [Ikeda et al. 2009, Theorem 2] and [Abdullah 2011, Theorem 68] respectively.

Theorem 5.2. *Let G be a graph with assigned weights $w_{uv} = 1/\sqrt{d(u)d(v)}$ for each edge uv . Then the hitting time satisfies $H(x, y) \leq 3n^2 - 9n + \frac{15}{2}$.*

Proof. For the two terms in the right-hand side of (19), we have the estimates

$$(27) \quad d_u = \sum_{v \in \Gamma(u)} \frac{1}{\sqrt{d(u)d(v)}} \leq \frac{1}{2} \sum_{v \in \Gamma(u)} \left(\frac{1}{d(u)} + \frac{1}{d(v)} \right) \leq \frac{1}{2} + \frac{d(u)}{2} \leq \frac{n}{2}.$$

and, by using $\sum_{u \in V(G)} \sum_{v \in \Gamma(u)} \frac{1}{2}(1/d(u) + 1/d(v)) = n$,

$$(28) \quad \sum_{\substack{u \in V(G), u \neq y \\ u \notin \Gamma(y)}} d_u \leq \sum_{\substack{u \in V(G), u \neq y \\ u \notin \Gamma(y)}} \sum_{v \in \Gamma(u)} \frac{1}{2} \left(\frac{1}{d(u)} + \frac{1}{d(v)} \right) \leq n - 1 - \frac{d(y)}{2} \leq n - \frac{3}{2}.$$

Let $u_0 = x, u_1, \dots, u_l = y$ be a shortest path (achieving minimum l) connecting x and y . Then

$$(29) \quad d(x, y) \leq \sum_{i=0}^{l-1} \sqrt{d(u_i)d(u_{i+1})} \leq \sum_{i=0}^{l-1} \frac{d(u_i) + d(u_{i+1})}{2} \leq 3n - 5.$$

The last inequality follows from Lemma 5.1. By (19), we have

$$H(x, y) \leq (3n - 5)(n - \frac{3}{2}) + \frac{1}{2}n = 3n^2 - 9n + \frac{15}{2},$$

as claimed. □

Theorem 5.3. *Let G be a graph with assigned weights $w_{uv} = 1/\min\{d(u), d(v)\}$ for each edge uv . Then the hitting time satisfies $H(x, y) \leq 6n^2 - 18n + 14$.*

Proof. For the two terms in the right-hand side of (19), we have

$$(30) \quad d_u = \sum_{v \in \Gamma(u)} \frac{1}{\min\{d(u), d(v)\}} \leq d(u) \leq n - 1$$

and, similarly to (28),

$$(31) \quad \sum_{\substack{u \in V(G), u \neq y \\ u \notin \Gamma(y)}} d_u \leq \sum_{\substack{u \in V(G), u \neq y \\ u \notin \Gamma(y)}} \sum_{v \in \Gamma(u)} \left(\frac{1}{d(u)} + \frac{1}{d(v)} \right) \leq 2n - 3.$$

Similarly to (29), we have

$$d(x, y) \leq \sum_{i=0}^{l-1} \min\{d(u_i), d(u_{i+1})\} \leq \sum_{i=0}^{l-1} d(u_i) \leq 3n - 5.$$

The desired upper bound of $H(x, y)$ follows from (19). \square

Theorem 5.4. *Let G be a graph with assigned weights $w_{uv} = 1/\max\{d(u), d(v)\}$ for each edge uv . Then the hitting time satisfies $H(x, y) \leq 6n^2 - 23n + 23$.*

Proof. For the two terms in the right-hand side of (19), we have

$$d_u = \sum_{v \in \Gamma(u)} \frac{1}{\max\{d(u), d(v)\}} \leq 1 \quad \text{and} \quad \sum_{\substack{u \in V(G), u \neq y \\ u \notin \Gamma(y)}} d_u \leq n - 2.$$

Similar to (29), we have

$$d(x, y) \leq \sum_{i=0}^{l-1} \max\{d(u_i), d(u_{i+1})\} \leq \sum_{i=0}^{l-1} (d(u_i) + d(u_{i+1}) - 1) \leq 6n - 11.$$

The desired upper bound of $H(x, y)$ follows from (19). \square

References

- [Abdullah 2011] M. Abdullah, *The cover time of random walks on graphs*, Ph.D. thesis, King's College London, 2011. arXiv 1202.5569
- [Aldous and Fill 2014] D. Aldous and J. Fill, "Reversible Markov chains and random walks on graphs", Unfinished monograph, 2014, Available at <http://www.stat.berkeley.edu/~aldous/RWG/book.pdf>.
- [Chandra et al. 1989] A. K. Chandra, P. Raghavan, W. L. Ruzzo, R. Smolensky, and P. Tiwari, "The electrical resistance of a graph captures its commute and cover times", pp. 574–586 in *Proc. 21st ACM Symp. Theory of Computing* (Seattle, 1989), edited by D. S. Johnson, Association for Computing Machinery, New York, 1989.
- [Chang and Xu ≥ 2015] X. Chang and H. Xu, "Chung–Yau invariants and graphs with symmetric hitting times", preprint, Available at <http://pitt.edu/~haoxu/ReversibleGraph.pdf>.
- [Chung 2011] F. R. K. Chung, "PageRank as a discrete Green's function", pp. 285–302 in *Geometry and analysis, I* (Cambridge, MA, 2008), edited by L. Ji, Adv. Lect. Math. (ALM) **17**, Int. Press, Somerville, MA, 2011. MR 2012m:68344 Zbl 1255.68109
- [Chung and Yau 2000] F. Chung and S.-T. Yau, "Discrete Green's functions", *J. Combin. Theory Ser. A* **91**:1-2 (2000), 191–214. MR 2001g:05073 Zbl 0963.65120
- [Georgakopoulos 2012] A. Georgakopoulos, "On walk-regular graphs and graphs with symmetric hitting times", preprint, 2012. arXiv 1211.5689
- [Haiyan and Fuji 2004] C. Haiyan and Z. Fuji, "The expected hitting times for graphs with cutpoints", *Statist. Probab. Lett.* **66**:1 (2004), 9–17. MR 2005a:60065 Zbl 1113.60046
- [Ikeda et al. 2009] S. Ikeda, I. Kubo, and M. Yamashita, "The hitting and cover times of random walks on finite graphs using local degree information", *Theoret. Comput. Sci.* **410**:1 (2009), 94–100. MR 2010c:60143 Zbl 05509592

- [Lovász 1996] L. Lovász, “Random walks on graphs: A survey”, pp. 353–397 in *Combinatorics, Paul Erdős is eighty, II* (Keszthely, 1993), edited by D. Miklós et al., Bolyai Soc. Math. Stud. **2**, János Bolyai Math. Soc., Budapest, 1996. MR 97a:60097 Zbl 0854.60071
- [Moon 1973] J. W. Moon, “Random walks on random trees”, *J. Austral. Math. Soc.* **15** (1973), 42–53. MR 47 #6525 Zbl 0265.60065
- [Palacios and Tetali 1996] J. L. Palacios and P. Tetali, “A note on expected hitting times for birth and death chains”, *Statist. Probab. Lett.* **30**:2 (1996), 119–125. MR 97h:60069 Zbl 0883.60081
- [Tetali 1991] P. Tetali, “Random walks and the effective resistance of networks”, *J. Theoret. Probab.* **4**:1 (1991), 101–109. MR 92c:60097 Zbl 0722.60070
- [Xu and Yau 2013a] H. Xu and S.-T. Yau, “Discrete Green’s functions and random walks on graphs”, *J. Combin. Theory Ser. A* **120**:2 (2013), 483–499. MR 2995053 Zbl 1256.05225
- [Xu and Yau 2013b] H. Xu and S. T. Yau, “An explicit formula of hitting times for random walks on graphs”, preprint, 2013. To appear in *Pure Appl. Math. Q.* arXiv 1312.0065

Received February 24, 2014.

XIAO CHANG
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF PITTSBURGH
301 THACKERAY HALL
PITTSBURGH, PA 15260
UNITED STATES
xic58@pitt.edu

HAO XU
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF PITTSBURGH
301 THACKERAY HALL
PITTSBURGH, PA 15260
UNITED STATES
mathxuhao@gmail.com

SHING-TUNG YAU
DEPARTMENT OF MATHEMATICS
HARVARD UNIVERSITY
CAMBRIDGE, MA 02138
UNITED STATES
yau@math.harvard.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 273 No. 1 January 2015

Maximal estimates for Schrödinger equations with inverse-square potential	1
CHANGXING MIAO, JUNYONG ZHANG and JIQIANG ZHENG	
Vassiliev Invariants of Virtual Legendrian Knots	21
PATRICIA CAHN and ASA LEVI	
Some results on the generic vanishing of Koszul cohomology via deformation theory	47
JIE WANG	
Conformal metrics with constant curvature one and finitely many conical singularities on compact Riemann surfaces	75
QING CHEN, WEI WANG, YINGYI WU and BIN XU	
\mathbb{Q} -bases of the Néron–Severi groups of certain elliptic surfaces	101
MASAMICHI KURODA	
On a prime zeta function of a graph	123
TAKEHIRO HASEGAWA and SEIKEN SAITO	
On Whittaker modules for a Lie algebra arising from the 2-dimensional torus	147
SHAOBIN TAN, QING WANG and CHENGGANG XU	
Fréchet quantum supergroups	169
AXEL DE GOURSAC	
Generators of the Gauss–Picard modular group in three complex dimensions	197
BAOHUA XIE, JIEYAN WANG and YUEPING JIANG	
Complete characterization of isolated homogeneous hypersurface singularities	213
STEPHEN YAU and HUIQING ZUO	
A theorem of Mœglin and Waldspurger for covering groups	225
SHIV PRAKASH PATEL	
Spanning trees and random walks on weighted graphs	241
XIAO CHANG, HAO XU and SHING-TUNG YAU	



0030-8730(201501)273:1;1-6