

*Pacific
Journal of
Mathematics*

**UNIFORM BOUNDEDNESS OF S -UNITS
IN ARITHMETIC DYNAMICS**

HOLLY KRIEGER, AARON LEVIN, ZACHARY SCHERR,
THOMAS TUCKER, YU YASUFUKU AND MICHAEL E. ZIEVE

UNIFORM BOUNDEDNESS OF S -UNITS IN ARITHMETIC DYNAMICS

HOLLY KRIEGER, AARON LEVIN, ZACHARY SCHERR,
THOMAS TUCKER, YU YASUFUKU AND MICHAEL E. ZIEVE

Let K be a number field and let S be a finite set of places of K which contains all the archimedean places. For any $\phi(z) \in K(z)$ of degree $d \geq 2$ which is not a d -th power in $\bar{K}(z)$, Siegel's theorem implies that the image set $\phi(K)$ contains only finitely many S -units. We conjecture that the number of such S -units is bounded by a function of $|S|$ and d (independently of K , S and ϕ). We prove this conjecture for several classes of rational functions, and show that the full conjecture follows from the Bombieri–Lang conjecture.

1. Introduction

Let K be a number field, let S be a finite set of places of K which contains the set S_∞ of archimedean places of K , and write \mathfrak{o}_S for the ring of S -integers of K and \mathfrak{o}_S^* for the group of S -units of K . The genus-0 case of Siegel's theorem asserts that, for any $\phi(z) \in K(z)$ which has at least three poles in $\mathbb{P}^1(\bar{K})$, the image set $\phi(K)$ contains only finitely many S -integers. However, the number of S -integers in $\phi(K)$ cannot be bounded independently of $\phi(z)$, even if we restrict to functions $\phi(z)$ having a fixed degree, since $\psi(z) := \beta\phi(z)$ satisfies $\psi(K) = \beta\phi(K)$ for any $\beta \in K^*$.

Although the number of S -integers in $\phi(K)$ cannot be bounded in terms of only K , S , and $\deg \phi$, such a bound may be possible for the number of S -units in $\phi(K)$. In fact we conjecture that there is a bound depending only on $|S|$ and $\deg \phi$ (and not on K):

Conjecture 1.1. *For any integers $s \geq 1$ and $d \geq 2$, there is a constant $C = C(s, d)$ such that for any*

- *number field K ,*

The authors were partially supported by NSF grants DMS-1303770 (H.K.), DMS-1102563 (A.L.), DMS-1200749 (T.T.), and DMS-1162181 (M.Z.). The fifth author was partially supported by JSPS Grants-in-Aid 23740033.

MSC2010: primary 37P05, 37P15; secondary 11G99, 11R99.

Keywords: arithmetic dynamics, S -units, uniform boundedness.

- s -element set S of places of K with $S \supseteq S_\infty$,
- degree- d rational function $\phi(z) \in K(z)$ which is not a d -th power in $\bar{K}(z)$,

we have

$$|\phi(K) \cap \mathfrak{o}_S^*| \leq C.$$

We will prove [Conjecture 1.1](#) in case $\phi(z)$ is restricted to certain classes of rational functions, and we will also prove that the full conjecture is a consequence of a variant of the Caporaso–Harris–Mazur conjecture on uniform boundedness of rational points on curves of fixed genus.

We also consider a variant of [Conjecture 1.1](#), which addresses S -units in an orbit of ϕ rather than in the image set $\phi(K)$. Here, for any $\alpha \in \mathbb{P}^1(K)$, the orbit of α under $\phi(z)$ is the set

$$\mathcal{O}_\phi(\alpha) := \{\phi^n(\alpha) : n \geq 1\},$$

where $\phi^n = \phi \circ \dots \circ \phi$ denotes the n -fold composition of ϕ with itself. For any $\phi(z) \in K(z)$ of degree at least 2 such that $\phi^2(z) \notin K[z]$, Silverman [1993] showed that $\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S$ is finite. However, for any $\beta \in K^*$ the function $\psi(z) := \beta\phi(z/\beta)$ satisfies $\mathcal{O}_\psi(\alpha\beta) = \beta\mathcal{O}_\phi(\alpha)$, so the size of $\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S$ cannot be bounded independently of $\phi(z)$. We conjecture that there is a uniform bound on the number of S -units in an orbit:

Conjecture 1.2. *For any integers $s \geq 1$ and $d \geq 2$, there is a constant $C = C(s, d)$ such that for any*

- number field K ,
- s -element set S of places of K with $S \supseteq S_\infty$,
- degree- d rational function $\phi(z) \in K(z)$ which is not of the form $\beta z^{\pm d}$ with $\beta \in K^*$,
- $\alpha \in \mathbb{P}^1(K)$,

we have

$$|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq C.$$

It turns out that this conjecture is a consequence of [Conjecture 1.1](#):

Proposition 1.3. *If [Conjecture 1.1](#) is true then [Conjecture 1.2](#) is true.*

Remark 1.4. The hypotheses of [Conjectures 1.1](#) and [1.2](#) imply that $[K : \mathbb{Q}] \leq 2s$, since $S_\infty \subseteq S$.

In [Section 3](#) we prove the following preliminary results, which show that [Conjectures 1.1](#) and [1.2](#) would be true if we allowed the constants C in those conjectures to depend on K , S , and ϕ , rather than just on s and d . We note that in the case of [Conjecture 1.1](#) this simply says that $\phi(K) \cap \mathfrak{o}_S^*$ is finite. These results also indicate the special behavior of the functions excluded in the statements of these conjectures.

Proposition 1.5. *Let K be a number field, let S be a finite set of places of K with $S \supseteq S_\infty$, and let $\phi(z) \in K(z)$ be any rational function.*

- (a) *If $|\phi^{-1}(\{0, \infty\})| \neq 2$ then $\phi(K) \cap \mathfrak{o}_S^*$ is finite.*
- (b) *If $|\phi^{-1}(\{0, \infty\})| = 2$ then there is a finite set $S' \supseteq S$ for which $\phi(K) \cap \mathfrak{o}_{S'}^*$ is infinite.*

Proposition 1.6. *Let K be a number field, let S be a finite set of places of K with $S \supseteq S_\infty$, and let $\phi(z) \in K(z)$ have degree $d \geq 2$.*

- (a) *If $\phi(z)$ does not have the form $\beta z^{\pm d}$ with $\beta \in K^*$, then there is a constant $C(K, S, \phi)$ such that every $\alpha \in \mathbb{P}^1(K)$ satisfies $|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq C(K, S, \phi)$.*
- (b) *If $\phi(z) = \beta z^{\pm d}$ with $\beta \in K^*$, then there exist $\alpha \in \mathbb{P}^1(K)$ and a finite set $S' \supseteq S$ for which $\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_{S'}^*$ is infinite.*

We note that part (a) of each of these propositions follows from Siegel's theorem. For, if $|\phi^{-1}(\{0, \infty\})| > 2$ then $\psi(z) := \phi(z) + 1/\phi(z)$ has at least three poles so that $\psi(K) \cap \mathfrak{o}_S$ is finite; but $\psi(\beta)$ is in \mathfrak{o}_S whenever $\phi(\beta)$ is in \mathfrak{o}_S^* , so also $\phi(K) \cap \mathfrak{o}_S^*$ is finite. Next, if $\phi^{-1}(\{0, \infty\})$ is a two-element set other than $\{0, \infty\}$, then [Lemma 3.2](#) implies that $|\phi^{-2}(\{0, \infty\})| > 2$, so that $\phi^2(K) \cap \mathfrak{o}_S^*$ has size $N < \infty$, whence $|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq N + 1 = C(K, S, \phi)$.

In [Section 2](#) we prove [Conjectures 1.1](#) and [1.2](#) for some families of polynomial maps. The first family consists of monic polynomials in $\mathfrak{o}_S[z]$:

Theorem 1.7. *Let $s \geq 1$ and $d \geq 2$ be integers. There is a constant $C = C(s, d)$ such that for any*

- *number field K ,*
- *s -element set S of places of K with $S \supseteq S_\infty$,*
- *degree- d monic polynomial $\phi(z) \in \mathfrak{o}_S[z]$ which does not equal $(z - \beta)^d$ for any $\beta \in K$,*

we have

$$|\phi(K) \cap \mathfrak{o}_S^*| \leq C.$$

[Theorem 1.7](#) proves [Conjecture 1.1](#) for monic polynomials in $\mathfrak{o}_S[z]$; for such polynomials, [Conjecture 1.2](#) follows by applying [Theorem 1.7](#) to $\phi^2(z)$.

We also prove [Conjecture 1.2](#) for monic polynomials in $K[z]$ in which the coefficients of all but one term are in \mathfrak{o}_S , so long as this exceptional term does not have degree $d - 1$. We deduce this from the following more general result in v -adic dynamics.

Theorem 1.8. *Let K be a field with a nonarchimedean valuation v , and let*

$$\phi(z) = a_d z^d + \cdots + a_1 z + a_0 \in K[z]$$

be a polynomial satisfying

- $v(a_d) = 0$,
- there is exactly one integer i for which $v(a_i) < 0$, and this exceptional i satisfies $i \neq d - 1$.

Then for each $\alpha \in K$, the set $\{n \geq 1 \mid v(\phi^n(\alpha)) = 0\}$ contains at most one element.

As an immediate corollary, we have the stated case of [Conjecture 1.2](#):

Corollary 1.9. *Let K be a number field, and let S be a finite set of places of K with $S \supseteq S_\infty$. For any monic $\phi_0(z) \in \mathfrak{o}_S[z]$, any $\alpha, \beta \in K$ with $\beta \notin \mathfrak{o}_S$, and any integer i with $0 \leq i < \deg \phi_0 - 1$, the polynomial $\phi(z) := \phi_0(z) + \beta z^i$ satisfies*

$$|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq 1.$$

Remark 1.10. [Conjecture 1.2](#) also follows from Theorem 2 of [\[Levin 2012\]](#) for rational functions of the form

$$\phi(z) := \frac{z^d + \beta_{d-1}z^{d-1} + \cdots + \beta_1z}{\gamma_{d-1}z^{d-1} + \gamma_{d-2}z^{d-2} + \cdots + \gamma_1z + 1}$$

with $\beta_1, \dots, \beta_{d-1}, \gamma_1, \dots, \gamma_{d-1} \in \mathfrak{o}_S$ and $\phi(z) \neq z^d$. For that theorem gives a uniform bound on the number of elements of K in the backwards orbit of any element of \mathfrak{o}_S^* . This also bounds the number of S -units in $\mathcal{O}_\phi(\alpha)$ for any $\alpha \in K$, since if $\phi^n(\alpha) \in \mathfrak{o}_S^*$ then $\alpha, \phi(\alpha), \dots, \phi^{n-1}(\alpha)$ are elements of K in the backwards orbit of $\phi^n(\alpha)$.

We prove our conjectures for some further classes of rational functions in [Section 4](#).

In [Section 3](#) we show that our conjectures are consequences of the following variant of the deep conjecture of Caporaso, Harris and Mazur [\[Caporaso et al. 1997\]](#) concerning rational points on curves of a fixed genus.

Conjecture 1.11. *Fix integers $g \geq 2$ and $D \geq 1$. There is a constant $N = N(D, g)$ such that $|X(K)| \leq N$ for every smooth, projective, geometrically irreducible genus- g curve X defined over a degree- D number field K .*

Theorem 1.12. *If [Conjecture 1.11](#) is true, then [Conjecture 1.1](#) and [Conjecture 1.2](#) are true.*

Remark 1.13. [Conjecture 1.11](#) follows from the Bombieri–Lang conjecture [\[Pacelli 1997\]](#).

The referee provided the following geometric explanation of the difference between the questions of S -integers and S -units in the image set $\phi(K)$ of a rational function ϕ , indicating possible directions for future work. Writing $\phi(x/y) = \frac{f(x,y)}{g(x,y)}$ as the ratio of two coprime homogeneous polynomials, we see that the S -integral

points of $\phi(K)$ correspond to the S -integral points of the quasi-affine variety cut out by

$$zg(x, y) = f(x, y) \text{ in } \mathbb{P}^1 \times \mathbb{A}^1.$$

Similarly, the S -unit points in $\phi(K)$ correspond to the S -integral points of the variety defined by

$$zg(x, y) = wf(x, y) \text{ and } zw = 1 \text{ in } \mathbb{P}^1 \times \mathbb{A}^2.$$

It would be interesting to seek generalizations of [Conjecture 1.1](#) by considering more generally what sorts of families of varieties are likely to satisfy uniform boundedness statements for their S -integral points.

2. Special classes of rational functions

In this section we prove [Theorems 1.7](#) and [1.8](#).

Proof of [Theorem 1.7](#). Let K be a number field, let S be a finite set of places of K with $S \supseteq S_\infty$, and let $\phi(z) \in \mathfrak{o}_S[z]$ be monic of degree $d \geq 2$ with $\phi(z) \neq (z - \beta)^d$ for any $\beta \in K$. Then $\phi(z)$ has at least two distinct roots δ_1, δ_2 in \bar{K} . Let $K' = K(\delta_1, \delta_2)$ and let S' be the set of places of K' which lie over places in S , so that $|S'| \leq [K' : K]|S| \leq d(d-1)|S|$ and $\delta_i \in \mathfrak{o}_{S'}$. Then we can write

$$\phi(z) = (z - \delta_1)(z - \delta_2)\psi(z),$$

where $\psi(z)$ is a monic polynomial in $\mathfrak{o}_{S'}[z]$. Let $\gamma \in K$ satisfy $\phi(\gamma) \in \mathfrak{o}_S^*$. Then we must have $\gamma \in \mathfrak{o}_S$, so that both $u_i := \gamma - \delta_i$ and $\psi(\gamma)$ are in $\mathfrak{o}_{S'}$. Since $u_1 u_2 \psi(\gamma) = \phi(\gamma)$ is in \mathfrak{o}_S^* , it follows that $u_1, u_2 \in \mathfrak{o}_{S'}^*$. In addition we have

$$(2-1) \quad \frac{1}{\delta_2 - \delta_1} u_1 - \frac{1}{\delta_2 - \delta_1} u_2 = 1.$$

Moreover, γ is uniquely determined by u_1 , so the number of elements $\gamma \in \mathfrak{o}_S$ for which $\phi(\gamma) \in \mathfrak{o}_S^*$ is at most the number of solutions to (2-1) in elements $u_1, u_2 \in \mathfrak{o}_{S'}^*$. Finally, by [\[Evertse 1984\]](#), the number of such solutions is at most $C_1 C_2^{|S'|-1}$ for some absolute constants C_1, C_2 (in fact, we can take $C_1 = C_2 = 256$ [\[Beukers and Schlickewei 1996\]](#)). Therefore $|\phi(K) \cap \mathfrak{o}_S^*|$ is bounded by a function of $|S'|$, and hence by a function of $|S|$ and d . \square

Proof of [Theorem 1.8](#). Suppose that $\mathcal{O}_\phi(\alpha)$ contains a unit of the valuation ring, and let m be the least positive integer for which $v(\phi^m(\alpha)) = 0$. Writing $\gamma := \phi^m(\alpha)$, we will show by induction that $|\phi^n(\gamma)|_v = |a_i|_v^{d^{n-1}}$ for every $n \geq 1$. The strong triangle inequality implies that $|\phi(\gamma)|_v = |a_i|_v$, proving the base case $n = 1$. If $\delta := \phi^n(\gamma)$ satisfies $|\delta|_v = |a_i|_v^{d^{n-1}}$ for some $n \geq 1$, then $|a_i \delta^i|_v = |a_i|_v^{1+d^{n-1}}$ and $|a_j \delta^j|_v \leq |a_i|_v^{j d^{n-1}}$ for $j \neq i$, with equality when $j = d$. Our hypothesis $i < d - 1$

implies that $d^n > 1 + id^{n-1}$, so that $|\phi^{n+1}(\gamma)|_v = |a_i|_v^{d^n}$, which completes the induction. It follows that $v(\phi^n(\gamma)) < 0$ for every $n > 0$, so that $\mathcal{O}_\phi(\alpha)$ contains exactly one unit, which concludes the proof. \square

3. Connection with rational points on curves

In this section we prove [Theorem 1.12](#) and [Propositions 1.3, 1.5, and 1.6](#). We begin by relating S -units in the image set $\phi(K)$ of a rational function to rational points on certain curves.

Lemma 3.1. *Let K be a number field, let S be a finite set of places of K with $S \supseteq S_\infty$, and let $\phi(z) \in K(z)$ be a nonconstant rational function. For any prime p with $p > \deg \phi$, there are elements $\gamma_1, \dots, \gamma_t \in \mathfrak{o}_S^*$, where $t \leq p^{|S|}$, with the following properties:*

- For each i , the affine curve X_i defined by $y^p = \gamma_i \phi(z)$ is geometrically irreducible.
- We have $|\phi(K) \cap \mathfrak{o}_S^*| \leq \sum_{i=1}^t N_i$ where N_i is the number of points in $X_i(K)$ having nonzero y -coordinate.

Proof. First note that $y^p = \gamma \phi(z)$ is geometrically irreducible for any $\gamma \in K^*$, since $\gamma \phi(z)$ is not a p -th power in $\bar{K}(z)$. Dirichlet's S -unit theorem asserts that $\mathfrak{o}_S^* \cong \mu_K \times \mathbb{Z}^{|S|-1}$, where μ_K denotes the group of roots of unity in K . Since μ_K is cyclic, it follows that $\mathfrak{o}_S^* / (\mathfrak{o}_S^*)^p \cong (\mathbb{Z}/p\mathbb{Z})^r$ where $r \in \{|S| - 1, |S|\}$. Let Γ be a set of p^r elements in \mathfrak{o}_S^* whose images in $\mathfrak{o}_S^* / (\mathfrak{o}_S^*)^p$ are pairwise distinct. For any $\beta \in K$ such that $\phi(\beta) \in \mathfrak{o}_S^*$, we can write $\phi(\beta) = \gamma^{-1} \delta^p$ for some $\gamma \in \Gamma$ and $\delta \in \mathfrak{o}_S^*$. Then (δ, β) is a K -rational point on the curve $y^p = \gamma \phi(z)$ whose y -coordinate is nonzero. Since the z -coordinate of this point is β , the result follows. \square

Proof of Theorem 1.12. By [Proposition 1.3](#), it suffices to show that [Conjecture 1.11](#) implies [Conjecture 1.1](#). Let K be a number field, let S be a finite set of places of K with $S \supseteq S_\infty$, and let $\phi(z) \in K(z)$ have degree $d \geq 2$. Assume that $\phi(z)$ is not a d -th power in $\bar{K}(z)$, so that $m := |\phi^{-1}(\{0, \infty\})|$ is at least 3. Let p be the smallest prime for which $p > d$ and $(p - 1)(m - 2) > 2$. Then $p = 5$ if $d = 2$ and $m = 3$, and in all other cases $p < 2d$ by Bertrand's postulate. Let $\gamma_1, \dots, \gamma_t$ satisfy the conclusion of [Lemma 3.1](#), so that $\gamma_i \in K^*$ and $t \leq p^{|S|}$. Writing X_i for the curve $y^p = \gamma_i \phi(z)$, and N_i for the number of points in $X_i(K)$ having nonzero y -coordinate, it follows that $|\phi(K) \cap \mathfrak{o}_S^*| \leq \sum_{i=1}^t N_i$. Since every point on X_i having nonzero y -coordinate is nonsingular, we see that N_i is bounded above by the number of K -rational points on the unique smooth projective curve Y_i over K which is birational to X_i . Since $p > d$, the classical genus formula for Kummer covers [[Stichtenoth 2009](#), Proposition III.7.3] implies that the genus g of Y_i is

$(p-1)(m-2)/2$. Thus our choice of p ensures that

$$2 \leq g \leq \frac{1}{2}(\frac{5}{2}d - 1)(2d - 2).$$

If [Conjecture 1.11](#) is true then $|Y_i(K)|$ is bounded by a constant which depends only on the genus of $Y_i(K)$ and the degree $[K : \mathbb{Q}]$. Since the genus is bounded by a function of d , and the degree $[K : \mathbb{Q}]$ is bounded by a function of $|S|$ (by [Remark 1.4](#)), it follows that $|Y_i(K)|$ is bounded by a constant depending on d and $|S|$. Since $t \leq p^{|S|} \leq (5d/2)^{|S|}$, this proves that [Conjecture 1.11](#) implies [Conjecture 1.1](#). \square

Our proof of [Proposition 1.3](#) relies on the following well-known lemma.

Lemma 3.2. *Let $\phi(z) \in \mathbb{C}(z)$ be any rational function of degree $d \geq 2$ which is not of the form $\beta z^{\pm d}$ with $\beta \in \mathbb{C}^*$. Then $|\phi^{-2}(\{0, \infty\})| \geq 3$.*

Proof. Write $m := |\phi^{-2}(\{0, \infty\})|$, so we must show that $m \geq 3$. Plainly $m \geq |\phi^{-1}(\{0, \infty\})| \geq 2$, so the conclusion holds unless $|\phi^{-1}(\{0, \infty\})| = 2$. In this case ϕ is totally ramified over both 0 and ∞ , so the Riemann–Hurwitz formula (or writing down $\phi(z)$) implies that ϕ is unramified over all other points. Since $\phi(z)$ does not have the form $\beta z^{\pm d}$, we know that $\phi^{-1}(\{0, \infty\}) \neq \{0, \infty\}$, so that at least one point in $\phi^{-1}(\{0, \infty\})$ has d distinct ϕ -preimages. Since each point has at least one preimage, we conclude that $m \geq d + 1 \geq 3$, as desired. \square

Proof of [Proposition 1.3](#). If $\phi(z) \neq \beta z^{\pm d}$ then $\phi^2(z)$ has a total of at least three zeroes and poles by [Lemma 3.2](#), and hence is not a d^2 -th power in $\bar{K}(z)$. Thus [Conjecture 1.1](#) implies that $|\phi^2(K) \cap \mathfrak{o}_S^*| \leq C(s, d)$, so that

$$|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq C(s, d) + 1. \quad \square$$

Part (a) of [Proposition 1.5](#) follows from our proof of [Theorem 1.12](#), by using Faltings’ theorem [[1983](#)] instead of [Conjecture 1.11](#). We now give a more elementary proof of [Proposition 1.5](#).

Proof of [Proposition 1.5](#). If $|\phi^{-1}(\{0, \infty\})| > 2$, the function $\psi(z) := \phi(z) + 1/\phi(z)$ satisfies $|\psi^{-1}(\{0, \infty\})| \geq 3$, so $\psi(K) \cap \mathfrak{o}_S$ is finite by Siegel’s theorem; but $\psi(\beta)$ is in \mathfrak{o}_S whenever $\phi(\beta)$ is in \mathfrak{o}_S^* , so it follows that $\phi(K) \cap \mathfrak{o}_S^*$ is finite. Now assume that $|\phi^{-1}(\{0, \infty\})| = 2$, so that $\phi(z) = \gamma \mu(z)^d$ for some $d \geq 1$, some $\gamma \in K^*$, and some degree-one $\mu(z) \in K(z)$. Let S' be a finite set of places of K such that $\gamma \in \mathfrak{o}_{S'}^*$, $S' \supseteq S$, and $|S'| > 1$. Since $\mu(K)$ contains all but at most one element of K , it follows that $\phi(K)$ contains all but at most one element of $\gamma(\mathfrak{o}_{S'}^*)^d$. Since $\gamma \in \mathfrak{o}_{S'}^*$ and $|S'| > 1$, this shows that $\phi(K) \cap \mathfrak{o}_{S'}^*$ is infinite. \square

Proof of [Proposition 1.6](#). If $\phi(z)$ does not have the form $\beta z^{\pm d}$ then $|\phi^{-2}(\{0, \infty\})| \geq 3$ by [Lemma 3.2](#), so [Proposition 1.5](#) implies that $\phi^2(K) \cap \mathfrak{o}_S^*$ has size $N < \infty$, whence

$$|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq N + 1 = C(K, S, \phi).$$

Now consider $\phi(z) = \beta z^{\pm d}$ with $\beta \in K^*$ and $d \geq 2$. Any $\alpha \in K^*$ satisfies $\mathcal{O}_\phi(\alpha) \subseteq \mathfrak{o}_{S'}^*$, where S' is the union of S with the set of places v of K for which $|\alpha|_v \neq 1$ or $|\beta|_v \neq 1$. If $\alpha \in K^*$ is not a root of unity then $\mathcal{O}_\phi(\alpha)$ is infinite, so that $\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_{S'}^*$ is infinite. \square

4. Additional remarks

We make two final remarks. First, the proofs of Theorems 1.7 and 1.8 can be modified to treat some classes of Laurent polynomials. For example, let d and d' be distinct positive integers, and let $\phi(z) = (\gamma_d z^d + \cdots + \gamma_1 z + \gamma_0)/z^{d'}$ where $\gamma_i \in \mathfrak{o}_S$ and $\gamma_d, \gamma_0 \in \mathfrak{o}_S^*$. Suppose in addition that the numerator is not a d -th power in $\overline{K}[z]$. Then $|\phi(K) \cap \mathfrak{o}_S^*| \leq C(s, d)$ for any $\alpha \in \mathbb{P}^1(K)$. Indeed, since γ_0 and γ_d are assumed to be units, $\phi(\beta)$ cannot be in \mathfrak{o}_S^* if $|\beta|_v \neq 1$ for some $v \notin S$. Thus we need only consider $\beta \in \mathfrak{o}_S^*$, and now the desired bound follows from the proof of Theorem 1.7.

As another example, consider $\phi(z) = (\gamma_d z^d + \cdots + \gamma_1 z + \gamma_0)/z^{d'}$ where $d > d'$, $\gamma_i \in K$, and there is some $v \notin S$ for which $|\gamma_d|_v > \max(1, |\gamma_i|_v)$ for each $i < d$. Then $|\mathcal{O}_\phi(\alpha) \cap \mathfrak{o}_S^*| \leq 1$ for any $\alpha \in \mathbb{P}^1(K)$, as the orbit of an S -unit cannot contain another S -integer by the proof of Theorem 1.8. Both this class of examples and the previous class are quite special, but they serve as further evidence for Conjectures 1.1 and 1.2.

We conclude this paper by noting that the constant C that appears in Conjectures 1.1 and 1.2 must depend on both s and d . The necessity of dependence on s is clear. Dependence on d is also required, since by Lagrange interpolation one can construct polynomials $\phi(z) \in K[z]$ in which the first several $\phi^i(\alpha)$ take on any prescribed distinct values in K while also $\phi(z)$ has at least two zeroes (and hence is not $\beta z^{\pm d}$).

Acknowledgments

We thank ICERM, where collaboration for this project began at the 2012 ICERM workshop on Global Arithmetic Dynamics. We also thank the referee for a thorough report which improved the exposition and content of this paper.

References

- [Beukers and Schlickewei 1996] F. Beukers and H. P. Schlickewei, “The equation $x + y = 1$ in finitely generated groups”, *Acta Arith.* **78**:2 (1996), 189–199. MR 97k:11051 Zbl 0880.11034
- [Caporaso et al. 1997] L. Caporaso, J. Harris, and B. Mazur, “Uniformity of rational points”, *J. Amer. Math. Soc.* **10**:1 (1997), 1–35. MR 97d:14033 Zbl 0872.14017
- [Evertse 1984] J.-H. Evertse, “On equations in S -units and the Thue–Mahler equation”, *Invent. Math.* **75**:3 (1984), 561–584. MR 85f:11048 Zbl 0521.10015
- [Faltings 1983] G. Faltings, “Endlichkeitssätze für abelsche Varietäten über Zahlkörpern”, *Invent. Math.* **73**:3 (1983), 349–366. MR 85g:11026a Zbl 0588.14026

- [Levin 2012] A. Levin, “Rational preimages in families of dynamical systems”, *Monatsh. Math.* **168**:3–4 (2012), 473–501. MR 2993960 Zbl 06111280
- [Pacelli 1997] P. L. Pacelli, “Uniform boundedness for rational points”, *Duke Math. J.* **88**:1 (1997), 77–102. MR 98b:14020 Zbl 0935.14016
- [Silverman 1993] J. H. Silverman, “Integer points, Diophantine approximation, and iteration of rational maps”, *Duke Math. J.* **71**:3 (1993), 793–829. MR 95e:11070 Zbl 0811.11052
- [Stichtenoth 2009] H. Stichtenoth, *Algebraic function fields and codes*, 2nd ed., Graduate Texts in Mathematics **254**, Springer, Berlin, 2009. MR 2010d:14034 Zbl 1155.14022

Received June 19, 2014. Revised August 27, 2014.

HOLLY KRIEGER
DEPARTMENT OF MATHEMATICS
MASSACHUSETTS INSTITUTE OF TECHNOLOGY
CAMBRIDGE, MA 02139
UNITED STATES
hkrieger@math.mit.edu

AARON LEVIN
DEPARTMENT OF MATHEMATICS
MICHIGAN STATE UNIVERSITY
EAST LANSING, MI 48824
UNITED STATES
adlevin@math.msu.edu

ZACHARY SCHERR
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF PENNSYLVANIA
DAVID RITTENHOUSE LAB
PHILADELPHIA, PA 19104–6395
UNITED STATES
zscherr@math.upenn.edu

THOMAS TUCKER
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ROCHESTER
ROCHESTER, NY 14627
UNITED STATES
thomas.tucker@rochester.edu

YU YASUFUKU
DEPARTMENT OF MATHEMATICS
COLLEGE OF SCIENCE AND TECHNOLOGY
NIHON UNIVERSITY
CHIYODA-KU, TOKYO 101-8308
JAPAN
yasufuku@math.cst.nihon-u.ac.jp

MICHAEL E. ZIEVE
MATHEMATICAL SCIENCES CENTER
TSINGHUA UNIVERSITY
BEIJING, 100084
CHINA

and

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MICHIGAN
ANN ARBOR, MI 48109-1043
UNITED STATES

zieve@umich.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 274 No. 1 March 2015

Unimodal sequences and “strange” functions: a family of quantum modular forms	1
KATHRIN BRINGMANN, AMANDA FOLSOM and ROBERT C. RHOADES	
Congruence primes for Ikeda lifts and the Ikeda ideal	27
JIM BROWN and RODNEY KEATON	
Constant mean curvature, flux conservation, and symmetry	53
NICK EDELEN and BRUCE SOLOMON	
The cylindrical contact homology of universally tight sutured contact solid tori	73
ROMAN GOLOVKO	
Uniform boundedness of S -units in arithmetic dynamics	97
HOLLY KRIEGER, AARON LEVIN, ZACHARY SCHERR, THOMAS TUCKER, YU YASUFUKU and MICHAEL E. ZIEVE	
A counterexample to the energy identity for sequences of α -harmonic maps	107
YUXIANG LI and YOUDE WANG	
Theory of newforms of half-integral weight	125
MURUGESAN MANICKAM, JABAN MEHER and BALAKRISHNAN RAMAKRISHNAN	
Algebraic families of hyperelliptic curves violating the Hasse principle	141
NGUYEN NGOC DONG QUAN	
F -zips with additional structure	183
RICHARD PINK, TORSTEN WEDHORN and PAUL ZIEGLER	
Mean values of L -functions over function fields	237
JEFFREY LIN THUNDER	