

*Pacific  
Journal of  
Mathematics*

**$L^p$  REGULARITY OF WEIGHTED SZEGŐ PROJECTIONS  
ON THE UNIT DISC**

SAMANGI MUNASINGHE AND YUNUS E. ZEYTUNCU

# $L^p$ REGULARITY OF WEIGHTED SZEGŐ PROJECTIONS ON THE UNIT DISC

SAMANGI MUNASINGHE AND YUNUS E. ZEYTUNCU

We present a family of weights on the unit disc for which the corresponding weighted Szegő projection operators are irregular on  $L^p$  spaces. We further investigate the dual spaces of weighted Hardy spaces corresponding to this family.

## 1. Introduction

**1.1. Classical setting.** Let  $\mathbb{D}$  denote the unit disc in  $\mathbb{C}$  and  $\mathbb{T}$  the unit circle. Let  $\mathcal{O}(\mathbb{D})$  denote the set of holomorphic functions on  $\mathbb{D}$ . For  $1 \leq p < \infty$ , the ordinary Hardy space is defined as

$$\mathcal{H}^p(\mathbb{T}) = \{f \in \mathcal{O}(\mathbb{D}) \text{ and } \|f\|_{\mathcal{H}^p} < \infty\},$$

where

$$\|f\|_{\mathcal{H}^p}^p = \sup_{0 \leq r < 1} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta.$$

It is known (see [Duren 1970]) that functions in  $\mathcal{H}^p(\mathbb{T})$  have boundary limits almost everywhere, i.e., for almost every  $\theta \in [0, 2\pi]$

$$f(e^{i\theta}) = \lim_{r \rightarrow 1^-} f(re^{i\theta})$$

exists. Moreover,

$$\|f\|_{L^p(\mathbb{T})} = \|f\|_{\mathcal{H}^p(\mathbb{T})},$$

where  $L^p(\mathbb{T})$  is defined using the standard Lebesgue measure (denoted by  $d\theta$ ) on the unit circle. It is also known that  $\mathcal{H}^p(\mathbb{T})$  is a closed subspace of  $L^p(\mathbb{T})$ . In particular, for  $p = 2$ , the orthogonal projection operator, called the Szegő projection operator exists;

$$S : L^2(\mathbb{T}) \longrightarrow \mathcal{H}^2(\mathbb{T}).$$

*MSC2010:* primary 32A25; secondary 32A35, 42B30.

*Keywords:* Szegő projection,  $A_p$  weights, Weighted Hardy spaces, Dual spaces.

The operator  $S$  is an integral operator with the kernel  $S(z, w)$  (called the Szegő kernel), and for  $f \in L^2(\mathbb{T})$ ,

$$Sf(z) = \int_{\mathbb{T}} S(z, w)f(w) d\theta.$$

It follows from the general theory of reproducing kernels that for any orthonormal basis  $\{e_n(z)\}_{n=0}^\infty$  for  $\mathcal{H}^2(\mathbb{T})$ , the Szegő kernel is given by

$$S(z, w) = \sum_{n=0}^\infty e_n(z)\overline{e_n(w)}.$$

**1.2. Weighted setting.** Let  $g(z)$  be a holomorphic function on  $\mathbb{D}$  that is continuous on  $\overline{\mathbb{D}}$  and has no zeros inside  $\mathbb{D}$ . We set  $\mu(z) = |g(z)|^2$  and define weighted Hardy spaces and weighted Szegő projections using the function  $\mu(z)$  as a weight on  $\mathbb{T}$ .

For  $1 \leq p < \infty$ , we define the weighted Lebesgue and Hardy spaces with respect to  $\mu$  as

$$L^p(\mathbb{T}, \mu) = \{f \text{ measurable function on } \mathbb{D} \text{ and } \|f\|_{p,\mu} < \infty\},$$

where

$$\|f\|_{p,\mu}^p = \int_{\mathbb{T}} |f(w)|^p \mu(w) d\theta = \int_{\mathbb{T}} |f(w)(g(w))^{2/p}|^p d\theta,$$

and

$$\mathcal{H}^p(\mathbb{T}, \mu) = \{f \in \mathcal{O}(\mathbb{D}) \text{ such that } \|f\|_{\mathcal{H}^p,\mu} < \infty\},$$

where

$$\|f\|_{\mathcal{H}^p,\mu}^p = \sup_{0 \leq r < 1} \int_0^{2\pi} |f(re^{i\theta})(g(re^{i\theta}))^{2/p}|^p d\theta.$$

Note that,  $f \in \mathcal{H}^p(\mathbb{T}, \mu)$  implies  $f(z)(g(z))^{2/p} \in \mathcal{H}^p(\mathbb{T})$ , which in turn gives that  $f(z)(g(z))^{2/p}$  has almost everywhere boundary limits. Hence so does  $f(z)$ . Additionally,  $\|f\|_{\mathcal{H}^p,\mu} = \|f\|_{p,\mu}$ . Furthermore,  $L^p(\mathbb{T}, \mu)$  is a Banach space and  $\mathcal{H}^p(\mathbb{T}, \mu)$  is a closed subspace of  $L^p(\mathbb{T}, \mu)$ .

In particular, again when  $p = 2$ , we obtain the weighted Szegő projection

$$S_\mu : L^2(\mathbb{T}, \mu) \longrightarrow \mathcal{H}^2(\mathbb{T}, \mu).$$

Following the similar theory, we note that  $S_\mu$  is an integral operator

$$S_\mu f(z) = \int_{\mathbb{T}} S_\mu(z, w)f(w)\mu(w) d\theta.$$

If  $\{f_n(z)\}_{n=0}^\infty$  is an orthonormal basis for  $\mathcal{H}^2(\mathbb{T}, \mu)$  then

$$S_\mu(z, w) = \sum_{n=0}^\infty f_n(z) \overline{f_n(w)}.$$

We are interested in the action of  $S_\mu$  on  $L^p(\mathbb{T}, \mu)$ . By definition,  $S_\mu$  is a bounded operator from  $L^2(\mathbb{T}, \mu)$  to  $L^2(\mathbb{T}, \mu)$ . The problem we investigate is the boundedness of  $S_\mu$  from  $L^p(\mathbb{T}, \mu)$  to  $L^p(\mathbb{T}, \mu)$  for other values of  $p \in (1, \infty)$ . Note that for any given weight  $\mu$  as above, we can associate an interval  $I_\mu \subset (1, \infty)$  such that  $S_\mu$  is bounded from  $L^p(\mathbb{T}, \mu)$  to  $L^p(\mathbb{T}, \mu)$  if and only if  $p \in I_\mu$ . By definition,  $2 \in I_\mu$ , and by duality and interpolation,  $I_\mu$  is a conjugate symmetric interval around 2. Namely, if some  $p_0 > 2$  is in  $I_\mu$ , so is  $q_0$  where  $1/q_0 + 1/p_0 = 1$ .

In the classical setting, i.e.,  $\mu \equiv 1$ , the Szegő projection operator is bounded from  $L^p(\mathbb{T})$  to  $L^p(\mathbb{T})$  for any  $1 < p < \infty$ , see [Zhu 2007, page 257].

The purpose of this note is to construct weights  $\mu$  on  $\mathbb{T}$  for which the corresponding interval  $I_\mu$  can be any open interval larger than  $\{2\}$  but smaller than  $(1, \infty)$ .

**Theorem 1.** *For any given  $p_0 > 2$ , there exists a weight  $\mu$  on  $\mathbb{T}$  such that  $I_\mu = (q_0, p_0)$  where  $1/q_0 + 1/p_0 = 1$ , i.e., the weighted Szegő projection  $S_\mu$  is bounded on  $L^p(\mathbb{T}, \mu)$  if and only if  $q_0 < p < p_0$ .*

Our proof of this theorem is similar to the proof of the analogous statement for weighted Bergman projections in [Zeytuncu 2013] with modifications from Bergman kernels to Szegő kernels. The main ingredient is the theory of  $A_p$  weights on  $\mathbb{T}$ .

When the weighted Szegő projection  $S_\mu$  is bounded on  $L^p(\mathbb{T}, \mu)$  for some  $p$ , one can identify the dual space of the weighted Hardy space  $\mathcal{H}^p(\mathbb{T}, \mu)$ . However, when  $S_\mu$  fails to be bounded, a different approach is needed to identify the dual spaces. In the third section, we address this issue and describe the dual spaces of weighted Hardy spaces.

The following notation is used in the rest of the note. We write  $f(z) \simeq g(z)$  when  $c \cdot g(z) \leq f(z) \leq C \cdot g(z)$  for some positive constants  $c$  and  $C$  which are independent of  $z$ . Similarly we write  $f(z) \lesssim g(z)$  when  $f(z) \leq C \cdot g(z)$  for some positive constant  $C$ . We use  $d\theta$  for the Lebesgue measure on the unit circle  $\mathbb{T}$ . When we integrate functions (that are also defined on the unit disc) on  $\mathbb{T}$ , instead of writing  $e^{i\theta}$ , we keep  $z$  and  $w$  as the variables.

## 2. Proof of Theorem 1

**2.1. Relation between weighted kernels.** The particular choice of  $\mu(z)$  indicates the following relation between the weighted Szegő kernels  $S_\mu(z, w)$  and the ordinary Szegő kernel  $S(z, w)$ .

**Proposition 2.** For  $\mu(z) = |g(z)|^2$  as above, the following relation holds

$$(1) \quad S(z, w) = g(z)S_\mu(z, w)\overline{g(w)}.$$

*Proof.* Let  $\{e_n(z)\}_{n=0}^\infty$  be an orthonormal basis for  $\mathcal{H}^2(\mathbb{T})$ . Since  $g(z)$  does not vanish inside  $\mathbb{D}$ , each  $e_n(z)/g(z)$  is a holomorphic function on  $\mathbb{D}$  and is in  $\mathcal{H}^2(\mathbb{T}, |g|^2)$  by construction. Following the orthonormal properties of the  $e_n(z)$  we have

$$\left\langle \frac{e_n(z)}{g(z)}, \frac{e_m(z)}{g(z)} \right\rangle_\mu = \langle e_n(z), e_m(z) \rangle = \delta_{n,m},$$

where  $\delta_{n,m}$  is the Kronecker delta.

Also for any  $f$  in  $\mathcal{H}^2(\mathbb{T}, |g|^2)$ ,  $(f \cdot g)$  is in  $\mathcal{H}^2(\mathbb{T})$  and hence can be written as a linear combination of the  $e_n(z)$ . Consequently so can  $f$ , using the quotients  $e_n(z)/g(z)$ . Hence,  $\{e_n(z)/g(z)\}_{n=0}^\infty$  is an orthonormal basis for  $\mathcal{H}^2(\mathbb{T}, |g|^2)$ .

Therefore, using the basis representation of the Szegő kernels we obtain

$$\begin{aligned} S(z, w) &= \sum_{n=0}^\infty e_n(z)\overline{e_n(w)} = g(z) \left( \sum_{n=0}^\infty \frac{e_n(z)}{g(z)} \frac{\overline{e_n(w)}}{g(w)} \right) \overline{g(w)} \\ &= g(z)S_\mu(z, w)\overline{g(w)}. \end{aligned} \quad \square$$

**2.2.  $A_p$  weights on  $\mathbb{T}$ .** For  $p \in (1, \infty)$ , a weight  $\mu$  on  $\mathbb{T}$  is said to be in  $A_p(\mathbb{T})$  if

$$\sup_{I \subset \mathbb{T}} \left( \frac{1}{|I|} \int_I \mu(\theta) d\theta \right) \left( \frac{1}{|I|} \int_I \mu(\theta)^{\frac{-1}{p-1}} d\theta \right)^{p-1} < \infty,$$

where  $I$  denotes intervals in  $\mathbb{T}$ .

These weights are used to characterize the  $L^p$  regularity of the ordinary Szegő projection on weighted spaces. The following result appears in [Garnett 1981] and is used in [Lanzani and Stein 2004, Equation (2.3)] in connection with a conformal map based approach to the investigation of the unweighted Szegő projection for a general domain.

**Theorem 3.** The ordinary Szegő projection  $S$  is bounded from  $L^p(\mathbb{T}, \mu)$  to  $L^p(\mathbb{T}, \mu)$  if and only if  $\mu \in A_p(\mathbb{T})$ .

*Proof.* This result is an immediate consequence of the fact that the Szegő kernel of the unit disc agrees with the Cauchy kernel (see [Kerzman and Stein 1978]) together with the classical weighted theory for the latter, see also [Garnett 1981].  $\square$

The following theorem follows from Equation (1) and Theorem 3.

**Proposition 4.** For  $1 < p < \infty$  and  $\mu(z) = |g(z)|^2$  as above, the following are equivalent.

$$(1) \quad S_\mu \text{ is bounded from } L^p(\mathbb{T}, |g|^2) \text{ to } L^p(\mathbb{T}, |g|^2).$$

- (2)  $\mathcal{S}$  is bounded from  $L^p(\mathbb{T}, |g|^{2-p})$  to  $L^p(\mathbb{T}, |g|^{2-p})$ .  
 (3)  $|g|^{2-p} \in A_p(\mathbb{T})$ .

*Proof.* **Theorem 3** gives the equivalence of (2) and (3). We show the equivalence of (1) and (2). Using the relation between the kernels from the previous proposition, we obtain the following relation between the corresponding operators:

$$g(z)(\mathcal{S}_\mu f)(z) = (\mathcal{S}(f \cdot g))(z) \quad \text{for } f \in L^2(\mathbb{T}, |g|^2).$$

Indeed, suppose (2) is true. Then

$$\begin{aligned} \|\mathcal{S}_\mu f\|_{p, |g|^2}^p &= \int_{\mathbb{T}} |(\mathcal{S}_\mu f)(w)|^p |g(w)|^2 d\theta \\ &= \int_{\mathbb{T}} |(\mathcal{S}(f \cdot g))(w)|^p |g(w)|^{2-p} d\theta = \|\mathcal{S}(f \cdot g)\|_{p, |g|^{2-p}}^p \\ &\lesssim \|f \cdot g\|_{p, |g|^{2-p}}^p = \|f\|_{p, |g|^2}^p, \end{aligned}$$

which proves (1).

Now when (1) is true,

$$\begin{aligned} \|\mathcal{S}f\|_{p, |g|^{2-p}}^p &= \int_{\mathbb{T}} |(\mathcal{S}f)(w)|^p |g(w)|^{2-p} d\theta \\ &= \int_{\mathbb{T}} |(\mathcal{S}_\mu(f/g))(w)|^p |g(w)|^2 d\theta = \|\mathcal{S}_\mu(f/g)\|_{p, |g|^2}^p \\ &\lesssim \|f/g\|_{p, |g|^2}^p = \|f\|_{p, |g|^{2-p}}^p \end{aligned}$$

and hence (2) is true. □

We can now present a family of weights that behave as claimed in **Theorem 1**.

**Theorem 5.** For  $\alpha \geq 0$ , let  $g_\alpha(z) = (z - 1)^\alpha$  and  $\mu_\alpha(z) = |g_\alpha(z)|^2$ . Then the weighted Szegő projection operator  $\mathcal{S}_{\mu_\alpha}$  is bounded on  $L^p(\mathbb{T}, \mu_\alpha)$  if and only if  $p \in (\frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha})$ .

**Remark 6.** **Theorem 5** is a quantitative version of **Theorem 1** and therefore we obtain a proof of **Theorem 1** when we prove **Theorem 5**.

**Remark 7.** Note that as  $\alpha \rightarrow 0^+$  the interval  $(\frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha})$  approaches  $(1, \infty)$  and as  $\alpha \rightarrow \infty$  the interval  $(\frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha})$  approaches  $\{2\}$ . Hence, any conjugate symmetric interval around 2 can be achieved as the boundedness range of a weighted Szegő projection.

*Proof of Theorem 5.* First note that on intervals  $I$  with  $\theta = 0 \notin I$ , the weight  $|g_\alpha(z)|^{2-p} = |z - 1|^{\alpha(2-p)} \simeq C$ . Therefore, both integrals in the  $A_p(\mathbb{T})$  condition are finite and hence so is the supremum over all such intervals when  $p$  is in the given range. On intervals that contain  $z = 0$  we have the following.

*Step 1.* We show that for the weights  $\omega(z) = |g_\alpha(z)|^{2-p} = |z-1|^{\alpha(2-p)}$ , the second integral in the  $A_p(\mathbb{T})$  condition diverges for arcs  $I = (-\epsilon, \epsilon)$  if and only if  $p$  is outside the given region.

For intervals  $I = (-\epsilon, \epsilon)$  with small  $\epsilon$  and  $p \leq \frac{2\alpha+1}{\alpha+1}$ ,

$$\begin{aligned} \int_I \omega(z)^{\frac{1}{1-p}} d\theta &= \int_{-\epsilon}^{\epsilon} |e^{i\theta} - 1|^{\frac{\alpha(2-p)}{(1-p)}} d\theta \\ &= \int_{-\epsilon}^{\epsilon} (\sqrt{2(1 - \cos(\theta))})^{\frac{\alpha(2-p)}{(1-p)}} d\theta \\ &\simeq \int_{-\epsilon}^{\epsilon} \theta^{\frac{\alpha(2-p)}{(1-p)}} d\theta = \infty, \end{aligned}$$

because  $\alpha(2-p)/(1-p) \leq -1$ . Hence  $\omega \notin A_p(\mathbb{T})$  for such  $p$ .

Also, when  $p \geq \frac{2\alpha+1}{\alpha}$ ,

$$\int_I \omega(z) d\theta \simeq \int_{-\epsilon}^{\epsilon} \theta^{\alpha(2-p)} d\theta = \infty,$$

because  $\alpha(2-p) \leq -1$ . Hence  $\omega \notin A_p(\mathbb{T})$  for  $p \geq \frac{2\alpha+1}{\alpha}$  either.

The same calculations show convergence of all integrals for  $p$  in the desired range.

*Step 2.* We show that for  $p \in (\frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha})$  the integrals in the  $A_p$  condition are finite over any (general) interval  $I = (\theta_0 - R, \theta_0 + R)$  with  $\theta_0 \neq 0$ . We consider two cases.

*Case 1.*  $I \cap \text{Arc}(0, 2R) = \emptyset$ .

On such intervals  $I$ ,  $3R < \theta_0$  and so  $2\theta_0/3 \leq \theta_0 - R \leq \theta \leq \theta_0 + R \leq 4\theta_0/3$  giving  $\theta \simeq \theta_0$ . So,  $\omega = |z-1|^{\alpha(2-p)} \simeq \theta_0^{\alpha(2-p)}$ . Therefore,

$$\frac{1}{|I|} \int_I \omega(z) d\theta \lesssim \frac{1}{2R} \int_I \theta_0^{\alpha(2-p)} d\theta = \theta_0^{\alpha(2-p)}.$$

and

$$\begin{aligned} \left( \frac{1}{|I|} \int_I \omega(z)^{\frac{1}{1-p}} d\theta \right)^{p-1} &\lesssim \left( \frac{1}{2R} \int_I \theta_0^{\frac{\alpha(2-p)}{1-p}} d\theta \right)^{p-1} \\ &= \theta_0^{-\alpha(2-p)}. \end{aligned}$$

Hence the supremum over all such intervals is finite.

*Case 2.*  $I \cap \text{Arc}(0, 2R) \neq \emptyset$ .

In this case, since  $I \subset \text{Arc}(0, 4R)$  and  $\alpha(2-p) + 1 > 0$  when  $2\alpha + 1/\alpha > p$ , we have

$$\frac{1}{|I|} \int_I \omega(z) d\theta \simeq \frac{2}{8R} \int_0^{4R} \theta^{\alpha(2-p)} d\theta = \frac{1}{4R} \frac{\theta^{\alpha(2-p)+1}}{\alpha(2-p)+1} \Big|_0^{4R} = \frac{4R^{\alpha(2-p)}}{\alpha(2-p)+1}.$$

Also since  $\alpha(2 - p)/(1 - p) + 1 > 0$  when  $2\alpha + 1/(\alpha + 1) < p$ ,

$$\begin{aligned} \left( \frac{1}{|I|} \int_I \omega(z)^{\frac{1}{1-p}} d\theta \right)^{p-1} &\simeq \left( \frac{2}{8R} \int_0^{4R} \theta^{\frac{\alpha(2-p)}{1-p}} d\theta \right)^{p-1} = \left( \frac{1}{4R} \frac{\theta^{\frac{\alpha(2-p)}{1-p} + 1}}{\frac{\alpha(2-p)}{1-p} + 1} \right)^{p-1} \\ &\simeq \left( \frac{R^{\frac{\alpha(2-p)}{1-p}}}{\frac{\alpha(2-p)}{1-p} + 1} \right)^{p-1} = \frac{2R^{-\alpha(2-p)}}{\left( \frac{\alpha(2-p)}{1-p} + 1 \right)^{p-1}}. \end{aligned}$$

Therefore, the supremum over all intervals of the type in case 2 are also finite and  $\omega = |g|^{2-p} \in A^p(\mathbb{T})$  if and only if  $p \in \left( \frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha} \right)$ . □

**Remark 8.** The analog of [Theorem 1](#) for domains in  $\mathbb{C}^n$  ( $n \geq 2$ ) is an open problem. See [\[Békollé and Bonami 1995\]](#) for a partial result. Also see [\[Lanzani and Stein 2013\]](#) for the regularity on strongly pseudoconvex domains.

### 3. Duality

In this section, we investigate the duals of Hardy spaces corresponding to weights from the previous section. For  $\alpha \geq 0$  and  $\mu_\alpha(z) = |z - 1|^{2\alpha}$ , a consequence of [Theorem 5](#) is the following.

**Theorem 9.** *Let  $\alpha \geq 0$  and  $\mu_\alpha(z) = |z - 1|^{2\alpha}$ . For any  $p \in \left( \frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha} \right)$ , the dual space of the weighted Hardy space  $\mathcal{H}^p(\mathbb{T}, |z - 1|^{2\alpha})$  can be identified with  $\mathcal{H}^q(\mathbb{T}, |z - 1|^{2\alpha})$ , where  $1/p + 1/q = 1$ , under the pairing*

$$\langle f, h \rangle = \int_{\mathbb{T}} f(z) \overline{h(z)} |z - 1|^{2\alpha} d\theta.$$

*Proof.* This is a standard argument; however, we present a proof here for completeness. For a given function  $h \in \mathcal{H}^q(\mathbb{T}, |z - 1|^{2\alpha})$ , we define a linear functional on  $\mathcal{H}^p(\mathbb{T}, |z - 1|^{2\alpha})$  by

$$\mathbf{G}(f) = \int_{\mathbb{T}} f(z) \overline{h(z)} |z - 1|^{2\alpha} d\theta.$$

It is clear that, by Hölder’s inequality,  $\mathbf{G}$  is a bounded functional with operator norm less than  $\|h\|_{\mathcal{H}^q(\mathbb{T}, |z-1|^{2\alpha})}$ .

Conversely, let  $\mathbf{G}$  be a bounded linear functional on  $\mathcal{H}^p(\mathbb{T}, |z - 1|^{2\alpha})$ . By the Hahn–Banach theorem,  $\mathbf{G}$  extends to a bounded linear functional on  $L^p(\mathbb{T}, |z - 1|^{2\alpha})$ . Now using the duality of  $L^p$  spaces, we find a function  $h \in L^q(\mathbb{T}, |z - 1|^{2\alpha})$  such that

$$\mathbf{G}(f) = \int_{\mathbb{T}} f(z) \overline{h(z)} |z - 1|^{2\alpha} dz \quad \text{for } f \in L^p(\mathbb{T}, |z - 1|^{2\alpha}).$$



When we restrict  $\mathbf{G}$  to  $L^p(\mathbb{T}, |z - 1|^{2\alpha}) \cap \mathcal{H}^2(\mathbb{T}, |z - 1|^{2\alpha})$  and use self-adjointness of  $\mathbf{S}_{\mu_\alpha}$  we get the following.

$$\begin{aligned} \mathbf{G}(f) &= \int_{\mathbb{T}} f(z)\overline{h(z)}|z - 1|^{2\alpha} d\theta \\ &= \int_{\mathbb{T}} (\mathbf{S}_{\mu_\alpha} f)(z)\overline{h(z)}|z - 1|^{2\alpha} d\theta \\ &= \int_{\mathbb{T}} f(z)\overline{(\mathbf{S}_{\mu_\alpha} h)(z)}|z - 1|^{2\alpha} d\theta \end{aligned}$$

for  $f \in L^p(\mathbb{T}, |z - 1|^{2\alpha}) \cap \mathcal{H}^2(\mathbb{T}, |z - 1|^{2\alpha})$ .

Since the intersection of these two spaces is dense in  $\mathcal{H}^p(\mathbb{T}, |z - 1|^{2\alpha})$ , we note that  $\mathbf{G}$  is represented by the function  $(\mathbf{S}_{\mu_\alpha} h)(z)$  and  $\mathbf{S}_{\mu_\alpha} h \in \mathcal{H}^q(\mathbb{T}, |z - 1|^{2\alpha})$  by [Theorem 5](#). □

A natural question arises after this statement. How can we identify the dual space of the weighted Hardy space,  $\mathcal{H}^p(\mathbb{T}, |z - 1|^{2\alpha})$ , when  $p \notin (\frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha})$ ?

The answer to this question follows from the following result on the boundedness of the weighted Szegő projection,  $\mathbf{S}_{\mu_\alpha}$ . Similar results for weighted Bergman projections have been presented recently in [\[Arroussi and Pau 2014\]](#) and [\[Constantin and Peláez 2015\]](#).

**Proposition 10.** *Let  $\alpha \geq 0$  and  $\mu_\alpha = |z - 1|^{2\alpha}$ . For any  $1 < p < \infty$ , the weighted Szegő projection  $\mathbf{S}_{\mu_\alpha}$  is bounded on  $L^p(\mathbb{T}, |z - 1|^{\alpha p})$ .*

**Remark 11.** Note that as  $p$  varies, changes occur not only in the integrability scale but also in the measure.

*Proof.* The proof follows from the relation between the kernels in [Proposition 2](#) and the fact that the unweighted Szegő projection  $\mathbf{S}$  is bounded on  $L^p(\mathbb{T})$  for  $1 < p < \infty$ .

Let us take  $f(z) \in L^p(\mathbb{T}, |z - 1|^{\alpha p})$  and set

$$\tilde{f}(z) = f(z) \frac{|z - 1|^{2\alpha}}{(z - 1)^\alpha},$$

then we have  $\tilde{f} \in L^p(\mathbb{T})$ . Using this notation, we notice

$$\begin{aligned} \mathbf{S}_{\mu_\alpha} f(z) &= \int_{\mathbb{T}} S_{\mu_\alpha}(z, w) f(w) |w - 1|^{2\alpha} d\theta \\ &= \frac{(z - 1)^\alpha}{(z - 1)^\alpha} \int_{\mathbb{T}} S_{\mu_\alpha}(z, w) (w - 1)^\alpha f(w) \frac{|w - 1|^{2\alpha}}{(w - 1)^\alpha} d\theta \\ &= \frac{1}{(z - 1)^\alpha} \int_{\mathbb{T}} S(z, w) \tilde{f}(w) d\theta \\ &= \frac{1}{(z - 1)^\alpha} \mathbf{S}(\tilde{f}(w))(z), \end{aligned}$$

where we invoke [Proposition 2](#) when we pass from the second to the third line. Next by using the fact that the unweighted Szegő projection operator  $S$  is bounded on  $L^p(\mathbb{T})$ , we obtain the following.

$$\begin{aligned} \|S_{\mu_\alpha} f\|_{L^p(\mathbb{T}, |z-1|^{\alpha p})} &= \int_{\mathbb{T}} |z-1|^{\alpha p} \frac{1}{|z-1|^{\alpha p}} |S(\tilde{f}(w))(z)|^p d\theta \\ &= \|S(\tilde{f}(w))\|_{L^p(\mathbb{T})}^p \lesssim \|\tilde{f}(w)\|_{L^p(\mathbb{T})}^p \\ &= \left\| f(w) \frac{|w-1|^{2\alpha}}{(w-1)^\alpha} \right\|_{L^p(\mathbb{T})}^p \\ &= \|f\|_{L^p(\mathbb{T}, |z-1|^{\alpha p})}^p. \end{aligned}$$

This finishes the proof of the proposition. □

Now we can answer the duality question by using [Proposition 10](#). Following the same argument as in the proof of [Theorem 9](#), we obtain the following statement.

**Theorem 12.** *Let  $\alpha \geq 0$  and  $\mu_\alpha = |z-1|^{2\alpha}$ . Then for any  $p \in (1, \infty)$ , the dual space of the weighted Hardy space  $\mathcal{H}^p(\mathbb{T}, |z-1|^{\alpha p})$  can be identified with  $\mathcal{H}^q(\mathbb{T}, |z-1|^{\alpha q})$ , where  $1/p + 1/q = 1$ , under the pairing*

$$\langle f, h \rangle = \int_{\mathbb{T}} f(z) \overline{h(z)} |z-1|^{2\alpha} d\theta.$$

At first, the two duality results in [Theorem 9](#) and [Theorem 12](#) may seem confusing for  $p \in (\frac{2\alpha+1}{\alpha+1}, \frac{2\alpha+1}{\alpha})$ . However, the main point is to note the difference in the exponents of the weights and the way the pairing is defined. We illustrate these two results in the following example.

**Example 13.** Let us take  $\alpha = 1/2$ . Then  $S_{|z-1|}$  is bounded on  $L^p(\mathbb{T}, |z-1|)$  for  $p \in (4/3, 4)$ . In particular, for any  $p \in (4/3, 4)$ , the dual space of  $\mathcal{H}^p(\mathbb{T}, |z-1|)$  can be identified with  $\mathcal{H}^q(\mathbb{T}, |z-1|)$ , where  $1/p + 1/q = 1$ , under the pairing

$$\langle f, h \rangle_{|z-1|} = \int_{\mathbb{T}} f(z) \overline{h(z)} |z-1| d\theta.$$

On the other hand, using the second duality result for any  $p > 1$ , the dual space of  $\mathcal{H}^p(\mathbb{T}, |z-1|)$  can be identified with  $\mathcal{H}^q(\mathbb{T}, |z-1|^{q/p})$  when  $1/p + 1/q = 1$ , under the pairing

$$\langle f, h \rangle_{|z-1|^{2/p}} = \int_{\mathbb{T}} f(z) \overline{h(z)} |z-1|^{2/p} d\theta.$$

### Acknowledgments

We thank the anonymous referee for constructive comments on the proofs of [Proposition 10](#) and [Theorem 12](#) and also for the useful editorial remarks on the

exposition of the article. This study started when the first author visited the second author at Texas A&M University; we thank the Department of Mathematics for the hospitality. The first author also wishes to acknowledge the support from the WKU Office of Research for this visit.

## References

- [Arroussi and Pau 2014] H. Arroussi and J. Pau, “Reproducing kernel estimates, bounded projections and duality on large weighted Bergman spaces”, *J. Geom. Anal.* (online publication July 2014).
- [Békollé and Bonami 1995] D. Békollé and A. Bonami, “Estimates for the Bergman and Szegő projections in two symmetric domains of  $\mathbb{C}^n$ ”, *Colloq. Math.* **68**:1 (1995), 81–100. MR 95k:32022 Zbl 0863.47018
- [Constantin and Peláez 2015] O. Constantin and J. Á. Peláez, “Boundedness of the Bergman projection on  $L^p$ -spaces with exponential weights”, *Bull. Sci. Math.* **139**:3 (2015), 245–268. MR 3336683 Zbl 06352015
- [Duren 1970] P. L. Duren, *Theory of  $H^p$  spaces*, Pure and Applied Mathematics **38**, Academic Press, New York, 1970. MR 42 #3552 Zbl 0215.20203
- [Garnett 1981] J. B. Garnett, *Bounded analytic functions*, Pure and Applied Mathematics **96**, Academic Press, New York, 1981. MR 83g:30037 Zbl 0469.30024
- [Kerzman and Stein 1978] N. Kerzman and E. M. Stein, “The Cauchy kernel, the Szegő kernel, and the Riemann mapping function”, *Math. Ann.* **236**:1 (1978), 85–93. MR 58 #6199 Zbl 0419.30012
- [Lanzani and Stein 2004] L. Lanzani and E. M. Stein, “Szegő and Bergman projections on non-smooth planar domains”, *J. Geom. Anal.* **14**:1 (2004), 63–86. MR 2004m:30063 Zbl 1046.30023
- [Lanzani and Stein 2013] L. Lanzani and E. M. Stein, “Cauchy-type integrals in several complex variables”, *Bull. Math. Sci.* **3**:2 (2013), 241–285. MR 3084008 Zbl 1277.32002
- [Zeytuncu 2013] Y. E. Zeytuncu, “ $L^p$  regularity of weighted Bergman projections”, *Trans. Amer. Math. Soc.* **365**:6 (2013), 2959–2976. MR 3034455 Zbl 1278.32007
- [Zhu 2007] K. Zhu, *Operator theory in function spaces*, 2nd ed., Mathematical Surveys and Monographs **138**, American Mathematical Society, Providence, RI, 2007. MR 2008i:47064 Zbl 1123.47001

Received April 14, 2014. Revised October 29, 2014.

SAMANGI MUNASINGHE  
DEPARTMENT OF MATHEMATICS  
WESTERN KENTUCKY UNIVERSITY  
BOWLING GREEN, KY 42101  
UNITED STATES  
[samangi.munasinghe@wku.edu](mailto:samangi.munasinghe@wku.edu)

YUNUS E. ZEYTUNCU  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF MICHIGAN-DEARBORN  
DEARBORN, MI 48128  
UNITED STATES  
[zeytuncu@umich.edu](mailto:zeytuncu@umich.edu)

# PACIFIC JOURNAL OF MATHEMATICS

[msp.org/pjm](http://msp.org/pjm)

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 276    No. 2    August 2015

---

Free evolution on algebras with two states, II MICHAEL ANSHELEVICH	257
Systems of parameters and holonomicity of $A$ -hypergeometric systems CHRISTINE BERKESCH ZAMAERE, STEPHEN GRIFFETH and EZRA MILLER	281
Complex interpolation and twisted Hilbert spaces FÉLIX CABELLO SÁNCHEZ, JESÚS M. F. CASTILLO and NIGEL J. KALTON	287
The ramification group filtrations of certain function field extensions JEFFREY A. CASTAÑEDA and QINGQUAN WU	309
A mean field type flow, II: Existence and convergence JEAN-BAPTISTE CASTÉRAS	321
Isometric embedding of negatively curved complete surfaces in Lorentz–Minkowski space BING-LONG CHEN and LE YIN	347
The complex Monge–Ampère equation on some compact Hermitian manifolds JIANCHUN CHU	369
Topological and physical link theory are distinct ALEXANDER COWARD and JOEL HASS	387
The measures of asymmetry for coproducts of convex bodies QI GUO, JINFENG GUO and XUNLI SU	401
Regularity and analyticity of solutions in a direction for elliptic equations YONGYANG JIN, DONGSHENG LI and XU-JIA WANG	419
On the density theorem for the subdifferential of convex functions on Hadamard spaces MINA MOVAHEDI, DARYOUSH BEHMARDI and SEYEDEHSOMAYEH HOSSEINI	437
$L^p$ regularity of weighted Szegő projections on the unit disc SAMANGI MUNASINGHE and YUNUS E. ZEYTUNCU	449
Topology of complete Finsler manifolds admitting convex functions SORIN V. SABAU and KATSUHIRO SHIOHAMA	459
Variations of the telescope conjecture and Bousfield lattices for localized categories of spectra F. LUKE WOLCOTT	483