

*Pacific
Journal of
Mathematics*

JET SCHEMES OF THE CLOSURE OF NILPOTENT ORBITS

ANNE MOREAU AND RUPERT WEI TZE YU

JET SCHEMES OF THE CLOSURE OF NILPOTENT ORBITS

ANNE MOREAU AND RUPERT WEI TZE YU

We study in this paper the jet schemes of the closure of nilpotent orbits in a finite-dimensional complex reductive Lie algebra. For the nilpotent cone, which is the closure of the regular nilpotent orbit, all the jet schemes are irreducible. This was first observed by Eisenbud and Frenkel, and follows from a strong result of Mustață (2001). Using induction and restriction of “little” nilpotent orbits in reductive Lie algebras, we show that for a large number of nilpotent orbits, the jet schemes of their closures are reducible. As a consequence, we obtain certain geometric properties of these nilpotent orbit closures.

1. Introduction	137
2. Generalities on jet schemes	142
3. Nilpotent orbit closures	148
4. Little nilpotent orbits	152
5. Induced nilpotent orbits	156
6. Consequence of Theorem 5.6	163
7. Applications, remarks and comments	166
Appendix A. Nilpotent orbits in classical simple Lie algebras	170
Appendix B. Statistics in types B, C, and D	172
Appendix C. Tables for exceptional types	173
Acknowledgments	181
References	181

1. Introduction

Throughout this paper, the ground field will be the field \mathbb{C} of complex numbers. We shall work with the Zariski topology, and by *variety* we mean a reduced, irreducible, and separated scheme of finite type over \mathbb{C} .

For X a scheme of finite type over \mathbb{C} and $m \in \mathbb{N}$, we denote by $\mathcal{J}_m(X)$ the m -th jet scheme of X . It is a scheme of finite type over \mathbb{C} whose \mathbb{C} -valued points are naturally in bijection with the $\mathbb{C}[t]/(t^{m+1})$ -valued points of X ; see, e.g., [Mustață 2001; Ein and Mustață 2009; Ishii 2011]. We have $\mathcal{J}_0(X) \simeq X$ and $\mathcal{J}_1(X) \simeq TX$,

MSC2010: 14L30, 17B20, 17B08.

Keywords: nilpotent orbits, jet schemes, induction.

where TX is the total tangent bundle of X ; see [Section 2](#) for more details about generalities on jet schemes. From Nash [\[1995\]](#), it is known that the geometry of the jet schemes is deeply related to the singularities of X . As an illustration of that phenomenon, we have the following result, first conjectured by Eisenbud and Frenkel [\[Mustață 2001, Introduction\]](#), which will be important for us.

Theorem 1 [\[Mustață 2001, Theorem 1\]](#). *Let X be an irreducible scheme of finite type over \mathbb{C} . If X is locally a complete intersection, then $\mathcal{J}_m(X)$ is irreducible for every $m \in \mathbb{N}$ if and only if X has rational singularities.*

According to Kolchin [\[1973\]](#), in contrast to the above theorem, the arc space $\mathcal{J}_\infty(X) = \varinjlim \mathcal{J}_m(X)$ of X is always irreducible when X is irreducible. In this paper, we shall be interested in the irreducibility of the jet schemes for the closure of nilpotent orbits in a complex reductive Lie algebra.

Let G be a complex connected reductive algebraic group, \mathfrak{g} its Lie algebra, and $\mathcal{N}(\mathfrak{g})$ the nilpotent cone of \mathfrak{g} . It is the subscheme of \mathfrak{g} associated to the augmentation ideal of $\mathbb{C}[\mathfrak{g}]^G$. It is a finite union of nilpotent G -orbits, and there is a unique nilpotent orbit of \mathfrak{g} , called the regular nilpotent orbit and denoted by \mathcal{O}_{reg} , such that $\mathcal{N}(\mathfrak{g}) = \overline{\mathcal{O}_{\text{reg}}}$.

According to Kostant [\[1963\]](#), the nilpotent cone is a complete intersection which is irreducible, reduced, and normal. Furthermore, by [\[Hesselink 1976\]](#), it has rational singularities. Hence by [Theorem 1](#), the jet scheme $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ is irreducible for every $m \geq 1$. In fact, by [\[Mustață 2001, Propositions 1.4 and 1.5\]](#), $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ is also a complete intersection which is reduced for every $m \geq 1$.

In [\[op. cit., Appendix\]](#), Eisenbud and Frenkel used these results to extend certain results of Kostant [\[1963\]](#) in the setting of jet schemes. In particular, they proved that $\mathbb{C}[\mathcal{J}_m(\mathfrak{g})]$ is free over the ring $\mathbb{C}[\mathcal{J}_m(\mathfrak{g})]^{\mathcal{J}_m(G)}$ of $\mathcal{J}_m(G)$ -invariants of $\mathbb{C}[\mathcal{J}_m(\mathfrak{g})]$.

Other nilpotent orbit closures do not share these geometric properties in general. Indeed, according to a recent result of Namikawa [\[2013\]](#), for a nonzero and non-regular nilpotent orbit \mathcal{O} , $\overline{\mathcal{O}}$ is not a complete intersection. In addition, $\overline{\mathcal{O}}$ does not always have rational singularities since it is not always normal; see, e.g., [\[Levasseur and Smith 1988; Kraft and Procesi 1982; Kraft 1989; Broer 1998; Sommers 2003\]](#).

Thus, it is quite natural to ask the following question.

Question 1. Let \mathcal{O} be a nilpotent orbit of \mathfrak{g} , and $m \in \mathbb{N}^*$. Is $\mathcal{J}_m(\overline{\mathcal{O}})$ irreducible?

Answering [Question 1](#) is the main purpose of this paper. For the zero orbit and the regular orbit, the answer is positive for every $m \in \mathbb{N}$. Outside these extreme cases, we will see that these jet schemes are rarely irreducible.

Motivations. Since $\overline{\mathcal{O}}$ is not a complete intersection for \mathcal{O} nonzero and nonregular, [Theorem 1](#) cannot be applied directly to answer [Question 1](#). Very recently, Brion and Fu [\[2015\]](#) gave another proof of Namikawa's result, which is more uniform and

slightly shorter. An interesting question, posed by Michel Brion to the first author, is whether jet schemes can be used to provide another proof of Namikawa's result.

Let us explain how we can tackle this problem using jet schemes. Let \mathcal{O} be a nilpotent orbit of \mathfrak{g} . The singular locus of $\bar{\mathcal{O}}$ is exactly $\bar{\mathcal{O}} \setminus \mathcal{O}$. This follows from [Kaledin 2006, Lemma 1.4; Panyushev 1991]; see also [Henderson 2014, Section 2] for a recent review. Moreover, we have

$$\text{codim}_{\bar{\mathcal{O}}}(\bar{\mathcal{O}} \setminus \mathcal{O}) \geq 2.$$

For the nilpotent cone, we have precisely $\text{codim}_{\mathcal{N}(\mathfrak{g})}(\mathcal{N}(\mathfrak{g}) \setminus \mathcal{O}_{\text{reg}}) = 2$, and the equality $\mathcal{N}(\mathfrak{g})_{\text{reg}} = \mathcal{O}_{\text{reg}}$ is a consequence of [Kostant 1963, Theorem 9] (thus the notation \mathcal{O}_{reg} does not bear any confusion).

So, if we assume that $\bar{\mathcal{O}}$ is a complete intersection, then $\bar{\mathcal{O}}$ is normal and so it has rational singularities by [Hinich 1991] or [Panyushev 1991]. Hence, in that event, Mustață's Theorem implies that $\mathcal{J}_m(\bar{\mathcal{O}})$ is irreducible for every $m \geq 1$. So if we can show that $\mathcal{J}_m(\bar{\mathcal{O}})$ is reducible for some $m \geq 1$, then we would obtain a contradiction.¹ The above was our original motivation to look into [Question 1](#).

It may happen that a variety X is not a complete intersection, that X has rational singularities, and that nonetheless $\mathcal{J}_m(X)$ is irreducible for every $m \geq 1$. The cone over the Segre embedding

$$\mathbb{P}^1 \times \mathbb{P}^{n-1} \hookrightarrow \mathbb{P}^{2n-1}, \quad n \geq 2,$$

shows that this situation is possible; see [Mustață 2001, Example 4.7]. We do not know so far whether this situation may happen in the context of nilpotent orbit closures.

More generally, following Nash's philosophy, it would be interesting to understand what kind of properties on the singularities of $\bar{\mathcal{O}}$ we can deduce from the study of $\mathcal{J}_m(\bar{\mathcal{O}})$, $m \geq 1$. The fact that $\bar{\mathcal{O}}$ is not a complete intersection (with \mathcal{O} nonzero and nonregular) whenever $\mathcal{J}_m(\bar{\mathcal{O}})$ is reducible for some $m \geq 1$ is one illustration of such a phenomenon.

Nilpotent orbit closures also form an interesting family of varieties, providing examples and counterexamples in the context of jet schemes. For instance, [Examples 7.6](#) and [7.7](#) illustrate that the locally complete intersection hypothesis cannot be removed from [Lemma 2.7\(3\)](#), and [Theorem 2.8\(3\)](#). Another example is that the normality is not conserved when we pass to jet schemes. By Kostant, the nilpotent cone $\mathcal{N}(\mathfrak{g})$ is normal, and we show in [Proposition 7.3](#) that $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$, $m \geq 1$, is not normal for a simple Lie algebra \mathfrak{g} .

¹There are other approaches that use jet schemes to show that $\bar{\mathcal{O}}$ is not a complete intersection; see [Example 7.2](#).

Main results. Let us describe the main techniques used to study [Question 1](#) and summarize the main results of the paper. To avoid technical details, we shall assume here that \mathfrak{g} is simple.

Let X be an irreducible variety, and $m \in \mathbb{N}$. Then $\mathcal{J}_m(X)$ is irreducible if and only if

$$\pi_{X,m}^{-1}(X_{\text{sing}}) \subset \overline{\pi_{X,m}^{-1}(X_{\text{reg}})},$$

where $\pi_{X,m} : \mathcal{J}_m(X) \rightarrow X$ is the canonical projection from $\mathcal{J}_m(X)$ onto X (see [Section 2](#)), X_{reg} is the smooth part of X , and X_{sing} is its complement (see [Lemma 2.7](#)). This is our starting point.

For \mathcal{O} a nilpotent orbit of \mathfrak{g} , the singular locus of $\overline{\mathcal{O}}$ is $\overline{\mathcal{O}} \setminus \mathcal{O}$ (see [Section 3](#)). The above criterion leads us to the following two conditions which will be central in our paper (see [Definition 3.3](#)).

Definition 1. Let \mathcal{O} be a nilpotent orbit of \mathfrak{g} .

- (1) We say that \mathcal{O} verifies RC_1 if $\pi_{\overline{\mathcal{O}},1}^{-1}(0)$ is not contained in the closure of $\pi_{\overline{\mathcal{O}},1}^{-1}(\mathcal{O})$.
- (2) Let $m \in \mathbb{N}^*$. We say that \mathcal{O} verifies $\text{RC}_2(m)$ if for some nilpotent orbit \mathcal{O}' contained in $\overline{\mathcal{O}} \setminus \mathcal{O}$, we have $\dim \pi_{\overline{\mathcal{O}},m}^{-1}(\mathcal{O}') \geq \dim \pi_{\overline{\mathcal{O}},m}^{-1}(\mathcal{O})$.

Here the letters RC stand for “reducibility condition”.

It follows readily (see [Lemma 3.4](#)) that if a nilpotent orbit \mathcal{O} of \mathfrak{g} verifies RC_1 , then $\mathcal{J}_1(\overline{\mathcal{O}})$ is reducible. Similarly, if a nilpotent orbit \mathcal{O} of \mathfrak{g} verifies $\text{RC}_2(m)$ for some $m \in \mathbb{N}^*$, then $\mathcal{J}_m(\overline{\mathcal{O}})$ is reducible.

We have a characterization for the condition RC_1 (see [Proposition 3.6](#)) which allows us, for example, to show that the nilpotent orbits of $\mathfrak{sl}_{2p}(\mathbb{C})$, with $p \geq 2$, associated with partitions of the form (2^p) verify RC_1 (see [Example 3.7](#)). Note that these orbits do not verify $\text{RC}_2(1)$ (see again [Example 3.7](#)).

A nilpotent orbit \mathcal{O} is called *little* if $0 < 2 \dim \mathcal{O} \leq \dim \mathfrak{g}$ (see [Definition 4.1](#)). For example, the minimal nilpotent orbit of \mathfrak{g} is little (see [Corollary 4.3](#)), and the nilpotent orbits of $\mathfrak{sl}_n(\mathbb{C})$ associated with partitions of the form $(2^p, 1^q)$, with $p, q \in \mathbb{N}^*$, are little (see [Example 4.4](#)). There are many other examples (see [Section 4](#)). Little nilpotent orbits verify both RC_1 and $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$ (see [Proposition 4.2](#)), and they turn out to be useful to study the reducibility of jet schemes of many other orbits via “restriction” or “induction” of orbits.

Firstly, by “restriction” to some Levi subalgebras of \mathfrak{g} (see [Proposition 4.6](#)), we can obtain from nilpotent orbits \mathcal{O} which verify $0 < 2 \dim \mathcal{O} < \dim \mathfrak{g}$ examples of nilpotent orbits which verify RC_1 (and that are not necessarily little); see [Table 1](#). More precisely, as we shall see (in a slightly more general context) in [Proposition 4.6](#), we have:

Proposition 1. *Let \mathfrak{l} be a Levi subalgebra of \mathfrak{g} with a center of dimension one, and such that $\mathfrak{a} := [\mathfrak{l}, \mathfrak{l}]$ is simple. Denote by A the connected subgroup of G whose*

Lie algebra is \mathfrak{a} . Let e be a nilpotent element of \mathfrak{a} and suppose that the following conditions are satisfied:

- (i) \mathfrak{a} contains a regular semisimple element of \mathfrak{g} ,
- (ii) $2 \dim G \cdot e < \dim \mathfrak{g}$.

Then $A \cdot e$ verifies RC_1 .

Secondly, by “induction”, we can reach from nilpotent orbits of reductive Lie subalgebras of \mathfrak{g} many nilpotent orbits of \mathfrak{g} . Here, we consider induction in the sense of Lusztig and Spaltenstein [1979]. We refer the reader to Section 5 for the precise definition of a nilpotent orbit of \mathfrak{g} induced from another one in some proper Levi subalgebra \mathfrak{l} of \mathfrak{g} . Our next statement says that condition $\text{RC}_2(m)$, for $m \in \mathbb{N}^*$, passes through induction.

Theorem 2. *Let \mathfrak{l} be a Levi subalgebra of \mathfrak{g} , $\mathcal{O}_{\mathfrak{l}}$ a nilpotent orbit of \mathfrak{l} and $\mathcal{O}_{\mathfrak{g}}$ the induced nilpotent orbit of \mathfrak{g} from $\mathcal{O}_{\mathfrak{l}}$. If $\mathcal{O}_{\mathfrak{l}}$ verifies $\text{RC}_2(m)$ for some $m \in \mathbb{N}^*$, then $\mathcal{O}_{\mathfrak{g}}$ also verifies $\text{RC}_2(m)$.*

From this result, we are able to deal with a large number of nilpotent orbits. First of all, any nilpotent orbit induced from a nilpotent orbit that has a little factor verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$ (see Theorem 6.1). In particular, if \mathfrak{g} is not of type A_1 , $B_2 = C_2$, or G_2 , then the subregular nilpotent orbit $\mathcal{O}_{\text{subreg}}$ of \mathfrak{g} verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$ (see Corollary 6.2), and so $\mathcal{I}_m(\overline{\mathcal{O}_{\text{subreg}}})$ is reducible for every $m \in \mathbb{N}^*$.

It turns out that many nilpotent orbits can be induced from a nilpotent orbit that has a little factor. This allows us to obtain the following result when \mathfrak{g} is of type A (see Theorem 6.5).

Theorem 3. *Any nilpotent orbit of $\mathfrak{sl}_n(\mathbb{C})$ associated with a nonrectangular partition of n verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*

For the other simple Lie algebras of classical types, we have the following (see Theorem 6.7).

Theorem 4. *Let $n \in \mathbb{N}^*$, $\lambda = (\lambda_1, \dots, \lambda_t)$ be a partition of n , and $\lambda_{t+1} = 0$. Suppose that there exist $1 \leq k < \ell \leq t$ such that $\lambda_k \geq \lambda_{k+1} + 2$ and $\lambda_{\ell} \geq \lambda_{\ell+1} + 2$.*

- (1) *If \mathcal{O} is a nilpotent orbit of $\mathfrak{so}_n(\mathbb{C})$ whose associated partition is λ , then \mathcal{O} verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*
- (2) *If n is even and \mathcal{O} is a nilpotent orbit of $\mathfrak{sp}_n(\mathbb{C})$ whose associated partition is λ , then \mathcal{O} verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*

While our result in the special linear case is exhaustive relative to induction, in the orthogonal and symplectic cases, other nilpotent orbits can be obtained by induction from a little orbit (see [Theorem 6.7](#) and [Remark 6.8](#)). For a simple Lie algebra of exceptional type, we have a list of nilpotent orbits which can be induced from a little one (see [Appendix C](#)).

Organization of the paper. In [Section 2](#), we state some basic properties on jet schemes with some proofs for the convenience of the reader.

In [Section 3](#), we recall some standard properties of nilpotent orbit closures, and of their jet schemes. We introduce here the two sufficient conditions RC_1 and $\text{RC}_2(m)$, $m \geq 1$, to study the reducibility of these jet schemes, and we state some first properties of these conditions.

[Section 4](#) is devoted to little nilpotent orbits. We show that little nilpotent orbits verify both RC_1 and $\text{RC}_2(m)$ for every $m \geq 1$, and we show how they can be used to prove condition RC_1 via the “restriction” of orbits (see [Proposition 4.6](#)).

In [Section 5](#), we study the induction of nilpotent orbits the sense of Luzstig and Spaltenstein [[1979](#)]. The main result is that condition $\text{RC}_2(m)$, for $m \geq 1$, passes through induction (see [Theorem 5.6](#)). We describe in [Section 6](#) how to use [Theorem 5.6](#) to obtain the reducibility of nilpotent orbit closures in simple Lie algebras according to their Dynkin type. The details of some of the conclusions are presented in [Appendices B](#) and [C](#).

We present in [Section 7](#) some applications of our results to geometric properties of nilpotent orbit closures. We also discuss in this section some open problems.

The standard notations relative to nilpotent orbits in classical simple Lie algebras are gathered together in [Appendix A](#). [Appendix B](#) contains some numerical data for classical simple Lie algebras, and [Appendix C](#) summarizes our conclusions for simple Lie algebras of exceptional type.

2. Generalities on jet schemes

In this section, we present some general facts on jet schemes. Our main references on the topic are [[Mustața 2001](#); [Ein and Mustața 2009](#); [Ishii 2011](#)], and [[de Fernex et al. 2013](#), Chapter 8].

Let X be a scheme of finite type over \mathbb{C} , and $m \in \mathbb{N}$.

Definition 2.1. An m -jet of X is a morphism

$$\text{Spec } \mathbb{C}[t]/(t^{m+1}) \longrightarrow X.$$

The set of all m -jets of X carries the structure of a scheme $\mathcal{J}_m(X)$, called the m -th jet scheme of X . It is a scheme of finite type over \mathbb{C} characterized by the following

functorial property: for every scheme Z over \mathbb{C} , we have

$$\mathrm{Hom}(Z, \mathcal{I}_m(X)) = \mathrm{Hom}(Z \times_{\mathrm{Spec} \mathbb{C}} \mathrm{Spec} \mathbb{C}[t]/(t^{m+1}), X).$$

The \mathbb{C} -points of $\mathcal{I}_m(X)$ are thus the $\mathbb{C}[t]/(t^{m+1})$ -points of X . From [Definition 2.1](#), we have for example that $\mathcal{I}_0(X) \simeq X$ and that $\mathcal{I}_1(X) \simeq \mathrm{TX}$ where TX denotes the total tangent bundle of X .

For $p \in \{0, \dots, m\}$, the canonical projection $\mathbb{C}[t]/(t^{m+1}) \rightarrow \mathbb{C}[t]/(t^{p+1})$ induces a *truncation morphism*,

$$\pi_{X,m,p} : \mathcal{I}_m(X) \rightarrow \mathcal{I}_p(X).$$

We shall simply denote by $\pi_{X,m}$ the morphism $\pi_{X,m,0}$,

$$\pi_{X,m} : \mathcal{I}_m(X) \rightarrow \mathcal{I}_0(X) \simeq X.$$

Also, the canonical injection $\mathbb{C} \hookrightarrow \mathbb{C}[t]/(t^{m+1})$ induces a morphism $\iota_{X,m} : X \rightarrow \mathcal{I}_m(X)$, and we have $\pi_{X,m} \circ \iota_{X,m} = \mathrm{Id}_X$. Hence $\iota_{X,m}$ is injective and $\pi_{X,m}$ is surjective. We shall always view X as a subscheme of $\mathcal{I}_m(X)$.

If $f : X \rightarrow Y$ is a morphism of schemes, then we naturally obtain a morphism $f_m : \mathcal{I}_m(X) \rightarrow \mathcal{I}_m(Y)$ making the following diagram commutative:

$$\begin{array}{ccc} \mathcal{I}_m(X) & \xrightarrow{f_m} & \mathcal{I}_m(Y) \\ \pi_{X,m} \downarrow & & \downarrow \pi_{Y,m} \\ X & \xrightarrow{f} & Y \end{array}$$

Remark 2.2. In the case where X is affine, we have the following explicit description of $\mathcal{I}_m(X)$.

Let $n \in \mathbb{N}^*$ and $X \subset \mathbb{C}^n$ be the affine subscheme defined by an ideal $I = (f_1, \dots, f_r)$ of $\mathbb{C}[x_1, \dots, x_n]$. Thus

$$X = \mathrm{Spec} \mathbb{C}[x_1, \dots, x_n]/I.$$

For $k \in \{1, \dots, r\}$, we extend f_k as a map from $(\mathbb{C}[t]/(t^{m+1}))^n$ to $\mathbb{C}[t]/(t^{m+1})$ via base extension. Then giving a morphism $\gamma : \mathrm{Spec} \mathbb{C}[t]/(t^{m+1}) \rightarrow X$ is equivalent to giving a morphism $\gamma^* : \mathbb{C}[x_1, \dots, x_n]/I \rightarrow \mathbb{C}[t]/(t^{m+1})$, or to giving

$$\gamma^*(x_i) = \sum_{j=0}^m \gamma_i^{(j)} t^j, \quad 1 \leq i \leq n$$

such that for any $k \in \{1, \dots, r\}$,

$$f_k(\gamma^*(x_1), \dots, \gamma^*(x_n)) = 0 \quad \text{in } \mathbb{C}[t]/(t^{m+1}).$$

For $k \in \{1, \dots, r\}$, there exist functions $f_k^{(0)}, \dots, f_k^{(m)}$, which depend only on f , in the variables $\boldsymbol{\gamma} = (\gamma_i^{(j)})$, for $1 \leq i \leq n$ and $0 \leq j \leq m$, such that

$$(1) \quad f_k(\boldsymbol{\gamma}^*(x_1), \dots, \boldsymbol{\gamma}^*(x_n)) = \sum_{j=0}^m f_k^{(j)}(\boldsymbol{\gamma}) t^j.$$

The jet scheme $\mathcal{J}_m(X)$ is then the closed subscheme in $\mathbb{C}^{(m+1)n}$ defined by the ideal generated by the polynomials $f_k^{(j)}$, where $k \in \{1, \dots, r\}$ and $j \in \{0, \dots, m\}$. More precisely,

$$\mathcal{J}_m(X) \simeq \text{Spec } \mathbb{C}[x_1^{(j)}, \dots, x_n^{(j)} : j=0, \dots, m] / (f_k^{(j)} : k=1, \dots, r; j=0, \dots, m).$$

In particular, if X is an n -dimensional vector space, then $\mathcal{J}_m(X) \simeq \mathbb{C}^{(m+1)n}$ and for $p \in \{0, \dots, m\}$, the projection $\mathcal{J}_m(X) \rightarrow \mathcal{J}_p(X)$ corresponds to the projection onto the first $(p+1)n$ coordinates.

Example 2.3. Let us consider a concrete example. Let

$$X = \text{Spec } \mathbb{C}[x, y, z] / (x^2 + yz) \subset \mathbb{C}^3,$$

and let us compute $\mathcal{J}_1(X)$ and $\mathcal{J}_2(X)$. We have

$$\begin{aligned} (x_0 + x_1 t + x_2 t^2)^2 + (y_0 + y_1 t + y_2 t^2)(z_0 + z_1 t + z_2 t^2) \\ = x_0^2 + y_0 z_0 + (2x_0 x_1 + y_0 z_1 + y_1 z_0)t \\ + (2x_0 x_2 + x_1^2 + y_0 z_2 + y_2 z_0 + y_1 z_1)t^2 \pmod{t^3}. \end{aligned}$$

Hence $\mathcal{J}_1(X)$ is the subscheme of

$$\mathcal{J}_1(\mathbb{C}^3) \simeq \mathbb{C}[x_0, y_0, z_0, x_1, y_1, z_1]$$

defined by the ideal

$$(x_0^2 + y_0 z_0, 2x_0 x_1 + y_0 z_1 + y_1 z_0),$$

and $\mathcal{J}_2(X)$ is the subscheme of

$$\mathcal{J}_2(\mathbb{C}^3) \simeq \mathbb{C}[x_0, y_0, z_0, x_1, y_1, z_1, x_2, y_2, z_2]$$

defined by the ideal

$$(x_0^2 + y_0 z_0, 2x_0 x_1 + y_0 z_1 + y_1 z_0, 2x_0 x_2 + x_1^2 + y_0 z_2 + y_1 z_1 + y_2 z_0).$$

We now list some basic properties that we need in the sequel. Their proofs can be found in [Ein and Mustařă 2009, Lemma 2.3, Remarks 2.8 and 2.10].

Lemma 2.4. (1) For every open subset U of X , we have $\mathcal{J}_m(U) = \pi_{X,m}^{-1}(U)$.

(2) For every scheme Y , we have a canonical isomorphism

$$\mathcal{J}_m(X \times Y) \simeq \mathcal{J}_m(X) \times \mathcal{J}_m(Y).$$

- (3) If G is a group scheme over \mathbb{C} , then $\mathcal{J}_m(G)$ is also a group scheme over \mathbb{C} . Moreover, if G acts on X , then $\mathcal{J}_m(G)$ acts on $\mathcal{J}_m(X)$.
- (4) If $f : X \rightarrow Y$ is a smooth surjective morphism between schemes, then f_m is also smooth and surjective for every $m \in \mathbb{N}^*$.

Geometric properties. It is known that the geometry of the jet schemes $\mathcal{J}_m(X)$, $m \geq 1$, is closely linked to that of X . More precisely, we can transport some geometric properties from $\mathcal{J}_m(X)$ to X .

The following proposition gives examples of such phenomena.

Proposition 2.5 [Mumford et al. 1994; Ishii 2011, Theorem 3.5]. *Let $m \in \mathbb{N}^*$. If $\mathcal{J}_m(X)$ is smooth (respectively, irreducible, reduced, normal, locally a complete intersection) for some m , then so is X .*

For smoothness, the converse is true, even with “every m ” instead of “for some m ”. In fact, for smooth varieties, we have the following more precise statement.

Proposition 2.6 [Ein and Mustařa 2009, Corollary 2.12]. *If X is a smooth variety of dimension n , then the truncation morphism $\pi_{m,p}$, for $p \in \{0, \dots, m\}$, is a locally trivial projection with fiber isomorphic to $\mathbb{C}^{(m-p)n}$. In particular, $\mathcal{J}_m(X)$ is a smooth variety of dimension $(m+1)n$.*

For the other properties stated in Proposition 2.5, the converse is not true in general. We refer to [Ishii 2011, §3] for counterexamples. We shall encounter other counterexamples in this paper in the setting of nilpotent orbit closures. In this setting, our main purpose is to study the irreducibility of jet schemes. The following lemma gives a necessary and sufficient condition for the converse of Proposition 2.5 to hold for irreducibility.

We denote by X_{reg} the smooth part of X , and by X_{sing} its complement.

Lemma 2.7. *Assume that X is an irreducible reduced scheme of finite type over \mathbb{C} , and let $m \in \mathbb{N}^*$.*

- (1) $\overline{\pi_{X,m}^{-1}(X_{\text{reg}})}$ is an irreducible component of $\mathcal{J}_m(X)$.
- (2) $\mathcal{J}_m(X)$ is irreducible if and only if $\pi_{X,m}^{-1}(X_{\text{sing}})$ is contained in $\overline{\pi_{X,m}^{-1}(X_{\text{reg}})}$.
- (3) If X is a complete intersection, then $\mathcal{J}_m(X)$ is irreducible if and only if $\dim \pi_{X,m}^{-1}(X_{\text{sing}}) < \dim \pi_{X,m}^{-1}(X_{\text{reg}})$.

In particular, if $\dim \pi_{X,m}^{-1}(X_{\text{sing}}) \geq \dim \overline{\pi_{X,m}^{-1}(X_{\text{reg}})}$, then $\mathcal{J}_m(X)$ is reducible.

Proof. Part (3) is proved in [Mustařa 2001, Proposition 1.4], and parts (1) and (2) follow from its proof. More precisely, since X_{reg} is smooth and irreducible, $\overline{\pi_{X,m}^{-1}(X_{\text{reg}})}$ is an irreducible closed subset of $\mathcal{J}_m(X)$ of dimension $(m+1) \dim X$;

see [Proposition 2.6](#). Then parts (1) and (2) follow easily from the fact that we have the decomposition

$$\mathcal{I}_m(X) = \pi_{X,m}^{-1}(X_{\text{sing}}) \cup \overline{\pi_{X,m}^{-1}(X_{\text{reg}})}$$

of closed subsets, and that $\pi_{X,m}^{-1}(X_{\text{sing}}) \not\supseteq \overline{\pi_{X,m}^{-1}(X_{\text{reg}})}$. □

There are also subtle connections between the geometry of $\mathcal{I}_m(X)$, $m \geq 1$, and the singularities of X which are important for us. In particular, according to [\[Mustařa 2001, Theorem 0.1, Propositions 1.5 and 4.12\]](#), we have the theorem:

Theorem 2.8 (Mustařa). *Let X be an irreducible variety over \mathbb{C} .*

- (1) *If X is locally a complete intersection, then $\mathcal{I}_m(X)$ is irreducible for every $m \geq 1$ if and only if X has rational singularities.*
- (2) *If X is locally a complete intersection and if $\mathcal{I}_m(X)$ is irreducible for some $m \geq 1$, then $\mathcal{I}_m(X)$ is also reduced.*
- (3) *If X is locally a complete intersection, then $(\mathcal{I}_1(X))_{\text{reg}} = \pi_{X,1}^{-1}(X_{\text{reg}})$.*

Let us give an easy counterexample to the converse implication of [Proposition 2.5](#) for normality. This example turns out to be a particular case of a more general situation that will be studied in [Proposition 7.3](#).

Example 2.9. Let X be as in [Example 2.3](#). Then X is a complete intersection and it is normal since the singular locus is reduced to $\{0\}$ which has codimension 2 in X . Next, it is not difficult to verify that $\mathcal{I}_1(X)$ is irreducible, reduced, and that it is a complete intersection. But $\mathcal{I}_1(X)$ is not normal. Indeed, by [Theorem 2.8\(3\)](#),

$$(\mathcal{I}_1(X))_{\text{sing}} = \pi_{X,1}^{-1}(\{0\}) \simeq \{0\} \times \mathbb{C}^3.$$

Hence, the singular locus of $\mathcal{I}_1(X)$ has codimension 1 in $\mathcal{I}_1(X)$ since

$$\dim \mathcal{I}_1(X) = 2 \dim X = 4.$$

Group actions. Let G be a connected algebraic group, acting on a variety X , and $m \in \mathbb{N}$. Denote by

$$\rho : G \times X \rightarrow X, \quad (g, x) \mapsto g \cdot x$$

the corresponding action. As stated in [Lemma 2.4](#), the morphism

$$\rho_m : \mathcal{I}_m(G \times X) \simeq \mathcal{I}_m(G) \times \mathcal{I}_m(X) \rightarrow \mathcal{I}_m(X)$$

defines an action of $\mathcal{I}_m(G)$ on $\mathcal{I}_m(X)$.

Recall that we embed X into $\mathcal{I}_m(X)$ through $\iota_{X,m}$. For $x \in X$, let us denote by G^x the stabilizer of x in G , and for $m \in \mathbb{N}$, we denote by $\mathcal{I}_m(G)^x$ its stabilizer in $\mathcal{I}_m(G)$. The following results are probably standard. Since we have not found any reference, we shall include their proofs.

Lemma 2.10. *Let $x \in X$. Then,*

$$\mathcal{I}_m(G) \cdot x = \mathcal{I}_m(G \cdot x), \quad \mathcal{I}_m(G^x) = \mathcal{I}_m(G)^x, \quad \pi_{G \cdot x, m}^{-1}(G \cdot x) = \mathcal{I}_m(G \cdot x).$$

Proof. The morphism $G \times \{x\} \rightarrow G \cdot x$, $(g, x) \mapsto g \cdot x$ is a submersion at all points of $G \times \{x\}$. Hence, according to [Hartshorne 1977, Chapter III, Proposition 10.4], it is a smooth morphism onto $G \cdot x$. So, by Lemma 2.4(4), the induced morphism $\mathcal{I}_m(G) \times \{x\} \rightarrow \mathcal{I}_m(G \cdot x)$ is also smooth and surjective. Consequently, we have the first equality $\mathcal{I}_m(G) \cdot x = \mathcal{I}_m(G \cdot x)$.

By applying the first equality to the algebraic group G^x , we get $\mathcal{I}_m(G^x) \cdot x = \mathcal{I}_m(G^x \cdot x)$, whence the inclusion $\mathcal{I}_m(G^x) \subset \mathcal{I}_m(G)^x$.

Conversely, let $\gamma : \text{Spec } \mathbb{C}[t]/(t^{m+1}) \rightarrow G$ be an element of $\mathcal{I}_m(G)^x$. Then $\rho_m(\gamma, x) = x$; hence, viewing x as a morphism $x : \text{Spec } \mathbb{C}[t]/(t^{m+1}) \rightarrow X$,

$$\rho(\gamma(\tau), x(\tau)) = x(\tau),$$

where τ is the unique element of $\text{Spec } \mathbb{C}[t]/(t^{m+1})$. Thus $\gamma(\tau) \in G^x$ and $x(\tau) = x$. So we have $\gamma \in \mathcal{I}_m(G^x)$, and the second equality follows.

The third equality is a direct consequence of Lemma 2.4(1) since $G \cdot x$ is open in its closure. □

Let \mathfrak{g} be the Lie algebra of G . We consider now the adjoint action of G on \mathfrak{g} . For the results we present here, we refer the reader to [Mustață 2001, Appendix]. Denote by

$$\mathfrak{g}_m := \mathfrak{g} \otimes_{\mathbb{C}} \mathbb{C}[t]/(t^{m+1})$$

the *generalized Takiff Lie algebra* whose Lie bracket is given by

$$[u \otimes x(t), v \otimes y(t)] = [u, v] \otimes x(t)y(t), \quad u, v \in \mathfrak{g}, \quad x(t), y(t) \in \mathbb{C}[t]/(t^{m+1}).$$

As Lie algebras, we have

$$\mathcal{I}_m(\mathfrak{g}) \simeq \mathfrak{g}_m \simeq \text{Lie}(\mathcal{I}_m(G)).$$

In the sequel, when there is no confusion, we shall use the notations \mathfrak{g}_m and G_m for $\mathcal{I}_m(\mathfrak{g})$ and $\mathcal{I}_m(G)$ respectively. If \mathfrak{a} is a Lie subalgebra of \mathfrak{g} , then $\mathcal{I}_m(\mathfrak{a}) \simeq \mathfrak{a}_m$ is a Lie subalgebra of \mathfrak{g}_m . In particular, for $x \in \mathfrak{g}$, we have $(\mathfrak{g}_m)^x = (\mathfrak{g}^x)_m$, where for any subalgebra \mathfrak{m} of \mathfrak{g}_k , with $k \geq 0$, \mathfrak{m}^x stands for the centralizer of x in \mathfrak{m} .

We can identify \mathfrak{g}_m with $\mathfrak{g}^{m+1} \simeq \mathcal{I}_m(\mathfrak{g})$ as a variety through the map

$$\mathfrak{g}^{m+1} \rightarrow \mathfrak{g}_m, \quad (x_0, x_1, \dots, x_m) \mapsto x_0 + x_1 \otimes t + \dots + x_m \otimes t^m.$$

Let G_m be a connected algebraic group whose Lie algebra is \mathfrak{g}_m . Let $\mathbb{C}[\mathfrak{g}_m]$ be the coordinate ring of \mathfrak{g}_m , and let $\mathbb{C}[\mathfrak{g}_m]^{G_m}$ be the subring of G_m -invariants. We conclude in this section with the following result.

Lemma 2.11. *For $f \in \mathbb{C}[\mathfrak{g}]^G$, the polynomials $f^{(0)}, \dots, f^{(m)}$, defined in [Remark 2.2](#), are elements of $\mathbb{C}[\mathfrak{g}_m]^{G_m}$.*

Proof. This is straightforward from the explicit description of the polynomials $f^{(0)}, \dots, f^{(m)}$ given in [Remark 2.2](#). \square

3. Nilpotent orbit closures

From now on, we let G to be a connected reductive algebraic group over \mathbb{C} , \mathfrak{g} its Lie algebra, and $\mathcal{N}(\mathfrak{g})$ the nilpotent cone of \mathfrak{g} . Recall that $\mathcal{N}(\mathfrak{g})$ is the subscheme of \mathfrak{g} defined by the augmentation ideal of $\mathbb{C}[\mathfrak{g}]^G$, and that $\mathcal{N}(\mathfrak{g}) = \overline{\mathcal{O}_{\text{reg}}}$ where \mathcal{O}_{reg} is the regular nilpotent orbit of \mathfrak{g} (see the introduction). As mentioned there, we are interested in this paper in the irreducibility of jet schemes of the closure of nilpotent orbits.

Recall that for an arbitrary nilpotent orbit \mathcal{O} of \mathfrak{g} , the singular locus of $\overline{\mathcal{O}}$ is $\overline{\mathcal{O}} \setminus \mathcal{O}$ and that $\text{codim}_{\overline{\mathcal{O}}}(\overline{\mathcal{O}} \setminus \mathcal{O}) \geq 2$ (see [Section 1](#)).

Definition 3.1. Let \mathcal{O} be a nonzero nilpotent orbit of \mathfrak{g} . Define $\mathfrak{g}_{\mathcal{O}}$ to be the smallest semisimple ideal of \mathfrak{g} containing \mathcal{O} .

More precisely, if $\mathfrak{g} \simeq \mathfrak{z}(\mathfrak{g}) \times \mathfrak{s}_1 \times \dots \times \mathfrak{s}_m$, with $\mathfrak{z}(\mathfrak{g})$ the center of \mathfrak{g} and $\mathfrak{s}_1, \dots, \mathfrak{s}_m$ the simple factors of \mathfrak{g} , then $\mathcal{O} = \mathcal{O}_1 \times \dots \times \mathcal{O}_m$, with \mathcal{O}_i a nilpotent orbit of \mathfrak{s}_i for $i = 1, \dots, m$, and

$$\mathfrak{g}_{\mathcal{O}} = \mathfrak{s}_{i_1} \times \dots \times \mathfrak{s}_{i_k},$$

where $\{i_1, \dots, i_k\}$ is the set of integers $j \in \{1, \dots, m\}$ such that \mathcal{O}_j is nonzero. In particular, if \mathcal{O} is zero, then $\mathfrak{g}_{\mathcal{O}} = 0$, and if \mathcal{O} is nonzero and \mathfrak{g} is simple, then $\mathfrak{g}_{\mathcal{O}} = \mathfrak{g}$.

For \mathcal{O} a nilpotent orbit of \mathfrak{g} , we denote by $\mathcal{I}_{\overline{\mathcal{O}}}$ the defining ideal of $\overline{\mathcal{O}}$ in $\mathfrak{g}_{\mathcal{O}}$. Thus,

$$\overline{\mathcal{O}} = \text{Spec } \mathbb{C}[\mathfrak{g}_{\mathcal{O}}]/\mathcal{I}_{\overline{\mathcal{O}}}.$$

Recall that $\overline{\mathcal{O}}$ is conical, so $\mathcal{I}_{\overline{\mathcal{O}}}$ is a homogeneous ideal.

Lemma 3.2. *Let \mathcal{O} be a nonzero nilpotent orbit of \mathfrak{g} . If f_1, \dots, f_s are homogeneous generators of $\mathcal{I}_{\overline{\mathcal{O}}}$, then the minimum degree of the f_i is exactly 2.*

Proof. By the above discussion, \mathcal{O} is a product of nilpotent orbits. We may therefore assume that $\mathfrak{g} = \mathfrak{g}_{\mathcal{O}}$ is simple.

Assume that for some $i \in \{1, \dots, s\}$, $\deg f_i = 1$. A contradiction is expected. Let \mathcal{V} be the intersection of all the hyperplanes \mathcal{H}_g , $g \in G$, defined by the linear form

$$g \cdot f_i : \mathfrak{g} \rightarrow \mathbb{C}, \quad x \mapsto f_i(g^{-1}(x)).$$

Since $\bar{\mathcal{O}}$ is G -invariant and is contained in the zero locus of f_i , $\bar{\mathcal{O}}$ is contained in \mathcal{V} . Thus \mathcal{V} is a nonzero G -invariant subspace of \mathfrak{g} which is different from \mathfrak{g} (because \mathcal{V} is contained in the hyperplane $\mathcal{H}_{1_{\mathcal{O}}}$), whence the contradiction since \mathfrak{g} is simple.

The Casimir element, $x \mapsto \langle x, x \rangle$ with $\langle \cdot, \cdot \rangle$ the Killing form of \mathfrak{g} , vanishes on the nilpotent cone of \mathfrak{g} . Hence it is contained in $\mathcal{I}_{\bar{\mathcal{O}}}$. Since it has degree 2, the minimal degree of the f_i is exactly 2. \square

To determine the reducibility of $\mathcal{J}_m(\bar{\mathcal{O}})$ for \mathcal{O} a (nonzero) nilpotent orbit of \mathfrak{g} , we introduce the two sufficient conditions below.

Definition 3.3. Let \mathcal{O} be a nilpotent orbit of \mathfrak{g} .

- (1) We say that \mathcal{O} verifies RC_1 if $\pi_{\bar{\mathcal{O}},1}^{-1}(0)$ is not contained in the closure of $\pi_{\bar{\mathcal{O}},1}^{-1}(\mathcal{O})$.
- (2) Let $m \in \mathbb{N}^*$. We say that \mathcal{O} verifies $\text{RC}_2(m)$ if for some nilpotent orbit \mathcal{O}' contained in $\bar{\mathcal{O}} \setminus \mathcal{O}$, we have $\dim \pi_{\bar{\mathcal{O}},m}^{-1}(\mathcal{O}') \geq \dim \pi_{\bar{\mathcal{O}},m}^{-1}(\mathcal{O}) = (m + 1) \dim \mathcal{O}$.

The following Lemma directly results from [Lemma 2.7\(2\)](#).

Lemma 3.4. Let \mathcal{O} be a nilpotent orbit of \mathfrak{g} .

- (1) If \mathcal{O} verifies RC_1 , then $\mathcal{J}_1(\bar{\mathcal{O}})$ is reducible.
- (2) If \mathcal{O} verifies $\text{RC}_2(m)$ for some $m \in \mathbb{N}^*$, then $\mathcal{J}_m(\bar{\mathcal{O}})$ is reducible.

The zero nilpotent orbit verifies neither RC_1 nor $\text{RC}_2(m)$ for $m \in \mathbb{N}^*$. Since $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ is irreducible for every $m \in \mathbb{N}^*$ (see [Section 1](#)), the same goes for the regular nilpotent orbit according to [Lemma 3.4](#).

In view of the conditions above, let us study the zero fiber of $\pi_{\bar{\mathcal{O}},1} : \mathcal{J}_1(\bar{\mathcal{O}}) \rightarrow \bar{\mathcal{O}}$. As in [Section 2](#), we identify $(\mathfrak{g}_{\mathcal{O}})_m$ with $(\mathfrak{g}_{\mathcal{O}})^{m+1} = \underbrace{\mathfrak{g}_{\mathcal{O}} \times \cdots \times \mathfrak{g}_{\mathcal{O}}}_{(m+1) \text{ factors}}$.

Lemma 3.5. Let \mathcal{O} be a nonzero nilpotent orbit of \mathfrak{g} , and $m \in \mathbb{N}^*$.

- (1) We have $\pi_{\bar{\mathcal{O}},1}^{-1}(0) \simeq \{0\} \times \mathfrak{g}_{\mathcal{O}}$. In particular, $\dim \pi_{\bar{\mathcal{O}},1}^{-1}(0) = \dim \mathfrak{g}_{\mathcal{O}}$.
- (2) If $m \geq 2$, then $\dim \pi_{\bar{\mathcal{O}},m}^{-1}(0) \geq \dim \mathcal{J}_{m-2}(\bar{\mathcal{O}}) + \dim \mathfrak{g}_{\mathcal{O}} \geq m \dim \mathcal{O} + \text{codim}_{\mathfrak{g}_{\mathcal{O}}}(\mathcal{O})$.

Part (2) of [Lemma 3.5](#) remains valid for an affine variety in \mathbb{C}^n defined by homogeneous polynomials of degree at least 2. The special case where all the generators have the same degree is treated in [\[Yuen 2007, Proposition 5.2\]](#).

Proof. Clearly we may assume that $\mathfrak{g}_{\mathcal{O}} = \mathfrak{g}$. Let f_1, \dots, f_r be homogeneous generators of $\mathcal{I}_{\bar{\mathcal{O}}}$ that we order so that $2 = d_1 \leq \dots \leq d_r$, with $d_i = \deg f_i$ for any $i = 1, \dots, r$ (see [Lemma 3.2](#)).

- (1) Through our identification, we can write

$$\pi_{\bar{\mathcal{O}},1}^{-1}(0) \simeq \{0\} \times \{x \in \mathfrak{g} \mid f_i(tx) = 0 \text{ mod } t^2 \text{ for any } i = 1, \dots, r\},$$

whence the statement since for any $x \in \mathfrak{g}$ and $i \in \{1, \dots, r\}$, we have $f_i(tx) = t^{d_i} f_i(x)$ and $d_i \geq 2$.

(2) Assume that $m \geq 2$. Let $(x_1, x_2, \dots, x_{m-1})$ be an element of $\mathcal{J}_{m-2}(\bar{\mathcal{O}})$, and let $x_m \in \mathfrak{g}$. Then for any $i \in \{1, \dots, r\}$, we get

$$f_i(tx_1 + t^2x_2 + \dots + t^m x_m) = f_i(tx_1 + t^2x_2 + \dots + t^{m-1}x_{m-1}) \pmod{t^{m+1}}$$

since f_i is homogeneous of degree at least 2. Hence,

$$f_i(tx_1 + t^2x_2 + \dots + t^m x_m) = t^{d_i} f_i(x_1 + tx_2 + \dots + t^{m-2}x_{m-1}) \pmod{t^{m+1}}.$$

But $f_i(x_1 + tx_2 + \dots + t^{m-2}x_{m-1}) = 0 \pmod{t^{m-1}}$ because $(x_1, x_2, \dots, x_{m-1}) \in \mathcal{J}_{m-2}(\bar{\mathcal{O}})$. So,

$$t^{d_i} f_i(x_1 + tx_2 + \dots + t^{m-2}x_{m-1}) = 0 \pmod{t^{m+1}}$$

since $d_i \geq 2$. In other words, $(0, x_1, x_2, \dots, x_m)$ is an element of $\pi_{\bar{\mathcal{O}}, m}^{-1}(0)$.

Thus we obtain an embedding from $\mathcal{J}_{m-2}(\bar{\mathcal{O}}) \times \mathfrak{g}$ into $\pi_{\bar{\mathcal{O}}, m}^{-1}(0)$ given by

$$\mathcal{J}_{m-2}(\bar{\mathcal{O}}) \times \mathfrak{g} \rightarrow \pi_{\bar{\mathcal{O}}, m}^{-1}(0), \quad ((x_1, x_2, \dots, x_{m-1}), x_m) \mapsto (0, x_1, x_2, \dots, x_{m-1}, x_m).$$

The assertions follows. \square

Let \mathcal{O} be a nonzero nilpotent orbit of \mathfrak{g} , and fix $e \in \mathcal{O}$. The tangent space at e to $\bar{\mathcal{O}}$ is the space $[e, \mathfrak{g}]$. Consider the morphism

$$\eta_{\mathfrak{g}, e} : G \times [e, \mathfrak{g}] \rightarrow \mathfrak{g}, \quad (g, x) \mapsto g(x).$$

Proposition 3.6. *The nonzero nilpotent orbit \mathcal{O} verifies RC_1 if and only if the closure of the image of $\eta_{\mathfrak{g}, e}$ is strictly contained in $\mathfrak{g}_{\mathcal{O}}$.*

Proof. Since $[e, \mathfrak{g}] = [e, \mathfrak{g}_{\mathcal{O}}]$, we may assume that $\mathfrak{g} = \mathfrak{g}_{\mathcal{O}}$. Thus, by the definition of condition RC_1 , we have to show that $\pi_{\bar{\mathcal{O}}, 1}^{-1}(0)$ is contained in

$$\overline{\pi_{\bar{\mathcal{O}}, 1}^{-1}(\mathcal{O})}$$

if and only if $\eta_{\mathfrak{g}, e}$ is dominant, i.e., $\overline{G \cdot [e, \mathfrak{g}]} = \mathfrak{g}$.

By Lemma 3.5(1), we have $\pi_{\bar{\mathcal{O}}, 1}^{-1}(0) \simeq \{0\} \times \mathfrak{g}$. On the other hand,

$$\pi_{\bar{\mathcal{O}}, 1}^{-1}(\mathcal{O}) = G \cdot (\{e\} \times [e, \mathfrak{g}]).$$

So, if $\pi_{\bar{\mathcal{O}}, 1}^{-1}(0) \subset \overline{\pi_{\bar{\mathcal{O}}, 1}^{-1}(\mathcal{O})}$, then

$$\{0\} \times \mathfrak{g} \subset \overline{G \cdot (\{e\} \times [e, \mathfrak{g}])} \subset \overline{G \cdot e} \times \overline{G \cdot [e, \mathfrak{g}]},$$

whence the inclusion $\mathfrak{g} \subset \overline{G \cdot [e, \mathfrak{g}]}$, and $\eta_{\mathfrak{g}, e}$ is dominant.

For the other direction, observe that

$$\overline{\pi_{\bar{\mathcal{O}}, 1}^{-1}(\mathcal{O})}$$

is a closed bicone of $\mathfrak{g} \times \mathfrak{g}$ since \mathcal{O} and $\overline{\mathcal{O}}$ are both subcones of \mathfrak{g} . Here, by *bicone*, we mean a subset of $\mathfrak{g} \times \mathfrak{g}$ stable under the natural $(\mathbb{C}^* \times \mathbb{C}^*)$ -action on $\mathfrak{g} \times \mathfrak{g}$. Therefore, if $\overline{G \cdot [e, \mathfrak{g}]} = \mathfrak{g}$, then

$$\overline{G \cdot (\{e\} \times [e, \mathfrak{g}])} = \overline{G \cdot (\mathbb{C}^* e \times [e, \mathfrak{g}])} \supset \{0\} \times \overline{G \cdot [e, \mathfrak{g}]} = \{0\} \times \mathfrak{g},$$

whence $\pi_{\overline{\mathcal{O}}, 1}^{-1}(0) \subset \overline{\pi_{\overline{\mathcal{O}}, 1}^{-1}(\mathcal{O})}$. □

Example 3.7. Let $p \in \mathbb{N}^*$ with $p \geq 2$, and $\mathfrak{g} = \mathfrak{sl}_{2p}(\mathbb{C})$. In the notation of [Appendix A](#), we claim that the nilpotent orbit $\mathcal{O}_{(2^p)}$ of \mathfrak{g} associated with the partition (2^p) verifies RC_1 . According to [Proposition 3.6](#), it suffices to prove that for the element

$$e := \begin{pmatrix} 0 & I_p \\ 0 & 0 \end{pmatrix} \in \mathcal{O}_{(2^p)},$$

the morphism $\eta_{\mathfrak{g}, e}$ is not dominant. We readily verify that $[e, \mathfrak{g}]$ consists of matrices of the form

$$\begin{pmatrix} A & C \\ 0 & -A \end{pmatrix}$$

with A and C of size p . In particular, $[e, \mathfrak{g}]$ is contained in the closed subset \mathcal{Z} of \mathfrak{g} consisting of the matrices whose characteristic polynomial is even. Since $\overline{G([e, \mathfrak{g}])}$ and \mathcal{Z} are both closed G -stable subsets of \mathfrak{g} , we get

$$\overline{G([e, \mathfrak{g}])} \subset \mathcal{Z}.$$

The diagonal matrix $\text{diag}(1, \dots, 1, -2p + 1)$ is in \mathfrak{g} but does not lie in \mathcal{Z} for $p \geq 2$. Hence, \mathcal{Z} is strictly contained in \mathfrak{g} , and $\eta_{\mathfrak{g}, e}$ is not dominant. Thus $\mathcal{O}_{(2^p)}$ verifies RC_1 .

According to [Lemma 3.4\(1\)](#), $\mathcal{J}_1(\overline{\mathcal{O}_{(2^p)}})$ is reducible. In fact, we can be more precise. By [\[Weyman 2002, Theorem 1\]](#) (see also [\[Weyman 1989\]](#) or [\[Weyman 2003, Proposition 8.2.15\]](#)), the defining ideal of $\overline{\mathcal{O}_{(2^p)}}$ is generated by the entries of the matrix X^2 as functions of $X \in \mathfrak{sl}_{2p}(\mathbb{C})$. It follows that $\mathcal{J}_1(\overline{\mathcal{O}_{(2^p)}})$ can be identified with the scheme of pairs $(X_0, X_1) \in \mathfrak{sl}_{2p}(\mathbb{C}) \times \mathfrak{sl}_{2p}(\mathbb{C})$ defined by the equations $X_0^2 = 0$ and $X_0 X_1 + X_1 X_0 = 0$. Using this identification, we obtain from direct computations that

- $\mathcal{J}_1(\overline{\mathcal{O}_{(2^p)}})$ has exactly one irreducible component of dimension $4p^2 = 2 \dim \mathcal{O}_{(2^p)}$,
- all the other irreducible components have dimension $4p^2 - 1$, and $\pi_{\overline{\mathcal{O}_{(2^p)}}, 1}^{-1}(0)$ is one of them.

Remark 3.8. Assume that $\mathfrak{g} = \mathfrak{g}_{\mathcal{O}}$. A nilpotent element e is distinguished if its centralizer is contained in the nilpotent cone. In particular, if e is distinguished, then the centralizer of an \mathfrak{sl}_2 -triple (e, h, f) in \mathfrak{g} is zero, and the theory of representations of \mathfrak{sl}_2 shows that $[e, \mathfrak{g}]$ contains \mathfrak{g}^h , and hence contains a Cartan subalgebra of \mathfrak{g} . Consequently, $G \cdot e$ does not verify RC_1 .

Remark 3.9. Assume that $\mathfrak{g} = \mathfrak{g}_{\mathcal{O}}$. Since $G \times [e, \mathfrak{g}]$ and \mathfrak{g} are irreducible varieties, $\eta_{\mathfrak{g},e}$ is dominant if and only if there is a nonempty open set U consisting of points $a \in G \times [e, \mathfrak{g}]$ such that $(d\eta_{\mathfrak{g},e})_a$ is surjective. The differential of $\eta_{\mathfrak{g},e}$ at $a = (g, [e, x])$, with $(g, x) \in G \times \mathfrak{g}$ is given by

$$\mathfrak{g} \times [e, \mathfrak{g}] \rightarrow \mathfrak{g}, \quad (v, [e, w]) \mapsto [v, [e, x]] + g([e, w]).$$

Let us endow $G \times [e, \mathfrak{g}]$ with the action of G by left multiplication on the first factor. Since $\eta_{\mathfrak{g},e}$ is G -equivariant, we may assume that a is of the form $a = (1_G, [e, x])$ with $x \in \mathfrak{g}$. Then $(d\eta_{\mathfrak{g},e})_a$ is surjective if and only if $[\mathfrak{g}, [e, x]] + [e, \mathfrak{g}] = \mathfrak{g}$.

Consequently, $\eta_{\mathfrak{g},e}$ is dominant if and only if there exists $x \in \mathfrak{g}$ such that $[\mathfrak{g}, [e, x]] + [e, \mathfrak{g}] = \mathfrak{g}$. This allows us to affirm in some cases that $\eta_{\mathfrak{g},e}$ is dominant. For example, for e in the nondistinguished nilpotent orbit $\mathcal{O}_{(3^2)}$ of $\mathfrak{sl}_6(\mathbb{C})$, the map $\eta_{\mathfrak{g},e}$ is dominant.

4. Little nilpotent orbits

We introduce in this section a family of nonzero nilpotent orbits which verify both RC_1 and $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$. This family turns out to be useful to study the reducibility of jet schemes of many other orbits.

[Lemma 3.5](#) leads us to the following definition.

Definition 4.1. Let \mathcal{O} be a nilpotent orbit of \mathfrak{g} and let $\mathfrak{g}_{\mathcal{O}}$ be as in [Definition 3.1](#). We say that \mathcal{O} is *little* if $0 < 2 \dim \mathcal{O} \leq \dim \mathfrak{g}_{\mathcal{O}}$.

In particular, neither the zero orbit nor the regular nilpotent orbit is little.

Proposition 4.2. *If \mathcal{O} is a little nilpotent orbit of \mathfrak{g} , then \mathcal{O} verifies RC_1 and $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*

Proof. Let \mathcal{O} be a little nilpotent orbit of \mathfrak{g} . As in the preceding proofs, we may assume that $\mathfrak{g} = \mathfrak{g}_{\mathcal{O}}$. According to [Lemma 3.5\(1\)](#), $\dim \pi_{\overline{\mathcal{O}},1}^{-1}(0) = \dim \mathfrak{g}$, and since $\pi_{\overline{\mathcal{O}},1}^{-1}(\mathcal{O})$ has dimension $2 \dim \mathcal{O} \leq \dim \mathfrak{g}$, it follows that \mathcal{O} verifies $\text{RC}_2(1)$ and RC_1 . Now let $m \geq 2$. According to [Lemma 3.5\(2\)](#), we have

$$\dim \pi_{\overline{\mathcal{O}},m}^{-1}(0) \geq m \dim \mathcal{O} + \text{codim}_{\mathfrak{g}}(\mathcal{O}) \geq (m+1) \dim \mathcal{O},$$

since $\text{codim}_{\mathfrak{g}}(\mathcal{O}) \geq \dim \mathcal{O}$ because \mathcal{O} is little. Hence \mathcal{O} verifies $\text{RC}_2(m)$. \square

When \mathfrak{g} is simple, there is a unique nonzero nilpotent orbit \mathcal{O}_{\min} , called the *minimal* nilpotent orbit of \mathfrak{g} , of minimal dimension and it is contained in the closure of all nonzero nilpotent orbits.

Corollary 4.3. *Assume that \mathfrak{g} is simple and not of type A_1 . Then \mathcal{O}_{\min} is little. In particular, $\mathcal{I}_m(\overline{\mathcal{O}_{\min}})$ is reducible for every $m \in \mathbb{N}^*$.*

Proof. Let $e \in \mathcal{O}_{\min}$ that we embed into an \mathfrak{sl}_2 -triple (e, h, f) of \mathfrak{g} , and consider the corresponding Dynkin grading,

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}(i) \quad \text{with } \mathfrak{g}(i) := \{x \in \mathfrak{g} \mid [h, x] = ix\}.$$

By [Collingwood and McGovern 1993, Lemma 4.1.3],

$$\dim \mathcal{O} = \dim \mathfrak{g} - \dim \mathfrak{g}(0) - \dim \mathfrak{g}(1).$$

In addition, since $e \in \mathcal{O}_{\min}$, we have $\dim \mathfrak{g}(2) = 1$ and $\mathfrak{g} = \sum_{-2 \leq i \leq 2} \mathfrak{g}(i)$ [Tauvel and Yu 2005, Proposition 34.4.1]. As a result,

$$\dim \mathfrak{g} - 2 \dim \mathcal{O} = \dim \mathfrak{g}(0) - 2.$$

The Levi subalgebra $\mathfrak{g}(0)$ contains a Cartan subalgebra which has dimension at least two by our hypothesis. Hence, $\dim \mathfrak{g} - 2 \dim \mathcal{O} \geq 0$, and so \mathcal{O}_{\min} is little. \square

For classical simple Lie algebras, there are explicit formulas (see Appendix A) for the dimension of nilpotent orbits. This allows us to readily obtain examples of little nilpotent orbits.

Example 4.4. Let $n \in \mathbb{N}^*$ and $p, q \in \mathbb{N}$.

- (i) A nilpotent orbit of $\mathfrak{sl}_n(\mathbb{C})$ corresponding to a rectangular partition is never little.
- (ii) The nilpotent orbit $\mathcal{O}_{(2^p, 1^q)}$ of $\mathfrak{sl}_{2p+q}(\mathbb{C})$ is little if and only if $p, q \in \mathbb{N}^*$.
- (iii) The nilpotent orbit $\mathcal{O}_{(p, 1^q)}$ of $\mathfrak{sl}_{p+q}(\mathbb{C})$ is little for $q \gg p$.

Explicit computations suggest that it is unlikely that there is a nice description of little nilpotent orbits in terms of partitions.

For the notation $\mathcal{P}_\varepsilon(n)$, $\varepsilon \in \{0, 1\}$, and \mathcal{O}_λ with $\lambda \in \mathcal{P}_\varepsilon(n)$, $n \in \mathbb{N}^*$, refer to Appendix A.

Example 4.5. Let $\lambda = (2^p, 1^q)$, with $p \in \mathbb{N}^*$ and $q \in \mathbb{N}$.

- (i) If p is even, then $\lambda \in \mathcal{P}_1(n)$, and the nilpotent orbit \mathcal{O}_λ of $\mathfrak{so}_{2p+q}(\mathbb{C})$ is little.
- (ii) If q is even, then $\lambda \in \mathcal{P}_{-1}(n)$, and the nilpotent orbit \mathcal{O}_λ of $\mathfrak{sp}_{2p+q}(\mathbb{C})$ is little if and only if $p \leq q(q+1)/2$.

The next proposition will allow us to produce new examples of nilpotent orbits which verify RC_1 by the “restriction” of certain little nilpotent orbits to Levi subalgebras.

Recall that for \mathcal{O} a nilpotent orbit of some reductive Lie algebra \mathfrak{a} , the semisimple Lie algebra $\mathfrak{a}_{\mathcal{O}}$ was defined in Definition 3.1.

Proposition 4.6. *Assume that \mathfrak{g} is simple. Let \mathfrak{l} be a Levi subalgebra of \mathfrak{g} with center $\mathfrak{z}(\mathfrak{l})$, and denote by A the connected subgroup of G whose Lie algebra is $\mathfrak{a} := [\mathfrak{l}, \mathfrak{l}]$. Let e be a nilpotent element of \mathfrak{a} and suppose that the following conditions are satisfied:*

- (i) \mathfrak{a} contains a regular semisimple element of \mathfrak{g} ,
- (ii) $\mathfrak{a}_{A \cdot e} = \mathfrak{a}$,
- (iii) $2 \dim G \cdot e \leq \dim \mathfrak{g} - \dim \mathfrak{z}(\mathfrak{l})$.

Then $A \cdot e$ verifies RC_1 .

Proof. Define the following maps

$$\theta : G \times \mathfrak{a} \rightarrow \mathfrak{g}, (g, x) \mapsto g(x), \quad \eta = \eta_{\mathfrak{g}, e} : G \times [e, \mathfrak{g}] \rightarrow \mathfrak{g}, (g, x) \mapsto g(x).$$

Observe that the image of each of these maps is irreducible. Moreover, for any $x \in \mathfrak{g}$, the map $g \mapsto (g^{-1}, g(x))$ defines a bijection between $G_\theta(x) := \{g \in G \mid g(x) \in \mathfrak{a}\}$ and $\theta^{-1}(\{x\})$. Similarly, we have a bijection between $G_\eta(x) := \{g \in G \mid g(x) \in [e, \mathfrak{g}]\}$ and $\eta^{-1}(\{x\})$. These bijections are isomorphisms of varieties.

Step 1. We shall first compute the dimension of the image of θ .

Let L be the connected subgroup of G whose Lie algebra is \mathfrak{l} . By condition (i), \mathfrak{a} contains regular semisimple elements of \mathfrak{g} . If s is such an element, then \mathfrak{g}^s is a Cartan subalgebra of \mathfrak{l} . Let $g \in G_\theta(s)$. Then $g(s) \in \mathfrak{a}$ and $\mathfrak{g}^{g(s)} = \mathfrak{g}(\mathfrak{g}^s)$ is another Cartan subalgebra of \mathfrak{l} . It follows that there exists $\tau \in L$ such that $\tau g \in N_G(\mathfrak{g}^s)$, with $N_G(\mathfrak{g}^s)$ the normalizer of \mathfrak{g}^s in G . Hence, $g \in LN_G(\mathfrak{g}^s)$. Thus, we have obtained the inclusion $G_\theta(s) \subset LN_G(\mathfrak{g}^s)$. On the other hand, since L normalizes \mathfrak{a} , we get $L \subset G_\theta(s)$ and therefore $\dim L \leq \dim G_\theta(s)$.

Let $C_G(\mathfrak{g}^s)$ and $C_L(\mathfrak{g}^s)$ be the centralizers of \mathfrak{g}^s in G and L , respectively. Since \mathfrak{g}^s is a Cartan subalgebra, $C_G(\mathfrak{g}^s)$ is connected, so $C_G(\mathfrak{g}^s) = C_L(\mathfrak{g}^s)$ is contained in L . It follows that $LN_G(\mathfrak{g}^s)$ is a finite union of right L -cosets. We deduce that

$$\dim \theta^{-1}(\{s\}) = \dim G_\theta(s) = \dim L = \dim \mathfrak{a} + \mathfrak{z}(\mathfrak{l}).$$

Since the set of regular semisimple elements in \mathfrak{g} is open and dense, we obtain that for s as above,

$$\dim \overline{\text{im } \theta} = \dim \mathfrak{g} + \dim \mathfrak{a} - \dim \theta^{-1}(\{s\}) = \dim \mathfrak{g} - \dim \mathfrak{z}(\mathfrak{l}).$$

Step 2. We now consider the image of η .

Let (e, h, f) be an \mathfrak{sl}_2 -triple of \mathfrak{g} . We easily check that $\mathfrak{c} := \mathbb{C}h \oplus \mathfrak{g}^e$ is a Lie subalgebra, and that \mathfrak{c} stabilizes $[e, \mathfrak{g}]$. Let C be the connected subgroup of G whose Lie algebra is \mathfrak{c} . Then C is contained in $G_\eta(x)$ for any $x \in [e, \mathfrak{g}]$. In particular, $\dim G_\eta(x) \geq \dim C = 1 + \dim \mathfrak{g}^e$ for $x \in [e, \mathfrak{g}]$, and so

$$\dim \overline{\text{im } \eta} \leq \dim \mathfrak{g} + \dim [e, \mathfrak{g}] - 1 - \dim \mathfrak{g}^e = 2 \dim G \cdot e - 1.$$

Step 3. By condition (iii) and Steps 1 and 2, we deduce that $\dim \overline{\text{im } \theta} > \dim \overline{\text{im } \eta}$. Thus $\overline{\text{im } \theta} \not\subset \overline{\text{im } \eta}$. We claim that this implies that $A \cdot e$ is RC_1 . Let us suppose on the contrary that $A \cdot e$ is not RC_1 . By condition (ii) and [Lemma 3.5\(1\)](#), $\pi_{A \cdot e, 1}^{-1}(0) = \{0\} \times \mathfrak{a}$. So, $\pi_{A \cdot e, 1}^{-1}(0)$ is contained in

$$\overline{\pi_{A \cdot e, 1}^{-1}(A \cdot e)}.$$

Recall from the end of [Section 2](#) the notation G_1 and A_1 for $\mathcal{J}_1(G)$ and $\mathcal{J}_1(A)$, respectively. It follows that

$$\{0\} \times G \cdot \mathfrak{a} \subset G_1 \cdot (\{0\} \times \mathfrak{a}) \subset G_1 \overline{A_1 \cdot e} \subset \overline{G_1 \cdot e},$$

whence

$$\{0\} \times \overline{G \cdot \mathfrak{a}} \subset \overline{G_1 \cdot e}.$$

Since $\overline{\pi_{G \cdot e, 1}^{-1}(G \cdot e)} = \overline{G_1 \cdot e}$ (see [Lemma 2.10](#)), it follows from the proof of [Proposition 3.6](#) that

$$\overline{G_1 \cdot e} \cap (\{0\} \times \mathfrak{g}) = \overline{\pi_{G \cdot e, 1}^{-1}(G \cdot e)} \cap (\{0\} \times \mathfrak{g}) = \{0\} \times \overline{G \cdot [e, \mathfrak{g}]}.$$

Hence we get $\overline{\text{im } \theta} \subset \overline{\text{im } \eta}$ and the contradiction. \square

Suppose that \mathfrak{g} is simple. Let us fix a Cartan subalgebra \mathfrak{h} of \mathfrak{g} . Denote by Δ the root system relative to $(\mathfrak{g}, \mathfrak{h})$ and let us fix a system of simple roots Π . Given $S \subset \Pi$, we denote $\Delta_S = \mathbb{Z}S \cap \Delta$ the subroot system generated by S , and

$$\mathfrak{l}_S = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta_S} \mathfrak{g}_\alpha$$

where \mathfrak{g}_α denotes the root subspace relative to α . Then \mathfrak{l}_S is a Levi subalgebra of \mathfrak{g} and any Levi subalgebra of \mathfrak{g} is conjugate to one in this form.

Given $S \subset \Pi$, set $\mathfrak{t} = [\mathfrak{l}_S, \mathfrak{l}_S] \cap \mathfrak{h}$. Then, \mathfrak{l}_S verifies condition (i) if and only if $\mathfrak{t} \not\subset \bigcup_{\alpha \in \Delta} \ker \alpha$. To check the latter condition, it is enough to verify that for every $\alpha \in \Delta$, there is $\beta \in S$ such that $\langle \beta^\vee, \alpha \rangle \neq 0$.

Thus not all Levi subalgebras of \mathfrak{g} verify condition (i) of [Proposition 4.6](#). For example, if \mathfrak{g} is simple of type B_ℓ , then a (maximal) Levi subalgebra whose semisimple part is simple of type $B_{\ell-1}$ does not verify the condition. The same goes for a Levi subalgebra in type C_ℓ whose semisimple part is simple of type $C_{\ell-1}$.

However, if \mathfrak{g} is simple of type D_ℓ and if \mathfrak{l} is a Levi subalgebra whose semisimple part is simple of type $D_{\ell-1}$, then \mathfrak{l} verifies the condition (i). Likewise, if \mathfrak{g} is simple of type E_7 and if \mathfrak{l} is a Levi subalgebra whose semisimple part is simple of type E_6 , then \mathfrak{l} verifies the condition (i). Applying [Proposition 4.6](#), we obtain examples of nilpotent orbits in types D or E_6 which verify RC_1 that are not little.

We list in [Table 1](#) some nilpotent orbits that we obtain in this way. In all the examples presented in the table, the center of the Levi subalgebra is 1-dimensional, and \mathfrak{a} is simple. The first and second columns give the type of the simple Lie

\mathfrak{g}	\mathfrak{a}	$G \cdot e$	$A \cdot e$
D_6	D_5	$(3, 2^2, 1^5)$	$(3, 2^2, 1^3)$
D_7	D_6	$(3^2, 1^8)$	$(3^2, 1^6)$
D_9	D_8	$(3^2, 2^2, 1^8)$	$(3^2, 2^2, 1^6)$
D_{10}	D_9	$(3^3, 1^{11})$	$(3^3, 1^9)$
D_{10}	D_9	$(4^2, 1^{12})$	$(4^2, 1^{10})$
D_{10}	D_9	$(5, 2^2, 1^{11})$	$(5, 2^2, 1^9)$
D_{10}	D_9	$(5, 3, 1^{12})$	$(5, 3, 1^{10})$
E_7	E_6	$(3A_1)'$	$3A_1$
E_7	E_6	A_2	A_2

Table 1. Examples of nonlittle nilpotent orbits satisfying RC_1 obtained by restriction.

algebras \mathfrak{g} and \mathfrak{a} . Condition (ii) is verified in view of the discussion above. We describe the nilpotent orbits $G \cdot e$ and $A \cdot e$ in the third and fourth columns, respectively. The description for an orbit in \mathfrak{g} of type D is given in terms of partitions (see [Appendix A](#)), while for an orbit in \mathfrak{g} of type E_6 or E_7 , it is given by its Bala–Carter label.

Remark 4.7. (1) The first and last lines of [Table 1](#) provide examples of a *rigid*² nilpotent orbit which verifies RC_1 and which is not little.

(2) Propositions [3.6](#), [4.2](#), and [4.6](#), together with [Remark 3.9](#), allow us to classify all nilpotent orbits verifying RC_1 in simple Lie algebras of exceptional type. They are listed in [Appendix C](#).

5. Induced nilpotent orbits

Let \mathfrak{l} be a proper Levi subalgebra of \mathfrak{g} , and let \mathfrak{p} be a parabolic subalgebra of \mathfrak{g} with Levi decomposition $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$ so that \mathfrak{u} is the nilpotent radical of \mathfrak{p} . Let $P, L,$ and U be the connected closed subgroups of G whose Lie algebra are $\mathfrak{p}, \mathfrak{l},$ and $\mathfrak{u},$ respectively. Then $P = LU$.

The following definitions and results on induced nilpotent orbits are mostly extracted from [[Richardson 1974](#); [Lusztig and Spaltenstein 1979](#)]. We refer to [[Collingwood and McGovern 1993](#), Chapter 7] for a recent survey.

Theorem 5.1. *Let $\mathcal{O}_\mathfrak{l}$ be a nilpotent orbit of \mathfrak{l} . There exists a unique nilpotent orbit $\mathcal{O}_\mathfrak{g}$ in \mathfrak{g} whose intersection with $\mathcal{O}_\mathfrak{l} + \mathfrak{u}$ is a dense open subset of $\mathcal{O}_\mathfrak{l} + \mathfrak{u}$. Moreover, the intersection of $\mathcal{O}_\mathfrak{g}$ with $\mathcal{O}_\mathfrak{l} + \mathfrak{u}$ consists of a single P -orbit and $\text{codim}_\mathfrak{g}(\mathcal{O}_\mathfrak{g}) = \text{codim}_\mathfrak{l}(\mathcal{O}_\mathfrak{l})$.*

²See [Section 5](#) for the notion of rigid nilpotent orbit, and [Appendices A](#) and [C](#) for the description of rigid nilpotent orbits in simple Lie algebras.

The nilpotent orbit $\mathcal{O}_{\mathfrak{g}}$ only depends on \mathfrak{l} , and not on the choice of a parabolic subalgebra \mathfrak{p} containing it. The nilpotent orbit $\mathcal{O}_{\mathfrak{g}}$ is called the *induced nilpotent orbit of \mathfrak{g} from $\mathcal{O}_{\mathfrak{l}}$* , and it is denoted by $\text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}_{\mathfrak{l}})$. A nilpotent orbit which is not induced in a proper way from another one is called *rigid*. In type A, only the zero orbit is rigid.

Remark 5.2. (1) Let $\mathfrak{s}_1, \dots, \mathfrak{s}_n$ be the simple factors of $[\mathfrak{g}, \mathfrak{g}]$ and denote by $\mathfrak{z}(\mathfrak{g})$ the center of \mathfrak{g} . Then there are Levi subalgebras $\mathfrak{r}_1, \dots, \mathfrak{r}_n$ of $\mathfrak{s}_1, \dots, \mathfrak{s}_n$, respectively, such that

$$\mathfrak{l} = \mathfrak{z}(\mathfrak{g}) \times \mathfrak{r}_1 \times \dots \times \mathfrak{r}_n.$$

If $\mathcal{O}_{\mathfrak{l}}$ is a nilpotent orbit of \mathfrak{l} , then $\mathcal{O}_{\mathfrak{l}} = \mathcal{O}_{\mathfrak{r}_1} \times \dots \times \mathcal{O}_{\mathfrak{r}_n}$, where $\mathcal{O}_{\mathfrak{r}_1}, \dots, \mathcal{O}_{\mathfrak{r}_n}$ are nilpotent orbits in the semisimple parts of $\mathfrak{r}_1, \dots, \mathfrak{r}_n$, respectively. Then

$$\text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}_{\mathfrak{l}}) = \text{Ind}_{\mathfrak{r}_1}^{\mathfrak{s}_1}(\mathcal{O}_{\mathfrak{r}_1}) \times \dots \times \text{Ind}_{\mathfrak{r}_n}^{\mathfrak{s}_n}(\mathcal{O}_{\mathfrak{r}_n}) = \text{Ind}_{[\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{l}}^{[\mathfrak{g}, \mathfrak{g}]}(\mathcal{O}_{\mathfrak{l}}).$$

(2) The induction property is transitive in the following sense [Collingwood and McGovern 1993, Proposition 7.1.4]: if \mathfrak{l}_1 and \mathfrak{l}_2 are two Levi subalgebras of \mathfrak{g} with $\mathfrak{l}_1 \subset \mathfrak{l}_2$, then

$$\text{Ind}_{\mathfrak{l}_2}^{\mathfrak{g}}(\text{Ind}_{\mathfrak{l}_1}^{\mathfrak{l}_2}(\mathcal{O}_{\mathfrak{l}_1})) = \text{Ind}_{\mathfrak{l}_1}^{\mathfrak{g}}(\mathcal{O}_{\mathfrak{l}_1}).$$

(3) If $\Omega_{\mathfrak{l}}$ is an L -orbit in $\overline{\mathcal{O}_{\mathfrak{l}}} \setminus \mathcal{O}_{\mathfrak{l}}$, then the induced nilpotent orbit of \mathfrak{g} from $\Omega_{\mathfrak{l}}$ is contained in $\overline{\mathcal{O}_{\mathfrak{g}}} \setminus \mathcal{O}_{\mathfrak{g}}$.

Let $\mathcal{O}_{\mathfrak{l}}$ be a nilpotent orbit of \mathfrak{l} and denote by $\mathcal{O}_{\mathfrak{g}}$ the induced nilpotent orbit of \mathfrak{g} from $\mathcal{O}_{\mathfrak{l}}$. According to Theorem 5.1, $\mathcal{O}_{\mathfrak{g}} \cap (\mathcal{O}_{\mathfrak{l}} + \mathfrak{u})$ is a single P -orbit that we shall denote by $\mathcal{O}_{\mathfrak{p}}$; that is,

$$\mathcal{O}_{\mathfrak{p}} := \mathcal{O}_{\mathfrak{g}} \cap (\mathcal{O}_{\mathfrak{l}} + \mathfrak{u}).$$

Lemma 5.3. *The orbits satisfy*

$$\overline{\mathcal{O}_{\mathfrak{p}}} = \overline{\mathcal{O}_{\mathfrak{l}}} + \mathfrak{u}, \quad \overline{\mathcal{O}_{\mathfrak{p}}} \cap \mathcal{O}_{\mathfrak{g}} = \mathcal{O}_{\mathfrak{p}}, \quad \overline{\mathcal{O}_{\mathfrak{g}}} = G \cdot (\overline{\mathcal{O}_{\mathfrak{l}}} + \mathfrak{u}).$$

Proof. The first equality is obvious since $\mathcal{O}_{\mathfrak{p}}$ is dense in $\mathcal{O}_{\mathfrak{l}} + \mathfrak{u}$ by definition.

Next, the inclusion $\mathcal{O}_{\mathfrak{p}} \subset \overline{\mathcal{O}_{\mathfrak{p}}} \cap \mathcal{O}_{\mathfrak{g}}$ is clear. To show the other inclusion, assume that there is $x \in \overline{\mathcal{O}_{\mathfrak{p}}} \cap \mathcal{O}_{\mathfrak{g}}$, with $x \notin \mathcal{O}_{\mathfrak{p}}$. A contradiction is expected. Since $x \in \overline{\mathcal{O}_{\mathfrak{p}}} \setminus \mathcal{O}_{\mathfrak{p}}$, $\dim P \cdot x < \dim P \cdot e$. Hence,

$$\dim \mathfrak{g}^x \geq \dim \mathfrak{p}^x > \dim \mathfrak{p}^e = \dim \mathfrak{g}^e.$$

As a consequence, x is not in $\mathcal{O}_{\mathfrak{g}}$, whence the contradiction.

A proof of the last equality can be found in [op. cit., Theorem 7.1.3]. □

We have the following generalization.

Lemma 5.4. *The jet schemes satisfy*

- (1) $\overline{\mathcal{J}_m(\mathcal{O}_p)} = \overline{\mathcal{J}_m(\mathcal{O}_l)} + \mathfrak{u}_m$,
- (2) $\overline{\mathcal{J}_m(\mathcal{O}_p)} \cap \mathcal{J}_m(\mathcal{O}_g) = \mathcal{J}_m(\mathcal{O}_p) = (\mathcal{J}_m(\overline{\mathcal{O}_l}) + \mathfrak{u}_m) \cap \mathcal{J}_m(\mathcal{O}_g)$,
- (3) $\overline{\mathcal{J}_m(\mathcal{O}_g)}$ is the closure of $G_m \cdot \overline{\mathcal{J}_m(\mathcal{O}_p)}$.

Proof. (1) Since $\mathcal{O}_p \subset \mathcal{O}_l + \mathfrak{u}$, we get $\mathcal{J}_m(\mathcal{O}_p) \subset \overline{\mathcal{J}_m(\mathcal{O}_l)} + \mathfrak{u}_m$ because $\overline{\mathcal{J}_m(\mathcal{O}_l)} + \mathfrak{u}_m$ is closed. Let $e' \in \mathcal{O}_l$ and $x \in \mathfrak{u}$ be such that $e := e' + x$ is in \mathcal{O}_p . From the above inclusion, we deduce that

$$\dim \mathfrak{p} - \dim \mathfrak{p}^e \leq \dim \mathfrak{l} - \dim \mathfrak{l}^{e'} + \dim \mathfrak{u} = \dim \mathfrak{p} - \dim \mathfrak{g}^e,$$

because $\dim \mathfrak{l}^{e'} = \dim \mathfrak{g}^e$ by [Theorem 5.1](#). Since $\dim \mathfrak{p}^e \leq \dim \mathfrak{g}^e$, we get $\mathfrak{p}^e = \mathfrak{g}^e$, whence $\dim \overline{\mathcal{J}_m(\mathcal{O}_p)} = \dim(\mathcal{J}_m(\mathcal{O}_l) + \mathfrak{u}_m)$ by [Lemma 2.10](#) and [Proposition 2.6](#). So $\overline{\mathcal{J}_m(\mathcal{O}_p)}$ and $\overline{\mathcal{J}_m(\mathcal{O}_l)} + \mathfrak{u}_m$ are irreducible varieties of the same dimension, and the equality follows.

(2) Taking into account [Lemma 2.10](#) and [Proposition 2.6](#), the result follows from the same arguments as in the proof of the second equality of [Lemma 5.3](#).

(3) By [Lemma 2.10](#),

$$\mathcal{J}_m(\mathcal{O}_g) = G_m \cdot \mathcal{J}_m(\mathcal{O}_p) \subset G_m \cdot \overline{\mathcal{J}_m(\mathcal{O}_p)}.$$

As a result, the jet scheme $\overline{\mathcal{J}_m(\mathcal{O}_g)}$ is in the closure of $G_m \cdot \overline{\mathcal{J}_m(\mathcal{O}_p)}$. On the other hand, since $\overline{\mathcal{J}_m(\mathcal{O}_g)}$ is G_m -stable, we get

$$G_m \cdot \overline{\mathcal{J}_m(\mathcal{O}_p)} \subset \overline{\mathcal{J}_m(\mathcal{O}_g)}.$$

So the closure of $G_m \cdot \overline{\mathcal{J}_m(\mathcal{O}_p)}$ is contained in $\overline{\mathcal{J}_m(\mathcal{O}_g)}$, whence the expected equality. \square

Question 5.5. For $m = 0$, $G_m \cdot \overline{\mathcal{J}_m(\mathcal{O}_p)}$ is closed (see [Lemma 5.3](#)) essentially because G/P is compact. For $m \geq 1$, G_m/P_m is a trivial fibration over G/P with m -dimensional affine fiber. Can we show nevertheless that $G_m \cdot (\overline{\mathcal{J}_m(\mathcal{O}_l)} + \mathfrak{u}_m)$ is closed, in other words that $\overline{\mathcal{J}_m(\mathcal{O}_g)} = G_m \cdot (\overline{\mathcal{J}_m(\mathcal{O}_l)} + \mathfrak{u}_m)$?

Theorem 5.6. *Let \mathfrak{l} be a Levi subalgebra of \mathfrak{g} , \mathcal{O}_l a nilpotent orbit of \mathfrak{l} , and \mathcal{O}_g the induced nilpotent orbit of \mathfrak{g} from \mathcal{O}_l . If \mathcal{O}_l verifies $\text{RC}_2(m)$ for some $m \in \mathbb{N}^*$, then \mathcal{O}_g also verifies $\text{RC}_2(m)$.*

The rest of the section will be devoted to the proof of [Theorem 5.6](#).

Definition 5.7. Let \mathfrak{l} be a Levi subalgebra of \mathfrak{g} . We say that \mathfrak{l} is a *maximal Levi subalgebra* of \mathfrak{g} if the center of $[\mathfrak{g}, \mathfrak{g}] \cap \mathfrak{l}$ has dimension one.

Let us first assume that \mathfrak{g} is simple and that \mathfrak{l} is a maximal Levi subalgebra of \mathfrak{g} . Thus, the center $\mathfrak{z}(\mathfrak{l})$ of \mathfrak{l} has dimension one. Let us fix a Cartan subalgebra \mathfrak{h} in \mathfrak{l} and Δ the root system relative to $(\mathfrak{g}, \mathfrak{h})$. There exists a simple root system Π and

a subset $\Pi' \subseteq \Pi$ verifying $\text{card}(\Pi \setminus \Pi') = 1$ such that \mathfrak{l} is the sum of \mathfrak{h} and all the α -root spaces for α in the root subsystem generated by Π' . Define z to be the element in \mathfrak{h} such that

$$\alpha(z) = \begin{cases} 0 & \text{if } \alpha \in \Pi', \\ 1 & \text{if } \alpha \in \Pi \setminus \Pi'. \end{cases}$$

Then z is a generator of $\mathfrak{z}(\mathfrak{l})$ and all the eigenvalues of $\text{ad } z$ are integers.

Let $m \in \mathbb{N}$. Then $\text{ad } z$ induces a \mathbb{Z} -grading on \mathfrak{g}_m ,

$$\mathfrak{g}_m = \bigoplus_{k \in \mathbb{Z}} \mathfrak{g}_m(k) \quad \text{with} \quad \mathfrak{g}_m(k) := \{y \in \mathfrak{g}_m \mid [z, y] = ky\}.$$

Set

$$\mathfrak{p} = \bigoplus_{k \geq 0} \mathfrak{g}_0(k) \quad \text{and} \quad \mathfrak{u} = \bigoplus_{k > 0} \mathfrak{g}_0(k).$$

Then \mathfrak{p} is a parabolic subalgebra of \mathfrak{g} , where $\mathfrak{l} = \mathfrak{g}_0(0)$ is a Levi factor and whose nilpotent radical is \mathfrak{u} . Denote by P , L , and U the connected closed subgroups of G whose Lie algebras are \mathfrak{p} , \mathfrak{l} , and \mathfrak{u} , respectively.

Observe that

$$\mathfrak{l}_m = \mathfrak{z}(\mathfrak{l})_m \oplus [\mathfrak{l}_m, \mathfrak{l}_m] = \mathfrak{g}_m(0), \quad \mathfrak{p}_m = \bigoplus_{k \geq 0} \mathfrak{g}_m(k), \quad \mathfrak{u}_m = \bigoplus_{k > 0} \mathfrak{g}_m(k).$$

Remark 5.8. Clearly, for any nonzero integer k , we have $[z, \mathfrak{g}_m(k)] = \mathfrak{g}_m(k)$. In particular, $\mathfrak{g}_m(0) = (\mathfrak{g}_m)^z = \mathfrak{u}_{\mathfrak{g}_m}(\mathbb{C}z)$ where $\mathfrak{u}_{\mathfrak{g}_m}(\mathbb{C}z)$ is the normalizer of z in \mathfrak{g}_m . Also, if $x \in \mathfrak{g}_m(k)$, with $k \in \mathbb{N}^*$, then x is ad -nilpotent, and $e^{\text{ad } x} z = z + [x, z] = z - kx$.

Lemma 5.9. *Let $\lambda \in \mathbb{C}^*$, $x \in \mathfrak{g}_m(0)$, and $y \in \mathfrak{u}_m$. If x is ad -nilpotent in \mathfrak{g}_m then there exists $\tau \in U_m$ such that $\tau(\lambda z + x + y) = \lambda z + x$.*

Proof. For some $p > 0$, $y = y_p + t$ with $y_p \in \mathfrak{g}_m(p)$ and $t \in \sum_{k \geq p+1} \mathfrak{g}_m(k)$. Since x is ad -nilpotent, the sequence $((\text{ad } x)^n \mathfrak{g}_m(p))_{n \in \mathbb{N}}$ is decreasing and $(\text{ad } x)^n \mathfrak{g}_m(p) = \{0\}$ for $n \geq \dim \mathfrak{g}_m(p)$. Let $q \in \mathbb{N}$ be such that $y_p \in (\text{ad } x)^q \mathfrak{g}_m(p)$. Then

$$\begin{aligned} e^{(1/p\lambda) \text{ad } y_p} (\lambda z + x + y) &= \lambda z + e^{(1/p\lambda) \text{ad } y_p} x + e^{(1/p\lambda) \text{ad } y_p} t \\ &= \lambda z + x + (1/p\lambda)[y_p, x] + t' = \lambda z + x + y' \end{aligned}$$

with $t' \in \sum_{k \geq p+1} \mathfrak{g}_m(k)$, $y' := (1/p\lambda)[y_p, x] + t'$, and

$$(1/p\lambda)[y_p, x] \in (\text{ad } x)^{q+1} \mathfrak{g}_m(p).$$

Therefore we may start again with y' . After a finite number of steps, we come to an element in $\sum_{k \geq p+1} \mathfrak{g}_m(k)$. Then we start again with $p+1$ instead of p and, after a finite number of steps, we come to an element of the expected form, $\lambda z + x$. \square

Lemma 5.10. *Let Ω be an L -orbit contained in $\overline{\mathcal{O}}_l$ and let X be an irreducible component of $\pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega})$. Then*

$$\dim G_m \cdot (\mathfrak{z}(l) + X + \mathfrak{u}_m) = \dim X + 2 \dim \mathfrak{u}_m + 1.$$

Proof. Set

$$C := \mathfrak{z}(l) + X + \mathfrak{u}_m.$$

Since Ω and $\overline{\Omega}$ are L -stable, $\pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega})$ is L_m -stable and so is X . In addition, $\mathfrak{z}(l)$ is L_m -stable too. Hence, C is P_m -stable because

$$P_m \cdot C = L_m U_m \cdot (\mathfrak{z}(l) + X + \mathfrak{u}_m) = L_m \cdot (\mathfrak{z}(l) + X + \mathfrak{u}_m) \subset C.$$

Observe also that the elements of X are all ad-nilpotent.

Consider the action of P_m on $G_m \times C$ given by $\rho \cdot (\sigma, c) = (\sigma \rho^{-1}, \rho(c))$. Denote by $\overline{(\sigma, c)}$ the P_m -orbit of $(\sigma, c) \in G_m \times C$ with respect to this action, and denote by $G_m \times_{P_m} C$ the corresponding quotient space. The natural morphism

$$G_m \times C \rightarrow \mathfrak{g}, \quad (\sigma, c) \mapsto \sigma(c)$$

factors through the quotient and we obtain a morphism

$$\psi : G_m \times_{P_m} C \rightarrow \mathfrak{g}$$

whose image is $G_m \cdot C$. Since X and \mathfrak{u}_m are both closed cones, $z = 1_{G_m}(z)$ lies in the image of ψ and

$$\psi^{-1}(z) = \{\overline{(\sigma, c)} \in G_m \times_{P_m} C \mid \sigma(c) = z\}.$$

Let $\overline{(\sigma, c)} \in \psi^{-1}(z)$. Because z is ad-semisimple, c is also ad-semisimple. Since all elements of X are ad-nilpotent, we deduce that c does not belong to $X + \mathfrak{u}_m$. Also, since $U_m \subset P_m$, we may assume by [Lemma 5.9](#) that c is of the form $\lambda z + x$ with $\lambda \in \mathbb{C}^*$ and $x \in X$. Since $x \in \mathfrak{g}_m(0) = (\mathfrak{g}_m)^z$, we deduce from the uniqueness of the Jordan decomposition that $c = \lambda z$. In particular, σ is in the normalizer $N_G(\mathbb{C}z)$ of z in G , and $c = \sigma^{-1}(z)$.

According to [Remark 5.8](#), the identity component of the centralizer $C_{G_m}(z)$ of z in G_m is contained in P_m and has finite index in $N_{G_m}(\mathbb{C}z)$. Consequently, $\psi^{-1}(z)$ is a finite set. Thus, we get that $\dim G_m \cdot C = \dim G_m \times_{P_m} C$ because they are both irreducible subsets. To conclude, it suffices to observe that $\dim G_m - \dim P_m = \dim \mathfrak{u}_m$ and $\dim C = 1 + \dim X + \dim \mathfrak{u}_m$ since $\mathfrak{z}(l) = \mathbb{C}z$. \square

Since \mathfrak{g} is simple, its Killing form $\langle \cdot, \cdot \rangle$ is nondegenerate. Let us denote by ϕ the element of $\mathbb{C}[\mathfrak{g}]^G$ defined for all $x \in \mathfrak{g}$ by

$$\phi(x) = \langle x, x \rangle.$$

By our choice of z , $\phi(z)$ is a nonzero positive integer. Set

$$\mathcal{C} := \mathfrak{z}(\mathfrak{l}) + \overline{\mathcal{O}}_l + \mathfrak{u}$$

Lemma 5.11. *The nullvariety in \mathcal{C} of ϕ is $\overline{\mathcal{O}}_l + \mathfrak{u}$.*

Proof. First of all, $\overline{\mathcal{O}}_l + \mathfrak{u}$ is contained in the nullvariety in \mathcal{C} of ϕ . For the other inclusion, let $u = \lambda z + x + y$ be in \mathcal{C} , with $\lambda \in \mathbb{C}$, $x \in \overline{\mathcal{O}}_l$, and $y \in \mathfrak{u}$, such that $\phi(u) = 0$. We have

$$0 = \phi(u) = \langle \lambda z + x + y, \lambda z + x + y \rangle = \lambda^2 \langle z, z \rangle + \langle x, x \rangle = \lambda^2 \langle z, z \rangle$$

since \mathfrak{u} is orthogonal to \mathfrak{p} , $\mathfrak{z}(\mathfrak{l})$ is orthogonal to $[\mathfrak{l}, \mathfrak{l}] \oplus \mathfrak{u}$, and $\langle x, x \rangle = \phi(x) = 0$. Hence $\lambda = 0$ since $\phi(z) \neq 0$. So, u lies in $\overline{\mathcal{O}}_l + \mathfrak{u}$, whence the other inclusion. \square

Let $\phi^{(0)}, \dots, \phi^{(m)} \in \mathbb{C}[\mathfrak{g}_m]$ be the polynomials as defined in [Remark 2.2](#) relative to ϕ . According to [Lemma 2.11](#), they are G_m -invariant. In particular, $\phi^{(0)}$ is G_m -invariant.

Lemma 5.12. *Let Ω_l be an L -orbit contained in $\overline{\mathcal{O}}_l$ and set $\Omega_{\mathfrak{g}} := \text{Ind}_l^{\mathfrak{g}}(\Omega_l)$. Then:*

- (1) *the nullvariety in $G_m \cdot (\mathfrak{z}(\mathfrak{l}) + \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + \mathfrak{u}_m)$ of $\phi^{(0)}$ is contained in $\pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\Omega}_{\mathfrak{g}})$,*
- (2) *$\dim \pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\Omega}_{\mathfrak{g}}) \geq \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + 2 \dim \mathfrak{u}_m$.*

Proof. Let us denote by Y the nullvariety in $G_m \cdot (\mathfrak{z}(\mathfrak{l}) + \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + \mathfrak{u}_m)$ of $\phi^{(0)}$. First of all, observe that Y contains 0 because each of the spaces $\mathfrak{z}(\mathfrak{l})$, $\pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l)$, and \mathfrak{u}_m is a closed cone. In particular, Y is nonempty.

(1) Let $u = g \cdot (\lambda z + x + y)$ be in Y , with $g \in G_m$, $\lambda \in \mathbb{C}$, $x \in \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l)$, and $y \in \mathfrak{u}_m$, such that $\phi^{(0)}(u) = 0$. Since $\phi^{(0)}$ is G_m -invariant, setting $x_0 := \pi_{\overline{\mathcal{O}}_l, m}^{-1}(x)$ and $y_0 := \pi_{\mathfrak{u}, m}(y)$, we get

$$0 = \phi^{(0)}(u) = \phi^{(0)}(\lambda z + x + y) = \phi(\lambda z + x_0 + y_0) = \lambda^2 \phi(z)$$

by the computations of the proof of [Lemma 5.11](#). Hence $\lambda = 0$ since $\phi(z) \neq 0$. So u lies in $G_m \cdot (\pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + \mathfrak{u}_m)$. But

$$G_m \cdot (\pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + \mathfrak{u}_m) \subset G_m \cdot (\mathcal{I}_m(\overline{\mathcal{O}}_l) + \mathfrak{u}_m) \subset G_m \cdot \mathcal{I}_m(\overline{\mathcal{O}}_{\mathfrak{g}}) = \mathcal{I}_m(\overline{\mathcal{O}}_{\mathfrak{g}})$$

because $\mathcal{I}_m(\overline{\mathcal{O}}_{\mathfrak{g}})$ is G_m -invariant. Thus Y is contained in $\mathcal{I}_m(\overline{\mathcal{O}}_{\mathfrak{g}})$. Then it remains to observe that for $u \in Y$,

$$\pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}(u) \in G \cdot (\overline{\Omega}_l + \mathfrak{u}) = \overline{\Omega}_{\mathfrak{g}}$$

by [Lemma 5.3](#). In conclusion, Y is contained in $\pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\Omega}_{\mathfrak{g}})$.

(2) Let X be an irreducible component of $\pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l)$ of maximal dimension, and let Y' be the nullvariety in $G_m \cdot (\mathfrak{z}(\mathfrak{l}) + X + \mathfrak{u}_m)$ of $\phi^{(0)}$. The function $\phi^{(0)}$ is not identically zero on $G_m \cdot (\mathfrak{z}(\mathfrak{l}) + X + \mathfrak{u}_m)$ since $z \in G_m \cdot (\mathfrak{z}(\mathfrak{l}) + X + \mathfrak{u}_m)$ and

$\phi^{(0)}(z) = \phi(z) \neq 0$. Since Y' is irreducible, we deduce by [Lemma 5.10](#) and our choice of X that

$$\begin{aligned} \dim Y' &= \dim G_m \cdot (\mathfrak{z}(\mathfrak{l}) + X + \mathfrak{u}_m) - 1 \\ &= \dim X + 2 \dim \mathfrak{u}_m = \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + 2 \dim \mathfrak{u}_m, \end{aligned}$$

whence the statement, by (1). \square

Proposition 5.13. *If for some L -orbit Ω_l in $\overline{\mathcal{O}}_l$, we have*

$$\dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) \geq \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\mathcal{O}_l),$$

then

$$\dim \pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\Omega}_{\mathfrak{g}}) \geq \dim \pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\mathcal{O}_{\mathfrak{g}}),$$

where $\Omega_{\mathfrak{g}}$ is the induced nilpotent orbit of \mathfrak{g} from Ω_l .

Proof. Assume that for some L -orbit Ω_l in $\overline{\mathcal{O}}_l$, we have

$$\dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) \geq \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\mathcal{O}_l).$$

Then by [Lemma 5.12](#),

$$\begin{aligned} \dim \pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\Omega}_{\mathfrak{g}}) &\geq \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l) + 2 \dim \mathfrak{u}_m \\ &\geq \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\mathcal{O}_l) + 2 \dim \mathfrak{u}_m = (m+1) \dim \mathcal{O}_l + 2(m+1) \dim \mathfrak{u}. \end{aligned}$$

To conclude, it remains to observe that $\pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\mathcal{O}_{\mathfrak{g}})$ has dimension

$$(m+1) \dim \mathcal{O}_l + 2(m+1) \dim \mathfrak{u}$$

because $\dim \mathcal{O}_{\mathfrak{g}} = 2 \dim \mathfrak{u} + \dim \mathcal{O}_l$ from [Theorem 5.1](#). \square

Remark 5.14. The above proof actually shows that $\pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\Omega}_{\mathfrak{g}})$ has dimension at least $2(m+1) \dim \mathfrak{u} + \dim \pi_{\overline{\mathcal{O}}_l, m}^{-1}(\overline{\Omega}_l)$ even if Ω_l does not verify the hypothesis of the proposition. This can be used in practice to give an estimate of $\dim \pi_{\overline{\mathcal{O}}_{\mathfrak{g}}, m}^{-1}(\overline{\mathcal{O}}_{\mathfrak{g}} \setminus \mathcal{O}_{\mathfrak{g}})$.

We are now in a position to prove the main result of the section.

Proof of Theorem 5.6. Let \mathfrak{l} be a Levi subalgebra of \mathfrak{g} . Then there is a finite sequence of Levi subalgebras

$$\mathfrak{l} = \mathfrak{l}_0 \subset \mathfrak{l}_1 \subset \mathfrak{l}_2 \subset \cdots \subset \mathfrak{l}_k = \mathfrak{g}$$

such that \mathfrak{l}_{i-1} is a maximal Levi subalgebra of \mathfrak{l}_i for every $i \in \{1, \dots, k\}$.

Let \mathcal{O}_l be a nilpotent orbit of $\mathfrak{l} = \mathfrak{l}_0$ verifying $\text{RC}_2(m)$ for some $m \in \mathbb{N}$, and set for $i \in \{1, \dots, k\}$,

$$\mathcal{O}_{\mathfrak{l}_i} = \text{Ind}_{\mathfrak{l}_{i-1}}^{\mathfrak{l}_i}(\mathcal{O}_{\mathfrak{l}_{i-1}}).$$

Since induction is transitive (see [Remark 5.2\(2\)](#)),

$$\mathcal{O}_{\mathfrak{g}} := \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}_l) = \text{Ind}_{\mathfrak{l}_{k-1}}^{\mathfrak{g}}(\text{Ind}_{\mathfrak{l}_{k-2}}^{\mathfrak{l}_{k-1}}(\cdots(\text{Ind}_{\mathfrak{l}_0}^{\mathfrak{l}_1}(\mathcal{O}_l))))).$$

So, in order to proof [Theorem 5.6](#), we may assume that \mathfrak{l} is maximal in \mathfrak{g} . Let us write $\mathcal{O}_{\mathfrak{l}}$ as a product $\mathcal{O}_{\mathfrak{l}} = \mathcal{O}_{\tau_1} \times \cdots \times \mathcal{O}_{\tau_n}$, with the τ_j as in [Remark 5.2\(1\)](#). Since $\mathcal{O}_{\mathfrak{l}}$ verifies $\text{RC}_2(m)$, \mathcal{O}_{τ_j} verifies $\text{RC}_2(m)$ for some $j \in \{1, \dots, n\}$. Since \mathfrak{l} is maximal in \mathfrak{g} , either $\tau_j = \mathfrak{s}_j$ and $\text{Ind}_{\tau_j}^{\mathfrak{s}_j}(\mathcal{O}_{\tau_j})$ obviously verifies $\text{RC}_2(m)$ too, or τ_j is maximal in \mathfrak{s}_j and by [Proposition 5.13](#), $\text{Ind}_{\tau_j}^{\mathfrak{s}_j}(\mathcal{O}_{\tau_j})$ verifies $\text{RC}_2(m)$ as well. Indeed, since \mathcal{O}_{τ_j} verifies $\text{RC}_2(m)$, for some Ω_{τ_j} in $\overline{\mathcal{O}_{\tau_j}} \setminus \mathcal{O}_{\tau_j}$,

$$\dim \pi_{\overline{\mathcal{O}_{\tau_j}, m}}^{-1}(\overline{\Omega_{\tau_j}}) \geq \dim \pi_{\overline{\mathcal{O}_{\tau_j}, m}}^{-1}(\mathcal{O}_{\tau_j})$$

and [Proposition 5.13](#) applies. In both cases, by [Remark 5.2\(3\)](#), we conclude that $\mathcal{O}_{\mathfrak{g}} := \text{Ind}_{\mathfrak{l}}^{\mathfrak{g}}(\mathcal{O}_{\mathfrak{l}})$ verifies $\text{RC}_2(m)$. □

6. Consequence of [Theorem 5.6](#)

[Theorem 5.6](#) allows us to answer the reducibility problem for many nilpotent orbits.

Recall from the beginning of [Section 3](#) that if \mathcal{O} is a nilpotent orbit of a reductive Lie algebra \mathfrak{g} with simple factors $\mathfrak{s}_1, \dots, \mathfrak{s}_m$, then $\mathcal{O} = \mathcal{O}_1 \times \cdots \times \mathcal{O}_m$ where \mathcal{O}_i is a nilpotent orbit of \mathfrak{s}_i . We shall say that \mathcal{O} has a little factor if there exists i such that \mathcal{O}_i is a little nilpotent orbit of \mathfrak{s}_i .

The following result is a direct consequence of [Theorem 5.6](#) and [Proposition 4.2](#).

Theorem 6.1. *Any nilpotent orbit induced from a nilpotent orbit that has a little factor verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*

When \mathfrak{g} is simple, there is a unique nilpotent orbit $\mathcal{O}_{\text{subreg}}$ of \mathfrak{g} , called the subregular nilpotent orbit, such that $\mathcal{N}(\mathfrak{g}) \setminus \mathcal{O}_{\text{reg}} = \overline{\mathcal{O}_{\text{subreg}}}$. It has codimension $\text{rk } \mathfrak{g} + 2$ in \mathfrak{g} .

Corollary 6.2. *Assume that \mathfrak{g} simple and not of type $A_1, B_2 = C_2$, or G_2 . Then the subregular nilpotent orbit $\mathcal{O}_{\text{subreg}}$ of \mathfrak{g} verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$. In particular, $\mathcal{I}_m(\overline{\mathcal{O}_{\text{subreg}}})$ is reducible for every $m \in \mathbb{N}^*$.*

Proof. Assume first that \mathfrak{g} has type A_2 . Then $\mathfrak{g} = \mathfrak{sl}_3(\mathbb{C})$ and $\mathcal{O}_{\text{subreg}} = \mathcal{O}_{\text{min}} = \mathcal{O}_{(2,1)}$. Hence, $\mathcal{O}_{\text{subreg}}$ is little and verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$ according to [Corollary 4.3](#).

Assume now that \mathfrak{g} is simple with $\text{rank} \geq 3$. Then there exists a Levi subalgebra \mathfrak{l} of \mathfrak{g} such that $[\mathfrak{l}, \mathfrak{l}]$ is simple of type A_2 , and the subregular nilpotent orbit of \mathfrak{g} is induced from that of $[\mathfrak{l}, \mathfrak{l}]$ for dimension reasons (see [Theorem 5.1](#)). Therefore, the theorem follows from the case $\mathfrak{sl}_3(\mathbb{C})$ and [Theorem 6.1](#). □

Remark 6.3. Outside types A and B, the subregular nilpotent orbit of a simple Lie algebra is distinguished. Thus [Corollary 6.2](#) provides examples of distinguished nilpotent orbits which verify $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$. In particular, according to [Remark 3.8](#), these nilpotent orbits verify $\text{RC}_2(1)$ but not RC_1 .

Remark 6.4. For $\mathfrak{g} = \mathfrak{sp}_4(\mathbb{C}) \simeq \mathfrak{so}_5(\mathbb{C})$, we can show that $\mathcal{I}_1(\overline{\mathcal{O}_{\text{subreg}}})$ is irreducible.

Let us detail this example where the computations are explicit. Let $\mathfrak{g} = \mathfrak{sp}_4(\mathbb{C})$. The subregular nilpotent orbit is $\mathcal{O}_{(2^2)}$. By [Appendix A](#), it has dimension 6, and its singular locus is the union of two nilpotent orbits, $\mathcal{O}_{(2,1^2)} = \mathcal{O}_{\min}$ and the zero orbit.

Using [\[Weyman 2002, Theorem 1\]](#) — see also [\[Weyman 1989\]](#) or [\[Weyman 2003, Proposition 8.2.15\]](#) — and the realization of $\mathfrak{sp}_4(\mathbb{C})$ as the set of anti-self-adjoint matrices for the symplectic form, we can show that the defining ideal of $\overline{\mathcal{O}_{(2^2)}}$ is generated by the entries of the matrix X^2 as functions of $X \in \mathfrak{sp}_4(\mathbb{C})$.³ It follows that $\mathcal{I}_1(\overline{\mathcal{O}_{(2^2)}})$ can be identified with the scheme of pairs $(X_0, X_1) \in \mathfrak{sp}_4(\mathbb{C}) \times \mathfrak{sp}_4(\mathbb{C})$ defined by the equations $X_0^2 = 0$ and $X_0X_1 + X_1X_0 = 0$.

Using this identification, we obtain from direct computations that

$$\dim \pi_{\overline{\mathcal{O}_{(2^2)},1}}^{-1}(\mathcal{O}_{(2,1^2)}) = 11 \quad \text{and} \quad \dim \pi_{\overline{\mathcal{O}_{(2^2)},1}}^{-1}(0) = 10.$$

Furthermore, there are no smooth points of $\mathcal{I}_1(\overline{\mathcal{O}_{(2^2)}})$ in

$$\pi_{\overline{\mathcal{O}_{(2^2)},1}}^{-1}(\mathcal{O}_{(2,1^2)}) \cup \pi_{\overline{\mathcal{O}_{(2^2)},1}}^{-1}(0).$$

To see this, we have computed the dimension of the tangent space to $\mathcal{I}_1(\overline{\mathcal{O}_{(2^2)}})$ at generic points in $\pi_{\overline{\mathcal{O}_{(2^2)},1}}^{-1}(\mathcal{O}_{(2,1^2)})$ and $\pi_{\overline{\mathcal{O}_{(2^2)},1}}^{-1}(0)$, and the smallest dimensions turn out to be 13 and 14, respectively.

Now, if $\mathcal{I}_1(\overline{\mathcal{O}_{(2^2)}})$ were reducible, it would have an irreducible component of dimension 10 or 11 by the above equalities. This is not possible according to the computations of the tangent space dimensions. Hence, $\mathcal{I}_1(\overline{\mathcal{O}_{(2^2)}})$ is irreducible.

Classical types. We now summarize our conclusions for the case where \mathfrak{g} is simple of classical type. We refer to [Appendix A](#) for the notation relative to the induction of nilpotent orbits in the classical cases.

Theorem 6.5 (Type A). *Let $n \in \mathbb{N}^*$, $n \geq 2$, and let $\lambda \in \mathcal{P}(n)$ be nonrectangular. Then the nilpotent orbit \mathcal{O}_λ of $\mathfrak{sl}_n(\mathbb{C})$ verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$. In particular, $\mathcal{I}_m(\overline{\mathcal{O}_\lambda})$ is reducible for every $m \in \mathbb{N}^*$.*

Proof. Suppose that $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathcal{P}(n)$ is nonrectangular, with $1 < r < n$. Then there exists $1 \leq p < r$ such that $\lambda_p > \lambda_{p+1}$. It follows that

$$\lambda = \text{Ind}_{(n-p-r, p+r)}^n \mathbf{A},$$

where

$$\mathbf{A} = ((\lambda_1 - 2, \dots, \lambda_p - 2, \lambda_{p+1} - 1, \dots, \lambda_r - 1), (2^p, 1^{r-p})).$$

³Here, we have used the computer program Macaulay2 to check that these equations indeed generate a reduced ideal.

Thus any nonrectangular partition of n can be induced from a partition of the form $(2^p, 1^q)$ with $p, q \in \mathbb{N}^*$. According to [Example 4.4](#), $\mathcal{O}_{(2^p, 1^q)}$ is little for $p, q \in \mathbb{N}^*$. Hence the theorem follows from [Theorem 6.1](#). \square

Remark 6.6. It is not difficult to see that rectangular partitions can only be induced from a rectangular one. So they cannot be induced from a nilpotent orbit that has a little factor (see [Example 4.4](#)).

In fact, for the rectangular case, the theorem is not true. First of all, it is obviously not true for $\lambda = (n)$ and $\lambda = (1^n)$. Let us look at some special cases.

(1) Let $\lambda = (2^p)$ with $2p = n$. Then we saw in [Example 3.7](#) that \mathcal{O}_λ is RC_1 , and that all the irreducible components of $\mathcal{J}_1(\overline{\mathcal{O}}_\lambda)$ different from $\pi_{\overline{\mathcal{O}}_\lambda, 1}^{-1}(\mathcal{O}_\lambda)$ have codimension one. In particular, it is not $\text{RC}_2(1)$.

(2) Let $\lambda = (3^2)$. By [[Weyman 2002](#)] — see also [[Weyman 1989](#)] or [[Weyman 2003](#), Proposition 8.2.15] — the defining ideal of $\overline{\mathcal{O}}_\lambda$ is generated by $\text{tr}(X^2)$ and the entries of the matrix X^3 as functions of $X \in \mathfrak{sl}_6(\mathbb{C})$. By [Appendix A](#), the singular locus of $\overline{\mathcal{O}}_\lambda$ is the finite union of the nilpotent orbits \mathcal{O}_μ with

$$\mu \in \{(3, 2, 1), (3, 1^3), (2^3), (2^2, 1^2), (2, 1^4), (1^6)\} \subset \mathcal{P}(6),$$

and the respective dimensions of $\pi_{\overline{\mathcal{O}}_\lambda, 1}^{-1}(\mathcal{O}_\mu)$ are 47, 44, 44, 47, 44, 35. Note that $\mathcal{J}_1(\overline{\mathcal{O}}_\lambda)$ has dimension 48. Next, we obtain that the respective dimensions of the tangent space to $\mathcal{J}_1(\overline{\mathcal{O}}_\lambda)$ at generic points in $\pi_{\overline{\mathcal{O}}_\lambda, 1}^{-1}(\mathcal{O}_\mu)$, with μ running through the above set, are 49, 51, 51, 48, 52, 69. Arguing as in [Remark 6.4](#), we conclude that $\mathcal{J}_1(\overline{\mathcal{O}})$ is irreducible.

Therefore, from [Remark 6.6](#)(1) and (2), we have complete answers for the reducibility of $\mathcal{J}_1(\overline{\mathcal{O}})$ for any nilpotent orbit \mathcal{O} in $\mathfrak{sl}_n(\mathbb{C})$, for $n \leq 7$, and for any nilpotent orbit \mathcal{O} in $\mathfrak{sl}_p(\mathbb{C})$, with p a prime number.

In the other classical simple Lie algebras, we have the following result.

Theorem 6.7 (Types B, C, D). *Let $\lambda = (\lambda_1, \dots, \lambda_t) \in \mathcal{P}_\varepsilon(n)$ with $\varepsilon \in \{+1, -1\}$, and set $\lambda_{t+1} = 0$.*

- (1) *Suppose that $\varepsilon = +1$ and there exist $1 \leq k < \ell \leq t$ such that $\lambda_k \geq \lambda_{k+1} + 2$ and $\lambda_\ell \geq \lambda_{\ell+1} + 2$, then the nilpotent orbit \mathcal{O}_λ of $\mathfrak{so}_n(\mathbb{C})$ verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*
- (2) *Suppose that $\varepsilon = -1$ and there exist $1 \leq k < \ell \leq t$ such that $\lambda_k \geq \lambda_{k+1} + 2$ and $\lambda_\ell \geq \lambda_{\ell+1} + 2$, then the nilpotent orbit \mathcal{O}_λ of $\mathfrak{sp}_n(\mathbb{C})$ verifies $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$.*
- (3) *Suppose that $\varepsilon = +1$ and that λ is very even. Then both $\mathcal{O}_\lambda^{\text{I}}$ and $\mathcal{O}_\lambda^{\text{II}}$ verify $\text{RC}_2(m)$ for every $m \in \mathbb{N}^*$. (See [Appendix A](#) for the definition of “very even”.)*

In particular, $\mathcal{J}_m(\overline{\mathcal{O}}_\lambda)$ is reducible for every $m \in \mathbb{N}^$.*

Proof. Let $\lambda = (\lambda_1, \dots, \lambda_t) \in \mathcal{P}_\varepsilon(n)$, set $\lambda_{t+1} = 0$, and suppose that there exist $1 \leq k < \ell \leq t$ such that $\lambda_k \geq \lambda_{k+1} + 2$ and $\lambda_\ell \geq \lambda_{\ell+1} + 2$ as in the theorem. Then

$$\lambda = \text{Ind}_{(\ell+k, n-2(\ell+k))}^{n, \varepsilon} \Gamma$$

where

$$\Gamma := ((2^k, 1^{\ell-k}); (\lambda_1 - 4, \dots, \lambda_k - 4, \lambda_{k+1} - 2, \dots, \lambda_\ell - 2, \lambda_{\ell+1}, \dots, \lambda_t)).$$

So λ is induced from a partition in $\mathcal{P}(n)$ of the form $(2^p, 1^q)$, with $p, q \in \mathbb{N}^*$. By [Example 4.4](#), the partition $(2^p, 1^q)$ is little. This concludes the proof of parts (1) and (2) according to [Theorem 6.1](#).

Finally, if $\lambda \in \mathcal{P}_1(n)$ is very even, then \mathcal{O}_λ is induced from the nilpotent orbit $\mathcal{O}_{(2^t)}$ of $\mathfrak{so}_{2t}(\mathbb{C})$ which is little by [Example 4.5](#). Again, we conclude, thanks to [Theorem 6.1](#). \square

Remark 6.8. Unlike the type A case, in types B, C, D, orbits other than the ones considered in [Theorem 6.7](#) can be induced from little ones. For example, for $\lambda, p, q \in \mathbb{N}^*$ with p even, we have $\lambda = ((2\lambda)^p, (2\lambda - 1)^q) \in \mathcal{P}_1(2\lambda(p+q) - q)$ and λ does not verify the conditions of [Theorem 6.7](#). However, we have

$$\lambda = ((2\lambda)^p, (2\lambda - 1)^q) = \text{Ind}_{((\lambda-1)(p+q), 2p+q)}^{2\lambda(p+q)-q, 1} ((\lambda - 1)^{p+q}, (2^p, 1^q))$$

Since the nilpotent orbit of $\mathfrak{so}_{2p+q}(\mathbb{C})$ corresponding to the partition $(2^p, 1^q)$ is little (see [Example 4.5](#)), \mathcal{O}_λ verifies $\text{RC}_2(m)$ for all $m \in \mathbb{N}^*$.

Unfortunately, in types B, C, D, we have not found a nice exhaustive description of nilpotent orbits that can be reached by induction from a little nilpotent orbit. Computations using GAP4 show that a big proportion of partitions can be induced from little ones. See [Appendix B](#) for some numerical data.

Exceptional types. Our conclusions for the exceptional types are summarized in [Appendix C](#). More precisely, we can find in [Appendix C](#) the list of nilpotent orbits in a simple Lie algebra of exceptional type which can be induced from a little one.

7. Applications, remarks and comments

We give in this section applications to geometric properties of nilpotent orbit closures.

Nilpotent orbits closures and complete intersections. Let \mathcal{O} be a nilpotent orbit of the reductive Lie algebra \mathfrak{g} .

Theorem 7.1. *If \mathcal{O} verifies RC_1 or $\text{RC}_2(m)$ for some $m \geq 1$, then $\bar{\mathcal{O}}$ is not a complete intersection.*

Proof. Since the singular locus of $\bar{\mathcal{O}}$ is $\bar{\mathcal{O}} \setminus \mathcal{O}$ (see [Section 1](#)), it has codimension at least two in $\bar{\mathcal{O}}$. Hence, $\bar{\mathcal{O}}$ is normal if it is a complete intersection. If so, by [\[Hinich 1991\]](#) or [\[Panyushev 1991\]](#), it has rational singularities. The theorem is then of direct consequence of [Theorem 2.8](#). \square

In [\[Namikawa 2013; Brion and Fu 2015\]](#), the authors use symplectic resolutions of singularities of nilpotent orbit closures to prove the above corollary for arbitrary nilpotent orbits in \mathfrak{g} . The foregoing provides an alternative method to obtain that result through jet schemes in a large number of cases (see [Section 6](#)). There are other approaches in the jet scheme setting to show that $\bar{\mathcal{O}}$ is not a complete intersection. Let us give an example.

Example 7.2. The computations described in [Remark 6.6\(2\)](#), show that for generic

$$x \in \pi_{\bar{\mathcal{O}}(3^2), 1}^{-1}(\mathcal{O}(2^2, 1^2)),$$

the tangent space at x of $\mathcal{J}_1(\bar{\mathcal{O}}(3^2))$ has dimension $48 = \dim \mathcal{J}_1(\bar{\mathcal{O}}(3^2))$. Hence, such an x is a smooth point of $\mathcal{J}_1(\bar{\mathcal{O}}(3^2))$, because $\mathcal{J}_1(\bar{\mathcal{O}}(3^2))$ is irreducible, which does not belong to $\pi_{\bar{\mathcal{O}}(3^2), 1}^{-1}(\mathcal{O}(3^2))$. So,

$$(\mathcal{J}_1(\bar{\mathcal{O}}(3^2)))_{\text{reg}} \neq \pi_{\bar{\mathcal{O}}(3^2), 1}^{-1}(\mathcal{O}(3^2))$$

and by [Theorem 2.8\(3\)](#), $\bar{\mathcal{O}}(3^2)$ is not a complete intersection.

Unfortunately, these arguments cannot be used for the nilpotent orbit $\mathcal{O}(2^2)$ of $\mathfrak{sp}_4(\mathbb{C})$ because, in this case, the computations of [Remark 6.4](#) show that we exactly have $(\mathcal{J}_1(\bar{\mathcal{O}}(2^2)))_{\text{reg}} = \pi_{\bar{\mathcal{O}}(2^2), 1}^{-1}(\mathcal{O}(2^2))$.

Examples and counterexamples. Our results provide many examples showing that the converse of [Proposition 2.5](#) for irreducibility is not true. Since the nilpotent cone $\mathcal{N}(\mathfrak{g})$ is normal, the following result illustrates that the converse of [Proposition 2.5](#) for normality is also not true.

Proposition 7.3. *Assume that \mathfrak{g} simple, and let $m \in \mathbb{N}$. Then $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ is normal if and only if $m = 0$.*

Proof. Since $\mathcal{J}_0(\mathcal{N}(\mathfrak{g})) \simeq \mathcal{N}(\mathfrak{g})$ is normal, we have to show that for any $m \in \mathbb{N}^*$, $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ is not normal.

Fix $m \in \mathbb{N}^*$. Let ℓ be the rank of \mathfrak{g} , and let p_1, \dots, p_ℓ be homogeneous generators of $\mathbb{C}[\mathfrak{g}]^G$ so that

$$\mathcal{N}(\mathfrak{g}) = \text{Spec } \mathbb{C}[\mathfrak{g}]/(p_1, \dots, p_\ell).$$

By [Remark 2.2](#), we get

$$\mathcal{J}_m(\mathcal{N}(\mathfrak{g})) \simeq \text{Spec } \mathbb{C}[\mathfrak{g}_m]/(p_i^{(j)} \mid i = 1, \dots, \ell, j = 0, \dots, m).$$

Since $\mathcal{N}(\mathfrak{g})$ is a complete intersection with rational singularities, $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ is irreducible and reduced by [Theorem 2.8](#). So, it is generically reduced. Furthermore,

$(\mathcal{J}_m(\mathcal{N}(\mathfrak{g})))_{\text{reg}}$ consists of the set of $x = x_0 + x_1t + \dots + x_mt^m \in \mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$ such that, for $i = 1, \dots, \ell$ and $j = 0, \dots, m$,

$$(2) \quad dp_i^{(j)}(x_0, x_1, \dots, x_m) \text{ are linearly independent.}$$

Let $x_0 + x_1t + \dots + x_mt^m \in \mathfrak{g}_m$. By [Raïs and Tauvel 1992, Lemma 3.3(i)], the vectors $dp_i^{(j)}(x_0, x_1, \dots, x_m)$ for $i \in \{1, \dots, \ell\}$ and $j \in \{0, \dots, m\}$ are linearly independent if and only if the vectors $dp_1(x_0), \dots, dp_\ell(x_0)$ are linearly independent. But by [Kostant 1963], the later condition is satisfied if and only if x_0 is a regular element of \mathfrak{g} . Therefore by (2),

$$(3) \quad (\mathcal{J}_m(\mathcal{N}(\mathfrak{g})))_{\text{reg}} = \pi_{\mathcal{N}(\mathfrak{g}),m}^{-1}(\mathcal{O}_{\text{reg}}) \quad \text{and} \quad (\mathcal{J}_m(\mathcal{N}(\mathfrak{g})))_{\text{sing}} = \pi_{\mathcal{N}(\mathfrak{g}),m}^{-1}(\overline{\mathcal{O}_{\text{subreg}}})$$

since $\mathcal{N}(\mathfrak{g}) \setminus \mathcal{O}_{\text{reg}} = \overline{\mathcal{O}_{\text{subreg}}}$. Then by Serre's criterion, it is enough to show that $\pi_{\mathcal{N}(\mathfrak{g}),m}^{-1}(\overline{\mathcal{O}_{\text{subreg}}})$ has codimension one in $\mathcal{J}_m(\mathcal{N}(\mathfrak{g}))$, or else that

$$(4) \quad \dim \pi_{\mathcal{N}(\mathfrak{g}),m}^{-1}(\overline{\mathcal{O}_{\text{subreg}}}) \geq \dim \mathcal{J}_m(\mathcal{N}(\mathfrak{g})) - 1.$$

The zero orbit of $\mathfrak{sl}_2(\mathbb{C})$ has codimension 2 in $\mathcal{N}(\mathfrak{sl}_2(\mathbb{C}))$. Hence, for dimension reasons, $\mathcal{O}_{\text{subreg}}$ is the induced nilpotent orbit from 0 in any Levi subalgebra \mathfrak{l} of \mathfrak{g} with semisimple part $[\mathfrak{l}, \mathfrak{l}]$ isomorphic to $\mathfrak{sl}_2(\mathbb{C})$. So by Remark 5.14, in order to prove (4), it suffices to show the statement for $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$.

If $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{C})$, then $\overline{\mathcal{O}_{\text{subreg}}} = 0$, but by Lemma 3.5(2),

$$\begin{aligned} \dim \pi_{\mathcal{N}(\mathfrak{sl}_2(\mathbb{C})),m}^{-1}(0) &\geq \dim \mathcal{J}_{m-2}(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C}))) + \dim \mathfrak{sl}_2(\mathbb{C}) \\ &= 2(m-1) + 3 = 2m + 1, \end{aligned}$$

whence the expected result since $\dim \mathcal{J}_m(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C}))) = 2(m+1) = 2m + 2$. \square

Remark 7.4. For $m = 1$, (3) is also a consequence of Theorem 2.8(3).

We now give an example illustrating the fact that the converse of Proposition 2.5 is also not true for reducedness.

Example 7.5. The scheme $\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C})))$ is irreducible and reduced. We readily obtain from the description of $\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C})))$ given in Example 2.3 that $\mathcal{J}_1(\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C}))))$ is defined by the ideal \mathcal{J} of

$$\mathbb{C}[x_0, y_0, z_0, x_1, y_1, z_1, x'_0, y'_0, z'_0, x'_1, y'_1, z'_1]$$

generated by the polynomials

$$\begin{aligned} x_0^2 + y_0z_0, \quad 2x_0x_1 + y_0z_1 + y_1z_0, \\ 2x_0x'_0 + y_0z'_0 + z_0y'_0, \quad 2x_0x'_1 + 2x_1x'_0 + y_0z'_1 + y_1z'_0 + z_1y'_0 + z_0y'_1. \end{aligned}$$

A computation made with the program Macaulay2 shows that \mathcal{J} is not radical, and that the radical of \mathcal{J} is the intersection of two prime ideals. So, $\mathcal{J}_1(\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C}))))$ is neither reduced nor irreducible.

Example 7.5 gives another piece of evidence that $\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C})))$ does not have rational singularities (see [Proposition 7.3](#)). Indeed, if it did, then by [Theorem 2.8](#), $\mathcal{J}_1(\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C}))))$ would be irreducible (and reduced) because $\mathcal{J}_1(\mathcal{N}(\mathfrak{sl}_2(\mathbb{C})))$ is a complete intersection.

We now turn to other interesting phenomena.

Example 7.6. As has been observed in [Example 3.7](#), for the nilpotent orbit $\mathcal{O}_{(2^p)}$ of $\mathfrak{sl}_{2p}(\mathbb{C})$, with $p \geq 2$, $\mathcal{J}_1(\overline{\mathcal{O}_{(2^p)}})$ is reducible and

$$\dim \pi_{\overline{\mathcal{O}_{(2^p)}}, 1}^{-1}((\overline{\mathcal{O}_{(2^p)}})_{\text{sing}}) < \dim \pi_{\overline{\mathcal{O}_{(2^p)}}, 1}^{-1}(\mathcal{O}_{(2^p)}).$$

This shows that [Lemma 2.7\(3\)](#), does not hold in general if X is not a complete intersection.

Example 7.7. As has been observed in [Remark 6.6\(2\)](#), for the nilpotent orbit $\mathcal{O}_{(3^2)}$ of $\mathfrak{sl}_6(\mathbb{C})$, $\mathcal{J}_1(\overline{\mathcal{O}_{(3^2)}})$ is irreducible and

$$(\mathcal{J}_1(\overline{\mathcal{O}_{(3^2)}}))_{\text{reg}} \neq \pi_{\overline{\mathcal{O}_{(3^2)}}, 1}^{-1}(\mathcal{O}_{(3^2)}).$$

This shows that [Theorem 2.8\(3\)](#) is not true for varieties that are not locally complete intersection.

Example 7.8. For the nilpotent orbit $\mathcal{O}_{(2^2)}$ of $\mathfrak{sp}_4(\mathbb{C})$, we have observed (see [Remark 6.4](#)) that

$$(\mathcal{J}_1(\overline{\mathcal{O}_{(2^2)}}))_{\text{reg}} = \pi_{\overline{\mathcal{O}_{(2^2)}}, 1}^{-1}(\mathcal{O}_{(2^2)}).$$

This shows that the equality of [Theorem 2.8\(3\)](#) may hold even if X is not locally a complete intersection.

Questions and remarks. Although we have determined the reducibility of the closure of many nilpotent orbits, we would like to complete the cases where our methods do not apply. Here are some open questions.

Question 7.9. We have seen that jet schemes of nilpotent orbits in $\mathfrak{sl}_n(\mathbb{C})$ corresponding to rectangular partitions can be irreducible or reducible. Is there an explicit characterization?

Question 7.10. In all our examples of nilpotent orbits \mathcal{O} with $\mathcal{J}_1(\overline{\mathcal{O}})$ reducible, the orbit \mathcal{O} verifies RC_1 or $\text{RC}_2(1)$. Are these conditions necessary or are there examples of \mathcal{O} for which $\mathcal{J}_1(\overline{\mathcal{O}})$ is reducible and that verify neither RC_1 nor $\text{RC}_2(1)$?

We have used the reducibility of jet schemes to study the property of complete intersection for nilpotent orbit closures. It is very likely that other geometric properties of nilpotent orbit closures can be studied using jet schemes.

Appendix A: Nilpotent orbits in classical simple Lie algebras

We fix in this appendix some notation and basic results, relative to nilpotent orbits in simple Lie algebras of classical type. Our main references are [Collingwood and McGovern 1993; Kempken 1983]. The results concerning the induction of nilpotent orbits are mostly taken from [loc. cit.].

Let $n \in \mathbb{N}^*$, and denote by $\mathcal{P}(n)$ the set of partitions of n . As a rule, unless otherwise specified, we write an element λ of $\mathcal{P}(n)$ as a decreasing sequence $\lambda = (\lambda_1, \dots, \lambda_r)$ omitting the zeroes. Thus,

$$\lambda_1 \geq \dots \geq \lambda_r \geq 1 \quad \text{and} \quad \lambda_1 + \dots + \lambda_r = n.$$

We shall denote the dual partition of a partition $\lambda \in \mathcal{P}(n)$ by ${}^t\lambda$. The *concatenation* of two partitions λ and λ' will be the rearrangement of the parts in decreasing order, and shall be denoted by $\lambda \smile \lambda'$.

Let us denote by \geq the partial order on $\mathcal{P}(n)$ relative to dominance. More precisely, given $\lambda = (\lambda_1, \dots, \lambda_r)$, $\mu = (\mu_1, \dots, \mu_s) \in \mathcal{P}(n)$, we have $\lambda \geq \mu$ if

$$\sum_{i=1}^k \lambda_i \geq \sum_{i=1}^k \mu_i$$

for $1 \leq k \leq \min(r, s)$.

Case $\mathfrak{sl}_n(\mathbb{C})$. According to [Collingwood and McGovern 1993, Theorem 5.1.1], nilpotent orbits of $\mathfrak{sl}_n(\mathbb{C})$ are parametrized by $\mathcal{P}(n)$. For $\lambda \in \mathcal{P}(n)$, we shall denote by \mathcal{O}_λ the corresponding nilpotent orbit of $\mathfrak{sl}_n(\mathbb{C})$, and if we write ${}^t\lambda = (d_1, \dots, d_s)$, then

$$\dim \mathcal{O}_\lambda = n^2 - \sum_{i=1}^s d_i^2.$$

Also, if $\lambda, \mu \in \mathcal{P}(n)$, then $\mathcal{O}_\mu \subset \overline{\mathcal{O}_\lambda}$ if and only if $\mu \leq \lambda$.

The Levi subalgebras of $\mathfrak{sl}_n(\mathbb{C})$ are parametrized by compositions of n . Let $\mathbf{m} = (m_1, \dots, m_r)$ be a composition of n and

$$\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)}) \in \mathcal{P}(m_1) \times \dots \times \mathcal{P}(m_r).$$

It corresponds to a nilpotent orbit in the Levi subalgebra associated to the composition \mathbf{m} . Set

$$\boldsymbol{\mu} := {}^t\lambda^{(1)} \smile \dots \smile {}^t\lambda^{(r)} \quad \text{and} \quad \mathbf{v} = {}^t\boldsymbol{\mu}.$$

Then the partition associated to the induced nilpotent orbit from $\mathcal{O}_{(\lambda^{(1)}, \dots, \lambda^{(r)})}$ is \mathbf{v} . Note that we have $v_i = \lambda_i^{(1)} + \dots + \lambda_i^{(k)}$ which is much simpler to compute in practice. We shall denote \mathbf{v} by $\text{Ind}_{\mathbf{m}}^n(\lambda^{(1)}, \dots, \lambda^{(r)})$ and we shall say that \mathbf{v} is *induced from* $(\lambda^{(1)}, \dots, \lambda^{(r)})$.

Case $\mathfrak{so}_n(\mathbb{C})$. For $n \in \mathbb{N}^*$, set

$$\mathcal{P}_1(n) := \{\lambda \in \mathcal{P}(n) \mid \text{number of parts of each even number in } \lambda \text{ is even}\}.$$

According to [Collingwood and McGovern 1993, Theorems 5.1.2 and 5.1.4], nilpotent orbits of $\mathfrak{so}_n(\mathbb{C})$ are parametrized by $\mathcal{P}(n)$, with the exception that each *very even* partition $\lambda \in \mathcal{P}_1(n)$ (i.e., λ has only even parts) corresponds to two nilpotent orbits. For $\lambda \in \mathcal{P}_1(n)$, not very even, we shall denote by \mathcal{O}_λ the corresponding nilpotent orbit of $\mathfrak{so}_n(\mathbb{C})$. For very even $\lambda \in \mathcal{P}_1(n)$, we shall denote by \mathcal{O}_λ^I and \mathcal{O}_λ^{II} the two corresponding nilpotent orbits of $\mathfrak{so}_n(\mathbb{C})$. In fact, their union form a single $\mathcal{O}_n(\mathbb{C})$ -orbit.

Let $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathcal{P}_1(n)$ and ${}^t\lambda = (d_1, \dots, d_s)$. Then

$$\dim \mathcal{O}_\lambda^* = \frac{n(n-1)}{2} - \frac{1}{2} \left(\sum_{i=1}^s d_i^2 - \#\{i \mid \lambda_i \text{ odd}\} \right)$$

where \mathcal{O}_λ^* is either \mathcal{O}_λ , \mathcal{O}_λ^I , or \mathcal{O}_λ^{II} according to whether λ is very even or not. Using the same notation, if $\lambda, \mu \in \mathcal{P}_1(n)$, then $\overline{\mathcal{O}_\mu^*} \subsetneq \overline{\mathcal{O}_\lambda^*}$ if and only if $\mu < \lambda$.

Given $\lambda \in \mathcal{P}(n)$, there exists a unique $\lambda^+ \in \mathcal{P}_1(n)$ such that $\lambda^+ \leq \lambda$, and if $\mu \in \mathcal{P}_1(n)$ satisfies $\mu \leq \lambda$, then $\mu \leq \lambda^+$. More precisely, let $\lambda = (\lambda_1, \dots, \lambda_n)$ (adding zeroes if necessary). If $\lambda \in \mathcal{P}_1(n)$, then $\lambda^+ = \lambda$, and if $\lambda \notin \mathcal{P}_1(n)$, set

$$\lambda' = (\lambda_1, \dots, \lambda_r, \lambda_{r+1} - 1, \lambda_{r+2}, \dots, \lambda_{s-1}, \lambda_s + 1, \lambda_{s+1}, \dots, \lambda_n)$$

where r is maximum such that $(\lambda_1, \dots, \lambda_r) \in \mathcal{P}_1(\lambda_1 + \dots + \lambda_r)$, and s is the index of the first even part in $(\lambda_{r+2}, \dots, \lambda_n)$. Note that $r = 0$ if such a maximum does not exist, while s is always defined. If λ' is not in $\mathcal{P}_1(n)$, then we repeat the process until we obtain an element of $\mathcal{P}_1(n)$ which will be our λ^+ .

The Levi subalgebras in $\mathfrak{so}_n(\mathbb{C})$ are parametrized by

$$\mathcal{L}(n) := \left\{ (p_1, \dots, p_k; r) \mid 2 \sum_{i=1}^k p_i + r = n \right\}.$$

Let $(p_1, \dots, p_k; r) \in \mathcal{L}(n)$, $(\lambda^{(1)}, \dots, \lambda^{(k)}) \in \mathcal{P}(p_1) \times \dots \times \mathcal{P}(p_k)$ and $\mu \in \mathcal{P}(r)$, and set

$$\mathbf{v} := \text{Ind}_{(p_1, \dots, p_k, r, p_k, \dots, p_1)}^n (\lambda^{(1)}, \dots, \lambda^{(k)}, \mu, \lambda^{(k)}, \dots, \lambda^{(1)})$$

in the notation of the $\mathfrak{sl}_n(\mathbb{C})$ case. Thus \mathbf{v} is the partition associated to the nilpotent orbit in $\mathfrak{sl}_n(\mathbb{C})$ induced from the nilpotent orbit in the Levi subalgebra of $\mathfrak{sl}_n(\mathbb{C})$ associated to the composition $(p_1, \dots, p_k, r, p_k, \dots, p_1)$ and the multipartition $(\lambda^{(1)}, \dots, \lambda^{(k)}, \mu, \lambda^{(k)}, \dots, \lambda^{(1)})$. The partition associated to the nilpotent orbit induced from $(\lambda^{(1)}, \dots, \lambda^{(k)}; \mu)$ is \mathbf{v}^+ . We shall denote \mathbf{v}^+ by

$$\text{Ind}_{(p_1, \dots, p_k; r)}^{n,+} (\lambda^{(1)}, \dots, \lambda^{(k)}; \mu).$$

The partition $\lambda \in \mathcal{P}_1(n)$ corresponds to a rigid orbit if and only if

- (i) $\lambda_i - \lambda_{i+1} \leq 1$ for all i , so the last part of λ is 1;
- (ii) no odd number occurs exactly twice in λ .

Note that in the case that λ is a very even partition, \mathfrak{v}^+ is also very even, and we obtain both nilpotent orbits corresponding to \mathfrak{v}^+ via induction of the nilpotent orbits corresponding to λ ; see [Collingwood and McGovern 1993, Theorem 7.3.3(iii)].

Case $\mathfrak{sp}_{2n}(\mathbb{C})$. For $n \in \mathbb{N}^*$, set

$$\mathcal{P}_{-1}(2n) := \{\lambda \in \mathcal{P}(2n) \mid \text{number of parts of each odd number is even}\}.$$

According to [op. cit., Theorem 5.1.3], nilpotent orbits of $\mathfrak{sp}_{2n}(\mathbb{C})$ are parametrized by $\mathcal{P}_{-1}(2n)$. For $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathcal{P}_{-1}(2n)$, we shall denote by \mathcal{O}_λ the corresponding nilpotent orbit of $\mathfrak{sp}_{2n}(\mathbb{C})$, and if we write ${}^t\lambda = (d_1, \dots, d_s)$, then

$$\dim \mathcal{O}_\lambda = n(2n + 1) - \frac{1}{2} \left(\sum_{i=1}^s d_i^2 + \#\{i \mid \lambda_i \text{ odd}\} \right).$$

As in the case of $\mathfrak{sl}_n(\mathbb{C})$, if $\lambda, \mu \in \mathcal{P}_{-1}(2n)$, then $\mathcal{O}_\mu \subset \overline{\mathcal{O}_\lambda}$ if and only if $\mu \leq \lambda$.

Given $\lambda \in \mathcal{P}(2n)$, there exists a unique $\lambda^- \in \mathcal{P}_{-1}(2n)$ such that $\lambda^- \leq \lambda$, and if $\mu \in \mathcal{P}_{-1}(2n)$ satisfies $\mu \leq \lambda$, then $\mu \leq \lambda^-$. The construction of λ^- is the same as in the orthogonal case except that s is the index of the first odd part in $(\lambda_{r+2}, \dots, \lambda_{2n})$.

As in the orthogonal case, Levi subalgebras are parametrized by $\mathcal{L}(2n)$. Let us conserve the same notations as in the orthogonal case. The partition associated to the nilpotent orbit induced from $(\lambda^{(1)}, \dots, \lambda^{(k)}; \mu)$ is \mathfrak{v}^- . We shall denote \mathfrak{v}^- by

$$\text{Ind}_{(p_1, \dots, p_k; r)}^{2n, -}(\lambda^{(1)}, \dots, \lambda^{(k)}; \mu).$$

The partition $\lambda \in \mathcal{P}_{-1}(2n)$ corresponds to a rigid orbit if and only if

- (i) $\lambda_i - \lambda_{i+1} \leq 1$ for all i , so the last part of λ is 1;
- (ii) no even number occurs exactly twice in λ .

Appendix B: Statistics in types B, C, and D

As mentioned in Remark 6.8, many nilpotent orbits in $\mathfrak{so}_n(\mathbb{C})$ and $\mathfrak{sp}_{2n}(\mathbb{C})$ can be obtained by induction from little nilpotent orbits. In particular, these induced orbits verify $\text{RC}_2(m)$ for all $m \in \mathbb{N}^*$. Computations using GAP4 gave us the following numerical data supporting our claim.

For $\varepsilon \in \{-1, +1\}$ and $n \in \mathbb{N}^*$, we denote by $\mathcal{P}_\varepsilon^\ell(n)$ the set of partitions in $\mathcal{P}_\varepsilon(n)$ induced from little ones.

Case $\mathfrak{so}_n(\mathbb{C})$.

n	$\#\mathcal{P}_1^\ell(n)$	$\#\mathcal{P}_1(n)$	n	$\#\mathcal{P}_1^\ell(n)$	$\#\mathcal{P}_1(n)$	n	$\#\mathcal{P}_1^\ell(n)$	$\#\mathcal{P}_1(n)$
2	0	1	19	111	130	36	2370	2741
3	0	2	20	130	161	37	2821	3206
4	1	3	21	170	196	38	3265	3740
5	1	4	22	195	236	39	3852	4368
6	2	5	23	250	287	40	4460	5096
7	4	7	24	291	350	41	5242	5922
8	6	10	25	367	420	42	6064	6868
9	9	13	26	423	501	43	7086	7967
10	10	16	27	527	602	44	8182	9233
11	16	21	28	609	722	45	9536	10670
12	20	28	29	751	858	46	10986	12306
13	27	35	30	869	1016	47	12748	14193
14	32	43	31	1055	1206	48	14667	16357
15	45	55	32	1223	1431	49	16974	18803
16	52	70	33	1474	1687	50	19485	21581
17	73	86	34	1710	1981	51	22464	24766
18	83	105	35	2039	2331			


Case $\mathfrak{sp}_{2n}(\mathbb{C})$.

n	$\#\mathcal{P}_{-1}^\ell(2n)$	$\#\mathcal{P}_{-1}(2n)$	n	$\#\mathcal{P}_{-1}^\ell(2n)$	$\#\mathcal{P}_{-1}(2n)$
1	0	2	13	594	728
2	1	4	14	857	1040
3	3	8	15	1223	1472
4	9	14	16	1726	2062
5	15	24	17	2421	2864
6	28	40	18	3378	3948
7	45	64	19	4652	5400
8	77	100	20	6374	7336
9	119	154	21	8677	9904
10	182	232	22	11728	13288
11	273	344	23	15755	17728
12	409	504	24	21061	23528

Appendix C: Tables for exceptional types


We list on the next few pages nilpotent orbits in a simple Lie algebra of exceptional type specifying when possible whether they are RC_1 or $\text{RC}_2(m)$. Condition RC_1 is

Type G_2 .




\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
A_1 [0, 1]	6	$\checkmark \leftarrow$ little	$\checkmark \leftarrow$ little	\checkmark
\tilde{A}_1 [1, 0]	8	\times	?	\checkmark
$G_2(a_1)$ [2, 0]	10	\times	?	\times

Type F_4 .




\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
A_1 [1, 0, 0, 0]	16	$\checkmark \leftarrow$ little	$\checkmark \leftarrow$ little	\checkmark
\tilde{A}_1 [0, 0, 0, 1]	22	$\checkmark \leftarrow$ little	$\checkmark \leftarrow$ little	\checkmark
$A_1 + \tilde{A}_1$ [0, 1, 0, 0]	28	\times	?	\checkmark
A_2 [2, 0, 0, 0]	30	\times	?	\times
\tilde{A}_2 [0, 0, 0, 2]	30	\times	?	\times
$A_2 + \tilde{A}_1$ [0, 0, 1, 0]	34	\times	?	\checkmark
B_2 [2, 0, 0, 1]	36	\times	$\checkmark \leftarrow \mathcal{O}_{\min}^{\{2,3,4\}}$	\times
$\tilde{A}_2 + A_1$ [0, 1, 0, 1]	36	\times	?	\checkmark
$C_3(a_1)$ [1, 0, 1, 0]	38	\times	$\checkmark \leftarrow \mathcal{O}_{\min}^{\{1,2,3\}}$	\times
$F_4(a_3)$ [0, 2, 0, 0]	40	\times	$\checkmark \leftarrow \mathcal{O}_{[0,1,0]}^{\{2,3,4\}}$	\times
B_3 [2, 2, 0, 0]	42	\times	?	\times
C_3 [1, 0, 1, 2]	42	\times	?	\times
$F_4(a_2)$ [0, 2, 0, 2]	44	\times	$\checkmark \leftarrow \mathcal{O}_{(2,1),(1^2)}^{\{1,2,4\}}$	\times
$F_4(a_1)$ [2, 2, 0, 2]	46	\times	$\checkmark \leftarrow \mathcal{O}_{(2,1),(2)}^{\{1,2,4\}}$	\times

Type E₆. The notation $\text{Res}_{E_6}^{E_7} \mathcal{O}$ means that the orbit is obtained by restriction from the little nilpotent orbit \mathcal{O} in E_7 as explained in Table 1.




	\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
A_1	$[0, 1, 0, 0, 0, 0]$	22	$\checkmark \leftarrow$ little	$\checkmark \leftarrow$ little	\checkmark
$2A_1$	$[1, 0, 0, 0, 0, 1]$	32	$\checkmark \leftarrow$ little	$\checkmark \leftarrow$ little	\times
$3A_1$	$[0, 0, 0, 1, 0, 0]$	40	$\checkmark \leftarrow \text{Res}_{E_6}^{E_7} \mathcal{O}_{(3A_1)'}$?	\checkmark
A_2	$[0, 2, 0, 0, 0, 0]$	42	$\checkmark \leftarrow \text{Res}_{E_6}^{E_7} \mathcal{O}_{A_2}$?	\times
$A_2 + A_1$	$[1, 1, 0, 0, 0, 1]$	46	\times	$\checkmark \leftarrow \mathcal{O}_{\min}^{\{1,2,3,4,5\}}$	\times
$2A_2$	$[2, 0, 0, 0, 0, 2]$	48	\times	$\checkmark \leftarrow \mathcal{O}_{(3,1^7)}^{\{1,2,3,4,5\}}$	\times
$A_2 + 2A_1$	$[0, 0, 1, 0, 1, 0]$	50	\times	?	\times
A_3	$[1, 2, 0, 0, 0, 1]$	52	\times	$\checkmark \leftarrow \mathcal{O}_{(2,1^4)}^{\{1,3,4,5,6\}}$	\times
$2A_2 + A_1$	$[1, 0, 0, 1, 0, 1]$	54	\times	?	\checkmark
$A_3 + A_1$	$[0, 1, 1, 0, 1, 0]$	56	\times	?	\times
$D_4(a_1)$	$[0, 0, 0, 2, 0, 0]$	58	\times	$\checkmark \leftarrow \mathcal{O}_{(2^2, 1^2)}^{\{1,3,4,5,6\}}$	\times
A_4	$[2, 2, 0, 0, 0, 2]$	60	\times	$\checkmark \leftarrow \mathcal{O}_{(3,1^3)}^{\{1,3,4,5,6\}}$	\times
D_4	$[0, 2, 0, 2, 0, 0]$	60	\times	?	\times
$A_4 + A_1$	$[1, 1, 1, 0, 1, 1]$	62	\times	$\checkmark \leftarrow \mathcal{O}_{((2,1^2); (1^2))}^{\{1,2,3,4,6\}}$	\times
A_5	$[2, 1, 1, 0, 1, 2]$	64	\times	?	\times
$D_5(a_1)$	$[1, 2, 1, 0, 1, 1]$	64	\times	$\checkmark \leftarrow \mathcal{O}_{(3,2,1)}^{\{1,3,4,5,6\}}$	\times
$E_6(a_3)$	$[2, 0, 0, 2, 0, 2]$	66	\times	$\checkmark \leftarrow \mathcal{O}_{(4,1^2)}^{\{1,3,4,5,6\}}$	\times
D_5	$[2, 2, 0, 2, 0, 2]$	68	\times	$\checkmark \leftarrow \mathcal{O}_{(4,2)}^{\{1,3,4,5,6\}}$	\times
$E_6(a_1)$	$[2, 2, 2, 0, 2, 2]$	70	\times	$\checkmark \leftarrow \mathcal{O}_{(5,1)}^{\{1,3,4,5,6\}}$	\times

Type E7. Note that the characteristics $[0, 0, 0, 0, 2, 0]$ and $[0, 0, 0, 0, 0, 2]$ of nilpotent orbits in D_6 correspond to the very even partition (2^6) .




	\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
A_1	$[1, 0, 0, 0, 0, 0, 0]$	34	✓ ← little	✓ ← little	✓
$2A_1$	$[0, 0, 0, 0, 0, 1, 0]$	52	✓ ← little	✓ ← little	✓
$(3A_1)''$	$[0, 0, 0, 0, 0, 0, 2]$	54	✓ ← little	✓ ← little	×
$(3A_1)'$	$[0, 0, 1, 0, 0, 0, 0]$	64	✓ ← little	✓ ← little	✓
A_2	$[2, 0, 0, 0, 0, 0, 0]$	66	✓ ← little	✓ ← little	×
$4A_1$	$[0, 1, 0, 0, 0, 0, 1]$	70	×	?	✓
$A_2 + A_1$	$[1, 0, 0, 0, 0, 1, 0]$	76	×	✓ ← $\mathcal{O}_{[0,1,0,0,0,0]}^{\{1,2,3,4,5,6\}}$	×
$A_2 + 2A_1$	$[0, 0, 0, 1, 0, 0, 0]$	82	×	?	✓
A_3	$[2, 0, 0, 0, 0, 1, 0]$	84	×	✓ ← $\mathcal{O}_{(2^2, 1^8)}^{\{2,3,4,5,6,7\}}$	×
$2A_2$	$[0, 0, 0, 0, 0, 2, 0]$	84	×	?	×
$A_2 + 3A_1$	$[0, 2, 0, 0, 0, 0, 0]$	84	×	?	×
$(A_3 + A_1)''$	$[2, 0, 0, 0, 0, 0, 2]$	86	×	✓ ← $\mathcal{O}_{(3, 1^9)}^{\{2,3,4,5,6,7\}}$	×
$2A_2 + A_1$	$[0, 0, 1, 0, 0, 1, 0]$	90	×	?	✓
$(A_3 + A_1)'$	$[1, 0, 0, 1, 0, 0, 0]$	92	×	?	✓
$D_4(a_1)$	$[0, 0, 2, 0, 0, 0, 0]$	94	×	✓ ← $\mathcal{O}_{(2^4, 1^4)}^{\{2,3,4,5,6,7\}}$	×
$A_3 + 2A_1$	$[1, 0, 0, 0, 1, 0, 1]$	94	×	?	×
D_4	$[2, 0, 2, 0, 0, 0, 0]$	96	×	✓ ← $\mathcal{O}_{[0,0,0,0,2,0]}^{\{2,3,4,5,6,7\}}$	×
$D_4(a_1) + A_1$	$[0, 1, 1, 0, 0, 0, 1]$	96	×	✓ ← $\mathcal{O}_{[0,0,0,0,0,2]}^{\{2,3,4,5,6,7\}}$	×
$A_3 + A_2$	$[0, 0, 0, 1, 0, 1, 0]$	98	×	✓ ← $\mathcal{O}_{(3, 2^2, 1^5)}^{\{2,3,4,5,6,7\}}$	×
A_4	$[2, 0, 0, 0, 0, 2, 0]$	100	×	✓ ← $\mathcal{O}_{(3^2, 1^6)}^{\{2,3,4,5,6,7\}}$	×
$A_3 + A_2 + A_1$	$[0, 0, 0, 0, 2, 0, 0]$	100	×	?	×
$(A_5)''$	$[2, 0, 0, 0, 0, 2, 2]$	102	×	✓ ← $\mathcal{O}_{(5, 1^7)}^{\{2,3,4,5,6,7\}}$	×

Type E₇ (continued). The characteristics $[0, 2, 0, 0, 2, 0]$ and $[0, 2, 0, 0, 0, 2]$ of nilpotent orbits in D_6 correspond to the very even partition $(4^2, 2^2)$, while the characteristics $[0, 2, 0, 2, 2, 0]$ and $[0, 2, 0, 2, 0, 2]$ correspond to (6^2) .




	\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
$D_4 + A_1$	$[2, 1, 1, 0, 0, 0, 1]$	102	×	?	×
$A_4 + A_1$	$[1, 0, 0, 1, 0, 1, 0]$	104	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,3,4,5,6,7\} \\ (2^2, 1^3)}}$	×
$D_5(a_1)$	$[2, 0, 0, 1, 0, 1, 0]$	106	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{2,3,4,5,6,7\} \\ (3^2, 2^2, 1^2)}}$	×
$A_4 + A_2$	$[0, 0, 0, 2, 0, 0, 0]$	106	×	?	×
$(A_5)'$	$[1, 0, 0, 1, 0, 2, 0]$	108	×	?	×
$A_5 + A_1$	$[1, 0, 0, 1, 0, 1, 2]$	108	×	?	×
$D_5(a_1) + A_1$	$[2, 0, 0, 0, 2, 0, 0]$	108	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,2,3,4,6,7\} \\ (2, 1^3), (1^3)}}$	×
$D_5(a_2)$	$[0, 1, 1, 0, 1, 0, 2]$	110	×	?	×
$E_6(a_3)$	$[0, 0, 2, 0, 0, 2, 0]$	110	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,2,3,5,6,7\} \\ (2, 1), (1^2), (1^4)}}$	×
D_5	$[2, 0, 2, 0, 0, 2, 0]$	112	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{2,3,4,5,6,7\} \\ [0, 2, 0, 0, 2, 0]}}$	×
$E_7(a_5)$	$[0, 0, 0, 2, 0, 0, 2]$	112	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{2,3,4,5,6,7\} \\ [0, 2, 0, 0, 0, 2]}}$	×
A_6	$[0, 0, 0, 2, 0, 2, 0]$	114	×	?	×
$D_5 + A_1$	$[2, 1, 1, 0, 1, 1, 0]$	114	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,2,3,4,6,7\} \\ (3, 1^2), (1^3)}}$	×
$D_6(a_1)$	$[2, 1, 1, 0, 1, 0, 2]$	114	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,2,4,5,6,7\} \\ (2), (3, 1^3)}}$	×
$E_7(a_4)$	$[2, 0, 0, 2, 0, 0, 2]$	116	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,3,4,5,6,7\} \\ (3^2, 1)}}$	×
D_6	$[2, 1, 1, 0, 1, 2, 2]$	118	×	?	×
$E_6(a_1)$	$[2, 0, 0, 2, 0, 2, 0]$	118	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{2,3,4,5,6,7\} \\ (5^2, 1^2)}}$	×
E_6	$[2, 0, 2, 2, 0, 2, 0]$	120	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{2,3,4,5,6,7\} \\ [0, 2, 0, 2, 2, 0]}}$	×
$E_7(a_3)$	$[2, 0, 0, 2, 0, 2, 2]$	120	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{2,3,4,5,6,7\} \\ [0, 2, 0, 2, 0, 2]}}$	×
$E_7(a_2)$	$[2, 2, 2, 0, 2, 0, 2]$	122	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,3,4,5,6,7\} \\ (5, 2)}}$	×
$E_7(a_1)$	$[2, 2, 2, 0, 2, 2, 2]$	124	×	$\checkmark \leftarrow \mathcal{O}_{\substack{\{1,3,4,5,6,7\} \\ (6, 1)}}$	×

Type E₈.




	\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
A_1	$[0, 0, 0, 0, 0, 0, 0, 1]$	58	✓ ← little	✓ ← little	✓
$2A_1$	$[1, 0, 0, 0, 0, 0, 0, 0]$	92	✓ ← little	✓ ← little	✓
$3A_1$	$[0, 0, 0, 0, 0, 0, 1, 0]$	112	✓ ← little	✓ ← little	✓
A_2	$[0, 0, 0, 0, 0, 0, 0, 2]$	114	✓ ← little	✓ ← little	×
$4A_1$	$[0, 1, 0, 0, 0, 0, 0, 0]$	128	×	?	✓
$A_2 + A_1$	$[1, 0, 0, 0, 0, 0, 0, 1]$	136	×	?	✓
$A_2 + 2A_1$	$[0, 0, 0, 0, 0, 1, 0, 0]$	146	×	?	✓
A_3	$[1, 0, 0, 0, 0, 0, 0, 2]$	148	×	✓ ← $\mathcal{O}_{[1,0,0,0,0,0,0]}^{\{1,2,3,4,5,6,7\}}$	×
$A_2 + 3A_1$	$[0, 0, 1, 0, 0, 0, 0, 0]$	154	×	?	✓
$2A_2$	$[2, 0, 0, 0, 0, 0, 0, 0]$	156	×	?	×
$2A_2 + A_1$	$[1, 0, 0, 0, 0, 0, 1, 0]$	162	×	?	✓
$A_3 + A_1$	$[0, 0, 0, 0, 0, 1, 0, 1]$	164	×	?	✓
$D_4(a_1)$	$[0, 0, 0, 0, 0, 0, 2, 0]$	166	×	✓ ← $\mathcal{O}_{[0,0,0,0,0,1,0]}^{\{1,2,3,4,5,6,7\}}$	×
D_4	$[0, 0, 0, 0, 0, 0, 2, 2]$	168	×	✓ ← $\mathcal{O}_{[0,0,0,0,0,0,2]}^{\{1,2,3,4,5,6,7\}}$	×
$2A_2 + 2A_1$	$[0, 0, 0, 0, 1, 0, 0, 0]$	168	×	?	✓
$A_3 + 2A_1$	$[0, 0, 1, 0, 0, 0, 0, 1]$	172	×	?	✓
$D_4(a_1) + A_1$	$[0, 1, 0, 0, 0, 0, 1, 0]$	176	×	?	✓
$A_3 + A_2$	$[1, 0, 0, 0, 0, 1, 0, 0]$	178	×	✓ ← $\mathcal{O}_{(2^2, 1^{10})}^{\{2,3,4,5,6,7,8\}}$	×
A_4	$[2, 0, 0, 0, 0, 0, 0, 2]$	180	×	✓ ← $\mathcal{O}_{(3, 1^{11})}^{\{2,3,4,5,6,7,8\}}$	×
$A_3 + A_2 + A_1$	$[0, 0, 0, 1, 0, 0, 0, 0]$	182	×	?	✓
$D_4 + A_1$	$[0, 1, 0, 0, 0, 0, 1, 2]$	184	×	?	×
$D_4(a_1) + A_2$	$[0, 2, 0, 0, 0, 0, 0, 0]$	184	×	?	×
$A_4 + A_1$	$[1, 0, 0, 0, 0, 1, 0, 1]$	188	×	✓ ← $\mathcal{O}_{[0,1,0,0,0,0]}^{\{1,2,3,4,5,6,8\}, (1^2)}$	×

Type E_8 (continued).



	\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
$2A_3$	$[1, 0, 0, 0, 1, 0, 0, 0]$	188	×	?	✓
$D_5(a_1)$	$[1, 0, 0, 0, 0, 1, 0, 2]$	190	×	✓ $\leftarrow \mathcal{O}_{(2^4, 1^6)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
$A_4 + 2A_1$	$[0, 0, 0, 1, 0, 0, 0, 1]$	192	×	?	×
$A_4 + A_2$	$[0, 0, 0, 0, 0, 2, 0, 0]$	194	×	?	×
A_5	$[2, 0, 0, 0, 0, 1, 0, 1]$	196	×	✓ $\leftarrow \mathcal{O}_{(3, 2^2, 1^7)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
$D_5(a_1) + A_1$	$[0, 0, 0, 1, 0, 0, 0, 2]$	196	×	?	×
$A_4 + A_2 + A_1$	$[0, 0, 1, 0, 0, 1, 0, 0]$	196	×	?	×
$D_4 + A_2$	$[0, 2, 0, 0, 0, 0, 0, 2]$	198	×	✓ $\leftarrow \mathcal{O}_{(2^6, 1^2)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
$E_6(a_3)$	$[2, 0, 0, 0, 0, 0, 2, 0]$	198	×	✓ $\leftarrow \mathcal{O}_{(3^2, 1^8)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
D_5	$[2, 0, 0, 0, 0, 0, 2, 2]$	200	×	✓ $\leftarrow \mathcal{O}_{(5, 1^9)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
$A_4 + A_3$	$[0, 0, 0, 1, 0, 0, 1, 0]$	200	×	?	✓
$A_5 + A_1$	$[1, 0, 0, 1, 0, 0, 0, 1]$	202	×	?	✓
$D_5(a_1) + A_2$	$[0, 0, 1, 0, 0, 1, 0, 1]$	202	×	?	✓
$D_6(a_2)$	$[0, 1, 1, 0, 0, 0, 1, 0]$	204	×	?	×
$E_6(a_3) + A_1$	$[1, 0, 0, 0, 1, 0, 1, 0]$	204	×	?	×
$E_7(a_5)$	$[0, 0, 0, 1, 0, 1, 0, 0]$	206	×	?	×
$D_5 + A_1$	$[1, 0, 0, 0, 1, 0, 1, 2]$	208	×	?	×
$E_8(a_7)$	$[0, 0, 0, 0, 2, 0, 0, 0]$	208	×	✓ $\leftarrow \mathcal{O}_{(3^2, 2^2, 1^4)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
A_6	$[2, 0, 0, 0, 0, 2, 0, 0]$	210	×	✓ $\leftarrow \mathcal{O}_{(3, 1^7), (1^3)}^{[1, 2, 3, 4, 5, 7, 8]}$	×
$D_6(a_1)$	$[0, 1, 1, 0, 0, 0, 1, 2]$	210	×	✓ $\leftarrow \mathcal{O}_{(3^2, 2^4)}^{[2, 3, 4, 5, 6, 7, 8]}$	×
$A_6 + A_1$	$[1, 0, 0, 1, 0, 1, 0, 0]$	212	×	?	×
$E_7(a_4)$	$[0, 0, 0, 1, 0, 1, 0, 2]$	212	×	✓ $\leftarrow \mathcal{O}_{[0, 0, 0, 1, 0, 1, 0]}^{[1, 2, 3, 4, 5, 6, 7]}$	×
$E_6(a_1)$	$[2, 0, 0, 0, 0, 2, 0, 2]$	214	×	✓ $\leftarrow \mathcal{O}_{(5, 3, 1^6)}^{[2, 3, 4, 5, 6, 7, 8]}$	×

Type E₈ (continued).



	\mathcal{O}	$\dim \mathcal{O}$	RC_1	RC_2	rigid
$D_5 + A_2$	$[0, 0, 0, 0, 2, 0, 0, 2]$	214	×	$\checkmark \leftarrow \mathcal{O}_{(2^3, 1^2)}^{\{1, 3, 4, 5, 6, 7, 8\}}$	×
D_6	$[2, 1, 1, 0, 0, 0, 1, 2]$	216	×	?	×
E_6	$[2, 0, 0, 0, 0, 2, 2, 2]$	216	×	$\checkmark \leftarrow \mathcal{O}_{(7, 1^7)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$D_7(a_2)$	$[1, 0, 0, 1, 0, 1, 0, 1]$	216	×	$\checkmark \leftarrow \mathcal{O}_{(1^2), (2^2, 1^3)}^{\{1, 2, 4, 5, 6, 7, 8\}}$	×
A_7	$[1, 0, 0, 1, 0, 1, 1, 0]$	218	×	?	×
$E_6(a_1) + A_1$	$[1, 0, 0, 1, 0, 1, 0, 2]$	218	×	$\checkmark \leftarrow \mathcal{O}_{(4^2, 2^2, 1^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_7(a_3)$	$[2, 0, 0, 1, 0, 1, 0, 2]$	220	×	$\checkmark \leftarrow \mathcal{O}_{(5, 3, 2^2, 1)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_8(b_6)$	$[0, 0, 0, 2, 0, 0, 0, 2]$	220	×	$\checkmark \leftarrow \mathcal{O}_{(4^3, 3, 1^3)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$D_7(a_1)$	$[2, 0, 0, 0, 2, 0, 0, 2]$	222	×	$\checkmark \leftarrow \mathcal{O}_{(4^2, 3^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_6 + A_1$	$[1, 0, 0, 1, 0, 1, 2, 2]$	222	×	?	×
$E_7(a_2)$	$[0, 1, 1, 0, 1, 0, 2, 2]$	224	×	?	×
$E_8(a_6)$	$[0, 0, 0, 2, 0, 0, 2, 0]$	224	×	$\checkmark \leftarrow \mathcal{O}_{(5, 3^3)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
D_7	$[2, 1, 1, 0, 1, 1, 0, 1]$	226	×	?	×
$E_8(b_5)$	$[0, 0, 0, 2, 0, 0, 2, 2]$	226	×	$\checkmark \leftarrow \mathcal{O}_{(5^2, 2^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_7(a_1)$	$[2, 1, 1, 0, 1, 0, 2, 2]$	228	×	$\checkmark \leftarrow \mathcal{O}_{(7, 3, 2^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_8(a_5)$	$[2, 0, 0, 2, 0, 0, 2, 0]$	228	×	$\checkmark \leftarrow \mathcal{O}_{(5^2, 3, 1)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_8(b_4)$	$[2, 0, 0, 2, 0, 0, 2, 2]$	230	×	$\checkmark \leftarrow \mathcal{O}_{(6^2, 1^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
E_7	$[2, 1, 1, 0, 1, 2, 2, 2]$	232	×	?	×
$E_8(a_4)$	$[2, 0, 0, 2, 0, 2, 0, 2]$	232	×	$\checkmark \leftarrow \mathcal{O}_{(7, 5, 1^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_8(a_3)$	$[2, 0, 0, 2, 0, 2, 2, 2]$	234	×	$\checkmark \leftarrow \mathcal{O}_{(7^2)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_8(a_2)$	$[2, 2, 2, 0, 2, 0, 2, 2]$	236	×	$\checkmark \leftarrow \mathcal{O}_{(9, 5)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×
$E_8(a_1)$	$[2, 2, 2, 0, 2, 2, 2, 2]$	238	×	$\checkmark \leftarrow \mathcal{O}_{(11, 3)}^{\{2, 3, 4, 5, 6, 7, 8\}}$	×

checked using Propositions 3.6, 4.2, 4.6, and Remark 3.9. When condition RC_1 is obtained via Proposition 4.6, we give an example of the bigger simple Lie algebra and the little nilpotent orbit satisfying condition (iii) of Proposition 4.6 from which it is obtained.

As for determining whether condition $RC_2(m)$ is verified, our main method is to list the orbits induced by nilpotent orbits that have a little factor (Theorem 6.1). Thus they are $RC_2(m)$ for all $m \in \mathbb{N}^*$. Since induction is transitive, we can proceed by induction on the rank of the Lie algebra, where at each step, we only need to consider induction from orbits in maximal Levi subalgebras which are themselves induced from nilpotent orbits with a little factor. For an orbit verifying condition $RC_2(m)$, we give an example of a maximal Levi subalgebra \mathfrak{l} and an orbit in \mathfrak{l} induced from a nilpotent orbit with a little factor.

In both cases, if the orbit is little, then we just label it little. The subscript of an orbit indicates: its characteristics, the associated partition, or its Bala-Carter label. If a superscript of an orbit is present, it indicates the corresponding maximal Levi subalgebra.

We have omitted the zero orbit and the regular orbit because they are neither RC_1 nor $RC_2(m)$.

All the computations are done using the package `sla` of GAP4.

Acknowledgments

We thank Michel Brion for suggesting us this topic, and bringing to our attention Namikawa's result. We are also indebted to *Mathematisches Forschungsinstitut Oberwolfach* (MFO) for its hospitality during our stay as "Research in pairs" in October, 2014.

We are grateful to Michaël Bulois for pointing out an error in a previous version, and to Jean-Yves Charbonnel and Michel Brion for their comments.

We are also thankful to the referee for careful reading and thoughtful suggestions.

This research is supported by the ANR Project GERCHER Grant number ANR-2010-BLAN-110-02.

References

- [Brion and Fu 2015] M. Brion and B. Fu, "Symplectic resolutions for conical symplectic varieties", *Int. Math. Res. Not.* **2015**:12 (2015), 4335–4343. [MR 3356756](#) [Zbl 06471154](#)
- [Broer 1998] A. Broer, "Normal nilpotent varieties in F_4 ", *J. Algebra* **207**:2 (1998), 427–448. [MR 99e:14056](#) [Zbl 0918.17003](#)
- [Collingwood and McGovern 1993] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold, New York, 1993. [MR 94j:17001](#) [Zbl 0972.17008](#)
- [Ein and Mustață 2009] L. Ein and M. Mustață, "Jet schemes and singularities", pp. 505–546 in *Algebraic geometry, part 2* (Seattle, 2005), edited by D. Abramovich et al., Proceedings of

- Symposia in Pure Mathematics **80**:2, Amer. Math. Soc., Providence, RI, 2009. [MR 2010h:14004](#)
[Zbl 1181.14019](#)
- [de Fernex et al. 2013] T. de Fernex, L. Ein, and M. Mustață, “Vanishing theorems and singularities in birational geometry”, book in preparation, 2013, <http://www.math.utah.edu/~defernex/book.pdf>.
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, New York, 1977. [MR 57 #3116](#) [Zbl 0367.14001](#)
- [Henderson 2014] A. Henderson, “Singularities of nilpotent orbit closures”, preprint, 2014. [arXiv 1408.3888v1](#)
- [Hesselink 1976] W. H. Hesselink, “Cohomology and the resolution of the nilpotent variety”, *Math. Ann.* **223**:3 (1976), 249–252. [MR 54 #5253](#) [Zbl 0318.14007](#)
- [Hinich 1991] V. Hinich, “On the singularities of nilpotent orbits”, *Israel J. Math.* **73**:3 (1991), 297–308. [MR 92m:14005](#) [Zbl 0766.14038](#)
- [Ishii 2011] S. Ishii, “Geometric properties of jet schemes”, *Comm. Algebra* **39**:5 (2011), 1872–1882. [MR 2012g:14017](#) [Zbl 1230.14052](#)
- [Kaledin 2006] D. Kaledin, “Symplectic singularities from the Poisson point of view”, *J. Reine Angew. Math.* **600** (2006), 135–156. [MR 2007j:32030](#) [Zbl 1121.53056](#)
- [Kempken 1983] G. Kempken, “Induced conjugacy classes in classical Lie algebras”, *Abh. Math. Sem. Univ. Hamburg* **53** (1983), 53–83. [MR 85i:17009](#) [Zbl 0495.17003](#)
- [Kolchin 1973] E. R. Kolchin, *Differential algebra and algebraic groups*, Pure and Applied Mathematics **54**, Academic Press, New York, 1973. [MR 58 #27929](#) [Zbl 0264.12102](#)
- [Kostant 1963] B. Kostant, “Lie group representations on polynomial rings”, *Amer. J. Math.* **85** (1963), 327–404. [MR 28 #1252](#) [Zbl 0124.26802](#)
- [Kraft 1989] H. Kraft, “Closures of conjugacy classes in G_2 ”, *J. Algebra* **126**:2 (1989), 454–465. [MR 91a:17020](#) [Zbl 0693.17004](#)
- [Kraft and Procesi 1982] H. Kraft and C. Procesi, “On the geometry of conjugacy classes in classical groups”, *Comment. Math. Helv.* **57**:4 (1982), 539–602. [MR 85b:14065](#) [Zbl 0511.14023](#)
- [Levasseur and Smith 1988] T. Levasseur and S. P. Smith, “Primitive ideals and nilpotent orbits in type G_2 ”, *J. Algebra* **114**:1 (1988), 81–105. [MR 89f:17013](#) [Zbl 0644.17005](#)
- [Lusztig and Spaltenstein 1979] G. Lusztig and N. Spaltenstein, “Induced unipotent classes”, *J. London Math. Soc. (2)* **19**:1 (1979), 41–52. [MR 82g:20070](#) [Zbl 0407.20035](#)
- [Mumford et al. 1994] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, 3rd ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete (2)* **34**, Springer, Berlin, 1994. [MR 95m:14012](#)
[Zbl 0797.14004](#)
- [Mustață 2001] M. Mustață, “Jet schemes of locally complete intersection canonical singularities”, *Invent. Math.* **145**:3 (2001), 397–424. [MR 2002f:14005](#) [Zbl 1091.14004](#)
- [Namikawa 2013] Y. Namikawa, “On the structure of homogeneous symplectic varieties of complete intersection”, *Invent. Math.* **193**:1 (2013), 159–185. [MR 3069115](#) [Zbl 1285.14051](#)
- [Nash 1995] J. F. Nash, Jr., “Arc structure of singularities”, *Duke Math. J.* **81**:1 (1995), 31–38. [MR 98f:14011](#) [Zbl 0880.14010](#)
- [Panyushev 1991] D. I. Panyushev, “Рациональность особенностей и Горенштейновость нильпотентных орбит”, *Funktional. Anal. i Prilozhen.* **25**:3 (1991), 76–78. Translated as “Rationality of singularities and the Gorenstein property of nilpotent orbits” in *Funct. Anal. Appl.* **25**:3 (1991), 225–226. [MR 92i:14047](#) [Zbl 0749.14030](#)
- [Raïs and Tauvel 1992] M. Raïs and P. Tauvel, “Indice et polynômes invariants pour certaines algèbres de Lie”, *J. Reine Angew. Math.* **425** (1992), 123–140. [MR 93g:17012](#) [Zbl 0743.17008](#)

- [Richardson 1974] R. W. Richardson, Jr., “Conjugacy classes in parabolic subgroups of semisimple algebraic groups”, *Bull. London Math. Soc.* **6** (1974), 21–24. MR 48 #8648 Zbl 0287.20036
- [Sommers 2003] E. Sommers, “Normality of nilpotent varieties in E_6 ”, *J. Algebra* **270**:1 (2003), 288–306. MR 2004i:20085 Zbl 1041.22009
- [Tauvel and Yu 2005] P. Tauvel and R. W. T. Yu, *Lie algebras and algebraic groups*, Springer, Berlin, 2005. MR 2006c:17001 Zbl 1068.17001
- [Weyman 1989] J. Weyman, “The equations of conjugacy classes of nilpotent matrices”, *Invent. Math.* **98**:2 (1989), 229–245. MR 91g:20070 Zbl 0717.20033
- [Weyman 2002] J. Weyman, “Two results on equations of nilpotent orbits”, *J. Algebraic Geom.* **11**:4 (2002), 791–800. MR 2003k:20071 Zbl 1009.20052
- [Weyman 2003] J. Weyman, *Cohomology of vector bundles and syzygies*, Cambridge Tracts in Mathematics **149**, Cambridge University Press, 2003. MR 2004d:13020 Zbl 1075.13007
- [Yuen 2007] C. Yuen, “Jet schemes of determinantal varieties”, pp. 261–270 in *Algebra, geometry and their interactions*, edited by A. Corso et al., Contemporary Mathematics **448**, Amer. Math. Soc., Providence, RI, 2007. MR 2009b:14093 Zbl 1142.14032

Received April 21, 2015. Revised August 3, 2015.

ANNE MOREAU
LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS
UNIVERSITE DE POITIERS
TEÉLÉPORT 2 - BP 30179
BOULEVARD MARIE ET PIERRE CURIE
86962 FUTUROSCOPE (POITIERS) CHASSENEUIL CEDEX
FRANCE
anne.moreau@math.univ-poitiers.fr

RUPERT WEI TZE YU
LABORATOIRE DE MATHÉMATIQUES DE REIMS (LMR) - EA 4535
UNIVERSITÉ DE REIMS CHAMPAGNE ARDENNE
U.F.R. SCIENCES EXACTES ET NATURELLES
MOULIN DE LA HOUSSE - BP 1039
51687 REIMS CEDEX 2
FRANCE
rupert.yu@univ-reims.fr

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 281 No. 1 March 2016

Compatible systems of symplectic Galois representations and the inverse Galois problem II: Transvections and huge image	1
SARA ARIAS-DE-REYNA, LUIS DIEULEFAIT and GABOR WIESE	
On the number of lines in the limit set for discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$	17
WALDEMAR BARRERA, ANGEL CANO and JUÁN NAVARRETE	
Galois theory, functional Lindemann–Weierstrass, and Manin maps	51
DANIEL BERTRAND and ANAND PILLAY	
Morse area and Scharlemann–Thompson width for hyperbolic 3-manifolds	83
DIANE HOFFOSS and JOSEPH MAHER	
Ricci tensor of real hypersurfaces	103
MAYUKO KON	
Monotonicity formulae and vanishing theorems	125
JINTANG LI	
Jet schemes of the closure of nilpotent orbits	137
ANNE MOREAU and RUPERT WEI TZE YU	
Components of spaces of curves with constrained curvature on flat surfaces	185
NICOLAU C. SALDANHA and PEDRO ZÜHLKE	
A note on minimal graphs over certain unbounded domains of Hadamard manifolds	243
MIRIAM TELICHEVESKY	