

*Pacific
Journal of
Mathematics*

**CARTAN-FUBINI TYPE RIGIDITY OF DOUBLE COVERING
MORPHISMS OF QUADRATIC MANIFOLDS**

HOSUNG KIM

Volume 282 No. 2

June 2016

CARTAN–FUBINI TYPE RIGIDITY OF DOUBLE COVERING MORPHISMS OF QUADRATIC MANIFOLDS

HOSUNG KIM

Let $Z \subset \mathbb{P}^N$ be a Fano manifold whose Picard group is generated by the hyperplane section class. Assume that Z is covered by lines and $i(Z) \geq 3$. Let $\phi : X^Z \rightarrow Z$ be a double cover, branched along a smooth hypersurface section of degree $2m$, $1 \leq m \leq i(Z) - 2$. We describe the defining ideal of the variety of minimal rational tangents at a general point. As an application, we show that if $Z \subset \mathbb{P}^N$ is defined by quadratic equations and $2 \leq m \leq i(Z) - 2$, then the morphism ϕ satisfies the Cartan–Fubini type rigidity property.

1. Introduction

Throughout the paper, we will work over the field of complex numbers. Let X be a Fano manifold of Picard number 1. The index of X is the integer $i(X)$ such that $-K_X = i(X)L$ where L is the ample generator of the Picard group of X . For a general point $x \in X$, a rational curve through x is called a *minimal rational curve* if it has minimal K_X^{-1} -degree among all rational curves through x . Denote by \mathcal{K}_x the normalized space of minimal rational curves through x . It is known (e.g., [Kollár 1996, II.3.11.5]) that \mathcal{K}_x is a disjoint union of finitely many nonsingular projective varieties of dimension $i(X) - 2$. The rational morphism $\mathcal{K}_x \dashrightarrow \mathbb{P}T_x(X)$, sending a member of \mathcal{K}_x which is smooth at x to its tangent direction, can be extended to a birational morphism $\tau_x : \mathcal{K}_x \rightarrow \mathbb{P}T_x(X)$; see [Hwang and Mok 2004; Kebekus 2002]. We denote the image of τ_x by \mathcal{C}_x and call it the *variety of minimal rational tangents (VMRT)* at x . The projective geometry of $\mathcal{C}_x \subset \mathbb{P}T_x(X)$ helps us to understand the geometry of X . This is the motivation for the study of the VMRT for various examples of X . For example, let $\phi : X \rightarrow \mathbb{P}^n$ be a double cover branched on a smooth hypersurface of degree $2m$, $2 \leq m \leq n - 1$. Then for a general point $x \in X$, the VMRT $\mathcal{C}_x \subset \mathbb{P}T_x(X)$ is a complete intersection of multidegree $(m + 1, \dots, 2m)$, and this description implies a certain rigidity property of ϕ [Hwang and Kim 2013].

This work was supported by the National Institute for Mathematical Sciences (NIMS) grant funded by the Korea government (No. C21501).

MSC2010: 14J45.

Keywords: double covers of Fano manifolds, varieties of minimal rational tangents, Cartan–Fubini type rigidity.

Let $Z \subset \mathbb{P}^N$ be a Fano manifold whose Picard group is generated by the hyperplane section class. For each point $y \in Z$, we denote by $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$ the space of tangent directions of lines in Z through y . We say that $Z \subset \mathbb{P}^N$ is covered by lines if $\mathcal{L}_y(Z)$ is nonempty for each $y \in Z$. If $Z \subset \mathbb{P}^N$ is covered by lines, then minimal rational curves on Z are lines in \mathbb{P}^N contained in Z , and for general $y \in Z$, $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$ coincides with the VMRT $\mathcal{C}_y \subset \mathbb{P}T_y(Z)$, which is smooth of dimension $i(Z) - 2$; see [Hwang 2001, Proposition 1.5].

Our first result is the following theorem.

Theorem 1.1. *Let $Z \subset \mathbb{P}^N$ be a Fano manifold whose Picard group is generated by the hyperplane section class. Assume that Z is covered by lines and $i(Z) \geq 3$. Let $Y \subset \mathbb{P}^N$ be a hypersurface of degree $2m$, $1 \leq m \leq i(Z) - 2$, with smooth intersection $Y \cap Z$. Let $\phi : X^Z \rightarrow Z$ be a double cover branched along $Y \cap Z$. Then for a general point $x \in X^Z$, the VMRT \mathcal{C}_x is smooth of dimension $i(Z) - m - 2$ and the differential $d\phi_x : \mathbb{P}T_x(X^Z) \rightarrow \mathbb{P}T_{\phi(x)}(Z)$ sends the VMRT $\mathcal{C}_x \subset \mathbb{P}T_x(X^Z)$ isomorphically to an intersection of $\mathcal{L}_{\phi(x)}(Z) \subset \mathbb{P}T_{\phi(x)}(Z)$ and m hypersurfaces in $\mathbb{P}T_{\phi(x)}(Z)$ of degrees $m + 1, \dots, 2m$ respectively.*

In order to prove the above theorem, we first show that for a certain choice of Y , the statements in Theorem 1.1 hold by identifying minimal rational curves on X^Z with ECO (even contact order) lines with respect to Y (see Definition 2.4) contained in Z . For arbitrary Y , we use a flatness argument.

We are going to show an application of Theorem 1.1. First, let us introduce the definition of Cartan–Fubini type rigidity (CF-rigidity) which was initially defined by Jun-Muk Hwang.

Definition 1.2 (Cartan–Fubini type rigidity). Let X_1 and X_2 be Fano manifolds of Picard number 1 such that $2 \leq \dim X_1 \leq \dim X_2$, and for general $x_1 \in X_1$ and $x_2 \in X_2$, $0 \leq \dim \mathcal{K}_{x_1} \leq \dim \mathcal{K}_{x_2}$. We say that a morphism $\phi : X_1 \rightarrow X_2$ is *CF-rigid* if for any connected open subset (in classical topology) U of X_1 and any biholomorphic immersion $\psi : U \rightarrow X_2$ such that for any member C of \mathcal{K}_x , $x \in U$, $\psi(C \cap U)$ is contained in a minimal rational curve of X_2 , then there exists $\Gamma \in \text{Aut}(X_2)$ such that $\psi = \Gamma \circ \phi|_U$.

The next theorem is on the CF-rigidity of the identity morphism, which was essentially proved in [Hwang and Mok 2001].

Theorem 1.3 (Cartan–Fubini type extension theorem). *Let X be a Fano manifold of Picard number 1 and suppose that $\dim \mathcal{K}_x \geq 1$ for general $x \in X$. Then the identity morphism on X is CF-rigid, i.e., for any connected open subset (in the standard topology) U of X and any biholomorphic immersion $\psi : U \rightarrow X$ such that for any member C of \mathcal{K}_x , $x \in U$, $\psi(C \cap U)$ is contained in a minimal rational curve of X , then there exists $\Gamma \in \text{Aut}(X)$ such that $\psi = \Gamma|_U$.*

Let X be a Fano manifold of Picard number 1 and let $\phi : X \rightarrow \mathbb{P}^n$ is a surjective holomorphic map sending minimal rational curves on X to lines in \mathbb{P}^n . [Hwang and Kim 2013, Theorem 5.4] says that if the VMRT $\mathcal{C}_x \subset \mathbb{P}T_x(X)$ is not contained in a hyperquadric, then $\phi : X \rightarrow \mathbb{P}^n$ is CF-rigid. For example, the double covering morphism $\phi : X \rightarrow \mathbb{P}^n$ branched along a smooth hypersurface of degree $2m$, with $2 \leq m \leq n - 1$, is CF-rigid.

Next is an application of Theorem 1.1 on CF-rigidity.

Theorem 1.4. *In the setting of Theorem 1.1, assume that $Z \subset \mathbb{P}^N$ is a quadratic manifold (i.e., scheme theoretically defined by quadratic equations) with $i(Z) \geq 4$, and $2 \leq m \leq i(Z) - 2$. Then $\phi : X^Z \rightarrow Z$ is CF-rigid. In other words, for any connected open subset (in classical topology) $U \subset X^Z$ and any biholomorphic immersion $\psi : U \rightarrow Z$ such that for any member C of \mathcal{K}_x , $x \in U$, the image $\psi(C \cap U) \subset Z$ is contained in a line in \mathbb{P}^N , there exists $\Gamma \in \text{Aut}(Z)$ such that $\psi = \Gamma \circ \phi|_U$.*

The key point is that for $Z \subset \mathbb{P}^N$ in Theorem 1.4, its VMRT $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$ is also a quadratic manifold for a general point $y \in Z$; see [Ionescu and Russo 2013, Theorem 2.4]. Using this observation and Theorem 1.3, we shall prove Theorem 1.4.

The organization of this paper is as follows. In Section 2, we will review some basic facts on ECO lines. In Section 3, we will study minimal rational curves on double covers of certain Fano manifolds covered by lines. Theorem 1.1 will be proved in Section 4. In the final section, we will prove Theorem 1.4 and present its applications.

2. ECO lines

The aim of this section is to give a brief review of basic facts on ECO lines which will be used in the proof of Theorem 1.1. For more details, see [Hwang and Kim 2013].

Definition 2.1. A homogeneous polynomial of degree $2m$, $m \geq 1$, in the polynomial ring $\mathbb{C}[s, t]$ with two variables s and t is an *ECO (even contact order) polynomial* if it can be written as the square of a homogeneous polynomial of degree m in $\mathbb{C}[s, t]$.

Proposition 2.2 [op. cit., Proposition 3.3]. *For each $m > 0$, there exist m unique polynomials in the variables t_1, \dots, t_m ,*

$$A_k(t_1, \dots, t_m) \in \mathbb{C}[t_1, \dots, t_m], \quad m + 1 \leq k \leq 2m$$

with the following properties:

- (i) $A_k(t_1, \dots, t_m)$ is weighted homogeneous of degree k with respect to $\text{wt}(t_i) = i$ for each $i = 1, \dots, m$;

(ii) *the polynomial in two variables (s, t)*

$$s^{2d} + a_1 s^{2d-1} t + \cdots + a_{2d-1} s t^{2d-1} + a_{2d} t^{2d}$$

is an ECO polynomial if and only if $a_k = A_k(a_1, \dots, a_d)$ for each $d+1 \leq k \leq 2d$. In particular, the polynomial

$$s^{2d} + a_{d+1} s^{d-1} t^{d+1} + \cdots + a_{2d-1} s t^{2d-1} + a_{2d} t^{2d}$$

is an ECO polynomial if and only if $a_{d+1} = \cdots = a_{2d} = 0$.

Definition 2.3. Let $f(t_0, \dots, t_N)$ be a homogeneous polynomial of degree $2m$ in variables t_0, \dots, t_N . Write

$$f(1, y_1 + \lambda z_1, \dots, y_N + \lambda z_N) = a_0^f(y; z) + a_1^f(y; z) \lambda + \cdots + a_{2m}^f(y; z) \lambda^{2m},$$

where each $a_k^f(y; z) = a_k^f(y_1, \dots, y_N; z_1, \dots, z_N)$ is a polynomial in $2N$ variables $y_1, \dots, y_N, z_1, \dots, z_N$. Let A_k be as in Proposition 2.2 and set

$$B_k^f(y; z) := \frac{a_k^f(y; z)}{a_0^f(y; z)} - A_k \left(\frac{a_1^f(y; z)}{a_0^f(y; z)}, \dots, \frac{a_m^f(y; z)}{a_0^f(y; z)} \right).$$

We remark that for a fixed y , $a_k^f(y; z)$ is a homogeneous polynomial in variables z_1, \dots, z_N of degree k . Furthermore for a fixed y with

$$a_0^f(y; z) = f(1, y_1, \dots, y_N) \neq 0,$$

each $B_k^f(y; z)$ is a homogeneous polynomial of degree k in variables z_1, \dots, z_N .

Definition 2.4. Let $Y \subset \mathbb{P}^N$ be a hypersurface of even degree $2m$. A line $\ell \subset \mathbb{P}^N$ is called an *ECO (even contact order) line* with respect to Y if $\ell \not\subset Y$ and the local intersection number at each point of $\ell \cap Y$ is even. For each point $y \in \mathbb{P}^N \setminus Y$, we denote by $\mathcal{E}_y^Y \subset \mathbb{P}T_y(\mathbb{P}^N)$ the space of tangent directions of ECO lines with respect to Y passing through y .

Proposition 2.5 [Hwang and Kim 2013, Proposition 3.8]. *Choose a homogeneous coordinate system t_0, \dots, t_N on \mathbb{P}^N . We denote by $\mathbb{P}_\infty^{N-1} \subset \mathbb{P}^N$ the hyperplane defined by $t_0 = 0$ and choose a homogeneous coordinate system z_1, \dots, z_N on \mathbb{P}_∞^{N-1} given by the restrictions of t_1, \dots, t_N respectively. Let $f(t_0, \dots, t_N)$ be a homogeneous polynomial of degree $2m$ and let $Y \subset \mathbb{P}^N$ be its associated hypersurface. For each point $y = [1 : y_1 : \cdots : y_N] \in \mathbb{P}^N \setminus Y \cup \mathbb{P}_\infty^{N-1}$, define the projective isomorphism*

$$v_y : \mathbb{P}_\infty^{N-1} \rightarrow \mathbb{P}T_y(\mathbb{P}^N)$$

by sending $[z_1 : \cdots : z_N] \in \mathbb{P}_\infty^{N-1}$ to the tangent direction of the line

$$\{(y_1 + \lambda z_1, \dots, y_N + \lambda z_N) \mid \lambda \in \mathbb{C}\}$$

at the point y . Then the variety $v_y^{-1}(\mathcal{E}_y^Y) \subset \mathbb{P}_\infty^{N-1}$ is set-theoretically the intersection of m hypersurfaces in \mathbb{P}_∞^{N-1} defined by polynomials in $\{B_k^f(y; z) \mid m + 1 \leq k \leq 2m\}$.

3. Minimal rational curves on double covers of prime Fano manifolds

Definition 3.1. Let $Z \subset \mathbb{P}^N$ be a Fano manifold whose Picard group is generated by the hyperplane section class H . Assume that Z is covered by lines and $i(Z) \geq 3$. Let $Y \subset \mathbb{P}^N$ be a hypersurface of degree $2m$, $1 \leq m \leq i(Z) - 2$, defining smooth hypersurface section $B := Y \cap Z \subset Z$. Let

$$\phi : X^Z \rightarrow Z$$

be a double cover branched along B . From the adjunction formula

$$K_{X^Z} = \phi^*(K_Z + \frac{1}{2}(B)) = \phi^*((-i(Z) + m)H),$$

X^Z is a Fano manifold of index $i(X^Z) = i(Z) - m$ and its Picard group is generated by $\phi^*(H)$.

Proposition 3.2. *In the setting of Definition 3.1, an irreducible reduced curve $C \subset X^Z$ with $\phi(C) \not\subset B$ is a minimal rational curve if and only if its image curve $\phi(C)$ is an ECO line with respect to Y . Moreover for any minimal rational curve $C \subset X^Z$ with $\phi(C) \not\subset B$, $\phi|_C : C \rightarrow \phi(C)$ is an isomorphism.*

Proof. We first observe:

Claim: An irreducible reduced curve $C \subset X^Z$ with $\phi(C) \not\subset B$ has ϕ^*H -degree 1 if and only if its image curve $\phi(C) \subset Z$ is an ECO line with respect to Y . Moreover for any ϕ^*H -degree 1 curve $C \subset X^Z$ with $\phi(C) \not\subset B$, $\phi|_C : C \rightarrow \phi(C)$ is an isomorphism.

Proof of the claim. Let $C \subset X^Z$ be an irreducible reduced curve such that the image $\phi(C)$ is an ECO line with respect to Y . Suppose that $\phi|_C : C \rightarrow \phi(C)$ is not birational. For a point $z \in \phi(C) \cap Y$, let t be a local uniformizing parameter on $\phi(C)$ at z and let r_z be the local intersection number of $\phi(C)$ and Y at z . Then C is analytically defined by the equation $s^2 = t^{r_z}$. Since r_z is even for any choice of $z \in \phi(C) \cap Y$, the composition of the normalization morphism $\tilde{C} \rightarrow C$ and the covering morphism $\phi|_C : C \rightarrow \phi(C)$ induces a morphism $\tilde{C} \rightarrow \phi(C)$ of degree 2 without ramification point, a contradiction. Thus $\phi|_C : C \rightarrow \phi(C)$ is birational and C has ϕ^*H -degree 1.

Conversely, if C is an irreducible reduced curve of ϕ^*H -degree 1, then $\phi(C) \subset Z$ with $\phi(C) \not\subset B$ and $\phi|_C : C \rightarrow \phi(C)$ must be birational. Thus $\phi^{-1}(\phi(C))$ has an irreducible component C' different from C with $\phi(C \cap C') = \phi(C) \cap B$. By the same argument as before, if the local intersection number r_z at $z \in \phi(C) \cap Y$ is odd, the germ of $\phi^{-1}(\phi(C))$ over z , defined by $s^2 = t^{r_z}$, is irreducible, a contradiction. Thus

r_z is even for all $z \in \phi(C) \cap Y$ which implies that $\phi(C)$ is an ECO line. Moreover, C must be smooth and the morphism $\phi|_C : C \rightarrow \phi(C)$ is an isomorphism. \square

Let us go back to the proof of our proposition. By the claim above, we only need to show that for a general point $x \in X^Z$, there exists a ϕ^*H -degree 1 curve through x . From $\dim \mathcal{L}_{\phi(x)}(Z) = i(Z) - 2 \geq m$ and Proposition 2.5, it is induced that there exists an ECO line ℓ with respect to Y through $\phi(x)$ and contained in Z . Take one such ECO line ℓ . The claim above shows that the inverse image $\phi^{-1}(\ell)$ consists of two smooth rational curves of degree 1 with respect to ϕ^*H . Clearly, one of those two curves passes through x . \square

4. Defining equations of VMRT

In order to find the defining ideal of the VMRT $\mathcal{C}_x \subset \mathbb{P}T_x(X^Z)$, we proceed in a manner analogous to [Hwang and Kim 2013].

Notation 4.1. Let $Y \subset \mathbb{P}^N$ be a hypersurface of even degree $2m$, $m \geq 1$, and let $Z \subset \mathbb{P}^N$ be a projective submanifold which is not contained in Y . For each point $y \in Z \setminus Z \cap Y$, we denote by $\mathcal{E}_y^Y(Z) \subset \mathbb{P}T_y(Z)$ the space of tangent directions of ECO lines with respect to Y contained in Z .

Proposition 4.2. *In the setting of Definition 3.1, for a general point $x \in X^Z$, the tangent morphism $\tau_x : \mathcal{K}_x \rightarrow \mathbb{P}T_x(X^Z)$, sending each member of \mathcal{K}_x to its tangent direction, is an embedding. In particular the VMRT $\mathcal{C}_x = \text{Im}(\tau_x) \subset \mathbb{P}T_x(X^Z)$ is a nonsingular projective variety with finitely many components of dimension $i(Z) - m - 2$, isomorphic to $\mathcal{E}_{\phi(x)}^Y(Z) \subset \mathbb{P}T_{\phi(x)}(Z)$ via the differential morphism $d\phi_x : \mathbb{P}T_x(X^Z) \rightarrow \mathbb{P}T_{\phi(x)}(Z)$.*

Proof. From Proposition 3.2, the differential $d\phi_x : \mathbb{P}T_x(X^Z) \rightarrow \mathbb{P}T_{\phi(x)}(Z)$ sends the VMRT $\mathcal{C}_x \subset \mathbb{P}T_x(X^Z)$ isomorphically to the variety $\mathcal{E}_{\phi(x)}^Y(Z) \subset \mathbb{P}T_{\phi(x)}(Z)$.

We note that $\tau_x : \mathcal{K}_x \rightarrow \mathbb{P}T_x(X^Z)$ is the normalization morphism of its image, which is equal to \mathcal{C}_x . Thus we only need to show that τ_x is an embedding because \mathcal{K}_x is a smooth projective variety of dimension $i(X) - 2 = i(Z) - m - 2$.

Assume that there are two distinct members C_1 and C_2 of \mathcal{K}_x such that $\tau_x([C_1]) = \tau_x([C_2])$. Thus $\phi(C_1)$ and $\phi(C_2)$ are lines on \mathbb{P}^N passing through $\phi(x)$ with the same tangent direction at $\phi(x)$, which implies that $\phi(C_1) = \phi(C_2)$ is a line; denote it by ℓ . Therefore $\phi^{-1}(\ell) = C_1 \cup C_2$, and hence C_1 and C_2 meets only over the points on $\ell \cap B$, a contradiction because $x \in C_1 \cap C_2$ but $\phi(x) \notin B$ by the general condition on x . Thus we have shown that τ_x is injective.

Since we know that \mathcal{K}_x is nonsingular, to prove that τ_x is an embedding, it remains to show that τ_x is an immersion. By [Hwang 2001, Proposition 1.4], this is equivalent to showing that for any member $C \subset X^Z$ of \mathcal{K}_x , the normal bundle N_{C/X^Z} satisfies

$$N_{C/XZ} = \mathcal{O}_{\mathbb{P}^1}(1)^{i(Z)-m-2} \oplus \mathcal{O}_{\mathbb{P}^1}^{n+m-i(Z)+1}.$$

By the generality of x , we can write

$$N_{C/XZ} = \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^1}(a_{n-1})$$

for integers $a_1 \geq \cdots \geq a_{n-1} \geq 0$ satisfying $\sum_i a_i = i(Z) - m - 2$. Since ϕ is unramified at general points of C and $\phi|_C : C \rightarrow \ell := \phi(C)$ is an isomorphism, we have an injective sheaf homomorphism

$$\phi_* : N_{C/XZ} \rightarrow N_{\ell/\mathbb{P}^N} = \mathcal{O}(1)^{N-1}.$$

Thus $a_1 \leq 1$; hence, $a_1 = \cdots = a_{i(Z)-m-2} = 1$ and $a_{i(Z)-m-1} = \cdots = a_{n-1} = 0$. \square

Proposition 4.3. *Let $Z \subset \mathbb{P}^N$ be a Fano manifold whose Picard group is generated by the hyperplane section class. Assume that Z is covered by lines and $i(Z) \geq 3$. Then for each m with $1 \leq m \leq i(Z) - 2$, there exists a hypersurface Y of degree $2m$ with smooth $Y \cap Z$ such that for a general point $y \in Z$, $\mathcal{E}_y^Y(Z) \subset \mathbb{P}T_y(Z)$ is scheme-theoretically the intersection of $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$ with m hypersurfaces in $\mathbb{P}T_y(Z)$ of degrees $m + 1, \dots, 2m$, respectively.*

Proof. Take a point $\hat{y} \in Z$ such that $\mathcal{L}_{\hat{y}}(Z)$ is smooth of dimension $i(Z) - 2$. Choose a homogeneous coordinate system t_0, \dots, t_N so that $\hat{y} = [1 : 0 : \cdots : 0] \in Z$ and the hyperplane section $Z \cap (t_0 = 0)$ is smooth. Choose homogeneous polynomials $\{b_k(t_1, \dots, t_N) \mid m + 1 \leq k \leq 2m\}$ with $\deg b_k = k$ so that each of the following is smooth:

- (i) the intersection of $Z \cap (t_0 \neq 0)$ with the hypersurface in \mathbb{P}^N defined by

$$1 + b_{m+1}(t_1, \dots, t_N) + \cdots + b_{2m}(t_1, \dots, t_N) = 0,$$

- (ii) the intersection of $Z \cap (t_0 = 0)$ with the hypersurface in \mathbb{P}^N defined by

$$b_{2m}(t_1, \dots, t_N) = 0.$$

Set

$$f(t_0, t_1, \dots, t_N) := t_0^{2m} + t_0^{m-1}b_{m+1}(t_1, \dots, t_N) + \cdots + t_0b_{2m-1}(t_1, \dots, t_N) + b_{2m}(t_1, \dots, t_N).$$

The assumptions (i) and (ii) imply that the hypersurface section of Z defined by $f(t_0, \dots, t_N)$ is smooth. From Proposition 2.2(ii) we obtain the equalities

$$B_k^f(y; z) = b_k(z_1, \dots, z_N), \quad m + 1 \leq k \leq 2m.$$

Since $v_y^{-1}(\mathcal{L}_y(Z))$ is smooth of dimension $i(Z) - 2 \geq m$, it follows that for general $\{b_k(z_1, \dots, z_N) \mid m + 1 \leq k \leq 2m\}$, the scheme-theoretical intersection of $v_y^{-1}(\mathcal{L}_y(Z))$ and the m hypersurfaces defined by $B_k^f(y; z)$, $m + 1 \leq k \leq 2m$,

is smooth of dimension $i(Z) - m - 2$. Therefore our proposition comes from Proposition 2.5. \square

Proof of Theorem 1.1. By Propositions 4.2 and 4.3, the theorem holds for a general hypersurface $Y \subset \mathbb{P}^N$ with smooth intersection $Y \cap Z$. In order to prove it for arbitrary hypersurface $Y \subset \mathbb{P}^N$ with smooth intersection $Y \cap Z$, choose a deformation $\{Y_t \mid |t| < \epsilon\}$ of $Y = Y_0$ with smooth $Y_t \cap Z$ such that for a Zariski open subset $U_t \subset Z \setminus Y_t \cap Z$, the varieties $\mathcal{E}_y^{Y_t}(Z) \subset \mathbb{P}T_y(Z)$, $y \in U_t$, are

- (i) smooth of dimension $i(Z) - m - 2$ for any t , and
- (ii) the intersection of m hypersurface sections of $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$ defined by hypersurfaces in $\mathbb{P}T_y(Z)$ of degrees $m + 1, \dots, 2m$, respectively. (The intersection is scheme-theoretic for any $t \neq 0$ and set-theoretic for $t = 0$.)

By shrinking ϵ if necessary, the intersection $\bigcap_t U_t$ is nonempty. Let V be a Zariski open subset of Z such that the variety $\mathcal{L}_y(Z)$ is smooth of dimension $i(Z) - 2$. Pick a point $y \in (\bigcap_t U_t) \cap V$. We can construct a smooth family $\{\phi_t : X_t^Z \rightarrow Z \mid |t| < \epsilon\}$ of double covers of Z branched along the $Z \cap Y_t$. Choose $x_t \in \phi_t^{-1}(y)$ in a continuous way. The family $\{\mathcal{K}_{x_t} \mid |t| < \epsilon\}$ is a flat family of nonsingular projective subvarieties; see, e.g., [Kollár 1996, II.3.11.5]. Via Proposition 4.2, this implies that $\{\mathcal{E}_y^{Y_t}(Z) \mid |t| < \epsilon\}$ is a flat family of nonsingular projective subvarieties of $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$. From condition (ii) and the flatness, we conclude that $\mathcal{E}_y^{Y_0}(Z)$ is also scheme-theoretically the intersection of $\mathcal{L}_y(Z) \subset \mathbb{P}T_y(Z)$ with m hypersurfaces in $\mathbb{P}T_y(Z)$ of degrees $m + 1, \dots, 2m$, respectively. \square

5. Rigidity and Extension

The following fact is obvious, but plays an important role in the proof of Theorem 1.4.

Lemma 5.1. *Let R be the polynomial ring $\mathbb{C}[z_1, \dots, z_N]$ in variables z_1, \dots, z_N . Consider R as a graded ring with $\deg z_i = 1$. Let I, J , and K be homogeneous ideals of R such that I and K are generated by homogeneous polynomials of degree 2, and J is generated by homogeneous polynomials of degrees ≥ 3 . If $h : R \rightarrow R$ is an automorphism of the graded ring R with $h(K) \subset I + J$, then $h(K) \subset I$.*

Proof of Theorem 1.4. By shrinking U if necessary, we may assume that

- the restriction $\phi|_U : U \rightarrow Z$ is an embedding,
- for any $x \in U$, $\mathcal{L}_{\phi(x)}(Z) \subset \mathbb{P}T_{\phi(x)}(Z)$ is a smooth quadratic manifold of dimension $i(Z) - 2$; see [Ionescu and Russo 2013, Theorem 2.4],
- for any $x \in U$, the space $\mathcal{E}_{\phi(x)}^Y \subset \mathbb{P}T_{\phi(x)}(Z)$ is scheme-theoretically the intersection of $\mathcal{L}_{\phi(x)}(Z) \subset \mathbb{P}T_{\phi(x)}(Z)$ and m hypersurfaces in $\mathbb{P}T_{\phi(x)}(Z)$ of degrees $m + 1, \dots, 2m$, respectively.

Set $U_1 = \phi(U)$ and $U_2 = \psi(U)$. Then we get a biholomorphism

$$\gamma := \psi \circ \phi|_U^{-1} : U_1 \rightarrow U_2.$$

We note that for any $y \in U_1$, the differential $d\gamma_y : \mathbb{P}T_y(Z) \rightarrow \mathbb{P}T_{\gamma(y)}(Z)$ is an isomorphism sending $\mathcal{E}_y^Y(Z) \subset \mathbb{P}T_y(Z)$ into $\mathcal{L}_{\gamma(y)}(Z) \subset \mathbb{P}T_{\gamma(y)}(Z)$. From Lemma 5.1 and Theorem 1.1, it follows that $d_y\gamma(\mathcal{L}_y(Z)) \subset \mathcal{L}_{\gamma(y)}(Z)$. By shrinking U if necessary, we may assume that $\mathcal{L}_{\gamma(y)} \subset \mathbb{P}T_{\gamma(y)}(Z)$ is also a smooth quadratic manifold of dimension $i(Z) - 2$. Therefore it follows that $d_y\gamma(\mathcal{L}_y(Z)) = \mathcal{L}_{\gamma(y)}(Z)$. We finish the proof by applying [Hwang 2001, Theorem 3.2]. \square

The next corollary is an algebraic version of Theorem 1.4

Corollary 5.2. *In the setting of Theorem 1.4, let \hat{X} be a projective variety with generically finite surjective morphisms $g : \hat{X} \rightarrow X^Z$ and $h : \hat{X} \rightarrow Z$ such that for a minimal rational curve C through a general point of X^Z , there exists an irreducible component C' of $g^{-1}(C)$ whose image $h(C') \subset Z \subset \mathbb{P}^N$ is a line. Then there exists an automorphism $\Gamma : Z \rightarrow Z$ such that $h = \Gamma \circ \phi \circ g$.*

Next, Theorems 5.3 and 5.4 can be proved by the same arguments as in the proof of Theorems 1.7 and 1.9 in [Hwang and Kim 2013], respectively. We include their proof for the reader’s convenience.

Theorem 5.3. *Let $Z \subset \mathbb{P}^N$ be a quadratic Fano manifold such that its Picard group is generated by the hyperplane section class and $i(Z) \geq 4$. Let $Y_1, Y_2 \subset \mathbb{P}^N$, $N \geq 3$, be two hypersurfaces of degree $2(i(Z) - 2)$ with smooth intersections $Y_1 \cap Z$ and $Y_2 \cap Z$. Let $\phi_1 : X_1 \rightarrow Z$ and $\phi_2 : X_2 \rightarrow Z$ be double covers of Z branched along $Y_1 \cap Z$ and $Y_2 \cap Z$, respectively. Suppose there exists a finite morphism $f : X_1 \rightarrow X_2$. Then f is an isomorphism.*

Proof. Put $m = i(Z) - 2$ in the proof of Proposition 4.2. Then minimal rational curves on X_i , $i = 1, 2$, have trivial normal bundles and rational curves through general points with trivial normal bundles are minimal rational curves. By [Hwang and Mok 2003, Proposition 6], for a general minimal rational curve $C \subset X_2$, each irreducible component of $f^{-1}(C)$ is a minimal rational curve in X_1 . In other words, f sends minimal rational curves of X_1 through a general point to those of X_2 . Putting

$$\hat{X} = X_1, \quad X = X_2, \quad g = f, \quad \phi = \phi_2, \quad \text{and} \quad h = \phi_1$$

in Corollary 5.2, we see that $\phi_1 = \Gamma \circ \phi_2 \circ f$ for some automorphism Γ of Z . Thus f must be birational, and hence an isomorphism. \square

The next theorem is a stronger version of Theorem 1.3.

Theorem 5.4. *Let $Z \subset \mathbb{P}^N$ be a quadratic Fano manifold such that its Picard group is generated by the hyperplane section class and $i(Z) \geq 4$. Let $Y_1, Y_2 \subset \mathbb{P}^N$ be two*

hypersurfaces of degree $2m$, $2 \leq m \leq i(Z) - 2$, with smooth $Y_1 \cap Z$ and $Y_2 \cap Z$. Let $\phi_1 : X_1^Z \rightarrow Z$ and $\phi_2 : X_2^Z \rightarrow Z$ be double covers of Z branched along $Y_1 \cap Z$ and $Y_2 \cap Z$ respectively, and let $U_1 \subset X_1^Z$ and $U_2 \subset X_2^Z$ be two connected open subsets. Suppose that we are given a biholomorphic map $\gamma : U_1 \rightarrow U_2$ such that for any minimal rational curve $C_1 \subset X_1^Z$, there exists a minimal rational curve $C_2 \subset X_2^Z$ with $\gamma(U_1 \cap C_1) = U_2 \cap C_2$. Then we can find a biregular morphism $\Gamma : X_1^Z \rightarrow X_2^Z$ with $\Gamma|_{U_1} = \gamma$.

Sketch of the proof. Applying Theorem 1.4 to $\psi := \phi_2 \circ \gamma : U_1 \rightarrow \phi_2(U_2) \subset Z$ and $\phi := \phi_1$, we have $\Gamma' \in \text{Aut}(Z)$ such that $\Gamma' \circ \phi_1|_{U_1} = \phi_2 \circ \gamma$. By the assumption on γ and Proposition 3.2, for a general point $y \in \phi_1(U_1)$, we have $d\Gamma'(\mathcal{E}_y^{Y_1}) = \mathcal{E}_{\Gamma'(y)}^{Y_2}$, which implies that a general ECO line with respect to Y_2 contained in Z should be an ECO line with respect to Y' .

Since the Picard group of Z is generated by the hyperplane section class and $\Gamma' \in \text{Aut}(Z)$, there exists a hypersurface $Y' \subset \mathbb{P}^N$ of degree $2m$ such that $\Gamma'(Y_1 \cap Z) = Y' \cap Z$. Suppose $Y' \cap Z \neq Y_2 \cap Z$. By the similar arguments in [Hwang and Kim 2013, Proposition 2.5], we can show that a general ECO line with respect to Y_2 contained in Z cannot be an ECO line with respect to Y' , a contradiction. Therefore $Y' \cap Z = Y_2 \cap Z$.

Thus replacing $Y_1 \cap Z$ by $\Gamma(Y_1 \cap Z)$ and ϕ_1 by $\Gamma' \circ \phi_1$, we may assume that $Y_1 \cap Z = Y_2 \cap Z$ and $\phi_1(U_1) = \phi_2(U_2)$. By the uniqueness of double covering, it follows that there exists a biregular morphism $\Gamma : X_1^Z \rightarrow X_2^Z$ with $\Gamma|_{U_1} = \gamma$. \square

Acknowledgments

The author would like to thank Jun-Muk Hwang for a number of discussions on this subject. She is also grateful to the referee for valuable comments to modify the original version.

References

- [Hwang 2001] J.-M. Hwang, “Geometry of minimal rational curves on Fano manifolds”, pp. 335–393 in *School on Vanishing Theorems and Effective Results in Algebraic Geometry* (Trieste, 2000), edited by J. P. Demailly et al., ICTP Lecture Notes **6**, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001. MR 2003g:14054 Zbl 1086.14506
- [Hwang and Kim 2013] J.-M. Hwang and H. Kim, “Varieties of minimal rational tangents on double covers of projective space”, *Math. Z.* **275**:1–2 (2013), 109–125. MR 3101799 Zbl 1282.14070
- [Hwang and Mok 2001] J.-M. Hwang and N. Mok, “Cartan–Fubini type extension of holomorphic maps for Fano manifolds of Picard number 1”, *J. Math. Pures Appl.* (9) **80**:6 (2001), 563–575. MR 2002j:32023 Zbl 1033.32013
- [Hwang and Mok 2003] J.-M. Hwang and N. Mok, “Finite morphisms onto Fano manifolds of Picard number 1 which have rational curves with trivial normal bundles”, *J. Algebraic Geom.* **12**:4 (2003), 627–651. MR 2005f:14086 Zbl 1038.14018

- [Hwang and Mok 2004] J.-M. Hwang and N. Mok, “Birationality of the tangent map for minimal rational curves”, *Asian J. Math.* **8**:1 (2004), 51–63. MR 2006c:14065 Zbl 1072.14015
- [Ionescu and Russo 2013] P. Ionescu and F. Russo, “Manifolds covered by lines and the Hartshorne conjecture for quadratic manifolds”, *Amer. J. Math.* **135**:2 (2013), 349–360. MR 3038714 Zbl 1271.14055
- [Kebekus 2002] S. Kebekus, “Families of singular rational curves”, *J. Algebraic Geom.* **11**:2 (2002), 245–256. MR 2003e:14011 Zbl 1054.14035
- [Kollár 1996] J. Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik (3) **32**, Springer, Berlin, 1996. MR 98c:14001

Received November 27, 2014. Revised August 22, 2015.

HOSUNG KIM
NATIONAL INSTITUTE FOR MATHEMATICAL SCIENCE
DAEJEON 34047
REPUBLIC OF KOREA
hosung@nims.re.kr

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 282 No. 2 June 2016

Exhausting curve complexes by finite rigid sets	257
JAVIER ARAMAYONA and CHRISTOPHER J. LEININGER	
A variational characterization of flat spaces in dimension three	285
GIOVANNI CATINO, PAOLO MASTROLIA and DARIO D. MONTICELLI	
Estimates of the gaps between consecutive eigenvalues of Laplacian	293
DAGUANG CHEN, TAO ZHENG and HONGCANG YANG	
Liouville type theorems for the p -harmonic functions on certain manifolds	313
JINGYI CHEN and YUE WANG	
Cartan–Fubini type rigidity of double covering morphisms of quadratic manifolds	329
HOSUNG KIM	
On the uniform squeezing property of bounded convex domains in \mathbb{C}^n	341
KANG-TAE KIM and LIYOU ZHANG	
Lefschetz pencils and finitely presented groups	359
RYOMA KOBAYASHI and NAOYUKI MONDEN	
Knot homotopy in subspaces of the 3-sphere	389
YUYA KODA and MAKOTO OZAWA	
On the relationship of continuity and boundary regularity in prescribed mean curvature Dirichlet problems	415
KIRK E. LANCASTER and JARON MELIN	
Bridge spheres for the unknot are topologically minimal	437
JUNG HOON LEE	
On the geometric construction of cohomology classes for cocompact discrete subgroups of $SL_n(\mathbb{R})$ and $SL_n(\mathbb{C})$	445
SUSANNE SCHIMPF	
On Blaschke’s conjecture	479
XIAOLE SU, HONGWEI SUN and YUSHENG WANG	
The role of the Jacobi identity in solving the Maurer–Cartan structure equation	487
ORI YUDILEVICH	



0030-8730(2016)282:2;1-3