

*Pacific  
Journal of  
Mathematics*

**CONJUGACY AND ELEMENT-CONJUGACY  
OF HOMOMORPHISMS OF COMPACT LIE GROUPS**

YINGJUE FANG, GANG HAN AND BINYONG SUN

# CONJUGACY AND ELEMENT-CONJUGACY OF HOMOMORPHISMS OF COMPACT LIE GROUPS

YINGJUE FANG, GANG HAN AND BINYONG SUN

Let  $G$  be a connected compact Lie group. Among other things, we prove that the following are equivalent. (a) For all connected compact Lie groups  $H$  and all continuous homomorphisms  $\phi, \phi' : H \rightarrow G$ , if  $\phi(h)$  and  $\phi'(h)$  are conjugate in  $G$  for all  $h \in H$ , then  $\phi$  and  $\phi'$  are  $G$ -conjugate. (b) The Lie algebra of  $G$  contains no simple ideal of type  $D_n$  ( $n \geq 4$ ),  $E_6$ ,  $E_7$ , or  $E_8$ .

## 1. Introduction

Let  $G$  and  $H$  be two topological groups, and let  $\phi, \phi' : H \rightarrow G$  be two continuous homomorphisms. We say that  $\phi$  and  $\phi'$  are conjugate if there is an element  $g \in G$  such that

$$g\phi(h)g^{-1} = \phi'(h) \quad \text{for all } h \in H.$$

We say that they are element-conjugate if for every  $h \in H$ , there is a  $g \in G$  such that

$$g\phi(h)g^{-1} = \phi'(h).$$

Clearly, conjugate homomorphisms are element-conjugate. Conversely, we are interested to know to what extent the following statement holds.

(1) If  $\phi$  and  $\phi'$  are element-conjugate, then they are conjugate.

This is closely related to the failure of multiplicity one for the cuspidal spectrum of reductive groups over number fields (see [Blasius 1994, Section 1.1; Lapid 1999, Section 3; Arthur 2002, page 471; Lafforgue 2014, Section 0.8]). Some counterexamples to (1) are used to construct nonisometric pairs of isospectral manifolds (see [Larsen 1996, Theorem 2.7]).

We say that  $G$  is  $H$ -acceptable if (1) holds for all continuous homomorphisms  $\phi$  and  $\phi'$ . M. Larsen [1994] defined  $G$  to be acceptable if it is  $H$ -acceptable whenever  $H$  is finite. In [Larsen 1996], he classified acceptable, connected, simply connected compact Lie groups. In this paper, we are more concerned with the

---

B. Sun was supported by NSFC Grants 11321101 and 11525105.

MSC2010: 22E46.

Keywords: element-conjugacy, conjugacy, compact Lie group.

statement (1) when both  $G$  and  $H$  are connected compact Lie groups. This is more relevant to the classification of reductive subalgebras of semisimple Lie algebras, as studied in [Dynkin 1952; Liebeck and Seitz 1996; Malcev 1944; Minchenko 2006].

It was essentially known to Dynkin how counterexamples to (1) can be constructed when  $G$  is a simple compact Lie group of type  $D_n$ , with  $n \geq 4$  (see [Dynkin 1952, Theorem 1.4]; see also [Wang 2012]). M. Liebeck and G. Seitz [1996] found a counterexample when  $G$  is simple of type  $E_8$ , and  $H$  is simple of type  $A_2$ , in the setting of algebraic groups. For Lie algebra homomorphisms from a semisimple Lie algebra to a simple Lie algebra of type  $E_6$ ,  $E_7$  or  $E_8$ , all counterexamples of the Lie algebra analogue of (1) are listed in [Minchenko 2006, Table 9] (see Lemma 3.6).

Based on the main result of [Minchenko 2006], we prove the following theorem for connected compact Lie groups.

**Theorem 1.1.** *Let  $G$  be a connected compact Lie group. Then the following are equivalent. (a) The Lie algebra of  $G$  contains no simple ideal of type  $D_n$  ( $n \geq 4$ ),  $E_6$ ,  $E_7$  or  $E_8$ . (b) The group  $G$  is  $H$ -acceptable for all connected compact Lie groups  $H$ .*

As a byproduct of the proof of Theorem 1.1, we get the following theorem for classical Lie groups.

**Theorem 1.2.** *Let  $G$  be a classical Lie group, that is,  $G$  is  $GL_n(\mathbb{R})$ ,  $GL_n(\mathbb{C})$ ,  $GL_n(\mathbb{H})$ ,  $U(p, q)$ ,  $O(p, q)$ ,  $O_n(\mathbb{C})$ ,  $Sp_{2n}(\mathbb{R})$ ,  $Sp_{2n}(\mathbb{C})$ ,  $Sp(p, q)$  or  $O^*(2n)$ , where  $p, q, n \geq 0$ . Then  $G$  is  $H$ -acceptable for all compact Hausdorff topological groups  $H$ .*

By convention, when  $n = 0$  or  $p + q = 0$ , the corresponding classical group of the above theorem is the trivial group, and the theorem is trivial in this case.

## 2. Classical Lie groups

In the rest of the paper, let  $H$  be a compact Hausdorff topological group. By abuse of notation, we do not distinguish a representation from its underlying vector space. All representations are assumed to be complex, finite-dimensional and continuous.

**Classical complex groups.** We begin with the following classical result.

**Proposition 2.1.** *The complex general linear group  $GL_n(\mathbb{C})$  ( $n \geq 0$ ) is  $H$ -acceptable.*

*Proof.* This is well known. Let  $\phi, \phi' : H \rightarrow GL_n(\mathbb{C})$  be two continuous homomorphisms, to be viewed as two  $n$ -dimensional representations of  $H$ . If they are element-conjugate, then they have the same character. By the classical character theory of representations of compact groups, these two representations are isomorphic. This is the same as saying that the homomorphisms  $\phi$  and  $\phi'$  are conjugate.  $\square$

We define an orthogonal representation of  $H$  to be a representation  $V$  of  $H$  together with an  $H$ -invariant orthogonal form  $\langle \cdot, \cdot \rangle$  on it. Here by an orthogonal form, we mean a nondegenerate symmetric bilinear form. Two orthogonal representations  $(V, \langle \cdot, \cdot \rangle)$  and  $(V', \langle \cdot, \cdot \rangle')$  are called isomorphic if there is an  $H$ -intertwining linear isomorphism from  $V$  to  $V'$  which sends  $\langle \cdot, \cdot \rangle$  to  $\langle \cdot, \cdot \rangle'$ .

Similarly, we define the notions of symplectic representations and isomorphisms of symplectic representations. The next result is well known (see [Malcev 1944]).

**Proposition 2.2.** *Let  $(V, \langle \cdot, \cdot \rangle)$  and  $(V', \langle \cdot, \cdot \rangle')$  be two orthogonal (symplectic) representations of  $H$ . If  $V$  and  $V'$  are isomorphic as representations of  $H$ , then  $(V, \langle \cdot, \cdot \rangle)$  and  $(V', \langle \cdot, \cdot \rangle')$  are isomorphic as orthogonal (symplectic) representations.*

It is clear that Propositions 2.1 and 2.2 imply the following result, which is stated and proved in [Larsen 1994, Propositions 2.3 and 2.4], in the setting that  $H$  is a finite group.

**Proposition 2.3.** *The complex orthogonal group  $O_n(\mathbb{C})$  and the complex symplectic group  $Sp_{2n}(\mathbb{C})$  ( $n \geq 0$ ) are  $H$ -acceptable.*

**Maximal compact subgroups.** Let  $G$  be a Lie group with finitely many connected components. Let  $K$  be a maximal compact subgroup of  $G$ , which always exists and is unique up to conjugation [Borel 1998, Chapter VII, Theorem 1.2(i)]. Write  $i_K : K \rightarrow G$  for the inclusion map.

**Lemma 2.4.** *Let  $\phi_K, \phi'_K : H \rightarrow K$  be two continuous homomorphisms. Write  $\phi := i_K \circ \phi_K$  and  $\phi' := i_K \circ \phi'_K$ . Then*

- (a)  $\phi$  and  $\phi'$  are conjugate if and only if  $\phi_K$  and  $\phi'_K$  are conjugate, and
- (b)  $\phi$  and  $\phi'$  are element-conjugate if and only if  $\phi_K$  and  $\phi'_K$  are element-conjugate.

*Proof.* We only prove (a), since (b) can be proved by the same method, and is also implied by (a). The “if” part of (a) is obvious. We prove the “only if” part below.

Write “Ad” for the conjugation action. By [Borel 1998, Chapter VII, Theorem 1.2(ii)], there is a closed analytic submanifold  $E$  of  $G$  such that

$$(2) \quad \text{Ad}_k(E) = E \quad \text{for all } k \in K,$$

and every  $g \in G$  is uniquely of the form

$$(3) \quad g = ke, \quad \text{with } k \in K, e \in E.$$

Assume that  $\phi$  and  $\phi'$  are conjugate, i.e., there is an element  $g$  in  $G$  such that

$$\phi' = \text{Ad}_g \circ \phi.$$

Write  $g = ke$  as in (3). Then

$$\text{Ad}_{k^{-1}} \circ \phi' = \text{Ad}_e \circ \phi.$$

Let  $h \in H$  and put

$$k_1 := (\text{Ad}_{k^{-1}} \circ \phi')(h), \quad k_2 := \phi(h).$$

Then  $k_1, k_2 \in K$  and

$$k_1 = \text{Ad}_e(k_2),$$

or the same,

$$k_1 e = k_2 \text{Ad}_{k_2^{-1}}(e).$$

Now (2) and the uniqueness of the decomposition (3) imply that  $k_1 = k_2$ . This proves that  $\text{Ad}_{k^{-1}} \circ \phi' = \phi$ , and thus  $\phi_K$  and  $\phi'_K$  are conjugate.  $\square$

The following result generalizes [Larsen 1994, Proposition 1.7].

**Proposition 2.5.** *The group  $G$  is  $H$ -acceptable if and only if so is  $K$ .*

*Proof.* Let us prove the “if” part first. Assume that  $K$  is  $H$ -acceptable, and let  $\phi, \phi' : H \rightarrow G$  be two continuous homomorphisms which are element-conjugate. We need to prove that  $\phi$  and  $\phi'$  are conjugate. Since every compact subgroup of  $G$  is conjugate to a subgroup of  $K$ , we assume without loss of generality that the images of  $\phi$  and  $\phi'$  are both contained in  $K$ . Let  $\phi_K$  and  $\phi'_K$  be as in Lemma 2.4 so that  $\phi := i_K \circ \phi_K$  and  $\phi' := i_K \circ \phi'_K$ . Then Lemma 2.4 implies that  $\phi_K$  and  $\phi'_K$  are element-conjugate, and they are conjugate since  $K$  is  $H$ -acceptable. This implies that  $\phi$  and  $\phi'$  are conjugate.

To prove the “only if” part, we assume that  $G$  is  $H$ -acceptable. Write  $\phi, \phi', \phi_K$  and  $\phi'_K$  as before. Assume that  $\phi_K$  and  $\phi'_K$  are element-conjugate. Then  $\phi$  and  $\phi'$  are element-conjugate, and therefore conjugate since  $G$  is  $H$ -acceptable. Now Lemma 2.4 implies that  $\phi_K$  and  $\phi'_K$  are conjugate. This proves that  $K$  is  $H$ -acceptable.  $\square$

**Corollary 2.6.** *The compact groups  $U(n)$ ,  $O(n)$  and  $Sp(n)$ , where  $n \geq 0$ , are  $H$ -acceptable.*

*Proof.* Note that  $U(n)$ ,  $O(n)$  or  $Sp(n)$  is a maximal compact subgroup of  $GL_n(\mathbb{C})$ ,  $O_n(\mathbb{C})$  or  $Sp_{2n}(\mathbb{C})$ , respectively. Therefore the corollary is a consequence of Propositions 2.1, 2.3 and 2.5.  $\square$

The following lemma is obvious.

**Lemma 2.7.** *Let  $G_1$  and  $G_2$  be two topological groups. Then  $G_1 \times G_2$  is  $H$ -acceptable if and only if so are both  $G_1$  and  $G_2$ .*

Now we come to the proof of [Theorem 1.2](#). When  $G$  is  $\mathrm{GL}_n(\mathbb{R})$ ,  $\mathrm{GL}_n(\mathbb{C})$ ,  $\mathrm{GL}_n(\mathbb{H})$ ,  $\mathrm{O}(p, q)$ ,  $\mathrm{O}_n(\mathbb{C})$ ,  $\mathrm{U}(p, q)$ ,  $\mathrm{Sp}(p, q)$ ,  $\mathrm{Sp}_{2n}(\mathbb{R})$ ,  $\mathrm{Sp}_{2n}(\mathbb{C})$  or  $\mathrm{O}^*(2n)$ , its maximal compact subgroup is  $\mathrm{O}(n)$ ,  $\mathrm{U}(n)$ ,  $\mathrm{Sp}(n)$ ,  $\mathrm{O}(p) \times \mathrm{O}(q)$ ,  $\mathrm{O}(n)$ ,  $\mathrm{U}(p) \times \mathrm{U}(q)$ ,  $\mathrm{Sp}(p) \times \mathrm{Sp}(q)$ ,  $\mathrm{U}(n)$ ,  $\mathrm{Sp}(n)$  or  $\mathrm{U}(n)$ , respectively. By [Corollary 2.6](#) and [Lemma 2.7](#), all these compact groups are  $H$ -acceptable. Therefore  $G$  is  $H$ -acceptable by [Proposition 2.5](#). This proves [Theorem 1.2](#).

We record the following two results for later use.

**Corollary 2.8.** *The compact groups  $\mathrm{SU}(n)$  and  $\mathrm{SO}(2n + 1)$ , where  $n \geq 0$ , are  $H$ -acceptable.*

*Proof.* By [Corollary 2.6](#), the groups  $\mathrm{U}(n)$  and  $\mathrm{O}(2n + 1)$  are  $H$ -acceptable. Then the corollary follows by noting that every inner automorphism of  $\mathrm{U}(n)$  or  $\mathrm{O}(2n + 1)$  restricts to an inner automorphism of  $\mathrm{SU}(n)$  or  $\mathrm{SO}(2n + 1)$ , respectively.  $\square$

The following lemma is obvious.

**Lemma 2.9.** *If  $G$  is commutative, then it is  $H$ -acceptable.*

### 3. A proof of [Theorem 1.1](#)

In this section, we concentrate on connected compact Lie groups.

**Lemma 3.1.** *Let  $\phi, \phi' : H \rightarrow G$  be two continuous homomorphisms of connected compact Lie groups. Then they are element-conjugate if and only if  $\phi|_S$  and  $\phi'|_S$  are conjugate, where  $S$  is a maximal torus in  $H$ .*

*Proof.* This is because  $S$  is topologically cyclic, and every element of  $H$  is  $H$ -conjugate to an element of  $S$ .  $\square$

**Lemma 3.2.** *Let  $\rho : \tilde{G} \rightarrow G$  be a surjective continuous homomorphism with finite kernel of connected compact Lie groups. Let  $\tilde{\phi}, \tilde{\phi}' : H \rightarrow \tilde{G}$  be two continuous homomorphisms of connected compact Lie groups. Then  $\tilde{\phi}$  and  $\tilde{\phi}'$  are conjugate if and only if  $\rho \circ \tilde{\phi}$  and  $\rho \circ \tilde{\phi}'$  are conjugate, and  $\tilde{\phi}$  and  $\tilde{\phi}'$  are element-conjugate if and only if  $\rho \circ \tilde{\phi}$  and  $\rho \circ \tilde{\phi}'$  are element-conjugate.*

*Proof.* The first assertion easily follows from the observation that

$$\tilde{\phi} = \tilde{\phi}' \quad \text{if and only if} \quad \rho \circ \tilde{\phi} = \rho \circ \tilde{\phi}'.$$

We leave the details to the reader. The second assertion is a consequence of the first one and [Lemma 3.1](#).  $\square$

**Lemma 3.3.** *Let  $G$  and  $G'$  be two connected compact Lie groups with isomorphic Lie algebras. If  $G$  is  $H$ -acceptable for all connected compact Lie groups  $H$ , then so is  $G'$ .*

*Proof.* Note that  $G$  and  $G'$  have a common finite fold covering group, that is, there is a connected compact Lie group  $\tilde{G}$ , and surjective continuous homomorphisms  $\rho : \tilde{G} \rightarrow G$  and  $\rho' : \tilde{G} \rightarrow G'$  with finite kernels.

Assume that  $G$  is  $H$ -acceptable for all connected compact Lie groups  $H$ . Let  $\tilde{\phi}_1, \tilde{\phi}_2 : H \rightarrow \tilde{G}$  be two element-conjugate continuous homomorphisms. Then  $\rho \circ \tilde{\phi}_1$  and  $\rho \circ \tilde{\phi}_2$  are element-conjugate. Therefore by the assumption on  $G$ ,  $\rho \circ \tilde{\phi}_1$  and  $\rho \circ \tilde{\phi}_2$  are conjugate. Therefore by Lemma 3.2,  $\tilde{\phi}_1$  and  $\tilde{\phi}_2$  are conjugate. This shows that  $\tilde{G}$  is  $H$ -acceptable for all connected compact Lie groups  $H$ .

To prove that  $G'$  is  $H$ -acceptable for all connected compact Lie groups  $H$ , we let  $\phi'_1, \phi'_2 : H \rightarrow G'$  be two element-conjugate continuous homomorphisms. Then there is a connected compact Lie group  $\tilde{H}$ , with a surjective continuous homomorphism  $\rho_H : \tilde{H} \rightarrow H$  with finite kernel, and two continuous homomorphisms  $\tilde{\phi}'_1, \tilde{\phi}'_2 : \tilde{H} \rightarrow \tilde{G}$  such that the diagram

$$\begin{array}{ccc} \tilde{H} & \xrightarrow{\tilde{\phi}'_i} & \tilde{G} \\ \rho_H \downarrow & & \downarrow \rho' \\ H & \xrightarrow{\phi'_i} & G' \end{array}$$

commutes ( $i = 1, 2$ ).

Since  $\phi'_1$  and  $\phi'_2$  are element-conjugate, we know that  $\phi'_1 \circ \rho_H$  and  $\phi'_2 \circ \rho_H$  are element-conjugate, or equivalently,  $\rho' \circ \tilde{\phi}'_1$  and  $\rho' \circ \tilde{\phi}'_2$  are element-conjugate. Then by Lemma 3.2,  $\tilde{\phi}'_1$  and  $\tilde{\phi}'_2$  are element-conjugate. Therefore  $\tilde{\phi}'_1$  and  $\tilde{\phi}'_2$  are conjugate since  $\tilde{G}$  is  $\tilde{H}$ -acceptable. Thus  $\rho' \circ \tilde{\phi}'_1$  and  $\rho' \circ \tilde{\phi}'_2$  are conjugate, or equivalently,  $\phi'_1 \circ \rho_H$  and  $\phi'_2 \circ \rho_H$  are conjugate. This implies that  $\phi'_1$  and  $\phi'_2$  are conjugate. Thus  $G'$  is  $H$ -acceptable.  $\square$

We say that two homomorphisms  $\phi, \phi' : \mathfrak{h} \rightarrow \mathfrak{g}$  between two finite-dimensional complex Lie algebras are conjugate if there is an inner automorphism  $\varphi$  of  $\mathfrak{g}$  such that  $\varphi \circ \phi = \phi'$ .

**Lemma 3.4** [Minchenko 2006, Theorem 3; Dynkin 1952, Theorem 1.1]. *Let  $\mathfrak{g}$  be a simple complex Lie algebra of type  $G_2$  or  $F_4$ . Let  $\mathfrak{h}$  be a reductive complex Lie algebra, and let  $\phi, \phi' : \mathfrak{h} \rightarrow \mathfrak{g}$  be two injective Lie algebra homomorphisms whose images are reductive Lie subalgebras of  $\mathfrak{g}$ . If  $\phi|_{\mathfrak{s}}$  and  $\phi'|_{\mathfrak{s}}$  are conjugate, then  $\phi$  and  $\phi'$  are conjugate, where  $\mathfrak{s}$  is a Cartan subalgebra of  $\mathfrak{h}$ .*

Recall that a Lie subalgebra of a finite-dimensional complex Lie algebra  $\mathfrak{g}$  is said to be reductive if its adjoint representation on  $\mathfrak{g}$  is completely reducible.

Lemma 3.4 has the following consequence.

**Proposition 3.5.** *Let  $G$  be a connected compact Lie group whose complexified Lie algebra is simple of type  $G_2$  or  $F_4$ . Then  $G$  is  $H$ -acceptable for all connected compact Lie groups  $H$ .*

*Proof.* Let  $\phi, \phi' : H \rightarrow G$  be two element-conjugate homomorphisms. We want to show that they are conjugate. Note that  $\phi$  and  $\phi'$  have the same kernel. Replacing  $H$  by its quotient by the kernel, we assume without loss of generality that both  $\phi$  and  $\phi'$  are injective. Let  $S$  be a maximal torus in  $H$ . Write  $c_G : G \rightarrow G_{\mathbb{C}}$  for the universal complexification of  $G$ , which is injective (see [Hochschild 1966]). Write  $\mathfrak{s}, \mathfrak{h}$  and  $\mathfrak{g}$  for the complexified Lie algebras of  $S, H$  and  $G$ , respectively.

By Lemma 3.1,  $\phi|_S$  and  $\phi'|_S$  are conjugate. Therefore their complexified differentials

$$d(\phi|_S) : \mathfrak{s} \rightarrow \mathfrak{g} \quad \text{and} \quad d(\phi'|_S) : \mathfrak{s} \rightarrow \mathfrak{g}$$

are conjugate. Then Lemma 3.4 implies that the complexified differentials

$$d(\phi) : \mathfrak{h} \rightarrow \mathfrak{g} \quad \text{and} \quad d(\phi') : \mathfrak{h} \rightarrow \mathfrak{g}$$

are conjugate. This implies that the homomorphisms

$$c_G \circ \phi : H \rightarrow G_{\mathbb{C}} \quad \text{and} \quad c_G \circ \phi' : H \rightarrow G_{\mathbb{C}}$$

are conjugate. Since  $G$  is a maximal compact subgroup of  $G_{\mathbb{C}}$ , Lemma 2.4 implies that  $\phi$  and  $\phi'$  are conjugate. This proves the proposition.  $\square$

**Lemma 3.6.** *Let  $\mathfrak{g}$  be a simple complex Lie algebra of type  $D_n$  ( $n \geq 4$ ),  $E_6, E_7$  or  $E_8$ . Then there are a semisimple complex Lie algebra  $\mathfrak{h}$  and two nonconjugate injective Lie algebra homomorphisms  $\phi, \phi' : \mathfrak{h} \rightarrow \mathfrak{g}$  such that  $\phi|_{\mathfrak{s}}$  and  $\phi'|_{\mathfrak{s}}$  are conjugate. Here  $\mathfrak{s}$  is a Cartan subalgebra of  $\mathfrak{h}$ .*

*Proof.* The lemma is a consequence of [Dynkin 1952, Theorem 1.4] when  $\mathfrak{g}$  has type  $D_n$  ( $n \geq 4$ ) and a consequence of [Minchenko 2006, Theorem 7] when  $\mathfrak{g}$  has type  $E_6, E_7$  or  $E_8$ .  $\square$

Lemma 3.6 has the following consequence.

**Proposition 3.7.** *Let  $G$  be a connected compact Lie group whose complexified Lie algebra is simple of type  $D_n$  ( $n \geq 4$ ),  $E_6, E_7$  or  $E_8$ . Then  $G$  is not  $H$ -acceptable for some connected compact Lie group  $H$ .*

*Proof.* Write  $\mathfrak{g}$  for the complexified Lie algebra of  $G$ . Let  $\mathfrak{h}, \mathfrak{s}$ , and  $\phi, \phi' : \mathfrak{h} \rightarrow \mathfrak{g}$  be as in Lemma 3.6. As in the proof of Proposition 3.5, write  $c_G : G \rightarrow G_{\mathbb{C}}$  for the universal complexification of  $G$ . Let  $H_{\mathbb{C}}$  be a simply connected, connected complex Lie group whose Lie algebra is identified with  $\mathfrak{h}$ . Then  $\phi, \phi'$  integrate to holomorphic homomorphisms

$$(4) \quad \psi_{\mathbb{C}}, \psi'_{\mathbb{C}} : H_{\mathbb{C}} \rightarrow G_{\mathbb{C}}.$$

Take a maximal compact subgroup  $H$  of  $H_{\mathbb{C}}$ , and a maximal torus  $S$  in  $H$  such that the complexified Lie algebra of  $S$  equals  $\mathfrak{s}$ . Replacing  $\phi$  and  $\phi'$  by their conjugations by appropriate elements of  $G_{\mathbb{C}}$ , we assume without loss of generality



that  $\psi_{\mathbb{C}}(H) \subset G$  and  $\psi'_{\mathbb{C}}(H) \subset G$ . Then the homomorphisms in (4) restrict to two homomorphisms

$$\psi, \psi' : H \rightarrow G.$$

Since  $\phi$  and  $\phi'$  are nonconjugate, we know that  $\psi_{\mathbb{C}}$  and  $\psi'_{\mathbb{C}}$  are nonconjugate, which implies that  $\psi$  and  $\psi'$  are nonconjugate. On the other hand, since  $\phi|_S$  and  $\phi'|_S$  are conjugate, we know that  $\psi_{\mathbb{C}}|_S$  and  $\psi'_{\mathbb{C}}|_S$  are conjugate. Then Lemma 2.4 implies that  $\psi|_S$  and  $\psi'|_S$  are conjugate. This implies that  $\psi$  and  $\psi'$  are element-conjugate by Lemma 3.1. This proves the proposition.  $\square$

Finally, in view of Lemmas 3.3 and 2.7, Theorem 1.1 is a consequence of Lemma 2.9, Corollaries 2.6 and 2.8, and Propositions 3.5 and 3.7.

## References

- [Arthur 2002] J. Arthur, “A note on the automorphic Langlands group”, *Canad. Math. Bull.* **45**:4 (2002), 466–482. [MR 1941222](#) [Zbl 1031.11066](#)
- [Blasius 1994] D. Blasius, “On multiplicities for  $SL(n)$ ”, *Israel J. Math.* **88**:1-3 (1994), 237–251. [MR 1303497](#) [Zbl 0826.11023](#)
- [Borel 1998] A. Borel, *Semisimple groups and Riemannian symmetric spaces*, Texts and Readings in Mathematics **16**, Hindustan Book Agency, New Delhi, 1998. [MR 1661166](#) [Zbl 0954.53002](#)
- [Dynkin 1952] E. B. Dynkin, “Полупростые подалгебры полупростых алгебр Ли”, *Mat. Sbornik (N.S.)* **30**:72 (1952), 349–462. Translated as “Semisimple subalgebras of semisimple Lie algebras” in *Amer. Math. Soc. Transl. (2)* **6** (1957), 111–243. [MR 0047629](#) [Zbl 0077.03404](#)
- [Hochschild 1966] G. Hochschild, “Complexification of real analytic groups”, *Trans. Amer. Math. Soc.* **125** (1966), 406–413. [MR 0206141](#) [Zbl 0149.27603](#)
- [Lafforgue 2014] V. Lafforgue, “Introduction to chtoucas for reductive groups and to the global Langlands parameterization”, preprint, 2014. [arXiv 1404.6416](#)
- [Lapid 1999] E. M. Lapid, “Some results on multiplicities for  $SL(n)$ ”, *Israel J. Math.* **112** (1999), 157–186. [MR 1714998](#) [Zbl 0937.22002](#)
- [Larsen 1994] M. Larsen, “On the conjugacy of element-conjugate homomorphisms”, *Israel J. Math.* **88**:1-3 (1994), 253–277. [MR 1303498](#) [Zbl 0898.20025](#)
- [Larsen 1996] M. Larsen, “On the conjugacy of element-conjugate homomorphisms II”, *Quart. J. Math. Oxford Ser. (2)* **47**:185 (1996), 73–85. [MR 1380951](#) [Zbl 0898.20026](#)
- [Liebeck and Seitz 1996] M. W. Liebeck and G. M. Seitz, *Reductive subgroups of exceptional algebraic groups*, Memoirs of the American Mathematical Society **121**:580, American Mathematical Society, Providence, RI, 1996. [MR 1329942](#) [Zbl 0851.20045](#)
- [Malcev 1944] A. I. Malcev, “О полупростых подгруппах групп Ли”, *Izvestia Akad. Nauk SSSR* **8**:4 (1944), 143–174. Translated as “On semi-simple subgroups of Lie groups” in *Amer. Math. Soc. Transl.* **33** (1950). [MR 0011303](#) [Zbl 0061.04701](#)
- [Minchenko 2006] A. N. Minchenko, “Полупростые подалгебры особых алгебр Ли”, *Tr. Mosk. Mat. Obs.* **67** (2006), 256–293. Translated as “Semisimple subalgebras of exceptional Lie algebras” in *Trans. Mosc. Math. Soc.* **2006** (2006), 225–259. [MR 2301595](#) [Zbl 1152.17003](#)
- [Wang 2012] S. Wang, “On dimension data and local vs. global conjugacy”, pp. 365–382 in *Fifth International Congress of Chinese Mathematicians* (Beijing, 2010), edited by L. Ji et al., AMS/IP

Studies in Advanced Mathematics **51**, part 1, American Mathematical Society, Providence, RI, 2012.  
[MR 2908081](#) [Zbl 1246.22028](#)

Received September 29, 2015. Revised January 13, 2016.

YINGJUE FANG  
COLLEGE OF MATHEMATICS AND STATISTICS  
SHENZHEN UNIVERSITY  
SHENZHEN, 518060  
CHINA  
[joyfang@szu.edu.cn](mailto:joyfang@szu.edu.cn)

GANG HAN  
DEPARTMENT OF MATHEMATICS  
ZHEJIANG UNIVERSITY  
P.O. BOX 1511  
HANG ZHOU, 310027  
CHINA  
[mathgg@zju.edu.cn](mailto:mathgg@zju.edu.cn)

BINYONG SUN  
ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE  
CHINESE ACADEMY OF SCIENCES  
BEIJING, 100190  
CHINA  
[sun@math.ac.cn](mailto:sun@math.ac.cn)

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Igor Pak  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pak.pjm@gmail.com](mailto:pak.pjm@gmail.com)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2016 is US \$/year for the electronic version, and \$/year for print and electronic.

Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 283    No. 1    July 2016

---

A New family of simple $\mathfrak{gl}_{2n}(\mathbb{C})$ -modules JONATHAN NILSSON	1
Derived categories of representations of small categories over commutative noetherian rings BENJAMIN ANTIEAU and GREG STEVENSON	21
Vector bundles over a real elliptic curve INDRANIL BISWAS and FLORENT SCHAFFHAUSER	43
$\mathbb{Q}(\mathbb{N})$ -graded Lie superalgebras arising from fermionic-bosonic representations JIN CHENG	63
Conjugacy and element-conjugacy of homomorphisms of compact Lie groups YINGJUE FANG, GANG HAN and BINYONG SUN	75
Entire sign-changing solutions with finite energy to the fractional Yamabe equation DANILO GARRIDO and MONICA MUSSO	85
Calculation of local formal Mellin transforms ADAM GRAHAM-SQUIRE	115
The untwisting number of a knot KENAN INCE	139
A Plancherel formula for $L^2(G/H)$ for almost symmetric subgroups BENT ØRSTED and BIRGIT SPEH	157
Multiplicative reduction and the cyclotomic main conjecture for $GL_2$ CHRISTOPHER SKINNER	171
Commensurators of solvable $S$ -arithmetic groups DANIEL STUDENMUND	201
Gerstenhaber brackets on Hochschild cohomology of quantum symmetric algebras and their group extensions SARAH WITHERSPOON and GUODONG ZHOU	223