

*Pacific
Journal of
Mathematics*

**A PLANCHEREL FORMULA FOR $L^2(G/H)$ FOR ALMOST
SYMMETRIC SUBGROUPS**

BENT ØRSTED AND BIRGIT SPEH

Volume 283 No. 1

July 2016

A PLANCHEREL FORMULA FOR $L^2(G/H)$ FOR ALMOST SYMMETRIC SUBGROUPS

BENT ØRSTED AND BIRGIT SPEH

We study the Plancherel formula for a new class of homogeneous spaces for real reductive Lie groups; these spaces are fibered over non-Riemannian symmetric spaces, and they exhibit a phenomenon of uniform infinite multiplicities. The proof for this is new but rather elementary, and we give all details. As an application we use several results from the recent literature studying possible nontemperedness of homogeneous spaces; thus we provide examples of nontempered representations of the group appearing in the Plancherel formula for our homogeneous spaces. Several classes of examples are given, each building on different techniques and new results from the theory of symmetric spaces.

I. Introduction

Considerable efforts have been devoted to obtaining the Plancherel formula for homogeneous spaces of the form G/H with G a real reductive Lie group and H a symmetric subgroup, a program completed by T. Oshima, P. Delorme, E. van den Ban, and H. Schlichtkrull. This is a central theme in harmonic analysis, and there are a number of natural ways to extend such a program. One is to consider spherical spaces, i.e., where the homogeneous space admits an open orbit of a parabolic subgroup. In this paper we shall rather extend the interest to

- (1) square-integrable sections of homogeneous line bundles over symmetric spaces, and
- (2) spaces fibered over symmetric spaces.

Of course, these two questions are related, and we shall find several classes of spaces where rather explicit answers can be found. As an example consider $G = \mathrm{SL}(2, \mathbb{R})$ with H the connected diagonal subgroup; for each unitary character of H we may consider the space (1) and the corresponding Plancherel formula: This turns out to be independent of the character, and hence the space as in (2)

Research by Speh is partially supported by NSF grant DMS-0901024.

MSC2010: primary 22E46, 43A85; secondary 22E30.

Keywords: reductive Lie group, tempered representations, Plancherel formula.

above (in our case just the group G) has the same L^2 -content as the symmetric space, only with infinite multiplicity. It is perhaps a little surprising, that one may thus find embeddings of, e.g., the discrete series of G in a uniform way in each of the spaces of sections (1).

To be more specific, our motivation in undertaking this work was to understand the disintegration of the representation of a semisimple Lie group G on the space $L^2(G/H_{ss})$ where H_{ss} is a semisimple subgroup which differs from a symmetric subgroup by a noncompact central real factor. In this paper we study this situation for the simplest nonsymmetric subgroups H_{ss} from the point of view of harmonic analysis and obtain a Plancherel theorem for space $L^2(G/H_{ss})$ in terms of the one for G/H .

Recently Y. Benoist and T. Kobayashi [2015] proved general criteria to determine if for a semisimple subgroup H the spectrum of $L^2(G/H)$ contains nontempered representations; this they use to determine in many examples if $L^2(G/H)$ is tempered. Here a representation is called tempered if it appears in the usual Plancherel formula for $L^2(G)$. However these authors do not obtain any results concerning the multiplicity of the representations in the Plancherel formula. By obtaining a Plancherel formula for $L^2(G/H_{ss})$ we are in a position to determine exactly in our examples which nontempered representations appear in the spectrum, and also to show that they appear with infinite multiplicities.

We consider a noncompact subgroup $H = H_{ss}Z_H$ where H is a subgroup of finite index in the fixpoints of an involution of G and $Z_H \simeq \mathbb{R}$ is a subgroup of finite index of the center of H . Under these assumptions we show the following.

Theorem. *As a left regular representation of G*

$$L^2(G/H_{ss}) \simeq L^2(G/H) \otimes L^2(Z_H).$$

It is instructive to compare with the situation where the central subgroup is compact, e.g., the case of G a simple noncompact Lie group and K a maximal compact subgroup with a one-dimensional center Z . Here G/K is a noncompact Riemannian symmetric space of Hermitian type, and $L^2(G/K)$ has a different Plancherel decomposition than $L^2(G/K, \chi)$, the square-integrable sections of the line bundle induced from a nontrivial unitary character χ of Z . In particular the first space contains no discrete series representations, whereas the second space typically does. Compare with Proposition III.3 for our situation of a noncompact center.

A related problem for spherical varieties over non-Archimedean fields is discussed in [Sakellaridis and Venkatesh 2014, Section 9.5].

The paper is organized as follows: In Section II, we show that we can regard H as a subgroup of finite index in the Levi subgroup of a parabolic subgroup with abelian nilradical. In Section III we prove our main theorem above. In Section IV

we discuss some examples. In particular we note that we find several examples of nontempered homogeneous spaces, some of them new; quite possibly our method could extend to other instances of Plancherel theorems, such as cases of vector bundles (as opposed to the cases of line bundles treated here).

II. Notation and preliminaries

We introduce the notation and prove some preliminary results.

Notation and assumptions. Let G be a real linear semisimple connected algebraic group with maximal compact subgroup K and complexification $G_{\mathbb{C}}$. We consider G and $K \subset G_{\mathbb{C}}$ as subgroups.

Proposition II.1. *Suppose that $P = LN$ is a maximal parabolic subgroup with an abelian nilradical N . Then L is the fixpoint set of an involution*

$$\tau : G \rightarrow G.$$

Proof (due to Dan Barbasch). We consider a maximal split Cartan subgroup and its corresponding complex Cartan subalgebra. A parabolic subalgebra is given by removing some simple roots from the diagram. The only way to get an abelian nilradical is to remove a single simple root which appears with coefficient at most one 1 if we write the roots as linear combinations of simple roots. The involution τ is then conjugation by $\exp(i\pi\varpi)$ where ϖ is the coroot of the simple root which was removed. □

Let H be a subgroup of the Levi subgroup L of P which contains the connected component L^0 of L . Then $H = H_{ss}Z_H$ where Z_H is a one dimensional connected subgroup in the center of H and H_{ss} is semisimple or discrete.

Example 1. $G = \mathrm{SL}(2, \mathbb{R})$, L diagonal matrices which are the fixed points under the conjugation by the diagonal matrix of order 2 and determinant -1 . Alternatively we consider the adjoint representation. Then L is the stabilizer of a semisimple nontrivial element of order 2. It is also the fixed point set of the automorphism by the adjoint action of the matrix

$$\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} = \exp\left(\pi i \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix}\right).$$

We have to consider two subgroups H and H_{ss} :

- (a) $H = L$, $H_{ss} = \mathbb{Z}_2$, $Z_H = \mathbb{R}_+$. $G/H_{ss} = \mathrm{PSL}(2, \mathbb{R})$,
- (b) $H = L^0$, $H_{ss} = I$ and $G/H_{ss} = \mathrm{SL}(2, \mathbb{R})$.

Proposition II.2. *Suppose that F is the fixed point set of an involution $\tau : G \rightarrow G$. Assume in addition that it is a product $F = F_{ss}Z_F$ where Z_F is a subgroup of the*

center of F isomorphic to \mathbb{R}^+ and F_{ss} is a semisimple group. Then F is contained in the Levi subgroup of a maximal parabolic subgroup P with abelian nilradical N .

Proof. We choose maximally split Cartan subgroup $C \subset F$ with complexified Lie algebra $\mathfrak{h}_{\mathbb{C}}$. We choose the simple roots of $\mathfrak{h}_{\mathbb{C}}, \mathfrak{g}_{\mathbb{C}}$ so that they are simple roots in $\mathfrak{h}_{\mathbb{C}}, \mathfrak{f}_{\mathbb{C}}$. (In the lexicographical order we let \mathfrak{f} come before \mathfrak{g} , \mathfrak{f} the Lie algebra of F .) Then \mathfrak{f} is the Levi subalgebra of a maximal parabolic subalgebra $\mathfrak{p}_{\mathbb{C}} = \mathfrak{f}_{\mathbb{C}} \oplus \mathfrak{n}_{\mathbb{C}}$ of $\mathfrak{g}_{\mathbb{C}}$.

It remains to show that N is abelian. Since τ leaves C invariant the induced homomorphism of $\tau : N \rightarrow N$ is equal to -1 . Since τ induces a Lie algebra homomorphism and hence preserves the Lie bracket in $\mathfrak{n}_{\mathbb{C}}$, the results follows from the observation that $\tau(X) = -X$ and $\tau(Y) = -Y$ then $\tau([X, Y]) = [X, Y]$. \square

Note that in the setting above, we have a direct product decomposition

$$G/H \times Z_H = G/H_{ss}.$$

This will be useful later in connection with integration over this space, and in considering the corresponding L^2 -space.

About $L^2(G/H)$. Keep the assumptions on G, H, Z_H as above. We extend a unitary character $\chi \in \widehat{Z_H}$ to a character of H and consider the unitary induced representation $\text{Ind}_H^G \chi$ on $L^2(G/H)_{\chi^{-1}}$. Normalize Plancherel measures on Z_H and its dual group in the usual way.

Proposition II.3. *As a representation of G*

$$L^2(G/H_{ss}) = \int_{\chi \in \widehat{Z_H}} L^2(G/H)_{\chi^{-1}} d\chi,$$

Proof. For $f \in L^2(G/H_{ss})$ and $\chi \in \widehat{Z_H}$ define

$$F(\chi, g) = \int_{Z_H} f(gz)\chi(z)^{-1} dz,$$

Then for $z_0 \in Z_H$

$$F(\chi, gz_0) = F(\chi, g)\chi^{-1}(z_0),$$

so $F(\chi) \in L^2(G/H)_{\chi^{-1}}$. By Fourier analysis on Z_H we have

$$\int_{\chi \in \widehat{Z_H}} |F(\chi, g)|^2 d\chi = \int_{z \in Z_H} |f(gz)|^2 dz.$$

So

$$\int_{G/H_{ss}} |f(g')|^2 dg' = \int_{G/H} \int_{Z_H} |f(g'z)|^2 dz dg'$$

completes the proof. \square

III. Main results

In this section we relate the Plancherel formula for the left regular representation of G on $L^2(G/H_{ss})$ to the Plancherel formula for the left regular representation on $L^2(G/H)$. It turns out that these two spaces have the same content of unitary representations of G , only differing by their multiplicities.

Induction to the parabolic subgroup P .

Lemma III.1. *Let \widehat{N} the dual group of N . There exist finitely many open H orbits \mathbb{O}_i in \widehat{N} so that \widehat{N} is the closure of their union $\bigcup_i \mathbb{O}_i$.*

Proof. Here we refer to results by Wallach [2006]. Here he proves that our parabolic algebras are “very nice” since they have abelian nilradicals (see Corollary 6.4 of that reference). In particular there is only one open orbit of L on N .

Since our group H is a subgroup of finite index in L we will get a finite number of open orbits with dense union. Actually, the statement that “open orbit is generic” (i.e., “nice parabolic”) would suffice for our purposes here. \square

Let $\chi \in \widehat{Z}_H$. We consider again χ as a character of H and consider again the unitary induced representation $\text{Ind}_H^P \chi$.

Proposition III.2. *Let χ and $\tilde{\chi}$ be unitary characters of Z_H considered as characters of H . Then we have (equivalence of representations)*

$$\text{Ind}_H^P \chi = \text{Ind}_H^P \tilde{\chi}.$$

Proof. We denote the induced representations acting on functions $F \in L^2(N)$ by

$$\begin{aligned} \rho_\chi(n_0)F(n) &= F(n \cdot n_0), \\ \rho_\chi(h_0)F(n) &= \chi(h_0)F(h_0^{-1}nh_0). \end{aligned}$$

Using the Fourier transform we realize the representation $\text{Ind}_H^P \chi$ on $L^2(\widehat{N})$. It is a direct sum of irreducible representations on $L^2(\mathbb{O}_i)$ where

$$\begin{aligned} \hat{\rho}_\chi(n_0) &\text{ is a multiplication operator,} \\ \hat{\rho}_\chi(h_0)\hat{F}(\xi) &= \chi(h_0)J(h_0^t\xi)^{1/2}\hat{F}(h_0^t\xi). \end{aligned}$$

The other representation on the orbit is obtained by multiplication of the right hand side of the second equation with a character $\chi_1 = \tilde{\chi}\chi^{-1}$ of H . In each orbit we fix an element ξ_i .

We get an intertwining operator on each of the irreducible representations by

$$I(F)(\xi) = \chi_1(\xi)F(\xi).$$

Here $\xi = h\xi_i$ and $\chi_1(\xi) := \chi_1(h)$. \square

Example 2. Consider the group $P = HN$ with

$$H = \left\{ \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \mid a > 0 \right\}$$

and

$$N = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \mid b \in \mathbb{R} \right\}.$$

We note that there are three orbits of H on

$$\hat{N} = \left\{ \xi_t \mid \xi_t \left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \right) = e^{it \cdot b} \right\},$$

namely, $\mathbb{O}^+ = \{\xi_t \mid t > 0\}$, $\mathbb{O}^- = \{\xi_t \mid t < 0\}$ and $\mathbb{O}^1 = \{\xi_0\}$. The unitary representation ρ_1 of P induced from the trivial representation of H acts on $L^2(N)$ by

$$\rho_1 \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) F(x) = a^{1/2} F(ax)$$

and

$$\rho_1 \left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \right) F(x) = F(x + b).$$

To analyze this representation we consider the Fourier transform of $L^2(N)$. The representation is a direct sum of two unitary representations of functions whose Fourier transform has support in $\xi \in \mathbb{O}^+$ and in $\xi \in \mathbb{O}^-$.

We consider $\chi_s : a \rightarrow a^{is}$ as a character of H . After applying the Fourier transform the representation $\hat{\rho}_s$ induced from χ_s has the form

$$\hat{\rho}_s \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) \hat{F}(\xi) = a^{-1/2} a^{is} \hat{F}(a^{-1}\xi)$$

and

$$\hat{\rho}_s \left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \right) \hat{F}(\xi) = e^{ib\xi} \hat{F}(\xi).$$

The equivalence of the representations ρ_s and ρ_1 follows from the intertwining operator

$$\mathcal{I}_s : \rho_0 \rightarrow \rho_s \quad \text{defined by } \mathcal{I}_s \hat{F}(\xi) = \xi^{is} \hat{F}(\xi).$$

Induction to G .

Proposition III.3. *Let χ and $\tilde{\chi}$ be characters of Z_H considered as characters of H . As representations of G we have (equivalence)*

$$\text{Ind}_H^G \chi = \text{Ind}_H^G \tilde{\chi}.$$

Proof. By induction by stages (Proposition III.2) we have

$$\text{Ind}_H^G \chi = \text{Ind}_P^G \text{Ind}_H^P \chi = \text{Ind}_P^G \text{Ind}_H^P \tilde{\chi} = \text{Ind}_H^G \tilde{\chi}. \quad \square$$

Example 3. $G = \text{SU}(1, 1)$ and

$$H = A = \exp \mathbb{R} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

We identify G/K with the complex unit disk \mathcal{D} . We realize the discrete series representations D_n in the holomorphic functions on \mathcal{D} . Then we have the H -invariant distribution vector, giving the embedding into $L^2(G/H)$,

$$v^* = (1 + z^2)^{-n/2} \in D_n^{-\infty, H},$$

and similarly the distribution vector

$$v^* = (1 + z^2)^{-n/2} \left(\frac{1 - z}{1 + z} \right)^{i\lambda} \in D_n^{-\infty, H, \chi_\lambda},$$

transforming by the character χ_λ of H . So indeed every discrete series representation occurs in every $L^2(G/H)_{\chi_\lambda}$.

Theorem III.4. *As a left regular representation of G*

$$L^2(G/H_{ss}) \simeq (\text{Ind}_H^G 1) \otimes L^2(Z_H) \simeq L^2(G/H) \otimes L^2(Z_H).$$

Proof. This follows from Propositions II.3 and III.3. □

Corollary III.5. *All irreducible representations in the discrete spectrum of*

$$L^2(G/H_{ss})$$

have infinite multiplicity.

Definition. Following Benoist and Kobayashi we say that $L^2(G/H_{ss})$ is not tempered if the representations in the Plancherel formula for the right regular representation of G on $L^2(G/H_{ss})$ are not a subset of the representations of the Plancherel formula for G .

Corollary III.6. *$L^2(G/H_{ss})$ is tempered if and only if $L^2(G/H)$ is tempered.*

Example 1 (continued). $G = \text{SL}(2, \mathbb{R})$, H diagonal matrices, Then $X = G/H$ is a hyperboloid and

$$L^2(G/H) = \bigoplus_{\nu \in 2\mathbb{N}} D_\nu \oplus 2 \int_0^\infty \pi_{it},$$

where D_ν are the discrete series representations with parameter ν and π_{it} are the tempered spherical principal series representations with parameter it . Here $H_{ss} = \mathbb{Z}_2$, then $L^2(G/H_{ss}) = L^2(\text{PSL}(2, \mathbb{R}))$ and so the left regular representation contains the even discrete series representations with ∞ multiplicity.

If H is connected, then $L^2(G/H)$ contains all discrete series representations and so does the left regular representation of G on $L^2(G)$.

IV. More examples

We discuss in this section some interesting examples of groups G and H_{ss} , illustrating our results; one aspect is to find reductive spaces that are not tempered.

We use the Plancherel formula to determine if $L^2(G/H_{ss})$ is tempered. Some of our examples are also contained in [Benoist and Kobayashi 2015], where they are obtained with a different technique; others are new.

E. van den Ban and H. Schlichtkrull [2005] proved a Plancherel formula for $L^2(G/H)$ for a fixed point set H of an involution τ of G . They showed that only discrete series representations of $L^2(G/H)$ and principal series representations unitarily induced from a $\theta\tau$ invariant parabolic MAN , a discrete series representation π of $M/M \cap H$ and a unitary character of A contribute to the Plancherel formula. On the other hand the work of M. Flensted-Jensen and Oshima and Matsuki shows that the discrete spectrum of G/H is nontrivial if and only if

$$\text{rank } G/H = \text{rank } K/K \cap H.$$

A parametrization of the representations in the discrete spectrum was obtained by T. Matsuki and T. Oshima [1984]. See also [Schlichtkrull 1983]. We will make extensive use of these results in the proofs of our examples.

Remark 1. Induction by stages enlarges the set of pairs G, \tilde{H} for which $L^2(G/\tilde{H})$ is tempered. (See [Fell 1962, Theorem 4.2]; here the point is that induction preserves weak containment, so if we have groups $H \subset \tilde{H} \subset G$ so that $L^2(G/H)$ is tempered and we know that $L^2(\tilde{H}/H)$ contains the trivial representation weakly, then also $L^2(G/\tilde{H})$ is tempered.)

Remark 2. The nontempered representations in the discrete spectrum of $L^2(G/H_{ss})$ are automorphic representations [Burger and Sarnak 1991]. Most of these automorphic representations are known and have been constructed using other techniques for example in [Kudla and Rallis 1990; Howe and Piatetski-Shapiro 1979; Schlichtkrull 1983; Mœglin and Waldspurger 1989].

Example 4. Let $G = \text{SL}(2n, \mathbb{R})$, We take H as the connected component of $S(\text{GL}(p, \mathbb{R}) \times \text{GL}(q, \mathbb{R}))$. Then $H_{ss} = \text{SL}(p, \mathbb{R}) \times \text{SL}(q, \mathbb{R})$ where $p + q = 2n$. and

$$\text{rank } G/H = \text{rank } K/K \cap H = \min(p, q).$$

The results of van den Ban and Schlichtkrull show that all the representations in the continuous spectrum are unitarily induced from $\theta\tau$ -stable parabolic subgroups. It is easy to see that these parabolic subgroups are all cuspidal and thus the representations in the discrete spectrum determine whether $L^2(\text{SL}(n, \mathbb{R})/H)$ is tempered.

We recall the parametrization of the representations in the discrete spectrum. Using the decomposition $\mathfrak{h} \otimes \mathbb{C} \oplus \mathfrak{q} \otimes \mathbb{C}$ of $\mathfrak{gl}(2n, \mathbb{C})$ we conclude that the skew diagonal matrices in $\mathfrak{q} \otimes \mathbb{C}$ are a maximal abelian subspace of $\mathfrak{so}(2n, \mathbb{C}) \cap \mathfrak{q} \otimes \mathbb{C}$. By [Ōshima and Matsuki 1984] their centralizer L is the Levi subgroup of a θ stable parabolic subgroup. The representations in the discrete spectrum are cohomologically induced from a character of the subgroup L . If the commutator subgroup L does not contain a noncompact semisimple subgroup then the representations are tempered. (For this, see [Knapp and Vogan 1995, Chapter XI] or [Vogan and Zuckerman 1984, Theorem 6.16]). Thus we conclude:

- If $p = q = n$ the subgroup $[L, L]$ is a product of n compact tori. Hence all representation $L^2(\mathrm{SL}(2n, \mathbb{R})/(\mathrm{SL}(n, \mathbb{R}) \times \mathrm{SL}(n, \mathbb{R})))$ in the discrete spectrum are tempered and thus $L^2(\mathrm{SL}(2n, \mathbb{R})/H_{ss})$ is tempered.
- If $p - q \geq 2$ then L has a noncompact subgroup and hence the representations in the discrete spectrum of $L^2(G/H_{ss})$ are the Langlands subquotient of representations which is not unitarily induced. Hence

$$L^2(\mathrm{SL}(2n, \mathbb{R})/(\mathrm{SL}(n, \mathbb{R}) \times \mathrm{SL}(n, \mathbb{R})))$$

is not tempered.

- Using Remark 2 we can construct a large number of additional semisimple subgroups H_{ss} so that $L^2(\mathrm{SL}(2n, \mathbb{R})/H_{ss})$ is tempered.

C. Mœglin and J. L. Waldspurger [1989] show that these representations are in the residual spectrum of a congruence subgroup of $\mathrm{GL}(n, \mathbb{R})$. Similar considerations for general linear groups can be found in [Venkatesh 2005].

Example 5. $G = \mathrm{SO}(p, q)$, $p + q = 2n \geq 4$ with $p \geq q > 2$ and

$$H = \mathrm{SO}(1, 1) \times \mathrm{SO}(p - 1, q - 1) \quad \text{and} \quad H_{ss} = \mathrm{SO}(p - 1, q - 1).$$

Claim. $L^2(\mathrm{SO}(p, q)/\mathrm{SO}(p - 1, q - 1))$ is not tempered.

We have

$$\mathrm{rank} G/H = \mathrm{rank} K/K \cap H = 2.$$

We argue as in Example 2. The group $[L, L]$ has a factor isomorphic to

$$\mathrm{SO}(p - 2, q - 2),$$

and is hence is not compact. So there are nontempered representations in the discrete spectrum.

T. Kobayashi [1992] considered the case G/H_0 where $H = H_c \times H_0$. Here H_c is a compact orthogonal group and H_0 is a noncompact orthogonal group. He determined the parameter of the representations in the discrete spectrum of $L^2(G/H_0)$ and their multiplicities.

Example 6. $G = \mathrm{Sp}(n, \mathbb{R})$, $H = \mathrm{GL}(n, \mathbb{R})$, $H_{ss} = \mathrm{SL}(n, \mathbb{R})$.

Claim. $L^2(\mathrm{Sp}(n, \mathbb{R})/\mathrm{SL}(n, \mathbb{R}))$ is tempered.

The proof proceeds as follows:

Step 1: All the representations in the discrete spectrum are tempered.

Step 2: Each conjugacy class of parabolic subgroups contains a $\theta\tau$ -invariant parabolic subgroup MAN .

Step 3: All discrete series representations of $M/M \cap H$ are tempered.

For simplicity assume that the symplectic group is defined by the quadratic form defined by the matrix

$$\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix},$$

where I is the identity matrix. The subgroup $H = \mathrm{GL}(n, \mathbb{R})$ of G is the fixed point set of the automorphism τ defined by conjugation with

$$\begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix}.$$

The maximal compact subgroup K_H of H is $K \cap H = O(n)$. Furthermore

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p},$$

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}.$$

The one-dimensional torus T_0 in the center of K also defines a torus on $K/H \cap K$. Its Lie algebra \mathfrak{t}_0 is direct summand of the maximal abelian subalgebra \mathfrak{a}_k of $\mathfrak{q}_k = \mathfrak{k} \cap \mathfrak{q}$. Since T_0 defines the complex structure on the symmetric space G/K the centralizer of \mathfrak{a}_k in G is contained in K . Thus every representation in the discrete spectrum is tempered.

The $\theta\tau$ -stable parabolic subalgebras are determined by maximal abelian subspaces \mathfrak{i} in $\mathfrak{p} \cap \mathfrak{q}$. Now

$$\tilde{\mathfrak{h}} := \mathfrak{k} \cap \mathfrak{h} \oplus \mathfrak{p} \cap \mathfrak{q} = \mathfrak{gl}(n, \mathbb{R})$$

is the fixed point set of the involution $\theta\tau$ since the fixed point set of $\theta\tau$ is conjugate in $\mathrm{GL}(2n, \mathbb{R})$ to $\mathrm{GL}(n, \mathbb{R})$. This implies that there is a n -dimensional abelian split subalgebra $\tilde{\mathfrak{a}}_H$ in $\mathfrak{p} \cap \mathfrak{q}$ consisting of the matrices

$$\begin{pmatrix} 0 & D \\ D & 0 \end{pmatrix},$$

where D is a real diagonal matrix. Hence every conjugacy classes of parabolic subgroups contains a $\theta\tau$ -stable parabolic $P_s = M_s A_s N_s$ whose Levi subgroup is a centralizer of $\tilde{\mathfrak{a}}_H$.

Next we have to determine $M_s \cap H$. Note that M_s is a product of general linear groups and a symplectic group. The factors isomorphic to general linear groups are subgroups of \tilde{H} . Since $\tilde{H} \cap H = K \cap H$ is an orthogonal group, the intersection of the general linear subgroups of M_s with H are orthogonal groups and hence the corresponding symmetric space has no discrete spectrum. Thus we may assume that $M_s = \text{Sp}(m, \mathbb{R})$ with $m < n$. In this case $\theta\tau$ is an involution of M_s with fixed points $\tilde{H} \cap M_s = \text{GL}(m, \mathbb{R})$. Furthermore since θ and τ commute their restriction to M_s also defines an automorphisms of M_s . So the fixed point set of $\theta\tau|_{M_s}$ is conjugate to the fixpoint set of $\tau|_{M_s}$ in $\text{GL}(2n, \mathbb{R})$. Hence we conclude that $M_s \cap H$ is isomorphic to $\text{GL}(m, \mathbb{R})$. By Step 1 the representations in the discrete spectrum of $\text{Sp}(m, \mathbb{R})/\text{GL}(m, \mathbb{R})$ are tempered and thus by [van den Ban and Schlichtkrull 2005] all the representations in the continuous spectrum of $\text{Sp}(n, \mathbb{R})/\text{SL}(n, \mathbb{R})$ are tempered.

Example 7. Cayley-type spaces are considered in [Ólafsson and Ørsted 1999; Faraut and Korányi 1994]. These are

- (a) $G = \text{Sp}(n, \mathbb{R})$, $H = \text{GL}(n, \mathbb{R})$ and $H_{ss} = \text{SL}(n, \mathbb{R})$, $n > 1$;
- (b) $G = \text{SO}(2, n)$, $H = \text{SO}(1, 1)\text{SO}(1, n - 1)$ and $H_{ss} = \text{SO}(1, n)$, $n > 2$;
- (c) $G = \text{SU}(n, n)$, $H = \text{SL}(n, \mathbb{C})\mathbb{R}^+$ and $H_{ss} = \text{SL}(n, \mathbb{C})$;
- (d) $G = O^*(2n)$, $H = \mathbb{R}^+ \text{SU}^*(2n)$ and $H_{ss} = \text{SU}^*(2n)$;
- (e) $G = E_{7(-25)}$, $H = E_{6(-26)}\mathbb{R}^+$ and $H_{ss} = E_{6(-26)}$.

Claims.

- In Example 7(b)–(d) with n large enough $L^2(G/H_{ss})$ is not tempered.
- In Example 7(a) $L^2(G/H_{ss})$ is tempered.
- We expect that in Example 7(e) $L^2(G/H_{ss})$ is tempered.

Proof. The proof is based on case by case considerations of the spectrum of $L^2(G/H)$. Ólafsson and Ørsted [1999] proved that all these spaces are of equal rank and hence $L^2(G/H)$ has a discrete spectrum.

Case (b). The arguments in Example 5 show that the representations in the discrete spectrum of $L^2(\text{SO}(n, 2)/\text{SO}(n - 1, 1))$ are tempered if and only if $n \leq 2$. So we can conclude that $L^2(\text{SO}(n, 2)/\text{SO}(n - 1, 1))$ is not tempered if $3 \leq n$.

Case (c). It was proved in [Ólafsson and Ørsted 1988] that the discrete spectrum for $\text{SU}(n, n)/H$ contains some nontempered highest weight representations. Hence $L^2(\text{SU}(n, n)/\text{SL}(n, \mathbb{C}))$ is not tempered.

Case (a). This was proved in Example 6.

Case (d). We have $\text{rank}(G/H) = n$. The Levi of the θ -stable parabolic subgroup also contains a subgroup of type A_{2n-1} . Since it is not the maximal compact

subgroup, L has a noncompact subgroup. This implies that the discrete spectrum of $L^2(O^*(2n)/\mathrm{SU}^*(2n))$ is not tempered.

Case (e). We only prove that the discrete spectrum is tempered. The arguments are the same as in Example 4. Recall that

- (1) the rank of G/H is 3;
- (2) the maximal compact subgroup K of G is $E_6 \mathrm{SO}(2)$;
- (3) the maximal compact subgroup K_H of H is F_4 ;
- (4) $K/H \cap K$ has a one-dimensional compact torus T_0 as factor.

The centralizer of this torus T_0 is K . Its Lie algebra is a direct summand of the maximal abelian subalgebra \mathfrak{a}_k of $\mathfrak{q}_k = \mathfrak{k} \cap \mathfrak{q}$. Since T_0 defines the complex structure on the symmetric space G/K the centralizer of \mathfrak{a}_k in G is contained in K . Thus every representation in the discrete spectrum is tempered.

As in Example 6 we conclude that the fixed point set of $\theta\tau$ is a subgroup isomorphic to $H = E_{6(-26)}\mathbb{R}^+$, which has real rank 3. Hence every conjugacy classes of parabolic subgroups contains a $\theta\tau$ -stable parabolic $P_s = M_s A_s N_s$ whose Levi subgroup is a centralizer of $\tilde{\mathfrak{a}}_H$. \square

Example 8. $G = \mathrm{SL}(2n, \mathbb{C})$ and H_{ss} has a covering $T^1 \mathrm{SL}(p, \mathbb{C}) \times \mathrm{SL}(q, \mathbb{C})$, $p+q = 2n$ for a one dimensional torus T^1 . Then

$$L^2(\mathrm{SL}(n, \mathbb{C})/\mathrm{SL}(p, \mathbb{C}) \times \mathrm{SL}(q, \mathbb{C})) = \bigoplus_{\delta \in \hat{T}} L^2(\mathrm{SL}(n, \mathbb{C})/H_{ss}, \delta),$$

where $L^2(\mathrm{SL}(n, \mathbb{C})/H_{ss}, \delta)$ are the L^2 -sections of the line bundle defined by the character δ of H_{ss} . As in Example 2 we are in the equal rank case.

The same arguments as in Example 4 show:

- If $p = q = n$ the subgroup $[L, L]$ is compact. Hence all representations in the discrete spectrum of $L^2(\mathrm{SL}(2n, \mathbb{C})/H)$ are tempered, which implies that $L^2(\mathrm{SL}(2n, \mathbb{C})/H_{ss})$ is tempered.
- If $p - q \geq 2$ then $[L, L]$ is not compact and hence the representations in the discrete spectrum of $L^2(G/H_{ss})$ are the Langlands subquotients of representations which are not unitarily induced. Hence $L^2(\mathrm{SL}(2n, \mathbb{C})/\mathrm{SL}(n, \mathbb{C}) \times \mathrm{SL}(n, \mathbb{C}))$ is not tempered.

Acknowledgement

Speh would like to thank the Department of Mathematics of the University of Aarhus for their hospitality.

References

- [van den Ban and Schlichtkrull 2005] E. P. van den Ban and H. Schlichtkrull, “The Plancherel decomposition for a reductive symmetric space, II: Representation theory”, *Invent. Math.* **161**:3 (2005), 567–628. MR 2181716 Zbl 1078.22013
- [Benoist and Kobayashi 2015] Y. Benoist and T. Kobayashi, “Temperedness of reductive homogeneous spaces”, *J. Euro. Math. Soc.* **17**:12 (2015), 3015–3036.
- [Burger and Sarnak 1991] M. Burger and P. Sarnak, “Ramanujan duals, II”, *Invent. Math.* **106**:1 (1991), 1–11. MR 1123369 Zbl 0774.11021
- [Faraut and Korányi 1994] J. Faraut and A. Korányi, *Analysis on symmetric cones*, Oxford University Press, New York, NY, 1994. MR 1446489 Zbl 0841.43002
- [Fell 1962] J. M. G. Fell, “Weak containment and induced representations of groups”, *Canad. J. Math.* **14** (1962), 237–268. MR 0150241 Zbl 0138.07301
- [Howe and Piatetski-Shapiro 1979] R. Howe and I. I. Piatetski-Shapiro, “A counterexample to the ‘generalized Ramanujan conjecture’ for (quasi-) split groups”, pp. 315–322 in *Automorphic forms, representations and L-functions* (Corvallis, OR, 1977), edited by A. Borel and W. Casselman, Proceedings of Symposia in Pure Mathematics **33**:1, American Mathematical Society, Providence, RI, 1979. MR 546605 Zbl 0423.22018
- [Knapp and Vogan 1995] A. W. Knap and D. A. Vogan, Jr., *Cohomological induction and unitary representations*, Princeton Mathematical Series **45**, Princeton University Press, 1995. MR 1330919 Zbl 0863.22011
- [Kobayashi 1992] T. Kobayashi, *Singular unitary representations and discrete series for indefinite Stiefel manifolds $U(p, q; \mathbf{F})/U(p - m, q; \mathbf{F})$* , Memoirs of the American Mathematical Society **95**:462, American Mathematical Society, Providence, RI, 1992. MR 1098380 Zbl 0752.22007
- [Kudla and Rallis 1990] S. S. Kudla and S. Rallis, “Degenerate principal series and invariant distributions”, *Israel J. Math.* **69**:1 (1990), 25–45. MR 1046171 Zbl 0708.22005
- [Mœglin and Waldspurger 1989] C. Mœglin and J.-L. Waldspurger, “Le spectre résiduel de $GL(n)$ ”, *Ann. Sci. École Norm. Sup.* (4) **22**:4 (1989), 605–674. MR 1026752 Zbl 0696.10023
- [Ólafsson and Ørsted 1988] G. Ólafsson and B. Ørsted, “The holomorphic discrete series for affine symmetric spaces, I”, *J. Funct. Anal.* **81**:1 (1988), 126–159. MR 967894 Zbl 0678.22008
- [Ólafsson and Ørsted 1999] G. Ólafsson and B. Ørsted, “Causal compactification and Hardy spaces”, *Trans. Amer. Math. Soc.* **351**:9 (1999), 3771–3792. MR 1458309 Zbl 0928.43007
- [Ōshima and Matsuki 1984] T. Ōshima and T. Matsuki, “A description of discrete series for semisimple symmetric spaces”, pp. 331–390 in *Group representations and systems of differential equations* (Tokyo, 1982), edited by K. Okamoto, Advanced Studies in Pure Mathematics **4**, North-Holland, Amsterdam, 1984. MR 810636 Zbl 0577.22012
- [Sakellaridis and Venkatesh 2014] Y. Sakellaridis and A. Venkatesh, “Periods and harmonic analysis on spherical varieties”, preprint, 2014. arXiv 1203.0039v3
- [Schlichtkrull 1983] H. Schlichtkrull, “The Langlands parameters of Flensted-Jensen’s discrete series for semisimple symmetric spaces”, *J. Funct. Anal.* **50**:2 (1983), 133–150. MR 693225 Zbl 0507.22013
- [Venkatesh 2005] A. Venkatesh, “The Burger–Sarnak method and operations on the unitary dual of $GL(n)$ ”, *Represent. Theory* **9** (2005), 268–286. MR 2133760 Zbl 1077.22022
- [Vogan and Zuckerman 1984] D. A. Vogan, Jr. and G. J. Zuckerman, “Unitary representations with nonzero cohomology”, *Compositio Math.* **53**:1 (1984), 51–90. MR 762307 Zbl 0692.22008

[Wallach 2006] N. R. Wallach, “Holomorphic continuation of generalized Jacquet integrals for degenerate principal series”, *Represent. Theory* **10** (2006), 380–398. MR 2266697 Zbl 1135.22002

Received December 10, 2014. Revised July 14, 2015.

BENT ØRSTED
DEPARTMENT OF MATHEMATICS
AARHUS UNIVERSITY
NY MUNKEGADE
DK-8000 AARHUS C
DENMARK
orsted@math.au.dk

BIRGIT SPEH
DEPARTMENT OF MATHEMATICS
CORNELL UNIVERSITY
MALOTT HALL
ITHACA, NY 14853-4201
UNITED STATES
speh@math.cornell.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$/year for the electronic version, and \$/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 283 No. 1 July 2016

A New family of simple $\mathfrak{gl}_{2n}(\mathbb{C})$ -modules	1
JONATHAN NILSSON	
Derived categories of representations of small categories over commutative noetherian rings	21
BENJAMIN ANTIEAU and GREG STEVENSON	
Vector bundles over a real elliptic curve	43
INDRANIL BISWAS and FLORENT SCHAFFHAUSER	
$\mathbb{Q}(\mathbb{N})$ -graded Lie superalgebras arising from fermionic-bosonic representations	63
JIN CHENG	
Conjugacy and element-conjugacy of homomorphisms of compact Lie groups	75
YINGJUE FANG, GANG HAN and BINYONG SUN	
Entire sign-changing solutions with finite energy to the fractional Yamabe equation	85
DANILO GARRIDO and MONICA MUSSO	
Calculation of local formal Mellin transforms	115
ADAM GRAHAM-SQUIRE	
The untwisting number of a knot	139
KENAN INCE	
A Plancherel formula for $L^2(G/H)$ for almost symmetric subgroups	157
BENT ØRSTED and BIRGIT SPEH	
Multiplicative reduction and the cyclotomic main conjecture for GL_2	171
CHRISTOPHER SKINNER	
Commensurators of solvable S -arithmetic groups	201
DANIEL STUDENMUND	
Gerstenhaber brackets on Hochschild cohomology of quantum symmetric algebras and their group extensions	223
SARAH WITHERSPOON and GUODONG ZHOU	



0030-8730(201607)283:1;1-X