

*Pacific
Journal of
Mathematics*

DISCRIMINANTS AND THE MONOID OF QUADRATIC RINGS

JOHN VOIGHT

Volume 283 No. 2

August 2016

DISCRIMINANTS AND THE MONOID OF QUADRATIC RINGS

JOHN VOIGHT

We consider the natural monoid structure on the set of quadratic rings over an arbitrary base scheme and characterize this monoid in terms of discriminants.


Quadratic field extensions K of \mathbb{Q} are characterized by their discriminants. Indeed, there is a bijection

$$\left\{ \begin{array}{l} \text{separable quadratic algebras} \\ \text{over } \mathbb{Q} \text{ up to isomorphism} \end{array} \right\} \xrightarrow{\sim} \mathbb{Q}^\times / \mathbb{Q}^{\times 2}, \quad \mathbb{Q}[\sqrt{d}] = \mathbb{Q}[x]/(x^2 - d) \mapsto d\mathbb{Q}^{\times 2},$$

where a separable quadratic algebra over \mathbb{Q} is either a quadratic field extension or the algebra $\mathbb{Q}[\sqrt{1}] \simeq \mathbb{Q} \times \mathbb{Q}$ of discriminant 1. In particular, the set of isomorphism classes of separable quadratic extensions of \mathbb{Q} can be given the structure of an elementary abelian 2-group, with identity element the class of $\mathbb{Q} \times \mathbb{Q}$: we have, simply,

$$\mathbb{Q}[\sqrt{d_1}] * \mathbb{Q}[\sqrt{d_2}] = \mathbb{Q}[\sqrt{d_1 d_2}]$$

up to isomorphism. If $d_1, d_2, d_1 d_2 \in \mathbb{Q}^\times \setminus \mathbb{Q}^{\times 2}$ then $\mathbb{Q}(\sqrt{d_1 d_2})$ sits as the third quadratic subfield of the compositum $\mathbb{Q}(\sqrt{d_1}, \sqrt{d_2})$:



Indeed, if σ_1 is the nontrivial element of $\text{Gal}(\mathbb{Q}(\sqrt{d_1})/\mathbb{Q})$, then there is a unique extension of σ_1 to $\mathbb{Q}(\sqrt{d_1}, \sqrt{d_2})$ leaving $\mathbb{Q}(\sqrt{d_2})$ fixed, similarly with σ_2 , and $\mathbb{Q}(\sqrt{d_1 d_2})$ is the fixed field of the composition $\sigma_1 \sigma_2 = \sigma_2 \sigma_1$.

This characterization of quadratic extensions works over any base field F with $\text{char } F \neq 2$ and is summarized concisely in the Kummer theory isomorphism

$$H^1(\text{Gal}(\bar{F}/F), \{\pm 1\}) = \text{Hom}(\text{Gal}(\bar{F}/F), \{\pm 1\}) \simeq F^\times / F^{\times 2}.$$

MSC2010: primary 11E99; secondary 11R11, 11R29.

Keywords: quadratic forms, quadratic rings, monoids, discriminants, discriminant algebras.

On the other hand, over a field F with $\text{char } F = 2$, all separable quadratic extensions have trivial discriminant and instead they are classified by the (additive) Artin–Schreier group

$$F/\wp(F), \quad \text{where } \wp(F) = \{r + r^2 : r \in F\},$$

with the class of $a \in F$ in correspondence with the isomorphism class of the extension $F[x]/(x^2 - x + a)$. By similar considerations as above, we again find a natural structure of an elementary abelian 2-group on the set of isomorphism classes of separable quadratic extensions of F .

One can extend this correspondence between quadratic extensions and discriminants integrally, as follows: Let R be a commutative ring. An R -algebra is a ring B equipped with an embedding $R \hookrightarrow B$ of rings (mapping $1 \in R$ to $1 \in B$) whose image lies in the center of B ; we identify R with its image via this embedding. A *free quadratic R -algebra* (also called a *free quadratic ring over R*) is an R -algebra S (associative with 1) that is free of rank 2 as an R -module. Let S be a free quadratic R -algebra. Then $S/R \simeq \wedge^2 S \simeq R$ is projective, so there is an R -basis $1, x$ for S ; we find that $x^2 = tx - n$ for some $t, n \in R$ and that S is commutative. The map $\sigma : S \rightarrow S$ induced by $x \mapsto t - x$ is the unique *standard involution* on S , an R -linear (anti)automorphism such that $y\sigma(y) \in R$ for all $y \in S$. The class in $R/R^{\times 2}$ of the *discriminant* of S ,

$$d = d(S) = (x - \sigma(x))^2 = t^2 - 4n,$$

is independent of the choice of basis $1, x$. A discriminant d satisfies the congruence $d \equiv t^2 \pmod{4R}$, so for example if $R = \mathbb{Z}$ then $d \equiv 0, 1 \pmod{4}$.

Now suppose that R is an integrally closed domain of characteristic not 2. Then there is a bijection

$$\left\{ \begin{array}{l} \text{free quadratic rings over } R \\ \text{up to isomorphism} \end{array} \right\} \xrightarrow{\simeq} \{d \in R : d \text{ is a square in } R/4R\}/R^{\times 2},$$

$$S \mapsto d(S).$$

For example, over $R = \mathbb{Z}$, the free quadratic ring $S(d)$ over \mathbb{Z} of discriminant $d \in \mathbb{Z} = \mathbb{Z}/\mathbb{Z}^{\times 2}$ with $d \equiv 0, 1 \pmod{4}$ is given by

$$S(d) = \begin{cases} \mathbb{Z}[x]/(x^2) \hookrightarrow \mathbb{Q}[x]/(x^2) & \text{if } d = 0; \\ \mathbb{Z}[x]/(x^2 - \sqrt{d}x) \hookrightarrow \mathbb{Q} \times \mathbb{Q} & \text{if } d \neq 0 \text{ is a square;} \\ \mathbb{Z}[(d + \sqrt{d})/2] \hookrightarrow \mathbb{Q}(\sqrt{d}) & \text{otherwise.} \end{cases}$$

The set of discriminants under multiplication has the structure of a *commutative monoid*, a nonempty set equipped with a commutative binary operation and identity element. Hence, so does the set of isomorphism classes of free quadratic R -algebras, an operation we denote by $*$: the identity element is the class of

$R \times R \simeq R[x]/(x^2 - x)$. The class of the ring $R[x]/(x^2)$ with discriminant 0 is called an *absorbing element*.

More generally, a *quadratic R -algebra* or *quadratic ring over R* is an R -algebra S that is locally free of rank 2 as an R -module. By definition, a quadratic ring over R localized at any prime (or maximal) ideal of R is a free quadratic R -algebra. Being true locally, a quadratic R -algebra S is commutative and has a unique standard involution.

There is a natural description of quadratic R -algebras as a stack quotient, as follows: A free quadratic R -algebra equipped with a basis $1, x$ has multiplication table uniquely determined by $t, n \in R$ and has no automorphisms, so the functor that associates to a commutative ring R the set of free quadratic R -algebras with basis (up to isomorphism) is represented by two-dimensional affine space \mathbb{A}^2 (over \mathbb{Z}). The change of basis for a free quadratic R -algebra is of the form $x \mapsto u(x + r)$ with $u \in R^\times$ and $r \in R$, mapping

$$(t, n) \mapsto (u(t + 2r), u^2(n + tr + r^2)).$$

Therefore, we have a map from the set of free quadratic R -algebras with basis to the quotient of $\mathbb{A}^2(R)$ by $G(R) = (\mathbb{G}_m \times \mathbb{G}_a)(R)$ with the above action. Working over $\text{Spec } \mathbb{Z}$, the group scheme G is naturally a subgroup scheme of GL_2 , but it does not act linearly on \mathbb{A}^2 , and the Artin stack $[\mathbb{A}^2/G]$ has dimension zero over $\text{Spec } \mathbb{Z}$! Nevertheless, the set $[\mathbb{A}^2/G](R)$ is in bijection with the set of quadratic R -algebras up to isomorphism.

Recall that a commutative R -algebra S is *separable* if S is (faithfully) projective as an $S \otimes_R S$ -module via the map $x \otimes y \mapsto xy$. A free quadratic R -algebra S is separable if and only if $d(S) \in R^\times$; so, for example, the only separable (free) R -algebra over \mathbb{Z} is the ring $\mathbb{Z} \times \mathbb{Z}$ of discriminant 1, an impoverishment indeed! A separable quadratic R -algebra S is étale over R , and R is equal to the fixed subring of the standard involution of S over R . (In many contexts, one then says that S is Galois over R with Galois group $\mathbb{Z}/2\mathbb{Z}$, though authors differ on precise terminology. See Lenstra [2008] for one approach to Galois theory for schemes.) If S and T are separable free quadratic R -algebras, where R is a Dedekind domain of characteristic not 2, having standard involutions σ and τ , respectively, then the monoid product $S * T$ defined above (by transporting the monoid structure on the set of discriminants) is the fixed subring of $S \otimes_R T$ by $\sigma \otimes \tau$, in analogy with the case of fields.

The characterization of free quadratic R -algebras by their discriminants is an example of the parametrization of algebraic structures, corresponding to the Lie group A_1 in the language of Bhargava [2002] (perhaps indexed by A_0 in the schema of Knus, Ojanguren, Parimala and Tignol [Knus et al. 1998, §15]).

Results in this direction go back to Gauss's composition law for binary quadratic forms and have been extended in recent years by Bhargava [2006], Wood [2011] and

others. Indeed, several authors have considered the case of quadratic R -algebras, including Kanzaki [1973] and Small [1972]. In this article, we consider a very general instance of this monoidal correspondence between quadratic R -algebras and discriminants over an arbitrary base scheme.

Let X be a scheme. A *quadratic \mathcal{O}_X -algebra* is a coherent sheaf \mathcal{S} of \mathcal{O}_X -algebras that is locally free of rank 2 as a sheaf of \mathcal{O}_X -modules. Equivalently, a quadratic \mathcal{O}_X -algebra is specified by a finite, locally free morphism of schemes $\phi : Y \rightarrow X$ of degree 2 (sometimes called a *double cover*): the sheaf $\phi_*\mathcal{O}_Y$ is a sheaf of \mathcal{O}_X -algebras that is locally free of rank 2. If $f : X \rightarrow Z$ is a morphism of schemes and \mathcal{S} is a quadratic \mathcal{O}_Z -algebra, then the pullback $f^*\mathcal{S}$ is a quadratic \mathcal{O}_X -algebra. Let $\text{Quad}(X)$ denote the set of isomorphism classes of quadratic \mathcal{O}_X -algebras and, for an invertible \mathcal{O}_X -module \mathcal{L} , let $\text{Quad}(X; \mathcal{L}) \subseteq \text{Quad}(X)$ be the subset of those algebras \mathcal{S} such that there exists an isomorphism $\bigwedge^2 \mathcal{S} \simeq \mathcal{L}$ of \mathcal{O}_X -modules.

Our first result provides an axiomatic description of the monoid structure on the set $\text{Quad}(X)$ (see Theorem 3.27).

Theorem A. *There is a unique system of binary operations*

$$*_X : \text{Quad}(X) \times \text{Quad}(X) \rightarrow \text{Quad}(X),$$

one for each scheme X , such that:

- (i) $\text{Quad}(X)$ is a commutative monoid under $*_X$, with identity element the isomorphism class of $\mathcal{O}_X \times \mathcal{O}_X$.
- (ii) The association $X \mapsto (\text{Quad}(X), *_X)$ from schemes to commutative monoids is functorial in X : for each morphism $f : X \rightarrow Z$ of schemes, the diagram

$$\begin{array}{ccc} \text{Quad}(Z) \times \text{Quad}(Z) & \xrightarrow{*_Z} & \text{Quad}(Z) \\ \downarrow & & \downarrow f^* \\ \text{Quad}(X) \times \text{Quad}(X) & \xrightarrow{*_X} & \text{Quad}(X) \end{array}$$

is commutative.

- (iii) If $X = \text{Spec } R$ and S and T are separable quadratic R -algebras with standard involutions σ and τ , then $S *_X T$ is the fixed subring of $S \otimes_R T$ under $\sigma \otimes \tau$.

The binary operation is defined locally (Construction 3.14): if $X = \text{Spec } R$, and $S = R \oplus Rx$ and $T = R \oplus Ry$ are free quadratic R -algebras with $x^2 = tx - n$ and $y^2 = sy - m$, then we define the free quadratic R -algebra

$$S *_X T = R \oplus Rw,$$

where

$$w^2 = (st)w - (mt^2 + ns^2 - 4nm).$$

This explicit description (in the free case over an affine base) is given by Hahn [1994, Exercises 14–20, pp. 42–43].

A general investigation of the monoid structure on quadratic algebras goes back at least to Loos [1996], who gives via a universal construction a tensor product on the larger category of unital quadratic forms (quadratic forms representing 1); this category is equivalent to the category of quadratic algebras for forms on a finitely generated, projective module of rank 2 [Loos 1996, Proposition 1.6] as long as one takes morphisms as isomorphisms in the category [Loos 2007, §1.4]. (See also §6.1 of Loos [2007] for further treatment.) The existence of the monoid structure was also established in an unpublished letter of Deligne to Rost and Bhargava (March 2, 2005) by a different method: he associates to every R -algebra its discriminant algebra (a quadratic algebra) and extends the natural operation of addition of $\mathbb{Z}/2\mathbb{Z}$ -torsors from the étale case to the general case by geometric arguments. Our proof of Theorem A carries the same feel as these results, but it is accomplished in a more direct fashion and gives a characterization (in particular, uniqueness).

Recently, there has been renewed interest in the construction of discriminant algebras (sending an R -algebra A of rank n to a quadratic R -algebra) by Loos [2007], Rost [2002] and Biesel and Gioia [2015]. Indeed, Biesel and Gioia [2015, Section 8] describe the monoid operation in Theorem A over an affine base in the context of discriminant algebras. We hope that our theorem will have some applications in this context.

Our second result characterizes quadratic algebras in terms of discriminants. A *discriminant* (over X) is a pair (d, \mathcal{L}) such that \mathcal{L} is an invertible \mathcal{O}_X -module and $d \in (\mathcal{L}^\vee)^{\otimes 2}(X)$ is a global section that is a *square modulo 4*: there exists a global section $\bar{t} \in \mathcal{L}^\vee(X)/2\mathcal{L}^\vee(X) = \mathcal{L}^\vee \otimes \mathcal{O}_X/2\mathcal{O}_X$ such that

$$\bar{t} \otimes \bar{t} = \bar{d} \in (\mathcal{L}^\vee)^{\otimes 2}(X)/4(\mathcal{L}^\vee)^{\otimes 2}(X).$$

We can of course also think of $d \in (\mathcal{L}^\vee)^{\otimes 2}(X)$ as an \mathcal{O}_X -module homomorphism $d : \mathcal{L}^{\otimes 2} \rightarrow \mathcal{O}_X$; but such a global section is also equivalently given by a quadratic form $D : \mathcal{L} \rightarrow \mathcal{O}_X$ (see Section 2).

An isomorphism of discriminants $(d, \mathcal{L}), (d', \mathcal{L}')$ is an isomorphism $f : \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$ such that $(f^\vee)^{\otimes 2}(d') = d$. For example, if $X = \text{Spec } R$ for R a commutative ring and $\mathcal{L} = \mathcal{O}_X = \tilde{R}$, then, as above, a discriminant is specified by an element $d \in R$ such that $d \equiv t^2 \pmod{4R}$ for some $t \in R$ (noting that only $t \in R/2R$ matters), and two discriminants d, d' are isomorphic if and only if there exists $u \in R^\times$ such that $u^2 d' = d$. Thinking of a discriminant as a quadratic form $D : \mathcal{L} \rightarrow \mathcal{O}_X$, its image generates a locally principal ideal sheaf $\mathcal{I} \subseteq \mathcal{O}_X$, and the set of discriminants with given locally free image $\mathcal{I} \subseteq \mathcal{O}_X$, if nonempty, is a principal homogeneous space for the group $\mathcal{O}_X^\times / \mathcal{O}_X^{\times 2}$. Let $\text{Disc}(X)$ denote the set of isomorphism classes of

discriminants and $\text{Disc}(X; \mathcal{L}) \subseteq \text{Disc}(X)$ the subset with underlying line bundle \mathcal{L} . Then the tensor product

$$(d, \mathcal{L}) * (d', \mathcal{L}') = (d \otimes d', \mathcal{L} \otimes \mathcal{L}')$$

gives $\text{Disc}(X)$ and $\text{Disc}(X; \mathcal{O}_X)$ the structure of a commutative monoid with identity element the class of $(1, \mathcal{O}_X)$.

A quadratic \mathcal{O}_X -algebra \mathcal{S} has a discriminant $\text{disc}(\mathcal{S}) = (d(\mathcal{S}), \wedge^2 \mathcal{S})$, defined by

$$d(\mathcal{S}) : (\wedge^2 \mathcal{S})^{\otimes 2} \rightarrow \mathcal{O}_X,$$

$$(x \wedge y) \otimes (z \wedge w) \mapsto (x\sigma(y) - \sigma(x)y)(z\sigma(w) - \sigma(z)w),$$

where σ is the unique standard involution on \mathcal{S} . Although, a priori, the codomain of $d(\mathcal{S})$ is \mathcal{S} , in fact its image lies in \mathcal{O}_X : if $X = \text{Spec } R$ and $\mathcal{S} = \text{Spec } S$ and S is free with basis $1, x$, then $(\wedge^2 S)^{\otimes 2}$ is freely generated by $(1 \wedge x) \otimes (1 \wedge x)$ and

$$1 \wedge x \otimes 1 \wedge x \mapsto (x - \sigma(x))^2 \in R.$$

We have a natural forgetful map $\text{Disc}(X) \rightarrow \text{Pic}(X)$, where (d, \mathcal{L}) maps to the isomorphism class of \mathcal{L} .

We say a sequence $A \xrightarrow{f} B \xrightarrow{g} C$ of commutative monoids is *exact* if f is injective, g is surjective, and for all $z, w \in B$ we have

$$g(z) = g(w) \iff xz = yw \text{ for some } x, y \in A;$$

equivalently, the sequence is exact if and only if f is injective and g induces an isomorphism of monoids $B/f(A) \simeq C$. (For a review of monoids, see Section 1.)

We now describe the monoid $\text{Quad}(X)$. We begin with the statement that the forgetful map is compatible with the discriminant map, as follows.

Theorem B. *Let X be a scheme. Then the diagram of commutative monoids*

$$\begin{array}{ccccc} \text{Quad}(X; \mathcal{O}_X) & \longrightarrow & \text{Quad}(X) & \xrightarrow{\wedge^2} & \text{Pic}(X) \\ \downarrow \text{disc} & & \downarrow \text{disc} & & \parallel \\ \text{Disc}(X; \mathcal{O}_X) & \longrightarrow & \text{Disc}(X) & \longrightarrow & \text{Pic}(X) \end{array}$$

is functorial and commutative with exact rows and Zariski locally surjective columns.

By “Zariski locally surjective columns”, we mean that there is an (affine) open cover of X where (under pullback) the columns are surjective. (Considering the corresponding sheaves over X , we also obtain a surjective map of sheaves; see Theorem 3.28.)

We now turn to describe the morphism $\text{Quad}(X; \mathcal{O}_X) \rightarrow \text{Disc}(X; \mathcal{O}_X)$. For this purpose, we work locally and assume $X = \text{Spec } R$ for a commutative ring R ; we

abbreviate $\text{Quad}(\text{Spec } R) = \text{Quad}(R)$ and $\text{Quad}(\text{Spec } R; \mathcal{O}_{\text{Spec } R}) = \text{Quad}(R; R)$, and similarly with discriminants.

We would like to be able to fit the surjective map $\text{Quad}(R) \xrightarrow{\text{disc}} \text{Disc}(R)$ of monoids into an exact sequence by identifying its kernel, but unfortunately the fibers of this map vary over the codomain. Instead, we will describe the action of a subgroup of $\text{Quad}(R)$ on the fibers of the map disc ; this is a natural generalization, as the fibers of a group homomorphism are principal homogeneous spaces for the kernel K and are noncanonically isomorphic as a K -set to K with the regular representation.

Recalling the case of quadratic extensions of a field F with $\text{char } F = 2$, for a commutative ring R we define the *Artin–Schreier group* $\text{AS}(R)$ to be the additive quotient

$$\text{AS}(R) = \frac{R[4]}{\wp(R)[4]}, \quad \text{where } \wp(R)[4] = \{n = r + r^2 \in R : r \in R\} \cap R[4]$$

and $R[4] = \{a \in R : 4a = 0\}$. We have a map $i : \text{AS}(R) \rightarrow \text{Quad}(R; R) \hookrightarrow \text{Quad}(R)$ sending the class of $n \in \text{AS}(R)$ to the isomorphism class of the algebra $S = R[x]/(x^2 - x + n)$. The group $\text{AS}(R)$ is an elementary abelian 2-group since $2R[4] \subseteq \wp(R)$.

Our next main result is as follows (see Theorem 4.3).

Theorem C. *The fibers of the map $\text{disc} : \text{Quad}(R) \rightarrow \text{Disc}(R)$ have a unique action of the group $\text{AS}(R)$ compatible with the inclusion of monoids $\text{AS}(R) \hookrightarrow \text{Quad}(R)$. Moreover, the kernel of this action on the fiber $\text{disc}^{-1}(dR^{\times 2})$ contains $\text{ann}_R(d)[4]$.*

Roughly speaking, Theorems B and C together say that “a quadratic algebra is determined by its Steinitz class and its discriminant, locally up to an Artin–Schreier extension”. These theorems could be rephrased in terms of the Grothendieck group; however, due to the existence of an absorbing element, the group $K_0(\text{Quad}(X))$ is trivial for all schemes X .

The article is organized as follows. In Section 1, we briefly review the relevant notions from monoid theory. In Section 2, we consider the monoid of discriminants; in Section 3 we define the monoid of quadratic R -algebras and prove Theorems A and B. In Section 4 we prove Theorem C.

1. Monoids

To begin, we review standard terminology for monoids. A reference for the material in this section is Bergman [2015, Chapter 3]; more generally, see Burris and Sankappanavar [2012] and McKenzie, McNulty and Taylor [McKenzie et al. 1987].

A *semigroup* is a nonempty set A equipped with an associative binary operation

$$* : A \times A \rightarrow A.$$

A *monoid* is a semigroup with identity element 1 for $*$ (necessarily unique). Any semigroup without 1 can be augmented to a monoid. Natural examples of monoids abound: the natural numbers $\mathbb{N} = \mathbb{Z}_{\geq 0}$ under addition, a ring R under its multiplication, and the set of endomorphisms of an algebraic object (such as a variety) under composition. A *group* is a monoid equipped with an inverse map $^{-1} : A \rightarrow A$.

Let A be a semigroup. We say A is *commutative* if $xy = yx$ for all $x, y \in A$. An *absorbing element* of A is an element $0 \in A$ such that $0x = x0 = 0$ for all $x \in A$; a monoid has at most one absorbing element. Multiplicative notation for A will in general be more natural for us; however, we will occasionally write A additively with operation $+$, in which case the identity element will be denoted 0 and, to avoid confusion, A will have no absorbing element. An element $x \in A$ is (left) *cancellative* if $xy = xz$ implies $y = z$ for all $y, z \in A$.

A *homomorphism* of semigroups is a map $f : A \rightarrow B$ such that $f(xy) = f(x)f(y)$ for all $x, y \in A$, and a homomorphism of monoids is a homomorphism of semigroups such that $f(1_A) = 1_B$.

Let $f : A \rightarrow B$ be a homomorphism of monoids. Unlike groups, the kernel $\ker f = \{x \in A : f(x) = 1\}$ of a monoid homomorphism does not determine the structure of the image of f ; instead, we define the *kernel congruence* of f by

$$K_f = \{(x, y) : f(x) = f(y)\} \subseteq A \times A.$$

The set K_f defines a *congruence* on A , an equivalence relation compatible with the operation on A , i.e., if $(x, y), (z, w) \in K_f$ then $(xz, yw) \in K_f$. Conversely, given a congruence K on a monoid A , the set operation $[x] \cdot [y] = [x \cdot y]$ on equivalence classes $[x], [y] \in A/K$ is well-defined and the quotient map $A \rightarrow A/K$ via $x \mapsto [x]$ is a surjective homomorphism of monoids with kernel K ; any homomorphism $f : A \rightarrow B$ with $K_f \supseteq K$ factors through $A \rightarrow A/K$.

The image $f(A) = \{f(x) : x \in A\}$ is a submonoid of B , but if A and B are noncommutative then not every submonoid is eligible to be the kernel of a homomorphism (just as not every subgroup is normal). As we will be interested only in commutative monoids, and this assumption simplifies the presentation, suppose from now on that A and B are commutative. Then the set

$$I_f = \{(z, w) : f(x)z = f(y)w \text{ for some } x, y \in A\} \subseteq B \times B$$

is a congruence called the *image congruence*. (Without the hypothesis of commutativity, I_f is a relation that is reflexive and symmetric, but not necessarily transitive nor a congruence; if A and B are possibly nonabelian groups, then I_f is transitive and is a congruence if and only if $f(A)$ is a normal subgroup of B .) Note that if $0 \in f(A)$ then $I_f = B \times B$. We write $B/f(A) = B/I_f$.

A sequence

$$(1.1) \quad A \xrightarrow{f} B \xrightarrow{g} C$$

is *exact* if f is injective, g is surjective, and $K_g = I_f$, i.e.,

$$g(z) = g(w) \iff xz = yw \text{ for some } x, y \in A;$$

equivalently, (1.1) is exact if f is injective and g induces an isomorphism $B/f(A) = B/I_f \xrightarrow{\sim} C$. A sequence of groups (1.1) is exact as a sequence of groups if and only if it is exact as a sequence of monoids.

Remark 1.2. We will not make use of long exact sequences of monoids here, nor write the customary 0 or 1 at the ends of our short exact sequences. Indeed, the straightforward extension of the notion from groups to monoids using the definitions above (kernel congruence equals image congruence) has a defect: the sequence

$$\mathbb{N} \xrightarrow{f} \mathbb{Z} \xrightarrow{j} 0$$

of monoids under addition has $I_f = \mathbb{Z} \times \mathbb{Z} = K_j$ even though f is not surjective. (The map f is, however, an epimorphism in the category of monoids.)

We will also make use of sheaves of monoids over a scheme X . A sequence $\mathcal{A} \xrightarrow{f} \mathcal{B} \xrightarrow{g} \mathcal{C}$ of sheaves of monoids is *exact* if the sheaf associated to the presheaf $U \mapsto \mathcal{B}(U)/f(\mathcal{A}(U))$ is isomorphic to \mathcal{C} , or equivalently if the induced sequence $\mathcal{A}_x \xrightarrow{f_x} \mathcal{B}_x \xrightarrow{g_x} \mathcal{C}_x$ of monoid stalks is exact for all $x \in X$.

Like the formation of the integers from the natural numbers, one can construct the Grothendieck group $K_0(A)$ of a commutative monoid A , with the universal property that, for any monoid homomorphism $A \rightarrow G$ with G an abelian group, there exists a unique group homomorphism $K_0(A) \rightarrow G$ such that the diagram

$$\begin{array}{ccc} A & \longrightarrow & G \\ \downarrow & \exists! \nearrow & \uparrow \\ K_0(A) & & \end{array}$$

commutes. The group $K_0(A)$ is constructed as $A \times A$ under the equivalence relation $(x, x') \sim (y, y')$ if there exists $z \in A$ such that $xy'z = x'yz$. Note that if A has an absorbing element 0 then $K_0(A) = \{0\}$. The set of cancellative elements A_{canc} is the largest submonoid of A that can be embedded in a group, and the smallest such containing group is the Grothendieck group $K_0(A_{\text{canc}})$.

2. Discriminants

In this section, we define discriminants for quadratic rings over general schemes (Definition 2.11); for a discussion of discriminant modules overlapping the one

presented here, see Knus [1991, §III.3] and Loos [1996, §1.2]. We also relate seminondegenerate quadratic forms on line bundles by their images (Lemma 2.8) and factor the monoid of discriminants over the Picard group (Proposition 2.18).

Let X be a scheme. A *quadratic form* over X is a pair (\mathcal{M}, Q) where \mathcal{M} is a locally free \mathcal{O}_X -module of finite rank and $Q : \mathcal{M} \rightarrow \mathcal{O}_X$ is a *quadratic map*, i.e., for all open sets $U \subseteq X$ we have:

- (i) $Q(rx) = r^2Q(x)$ for all $r \in \mathcal{O}_X(U)$ and $x \in \mathcal{M}(U)$.
- (ii) The map $T : \mathcal{M}(U) \times \mathcal{M}(U) \rightarrow \mathcal{O}_X(U)$ defined by

$$T(x, y) = Q(x + y) - Q(x) - Q(y)$$

is $\mathcal{O}_X(U)$ -bilinear; we call T the *associated bilinear form*.

An *isometry* between quadratic forms $Q : \mathcal{M} \rightarrow \mathcal{O}_X$ and $Q' : \mathcal{M}' \rightarrow \mathcal{O}_X$ is an \mathcal{O}_X -module isomorphism $f : \mathcal{M} \xrightarrow{\sim} \mathcal{M}'$ such that $Q' \circ f = Q$. A *similarity* between quadratic forms Q and Q' is a commutative square:

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{Q} & \mathcal{O}_X \\ \wr \downarrow f & & g \downarrow \wr \\ \mathcal{M}' & \xrightarrow{Q'} & \mathcal{O}_X \end{array}$$

and so an isometry is just a similarity with $g = \text{id}$.

A quadratic form (\mathcal{M}, Q) is also equivalently specified by \mathcal{M} and an \mathcal{O}_X -module homomorphism $Q : \text{Sym}_2 \mathcal{M} \rightarrow \mathcal{O}_X$, or a global section

$$Q \in \text{Hom}(\text{Sym}_2 \mathcal{M}, \mathcal{O}_X) = (\text{Sym}_2 \mathcal{M})^\vee(X) \simeq \text{Sym}^2(\mathcal{M}^\vee)(X).$$

(Here, $\text{Sym}^2 \mathcal{M}$ denotes the second symmetric power of \mathcal{M} and $\text{Sym}_2 \mathcal{M}$ the submodule of symmetric second tensors of \mathcal{M} .)

A quadratic form $Q : \mathcal{M} \rightarrow \mathcal{O}_X$ with associated bilinear form $T : \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{O}_X$ induces a homomorphism of \mathcal{O}_X -modules $\mathcal{M} \rightarrow \mathcal{M}^\vee$ defined by

$$\mathcal{M}(U) \rightarrow \mathcal{M}^\vee(U), \quad y \mapsto (x \mapsto T(x, y)).$$

Following Knus [1991, (I.3.2)], we say $Q : \mathcal{M} \rightarrow \mathcal{O}_X$ is *nondegenerate* if the associated map $\mathcal{M} \rightarrow \mathcal{M}^\vee$ is injective and *nonsingular* (or *regular*) if the associated map $\mathcal{M} \rightarrow \mathcal{M}^\vee$ is an isomorphism; these properties hold for Q if and only if they hold on an affine open cover. On an open set $U = \text{Spec } R$ where $\mathcal{M}(U) = M$ is free of rank n , we define the *discriminant* of Q as $\text{disc}(Q) = \det(T) \in R/R^{\times 2}$, the determinant of the bilinear form T with respect to a basis of $M \simeq R^n$; then Q is nondegenerate if and only if $\text{disc}(Q)$ is a nonzerodivisor and nonsingular if and only if $\text{disc}(Q)$ is a unit in R . When, further, n is odd, we define the *half-discriminant* (see, e.g., Knus [1991, (IV.3.1.3)]) by a universal formula, and we say that Q is

seminondegnerate (resp. *seminonsingular* or *semiregular*) if the half-discriminant is a nonzerodivisor (resp. a unit), and we extend these notions globally to a quadratic form $Q : \mathcal{M} \rightarrow \mathcal{O}_X$ if they hold on an affine open cover.

Now let \mathcal{L} be an invertible \mathcal{O}_X -module (i.e., locally free of rank 1). A quadratic form on \mathcal{L} , the case of our primary concern, is a quadratic map $Q : \mathcal{L} \rightarrow \mathcal{O}_X$, but it is given equivalently by a global section $q \in \text{Sym}^2(\mathcal{L}^\vee)(X) \simeq (\mathcal{L}^\vee)^{\otimes 2}(X)$. Because we will use this identification frequently, we make it explicit. The identification is defined locally on X , so suppose $X = \text{Spec } R$ and L is a free module of rank 1 over R . Then $\text{Sym}^2(L^\vee) \simeq (L^\vee)^{\otimes 2} \simeq (L^{\otimes 2})^\vee$. Suppose $q \in \text{Sym}^2(L^\vee)$, so $q : L \otimes L \rightarrow R$ is an R -module homomorphism. Let $L = Re$ for some $e \in L$ and define the quadratic map $Q : L \rightarrow R$ by $Q(re) = r^2q(e \otimes e)$; this definition is independent of the choice of e . Conversely, if $Q : L \rightarrow R$ is a quadratic map, then again letting $L = Re$ we define the R -module homomorphism $q : L \otimes L \rightarrow R$ by $q(e \otimes e) = Q(e)$.

Remark 2.1. One must remember the domain \mathcal{L} in this identification. Indeed, if $i : \mathcal{T}^{\otimes 2} \simeq \mathcal{O}_X$ is an isomorphism of \mathcal{O}_X -modules, so that $\mathcal{T} \in \text{Pic}(X)[2]$, then i defines a quadratic form $I : \mathcal{T} \rightarrow \mathcal{O}_X$, called a *neutral form*, giving rise to an isomorphism

$$(\mathcal{L}^\vee)^{\otimes 2} \xrightarrow{\simeq} ((\mathcal{L} \otimes \mathcal{T})^\vee)^{\otimes 2}.$$

The notions of *(semi)nondegnerate* and *(semi)nonsingular* can be made quite explicit for quadratic forms of rank 1. These conditions are local, so let $Q : L \rightarrow R$ be a quadratic form with $L = Re$. Then Q is uniquely specified by the element $Q(e) = a \in R$ and the associated bilinear form is specified by $T(e, e) = 2a$, with $2a = \det(Q) \in R/R^{\times 2}$, because a different choice of basis $e' = ue$ gives $Q(e') = u^2Q(e) = au^2$ with $u \in R^\times$. We find that Q is nondegnerate if and only if $2a$ is a nonzerodivisor and is regular if $2a \in R^\times$, and Q is seminondegnerate if a is a nonzerodivisor and seminonsingular if a is a unit.

Remark 2.2. The slightly unpleasant term *seminondegnerate* is not standard in the literature. It is common to use the term *nondegnerate* instead, but we do not do this here to avoid potential confusion.

Given two quadratic forms $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ and $Q' : \mathcal{L}' \rightarrow \mathcal{O}_X$, corresponding to $q \in (\mathcal{L}^\vee)^{\otimes 2}(X)$ and $q' \in (\mathcal{L}'^\vee)^{\otimes 2}(X)$, from the element

$$q \otimes q' \in ((\mathcal{L}^\vee)^{\otimes 2} \otimes (\mathcal{L}'^\vee)^{\otimes 2})(X) \simeq ((\mathcal{L} \otimes \mathcal{L}')^\vee)^{\otimes 2}(X)$$

we define the corresponding tensor product $Q \otimes Q' : \mathcal{L} \otimes \mathcal{L}' \rightarrow \mathcal{O}_X$: following the identification above, over $X = \text{Spec } R$, if $L = Re$ and $L' = Re'$ then $L \otimes L' = R(e \otimes e')$ and $(Q \otimes Q')(e \otimes e') = Q(e)Q(e')$. The tensor product gives the set of similarity classes of quadratic forms of rank 1 over X the structure of a commutative monoid.

Remark 2.3. The definition of the tensor product of quadratic forms is more subtle in general for forms of arbitrary rank; here we find the correct notion because we can think of rank 1 quadratic forms as rank 1 symmetric bilinear forms.

Definition 2.4. Let $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ be a (rank 1) quadratic form. We say Q is *cancellative* if it is cancellative in the monoidal sense, as $q \in (\mathcal{L}^\vee)^{\otimes 2}(X)$: if $q' \in ((\mathcal{L}')^\vee)^{\otimes 2}$ and $q'' \in ((\mathcal{L}'')^\vee)^{\otimes 2}(X)$ have $q \otimes q'$ similar to $q \otimes q''$, then q' is similar to q'' .

We say Q is *locally cancellative* if for all $x \in X$ there exists an affine open neighborhood $U \ni x$ such that $Q|_U$ is cancellative.

Proposition 2.5. *A (rank 1) quadratic form $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ is locally cancellative if and only if Q is seminondegenerate.*

Moreover, a locally cancellative rank 1 quadratic form Q over X is cancellative. If X is affine, then Q is cancellative if and only if it is locally cancellative.

Proof. Both properties are local, so it suffices to check this over a ring $X = \text{Spec } R$ such that the quadratic forms involved are free. To a quadratic form $Q : L = Re \rightarrow R$, we have $Q(e) = a \in R$. Similarly, if Q' and Q'' are other rank 1 quadratic forms with $Q'(e') = a' \in R$ and $Q''(e'') = a'' \in R$, then $(Q \otimes Q')(e \otimes e') = aa'$ and $(Q \otimes Q'')(e \otimes e'') = aa''$. We have $Q' \sim Q''$ if and only if there exists $u \in R^\times$ such that $a' = ua''$.

Thus, if Q is seminondegenerate then a is a nonzerodivisor, so $Q \otimes Q' \sim Q \otimes Q''$ implies $aa' = uaa''$, which implies $a' = ua''$, which implies $Q' \sim Q''$, so Q is locally cancellative. Conversely, if Q is locally cancellative and a is a zero divisor, with $aa' = 0$ and $a' \neq 0$, then taking $Q'(e') = a'$ and $Q''(e'') = 0$ we have $Q \otimes Q' \sim Q \otimes Q''$, so $Q' \sim Q''$ and thus there exists $u \in R^\times$ such that $a' = u(0) = 0$, a contradiction.

Now for the second statement. Let $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ be locally cancellative. By the previous paragraph, Q is seminondegenerate. Suppose that $Q \otimes Q' \sim Q \otimes Q''$ for rank 1 quadratic forms Q' and Q'' ; we will show that $Q' \sim Q''$. Cancelling in $\text{Pic}(X)$, we may assume without loss of generality that $\mathcal{L}' = \mathcal{L}''$. Let $U = \text{Spec } R$ be an affine open subset of X such that $\mathcal{L}|_U = L = Re$ is free, as is $L' = Re' = L'' = Re''$. Let $a = Q(e)$ and, similarly, $a' = Q'(e')$ and $a'' = Q''(e'')$, so as in the previous paragraph we are given (a unique) $u \in R^\times$ such that $aa' = uaa''$. Since Q is locally cancellative, we have $a' = ua''$, which defines a similarity $Q' \sim Q''$. Repeating this on an open cover, the elements u glue to give an element $g \in \mathcal{O}_X(X)^\times$ and so, together with the identity map on $\mathcal{L}' = \mathcal{L}''$, we therefore have a similarity $Q' \sim Q''$.

The converse in the final statement follows immediately by taking $U = X$ if X is affine. □

The following corollary is then immediate.

Corollary 2.6. *The subset of locally cancellative quadratic forms over X is a submonoid of the monoid of rank 1 quadratic forms over X .*

Remark 2.7. The global notion of cancellative is not as robust as one may like. Kleiman [1979] gives an example of a scheme X and a global section $t \in \mathcal{O}_X(X)$ that is a nonzerodivisor such that it becomes a zerodivisor in an affine open $t|_U \in \mathcal{O}_X(U)$.

The similarity class of a locally cancellative quadratic form is determined by its image (“effective Cartier divisors on a scheme are the same as invertible sheaves with a choice of regular global section” [Stacks 2015, Tag 01X0]), as follows.

Lemma 2.8. *There is a (functorial) isomorphism of commutative monoids*

$$\left\{ \begin{array}{l} \text{similarity classes of rank 1,} \\ \text{locally cancellative quadratic forms} \\ Q : \mathcal{L} \rightarrow \mathcal{O}_X \\ \text{modulo neutral forms} \end{array} \right\} \xrightarrow{\sim} \left\{ \begin{array}{l} \text{locally free ideal sheaves} \\ \mathcal{I} \subseteq \mathcal{O}_X \\ \text{such that } [\mathcal{I}] \in 2 \text{ Pic}(X) \end{array} \right\},$$

where the similarity class of a quadratic form $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ maps to the ideal \mathcal{I} of \mathcal{O}_X generated by the values $Q(\mathcal{L})$.

Ideal sheaves are a monoid under multiplication, so the “monoid” part of Lemma 2.8 says that the tensor product $Q \otimes Q'$ of two quadratic forms Q and Q' maps to the product $\mathcal{I}\mathcal{I}'$ of their associated ideal sheaves \mathcal{I} and \mathcal{I}' .

Proof. First, we show the map is well-defined, which we may do locally. Let $Q : L \rightarrow R$ be a rank 1, locally cancellative quadratic form; then L is free, so we may write $L = Re$ and then $Q(L) = Q(e)R$. By Proposition 2.5, we know that $Q(e)$ is a nonzerodivisor, and this is independent of the choice of e (up to a unit of R). If $Q' : L' \rightarrow R$ is similar to Q , then there exist R -linear isomorphisms $f : L \rightarrow L'$ and $g : R \rightarrow R$ such that $Q'(f(x)) = g(Q(x))$ for all $x \in L$. Letting $e' = f(e)$ we have $L' = Re'$. The map g must be of the form $g(a) = ua$ for some $u \in R^\times$, so $Q'(L') = Q'(f(L)) = uQ(L) = uQ(e)R = Q(e)R$, and thus the image is a well-defined principal ideal.

Now let $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ be a locally cancellative quadratic form of rank 1, corresponding to the global section $q \in (\mathcal{L}^{\otimes 2})^\vee(X)$. We claim that $q : \mathcal{L}^{\otimes 2} \rightarrow \mathcal{O}_X$ is an isomorphism onto its image $\mathcal{I} = Q(\mathcal{L}) = q(\mathcal{L}^{\otimes 2})$. If $U = \text{Spec } R$ is an affine open set, where $\mathcal{L}|_U = Re$, then $q(L^{\otimes 2}) = q(e \otimes e)R$; if, further, U is such that q is cancellative over U , we have that $q(e \otimes e)$ is a nonzerodivisor, so q is injective on U (and $q(L)$ is free). By hypothesis, such affine open sets U cover X , so we have $[\mathcal{I}] = [\mathcal{L}^{\otimes 2}] \in \text{Pic}(X)$.

Next, let $q : \mathcal{L} \rightarrow \mathcal{O}_X$ and $q' : \mathcal{L}' \rightarrow \mathcal{O}_X$ be locally cancellative quadratic forms such that $q(\mathcal{L}) = q'(\mathcal{L}') = \mathcal{I}$. Since $[\mathcal{I}] = [\mathcal{L}^{\otimes 2}] = [(\mathcal{L}')^{\otimes 2}]$, tensoring q' by a neutral form we may assume that $f : \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$. Then, on any affine open set $U = \text{Spec } R \subseteq X$, where $\mathcal{L}|_U = Re$ and $\mathcal{L}'|_U = Re'$, we have $q(L) = q(e)R = q'(e')R$. Therefore, there exists $u \in R$ such that $q(e) = uq'(e')$, and $u' \in R$ such that $q'(e') = u'q(e)$. Thus $q(e)(1 - uu') = 0$. On an affine open set U where

q is cancellative, we have $uu' = 1$, so $u \in R^\times$. Moreover, the element u is unique, since if $q(e) = uq'(e') = vq'(e')$ then $(u - v)q'(e') = 0$ so, since q' is cancellative, we have $u = v$. Repeating this argument on an open cover where both \mathcal{L} and \mathcal{L}' are free, there exists (a unique) $u \in \mathcal{O}_X(X)^\times$ giving rise to an isomorphism $\mathcal{O}_X \xrightarrow{\sim} \mathcal{O}_X$ such that $q'f = uq$, so that q and q' are similar.

Finally, the map is surjective: we are given that there exists an invertible bundle \mathcal{L} such that $\mathcal{L}^{\otimes 2} \simeq \mathcal{I}$. The embedding $\mathcal{L}^{\otimes 2} \simeq \mathcal{I} \hookrightarrow \mathcal{O}_X$ then defines a locally cancellative quadratic form $Q : \mathcal{L} \rightarrow \mathcal{O}_X$ with values $Q(\mathcal{L}) = \mathcal{I}$, as can readily be checked locally. □

Remark 2.9. A (locally) cancellative rank 1 quadratic form might not pull back to a (locally) cancellative form under an arbitrary morphism of schemes.

To work with discriminants, we will work modulo 2 and 4 as follows: The multiplication-by-4 map on \mathcal{O}_X gives a closed immersion

$$X_{[4]} = X \times_{\text{Spec } \mathbb{Z}} \text{Spec } \mathbb{Z}/4\mathbb{Z} \hookrightarrow X$$

and the pullback $\mathcal{L}_{[4]} = \mathcal{L} \otimes_{\mathcal{O}_X} 4\mathcal{O}_X$ is an invertible $\mathcal{O}_{X_{[4]}}$ -module, equipped with a map $_{[4]} : \mathcal{L} \rightarrow \mathcal{L}_{[4]}$. We can also, further, work modulo 2, obtaining $\mathcal{L}_{[2]}$.

Let R be a commutative ring. Then squaring gives a well-defined map of sets

$$(2.10) \quad \text{sq} : R/2R \rightarrow R/4R, \quad \text{sq}(t + 2R) = t^2 + 4R.$$

The map sq is functorial in R and canonically defined, so we can sheafify: if \mathcal{L} is an invertible \mathcal{O}_X -module, there is a unique map

$$\text{sq} : \mathcal{L}_{[2]} \rightarrow (\mathcal{L}_{[4]})^{\otimes 2}$$

locally defined by (2.10). Explicitly, for an affine open set $U = \text{Spec } R$ of X , where $\mathcal{L}(U) = L = Re$, we may write

$$\mathcal{L}_{[2]}(U) = (R/2R)e \quad \text{and} \quad \mathcal{L}_{[4]}^{\otimes 2}(U) = (R/4R)(e \otimes e),$$

and

$$\text{sq}((t + 2R)e) = \text{sq}(t + 2R)(e \otimes e)$$

is well-defined (independent of the choice of e).

Definition 2.11. A *discriminant* over X is a pair (d, \mathcal{L}) where \mathcal{L} is an invertible \mathcal{O}_X -module and $d \in (\mathcal{L}^\vee)^{\otimes 2}(X)$ such that there exists $t \in \mathcal{L}_{[2]}^\vee(X)$ with

$$(2.12) \quad \text{sq}(t) = d_{[4]} \in (\mathcal{L}_{[4]}^\vee)^{\otimes 2}(X).$$

If 2 is invertible in X , then $X_{[4]}$ is the empty scheme and the square condition (2.12) is vacuously satisfied.

Definition 2.13. An *isomorphism* between discriminants (d, \mathcal{L}) and (d', \mathcal{L}') is an isomorphism $f : \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$ such that $(f^\vee)^{\otimes 2} : (\mathcal{L}'^\vee)^{\otimes 2} \rightarrow (\mathcal{L}^\vee)^{\otimes 2}$ has $(f^\vee)^{\otimes 2}(d') = d$.

Equivalently, an isomorphism between discriminants is an isometry — not a similarity! — between the corresponding quadratic forms.

In what follows, we will often abbreviate d for a discriminant (d, \mathcal{L}) and refer to \mathcal{L} as the underlying invertible sheaf.

Let $\text{Disc}(X)$ denote the set of discriminants over X up to isomorphism. For an invertible sheaf \mathcal{L} on X , let $\text{Disc}(X; \mathcal{L}) \subseteq \text{Disc}(X)$ denote the subset of isomorphism classes of discriminants d whose underlying invertible sheaf is (isomorphic to) \mathcal{L} . Define $\mathbf{Disc}(X)$ to be the sheaf associated to the presheaf $U \mapsto \text{Disc}(U)$.

Lemma 2.14. $\text{Disc}(X)$ has the structure of commutative monoid under tensor product, with identity element represented by the class of $(1, \mathcal{O}_X)$. Moreover, $\text{Disc}(X; \mathcal{O}_X)$ is a submonoid of $\text{Disc}(X)$ with absorbing element $(0, \mathcal{O}_X)$.

Proof. The binary operation of tensor product on quadratic forms restricts to a binary operation on discriminants: if (d, \mathcal{L}) and (d', \mathcal{L}') are discriminants, with $t \in \mathcal{L}_{[2]}^\vee(X)$ satisfying $\text{sq}(t) = d_{[4]} \in (\mathcal{L}_{[4]}^\vee)^{\otimes 2}(X)$ and similarly for (d', \mathcal{L}') , then

$$\begin{aligned} \text{sq}(t \otimes t') &= d_{[4]} \otimes d'_{[4]} = (d \otimes d')_{[4]} \in ((\mathcal{L}_{[4]}^\vee)^{\otimes 2} \otimes ((\mathcal{L}'_{[4]})^\vee)^{\otimes 2})(X) \\ &\simeq ((\mathcal{L} \otimes \mathcal{L}')_{[4]})^\vee{}^{\otimes 2}(X). \end{aligned}$$

This definition is independent of the choice of a representative discriminant in an isomorphism class, so we obtain a binary operation on $\text{Disc}(X)$. This operation is associative and commutative, and $(1, \mathcal{O}_X)$ is an identity by definition of the tensor product. The subset $\text{Disc}(X; \mathcal{O}_X)$ is closed under tensor product, and $(0, \mathcal{O}_X)$ is visibly an absorbing element. \square

Lemma 2.15. *There is a functorial monoid isomorphism*

$$(2.16) \quad \{d \in \mathcal{O}_X(X) : d \text{ is a square modulo } 4\mathcal{O}_X(X)\} / \mathcal{O}_X(X)^{\times 2} \xrightarrow{\sim} \text{Disc}(X; \mathcal{O}_X).$$

Proof. The explicit identification of the monoid homomorphism between elements

$$d \in \mathcal{O}_X(X) \simeq (\mathcal{O}_X^\vee)^{\otimes 2}(X)$$

and discriminants as quadratic forms is explained in the beginning of Section 2, with the discriminant condition passing through on both sides.

Suppose that f is an isomorphism between discriminants d and d' . Let $U = \text{Spec } R \subseteq X$ be an affine open subset. Then $d|_U : R \rightarrow R$ is a quadratic map with $d(r) = r^2 d(1)$ for all $r \in R$. The restriction $f|_U : \mathcal{O}_X(U) = R \rightarrow R$ is an isomorphism and so is identified with a unique element $u \in R^\times$; thus, in R we have

$d'|_U(f|_U(1)) = d'|_U(u) = u^2d'|_U(1) = d|_U(1)$; by gluing, there exists a (unique) global section $u \in \mathcal{O}_X(X)^\times$ such that $d = u^2d'$. \square

Example 2.17. If $X = \text{Spec } R$, where R is a PID or local ring, then, by Lemma 2.15, $\text{Disc}(R) = \text{Disc}(R; R)$ is canonically identified with

$$\{d \in R : d \text{ is a square in } R/4R\}/R^{\times 2}.$$

So, for $R = \mathbb{Z}$, since $\mathbb{Z}^{\times 2} = \{1\}$ we recover the usual set of discriminants as those integers $d \equiv 0, 1 \pmod{4}$.

Given a discriminant (d, \mathcal{L}) , we can forget the quadratic map d and consider only the isomorphism class of the \mathcal{O}_X -module \mathcal{L} ; this gives a map

$$p : \text{Disc}(X) \rightarrow \text{Pic}(X).$$

Proposition 2.18. *The sequence*

$$\text{Disc}(X; \mathcal{O}_X) \rightarrow \text{Disc}(X) \xrightarrow{p} \text{Pic}(X)$$

of commutative monoids is exact.

Proof. The map $p : \text{Disc}(X) \rightarrow \text{Pic}(X)$ is surjective because an invertible module \mathcal{L} has the zero quadratic form $d = 0$, which is a discriminant taking $t = 0$. (One can hardly do better in general, since it may be the case that $\mathcal{L}(X) = \{0\}$ has no nonzero global sections.)

Let $i : \text{Disc}(X; \mathcal{O}_X) \hookrightarrow \text{Disc}(X)$ be the inclusion map. We show that $I_i = K_p$. The inclusion $I_i \subseteq K_p$ is easy, so we show the reverse inclusion. Let d and d' be discriminants and suppose $([d], [d']) \in K_p$; then the underlying invertible sheaves of d and d' are isomorphic, so we may assume without loss of generality that $d, d' \in (\mathcal{L}^\vee)^{\otimes 2}(X)$. To show $([d], [d']) \in I_i$, we need to show that there exist $\delta, \delta' \in \text{Disc}(X; \mathcal{O}_X)$ such that $\delta \otimes d' = \delta' \otimes d$. For this purpose, we may take $\delta = \delta' = 0$. More generally, we could take any $\delta \in (d : d') = \{\delta \in \mathcal{O}_X(X) : \delta d' \in \langle d \rangle\} \subseteq \mathcal{O}_X(X)$. \square

3. Quadratic algebras

In this section, we give a monoid structure on the set of isomorphism classes of quadratic algebras. We begin by discussing the algebras over commutative rings, then work over a general base scheme. For more on quadratic rings and standard involutions, see Knus [1991, Chapter I, §1.3] and Voight [2011, §§1–2].

Let R be a commutative ring. An R -algebra is an associative ring B with 1 equipped with an embedding $R \hookrightarrow B$ of rings (mapping $1 \in R$ to $1 \in B$) whose image lies in the center of B ; we identify R with this image in B . In particular, B is necessarily faithful as an R -module. A homomorphism of R -algebras is required to preserve 1.

Definition 3.1. A *quadratic R -algebra* (or *quadratic ring over R*) is an R -algebra S that is finite locally free of rank 2 as an R -module.

By *finite locally free* we mean that there is a cover of $\text{Spec } R$ by standard open sets $D(f_i)$ with $i \in I$ such that the localization M_{f_i} is a free R_{f_i} -module for all $i \in I$. There are a number of other equivalent formulations of this condition [Stacks 2015, Tag 00NV], including that M is finitely presented and R -flat, that M is finite projective, and that M is finitely presented with the localization $M_{\mathfrak{p}}$ free for all primes $\mathfrak{p} \in \text{Spec}(R)$.

Let S be a quadratic R -algebra. Then S is commutative, and there is a unique *standard involution* on S , an R -linear homomorphism $\sigma : S \rightarrow S^{\text{op}} = S$ such that $\sigma(\sigma(x)) = x$ and $x\sigma(x) \in R$ for all $x \in S$ [Voight 2011, Lemma 2.9]. Consequently, we have a linear map $\text{trd} : S \rightarrow R$ defined by $\text{trd}(x) = x + \sigma(x)$ and a multiplicative map $\text{nrd} : S \rightarrow R$ defined by $\text{nrd}(x) = x\sigma(x) = \sigma(x)x$ with the property that $x^2 - \text{trd}(x)x + \text{nrd}(x) = 0$ for all $x \in S$.

Lemma 3.2. *If S is free as an R -module, then there is a basis $1, x$ for S as an R -module.*

Proof. Let x_1, x_2 be a basis for S ; then there exist $a_1, a_2 \in R$ such that $1 = a_1x_1 + a_2x_2$. Let $x_1^2 = b_1x_1 + b_2x_2$ and $x_1x_2 = c_1x_1 + c_2x_2$ with $b_1, b_2, c_1, c_2 \in R$. Then

$$x_1 = x_1 \cdot 1 = a_1x_1^2 + a_2x_1x_2 = (a_1b_1 + a_2c_1)x_1 + (a_1b_2 + a_2c_2)x_2.$$

Thus $a_1b_1 + a_2c_1 = 1$. Let $x = -c_1x_1 + b_1x_2$. Then

$$\det \begin{pmatrix} a_1 & a_2 \\ -c_1 & b_1 \end{pmatrix} = a_1b_1 + a_2c_1 = 1 \quad \text{and} \quad \begin{pmatrix} a_1 & a_2 \\ -c_1 & b_1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ x \end{pmatrix}$$

so $1, x$ is a basis for S as an R -module. □

Remark 3.3. Lemma 3.2 does not generally extend to R -algebras of higher rank. On the one hand, if S is finite locally free we always have $S \simeq R \oplus S/R$ as R -modules [Voight 2011, Lemma 1.3]; on the other hand, if S is free this need not imply that S/R is free. (However, S/R is still locally free, so for the purposes of making local arguments you can refine an open cover to one over which S/R is in fact free to then find a basis containing 1.)

If S is free over R with basis $1, x$, then the multiplication table is determined by the multiplication $x^2 = tx - n$; consequently, we have a bijection

$$(3.4) \quad \left\{ \begin{array}{l} \text{free quadratic } R\text{-algebras } S \text{ over } R, \\ \text{equipped with a basis } 1, x \end{array} \right\} \xrightarrow{\sim} R^2,$$

$$S \mapsto (\text{trd}(x), \text{nrd}(x)) = (x + \bar{x}, x\bar{x}) = (t, n).$$

A change of basis for a free quadratic R -algebra is of the form $x \mapsto u(x+r)$ with $u \in R^\times$ and $a \in R$, mapping

$$(3.5) \quad (t, n) \mapsto (u(t+2r), u^2(n+tr+r^2)).$$

We have identified $R \subseteq S$ as a subring; the quotient S/R is locally free of rank 1. Therefore, we have a canonical identification

$$(3.6) \quad S/R \xrightarrow{\sim} \wedge^2 S, \quad x+R \mapsto 1 \wedge x.$$

Lemma 3.7. *Let S be a quadratic R -algebra. Then the map*

$$(3.8) \quad \begin{aligned} d : (\wedge^2 S)^{\otimes 2} &\rightarrow R, \\ (x \wedge y) \otimes (z \wedge w) &\mapsto (x\sigma(y) - \sigma(x)y)(z\sigma(w) - \sigma(z)w), \end{aligned}$$

is a discriminant.

We have

$$(3.9) \quad \begin{aligned} d((1 \wedge x)^{\otimes 2}) &= (x - \sigma(x))^2 = (2x - \text{trd}(x))^2 \\ &= 4x^2 - 4x \text{trd}(x) + \text{trd}(x)^2 = \text{trd}(x)^2 - 4 \text{nr}(x) \end{aligned}$$

in the lemma, as one might expect. We accordingly call the quadratic map $d = d(S)$ in Lemma 3.7 the *discriminant* of S .

Proof. We define the map

$$t : \wedge^2(S/2S) \rightarrow R/2R, \quad t(1 \wedge x) = \text{trd}(x),$$

via the identification (3.6). The map t is well-defined since $\text{trd}(x+r) = \text{trd}(x)+2r \equiv \text{trd}(x) \pmod{2R}$. We then verify that

$$\text{sq}(t)((1 \wedge x)^{\otimes 2}) = t(1 \wedge x)^2 = \text{trd}(x)^2 \equiv \text{trd}(x)^2 - 4 \text{nr}(x) = d((1 \wedge x)^{\otimes 2}) \pmod{4R}$$

by (3.9). □

Recall that a commutative R -algebra B is *separable* if B is projective as a $B \otimes_R B$ -module via the map $x \otimes y \mapsto xy$. If $B \simeq R[x]/(f(x))$ with $f(x) \in R[x]$, then B is separable if and only if the ideal generated by $f(x)$ and its derivative $f'(x)$ is the unit ideal.

Lemma 3.10. *A quadratic R -algebra S is separable if and only if its discriminant d is an isomorphism.*

Proof. The map $d : (\wedge^2 S)^{\otimes 2} \rightarrow R$ is an isomorphism if and only if it is locally an isomorphism, so we reduce to the case where $S = R[x]/(x^2-tx+n) = R[x]/(f(x))$. Then, by (3.9), d is an isomorphism if and only if $\text{trd}(x)^2 - 4 \text{nr}(x) \in R^\times$ if and only if S is separable [Knus 1991, Chapter I, (7.3.4)]. □

Corollary 3.11. *If S is a separable quadratic R -algebra then $\wedge^2 S \in \text{Pic}(R)[2]$.*

Now let X be a scheme.

Definition 3.12. A quadratic \mathcal{O}_X -algebra is a sheaf \mathcal{S} of \mathcal{O}_X -algebras that is locally free of rank 2 as a sheaf of \mathcal{O}_X -modules: there is a basis of open sets U of X such that $\mathcal{S}(U)$ is free of rank 2 as an $\mathcal{O}_X(U)$ -module.

Equivalently, a quadratic \mathcal{O}_X -algebra is given by a *double cover* $\phi : Y \rightarrow X$, a finite locally free morphism of schemes of degree 2: the sheaf $\phi_* \mathcal{O}_Y$ is a sheaf of \mathcal{O}_X -algebras that is locally free of rank 2. By uniqueness of the standard involution, we obtain a *standard involution* on \mathcal{S} , a standard involution on $\mathcal{S}(U)$ for all open sets U (covering each by affine open sets where \mathcal{S} is free), and, in particular, maps trd and nrd on \mathcal{S} .

Analogous to Lemma 3.7, we have the following result.

Proposition 3.13. *Let \mathcal{S} be a quadratic \mathcal{O}_X -algebra. Then there exists a unique discriminant $d : (\wedge^2 \mathcal{S})^{\otimes 2} \rightarrow \mathcal{O}_X$ that coincides locally with the one defined by (3.8).*

Proof. Let $\mathcal{L} = \wedge^2 \mathcal{S}$. We must exhibit $t \in \mathcal{L}_{[2]}^\vee(X)$ such that

$$\text{sq}(t) = d_{[4]} \in (\mathcal{L}_{[4]}^\vee)^{\otimes 2}(X),$$

where [4] denotes working modulo 4, as in the previous section. We adapt the argument in Lemma 3.7 to a global setting. First working locally, let $U = \text{Spec } R \subseteq X$ be an open set where $\mathcal{S}(U) = S$ and $\mathcal{L}(U) = L = \wedge^2 S$. Since $\text{trd}(x+r) = \text{trd}(x) + 2r$ for $r \in R$ and $x \in S$, the map

$$t : \wedge^2(S/2S) \simeq L/2L \rightarrow R/2R, \quad x \wedge y \mapsto \text{trd}(x + y),$$

is well-defined (since $t(x \wedge x) = \text{trd}(2x) = 0$) and R -linear. This map does not depend on any choices, so repeating this on an open cover we obtain an element $t \in \mathcal{L}_{[2]}^\vee(X)$.

Now we verify that $\text{sq}(t) = d_{[4]} \in (\mathcal{L}_{[4]}^\vee)^{\otimes 2}(X)$. We may do so on an open cover, where \mathcal{S} is free, so let $S = R \oplus Rx$ and $\wedge^2 S = L = R(1 \wedge x)^{\otimes 2}$. Then

$$d((1 \wedge x)^{\otimes 2}) = \text{trd}(x)^2 - 4 \text{nrd}(x) \equiv \text{trd}(x)^2 \pmod{4R}.$$

On the other hand, by definition we have

$$\text{sq}(t)((1 \wedge x)^{\otimes 2}) \equiv t(1 \wedge x)^2 = \text{trd}(1 + x)^2 = (2 + \text{trd}(x))^2 \equiv \text{trd}(x)^2 \pmod{4R}.$$

The result follows. □

A quadratic \mathcal{O}_X -algebra \mathcal{S} is separable if and only if d induces an isomorphism of \mathcal{O}_X -modules $(\wedge^2 \mathcal{S})^{\otimes 2} \xrightarrow{\sim} \mathcal{O}_X$, as this is true on any affine open set.

Let $\text{Quad}(X)$ denote the set of isomorphism classes of quadratic \mathcal{O}_X -algebras and, for an invertible \mathcal{O}_X -module \mathcal{L} , let $\text{Quad}(X; \mathcal{L}) \subseteq \text{Quad}(X)$ be the subset of

those algebras \mathcal{S} such that there exists an isomorphism $\bigwedge^2 \mathcal{S} \simeq \mathcal{L}$ of \mathcal{O}_X -modules. Similarly, define $\mathbf{Quad}(X)$ to be the sheaf associated to the presheaf $U \mapsto \mathbf{Quad}(U)$.

We now give $\mathbf{Quad}(X)$ the structure of a commutative monoid.

Construction 3.14. Let $X = \text{Spec } R$ and let $S = R \oplus Rx$ and $T = R \oplus Ry$ be free quadratic R -algebras with choice of basis. Let $x^2 = tx - n$ and $y^2 = sy - m$, so

$$t = \text{trd}(x), \quad n = \text{nrđ}(x), \quad s = \text{trd}(y), \quad m = \text{nrđ}(y) \quad \text{with } t, n, s, m \in R.$$

Then we define the free quadratic R -algebra

$$S * T = R \oplus Rw,$$

where

$$(3.15) \quad w^2 = (st)w - (mt^2 + ns^2 - 4nm).$$

Construction 3.14 has been known for some time, e.g., it is given by Hahn [1994, Exercises 14–20, pp. 42–43]. (See the introduction for further context and references.)

Lemma 3.16. *Construction 3.14 is functorial with respect to the base ring R . The operation $*$ gives the set of free quadratic R -algebras with basis the structure of commutative monoid with identity element $R \times R = R[x]/(x^2 - x)$ and absorbing element $R[x]/(x^2)$.*

Proof. Functoriality is clear, and $S * T = T * S$ for all free quadratic R -algebras S and T by the symmetry of the construction. It is routine to check associativity. To check that $E = R \times R$ is the identity element for $*$ we simply substitute $s = 1$ and $m = 0$ to obtain $S * E = S$; a similar check works for the absorbing element. \square

Remark 3.17. Construction 3.14 generalizes the Kummer map, presented in the introduction. Indeed, suppose that R is a PID or local ring and $2 \in R^\times$. Then, by completing the square, any quadratic R -algebra S is of the form $S = R[x]/(x^2 - n) = R[\sqrt{n}]$, where $n = d(S)/4$. So, if $S = R[\sqrt{n}]$ and $T = R[\sqrt{m}]$, then

$$S * T = R[x]/(x^2 - 4nm) \simeq R[\sqrt{nm}].$$

At the same time, Construction 3.14 generalizes Artin–Schreier extensions of fields. Suppose that $R = k$ is a field of characteristic 2. Then every separable extension of k can be written in the form $k[x]/(x^2 - x + n)$, and

$$k[x]/(x^2 - x + n) * k[x]/(x^2 - x + m) = k[x]/(x^2 - x + (m + n)).$$

Since $4 = 0$, the discriminant of every such algebra has class 1 in $R/R^{\times 2}$.

In the above construction, if S and T are separable over R , so that they are (étale) Galois extensions of R [Lenstra 2008] (with the standard involutions σ and τ , respectively, as the nontrivial R -algebra automorphisms), then the algebra $S * T$

is the subalgebra of $S \otimes_R T$ fixed by the product of the involutions $\sigma \otimes \tau$ acting on $S \otimes_R T$ [Small 1972, Proposition 1].

In all cases, a direct calculation shows that (3.15) is satisfied by the element

$$w = x \otimes y + \sigma(x) \otimes \tau(y) \in S \otimes_R T;$$

this will figure in the proof of Theorem A. However, there is no reason why the R -algebra generated by w need be free of rank 2 over R ; for example, if R has characteristic 2 with $\sigma(x) = x$ and $\tau(y) = y$, then $w = 0$. Thus, Construction 3.14 can be thought of as a formal way to create a fixed subalgebra of $S \otimes_R T$ under the involution given by the product of standard involutions.

Lemma 3.18. *Construction 3.14 is functorial with respect to isomorphisms: if*

$$\begin{aligned} \phi : S = R \oplus Rx &\xrightarrow{\sim} S' = R \oplus Rx', \\ \psi : T = R \oplus Ry &\xrightarrow{\sim} T' = R \oplus Ry', \end{aligned}$$

are R -algebra isomorphisms of quadratic R -algebras, then there is a canonical isomorphism

$$\phi * \psi : S * T \xrightarrow{\sim} S' * T'.$$

Proof. There exist unique $u, v \in R^\times$ and $r, q \in R$ such that $\phi(x) = ux' + r$ and $\psi(y) = vy' + q$. Because ϕ is an R -algebra homomorphism, both $\phi(x)$ and x satisfy the same unique monic quadratic polynomial, and from

$$(ux' + r)^2 = t(ux' + r) - n$$

we conclude that

$$(x')^2 = u^{-1}(t - 2r)x' - u^{-2}(n - tr - r^2) = t'x - n',$$

so $t = ut' + 2r$ and $n = u^2n' + tr + r^2$. Similarly, we obtain $s = vs' + 2q$ and $m = v^2m' + sq + q^2$. We claim then that the map

$$\phi * \psi : S * T \xrightarrow{\sim} S' * T', \quad (\phi * \psi)(w) = (uv)w' + (qt + rs - 2qr),$$

is an isomorphism; for this we simply verify that

$$((uv)w' + (qt + rs - 2qr))^2 = st((uv)w' + (qt + rs - 2qr)) - (mt^2 + ns^2 - 4nm)$$

and the result follows. □

Lemma 3.19. *Let \mathcal{S} and \mathcal{T} be quadratic \mathcal{O}_X -algebras. Then there is a unique quadratic \mathcal{O}_X -algebra $\mathcal{S} * \mathcal{T}$ up to \mathcal{O}_X -algebra isomorphism with the property that, on any affine open set $U \subseteq X$ such that $S = \mathcal{S}(U)$ and $T = \mathcal{T}(U)$ are free, we have*

$$(\mathcal{S} * \mathcal{T})(U) \simeq S * T,$$

as in Construction 3.14.

Proof. This lemma is a standard application of gluing; we give the argument for completeness. Let $\{U_i = \text{Spec } R_i\}$ be an open affine cover of X on which

$$\mathcal{S}(U_i) = S_i = R_i \oplus R_i x_i \quad \text{and} \quad \mathcal{T}(U_i) = T_i = R_i \oplus R_i y_i$$

are free. We define $(\mathcal{S} * \mathcal{T})(U_i) = S_i * T_i = R_i \oplus R_i w_i$ according to Construction 3.14. We glue these according to the isomorphisms on \mathcal{S} and \mathcal{T} using Lemma 3.18, as follows: We have $U_i \cap U_j = U_j \cap U_i = \bigcup_k U_{ijk}$ covered by open sets $U_{ijk} = \text{Spec } R_{ik} \simeq \text{Spec } R_{jk}$ distinguished in U_i and U_j . Because \mathcal{S} is a sheaf, we have compatible isomorphisms

$$\phi_{ijk} : R_{ik} \oplus R_{ik} x_i = \mathcal{S}(\text{Spec } R_{ik}) \simeq \mathcal{S}(\text{Spec } R_{jk}) = R_{jk} \oplus R_{jk} x_j$$

for each such open set. Similarly, we obtain compatible isomorphisms ψ_{ijk} for \mathcal{T} over the same open cover. By Lemma 3.18, we obtain compatible isomorphisms

$$\phi_{ijk} * \psi_{ijk} : (\mathcal{S} * \mathcal{T})(\text{Spec } R_{ik}) \simeq (\mathcal{S} * \mathcal{T})(\text{Spec } R_{jk})$$

and can thereby glue on X to obtain a quadratic \mathcal{O}_X -algebra, unique up to \mathcal{O}_X -algebra isomorphism. □

Corollary 3.20. *Construction 3.14 gives $\text{Quad}(X)$ the structure of a commutative monoid, functorial in X , with identity element the isomorphism class of $\mathcal{O}_X \times \mathcal{O}_X$.*

Proof. Lemma 3.19 shows that Construction 3.14 extends to X and is well-defined on the set of isomorphism classes $\text{Quad}(X)$ of quadratic \mathcal{O}_X -algebras. To check that we obtain a functorial commutative monoid, it is enough to show this when X is affine, and this follows from Lemmas 3.16 and 3.18. □

Lemma 3.21. *If \mathcal{S} is a separable quadratic \mathcal{O}_X -algebra, then $\mathcal{S} * \mathcal{S} \simeq \mathcal{O}_X \times \mathcal{O}_X$.*

Proof. By gluing, it is enough to show this on an affine cover. Suppose that $S = R[x]/(x^2 - tx + n)$ has discriminant $d = t^2 - 4n$. Then, by definition, we have

$$S * S = R[w]/(w^2 - t^2 w + 2n(t^2 - 2n));$$

with the substitution $w \leftarrow w - 2n$, we find that $S * S \simeq R[w]/(w^2 - dw)$. Since $d \in R^\times$, the replacement $w \leftarrow wd^{-1}$ yields an isomorphism $S * S \simeq R \times R$. □

Remark 3.22. Given our description of the monoid product in the separable case, it follows that the submonoid of separable quadratic algebras is isomorphic to the group of isomorphism classes of étale quadratic covers $\check{H}_{\text{ét}}^1(X, \mathbb{Z}/2\mathbb{Z})$, a group killed by 2; for more, see Knus [1991, §III.4].

Lemma 3.23. *If $\mathcal{S}, \mathcal{T} \in \text{Quad}(X)$ then*

$$d(\mathcal{S} * \mathcal{T}) = d(\mathcal{S})d(\mathcal{T}) \in \text{Disc}(X)$$

and

$$\wedge^2(\mathcal{S} * \mathcal{T}) \simeq \wedge^2 \mathcal{S} \otimes \wedge^2 \mathcal{T}.$$

Proof. If $S = \mathcal{S}(U)$ and $T = \mathcal{T}(U)$ are as in Construction 3.14, then

$$\begin{aligned}
 (3.24) \quad d(S * T)((1 \wedge (x \otimes y))^{\otimes 2}) &= (st)^2 - 4(mt^2 + ns^2 - 4nm) \\
 &= (t^2 - 4n)(s^2 - 4m) \\
 &= d(S)((1 \wedge x)^{\otimes 2})d(T)((1 \wedge y)^{\otimes 2}).
 \end{aligned}$$

The first statement then follows. For the second, again on affine open sets we have the isomorphism

$$(3.25) \quad \wedge^2 S \otimes_R \wedge^2 T \rightarrow \wedge^2(S * T), \quad (1 \wedge x) \otimes (1 \wedge y) \mapsto 1 \wedge w,$$

which glues to give the desired isomorphism globally. □

Lemma 3.26. *The discriminant maps*

$$\text{disc} : \mathbf{Quad}(X) \rightarrow \mathbf{Disc}(X) \quad \text{and} \quad \text{disc} : \mathbf{Quad}(X; \mathcal{O}_X) \rightarrow \mathbf{Disc}(X; \mathcal{O}_X)$$

are surjective homomorphisms of sheaves of commutative monoids.

Proof. The fact that these maps are homomorphisms of sheaves of monoids follows locally from Lemma 3.23. We show these maps are surjective locally, and for that we may assume $X = \text{Spec } R$ and $L = Re$. We refer to Lemma 2.15 and Example 2.17: given $d \in R$ such that $d = t^2 - 4n$ with $t, n \in R$, we have the quadratic ring $R[x]/(x^2 - tx + n)$ of discriminant d . □

We are now ready to prove Theorem A.

Theorem 3.27. *Construction 3.14 is the unique system of binary operations*

$$* = *_X : \mathbf{Quad}(X) \times \mathbf{Quad}(X) \rightarrow \mathbf{Quad}(X),$$

one for each scheme X , such that:

(i) $\mathbf{Quad}(X)$ is a commutative monoid under $*$, with identity element the class of $\mathcal{O}_X \times \mathcal{O}_X$.

(ii) For each morphism $f : X \rightarrow Y$ of schemes, the diagram

$$\begin{array}{ccc}
 \mathbf{Quad}(Y) \times \mathbf{Quad}(Y) & \xrightarrow{*_Y} & \mathbf{Quad}(Y) \\
 \downarrow & & \downarrow f^* \\
 \mathbf{Quad}(X) \times \mathbf{Quad}(X) & \xrightarrow{*_X} & \mathbf{Quad}(X)
 \end{array}$$

is commutative.

(iii) If $X = \text{Spec } R$, and S and T are separable quadratic R -algebras with standard involutions σ and τ , then $S * T$ is the fixed subring of $S \otimes_R T$ under $\sigma \otimes \tau$.

Proof. By (3.4), the universal free quadratic algebra with basis is the algebra

$$S_{\text{univ}} = R_{\text{univ}}[x]/(x^2 - tx + n),$$

where $R_{\text{univ}} = \mathbb{Z}[t, n]$ is the polynomial ring in two variables over \mathbb{Z} ; in other words, for any commutative ring R and free quadratic R -algebra S with basis, there is a unique map $f : R_{\text{univ}} \rightarrow R$ such that $S = f^*S_{\text{univ}} = S_{\text{univ}} \otimes_{f, R_{\text{univ}}} R$. By (ii), then, following this argument on an open affine cover, we see that the monoid structure on $\text{Quad}(\text{Spec } R_{\text{univ}})$ determines the monoid structure for all schemes X .

Dropping subscripts, consider $S = R[x]/(x^2 - tx + n)$ and $T = R[y]/(y^2 - sy + m)$, where $R = \mathbb{Z}[t, n, s, m]$; we show there is a unique way to define $S * T$.

To begin, we claim that $S * T$ is free over R . As R -modules, we can write $S * T = R \oplus Iz$, where $I \subseteq F = \text{Frac}(R)$ is a projective R -submodule of F and the class $[I] \in \text{Pic}(R)$ is well-defined. But $\text{Pic}(\mathbb{Z}[t, n, s, m]) \simeq \text{Pic}(\mathbb{Z}) = \{0\}$ (\mathbb{Z} is *seminormal* [Gilmer and Heitmann 1980]), so $I \simeq R$.

Now let

$$D = (t^2 - 4n)(s^2 - 4m).$$

Then $S[1/D]$ and $T[1/D]$ are separable over $R[1/D]$, with involutions $\sigma(x) = t - x$ and $\tau(y) = s - y$. By (iii), the product $S[1/D] * T[1/D]$ is the subring of $S[1/D] \otimes_{R[1/D]} T[1/D]$ generated by

$$z = x \otimes y + \sigma(x) \otimes \tau(y) = 2(x \otimes y) - s(x \otimes 1) - t(1 \otimes y).$$

Then

$$\begin{aligned} z^2 &= x^2 \otimes y^2 + 2nm + \sigma(x)^2 \otimes \tau(y)^2 \\ &= (tx - n) \otimes (sy - m) + 2nm + (t\sigma(x) - n) \otimes (s\tau(y) - m) \\ &= ts(x \otimes y + \sigma(x) \otimes \tau(y)) - mt((x + \sigma(x)) \otimes 1) - ns(1 \otimes (y + \tau(y))) + 4nm \\ &= (st)z - (mt^2 + ns^2 - 4nm). \end{aligned}$$

In particular, $S[1/D] * T[1/D] \simeq R[1/D] \oplus R[1/D]z$.

By (ii), $(S * T)[1/D] \simeq S[1/D] * T[1/D]$, and we have $S * T \subseteq (S * T)[1/D]$. Since R is a UFD and $S * T$ is free over R , it is generated as an R -algebra by an element of the form $(az + b)/D^k$ for some $a, b \in R$ and $k \in \mathbb{Z}_{\geq 0}$. But, by (3.24), $d((S * T)[1/D]) = D$, so $d(S * T) = (a/D^k)^2 D \in R$; thus, $a/D^k \in R$. Since $\text{trd}((az + b)/D^k) = (ast + 2b)/D^k \in R$, we conclude that $2b/D^k \in R$; since D is not divisible by 2, by Gauss's lemma we have $b/D^k \in R$, so without loss of generality we may take $b = 0$ and suppose $S * T$ is generated by az for some $a \in R$. Since $R^\times = \{\pm 1\}$ and $(S * T)[1/D]$ is generated by z , we must have $a = D^k$ for some $k \in \mathbb{Z}_{\geq 0}$. Finally, we consider

$$\frac{\mathbb{Z}[x]}{(x^2 - tx + n)} * \frac{\mathbb{Z}[y]}{(y^2 - y)} = \frac{\mathbb{Z}[z]}{(z^2 - D^k tz + D^{2k}n)}$$

over $\mathbb{Z}[t, n]$. The algebra on the right has discriminant $D^{2k}(t^2 - 4n)$, but, by (i), it must be isomorphic to the algebra on the left of discriminant $(t^2 - 4n)$, so we must have $D^{2k} = 1$, so $k = 0$. Therefore $S * T = R \oplus Rz$. \square

Having given the monoid structure, we conclude this section by proving Theorem B.

Theorem 3.28. *Let X be a scheme. Then the following diagram of commutative monoids is functorial and commutative with exact rows and surjective columns:*

$$\begin{array}{ccccc}
 \mathbf{Quad}(X; \mathcal{O}_X) & \longrightarrow & \mathbf{Quad}(X) & \xrightarrow{\wedge^2} & \mathbf{Pic}(X) \\
 \downarrow \text{disc} & & \downarrow \text{disc} & & \parallel \\
 \mathbf{Disc}(X; \mathcal{O}_X) & \longrightarrow & \mathbf{Disc}(X) & \longrightarrow & \mathbf{Pic}(X)
 \end{array}$$

Proof. The exactness of the bottom row follows from Proposition 2.18. The exactness of the top row and commutativity of the diagram follows by the same (trivial) argument. Surjectivity follows from Lemma 3.26. \square

4. Proof of Theorem C

In this section, we prove Theorem C and conclude with some final discussion.

Let R be a commutative ring and let $R[4] = \{a \in R : 4a = 0\}$. Let

$$\wp(R) = \{r + r^2 : r \in R\}$$

and let $\wp(R)[4] = \wp(R) \cap R[4]$. Note that $4(r + r^2) = 0$ if and only if $(1 + 2r)^2 = 1$, so we have, equivalently,

$$\wp(R)[4] = \{r + r^2 : r \in R \text{ and } (1 + 2r)^2 = 1\}.$$

Lemma 4.1. $\wp(R)[4]$ is a subgroup of $R[4]$ under addition.

Proof. We have $0 = 0 + 0^2 \in \wp(R)[4]$. If $n = r + r^2 \in \wp(R)[4]$ and $m = s + s^2 \in \wp(R)[4]$ then

$$(r + s + 2rs) + (r + s + 2rs)^2 = (r + r^2) + (s + s^2) + 4(r + r^2)(s + s^2) = n + m$$

and $4(n + m) = 0$, so $n + m \in \wp(R)[4]$. Finally, if $n \in \wp(R)[4]$ then $-n = 3n \in \wp(R)[4]$ by the preceding sentence. \square

We define the Artin–Schreier group $AS(R)$ to be the quotient

$$AS(R) = \frac{R[4]}{\wp(R)[4]}.$$

Since $2R[4] \subseteq \wp(R)[4]$, the group $AS(R)$ is an elementary abelian 2-group.

We define a map $i : AS(R) \rightarrow \mathbf{Quad}(R; R)$ sending the class of $n \in AS(R)$ to the isomorphism class of the algebra $S = R[x]/(x^2 - x + n)$.

Proposition 4.2. *The map $i : \text{AS}(R) \rightarrow \text{Quad}(R, R)$ is a (well-defined) injective map of commutative monoids.*

Proof. Let $S = i(n) = R[x]/(x^2 - x + n)$ and $T = i(m) = R[y]/(y^2 - y + m)$ with $n, m \in \text{AS}(R)$. Then $S \simeq T$ if and only if $y = u(x + r)$ for some $u \in R^\times$ and $r \in R$, which, by (3.5), holds if and only if $u(1 + 2r) = 1$ and $u^2(n + r + r^2) = m$; these are further equivalent to $1 + 2r \in R^\times$ and

$$n + r + r^2 = m(1 + 2r)^2 = (1 + 4r + 4r^2)m.$$

But $4m = 0$, so $n + r + r^2 = m$ and, since $4n = 0$, we have $4(r + r^2) = 0$. Thus $S \simeq T$ if and only if $(1 + 2r)^2 = 1$ and $n + r + r^2 = m$, as desired. It follows from Construction 3.14 that $S * T = R[w]/(w^2 - w + (n + m))$, since $4nm = 0$, and $i(0) = R[w]/(w^2 - w)$ is the identity, so i is a homomorphism of monoids. \square

We now prove Theorem C, and recall $\text{Quad}(R; R)$ is the set of isomorphism classes of free quadratic R -algebras.

Theorem 4.3. *Let R be a commutative ring and let $d \in R$ be a discriminant. Then the fiber $\text{disc}^{-1}(d)$ of the map*

$$\text{disc} : \text{Quad}(R; R) \rightarrow \text{Disc}(R; R)$$

above d has a unique action of the group $\text{AS}(R)/\text{ann}_R(d)[4]$ compatible with the inclusion of monoids $\text{AS}(R) \hookrightarrow \text{Quad}(R; R)$.

Proof. A (free) quadratic R -algebra with basis $1, x$ such that $(x - \sigma(x))^2 = d$ has the form $S = R[x]/(x^2 - tx + n)$ with $t^2 - 4n = d$. Let $m \in R[4]$. Then by Construction 3.14 we have

$$S * (R[y]/(y^2 - y + m)) = (R[y]/(y^2 - y + m)) * S = R[w]/(w^2 - tw + dm + n)$$

since $4m = 0$, so $dm = (t^2 - 4n)m = t^2m$. Thus we have an action of $R[4]$ on the set of these quadratic R -algebras with basis and a free action of $R[4]/\text{ann}_R(d)[4]$. Two quadratic R -algebras S and S' are in the same orbit if and only if $t' = t$ and $n' = dm + n$ for some $m \in R[4]$ if and only if $t' = t$ and $n' - n \in dR[4]$; therefore, the orbits are indexed noncanonically by the set

$$\{t \in R : t^2 \equiv d \pmod{4R}\} \times R[4]/dR[4].$$

We now descend to isomorphism classes: by Proposition 4.2, the monoid multiplication $*$ is well-defined on isomorphism classes, giving the unique action of $\text{AS}(R)/\text{ann}_R(d)[4]$ on the fiber over d . \square

Under favorable hypotheses, the action of $\text{AS}(R)$ is free, so we make the following definition.

Definition 4.4. An element $t \in R$ is *SEC* (square even cancellative) if

- (i) t is a nonzerodivisor, and
- (ii) $r^2, 2r \in tR$ implies $r \in tR$ for all $r \in R$.

A quadratic R -algebra S is *SEC* if $\text{disc}(S)$ is a nonzerodivisor and there a basis $1, x$ for S such that $\text{trd}(x)$ is SEC. A quadratic \mathcal{O}_X -algebra \mathcal{S} is *SEC* if \mathcal{S} is SEC on an open affine cover of X .

Proposition 4.5. *Let R be a commutative ring. Then the action of $\text{AS}(R)$ on the set of SEC quadratic R -algebras of discriminant d is free.*

Proof. We continue as in the proof of Theorem 4.3. Let $m \in R[4]$ and let S be an SEC quadratic R -algebras with discriminant d . Let $1, x$ be a basis for $S = R[x]/(x^2 - tx + n)$ such that $t = \text{trd}(x)$ is SEC. To show the action is free, suppose that

$$(4.6) \quad S * (R[y]/(y^2 - y + m)) = R[w]/(w^2 - tw + dm + n) \simeq S;$$

we show that $m \in \wp(R)$. Equation (4.6) holds if and only if there exist $u \in R^\times$ and $r \in R$ such that $t = u(t + 2r)$ and $dm + n = u^2(n + tr + r^2)$. Consequently, $u^2d = d$. Since S is SEC, d is a nonzerodivisor, so $u^2 - 1 = 0$ and thus $t^2m = dm = tr + r^2$, so $r^2 = (tm - r)t$. We also have $2r = (1 - u)t$, so since t is SEC and $r^2, 2r \in tR$, we conclude that $r \in tR$. Let $r = at$ with $a \in R$. Then, substituting, we have $t^2(a^2 + a - m) = r^2 + tr - t^2m = 0$. Since t is a nonzerodivisor, we conclude that $a^2 + a - m = 0$, so $m \in \wp(R)$, as claimed. \square

Acknowledgements

The author would like to thank Asher Auel, Manjul Bhargava, James Borger, and Melanie Wood for helpful suggestions. The author is also indebted to the anonymous referee and Owen Biesel for very detailed and helpful comments and corrections. The author was supported by an NSF CAREER Award (DMS-1151047).

References

- [Bergman 2015] G. M. Bergman, *An invitation to general algebra and universal constructions*, 2nd ed., Springer, Cham, 2015. MR 3309721 Zbl 1317.08001
- [Bhargava 2002] M. Bhargava, “Gauss composition and generalizations”, pp. 1–8 in *Algorithmic number theory* (Sydney, 2002), edited by C. Fieker and D. R. Kohel, Lecture Notes in Computer Science **2369**, Springer, Berlin, 2002. MR 2041069 Zbl 1058.11030
- [Bhargava 2006] M. Bhargava, “Higher composition laws and applications”, pp. 271–294 in *International Congress of Mathematicians* (Madrid, 2006), vol. 3, edited by M. Sanz-Solé et al., European Mathematical Society, Zürich, 2006. MR 2275598 Zbl 1129.11049
- [Biesel and Gioia 2015] O. Biesel and A. Gioia, “A new discriminant algebra construction”, preprint, 2015. arXiv 1503.05318v1

- [Burriss and Sankappanavar 2012] S. N. Burriss and H. P. Sankappanavar, “A course in universal algebra: the millennium edition”, 2012, available at <https://www.math.uwaterloo.ca/~snburriss/htdocs/ualg.html>.
- [Gilmer and Heitmann 1980] R. Gilmer and R. C. Heitmann, “On $\text{Pic}(R[X])$ for R seminormal”, *J. Pure Appl. Algebra* **16**:3 (1980), 251–257. MR 558489 Zbl 0428.13003
- [Hahn 1994] A. J. Hahn, *Quadratic algebras, Clifford algebras, and arithmetic Witt groups*, Springer, New York, 1994. MR 1249481 Zbl 0791.11021
- [Kanzaki 1973] T. Kanzaki, “On the quadratic extensions and the extended Witt ring of a commutative ring”, *Nagoya Math. J.* **49** (1973), 127–141. MR 0332778 Zbl 0238.13018
- [Kleiman 1979] S. L. Kleiman, “Misconceptions about K_X ”, *Enseign. Math.* (2) **25**:3–4 (1979), 203–206. MR 570309 Zbl 0432.14013
- [Knus 1991] M.-A. Knus, *Quadratic and Hermitian forms over rings*, Grundlehren der Math. Wissenschaften **294**, Springer, Berlin, 1991. MR 1096299 Zbl 0756.11008
- [Knus et al. 1998] M.-A. Knus, A. Merkurjev, M. Rost, and J.-P. Tignol, *The book of involutions*, American Mathematical Society Colloquium Publications **44**, American Mathematical Society, Providence, RI, 1998. MR 1632779 Zbl 0955.16001
- [Lenstra 2008] H. W. Lenstra, “Galois theory for schemes”, Lecture notes, 3rd ed., Universiteit Leiden, 2008, available at <http://websites.math.leidenuniv.nl/algebra/GSchemes.pdf>.
- [Loos 1996] O. Loos, “Tensor products and discriminants of unital quadratic forms over commutative rings”, *Monatsh. Math.* **122**:1 (1996), 45–98. MR 1397923 Zbl 0872.11022
- [Loos 2007] O. Loos, “Discriminant algebras of finite rank algebras and quadratic trace modules”, *Math. Z.* **257**:3 (2007), 467–523. MR 2328810 Zbl 1136.11031
- [McKenzie et al. 1987] R. N. McKenzie, G. F. McNulty, and W. F. Taylor, *Algebras, lattices, varieties*, vol. 1, Wadsworth and Brooks/Cole, Monterey, CA, 1987. MR 883644 Zbl 0611.08001
- [Rost 2002] M. Rost, “The discriminant algebra of a cubic algebra”, preprint, 2002, available at <http://www.math.uni-bielefeld.de/~rost/data/cub-disc.pdf>.
- [Small 1972] C. Small, “The group of quadratic extensions”, *J. Pure Appl. Algebra* **2**:2 (1972), 83–105. MR 0306170 Zbl 0242.13005
- [Stacks 2015] P. Belmans, A. J. de Jong, et al., “The Stacks project”, electronic reference, Columbia University, 2015, available at <http://stacks.math.columbia.edu>.
- [Voight 2011] J. Voight, “Rings of low rank with a standard involution”, *Illinois J. Math.* **55**:3 (2011), 1135–1154. MR 3069299 Zbl 06185698
- [Wood 2011] M. M. Wood, “Gauss composition over an arbitrary base”, *Adv. Math.* **226**:2 (2011), 1756–1771. MR 2737799 Zbl 1262.11049

Received May 8, 2015. Revised February 12, 2016.

JOHN VOIGHT
DEPARTMENT OF MATHEMATICS
DARTMOUTH COLLEGE
6188 KEMENY HALL
HANOVER, NH 03755
UNITED STATES
jvoight@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 283 No. 2 August 2016

The fundamental theorem of tropical differential algebraic geometry	257
FUENSANTA AROCA, CRISTHIAN GARAY and ZEINAB TOGHANI	
A simple solution to the word problem for virtual braid groups	271
PAOLO BELLINGERI, BRUNO A. CISNEROS DE LA CRUZ and LUIS PARIS	
Completely contractive projections on operator algebras	289
DAVID P. BLECHER and MATTHEW NEAL	
Invariants of some compactified Picard modular surfaces and applications	325
AMIR DŽAMBIĆ	
Radial limits of bounded nonparametric prescribed mean curvature surfaces	341
MOZHGAN (NORA) ENTEKHABI and KIRK E. LANCASTER	
A remark on the Noetherian property of power series rings	353
BYUNG GYUN KANG and PHAN THANH TOAN	
Curves with prescribed intersection with boundary divisors in moduli spaces of curves	365
XIAO-LEI LIU	
Virtual rational Betti numbers of nilpotent-by-abelian groups	381
BEHROOZ MIRZAI and FATEMEH Y. MOKARI	
A planar Sobolev extension theorem for piecewise linear homeomorphisms	405
EMANUELA RADICI	
A combinatorial approach to Voiculescu's bi-free partial transforms	419
PAUL SKOUFRANIS	
Vector bundle valued differential forms on $\mathbb{N}Q$ -manifolds	449
LUCA VITAGLIANO	
Discriminants and the monoid of quadratic rings	483
JOHN VOIGHT	



0030-8730(201608)283:2;1-V