

*Pacific
Journal of
Mathematics*

**RECOGNIZING RIGHT-ANGLED COXETER GROUPS
USING INVOLUTIONS**

CHARLES CUNNINGHAM, ANDY EISENBERG,
ADAM PIGGOTT AND KIM RUANE

RECOGNIZING RIGHT-ANGLED COXETER GROUPS USING INVOLUTIONS

CHARLES CUNNINGHAM, ANDY EISENBERG,
ADAM PIGGOTT AND KIM RUANE

We consider the question of determining whether or not a given group (especially one generated by involutions) is a right-angled Coxeter group. We describe a group invariant, the *involution graph*, and we characterize the involution graphs of right-angled Coxeter groups. We use this characterization to describe a process for constructing candidate right-angled Coxeter presentations for a given group or proving that one cannot exist. We apply this process to a number of examples. Our new results imply several known results as corollaries. In particular, we provide an elementary proof of rigidity of the defining graph for a right-angled Coxeter group, and we recover an existing result stating that if Γ satisfies a particular graph condition (called *no SILs*), then $\text{Aut}^0(W_\Gamma)$ is a right-angled Coxeter group.

1. Introduction	41
2. A summary of the recognition algorithm	45
3. Applications and results	50
4. Details	63
5. Further research	75
Acknowledgements	76
References	77

1. Introduction

Given a finite simple graph Γ , the *right-angled Coxeter group defined by Γ* is the group $W = W_\Gamma$ generated by the vertices of Γ . The relations of W_Γ declare that the generators all have order 2, and adjacent vertices commute with each other. Right-angled Coxeter groups (commonly abbreviated RACG) have a rich combinatorial and geometric history [Davis 2008]. The particular presentation specified by Γ is called a *right-angled Coxeter system*. When encountering a group generated by involutions, a natural question is to ask whether or not this group might be a right-angled Coxeter group, and if so, how to identify the preferred presentation.

MSC2010: primary 20F55, 20F65; secondary 05C75.

Keywords: Coxeter group, involutions, graph theory, automorphisms.

The main objective of this paper is the development of a recognition procedure that successfully answers this question for certain families of groups. Although the procedure may be applied more generally, our applications focus primarily on two classes of examples. Given a right-angled Coxeter group W_Γ , we consider

- (1) extensions of W_Γ by subgroups of $\text{Out}^0(W_\Gamma)$, and
- (2) subgroups of W_Γ generated by chosen sets of involutions.

(Recall that $\text{Aut}^0(W_\Gamma)$ consists of the automorphisms of W_Γ which map each generator to a conjugate of itself, and $\text{Out}^0(W_\Gamma)$ is the quotient $\text{Aut}^0(W_\Gamma)/\text{Inn}(W_\Gamma)$.) In each of these cases, we give examples of groups which are right-angled Coxeter and examples which are not. For those cases which are right-angled Coxeter, our procedure produces the preferred presentations. We show:

Theorem 1.1 (p. 57). *Suppose χ_1, \dots, χ_k are pairwise commuting partial conjugations of the right-angled Coxeter group W_Γ such that whenever χ_i and χ_j have the same acting letter, their domains don't intersect. Then $G = W \rtimes \langle \chi_1, \dots, \chi_k \rangle$ is a right-angled Coxeter group. Further, writing $S_i \subseteq \{\chi_1, \dots, \chi_k\}$ for the set comprising those partial conjugations with acting letter a_i , we have*

$$\left\{ a_1 \prod_{\chi_i \in S_1} \chi_i, \dots, a_n \prod_{\chi_i \in S_n} \chi_i \right\} \cup \{ \chi_1, \dots, \chi_k \}$$

is a Coxeter generating set for G .

If a group G has only 2-torsion, and G is not a right-angled Coxeter group, then G is not a Coxeter group. So our procedure may in fact enable one to show that a given group is not a Coxeter group. Cunningham [2015] has used some of the methods described here to show that $\text{Out}^0(W_n)$ for $n \geq 4$ is not a Coxeter group. (W_n is the *universal Coxeter group* whose defining graph has n vertices and no edges.)

Given a group G , the *involution graph* Δ_G of G is the group invariant defined as follows: the vertices in Δ_G correspond to the conjugacy classes of involutions in G ; vertices are adjacent when there exist commuting representatives of the corresponding conjugacy classes. In general, this invariant is unwieldy. It may be infinite, and even when it's finite, it may be impossible to construct. Nevertheless, for certain classes of groups the invariant promises insights. Like any invariant, it can allow us to distinguish between groups. It also carries information on the automorphism group of G . Since an automorphism must permute conjugacy classes of involutions and must preserve commuting relations, $\text{Aut}(G)$ acts naturally on Δ_G . The kernel of this action is therefore a natural normal subgroup of $\text{Aut}(G)$, and has finite index in $\text{Aut}(G)$ when Δ_G is finite.

The involution graph for a right-angled Coxeter group W_Γ is easily constructed directly from Γ : the vertices in Δ_W correspond to cliques in Γ ; vertices are adjacent

when the union of the corresponding cliques is also a clique. When constructed in this manner, we denote the graph Γ_K and call it the *clique graph* for Γ . Tits [1988] proved that the kernel of the action $\text{Aut}(W) \circlearrowleft \Delta_W$ has a natural complement, which is therefore a finite subgroup of $\text{Aut}(\Delta_W)$. Thus the involution graphs of right-angled Coxeter groups are significantly more tractable than the involution graphs of arbitrary groups, and may be more convenient for certain purposes than the defining graph Γ . Aaron Meyers, in his undergraduate thesis under the supervision of Piggott, began to explore some properties of clique graphs and how to recover their base graphs. (As this work is unpublished, new proofs are given in the following sections.)

The reader may compare our use of the clique graph and involution graph to the use of the *clique graph*, *extension graph*, and *commutation graph* in [Kim and Koberda 2013] in the context of right-angled Artin groups. Our use of the term and notation for the clique graph comes from that reference. In addition, Kim and Koberda define the *extension graph* Γ^e of Γ and the *commutation graph* of a subset $S \subset A(\Gamma)$ of elements of the right-angled Artin group. The vertices of Γ^e are the words in the right-angled Artin group $A(\Gamma)$ which are conjugate to a vertex of Γ , and two such vertices are connected by an edge if they commute with one another. More generally, the commutation graph of S has vertices given by the elements of S , and two of these are connected by an edge if they commute with each other.

It is straightforward to define the extension and commutation graphs in the context of right-angled Coxeter groups. Note that the vertices of Γ^e are the individual group elements, not conjugacy classes, so that Γ^e is infinite whereas Δ_{W_Γ} is finite. Moreover, Γ^e does not contain words that are only conjugate to a product of pairwise commuting generators, so it is not the case that Δ_{W_Γ} is a quotient graph of Γ^e . [Kim and Koberda 2013, Theorem 1.3] states that, given graphs Λ and Γ , if Λ is contained in Γ^e , then $A(\Lambda) \leq A(\Gamma)$. The analogous statement about right-angled Coxeter groups is certainly false, and a counterexample is provided by

$$D_\infty = W_2 = \langle a, b \mid a^2 = b^2 = 1 \rangle.$$

The defining graph Γ consists of two vertices with no edges. The extension graph Γ^e has countably many vertices and no edges, but D_∞ cannot contain subgroups which are free products of more than two copies of $\mathbb{Z}/2\mathbb{Z}$. If we replace the extension graph with the involution graph Δ_{W_Γ} in [Kim and Koberda 2013], the claim would still be false: Δ_{W_Γ} contains cliques which are larger than any clique in Γ .

Finally, we note that the involution graph Δ_G of a group which is not a right-angled Coxeter group may not be a commutation graph on any subset $\{g_1, \dots, g_n\}$ of elements. A priori, it could be the case that there is no single collection of elements, one from each conjugacy class, which simultaneously exhibit all commuting and noncommuting relationships dictated by the involution graph. (When W_Γ is a right-angled Coxeter group, Δ_{W_Γ} is the commutation graph on the set of products

of pairwise commuting generators.) It may be that the techniques of [Kim and Koberda 2013] could be adapted to the case of right-angled Coxeter groups, but as the current paper focuses on the recognition problem, we have not considered questions of embeddability.

In Section 2, we summarize our recognition procedure, which attempts to construct right-angled Coxeter presentations for a given group. This procedure relies on many facts about clique graphs and involution graphs which, for clarity of exposition, are only stated in that section. Detailed proofs have been relegated to Section 4 at the end of the paper. Section 2 contains all necessary definitions and results to understand the applications in Section 3.

In Section 3, we apply our procedure to several first examples of potential right-angled Coxeter groups. Section 3A collects examples of families of groups which are right-angled Coxeter. Γ is said to contain a *separating intersection of links* (SIL) if, for some pair of vertices v and w with $d(v, w) \geq 2$, there is a connected component of $\Gamma \setminus (\text{Lk}(v) \cap \text{Lk}(w))$ which contains neither v nor w . Otherwise, we say Γ contains no SILs. Section 3A also gives a new, shortened proof of [Charney et al. 2010, Theorem 3.6]: that $\text{Aut}^0(W_\Gamma)$ is right-angled Coxeter if Γ contains no SILs. Section 3B shows several examples of groups which we prove cannot be right-angled Coxeter. This includes, in particular, an iterated extension

$$\underbrace{(W_\Gamma \rtimes \mathbb{Z}/2\mathbb{Z}) \rtimes \mathbb{Z}/2\mathbb{Z}}_H^G$$

in which H is not right-angled Coxeter, but G is. We also note that $\text{Aut}^0(W_3)$ is not right-angled Coxeter, answering a motivating question for the authors.

Section 3C states some results that essentially identify features of a given graph Λ which indicate that W_Λ has a semidirect product decomposition $W_\Lambda = W_\Gamma \rtimes H$, where $H \leq \text{Out}^0(W_\Gamma)$. The results of this section follow from those in Section 3A quite easily, and the semidirect product decompositions are certainly not unique.

Section 4 presents detailed proofs for many facts stated without proof in Section 2. In this section, we present a characterization of those finite graphs which arise as clique graphs (i.e., a characterization of those graphs which arise as the involution graphs of right-angled Coxeter groups). We present a collapsing procedure to recover Γ from Γ_K , and we establish the correctness of our recognition procedure for constructing right-angled Coxeter presentations.

Finally, in Section 5 we give many follow-up questions which may be approachable using our recognition procedure. These include the question of characterizing those subgroups $H \leq \text{Out}^0(W_\Gamma)$ such that $W_\Gamma \rtimes H$ is again right-angled Coxeter, and determining when the involution graph of a subgroup $H \leq G$ can be calculated easily from the involution graph of G .

2. A summary of the recognition algorithm

In this section, we present the definitions and basic properties of the clique graph, star poset, and involution graph constructions. We state one of our main theorems characterizing those finite graphs which arise as clique graphs, and we describe a procedure which recovers a graph Γ from its clique graph Γ_K . Finally, we prove several algebraic results about right-angled Coxeter groups which allow us to modify this procedure to seek right-angled Coxeter presentations of a given group. Many of the proofs of this section are elementary or nongeometric in nature, so they have been pushed to [Section 4](#) at the end of the paper, where the interested reader will find all of the details. In this section, we present only the definitions and statements of results necessary to understand the applications in [Section 3](#).

A finite simple graph $\Gamma = (V, E)$ is an ordered pair of finite sets. We require that V , the set of *vertices*, is nonempty and E , the set of *edges*, consists of 2-element subsets of V . We say $a, b \in V$ are *adjacent* if $\{a, b\} \in E$. All graphs we consider in this paper will be undirected and have finitely many vertices, no loops, and no parallel edges. We will use the notation

$$\text{Lk}(v) = \{w \in V \mid \{v, w\} \in E\}$$

for the *link* of v and

$$\text{St}(v) = \text{Lk}(v) \cup \{v\}$$

for the *star* of v .

Definition 2.1. Let Γ be a graph. A *clique* in Γ is a nonempty subset of pairwise adjacent vertices. The *clique graph* of Γ is the graph $\Gamma_K = (V_K, E_K)$ whose vertices correspond to the cliques of Γ . Two vertices of Γ_K are adjacent if the union of the corresponding cliques in Γ is also a clique. [Figure 1](#) depicts an example.

The relation $v \sim w$ when $\text{St}(v) = \text{St}(w)$ is an equivalence relation on $V(\Gamma)$. Write $[v]$ for the equivalence class of v . Declaring that $[v] \leq [w]$ if $\text{St}(v) \subseteq \text{St}(w)$ we define a partial ordering, and we write $\mathcal{P}(\Gamma)$ for the poset of star-equivalence classes of vertices in Γ .




Figure 1. A graph Γ (left) and its corresponding clique graph Γ_K (right).

Throughout this paper, we will write $\Gamma_1, \Gamma_2, \dots, \Gamma_r$ for the maximal cliques of Γ . If $I \subset \{1, 2, \dots, r\}$, then

$$\Gamma_I = \bigcap_{i \in I} \Gamma_i$$

is the corresponding intersection of maximal cliques.

Definition 2.2. A vertex $v \in \Gamma$ is called *minimal* if it is contained in a unique maximal clique. Given $J \subset \{1, 2, \dots, r\}$, we say v is *J-minimal* if there is no $J' \supset J$ such that $\Gamma_{J'} \subsetneq \Gamma_J$ and $v \in \Gamma_{J'}$.

Theorem 2.3 (p. 64). *Let Γ' be a graph. There exists a graph Γ such that $\Gamma' = \Gamma_K$ if and only if the following three conditions are satisfied:*

(1) *Maximal clique condition (MCC): For all I , there exists some k_I such that*

$$|\Gamma'_I| = 2^{k_I} - 1.$$

(2) *Minimal vertex condition (MVC): Each nonempty intersection Γ'_J contains some J -minimal vertex v_J .*

(3) *Inclusion-exclusion condition (IEC): For each J ,*

$$\sum_{I \supseteq J} (-1)^{|I \setminus J|+1} k_I \leq k_J.$$

Moreover, if Γ' is a clique graph, then the graph Γ such that $\Gamma' = \Gamma_K$ is unique. The following procedure, which we call the *collapsing procedure*, recovers Γ from Γ' . We may write $\Gamma = C(\Gamma')$.

Theorem 2.4 (p. 68). *Let Γ' be a graph which satisfies the [MCC](#), [MVC](#), and [IEC](#). Then there is a unique (up to isomorphism) graph Γ such that Γ' is isomorphic to Γ_K . Moreover, the following collapsing procedure produces the graph Γ if it exists.*

(1) *Initially, let $V = \{\}$.*

(2) *Let $[w] \in \mathcal{P}(\Gamma')$ be a class such that every class $[v]$ with $[w] < [v]$ has already been considered. Write*

$$S_w = \bigcup_{[v] \geq [w]} [v].$$

Then there is some k such that $|S_w| = 2^k - 1$. Let k' be the number of vertices of S_w which are already contained in V . Choose $k - k'$ vertices of $[w]$ to add to the vertex set V .

(3) *Repeat the previous step until all classes of $\mathcal{P}(\Gamma')$ have been considered.*

(4) *Return the graph $C(\Gamma')$ which is the induced subgraph of Γ' on the vertex set V .*

The set S_w forms a clique in Γ' which is an intersection of maximal cliques, so its size has the desired form by the MCC. The details can be found in Section 4A.

Definition 2.5. Let G be a (finitely generated) group. The *involution graph* Δ_G of G is defined as follows. The vertices are the conjugacy classes of involutions in G . Two vertices $[x]$ and $[y]$ are connected by an edge if there exist representatives gxg^{-1} and hyh^{-1} that commute with each other.

We make a few remarks. The particular conjugates which witness commutativity are chosen for each edge individually. A system of representatives of each conjugacy class which act as witnesses for every edge simultaneously is called a *full system of representatives*. Such a system need not exist in general, but a right-angled Coxeter group will always have a full system of representatives.

We have also said earlier that all graphs we consider do not have loops, although the involution graph as defined here may contain a loop if an involution commutes with a conjugate of itself. This may happen in general, but it will never happen in a right-angled Coxeter group. So, if the involution graph of a given G contains a loop, we may immediately conclude that G is not a right-angled Coxeter group.

Lemma 2.6. *Let Γ be a graph. Then $\Delta_{W_\Gamma} = \Gamma_K$.*

Proof. It is a well-known fact about right-angled Coxeter groups that the only nontrivial torsion elements have order 2, and that any involution is conjugate to some product of pairwise commuting generators. The set of products of pairwise commuting generators forms a full system of representatives for the involution graph (this follows essentially from the deletion condition), and two such products commute if and only if all the generators involved in each product pairwise commute, i.e., if the collection of all these generators forms a clique in Γ . \square

We recover the rigidity of right-angled Coxeter groups as an immediate consequence. This was originally proven in [Green 1990] (for a more general class of groups), and many other proofs have been presented for different classes of groups containing right-angled Coxeter groups as a subclass; see, for example, [Droms 1987; Laurence 1995; Radcliffe 2003].

Corollary 2.7. *The defining graph of a right-angled Coxeter group W_Γ is unique up to isomorphism.*

Proof. The involution graph is an algebraic invariant (it does not depend on the chosen right-angled Coxeter presentation). By the previous lemma, the involution graph Δ_{W_Γ} is a clique graph, and by Theorem 2.3 the collapsed graph $C(\Delta_{W_\Gamma})$ is unique (up to isomorphism). \square

At this point, we can essentially describe our recognition procedure for seeking a right-angled Coxeter presentation for a given group G . First, we form the involution graph Δ_G . If this is not a clique graph, then G is not a right-angled Coxeter group.

If it is, then we must find a full system of representatives for the vertices. If such a system does not exist, then G is not a right-angled Coxeter group. If we find a full system of representatives, then the collapsing procedure will produce a labeled graph $\Gamma = C(\Delta_G)$, which gives a map $W_\Gamma \rightarrow G$ by sending the generators of W_Γ to the labels of the corresponding vertices. If we can show the candidate map is an isomorphism, then G is a right-angled Coxeter group, and the labels of Γ form a right-angled Coxeter generating set. (On the other hand, if the candidate map is not an isomorphism, we cannot conclude that G is not a right-angled Coxeter group. We may have simply chosen the wrong full system of representatives for Δ_G .)

We must address one subtlety in this procedure. In [Theorem 2.4](#), we chose vertices from $[w]$ to add to the vertex set V arbitrarily. It only mattered that we had the right number of vertices from each intersection of maximal cliques. In the algebraic setting, this is not sufficient, as the following simple example shows.

Example 2.8. Let Γ be a triangle with vertices a, b, c . Then $\Gamma_K = \Delta_{W_\Gamma}$ is a clique of size 7 with the labels a, b, c, ab, ac, bc, abc . In the star poset $\mathcal{P}(\Gamma_K)$, all vertices are equivalent, so there is only one $[w]$ to consider. The collapsing procedure says to choose 3 vertices from this class at random. If we choose, for example, the vertices a, b, c , then the collapsing procedure recovers Γ . If we choose a, ab, abc , then we find a new right-angled Coxeter presentation for W_Γ . However, if we pick a, b, ab , then we don't get a right-angled Coxeter presentation (because there is an additional relation between these vertices).

Essentially, at this step in the collapsing procedure we are choosing which vertices of the involution graph represent generators and which represent products of generators. There are (generally) many different ways that we can make this choice, but we have to make use of some algebraic information to avoid choosing products as if they were generators. The following results are certainly of independent interest, but we will, in particular, use them to make intelligent choices during the collapsing procedure.

Since we wish to avoid choosing vertices whose labels have a nontrivial product relation, it would certainly help if we could solve the word problem in G . However, depending on how G is presented, such a solution may or may not be evident (if it even exists). For this reason, we pass to the abelianization G^{ab} , in which there is a solution to the word problem. If G is a right-angled Coxeter group, then

$$G^{\text{ab}} \cong (\mathbb{Z}/2\mathbb{Z})^n,$$

and a product relation among involutions in G must also occur in G^{ab} .

From this point forward, for $g \in G$, we will write \bar{g} for the image of g in the abelianization. An important fact about right-angled Coxeter groups is that the abelianization is injective on conjugacy classes of involutions.

Proposition 2.9. *Let W_Γ be a right-angled Coxeter group. Let $x, y \in W_\Gamma$ such that $x^2 = y^2 = 1$. Then $\bar{x} = \bar{y}$ in W_Γ^{ab} if and only if x and y are conjugate in W_Γ .*

Proof. The “if” direction is trivial. Now, suppose x and y are not conjugate in W_Γ . Since x, y are involutions, there are pairwise commuting generators a_1, a_2, \dots, a_k , pairwise commuting generators b_1, b_2, \dots, b_ℓ , and words g, h such that

$$x = ga_1a_2 \cdots a_k g^{-1} \quad \text{and} \quad y = hb_1b_2 \cdots b_\ell h^{-1}.$$

Without loss of generality, since x and y are not conjugate, there is a b_j that does not appear among the a_i . But since it is a generator, there is a $\mathbb{Z}/2\mathbb{Z}$ direct factor in W_Γ^{ab} corresponding to that b_j . Therefore, \bar{y} will have a 1 in this factor and \bar{x} will have a 0. Thus, $\bar{x} \neq \bar{y}$ in W_Γ^{ab} . \square

Corollary 2.10. *For a right-angled Coxeter group W_Γ , if H is a subgroup generated by distinct, commuting involutions, then $H \cong H^{\text{ab}}$ injects into W_Γ^{ab} .*

Proof. H is a finite subgroup of W_Γ and so is conjugate to a special subgroup H' . Each element of H' is a distinct product of commuting generators from W_Γ and so each gets sent to a distinct element of W_Γ^{ab} . Thus, no two elements of H' can be conjugate in W_Γ and so neither can any two elements of H . By [Proposition 2.9](#), H injects into W_Γ^{ab} . \square

Proposition 2.11 (p. 72). *If W_Γ is a right-angled Coxeter group, then in step 2 of the collapsing procedure in [Theorem 2.4](#), we can choose the $k - k'$ involutions of W_Γ so that the chosen elements do not exhibit a nontrivial product relation.*

This proposition, which is proved in [Section 4B](#), makes use of the available algebraic information to amend our collapsing procedure and avoid nontrivial product relations. We can make further use of the available algebraic information to improve upon the procedure. In general, we have no particular method (or hope of finding a method) to construct Δ_G for an arbitrary G . Each of the following steps seem to be generally insurmountable:

- (1) Identify all involutions in G .
- (2) Separate all involutions into their conjugacy classes.
- (3) Determine the presence or lack of each edge in Δ_G (i.e., find a pair of commuting representatives or prove that none exist).
- (4) Find a full system of representatives.
- (5) Identify a full system of representatives so that the candidate maps are isomorphisms.

For a right-angled Coxeter system, it happens that all of these steps are not just possible, but straightforward.

Proposition 2.12 (p. 74). *If W_Γ is a right-angled Coxeter group, then two conjugacy classes of involutions $[x]$ and $[y]$ are connected by an edge in Δ_{W_Γ} if and only if there exists another class $[z]$ such that $\bar{z} = \bar{x}\bar{y}$ in the abelianization.*

If we are given a group G , supposing we can identify the conjugacy classes of involutions (i.e., the vertices of Δ_G), we can identify hypothetical edges and nonedges by looking for such \bar{z} in G^{ab} . If G is a right-angled Coxeter group, then this will produce the correct involution graph, and the remainder of the procedure will (hopefully, if we pick a good full system of representatives) identify a right-angled Coxeter presentation. On the other hand, if this not-quite involution graph of G is not a clique graph, we can be certain that G is not a right-angled Coxeter group. At no point do we directly need to check that we have calculated the true involution graph of G . We summarize this discussion with the following amended collapsing procedure. For details (including a full description of how to do these calculations in the abelianization), refer to [Section 4B](#).

Theorem 2.13 (p. 74). *Suppose G is a group whose only torsion elements all have order 2, so that $G^{\text{ab}} \cong (\mathbb{Z}/2\mathbb{Z})^n$ for some n . If the following procedure returns TRUE, then G is a right-angled Coxeter group (and the procedure indicates a right-angled Coxeter presentation). If the procedure returns FALSE, then G is not a right-angled Coxeter group.*

- (1) *Determine all conjugacy classes of involutions in G , and let these be the vertices of a graph Γ' . If there are not finitely many, return FALSE.*
- (2) *Apply [Proposition 2.12](#) to construct the edges of Γ' .*
- (3) *If Γ' is not a clique graph, return FALSE.*
- (4) *Find a full system of representatives for the vertices of Γ' . If no such system exists, return FALSE.*
- (5) *Collapse as in [Theorem 2.4](#), using [Proposition 2.11](#) to ensure that nontrivial product relations are avoided. Write $C(\Gamma')$ for the resulting graph.*
- (6) *Let Γ be a graph isomorphic to $C(\Gamma')$ with generic vertex labels a_1, \dots, a_n . Let $\varphi: W_\Gamma \rightarrow G$ be the map which sends the generators of W_Γ to the word given by the corresponding labels of vertices in $C(\Gamma')$. If φ is an isomorphism, return TRUE.*
- (7) *Otherwise, return UNKNOWN.*

3. Applications and results

In this section, we apply the recognition procedure from [Section 2](#) to seek out right-angled Coxeter presentations for certain families of groups. We focus in particular on

- (1) semidirect products of a given right-angled Coxeter group W_Γ by certain subgroups of $\text{Out}^0(W_\Gamma)$, and
- (2) subgroups of a given W_Γ generated by chosen subsets of involutions.

In particular, we note that the families of groups that we consider are already generated by involutions, have no torsion of order other than 2, and are usually given by presentations which are nearly right-angled Coxeter.

If D is a union of connected components of $\Gamma \setminus \text{St}(a_i)$ for some i , then the automorphism of W_Γ determined by

$$\chi_{i,D}(a_j) = \begin{cases} a_i a_j a_i, & a_j \in D, \\ a_j, & \text{otherwise,} \end{cases}$$

is called the *partial conjugation with acting letter a_i and domain D* . (Note that this terminology is not entirely consistent in the literature. Other papers have reserved partial conjugation for the case in which D is a single connected component [Gutierrez et al. 2012; Charney et al. 2010], while Laurence [1995] used the term *locally inner automorphism* before the term partial conjugation became common. We have preferred here to allow for multiple connected components in the domain of a partial conjugation, and we would propose the term *elementary partial conjugation* for the case in which D consists of a single connected component.) The partial conjugations generate $\text{Out}^0(W_\Gamma)$.

In Section 3A, we present families of groups which our procedure shows to be right-angled Coxeter. One example is worked out in full detail to demonstrate the procedure. For the remaining results, we simply state the resulting right-angled Coxeter group and the isomorphism determined by our procedure. The reader is left to verify the details. Most of these results are about split extensions of a given W_Γ by a finite subgroup of $\text{Out}^0(W_\Gamma)$ generated by (pairwise commuting) partial conjugations.

In Section 3B, we present families of groups which our procedure shows cannot be right-angled Coxeter. Again, one example is worked out in full detail. We note one example which is of particular interest: we find a group W_Γ with two elements $x, y \in \text{Out}^0(W_\Gamma)$ such that $G = W_\Gamma \rtimes \langle x, y \rangle$ is a right-angled Coxeter group, but $H = W_\Gamma \rtimes \langle xy \rangle$ is not. In particular, we can realize G as the iterated semidirect product

$$G = (W_\Gamma \rtimes \langle xy \rangle) \rtimes \langle x \rangle,$$

where each extension has degree 2. So this gives, to our knowledge, the first example in which the existence of a right-angled Coxeter presentation is lost and then recovered by semidirect product extensions.

Finally, in Section 3C, we note that many of our examples of right-angled Coxeter families arise as semidirect products. By analyzing the properties of

the defining graphs of the groups arising from these semidirect products, we can identify semidirect product decompositions in many cases. Such decompositions are generally not unique, and we cannot at the moment provide an exhaustive list of graph features of Γ which indicate a semidirect product decomposition of W_Γ .

3A. Groups which are right-angled Coxeter.

Example 3.1. We begin with an explicit example in which we demonstrate the recognition procedure in detail. Consider the defining graph in [Figure 2](#).

Write $x = \chi_{1,\{2\}}$ for the partial conjugation with acting letter a_1 and domain $\{a_2\}$. We consider the group $G = W_\Gamma \rtimes \langle x \rangle$, which has the presentation

$$G = \langle a_1, a_2, a_3, a_4, x \mid a_i^2 = x^2 = 1, [a_1, a_4] = [a_2, a_4] = [a_3, a_4] = 1, \\ [a_1, x] = [a_3, x] = [a_4, x] = 1, xa_2x = a_1a_2a_1 \rangle.$$

This is not quite a right-angled Coxeter presentation, so we apply our procedure to see if we can find one.

First, we compute G^{ab} (removing any relations that become trivial and understanding that group presentations with additive notation are assumed to be abelian):

$$G^{\text{ab}} = \langle \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{a}_4, \bar{x} \mid 2\bar{a}_i = 2\bar{x} = 0 \rangle \\ \cong \langle \bar{a}_1 \rangle \times \langle \bar{a}_2 \rangle \times \langle \bar{a}_3 \rangle \times \langle \bar{a}_4 \rangle \times \langle \bar{x} \rangle \cong (\mathbb{Z}/2\mathbb{Z})^5.$$

The relation matrix

$$\begin{pmatrix} 2 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix}$$

is already in Smith normal form, and so our canonical abelianization map $G \rightarrow G^{\text{ab}}$ is given by $g \mapsto \bar{g}$.

We now want to list all conjugacy classes of involutions in G . The classes of involutions in W_Γ are evident by inspection of Γ : a_i for each i , and a_ja_4 for each $1 \leq j \leq 3$. The new generator x is also an involution, and the products of x with the other generators that commute with it give new involutions: xa_1, xa_3, xa_4 . There are two remaining conjugacy classes of involutions, namely xa_1a_2 and $xa_1a_2a_4$.




Figure 2. The defining graph Γ .




Figure 3. The involution graph Δ_G .

These are all of the conjugacy classes of involutions in G . We could try to prove this directly, but it will also end up following from the fact that our procedure in this case does in fact construct an explicit isomorphism with a right-angled Coxeter group. Thus, we can omit the details.

We claim that the graph in [Figure 3](#) is the involution graph Δ_G . The given system of representatives is a full system, and the commuting relations are straightforward to check. (If they weren't as straightforward, we could easily construct the edge relations given by [Proposition 2.12](#).)

The brackets in the involution graph represent conjugacy classes. Since we now have a full system of representatives, we may stop writing these brackets. For the remainder of the calculation, brackets around a vertex label will denote its star-equivalence class. Before calculating the star poset structure, we observe that this graph clearly satisfies the [MCC](#) and [MVC](#), and the [IEC](#) is straightforward to verify.

The equivalence classes in the star poset are the following (identified by the dashed ellipses in [Figure 3](#)):

$$\begin{aligned}
 [a_1] &= \{a_1, a_1a_4\}, & [a_2] &= \{a_2, a_2a_4, xa_1a_2, xa_1a_2a_4\}, \\
 [a_3] &= \{a_3, a_3a_4, xa_3, xa_3a_4\}, & [a_4] &= \{a_4\}, \\
 [x] &= \{x, xa_4\}, & [xa_1] &= \{xa_1, xa_1a_4\}.
 \end{aligned}$$

The Hasse diagram for this poset is depicted in [Figure 4](#).

The element $[a_4]$ is maximal in the poset structure and contains a single element. We add a_4 to V . Next, we consider $[x]$ (or $[xa_1]$; the order in which we consider these classes is irrelevant). The clique above $[x]$ has size 3, so 2 of its vertices




Figure 4. The Hasse diagram for the poset $\mathcal{P}(\Delta_G)$.




Figure 5. The collapsed graph Λ (left) and an isomorphic graph with generic labels (right).

must be added to V . We have already added 1, so we must pick one more from $[x]$. Examining the abelianization, $\langle \bar{a}_4, \bar{x} \rangle \cong (\mathbb{Z}/2\mathbb{Z})^2$ and either of \bar{x} or $\bar{x}\bar{a}_4$ will extend \bar{a}_4 into a basis. So we choose to add x to V . Similarly, we consider $[xa_1]$ and add xa_1 to V .

The remaining three classes are all minimal. Suppose we take $[a_2]$ next. The clique above $[a_2]$ has size 7, so we must choose 3 elements from it. We have already chosen 2, so we need to choose 1 more. Checking the abelianization again, we see that any choice of the 4 elements in $[a_2]$ will extend to a basis, and so we add a_2 to V . Similarly, from $[a_3]$, we add a_3 to V .

Finally, we consider $[a_1]$. The clique above $[a_1]$ has size 7, and we have already chosen 3 of these vertices, so we choose no more. This leaves us with $V = \{a_2, a_3, a_4, x, xa_1\}$. We take the induced subgraph Λ of Δ_G on these vertices; see Figure 5.

We now have a candidate map $\varphi : W_\Lambda \rightarrow G$. It is straightforward to check that the map ψ below is the inverse, and that φ and ψ are isomorphisms:

$$\begin{aligned} \varphi : \quad & b_1 \mapsto a_3, & b_2 \mapsto x, & b_3 \mapsto a_4, & b_4 \mapsto xa_1, & b_5 \mapsto a_2, \\ \psi : \quad & a_1 \mapsto b_2b_4, & a_2 \mapsto b_5, & a_3 \mapsto b_1, & a_4 \mapsto b_3, & x \mapsto b_2. \end{aligned}$$

Thus, G is a right-angled Coxeter group, completing the example.

In this example, we were extending a right-angled Coxeter group by a single partial conjugation. It turns out that this will always yield a right-angled Coxeter group, and in fact we can say much more.

Lemma 3.2. *Suppose W_Γ is a right-angled Coxeter group. If $\alpha_1, \dots, \alpha_k$ are partial conjugations of W with the same acting letter and pairwise disjoint domains, then $G = W \rtimes \langle \alpha_1, \dots, \alpha_k \rangle$ is a right-angled Coxeter group.*

Proof. Without loss of generality, we may assume each α_j has acting letter a_1 . Let D_i denote the domain of α_i for each $1 \leq i \leq k$. Now G is generated by the elements

$$\{a_1, \dots, a_n, \alpha_1, \dots, \alpha_k\}$$

with the relations

- (R1) $a_i^2 = 1$, for $1 \leq i \leq n$,
- (R2) $[a_i, a_j] = 1$, for $\{a_i, a_j\} \in E(\Gamma)$,
- (R3) $\alpha_i^2 = 1$, for $1 \leq i \leq k$,
- (R4) $[\alpha_i, \alpha_j] = 1$, for $1 \leq i < j \leq k$,
- (R5) $[\alpha_i, a_j] = 1$, for $a_j \notin D_j$,
- (R6) $\alpha_i a_j \alpha_i = a_1 a_j a_1$, for $a_j \in D_i$.

Let H be the group generated by

$$\{b_1, \dots, b_n, \beta_1, \dots, \beta_k\}$$

with the relations

- (S1) $b_i^2 = 1$, for $1 \leq i \leq n$,
- (S2) $[b_i, b_j] = 1$, for $\{a_i, a_j\} \in E(\Gamma)$,
- (S3) $\beta_i^2 = 1$, for $1 \leq i \leq k$,
- (S4) $[\beta_i, \beta_j] = 1$, for $1 \leq i < j \leq k$,
- (S5) $[\beta_i, b_j] = 1$, for $a_j \notin D_i$,
- (S6) $[b_1, b_i] = 1$, for $2 \leq i \leq n$ and $a_i \in D_1 \cup \dots \cup D_k$.

We note that the given presentation for H is a right-angled Coxeter presentation. We define maps

$$\begin{aligned} \hat{\varphi} : \{a_1, \dots, a_n, \alpha_1, \dots, \alpha_k\} &\rightarrow \{b_1, \dots, b_n, \beta_1, \dots, \beta_k\}, \\ a_1 &\mapsto b_1 \beta_1 \cdots \beta_k, \\ \alpha_i &\mapsto \beta_i \quad (1 \leq i \leq k), \\ a_i &\mapsto b_i \quad (2 \leq i \leq n), \\ \hat{\psi} : \{b_1, \dots, b_n, \beta_1, \dots, \beta_k\} &\rightarrow \{a_1, \dots, a_n, \alpha_1, \dots, \alpha_k\}, \\ b_1 &\mapsto a_1 \alpha_1 \cdots \alpha_k, \\ \beta_i &\mapsto \alpha_i \quad (1 \leq i \leq k), \\ b_i &\mapsto a_i \quad (2 \leq i \leq n). \end{aligned}$$

It is straightforward to check that $\hat{\varphi}$ and $\hat{\psi}$ preserve the relations (R1)–(R6) and (S1)–(S6), respectively, so they induce homomorphisms $\varphi : G \rightarrow H$ and $\psi : H \rightarrow G$.

(Note that the preservation of the relation (S6) uses the assumption that the domains D_i are pairwise disjoint.) Finally, it is straightforward to see that φ and ψ are inverses to each other, hence G and H are isomorphic. That is, G is a right-angled Coxeter group. \square

Suppose $H \leq \text{Out}^0(W_\Gamma)$ is generated by partial conjugations χ_1, \dots, χ_k . Having shown that the semidirect product extension of W_Γ by any single partial conjugation is again right-angled Coxeter, we might hope to show that $W_\Gamma \rtimes H$ is right-angled Coxeter by observing that this is isomorphic to taking the iterated semidirect products, each by a single χ_i :

$$W_\Gamma \rtimes H = (\cdots ((W_\Gamma \rtimes \langle \chi_1 \rangle) \rtimes \langle \chi_2 \rangle) \rtimes \cdots \rtimes \langle \chi_k \rangle).$$

However, there is a subtlety that ruins this argument, namely, that χ_2 will extend to some automorphism of $W_\Gamma \rtimes \langle \chi_1 \rangle$, but not necessarily to a partial conjugation. We cannot extend inductively, since we cannot ensure that we are always extending by single partial conjugations. The following lemma and theorem identify certain cases in which this inductive argument works.

Lemma 3.3. *Suppose W , Γ , $a_1, \alpha_1, \dots, \alpha_k$, H , and G are as in the lemma and proof above. Let γ be a partial conjugation of W with acting letter $a_2 \neq a_1$ and such that γ commutes with each of the automorphisms $\alpha_1, \dots, \alpha_k$. Then γ acts on G as a partial conjugation.*

Proof. Without loss of generality we may assume γ has acting letter a_2 and domain D . Recall that $a_2 = b_2$. To show that γ acts on G as a partial conjugation we consider the result of conjugation by γ on each of the generators $b_1, \dots, b_n, \beta_1, \dots, \beta_k$. Firstly we note: $\gamma\beta_i\gamma = \beta_i$ for $1 \leq i \leq k$; $\gamma b_i\gamma = b_i$ for $1 \leq i \leq n$ and $a_i \notin D$; $\gamma b_i\gamma = b_2 b_i b_2$ for $2 \leq i \leq n$ and $a_i \in D$. If $a_1 \notin D$, then $\gamma b_1\gamma = \gamma a_1\gamma = b_1$. Suppose $a_1 \in D$. Since γ commutes pairwise with $\alpha_1, \dots, \alpha_k$, we have $a_2 \notin D_1 \cup \cdots \cup D_k$. We compute

$$\begin{aligned} \gamma b_1\gamma &= \gamma a_1 \alpha_1 \cdots \alpha_k \gamma \\ &= \gamma a_1 \gamma \alpha_1 \cdots \alpha_k \\ &= a_2 a_1 a_2 \alpha_1 \cdots \alpha_k \\ &= a_2 a_1 \alpha_1 \cdots \alpha_k a_2 \\ &= b_2 b_1 b_2. \end{aligned}$$

Since γ is an automorphism of G , and γ takes each generator to either itself or the conjugate of itself by b_2 , we may conclude that γ is a partial conjugation of G .

Write $\varphi : \{a_1, \dots, a_n\} \rightarrow \{b_1, \dots, b_n\}$ for the map $\varphi(a_i) = b_i$. From the calculations above, the domain of γ acting on G is $\varphi(D)$. \square

Theorem 1.1. *Suppose χ_1, \dots, χ_k are pairwise commuting partial conjugations of the right-angled Coxeter group W_Γ such that whenever χ_i and χ_j have the same acting letter, their domains don't intersect. Then $G = W \rtimes \langle \chi_1, \dots, \chi_k \rangle$ is a right-angled Coxeter group. Further, writing $S_i \subseteq \{\chi_1, \dots, \chi_k\}$ for the set comprising those partial conjugations with acting letter a_i ,*

$$\left\{ a_1 \prod_{\chi_i \in S_1} \chi_i, \dots, a_n \prod_{\chi_i \in S_n} \chi_i \right\} \cup \{\chi_1, \dots, \chi_k\}$$

is a Coxeter generating set for G .

Proof. The proof is by induction, applying the lemmas above at each step. Let $\alpha_1, \dots, \alpha_{k_1}$ be those χ_i with acting letter 1. By assumption, they have pairwise disjoint domains. By [Lemma 3.2](#), $W_\Gamma \rtimes \langle \alpha_1, \dots, \alpha_{k_1} \rangle$ is a RACG.

Moreover, by [Lemma 3.3](#), the remaining χ_i still act like partial conjugations, and their domains do not intersect, since they didn't before the extension. Now take $\beta_1, \dots, \beta_{k_2}$ among the remaining χ_i to be those which have acting letter 2, and extend by $\langle \beta_1, \dots, \beta_{k_2} \rangle$.

Continuing inductively, we extend at the i -th step by all remaining partial conjugations with acting letter i . The result follows. \square

In [[Gutierrez et al. 2012](#)], the authors investigate the automorphism groups of graph products of cyclic groups. In the case that W is a right-angled Coxeter group, the authors recover a result of Tits [[1988](#)] which shows $\text{Aut}(W) = \text{Aut}^0(W) \rtimes \text{Aut}^1(W)$ with $\text{Aut}^1(W)$ finite. Thus $\text{Aut}^0(W)$ (sometimes denoted $\text{Aut}^{\text{PC}}(W)$), which is the subgroup of $\text{Aut}(W)$ generated by all partial conjugations of W , is a finite index subgroup of $\text{Aut}(W)$. They also show that $\text{Aut}^0(W)$ splits as $\text{Inn}(W) \rtimes \text{Out}^0(W)$. Finally, they give the following condition on Γ , called *no SILs*, which characterizes exactly when $\text{Out}^0(W)$ is finite and is thus isomorphic to \mathbb{Z}_2^n .

Definition 3.4. A graph Γ has a *separating intersection of links (SIL)* if, for some vertices v and w with $d(v, w) \geq 2$, there is a component of $\Gamma \setminus (\text{Lk}(v) \cap \text{Lk}(w))$ which contains neither v nor w . Otherwise, Γ is said to have *no SILs*.

$\text{Inn}(W_\Gamma)$ is known to be a right-angled Coxeter group. In the case that Γ has no SILs, $\text{Aut}^0(W_\Gamma)$ is a finite extension of $\text{Inn}(W_\Gamma)$. In [[Charney et al. 2010](#)], it is shown that $\text{Aut}^0(W_\Gamma)$ is again a right-angled Coxeter group in that case. We arrive at this same result as a direct application of the previous corollary.

Corollary 3.5. *If Γ contains no SILs, then $\text{Aut}^0(W)$ is a right-angled Coxeter group and thus $\text{Aut}(W)$ contains a right-angled Coxeter group as a subgroup of finite index.*

Proof. Without loss of generality we may assume W has trivial center. Suppose Γ contains no SILs. Then

$$\text{Aut}^0(W) = \text{Inn}(W) \rtimes \text{Out}^0(W) \cong W \rtimes \text{Out}^0(W),$$

and $\text{Out}^0(W)$ is generated by pairwise commuting partial conjugations which satisfy the condition in the corollary above. \square

In general, one should not expect $\text{Aut}(W)$ to be right-angled Coxeter. The elements of $\text{Aut}^1(W)$ include graph symmetries, which could then introduce torsion elements of order other than 2. One should not generally expect that $\text{Aut}^0(W)$ is a right-angled Coxeter group, but one might see the no SILs result as suggesting that we restrict our attention to extensions of right-angled Coxeter groups by finite subgroups of $\text{Out}^0(W)$ (although [Example 3.8](#) in the following section demonstrates that even this restriction is not sufficient).

3B. Groups which are not right-angled Coxeter.

Example 3.6. As in the previous section, we begin with an explicitly worked out example. Let G denote the group presented as

$$G = \langle a, b, c, x, y \mid a^2, b^2, c^2, x^2, y^2, xax = a, xbx = b, xcx = aca, \\ yay = a, yby = b, ycy = bcb \rangle.$$

Let $W = \langle a, b, c \rangle$ and $H = \langle x, y \rangle$. Then $W = \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$, $H \cong \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$, and $G \cong W \rtimes H$, where x and y act as a pair of noncommuting partial conjugations.

To construct Δ_G , we must understand the involutions in G . Since $G = W \rtimes H$, each $g \in G$ may be written uniquely in the form $g = wh$, where $w \in W$ and $h \in H$. Further, $g^2 = whwh = whw(h^{-1}h)h = ww^{h^{-1}}h^2$. Since every element in G can be uniquely written as a product of an element of G and an element of H , if g is an involution, then h is an involution and $w^{h^{-1}} = w^h = w^{-1}$. Because H is a right-angled Coxeter group (in fact, D_∞), every nontrivial involution in H is conjugate to either x or y ; it follows that, up to conjugation, we may suppose g has one of the forms

- (1) w such that $w^2 = 1$,
- (2) wx such that $w^x = w^{-1}$, or
- (3) wy such that $w^y = w^{-1}$.

Every element of the first type is conjugate to either a , b , or c . Now we'll try to list elements of the second type (elements of the third type will be analogous).

Suppose $g = wx$ with $w^x = w^{-1}$. We further suppose that, within the collection of words of this form in the conjugacy class of g , we choose the shortest possible w . The element w can be written uniquely in the form $u_0bu_1b \cdots u_{m-1}bu_m$, where

$m \geq 0$, each u_i is a geodesic word in $\{a, c\}^*$, and only u_0 and u_m may be trivial. Then $w^x = u_0^x b u_1^x b \cdots u_{m-1}^x b u_m^x = w^{-1}$ implies that $u_0^x = u_m^{-1}$, $u_1^x = u_{m-1}^{-1}$, and so on. We now consider a few subcases.

If $m > 0$ and u_0 is not trivial, then

$$\begin{aligned}
 u_0^{-1}(wx)u_0 &= u_0^{-1}(u_0 b u_1 b \cdots u_{m-1} b u_m x)u_0 \\
 &= b u_1 b \cdots u_{m-1} b u_m u_0^x x \\
 &= b u_1 b \cdots u_{m-1} b x.
 \end{aligned}$$

This contradicts the minimality of the length of w , so either $m = 0$ or u_0 is trivial. If u_0 is trivial and $m > 1$, then w begins and ends with b , so $|b(wx)b| < |wx|$. Again, this contradicts minimality, hence either $m = 0$ or $w = b$.

If $m = 0$, then $w = u_0 \in \langle a, c \rangle$ is geodesic and so is an alternating string of a and c . If $|w| > 1$ and $|w|$ is odd, then w begins and ends with the same letter. If w begins and ends with a , then $|awxa| = |awax| < |wx|$; if w begins and ends with c then w^x begins and ends with a ; hence $w^x \neq w^{-1}$. In either case, we have a contradiction, so $|w| = 1$, in which case $w = a$ or $w = c$, or else $|w|$ is even. If $w = (ac)^n$ and $n > 1$, then $|aca(wx)aca| < |wx|$; if $w = (ca)^n$ and $n > 1$, then $|cwx| < |wx|$. In both cases, we have a contradiction. Our only case left is $m = 0$, $n = 1$, which corresponds to $w = ac$ or $w = ca$. Therefore, our only nontrivial possibilities for w are $w = b, a, c, ac, ca$.

Note that $a(cax)a = acx$, so these cases fall into the same conjugacy classes. In summary, we have that each involution of the form wx is conjugate to exactly one of the elements x, ax, bx, acx . (We observe that the final option cx is not, in fact, an involution. In this case, $w = c$, and $w^x \neq w^{-1}$.) We also observe that none of these involutions are conjugate to each other since they all map to distinct elements in G^{ab} .

Similarly, each involution of the form wy is conjugate to exactly one of the elements y, ay, by, bcy . Therefore, the following is the complete list of conjugacy classes in G , and hence serves as the list of vertex labels in Δ_G :

$$[a], [b], [c], [x], [ax], [bx], [acx], [y], [ay], [by], [bcy].$$

We now consider pairs of distinct conjugacy classes, to see whether or not they should be adjacent in Δ_G . By [Proposition 2.12](#), we can just check the product relations among the images of the involutions in G^{ab} . We omit the actual calculation and show the resulting involution graph in [Figure 6](#).

Now Δ_G is not a clique graph, since, for example, the [IEC](#) fails. (The reader can check this directly for the maximal cliques labeled Γ_3 and Γ_4 in the figure.)

Example 3.7. $\text{Aut}^0(W_3)$ is not a right-angled Coxeter group. The details are very similar to the previous example (we extend by one further partial conjugation), and are omitted here. The involution graph is shown in [Figure 7](#).




Figure 6. An involution graph which cannot be a clique graph. The labeled triangles Γ_i are the maximal cliques.




Figure 7. The involution graph for $\text{Aut}^0(W_3)$.

Here we must give the following warning. The proof above relies on finding a portion of the involution graph which we know should not appear in any clique graph. In the example, it is the “triangle of triangles” configuration (see [Example 4.2](#)). This should not occur in the involution graph of a right-angled Coxeter group, essentially because it means that all three vertices of the central triangle must be generators (whereas, by construction of the involution graph in the case of right-angled Coxeter groups, we should expect two of the vertices to be generators and the third to be their product).

However, we must point out that, strictly speaking, there is no such thing as a “poison pill” subgraph — a subgraph which, by its presence, prevents the given graph from being a clique graph. Indeed, if Γ is any graph, then Γ is an induced subgraph of Γ_K . In this way, any finite graph may appear as an induced subgraph in some clique graph (even the “triangle of triangles”). In the example above, it




Figure 8. The defining graph Γ .

is important that we know the central triangles Γ_3 and Γ_4 to be not just induced subgraphs, but also maximal cliques.

In all of the previous results, we have only considered split extensions by subgroups $H \leq \text{Out}^0(W_\Gamma)$ which were generated by partial conjugations. In particular, if the partial conjugations commuted pairwise, then H was finite and the extension $G = W_\Gamma \rtimes H$ was right-angled Coxeter. On the other hand, in the example above, the partial conjugations did not commute, thus H was infinite and G was not right-angled Coxeter. One might wonder whether the existence of a right-angled Coxeter presentation for the extension G depends only on the finiteness of H . The following example answers this question in the negative.

Example 3.8. Let Γ be the graph shown in Figure 8. Let x be the partial conjugation with acting letter a_1 and domain $\{a_3, a_4\}$, and let y be the partial conjugation with acting letter a_2 and domain $\{a_3, a_5\}$. Since a_1 and a_2 commute, so do x and y . Now write $z = xy$ for the product, which is also an involution. It follows from Theorem 1.1 that $G = W_\Gamma \rtimes \langle x, y \rangle$ is a right-angled Coxeter group. Consider the subgroup $H = W_\Gamma \rtimes \langle z \rangle \leq G$. The defining graph for G and the involution graph for H are shown in Figure 9.

The reader could verify Δ_H in two ways — first, by directly calculating the involutions and checking their commuting relations; and second, using the defining graph of G to calculate Δ_G , and then picking out the subset of vertices in Δ_G which are labeled by elements in the subgroup H . (Note that this latter method




Figure 9. The defining graph of G (left) and the involution graph of H (right).

of constructing the involution graph of a subgroup will not work in general. It works for the current example because G is a right-angled Coxeter group and H is normal.)

We can realize G as the iterated semidirect product

$$G = (W_\Gamma \rtimes \langle z \rangle) \rtimes \langle x \rangle = H \rtimes \langle x \rangle.$$

This gives an example of a right-angled Coxeter group W_Γ with a degree-2 split extension H which is not right-angled Coxeter. Moreover, taking a further degree-2 extension G , we recover the right-angled Coxeter property.

3C. Semidirect product decompositions. Here we present some results which are unrelated to the problem of recognizing right-angled Coxeter groups. These results fall naturally out of the applications in [Section 3A](#), and they generally address our ability to recognize semidirect product decompositions of W_Γ by identifying features of Γ .

To give the basic idea of how to generate these results, we give the following alternate description of [Lemma 3.2](#). Suppose a_1, \dots, a_n are the vertices of Γ and $\alpha_1, \dots, \alpha_k$ are partial conjugations as in the lemma. We will suppose that a_1 is the acting letter and D_i is the domain of α_i . The lemma says that the group $G = W_\Gamma \rtimes \langle \alpha_1, \dots, \alpha_k \rangle$ is a right-angled Coxeter group, and the proof of the lemma gives the right-angled Coxeter generating set. We can directly construct the defining graph Λ for G from Γ as follows:

- (1) Add k new vertices labeled $\alpha_1, \dots, \alpha_k$, all connected to one another and to a_1 .
- (2) Connect each α_i to every a_j where $a_j \notin D_i$.
- (3) Relabel a_1 as $a_1\alpha_1\alpha_2 \cdots \alpha_k$ and connect this to each vertex in $D_1 \cup D_2 \cup \cdots \cup D_k$.

The vertices $a_1, \alpha_1, \dots, \alpha_k$ form a clique of size $k + 1$, and the union of the stars of these vertices cover all of Λ . The restriction in [Lemma 3.2](#) that the domains be pairwise disjoint implies the following: we can distinguish D_i as those elements in $\text{St}(a_1) \setminus \text{St}(\alpha_i)$ which are contained in $\text{St}(\alpha_j)$ for every $j \neq i$. The following corollary is immediate from this description.

Corollary 3.9. *Suppose Λ contains $k + 1$ vertices $a_1, \alpha_1, \dots, \alpha_k$ satisfying: the following properties:*

- (1) $a_1, \alpha_1, \dots, \alpha_k$ form a clique.
- (2) $\text{St}(a_1) \cup \bigcup_i \text{St}(\alpha_i) = \Lambda$.
- (3) the sets $D_i = (\text{St}(a_1) \setminus \text{St}(\alpha_i)) \cap \bigcap_{j \neq i} \text{St}(\alpha_j)$ are all nonempty.

Define Γ to be the graph obtained from Λ by removing the vertices $\alpha_1, \dots, \alpha_k$ and any edge from a_1 to any D_i . Then W_Λ can be realized as the semidirect product




Figure 10. $W_\Lambda = W_{\Gamma_1} \rtimes \langle x \rangle = W_{\Gamma_2} \rtimes \langle y \rangle$, where x, y act like the partial conjugations $x = \chi_{4, \{1\}}$ and $y = \chi_{2, \{1\}}$ on Γ_1, Γ_2 , respectively.

$W_\Gamma \rtimes H$, where $H \leq \text{Out}^0(W_\Gamma)$ is generated by the partial conjugations with acting letter a_1 and domains D_i .

Theorem 1.1 yields an analogous corollary, since in each case they tell how to build the defining graph of the extension from the original defining graph, and the process is always reversible. It is not uniquely reversible. A given right-angled Coxeter group will, in general, have many semidirect product decompositions. As an example, consider the decompositions shown in [Figure 10](#).

4. Details

In this section we explore the properties of the clique graph, the star poset, and the involution graph introduced in [Section 2](#). We present detailed proofs of these properties, including proofs establishing claims made in that section and the correctness of our collapsing algorithms.

4A. The clique graph and the star poset. Recall that, given a graph Γ , we write Γ_I for the intersections of maximal cliques in Γ . We begin by establishing a correspondence between the *maximal clique structure* of a graph Γ and its clique graph Γ_K . By *maximal clique structure*, we mean that there is a bijection between the maximal cliques of Γ and those of Γ_K , which respects intersections.

Proposition 4.1. *Suppose Γ is a finite graph with maximal cliques $\Gamma_1, \dots, \Gamma_r$. For any subset $I \subseteq \{1, 2, \dots, r\}$, write*

$$\Gamma_I = \bigcap_{i \in I} \Gamma_i.$$

Similarly, write $\Gamma_{K,1}, \dots, \Gamma_{K,s}$ for the maximal cliques of Γ_K , and write $\Gamma_{K,I}$ for the intersections of maximal cliques. Then, possibly after reindexing:

- (1) $r = s$.

- (2) Each $\Gamma_{K,J}$ contains at least one J -minimal vertex (namely, the vertex labeled by the clique Γ_J).
- (3) $\Gamma_{K,i} = (\Gamma_i)_K$ (that is, $(\Gamma_i)_K$ naturally injects as a labeled graph into Γ_K , and the image is precisely $\Gamma_{K,i}$).
- (4) $\Gamma_{K,I} = (\Gamma_I)_K$.
- (5) If Γ_I is a clique of size k , then $\Gamma_{K,I}$ is a clique of size $2^k - 1$.

Proof. (1) For each maximal clique Γ_i in Γ , there is a corresponding vertex v_i in Γ_K . This vertex is adjacent only to vertices representing subsets of Γ_i since Γ_i is maximal, and so v_i is contained in the unique maximal clique $\text{St}(v_i)$ in Γ_K . In particular, since each v_i, v_j can be in the same maximal clique of Γ_K , we have $r \leq s$.

Conversely, each vertex of the maximal clique $\Gamma_{K,i}$ is labeled by some clique of vertices in Γ . Since $\Gamma_{K,i}$ forms a clique, the collection of all vertices of Γ which appear in the labels of vertices of $\Gamma_{K,i}$ must form a clique Λ in Γ . It is clear that Λ is maximal, since $\Gamma_{K,i}$ is. Thus $\Lambda = \Gamma_j$ for some j . That is, $s \leq r$, establishing (1). The description we have just given of the cliques in Γ_K also establishes the correspondence in (3), and therefore in (4).

As noted in the claim, the clique Γ_J forms a vertex of Γ_K . It is straightforward to see that this vertex is J -minimal in $\Gamma_{K,J}$, establishing (2).

Finally, if Γ_I is a clique of size k , then every nonempty subset of vertices induces a clique, and so corresponds to a vertex in $\Gamma_{K,I}$. There are $2^k - 1$ of these subsets, which correspond to $2^k - 1$ vertices in $\Gamma_{K,I}$. \square

Let Γ be a finite graph with maximal cliques $\Gamma_1, \dots, \Gamma_r$. As before, write Γ_I for the intersections of the maximal cliques, and suppose $|\Gamma_I| = k_I$. Then

$$\sum_{I \supseteq J} (-1)^{|I \setminus J|+1} k_I \leq k_J.$$

This is a direct application of the inclusion-exclusion principle, since the left hand side of the inequality counts the number of vertices in $\Gamma_J \cap \bigcup_{i \notin J} \Gamma_i$ (while the right-hand side is, by definition, the total number of vertices in Γ_J). We have therefore established that any clique graph must satisfy the **MCC**, **MVC**, and **IEC**. This gives one direction of the characterization theorem:

Theorem 2.3. *Let Γ' be a graph. There exists a graph Γ such that $\Gamma' = \Gamma_K$ if and only if the following three conditions are satisfied:*

- (1) *Maximal clique condition (MCC): For all I , there exists some k_I such that*

$$|\Gamma'_I| = 2^{k_I} - 1.$$

- (2) *Minimal vertex condition (MVC): Each nonempty intersection Γ'_J contains some J -minimal vertex v_J .*




Figure 11. The triangle $\{a, b, c\}$ forms a maximal clique which fails the IEC. This was essentially the feature of Example 3.6 which prevented the group there from being right-angled Coxeter.

(3) *Inclusion-exclusion condition (IEC):* For each J ,

$$\sum_{I \supseteq J} (-1)^{|I \setminus J|+1} k_I \leq k_J.$$

If we are faced with some graph which we do not know to be a clique graph, we can check directly that the intersections of maximal cliques have sizes of the form $n_I = 2^{k_I} - 1$, and we can check directly that the system of integers k_I satisfies the inclusion-exclusion inequalities. Thus, determining whether a graph arises as a clique graph is reduced to checking a system of integer inequalities (once we establish the other direction of the theorem).

Example 4.2. Consider the graph in Figure 11. In this graph, all intersections of maximal cliques have sizes of the form $2^k - 1$, but the IEC fails. So the graph cannot arise as a clique graph.

We will establish the converse of Theorem 2.3 by proving that, for any graph which satisfies the MCC, MVC, and IEC, the proposed collapsing procedure of Theorem 2.4 produces the desired output. In order to evaluate the collapsing procedure, we must explore some properties of the star poset $\mathcal{P}(\Gamma)$.

Lemma 4.3. *Let $[v] \in \mathcal{P}(\Gamma)$. Then the vertices*

$$S = \bigcup_{[v] \leq [w]} [w]$$

form a clique in Γ . If this clique is maximal, then $[v]$ is minimal in $\mathcal{P}(\Gamma)$.

Proof. If $w, w' \in S$ are any vertices, then $w \in \text{St}(v) \subseteq \text{St}(w')$, so w and w' are adjacent. Thus S forms a clique.

We now suppose $[v]$ is not minimal. Then there is some $[w] < [v]$. In particular, $w \notin S$, but $w \in \text{St}(w) \subseteq \text{St}(s)$ for any $s \in S$, hence w is a vertex outside of S adjacent to all of S . Thus S is not maximal. □

Definition 4.4. For $[v] \in \mathcal{P}(\Gamma)$, we call the clique S defined in the lemma the *clique above $[v]$* . We will use the notation S_v if we need to keep track of the vertex v .




Figure 12. It is easy to check that each vertex is its own star-equivalence class, and that these equivalence classes are pairwise not comparable. In particular, each $[v]$ is minimal, and each $S_v = \{v\}$ is not a maximal clique.

The converse of the lemma (i.e., that minimality of $[v]$ implies maximality of S_v) is false in general. A simple example is given in Figure 12. However, we claim that the converse does hold for those Γ which are clique graphs. Namely:

Proposition 4.5. *Suppose Γ satisfies the MVC. Then $[v]$ is a minimal element of $\mathcal{P}(\Gamma)$ if and only if v is a minimal vertex of Γ . In this case, S_v is the unique maximal clique containing v .*

Proof. Suppose v is a minimal vertex of Γ . Then $\text{St}(v)$ is the unique maximal clique containing v . Since S_v is a clique containing v , it is clear that $S_v \subseteq \text{St}(v)$. Conversely, if $x \in \text{St}(v)$, then $\text{St}(v) \subseteq \text{St}(x)$, hence $[v] \leq [x]$ and $x \in S_v$. Thus $\text{St}(v) = S_v$ is maximal. By the previous lemma, since S_v is maximal, $[v]$ is minimal.

Conversely, suppose v is not minimal. Then v is contained in the intersection of two distinct maximal cliques, Γ_1 and Γ_2 . Since Γ_i are maximal cliques, they contain minimal vertices w_i . By the above argument, $[w_i] \leq [v]$, and this must be a strict inequality since, e.g., $w_2 \in \text{St}(v) \setminus \text{St}(w_1)$. Thus $[v]$ is not minimal. \square

Proposition 4.6. *For any finite graph Γ and $[v] \in \mathcal{P}(\Gamma)$, S_v is an intersection of maximal cliques.*

Proof. Let $\Gamma_1, \dots, \Gamma_k$ be all the maximal cliques of Γ containing S_v . It is clear that $S_v \subseteq \bigcap \Gamma_i$.

Conversely, let $v' \in \bigcap \Gamma_i$ and suppose $v' \notin S_v$. Since $\text{St}(v) \not\subseteq \text{St}(v')$, there is some $x \in \text{St}(v)$ which is not in $\text{St}(v')$. In particular, since $\bigcup \Gamma_i \subseteq \text{St}(v')$, we must have $x \notin \Gamma_i$ for any i . By construction of S_v , we must have $x \in \text{St}(w)$ for each $w \in S_v$. Now $S_v \cup \{x\}$ forms a clique which contains S_v and is not equal to Γ_i for any i , contradicting our assumption that the list of Γ_i contained all maximal cliques containing S_v . So there can exist no such v' , hence $S_v = \bigcap \Gamma_i$, proving the claim. \square

We observe that the previous two propositions say the following in the case of clique graphs (which must satisfy the MVC):

Corollary 4.7. *Suppose Γ_K is a clique graph.*

- (1) $[v]$ is minimal in $\mathcal{P}(\Gamma_K)$ if and only if v is minimal in Γ_K .
- (2) If $[v]$ is nonminimal, then S_v is the intersection of maximal cliques (and therefore has size of the form $2^k - 1$). In this case,

$$S_v = \bigcap_{\substack{[w] \text{ minimal} \\ [w] \leq [v]}} S_w.$$

This shows that the star poset also records information about the intersections of maximal cliques: any clique above $[v]$ is such an intersection. Finally, we prove the converse.

Proposition 4.8. *Suppose Γ_K is a clique graph. Then any intersection of maximal cliques is equal to S_v for some v .*

Proof. Since Γ_K is a clique graph, it satisfies the MVC. Let $\Gamma_{K,J}$ be any intersection of maximal cliques, and let $v \in \Gamma_{K,J}$ be a J -minimal vertex. Without loss of generality, let J be the maximal index set without changing the intersection. In particular, J is precisely the index set of all maximal cliques containing v , so that $\text{St}(v) = \bigcup_{j \in J} \Gamma_{K,j}$.

We claim that $S_v = \Gamma_{K,J}$. Let $u \in S_v$. By definition of S_v , $[v] \leq [u]$, so

$$\text{St}(u) \supset \text{St}(v) = \bigcup_{j \in J} \Gamma_{K,j}.$$

That is, u is adjacent to every vertex in $\Gamma_{K,j}$, for each $j \in J$. Since each $\Gamma_{K,j}$ is a maximal clique, this shows $u \in \Gamma_{K,j}$ for each $j \in J$. That is, $u \in \Gamma_{K,J}$.

Conversely, let $w \in \Gamma_{K,J}$. Then w is adjacent to all vertices in $\Gamma_{K,j}$ for $j \in J$, thus $\bigcup_{j \in J} \Gamma_{K,j} \subseteq \text{St}(w)$. That is, $\text{St}(v) \subseteq \text{St}(w)$, so $[v] \leq [w]$. By its definition, S_v contains w . \square

We note that the previous proof gives a nice description of the elements of each star-equivalence class.

Corollary 4.9. *Suppose Γ_K is a clique graph. Then any $[v] \in \mathcal{P}(\Gamma_K)$ consists precisely of the J -minimal vertices of Γ_K , where J is the largest index set such that $v \in \Gamma_J$.*

Proof. Clearly, all J -minimal vertices for the same index set J must have the same star (namely, $\bigcup_{j \in J} \Gamma_{K,j}$). Conversely, suppose v is J -minimal and $[v] = [w]$ for some w . Then $w \in \Gamma_j$ for each $j \in J$, and $w \notin \Gamma_i$ for any $i \notin J$. (Otherwise, all of Γ_i would be in $\text{St}(w)$, which we have assumed to be equal to $\text{St}(v)$, a contradiction.) Therefore, w is J -minimal. \square

These results establish that, for a clique graph, the cliques above vertices are precisely the intersections of maximal cliques, and every intersection of maximal

cliques is the clique above some vertex. (This is not, in general, a bijective correspondence. As remarked earlier, it may be that $\Gamma_{K,J} = \Gamma_{K,J'} = S_v$, where $J \neq J'$.) In our collapsing algorithm to recover Γ from Γ_K , we begin at the top of the poset (this is the deepest intersection of maximal cliques) and work downwards. The previous proposition ensures that the algorithm examines every intersection of maximal cliques as it traverses every element in the poset structure.

We now wish to prove the correctness of our collapsing procedure, which also establishes the other direction of [Theorem 2.3](#). Recall the procedure:

Theorem 2.4. *Let Γ' be a graph which satisfies the [MCC](#), [MVC](#), and [IEC](#). Then there is a unique (up to isomorphism) graph Γ such that Γ' is isomorphic to Γ_K . Moreover, the following collapsing procedure produces the graph Γ if it exists.*

- (1) Initially, let $V = \{\}$.
- (2) Let $[w] \in \mathcal{P}(\Gamma')$ be a class such that every class $[v]$ with $[w] < [v]$ has already been considered. Write

$$S_w = \bigcup_{[v] \geq [w]} [v].$$

Then there is some k such that $|S_w| = 2^k - 1$. Let k' be the number of vertices of S_w which are already contained in V . Choose $k - k'$ vertices of $[w]$ to add to the vertex set V .

- (3) Repeat the previous step until all classes of $\mathcal{P}(\Gamma')$ have been considered.
- (4) Return the graph $C(\Gamma')$ which is the induced subgraph of Γ' on the vertex set V .

We first must address a subtlety, namely, that we can carry out the choice in step 2 of the algorithm.

Proposition 4.10. *In step 2 of the collapsing procedure, $0 \leq k - k' \leq |[w]|$. So we are able to choose an appropriate number of vertices from $[w]$ to add to V .*

Proof. The clique S_w is some intersection of maximal cliques Γ'_j by [Proposition 4.5](#). From this clique, we have already chosen k' vertices, and every vertex among those already chosen comes from a larger poset element, which is therefore a strictly smaller intersection of maximal cliques. By the [IEC](#), the number of elements we could have chosen is at most $k_J = k$, hence $k' \leq k$.

Now $S_w = (\bigcup_{[w] < [v]} S_v) \cup [w]$. Because

$$|S_w| = 2^k - 1, \quad \left| \bigcup_{[w] < [v]} S_v \right| \leq 2^{k'} - 1, \quad \text{and} \quad |[w]| \leq 2^{|[w]|},$$

we have $2^k - 1 \leq 2^{k'} - 1 + 2^{|[w]|}$. Therefore, $2^k \leq 2^{k'} + 2^{|[w]|}$. But $2^x + 2^y \leq 2^{x+y}$ for all pairs of positive integers x, y . Thus $2^k \leq 2^{k'+|[w]|}$, and $k \leq k' + |[w]|$. \square

We also see that step 2 does not tell us explicitly which vertices of $[w]$ to add to V . We claim this choice does not matter:

Proposition 4.11. *Given Γ' , if the procedure above does not return FALSE, then the isomorphism type of the graph Γ does not depend on the choices made in step 2 of the collapsing procedure.*

Proof. Without loss of generality, we will suppose our choices differ by a single vertex. Suppose we are about to consider $[v]$ and have constructed the set V thus far. Let $v_1, \dots, v_{k+1} \in [v]$, where $k > 0$ is the number of vertices from $[v]$ which we must add to V . Let

$$\begin{aligned} V_1 &= V \cup \{v_1, \dots, v_k\}, \\ V_2 &= V \cup \{v_1, \dots, v_{k-1}, v_{k+1}\}. \end{aligned}$$

We observe that we can make all future choices the same (since we haven't changed the number of vertices we must pick from $[w]$ for any $[w] \leq [v]$), so that we create two final graphs Γ_1 and Γ_2 whose vertex sets differ only by switching v_k and v_{k+1} .

We now claim that the resulting graphs Γ_1 and Γ_2 are isomorphic. By the previous observation, the vertex sets of Γ_1 and Γ_2 differ only by switching v_k and v_{k+1} . So we can define a map $\varphi : \Gamma_1 \rightarrow \Gamma_2$ which sends each vertex other than v_k to itself, and which sends v_k to v_{k+1} . We claim that φ defines a graph isomorphism. Clearly any adjacency relation not involving v_k is preserved under φ . Suppose w is a vertex of Γ_1 adjacent to v_k . Then $w \in \text{St}(v_k) = \text{St}(v_{k+1})$, so w is adjacent to v_{k+1} . Thus φ is a graph homomorphism. By the same argument, the analogous map $\psi : \Gamma_2 \rightarrow \Gamma_1$ is also a graph homomorphism, and the two maps are clearly inverses. Hence Γ_1 is isomorphic to Γ_2 . The full result follows by induction. \square

This shows that the isomorphism type of an output graph $C(\Gamma')$ is determined. However, a priori it could be the case that there are two graphs Γ, Λ such that Γ_K and Λ_K are isomorphic, but the collapsing procedure applied to Γ_K always outputs the isomorphism type Γ . The following proposition says that the maximal clique structure of Γ' determines the maximal clique structure of the output $C(\Gamma')$. The theorem following the proposition establishes that the maximal clique structure (including information about the sizes of all intersections of maximal cliques) determines a graph up to isomorphism. By [Proposition 4.1](#), any graph whose clique graph is Γ' will have the same clique graph structure, and will therefore be isomorphic. These results together show that the collapsing procedure outputs the unique graph Γ up to isomorphism so that $\Gamma_K = \Gamma'$.

Proposition 4.12. *Let Γ' be a finite graph satisfying the [MCC](#), [MVC](#), and [IEC](#). In particular, this implies there is a system of integers k_I such that $|\Gamma'_I| = 2^{k_I} - 1$. Let $C(\Gamma') = \Gamma$. Then the maximal cliques of Γ correspond to the maximal cliques of Γ' , and $|\Gamma_I| = k_I$ for all I .*

Proof. By assumption, each Γ'_I contains an I -minimal vertex v'_I . We have $|\Gamma'_I| = 2^{k_I} - 1$, and the algorithm chooses exactly k_I vertices from $S_{v'_I}$. [Corollary 4.7](#) implies that the maximal cliques in Γ have sizes of the form k_i , and [Proposition 4.8](#) ensures that we have $|\Gamma_I| = k_I$ for all intersections of maximal cliques (since all intersections Γ'_I occur as the clique above some element in the poset). \square

We have shown now that, if the algorithm returns any graph, then it returns a graph with a certain number of maximal cliques, and the intersections of the maximal cliques have certain sizes. We now establish that a finite graph is determined up to isomorphism by the sizes of the intersections of maximal cliques.

Theorem 4.13. *Let Γ, Λ be finite graphs. Suppose both graphs have r maximal cliques which may be indexed in such a way that, for all index sets $I \subset \{1, 2, \dots, r\}$, $|\Gamma_I| = |\Lambda_I|$. That is, all intersections of maximal cliques have the same sizes in each graph. Then there is an isomorphism $\varphi : \Gamma \rightarrow \Lambda$ which maps Γ_i to Λ_i for each i .*

Proof. We first claim that the poset structures $\mathcal{P}(\Gamma)$ and $\mathcal{P}(\Lambda)$ are the same, and the corresponding equivalence classes have the same sizes. For each $v \in \Gamma$, let J_v be the maximal index set such that $v \in \Gamma_{J_v}$. Then $\text{St}(v) = \bigcup_{j \in J_v} \Gamma_j$. The equivalence class of v consists of the J_v -minimal vertices of Γ by [Corollary 4.9](#). By assumption, $|\Gamma_{J_v}| = |\Lambda_{J_v}|$. Moreover, the number of vertices which are in some further intersection is given by the inclusion-exclusion formula:

$$\sum_{J \supseteq J_v} (-1)^{|J \setminus J_v|+1} |\Gamma_J| = \sum_{J \supseteq J_v} (-1)^{|J \setminus J_v|+1} |\Lambda_J|.$$

That is, the number of J_v -minimal vertices in Γ and in Λ is the same. Since this is for any v , the sizes of star-equivalence classes of vertices in Γ and Λ are equal for every class. Each equivalence class is represented by some index set J (although not every index set represents a class).

An equivalence class represented by J is smaller in the poset structure than another represented by J' if and only if $J \subseteq J'$. Since this holds in both Γ and Λ , it follows that the poset structures are equivalent.

Now, we build a map $\varphi : \Gamma \rightarrow \Lambda$ by piecing together (arbitrary) bijections between each pair of corresponding equivalence classes. We observe that, by construction,

$$\varphi([v]) = [\varphi(v)].$$

We also observe that Γ_i is mapped to Λ_i for each i . Let $v \in \Gamma_i$, so that $i \in J_v$. By construction, $\varphi(v) \in \Lambda_{J_v}$, which is an intersection of maximal cliques including Λ_i . That is, $\varphi(v) \in \Lambda_i$. It follows that φ maps Γ_I to Λ_I for each I .

We must show that φ preserves adjacency. Suppose $v, w \in \Gamma$ are adjacent. Then the edge $\{v, w\}$ extends to some maximal clique Γ_i . Now φ maps Γ_i to Λ_i , so $\varphi(v)$ and $\varphi(w)$ are still adjacent. \square

This completes the proofs of [Theorem 2.4](#) and [Theorem 2.3](#).

4B. Calculations in the abelianization. We now discuss the modifications to the collapsing procedure to make use of algebraic information. Recall from the discussion in [Section 2](#) that, given a group G , we first form the involution graph Δ_G and try to find a full system of representatives (i.e., a labeling of the vertices of Δ_G which exhibit all commuting relations simultaneously). If Δ_G is a clique graph, the collapsing procedure will give a graph $\Gamma = C(\Delta_G)$ such that $\Gamma_K = \Delta_G$. Moreover, Γ will carry the labels of the vertices chosen during the collapsing, so that the choice of which vertices to keep and which to omit is essentially the choice of which elements of G will be the generators in a (hypothetical) right-angled Coxeter presentation. For this reason, we must take care when choosing our generator vertices to avoid choosing group elements which have a nontrivial product relation. We will now demonstrate a method of passing to the abelianization G^{ab} to determine product relations using straightforward calculations.

Suppose we are given a finitely presented group

$$G = \langle s_1, \dots, s_m \mid r_1, \dots, r_k \rangle.$$

Recall that, for $g \in G$, we write \bar{g} for the image of g in the abelianization. A presentation for G^{ab} is given by

$$G^{\text{ab}} \cong \langle \bar{s}_1, \bar{s}_2, \dots, \bar{s}_m \mid \bar{r}_1, \bar{r}_2, \dots, \bar{r}_k, [\bar{s}_i, \bar{s}_j] \text{ for } 1 \leq i, j \leq m \rangle.$$

Writing the group operation additively in G^{ab} , we can write the relations as linear combinations of the generators with integer coefficients:

$$\bar{r}_i = a_{i,1}\bar{s}_1 + a_{i,2}\bar{s}_2 + \dots + a_{i,m}\bar{s}_m.$$

The coefficients $(a_{i,j})$ form a $k \times m$ matrix R , called the *relations matrix* for G^{ab} .

We briefly recall the *Smith normal form*. Given the $k \times m$ integer matrix R , there exist $k \times k$ and $m \times m$ invertible matrices P, Q and a diagonal matrix S such that $R = PSQ$, and the diagonal elements of S are $\alpha_1, \dots, \alpha_r, 0, \dots, 0$ such that $\alpha_i \mid \alpha_{i+1}$. The diagonal matrix S is called the *Smith normal form* of R .

Interpreting S as the relation matrix for a presentation, we have that G^{ab} is in a canonical form as a direct product of cyclic groups. Normal forms are immediate and computations in G^{ab} are much easier. Moreover, we now have an effective quotient map from $G \rightarrow G^{\text{ab}}$ in this canonical form. Namely, for any $g \in G$ with $g = \prod s_j$, we have $\bar{g} = \sum \bar{s}_j = \sum_{i=1}^m b_i \bar{s}_i$. The vector-matrix product $(b_1 \ b_2 \ \dots \ b_m)Q$ will give the coefficients of \bar{g} in the Smith normal form presentation of G^{ab} . This makes product relations easy to compute.

We now apply this method to show that, in step 2 of the collapsing procedure, we can avoid nontrivial product relations.

Proposition 2.11. *If W_Γ is a right-angled Coxeter group, then in step 2 of the collapsing procedure in Theorem 2.4, we can choose the $k - k'$ involutions of W_Γ so that the chosen elements do not exhibit a nontrivial product relation.*

Proof. In step 2 of our collapsing procedure, we consider an equivalence class $[w]$ of Δ_{W_Γ} and the clique above it, S_w , where $|S_w| = 2^k - 1$ for some k . If (W_Γ, S) is a right-angled Coxeter system for W_Γ and the labels are distinct, pairwise commuting involutions, then $H = S_w \cup \{e\}$ is a finite subgroup isomorphic to $(\mathbb{Z}/2\mathbb{Z})^k$: the elements of S_w are all involutions which pairwise commute. Any product g of these elements is an involution and commutes with all other elements of S_w (so it is connected to all of S_w). Moreover, any h which commutes with all of S_w commutes with any product of elements in S_w (namely g), and so g is contained in any maximal clique containing all of S_w . Since S_w is an intersection of maximal cliques and g is in all of these cliques, g lies in S_w . So H is a subgroup.

By Corollary 2.10, this subgroup projects injectively as a vector subspace into W_Γ^{ab} . Inductively, we assume that there exists a choice of a right-angled Coxeter system (W_Γ, S) such that \bar{V} is a set of *standard basis elements* for $(W_\Gamma, S)^{\text{ab}} \cong (\mathbb{Z}/2\mathbb{Z})^k$, i.e., each element has only one nonzero component in the representation for the abelianization given by our choice of right-angled Coxeter system (W_Γ, S) . (The base case is $\bar{V} = \emptyset$ and any choice of (W_Γ, S) .)

It follows that $\overline{V \cap S_w}$ is a linearly independent set in the $\mathbb{Z}/2\mathbb{Z}$ -vector space W_Γ^{ab} . We can then choose $k - k'$ labels in $S_w - V$ to extend this linearly independent set to a basis \bar{B} of $\langle \overline{S_w} \rangle$. (It's possible that $k - k' = 0$.) Since H projects injectively, choosing a basis for $\langle \overline{S_w} \rangle$ is the same as choosing a basis for $\langle S_w \rangle$. We need to show that $\bar{V} \cup \bar{B}$ is linearly independent as well.

To clarify, we are now keeping track of *two* different representations of the abelianization. W_Γ^{ab} is the form calculated from the Smith normal form in step 0 of the procedure, and $(W_\Gamma, S)^{\text{ab}}$ is the form wherein each element of \bar{V} is a standard basis element. We will show that such a form must exist if W_Γ is a right-angled Coxeter group, but it will not be directly computable during the procedure itself. The existence of this form will be used to show that *any* choice of \bar{B} during our procedure will result in no nontrivial product relations.

Since H is a finite subgroup of (W_Γ, S) , it is conjugate to a special subgroup: $gHg^{-1} = \langle a_1, a_2, \dots, a_k \rangle$ for $\{a_1, a_2, \dots, a_k\} \subseteq S$. Consider $b \in B \subseteq S_w$. Then, reordering the vertices of S if necessary, $gbg^{-1} = a_1 a_2 \cdots a_m$ in (W_Γ, S) . By the deletion condition of right-angled Coxeter groups (see, for example, [Davis 2008]), a product $c_1 c_2 \cdots c_\ell$ of distinct commuting generators of (W_Γ, S) commutes with $a_1 a_2 \cdots a_m$ if and only if c_j commutes with a_i for each i, j . In particular, $[b] = [a_1 a_2 \cdots a_m] \leq [a_i]$ for each $1 \leq i \leq m$.

Suppose that $[b] \not\leq [a_i]$ for each i . Then the procedure has already considered

$[a_i]$, and a subset of \overline{V} is a basis for $\langle \overline{S_{a_i}} \rangle$, which contains $\overline{a_i}$. But by our inductive hypothesis, \overline{V} is a set of standard basis elements relative to $(W_\Gamma, S)^{\text{ab}}$; moreover $\overline{a_i}$ is also a standard basis element since $a_i \in S$. So the only way that $\overline{a_i} \in \langle \overline{V} \rangle$ is if $\overline{a_i} \in \overline{V}$ (and so by injectivity $g^{-1}a_i g \in V$). Thus, $b = g^{-1}a_1a_2 \cdots a_m g \in \langle V \rangle$, and so b would *not* be chosen by the procedure to linearly extend \overline{V} .

Therefore, there must be some i such that $[b] = [a_i]$. By reordering the vertices of S if necessary, $[b] = [a_1]$. But then $gbg^{-1} = a_1a_2 \cdots a_m$ and a_1 are involutions that commute with exactly the same involutions, and so

$$\varphi : W_\Gamma \rightarrow W_\Gamma, \quad \varphi(a_j) = \begin{cases} a_1a_2 \cdots a_m & \text{if } j = 1, \\ a_j & \text{otherwise,} \end{cases}$$

is an involutive automorphism (in fact a transvection) of (W_Γ, S) .

Now, $(W_\Gamma, \varphi(S))$ is also a right-angled Coxeter system for W_Γ with the exact same generators except for swapping a_1 and the product $a_1a_2 \cdots a_m$. The set \overline{V} consists of standard basis elements *not* including $\overline{a_1}$ and so is unchanged under the induced map $\overline{\varphi} : (W_\Gamma, S)^{\text{ab}} \rightarrow (W_\Gamma, \varphi(S))^{\text{ab}}$. Alternatively, $\varphi(b) = \varphi(g)^{-1}a_1\varphi(g)$ and so $\overline{\varphi(b)} = \overline{a_1}$. So if we let $(W_\Gamma, S') = (W_\Gamma, \varphi(S))$ be our new right-angled Coxeter system and let $V' = V \cup \{b\}$ be our new subset of labels from our chosen full set of representatives of Δ_{W_Γ} , then the inductive hypothesis is still satisfied. In particular, in our Smith normal form W_Γ^{ab} , the set $\overline{V'}$ is still linearly independent.

For each $b \in B$, we can perform this procedure in succession, making sure that for each b we choose different a_i such that $[b] = [a_i]$. If at any point this were not possible, it would mean that there was some $b_n = g^{-1}a_1a_2 \cdots a_m g$ (in the updated system (W_Γ, S') with V') such that each a_j either satisfies

- (1) $[b_n] \leq [a_j]$, in which case $\overline{a_j} \in \overline{V'}$ from a previous step in the procedure, or
- (2) $\overline{b_l} = \overline{a_j}$ for some $l < n$, in which case $\overline{a_j} \in \overline{V'}$ from a previous element of the basis.

In either case, since all of the $\overline{a_j}$ lie in $\overline{S_w}$, this would give a linear dependence in $\overline{S_w}$ among \overline{B} , which contradicts its choice as a basis.

Thus, by induction on both elements of the poset, and then within each class on the elements of each chosen basis, it will always be the case that \overline{V} will consist of elementary basis elements in $(W_\Gamma, S)^{\text{ab}}$ for some choice of system (W_Γ, S) . Since every generator a_i of S is in S_{a_i} , we have $\overline{a_i} \in \langle \overline{V} \cap \overline{S_{a_i}} \rangle$, but since \overline{V} are all elementary basis vectors, it must be that $\overline{a_i} \in \overline{V}$. Thus, at the end of the procedure, \overline{V} will always be the full standard basis for some system $(W_\Gamma, S)^{\text{ab}}$, and in particular, \overline{V} will always be a basis of W_Γ^{ab} .

Any nontrivial product relation among the elements of V would induce a linear dependence among their images in W_Γ^{ab} . But since \overline{V} is a basis, this can never happen. \square

Finally, we prove the proposition that allows us to hypothetically build edges in the involution graph of a given group by doing calculations in the abelianization:

Proposition 2.12. *If W_Γ is a right-angled Coxeter group, then two conjugacy classes of involutions $[x]$ and $[y]$ are connected by an edge in Δ_{W_Γ} if and only if there exists another class $[z]$ such that $\bar{z} = \bar{x}\bar{y}$ in the abelianization.*

Proof. Let (W_Γ, S) be a right-angled Coxeter system for W_Γ , and let $[x]$ and $[y]$ be conjugacy classes of involutions. Since x and y are involutions in a right-angled Coxeter group, they are each conjugate to a product of commuting generators. So there exist $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_m \in S$ and $g, h \in W_\Gamma$ such that

$$gxg^{-1} = a_1a_2 \cdots a_n \quad \text{and} \quad hyh^{-1} = b_1b_2 \cdots b_m,$$

where all of the a_i pairwise commute, and all of the b_j pairwise commute. Consider the product

$$w = a_1a_2 \cdots a_nb_1b_2 \cdots b_m = c_1c_2 \cdots c_k,$$

where the c_ℓ are the generators that appear among either the a_i or the b_j but not both. (The ones that appear in both cancel with each other since they can be brought to the front or back of their respective words.) In the abelianization W_Γ^{ab} , we have $\bar{x} = \overline{a_1a_2 \cdots a_n}$, $\bar{y} = \overline{b_1b_2 \cdots b_m}$, and $\bar{w} = \overline{c_1c_2 \cdots c_k}$.

Now suppose that $[x]$ and $[y]$ are connected by an edge in Δ_{W_Γ} . That means that some conjugates of x and y commute. This implies that the product z of those conjugates is an involution. But then $\bar{z} = \bar{x}\bar{y}$ in W_Γ^{ab} .

Conversely, suppose that there exists an involution z such that $\bar{z} = \bar{x}\bar{y}$. Since z is an involution, it must be conjugate to a product of distinct, commuting generators, each of which is mapped to its corresponding generator of W_Γ^{ab} and so can be recovered directly from \bar{z} . Thus, these generators must be exactly the c_i , and so they each pairwise commute. In particular, w is an involution, and gxg^{-1} and hyh^{-1} commute. Thus, $[x]$ and $[y]$ should be connected by an edge in Δ_{W_Γ} . \square

We have now established the correctness of our right-angled Coxeter recognition procedure:

Theorem 2.13. *Suppose G is a group whose only torsion elements all have order 2, so that $G^{\text{ab}} \cong (\mathbb{Z}/2\mathbb{Z})^n$ for some n . If the following procedure returns TRUE, then G is a right-angled Coxeter group (and the procedure indicates a right-angled Coxeter presentation). If it returns FALSE, then G is not a right-angled Coxeter group.*

- (1) *Determine all conjugacy classes of involutions in G , and let these be the vertices of a graph Γ' . If there are not finitely many, return FALSE.*
- (2) *Apply [Proposition 2.12](#) to construct the edges of Γ' .*
- (3) *If Γ' is not a clique graph, return FALSE.*

- (4) Find a full system of representatives for the vertices of Γ' . If no such system exists, return FALSE.
- (5) Collapse as in [Theorem 2.4](#), using [Proposition 2.11](#) to ensure that nontrivial product relations are avoided. Write $C(\Gamma')$ for the resulting graph.
- (6) Let Γ be a graph isomorphic to $C(\Gamma')$ with generic vertex labels a_1, \dots, a_n . Let $\varphi : W_\Gamma \rightarrow G$ be the map which sends the generators of W_Γ to the word given by the corresponding labels of vertices in $C(\Gamma')$. If φ is an isomorphism, return TRUE.
- (7) Otherwise, return UNKNOWN.

5. Further research

While we have used our decision procedure to successfully establish both positive and negative identification of right-angled Coxeter presentations among extensions of right-angled Coxeter groups, much work remains to be done. One might hope to eventually characterize all subgroups $H \leq \text{Out}^0(W_\Gamma)$ (or $H \leq \text{Aut}(W_\Gamma)$) such that $W_\Gamma \rtimes H$ is right-angled Coxeter. We note that subgroups $H \leq \text{Out}^0(W_\Gamma)$ are not necessarily generated by partial conjugations (they may be generated by products of partial conjugations). Even if we only considered those H generated by partial conjugations, we could not extend [Lemma 3.2](#) by induction. If x, y are two commuting partial conjugations of W_Γ , then

$$W_\Gamma \rtimes \langle x, y \rangle \cong (W_\Gamma \rtimes \langle x \rangle) \rtimes \langle y \rangle;$$

however, y may not act on $W_\Gamma \rtimes \langle x \rangle$ as a partial conjugation (it will generally act as a product of partial conjugations). [Theorem 1.1](#) extends the lemma by induction, but we have many more examples of right-angled Coxeter extensions which are not covered by this theorem. More work is required for a complete characterization.

As in [Section 3C](#), following a characterization of extensions $W_\Gamma \rtimes H$ which are right-angled Coxeter, we would also gain insight into semidirect product decompositions of right-angled Coxeter groups. Given a graph Λ , we could hope to obtain a complete list of graph features which identify W_Λ as $W_\Gamma \rtimes H$, where $H \leq \text{Out}^0(W_\Gamma)$. (We observe that this would not identify all semidirect product decompositions of right-angled Coxeter groups. There are certainly decompositions which are not of this form.)

We strongly suspect that, whenever $H \leq \text{Out}^0(W_\Gamma)$ is isomorphic to D_∞ , the product $W_\Gamma \rtimes H$ is not right-angled Coxeter. The first example of [Section 3B](#) is of this form. Much of the argument in that example rests on using a normal form to establish that the given list of classes of involutions is complete. A general proof would require substantially more work to prove that we can accurately build the involution graph in the general case.




Figure 13. Dashed lines represent edges that may be present in some cases.

In particular, in the case of universal right-angled Coxeter groups (those whose defining graphs have no edges), the outer automorphism groups act on a contractible simplicial complex called McCullough–Miller space [Piggott 2012]. This space is analogous to Culler–Vogtmann outer space for the case of free groups [Culler and Vogtmann 1986], and we can use the action to classify all conjugacy classes of involutions in the outer automorphism groups. An analogous structure does not currently exist for the outer automorphism group of a general right-angled Coxeter group, and such a theory would need to be developed in order to construct the involution graph and confirm our conjecture.

Nevertheless, we can provide the following heuristic about what ought to go wrong in such an extension. Consider, for simplicity, a D_∞ generated by two noncommuting partial conjugations. If $x = \chi_{i,D}$ and $y = \chi_{j,E}$ are the partial conjugations, let b be any vertex other than a_j which is outside $\text{St}(a_i) \cup D$. Then Figure 13 shows part of the involution graph of the extension.

In the figure, the edge from b to ya_j will be present if $b \in E$; the edge from b to a_j will be present if $b \in \text{St}(a_j)$. The figure as drawn so far cannot be a clique graph, because the central triangle is a maximal clique which does not satisfy the IEC. But even if other vertices were present which could turn the central triangle into a 7-clique (or larger) so that the condition would be satisfied, the collapsing procedure would need to choose all three vertices x, a_i, xa_i , which are not linearly independent in the abelianization. However, this only establishes that the given pattern of labeling vertices in the involution graph — a pattern which has produced full systems of labels in all other examples so far — does not give an isomorphism to a right-angled Coxeter group in this case. We have not sufficiently established that the extension could not have any right-angled Coxeter presentation.

Acknowledgements

This work was partially supported by a grant from the Simons Foundation (#317466 to Adam Piggott). The authors would also like to thank Mauricio Gutierrez and the anonymous referee for a careful reading and helpful comments and suggestions.

References

- [Charney et al. 2010] R. Charney, K. Ruane, N. Stambaugh, and A. Vijayan, “The automorphism group of a graph product with no SIL”, *Illinois J. Math.* **54**:1 (2010), 249–262. [MR 2012f:20106](#) [Zbl 1243.20047](#)
- [Culler and Vogtmann 1986] M. Culler and K. Vogtmann, “Moduli of graphs and automorphisms of free groups”, *Invent. Math.* **84**:1 (1986), 91–119. [MR 87f:20048](#) [Zbl 0589.20022](#)
- [Cunningham 2015] C. Cunningham, *Automorphisms of right-angled Coxeter groups*, Ph.D. thesis, Tufts University, May 2015.
- [Davis 2008] M. W. Davis, *The geometry and topology of Coxeter groups*, London Mathematical Society Monographs Series **32**, Princeton University Press, Princeton, NJ, 2008. [MR 2008k:20091](#) [Zbl 1142.20020](#)
- [Droms 1987] C. Droms, “Isomorphisms of graph groups”, *Proc. Amer. Math. Soc.* **100**:3 (1987), 407–408. [MR 88e:20033](#) [Zbl 0619.20015](#)
- [Green 1990] E. Green, *Graph products of groups*, Ph.D. thesis, University of Leeds, 1990.
- [Gutierrez et al. 2012] M. Gutierrez, A. Piggott, and K. Ruane, “On the automorphisms of a graph product of abelian groups”, *Groups Geom. Dyn.* **6**:1 (2012), 125–153. [MR 2888948](#) [Zbl 1242.20041](#)
- [Kim and Koberda 2013] S.-h. Kim and T. Koberda, “Embedability between right-angled Artin groups”, *Geom. Topol.* **17**:1 (2013), 493–530. [MR 3039768](#)
- [Laurence 1995] M. R. Laurence, “A generating set for the automorphism group of a graph group”, *J. London Math. Soc.* (2) **52**:2 (1995), 318–334. [MR 96k:20068](#) [Zbl 0836.20036](#)
- [Piggott 2012] A. Piggott, “The symmetries of McCullough–Miller space”, *Algebra Discrete Math.* **14**:2 (2012), 239–266. [MR 3099973](#) [Zbl 1288.20033](#)
- [Radcliffe 2003] D. G. Radcliffe, “Rigidity of graph products of groups”, *Algebr. Geom. Topol.* **3** (2003), 1079–1088. [MR 2004h:20056](#) [Zbl 1053.20025](#)
- [Tits 1988] J. Tits, “Sur le groupe des automorphismes de certains groupes de Coxeter”, *J. Algebra* **113**:2 (1988), 346–357. [MR 89b:20077](#)

Received July 20, 2015. Revised December 9, 2015.

CHARLES CUNNINGHAM
 DEPARTMENT OF MATHEMATICS
 BOWDOIN COLLEGE
 BRUNSWICK, ME 04011
 UNITED STATES
ccunning@bowdoin.edu

ADAM PIGGOTT
 DEPARTMENT OF MATHEMATICS
 BUCKNELL UNIVERSITY
 LEWISBURG, PA 17837
 UNITED STATES
ap030@bucknell.edu

ANDY EISENBERG
 DEPARTMENT OF MATHEMATICS
 OKLAHOMA STATE UNIVERSITY
 STILLWATER, OK 74078
 UNITED STATES
andy.eisenberg@okstate.edu

KIM RUANE
 DEPARTMENT OF MATHEMATICS
 TUFTS UNIVERSITY
 MEDFORD, MA 02155
 UNITED STATES
kim.ruane@tufts.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 284 No. 1 September 2016

Bitwist manifolds and two-bridge knots	1
JAMES W. CANNON, WILLIAM J. FLOYD, LEER LAMBERT, WALTER R. PARRY and JESSICA S. PURCELL	
Recognizing right-angled Coxeter groups using involutions	41
CHARLES CUNNINGHAM, ANDY EISENBERG, ADAM PIGGOTT and KIM RUANE	
On Yamabe-type problems on Riemannian manifolds with boundary	79
MARCO GHIMENTI, ANNA MARIA MICHELETTI and ANGELA PISTOIA	
Quantifying separability in virtually special groups	103
MARK F. HAGEN and PRIYAM PATEL	
Conformal designs and minimal conformal weight spaces of vertex operator superalgebras	121
TOMONORI HASHIKAWA	
Coaction functors	147
S. KALISZEWSKI, MAGNUS B. LANDSTAD and JOHN QUIGG	
Cohomology and extensions of braces	191
VICTORIA LEBED and LEANDRO VENDRAMIN	
Noncommutative differentials on Poisson–Lie groups and pre-Lie algebras	213
SHAHN MAJID and WEN-QING TAO	