

*Pacific
Journal of
Mathematics*

**AFFINE WEAKLY REGULAR
TENSOR TRIANGULATED CATEGORIES**

IVO DELL'AMBROGIO AND DONALD STANLEY

AFFINE WEAKLY REGULAR TENSOR TRIANGULATED CATEGORIES

IVO DELL'AMBROGIO AND DONALD STANLEY

We prove that the Balmer spectrum of a tensor triangulated category is homeomorphic to the Zariski spectrum of its graded central ring, provided the triangulated category is generated by its tensor unit and the graded central ring is noetherian and regular in a weak sense. There follows a classification of all thick subcategories, and the result extends to the compactly generated setting to yield a classification of all localizing subcategories as well as the analog of the telescope conjecture. This generalizes results of Shamir for commutative ring spectra.

1. Introduction and results

Let \mathcal{K} be an essentially small tensor triangulated category, with symmetric exact tensor product \otimes and tensor unit object $\mathbf{1}$. Balmer [2005] defined a topological space, the *spectrum* $\mathrm{Spc} \mathcal{K}$, that allows for the development of a geometric theory of \mathcal{K} , similarly to how the Zariski spectrum captures the intrinsic geometry of commutative rings; see the survey [Balmer 2010b]. Among other uses, Balmer's spectrum encodes the classification of the thick tensor ideals of \mathcal{K} in terms of certain subsets. It is therefore of interest to find an explicit description of the spectrum in the examples, but this is usually a difficult problem requiring some in-depth knowledge of each example at hand.

The goal of this note is to show that in some cases a concrete description of the spectrum can be obtained easily and completely formally. Let us denote by

$$R := \mathrm{End}_{\mathcal{K}}^*(\mathbf{1}) = \bigoplus_{i \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{K}}(\mathbf{1}, \Sigma^i \mathbf{1})$$

the graded endomorphism ring of the unit, where $\Sigma : \mathcal{K} \rightarrow \mathcal{K}$ is the suspension functor. In the terminology of [Balmer 2010a], this is the *graded central ring* of \mathcal{K} . It is a graded commutative ring and therefore we can consider its spectrum

Dell'Ambrogio was partially supported by the Labex CEMPI (ANR-11-LABX-0007-01) .

MSC2010: primary 18E30; secondary 55P42, 55U35.

Keywords: tensor triangulated category, thick subcategory, localizing subcategory, spectrum.

of homogeneous prime ideals, $\text{Spec } R$, equipped with the Zariski topology. As established in that paper, there is always a canonical continuous map

$$\rho : \text{Spc } \mathcal{K} \rightarrow \text{Spec } R$$

comparing the two spectra. Under some mild hypotheses, e.g., when R is noetherian, ρ can be shown to be surjective, but it is less frequently injective and, when it is, the proof of injectivity is typically much harder.

Here is our main result:

Theorem 1.1. *Assume that \mathcal{K} satisfies the two following conditions:*

- (a) \mathcal{K} is classically generated by $\mathbf{1}$, i.e., as a thick subcategory: $\text{Thick}(\mathbf{1}) = \mathcal{K}$.
- (b) R is a (graded) noetherian ring concentrated in even degrees and, for every homogeneous prime ideal \mathfrak{p} of R , the maximal ideal of the local ring $R_{\mathfrak{p}}$ is generated by a (finite) regular sequence of homogeneous non-zero-divisors.

Then the comparison map $\rho : \text{Spc } \mathcal{K} \xrightarrow{\sim} \text{Spec } R$ is a homeomorphism.

As in the title, we may refer to a tensor triangulated category \mathcal{K} satisfying hypotheses (a) and (b) as being *affine* and *weakly regular*, respectively. Note that R being noetherian implies that $R^0 = \text{End}_{\mathcal{K}}(\mathbf{1})$ is a noetherian ring and that R is a finitely generated R^0 -algebra, by [Goto and Yamagishi 1983].

The next result is an easy consequence of the theorem. Here $\text{Supp}_R H^*X$ denotes the (big) Zariski support of the cohomology graded R -module $H^*X := \text{Hom}_{\mathcal{K}}^*(\mathbf{1}, X)$.

Corollary 1.2. *If \mathcal{K} and R are as in the theorem, then there exists a canonical inclusion-preserving bijection*

$$\{\text{thick subcategories } \mathcal{C} \text{ of } \mathcal{K}\} \xleftrightarrow{\sim} \{\text{specialization closed subsets } V \text{ of } \text{Spec } R\}$$

mapping a thick subcategory \mathcal{C} to $V = \bigcup_{X \in \mathcal{C}} \text{Supp}_R H^*X$ and a specialization closed subset V to $\mathcal{C} = \{X \in \mathcal{K} \mid \text{Supp}_R H^*X \subseteq V\}$.

In many natural examples, \mathcal{K} occurs as the subcategory \mathcal{T}^c of compact objects in a compactly generated tensor triangulated category \mathcal{T} . By the latter we mean a compactly generated triangulated category \mathcal{T} equipped with a symmetric monoidal structure \otimes which preserves coproducts and exact triangles in both variables, and such that the compact objects form a tensor subcategory \mathcal{T}^c (that is, $\mathbf{1}$ is compact and the tensor product of two compact objects is again compact).

In this case, the same hypotheses allow us to classify also the localizing subcategories of \mathcal{T} , thanks to the stratification theory of compactly generated categories due to Benson, Iyengar and Krause [Benson et al. 2011]. The *support* $\text{supp}_R X \subseteq \text{Spec } R$ of an object $X \in \mathcal{T}$ is defined in [Benson et al. 2008], and can be described as the set

$$\text{supp}_R X = \{\mathfrak{p} \in \text{Spec } R \mid X \otimes K(\mathfrak{p}) \neq 0\},$$

where the *residue field object* $K(\mathfrak{p})$ of a prime ideal \mathfrak{p} is an object whose cohomology is the graded residue field of R at \mathfrak{p} ; see [Section 3](#).

Theorem 1.3. *Let \mathcal{T} be a compactly generated tensor triangulated category with compact objects $\mathcal{K} := \mathcal{T}^c$ and graded central ring R satisfying conditions (a) and (b). Then we have the following canonical inclusion-preserving bijection:*

$$\{\text{localizing subcategories } \mathcal{L} \subseteq \mathcal{T}\} \xrightleftharpoons{\sim} \{\text{subsets } S \subseteq \text{Spec } R\}.$$

The correspondence sends a localizing subcategory \mathcal{L} to $S = \bigcup_{X \in \mathcal{L}} \text{supp}_R X$, and an arbitrary subset S to $\mathcal{L} = \{X \in \mathcal{T} \mid \text{supp}_R X \subseteq S\}$. Moreover, the bijection restricts to localizing subcategories $\mathcal{L} = \text{Loc}(\mathcal{L} \cap \mathcal{K})$ which are generated by compact objects on the left and to specialization closed subsets $S = \bigcup_{\mathfrak{p} \in S} V(\mathfrak{p})$ on the right.

Note that here the affine condition (a) is equivalent to requiring that \mathcal{T} is generated by $\mathbf{1}$ as a localizing subcategory. As $\text{Supp}_R H^*X = \text{supp}_R X$ for all compact objects $X \in \mathcal{K}$, one sees easily that in the compactly generated case [Theorem 1.1](#) and [Corollary 1.2](#) are also a consequence of [Theorem 1.3](#).

The next corollary is another byproduct of stratification. Recall that a localizing subcategory $\mathcal{L} \subseteq \mathcal{T}$ is *smashing* if the inclusion functor $\mathcal{L} \hookrightarrow \mathcal{T}$ admits a coproduct-preserving right adjoint.

Corollary 1.4 (the telescope conjecture in the affine weakly regular case). *In the situation of [Theorem 1.3](#), every smashing subcategory of \mathcal{T} is generated by a set of compact objects of \mathcal{T} .*

A few special cases of our formal results had already been observed, such as when R is even periodic and of global dimension at most one; see [[Dell’Ambrogio and Tabuada 2012](#)]. We now consider some more concrete examples.

* * *

Example 1.5. Let A be a commutative dg algebra and $D(A)$ its derived category of dg modules. Then $D(A)$ is an affine compactly generated tensor triangulated category with respect to the standard tensor product $\otimes = \otimes_A^L$, and $R = H^*A$ is the cohomology algebra of A ; thus if the latter satisfies (b) all our results apply to $D(A)$. Actually, in this example we can improve our results a little by eliminating the hypothesis that R is even and that the elements of the regular sequences are non-zero-divisors:

Theorem 1.6. *Let A be a commutative dg algebra such that its graded cohomology ring $R = H^*A$ is noetherian and such that every local prime $\mathfrak{p}R_{\mathfrak{p}}$ is generated by a finite regular sequence. Then all the conclusions of [Theorems 1.1](#) and [1.3](#) and of [Corollaries 1.2](#) and [1.4](#) hold for $\mathcal{T} = D(A)$ and $\mathcal{K} = D(A)^c$.*

We can apply this, for instance, to a graded polynomial algebra with any choice of grading for the variables, seen as a strictly commutative formal dg algebra.

Example 1.7. Let A be a commutative S -algebra (a.k.a. commutative highly structured ring spectrum), and let $D(A)$ be its derived category. (This covers [Example 1.5](#), as commutative dga's can be seen as commutative S -algebras.) Then $D(A)$ is an affine compactly generated tensor triangulated category, and $R = \pi_* A$ is the stable homotopy algebra of A ; thus if the latter satisfies [\(b\)](#) all our results apply to $D(A)$. Shamir [\[2012\]](#) already treated this example under the additional hypothesis that $\pi_* A$ has finite Krull dimension. Working with ∞ -categories and E_∞ -rings, Mathew [\[2015, Theorem 1.4\]](#) established the classification of thick subcategories as in [Corollary 1.2](#) for the case when $\pi_* A$ is even periodic and $\pi_0 A$ regular noetherian. Remarkably, in the special case of S -algebras defined over \mathbb{Q} , Mathew was also able to prove the classification of thick subcategories only assuming $\pi_* A$ noetherian, i.e., without any regularity hypothesis; see [\[Mathew 2016, Theorem 1.4\]](#). (Note however that, thanks to [\[Mandell 2012\]](#), in order to apply our own results we really only need an E_4 -structure on a ring spectrum rather than a fully commutative E_∞ -structure.)

The next two well-known examples show that neither hypothesis [\(a\)](#) nor [\(b\)](#) can be weakened with impunity.

Example 1.8. The derived category $\mathcal{T} = D(\mathbb{P}_k^1)$ of the projective line over a field k is an example where $R = \text{End}^*(\mathbf{1}) \simeq k$ certainly satisfies [\(b\)](#) but [\(a\)](#) does not hold. Indeed ρ can be identified with the structure map $\mathbb{P}_k^1 \rightarrow \text{Spec } k$ and is therefore far from injective in this case; see [\[Balmer 2010a, Remark 8.2\]](#).

Example 1.9. If $\mathcal{T} = D(A)$ is the derived category of a commutative (ungraded) ring A , [Theorem 1.1](#) and the classification of thick subcategories always hold by a result of Thomason [\[1997\]](#) (see [\[Balmer 2010a, Proposition 8.1\]](#)); the classification of localizing subcategories and the telescope conjecture hold if A is noetherian by [\[Neeman 1992a\]](#). On the other hand, Keller [\[1994\]](#) found examples of *nonnoetherian* rings A for which the two latter results fail.

In view of these examples, it would be interesting to know how far our weak regularity hypothesis [\(b\)](#) can be weakened in general. Would noetherian suffice?

2. Preliminaries

Let \mathcal{K} be an essentially small tensor triangulated category.

For any two objects $X, Y \in \mathcal{K}$, consider the \mathbb{Z} -graded group $\text{Hom}_{\mathcal{K}}^*(X, Y) = \bigoplus_{i \in \mathbb{Z}} \text{Hom}_{\mathcal{K}}(X, \Sigma^i Y)$. Recall that the symmetric tensor product of \mathcal{K} canonically induces on $R := \text{Hom}_{\mathcal{K}}^*(\mathbf{1}, \mathbf{1})$ the structure of a graded commutative¹ ring, and

¹To be precise, graded commutativity means here that $fg = \epsilon^{|f||g|}gf$ for any two homogeneous elements $f \in \text{Hom}_{\mathcal{K}}(\mathbf{1}, \Sigma^{|f|}\mathbf{1})$ and $g \in \text{Hom}_{\mathcal{K}}(\mathbf{1}, \Sigma^{|g|}\mathbf{1})$, where $\epsilon \in R^0$ is a constant with $\epsilon^2 = 1$ induced by the symmetry isomorphism $\Sigma \mathbf{1} \otimes \Sigma \mathbf{1} \xrightarrow{\sim} \Sigma \mathbf{1} \otimes \Sigma \mathbf{1}$. In most cases we have $\epsilon = -1$, e.g., if \mathcal{K} admits a symmetric monoidal model, but usually no extra difficulty arises by allowing the general case. Of course, this is immaterial for R even.

on each $\text{Hom}_{\mathcal{K}}^*(X, Y)$ the structure of a (left and right) graded R -module. The composition of maps in \mathcal{K} and the tensor functor $-\otimes-$ are (graded) bilinear for this action. See [Balmer 2010a, §3] for details.

Since we are using cohomological gradings, we write H^*X for the R -module $\text{Hom}_{\mathcal{K}}^*(\mathbf{1}, X)$ and call it the *cohomology of X* .

Supports for graded modules. We denote by $\text{Spec } R$ the Zariski spectrum of all homogeneous prime ideals in R . If M is an R -module (always understood to be graded) and $\mathfrak{p} \in \text{Spec } R$, the graded localization of M at \mathfrak{p} is the R -module $M_{\mathfrak{p}}$ obtained by inverting the action of all the *homogeneous* elements in $R \setminus \mathfrak{p}$. The *big support* of M is the following subset of the spectrum:

$$\text{Supp}_R M = \{\mathfrak{p} \in \text{Spec } R \mid M_{\mathfrak{p}} \neq 0\}.$$

Since our graded ring R is noetherian we also dispose of the *small support*, defined in terms of the indecomposable injective R -modules $E(R/\mathfrak{p})$:

$$\text{supp}_R M = \{\mathfrak{p} \mid E(R/\mathfrak{p}) \text{ occurs in the minimal injective resolution of } M\}.$$

We recall from [Benson et al. 2008, §2] some well-known properties of supports. In general we have $\text{supp}_R M \subseteq \text{Supp}_R M$. If M is finitely generated, these two sets are equal and also coincide with the Zariski closed set $V(\text{Ann}_R M)$. For a general M , $\text{Supp}_R M$ is always *specialization closed*: if it contains any point \mathfrak{p} then it must contain its closure $V(\mathfrak{p}) = \{\mathfrak{q} \mid \mathfrak{p} \subseteq \mathfrak{q}\}$. In fact $\text{Supp}_R M$ is equal to the specialization closure of $\text{supp}_R M$: $\text{Supp}_R M = \bigcup_{\mathfrak{p} \in \text{supp}_R M} V(\mathfrak{p})$. The small support plays a fundamental role in the Benson–Iyengar–Krause stratification theory, but in this note it will only appear implicitly.

The next lemma follows by a standard induction on the length of the objects.

Lemma 2.1. *If $\mathcal{K} = \text{Thick}(\mathbf{1})$ is affine and R is noetherian, the graded R -module $\text{Hom}_{\mathcal{K}}^*(X, Y)$ is finitely generated for all $X, Y \in \mathcal{K}$. \square*

The comparison map of spectra. Recall from [Balmer 2005] that, as a set, the spectrum $\text{Spc } \mathcal{K}$ is defined to be the collection of all proper thick subcategories $\mathcal{P} \subsetneq \mathcal{K}$ which are *prime tensor ideals*: $X \otimes Y \in \mathcal{P} \iff X \in \mathcal{P} \text{ or } Y \in \mathcal{P}$. For every $\mathcal{P} \in \text{Spc } \mathcal{K}$, let $\rho_{\mathcal{K}}(\mathcal{P})$ denote the ideal of $\text{Spec } R$ generated by the set of homogeneous elements $\{f : \mathbf{1} \rightarrow \Sigma^{|f|}\mathbf{1} \mid \text{cone}(f) \notin \mathcal{P}\}$. By [Balmer 2010a, Theorem 5.3], the assignment $\mathcal{P} \mapsto \rho_{\mathcal{K}}(\mathcal{P})$ defines a continuous map $\rho_{\mathcal{K}} : \text{Spc } \mathcal{K} \rightarrow \text{Spec } R$, natural in \mathcal{K} . Moreover, the two spaces $\text{Spc } \mathcal{K}$ and $\text{Spec } R$ are *spectral* in the sense of Hochster [1969], and $\rho_{\mathcal{K}}$ is a *spectral map* in that the preimage of a compact open set is again compact.

Lemma 2.2. *If $\rho_{\mathcal{K}}$ is bijective then it is a homeomorphism.*

Proof. This is an immediate consequence of [Hochster 1969, Proposition 15], which says that for a spectral map of spectral topological spaces to be a homeomorphism it suffices that it is an order isomorphism for the specialization order of the two spaces. Recall that the specialization order is defined for the points of any topological space by $x \geq y \iff x \in \overline{\{y\}}$. Indeed $\rho := \rho_{\mathcal{K}}$ is inclusion reversing, $\mathcal{Q} \subseteq \mathcal{P} \iff \rho(\mathcal{Q}) \supseteq \rho(\mathcal{P})$, hence it maps the closure $\overline{\{\mathcal{P}\}} = \{\mathcal{Q} \mid \mathcal{Q} \subseteq \mathcal{P}\}$ in $\mathrm{Spc} \mathcal{K}$ of any point \mathcal{P} to the Zariski closure $V(\rho(\mathcal{P})) = \{\mathfrak{q} \mid \mathfrak{q} \supseteq \rho(\mathcal{P})\}$ in $\mathrm{Spec} R$ of the corresponding point. \square

Central localization. For every prime ideal \mathfrak{p} of the graded central ring R of \mathcal{K} , there exists by [Balmer 2010a, Theorem 3.6] a tensor triangulated category $\mathcal{K}_{\mathfrak{p}}$ having the same objects as \mathcal{K} and such that its graded Hom modules are the localizations

$$\mathrm{Hom}_{\mathcal{K}_{\mathfrak{p}}}^*(X, Y) = \mathrm{Hom}_{\mathcal{K}}^*(X, Y)_{\mathfrak{p}}.$$

In particular the graded central ring of $\mathcal{K}_{\mathfrak{p}}$ is the local ring $R_{\mathfrak{p}}$. There is a canonical exact functor $q_{\mathfrak{p}} : \mathcal{K} \rightarrow \mathcal{K}_{\mathfrak{p}}$, which is in fact the Verdier quotient by the thick tensor ideal generated by $\{\mathrm{cone}(f) \in \mathcal{K} \mid f \in R \setminus \mathfrak{p} \text{ homogeneous}\}$. For emphasis, we will sometimes write $X_{\mathfrak{p}}$ for $X = q_{\mathfrak{p}}X$ when considered as an object of $\mathcal{K}_{\mathfrak{p}}$.

Clearly if \mathcal{K} is generated by $\mathbf{1}$ then $\mathcal{K}_{\mathfrak{p}}$ is generated by $\mathbf{1}_{\mathfrak{p}}$. Later we will use the fact that if a tensor triangulated category is generated by its unit then every thick subcategory is automatically a tensor ideal.

Let $\ell_{\mathfrak{p}} : R \rightarrow R_{\mathfrak{p}}$ denote the localization map between the graded central rings of the two categories. By [Balmer 2010a, Theorem 5.4], we have a pullback square of spaces

$$(2.3) \quad \begin{array}{ccc} \mathrm{Spc}(\mathcal{K}_{\mathfrak{p}}) & \xhookrightarrow{\mathrm{Spc}(q_{\mathfrak{p}})} & \mathrm{Spc}(\mathcal{K}) \\ \rho_{\mathcal{K}_{\mathfrak{p}}} \downarrow & & \downarrow \rho_{\mathcal{K}} \\ \mathrm{Spec}(R_{\mathfrak{p}}) & \xhookrightarrow{\mathrm{Spec}(\ell_{\mathfrak{p}})} & \mathrm{Spec}(R) \end{array}$$

where the horizontal maps are injective.

Koszul objects. We adapt some convenient notation from [Benson et al. 2008]. For any object $X \in \mathcal{K}$ and homogeneous element $f \in R$, let $X // f := \mathrm{cone}(f \cdot X)$ be any choice of mapping cone for the map $f \cdot X : \Sigma^{-|f|} X \rightarrow X$ given by the R -action. If f_1, \dots, f_n is a finite sequence of homogeneous elements, define recursively $X_0 := X$ and $X_i := X // (f_1, \dots, f_i) := (X // (f_1, \dots, f_{i-1})) // f_i$ for $i \in \{1, \dots, n\}$. Thus by construction we have exact triangles

$$(2.4) \quad \Sigma^{-|f_i|} X_{i-1} \xrightarrow{f_i \cdot X_{i-1}} X_{i-1} \longrightarrow X_i \longrightarrow \Sigma^{-|f_i|+1} X_{i-1},$$

and moreover, since the tensor product is exact, we have isomorphisms

$$X // (f_1, \dots, f_i) \simeq X \otimes \mathbf{1} // f_1 \otimes \cdots \otimes \mathbf{1} // f_i$$

for all $i \in \{1, \dots, n\}$. In the following, we will perform this construction *inside the \mathfrak{p} -local category $\mathcal{K}_{\mathfrak{p}}$* .

We need the following triangular version of the Nakayama lemma, for \mathcal{K} affine.

Lemma 2.5. *If $X \in \mathcal{K}_{\mathfrak{p}}$ is any object and f_1, \dots, f_n is a set of homogeneous generators for $\mathfrak{p}R_{\mathfrak{p}}$, then in $\mathcal{K}_{\mathfrak{p}}$ we have $X = 0$ if and only if $X // (f_1, \dots, f_n) = 0$.*

Proof. Since \mathcal{K} and thus $\mathcal{K}_{\mathfrak{p}}$ are generated by their tensor unit, it suffices to show that $H^*X_{\mathfrak{p}} = 0$ if and only if $H^*(X // (f_1, \dots, f_n))_{\mathfrak{p}} = 0$, and the latter can be proved as in [Benson et al. 2008, Lemma 5.11 (3)]. We give the easy argument for completeness.

With the above notation, by taking cohomology $H^* = \text{Hom}_{\mathcal{K}_{\mathfrak{p}}}^*(\mathbf{1}_{\mathfrak{p}}, -)$ of the triangle (2.4) of $\mathcal{K}_{\mathfrak{p}}$ we obtain the long exact sequence of $R_{\mathfrak{p}}$ -modules

$$\dots \longrightarrow H^{*-|f_i|}X_{i-1} \xrightarrow{f_i} H^*X_{i-1} \longrightarrow H^*X_i \longrightarrow H^{*-|f_i|+1}X_{i-1} \longrightarrow \dots,$$

where each module is finitely generated by Lemma 2.1. Since $f_i \in \mathfrak{p}$, if $H^*X_{i-1} \neq 0$ the first map in the sequence is not invertible by the Nakayama lemma, hence $H^*X_i \neq 0$. The evident recursion shows that $H^*X \neq 0$ implies $H^*X_n \neq 0$. The very same exact sequences also show that if $H^*X = 0$ then $H^*X_n = 0$. \square

3. Thick subcategories

Assume from now on that \mathcal{K} satisfies conditions (a) and (b) of Theorem 1.1.

Residue field objects. By hypothesis, for every prime ideal $\mathfrak{p} \in \text{Spec } R$ there exists a regular sequence f_1, \dots, f_n of homogeneous non-zero-divisors of $R_{\mathfrak{p}}$ which generate the ideal $\mathfrak{p}R_{\mathfrak{p}}$. Choose one such sequence once and for all, and construct the associated Koszul object

$$K(\mathfrak{p}) := \mathbf{1}_{\mathfrak{p}} // (f_1, \dots, f_n) \simeq \mathbf{1}_{\mathfrak{p}} // f_1 \otimes \cdots \otimes \mathbf{1}_{\mathfrak{p}} // f_n$$

in the \mathfrak{p} -local tensor triangulated category $\mathcal{K}_{\mathfrak{p}}$.

Lemma 3.1. *For every object $X \in \mathcal{K}_{\mathfrak{p}}$ and every $i \in \{1, \dots, n\}$, each element f of the ideal $(f_1, \dots, f_i) \subset R_{\mathfrak{p}}$ acts as zero on $X // (f_1, \dots, f_i)$, i.e., $f \cdot X // (f_1, \dots, f_i) = 0$.*

Proof. Recall that, as an immediate consequence of the $R_{\mathfrak{p}}$ -bilinearity of the composition in $\mathcal{K}_{\mathfrak{p}}$, the elements of $R_{\mathfrak{p}}$ acting as zero on an object Y form an ideal (coinciding with the annihilator of the $R_{\mathfrak{p}}$ -module $\text{Hom}_{\mathcal{K}_{\mathfrak{p}}}^*(Y, Y)$). Thanks to the isomorphism $X // (f_1, \dots, f_i) \simeq X \otimes \mathbf{1}_{\mathfrak{p}} // f_1 \otimes \cdots \otimes \mathbf{1}_{\mathfrak{p}} // f_i$ and the $R_{\mathfrak{p}}$ -linearity of

the tensor product, it will therefore suffice to prove that f_i acts as zero on $\mathbf{1}_{\mathfrak{p}}//f_i$. Consider the commutative diagram

$$\begin{array}{ccccc}
 \Sigma^{-|f_i|}\mathbf{1}_{\mathfrak{p}} & \xrightarrow{f_i} & \mathbf{1}_{\mathfrak{p}} & \xrightarrow{g} & \mathbf{1}_{\mathfrak{p}}//f_i & \longrightarrow & \Sigma^{-|f_i|+1}\mathbf{1}_{\mathfrak{p}} \\
 & & \downarrow f_i & \searrow 0 & \downarrow f_i & \swarrow h & \\
 & & \Sigma^{|f_i|}\mathbf{1}_{\mathfrak{p}} & \xrightarrow{g} & \Sigma^{|f_i|}\mathbf{1}_{\mathfrak{p}}//f_i & &
 \end{array}$$

where the top row is the exact triangle defining $\mathbf{1}_{\mathfrak{p}}//f_i$. Being the composite of two consecutive maps in a triangle, gf_i is zero. Up to a suspension, this is also the diagonal map in the square. Hence $f_i \cdot \mathbf{1}_{\mathfrak{p}}//f_i$ factors through a map h as pictured. Since R is even by hypothesis, we have that $R_{\mathfrak{p}}$ is even, and we claim that also

$$(3.2) \quad H^n(\mathbf{1}_{\mathfrak{p}}//f_i) = 0 \quad \text{for all odd } n.$$

This implies $h = 0$ and therefore $f_i \cdot \mathbf{1}_{\mathfrak{p}}//f_i = 0$, as required. To prove the claim, note that the defining triangle of $\mathbf{1}_{\mathfrak{p}}//f_i$ induces the exact sequence

$$R_{\mathfrak{p}}^{n-|f_i|} \xrightarrow{f_i} R_{\mathfrak{p}}^n \longrightarrow H^n(\mathbf{1}_{\mathfrak{p}}//f_i) \longrightarrow R_{\mathfrak{p}}^{n-|f_i|+1} \xrightarrow{f_i} R_{\mathfrak{p}}^{n+1},$$

where the first and last maps are injective by the hypothesis that f_i is a non-zero-divisor in $R_{\mathfrak{p}}$. Thus (3.2), and even $H^*(\mathbf{1}_{\mathfrak{p}}//f_i) \simeq R_{\mathfrak{p}}/(f_i)$, follows immediately. \square

Corollary 3.3. $H^*(X \otimes K(\mathfrak{p}))$ is a graded $k(\mathfrak{p})$ -vector space for every $X \in \mathcal{K}_{\mathfrak{p}}$.

Proof. By Lemma 3.1 together with the R -linearity of the tensor product, each $f \in \mathfrak{p}R_{\mathfrak{p}}$ acts as zero on $X \otimes K(\mathfrak{p}) \simeq X \otimes \mathbf{1}_{\mathfrak{p}}//(f_1, \dots, f_n)$. Therefore all such f also act as zero on $H^*(X \otimes K(\mathfrak{p}))$ by the R -linearity of composition. \square

Lemma 3.4. *There is an isomorphism $H^*(\mathbf{1}_{\mathfrak{p}}//(f_1, \dots, f_i)) \simeq R_{\mathfrak{p}}/(f_1, \dots, f_i)$ of R -modules for all $i \in \{1, \dots, n\}$. In particular $H^*K(\mathfrak{p})$ is isomorphic to the residue field $k(\mathfrak{p}) := R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}$.*

Proof. Write $C_0 = \mathbf{1}_{\mathfrak{p}}$ and $C_i := \mathbf{1}_{\mathfrak{p}}//(f_1, \dots, f_i)$ for short. Then $K(\mathfrak{p}) = C_n$, and for all $i \in \{1, \dots, n\}$ we have exact triangles

$$\Sigma^{-|f_i|}C_{i-1} \xrightarrow{f_i \cdot C_{i-1}} C_{i-1} \longrightarrow C_i \longrightarrow \Sigma^{-|f_i|+1}C_{i-1}.$$

The claim follows by recursion on i . Indeed $H^*C_0 = R_{\mathfrak{p}}$, and assume that $H^*C_{i-1} \simeq R_{\mathfrak{p}}/(f_1, \dots, f_{i-1})$. Then the above triangle induces an exact sequence

$$H^{*-|f_i|}C_{i-1} \xrightarrow{f_i} H^*C_{i-1} \longrightarrow H^*C_i \longrightarrow H^{*-|f_i|+1}C_{i-1} \xrightarrow{f_i} H^{*+1}C_{i-1},$$

where the first and last maps are injective because by hypothesis f_i is a non-zero-divisor in the ring $R_{\mathfrak{p}}/(f_1, \dots, f_{i-1})$. We thus obtain a short exact sequence $0 \rightarrow f_i R_{\mathfrak{p}}/(f_1, \dots, f_{i-1}) \rightarrow R_{\mathfrak{p}}/(f_1, \dots, f_{i-1}) \rightarrow H^*C_i \rightarrow 0$, proving the claim for i . \square

Remark 3.5. Of the weak regularity hypothesis (b), the proof of Lemma 3.4 only uses that f_1, \dots, f_n is a regular sequence, while the proof of Lemma 3.1 only uses that the f_i are non-zero-divisors in $R_{\mathfrak{p}}$ and that the ring R is even. These are the only places where we make use of these assumptions (the noetherian hypothesis, on the other hand, will be needed on several occasions). Note that, although we already know by Corollary 3.3 that $H^*K(\mathfrak{p})$ is a $k(\mathfrak{p})$ -vector space, for the next proposition we also need it to be one-dimensional as per Lemma 3.4.

Proposition 3.6. *For all $\mathfrak{p} \in \text{Spec } R$ and $X \in \mathcal{K}_{\mathfrak{p}}$, the tensor product $X \otimes K(\mathfrak{p})$ decomposes into a coproduct of shifted copies of the residue field object:*

$$\coprod_{\alpha} \Sigma^{n_{\alpha}} K(\mathfrak{p}) \xrightarrow{\sim} X \otimes K(\mathfrak{p}).$$

Proof. By Corollary 3.3 we know that $H^*(X \otimes K(\mathfrak{p}))$ is a graded $k(\mathfrak{p})$ -vector space. Choose a graded basis $\{x_{\alpha}\}_{\alpha}$, corresponding to a morphism $\coprod_{\alpha} \Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} \rightarrow X \otimes K(\mathfrak{p})$. We will show that this map extends nontrivially to the Koszul object

$$\left(\coprod_{\alpha} \Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} \right) // (f_1, \dots, f_n) = \coprod_{\alpha} (\Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} // (f_1, \dots, f_n)).$$

For this, it will suffice to extend each individual map $x_{\alpha} : \Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} \rightarrow X \otimes K(\mathfrak{p})$. As before, we proceed recursively along the regular sequence f_1, \dots, f_n . Consider the commutative diagram

$$\begin{array}{ccccccc} \Sigma^{n_{\alpha}-|f_1|} \mathbf{1}_{\mathfrak{p}} & \xrightarrow{f_1} & \Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} & \longrightarrow & \Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} // f_1 & \longrightarrow & \\ \Sigma^{-|f_1|} x_{\alpha} \downarrow & \searrow 0 & \downarrow x_{\alpha} & \swarrow x_{\alpha}^1 & & & \\ \Sigma^{-|f_1|} X \otimes K(\mathfrak{p}) & \xrightarrow{f_1=0} & X \otimes K(\mathfrak{p}) & & & & \end{array}$$

where the top row is a rotation of the defining triangle for $\mathbf{1}_{\mathfrak{p}} // f_1$. The left-bottom composite vanishes because f_1 acts trivially on $X \otimes K(\mathfrak{p})$ by Lemma 3.1. Hence we obtain the map x_{α}^1 on the right. Note that $x_{\alpha}^1 \neq 0$ because $x_{\alpha} \neq 0$. Now we repeat the procedure for $i = 2, \dots, n$, using the triangle

$$\Sigma^{-|f_i|} \mathbf{1}_{\mathfrak{p}} // (f_1, \dots, f_{i-1}) \xrightarrow{f_i} \mathbf{1}_{\mathfrak{p}} // (f_1, \dots, f_{i-1}) \longrightarrow \mathbf{1}_{\mathfrak{p}} // (f_1, \dots, f_i) \longrightarrow$$

to extend x_{α}^{i-1} to a nonzero map $x_{\alpha}^i : \Sigma^{n_{\alpha}} \mathbf{1}_{\mathfrak{p}} // (f_1, \dots, f_i) \rightarrow X \otimes K(\mathfrak{p})$ hitting the same element in cohomology. In particular we obtain the announced extension $x_{\alpha}^n : \Sigma^{n_{\alpha}} K(\mathfrak{p}) \rightarrow X \otimes K(\mathfrak{p})$. As a nonzero map on a one-dimensional $k(\mathfrak{p})$ -vector space (Lemma 3.4), the induced map $H^*(x_{\alpha}^n)$ must be injective. Hence, collectively, the maps $\{x_{\alpha}^n\}_{\alpha}$ yield an isomorphism as required. \square

Proposition 3.7. *For every \mathfrak{p} , the thick subcategory $\text{Thick}(K(\mathfrak{p}))$ of $\mathcal{K}_{\mathfrak{p}}$ is minimal, meaning that it contains no proper nonzero thick subcategories.*

Proof. Note that for every nonzero object X of $\mathcal{K}_{\mathfrak{p}}$ we have $X \otimes K(\mathfrak{p}) \neq 0$. Indeed if $X \otimes K(\mathfrak{p}) = X // (f_1, \dots, f_n) = 0$ then $X_{\mathfrak{p}} = 0$ by [Lemma 2.5](#).

Let \mathcal{C} be a thick subcategory of $\text{Thick}(K(\mathfrak{p}))$. Because \mathcal{C} is a tensor ideal, if it contains a nonzero object X then it also contains $X \otimes K(\mathfrak{p})$, which is again nonzero by the above observation. Therefore \mathcal{C} must contain a shifted copy of $K(\mathfrak{p})$ by [Proposition 3.6](#), hence $\mathcal{C} = \text{Thick}(K(\mathfrak{p}))$. \square

Proof of Theorem 1.1. Now we show how to deduce our main result from the minimality of the thick subcategories $\text{Thick}(K(\mathfrak{p}))$. By [Lemma 2.2](#) it will suffice to show that the map $\rho_{\mathcal{K}} : \text{Spc } \mathcal{K} \rightarrow \text{Spec } R$ is bijective. Since R is graded noetherian, $\rho_{\mathcal{K}}$ is surjective by [[Balmer 2010a](#), Theorem 7.3]. It remains to prove it is injective.

Let $\mathfrak{p} \in \text{Spec } R$ be any homogeneous prime. We must show that the fiber of the comparison map $\rho_{\mathcal{K}} : \text{Spc } \mathcal{K} \rightarrow \text{Spec } R$ over \mathfrak{p} consists of a single prime tensor ideal. By the pullback square (2.3), every point of $\text{Spc } \mathcal{K}$ lying over \mathfrak{p} must belong to $\text{Spc } \mathcal{K}_{\mathfrak{p}}$. Hence it will suffice to show that the fiber of $\rho := \rho_{\mathcal{K}_{\mathfrak{p}}}$ over the maximal ideal $\mathfrak{m} := \mathfrak{p}R_{\mathfrak{p}}$ of $R_{\mathfrak{p}}$ consists of a single point. In fact a stronger statement is true: if $\mathcal{P} \in \text{Spc } \mathcal{K}_{\mathfrak{p}}$ is such that $\rho(\mathcal{P}) = \mathfrak{m}$, then $\mathcal{P} = \{0\}$. Let us prove this.

By definition of the comparison map we have

$$\rho(\mathcal{P}) = \langle \{f \in R_{\mathfrak{p}} \mid f \text{ is homogeneous and } \mathbf{1}_{\mathfrak{p}} // f \notin \mathcal{P}\} \rangle,$$

and as $\rho(\mathcal{P}) \subseteq \mathfrak{m}$ always holds by the maximality of \mathfrak{m} , the hypothesis $\rho(\mathcal{P}) = \mathfrak{m}$ precisely means that $\mathbf{1}_{\mathfrak{p}} // f \notin \mathcal{P}$ for all homogeneous elements $f \in \mathfrak{m}$. In particular $\mathbf{1}_{\mathfrak{p}} // f_i \notin \mathcal{P}$ for the elements f_i in the chosen regular sequence for \mathfrak{m} . As \mathcal{P} is a tensor prime, we deduce further that

$$(3.8) \quad K(\mathfrak{p}) \simeq \mathbf{1}_{\mathfrak{p}} // f_1 \otimes \cdots \otimes \mathbf{1}_{\mathfrak{p}} // f_n \notin \mathcal{P}.$$

Now let $X \in \mathcal{P}$ and assume that $X \neq 0$. Then $X \otimes K(\mathfrak{p}) \neq 0$ by [Lemma 2.5](#), hence

$$(3.9) \quad \text{Thick}(X \otimes K(\mathfrak{p})) = \text{Thick}(K(\mathfrak{p}))$$

by the minimality of $\text{Thick}(K(\mathfrak{p}))$, [Proposition 3.7](#). As \mathcal{P} is a thick tensor ideal we also have $X \otimes K(\mathfrak{p}) \in \mathcal{P}$ and therefore $K(\mathfrak{p}) \in \mathcal{P}$ by (3.9), but this contradicts (3.8). Therefore $X = 0$ and we conclude that $\mathcal{P} = \{0\}$, proving the claim. This concludes the proof of [Theorem 1.1](#).

Proof of Corollary 1.2. To deduce [Corollary 1.2](#) from the theorem, we must verify that the homeomorphism $\rho_{\mathcal{K}}$ identifies $\text{Supp}_R H^*X \subseteq \text{Spec } R$, the ring-theoretic support of an object $X \in \mathcal{K}$, with $\text{supp } X := \{\mathcal{P} \in \text{Spc } \mathcal{K} \mid X \notin \mathcal{P}\}$, the universal support datum of X :

Lemma 3.10. *We have $\text{Supp}_R H^*X = \rho_{\mathcal{K}}(\text{supp } X)$ for all $X \in \mathcal{K}$.*

Proof. Let $\mathfrak{p} = \rho_{\mathcal{K}}(\mathcal{P})$. It follows from (2.3) that $X \in \mathcal{P}$ if and only if $X_{\mathfrak{p}} \in \mathcal{P}_{\mathfrak{p}}$, where $\mathcal{P}_{\mathfrak{p}}$ denotes \mathcal{P} seen as an element of $\mathrm{Spc} \mathcal{K}_{\mathfrak{p}}$. We have just proved that $\rho_{\mathcal{K}_{\mathfrak{p}}} : \mathrm{Spc} \mathcal{K}_{\mathfrak{p}} \xrightarrow{\sim} R_{\mathfrak{p}}$ is a bijection sending $\{0\}$ to $\mathfrak{p}R_{\mathfrak{p}}$, so we must have $\mathcal{P}_{\mathfrak{p}} = \{0\}$. Therefore

$$\mathfrak{p} \in \mathrm{Supp}_R H^*X \iff H^*X_{\mathfrak{p}} \neq 0 \iff X_{\mathfrak{p}} \neq 0 \iff X_{\mathfrak{p}} \notin \mathcal{P}_{\mathfrak{p}} \iff \mathcal{P} \in \mathrm{supp} X. \quad \square$$

Now it suffices to appeal to the abstract classification theorem [Balmer 2005, Theorem 4.10]. Indeed, since R is noetherian, the space $\mathrm{Spec} R$ is noetherian and therefore its specialization closed subsets and its Thomason subsets coincide (cf. [Balmer 2005, Remark 4.11]). Moreover, since \mathcal{K} is generated by its tensor unit, all its objects are dualizable (because dualizable objects form a thick subcategory and $\mathbf{1}$ is dualizable) and therefore all its thick tensor ideals are radical (see [Balmer 2007, Proposition 2.4]). Hence by Theorem 1.1 and Lemma 3.10 the classification of [Balmer 2005, Theorem 4.10] immediately translates into the classification described in Corollary 1.2, as wished.

4. Localizing subcategories

Assume from now on that \mathcal{T} is a compactly generated tensor triangulated category such that its subcategory $\mathcal{K} := \mathcal{T}^c$ of compact objects satisfies hypotheses (a) and (b) of Theorem 1.1. Thus in particular \mathcal{T} is generated as a localizing subcategory by the tensor unit: $\mathrm{Loc}(\mathbf{1}) = \mathcal{T}$. It follows that every localizing subcategory of \mathcal{T} is automatically a tensor ideal.

Since \mathcal{T} is compactly generated, the (Verdier) \mathfrak{p} -localization functor $q_{\mathfrak{p}} : \mathcal{K} \rightarrow \mathcal{K}_{\mathfrak{p}}$ we used so far can be extended to a finite (Bousfield) localization functor

$$(-)_{\mathfrak{p}} : \mathcal{T} \rightarrow \mathcal{T}.$$

We briefly recall its properties, referring for all proofs to [Benson et al. 2011, §2] or [Dell'Ambrogio 2010, §2]. Let

$$\mathcal{L} = \mathrm{Loc}(\{\mathrm{cone}(f) \mid f \in R \setminus \mathfrak{p} \text{ homogeneous}\}).$$

Then the Verdier quotient $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{L} =: \mathcal{T}_{\mathfrak{p}}$ has a fully faithful right adjoint, $I : \mathcal{T}_{\mathfrak{p}} \hookrightarrow \mathcal{T}$, and the functor $(-)_{\mathfrak{p}}$ can be defined to be the composite $(-)_{\mathfrak{p}} := I \circ Q$. As \mathcal{L} is generated by a tensor ideal of dualizable objects, we have $X_{\mathfrak{p}} \cong X \otimes \mathbf{1}_{\mathfrak{p}}$ for all $X \in \mathcal{T}$. Moreover, the unit $X \rightarrow X_{\mathfrak{p}}$ of the (Q, I) -adjunction induces a natural map $\mathrm{Hom}_{\mathcal{T}}^*(Y, X)_{\mathfrak{p}} \rightarrow \mathrm{Hom}_{\mathcal{T}}^*(Y, X_{\mathfrak{p}})$ which is an isomorphism whenever $Y \in \mathcal{K}$ (see [Benson et al. 2011, Proposition 2.3] or [Dell'Ambrogio 2010, Theorem 2.33 (h)]). In particular we have the identification

$$(H^*X)_{\mathfrak{p}} \xrightarrow{\sim} H^*(X_{\mathfrak{p}})$$

for all $X \in \mathcal{T}$. It follows also that the restriction of Q to compact objects $X, Y \in \mathcal{K}$ agrees with q_p , so that we may identify \mathcal{K}_p with the full subcategory $I(\mathcal{K}_p)$ of \mathcal{T} (and thereby eliminate the slight ambiguity of the notation “ X_p ”).

Recall the residue field objects $K(\mathfrak{p})$ defined in [Section 3](#):

$$K(\mathfrak{p}) := \mathbf{1}_p // (f_1, \dots, f_n) \simeq \mathbf{1}_p // f_1 \otimes \cdots \otimes \mathbf{1}_p // f_n \in \mathcal{T}$$

(as before, f_1, \dots, f_n denotes the chosen regular sequence of non-zero-divisors generating the prime \mathfrak{p}).

The main point of this section is that the crucial minimality result of [Proposition 3.7](#) can be extended to localizing subcategories of \mathcal{T} , as we verify next.

Lemma 4.1. *For every object $X \in \mathcal{T}$ and every $i \in \{1, \dots, n\}$, each element f of $(f_1, \dots, f_i) \subset R$ acts as zero on $X_p // (f_1, \dots, f_i)$, i.e., $f \cdot X_p // (f_1, \dots, f_i) = 0$. In particular, the R -module $H^*(X \otimes K(\mathfrak{p}))$ is a graded $k(\mathfrak{p})$ -vector space.*

Proof. Exactly the same as for [Lemma 3.1](#) and [Corollary 3.3](#). (Use that $X \otimes K(\mathfrak{p}) = X_p \otimes K(\mathfrak{p})$ to work inside the big \mathfrak{p} -local category \mathcal{T}_p .) \square

Proposition 4.2. *For all $\mathfrak{p} \in \text{Spec } R$ and $X \in \mathcal{T}$, the tensor product $X \otimes K(\mathfrak{p})$ decomposes into a coproduct of shifted copies of the residue field object:*

$$\coprod_{\alpha} \Sigma^{n_{\alpha}} K(\mathfrak{p}) \xrightarrow{\sim} X \otimes K(\mathfrak{p}).$$

Proof. Exactly the same as for [Proposition 3.6](#), using [Lemma 4.1](#). \square

Proposition 4.3. *For every \mathfrak{p} , the localizing subcategory $\text{Loc}(K(\mathfrak{p}))$ of \mathcal{T} is minimal, meaning that it contains no proper nonzero localizing subcategories.*

Proof. This follows from [Proposition 4.2](#) precisely as in the proof of [Proposition 3.7](#), except that we cannot use [Lemma 2.5](#) to show that $X \otimes K(\mathfrak{p}) \neq 0$ for every nonzero object $X \in \text{Loc}(K(\mathfrak{p}))$. Instead, we may use the following argument.

First note that $X \otimes K(\mathfrak{q}) = 0$ for all $\mathfrak{q} \in \text{Spec } R \setminus \{\mathfrak{p}\}$. Indeed, this property holds for $X = K(\mathfrak{p})$ by [Lemma 4.1](#) (because if $\mathfrak{p} \neq \mathfrak{q}$ then some homogeneous element of R must act on $K(\mathfrak{p}) \otimes K(\mathfrak{q})$ both as zero and invertibly) and is stable under taking coproducts and mapping cones (as the latter are preserved by $- \otimes K(\mathfrak{p})$); hence it must hold for all objects of $\text{Loc}(K(\mathfrak{p}))$, as wished. Now combine this with [Proposition 4.5](#) below. \square

Lemma 4.4. *Let M be any nonzero module, possibly infinitely generated, over a noetherian \mathbb{Z} -graded commutative ring S . Then there exists a minimal prime in $\text{Supp}_S M := \{\mathfrak{p} \in \text{Spec } S \mid M_{\mathfrak{p}} \neq 0\}$, the big Zariski support of M .*

Proof. If $M \neq 0$ then $M_{\mathfrak{p}} \neq 0$ for some prime \mathfrak{p} , so the support is not empty. Moreover, it suffices to prove the claim for the nonzero module $M_{\mathfrak{p}}$ over $S_{\mathfrak{p}}$, because a minimal prime of $\text{Supp}_{S_{\mathfrak{p}}} M_{\mathfrak{p}}$ yields a minimal prime in $\text{Supp}_S M$; hence we may

assume that S is local. By Zorn's lemma it suffices to show that in $\text{Supp}_S M$ every chain of primes admits a minimum. Indeed, each such chain must stabilize, because a local commutative noetherian ring has finite Krull dimension. In the ungraded case, the latter is a well-known corollary of Krull's principal ideal theorem. A proof of the analogous result for graded rings can be found in [Bruns and Herzog 1993, Theorem 1.5.8] or [Park and Park 2011, Theorem 3.5]. \square

Proposition 4.5. *If an object $X \in \mathcal{T}$ is such that $X \otimes K(\mathfrak{p}) = 0$ for all $\mathfrak{p} \in \text{Spec } R$ then $X = 0$.*

Proof. We prove the contrapositive. Assume that $X \neq 0$. Then $H^*X \neq 0$, hence for some $\mathfrak{p} \in \text{Spec } R$ we must have $H^*(X_{\mathfrak{p}}) = (H^*X)_{\mathfrak{p}} \neq 0$ and therefore $X_{\mathfrak{p}} \neq 0$. By Lemma 4.4, we may choose a prime \mathfrak{p} which is minimal among the primes with this property. Thus the big support of the R -module $H^*X_{\mathfrak{p}}$ consists precisely of the prime \mathfrak{p} . We are going to recursively show that $X_i := X_{\mathfrak{p}} // (f_1, \dots, f_i)$ satisfies $\text{Supp}_R H^*X_i = \{\mathfrak{p}\}$ for all $i \in \{1, \dots, n\}$. Thus in particular $X \otimes K(\mathfrak{p}) = X_n \neq 0$, which proves the proposition. We already know $\text{Supp}_R H^*X_0 = \{\mathfrak{p}\}$ for $X_0 := X_{\mathfrak{p}}$, and suppose we have shown that $\text{Supp}_R H^*X_{i-1} = \{\mathfrak{p}\}$. The exact triangle

$$\Sigma^{-|f_i|} X_{i-1} \xrightarrow{f_i} X_{i-1} \longrightarrow X_i \longrightarrow \Sigma^{-|f_i|+1} X_{i-1}$$

implies that $\text{Supp}_R H^*X_i \subseteq \{\mathfrak{p}\}$. Hence $X_i \neq 0$ is equivalent to $\text{Supp}_R H^*X_i = \{\mathfrak{p}\}$. By the triangle again, if $X_i = 0$ were the case f_i would act invertibly on X_{i-1} and thus on H^*X_{i-1} . This implies $H^*X_{i-1} = (H^*X_{i-1})[f_i^{-1}]$, and since $f_i \in \mathfrak{p}$ we would conclude that $\mathfrak{p} \notin \text{Supp}_R H^*X_{i-1}$, in contradiction with the induction hypothesis. Therefore $X_i \neq 0$, as claimed. \square

Proof of Theorem 1.3. The result now follows easily from the machinery developed by Benson, Iyengar and Krause [Benson et al. 2008; 2011]. Indeed, by [Benson et al. 2011, Theorem 4.2], to obtain the claimed classification of localizing subcategories it suffices to verify that *the action of R stratifies \mathcal{T}* . By definition, this means that the following two axioms are satisfied:

- *The local-global principle:* For every object $X \in \mathcal{T}$ we have the equality

$$\text{Loc}(X) = \text{Loc}(\{\Gamma_{\mathfrak{p}}X \mid \mathfrak{p} \in \text{Spec } R\})$$

of localizing subcategories of \mathcal{T} .

- *Minimality:* For every $\mathfrak{p} \in \text{Spec } R$ the localizing subcategory $\Gamma_{\mathfrak{p}}\mathcal{T}$ of \mathcal{T} is minimal or zero.

The functors $\Gamma_{\mathfrak{p}} : \mathcal{T} \rightarrow \mathcal{T}$ are introduced in [Benson et al. 2008], but we don't need to know how they are defined. In our context, i.e., where \mathcal{T} is a tensor category and the action of R is the canonical one of the central ring, the local-global principle always holds by [Benson et al. 2011, Theorem 7.2] (see also [Stevenson 2013,

Theorem 6.8]). Moreover $\Gamma_{\mathfrak{p}}X = X \otimes \Gamma_{\mathfrak{p}}\mathbf{1}$ for all $X \in \mathcal{T}$, which implies that $\Gamma_{\mathfrak{p}}\mathcal{T} = \text{Loc}(\Gamma_{\mathfrak{p}}\mathbf{1})$ since \mathcal{T} is generated by $\mathbf{1}$. Therefore the remaining minimality condition follows from [Proposition 4.3](#), because $\text{Loc}(K(\mathfrak{p})) = \text{Loc}(\Gamma_{\mathfrak{p}}\mathbf{1})$ by [\[Benson et al. 2011, Lemma 3.8 \(2\)\]](#) (indeed, by construction $K(\mathfrak{p})$ is a particular instance of the objects collectively denoted by $\mathbf{1}(\mathfrak{p})$ in [\[loc. cit.\]](#)). This establishes the first bijection in [Theorem 1.3](#).

The claimed identification of the Benson–Iyengar–Krause support, $\text{supp}_R X = \{\mathfrak{p} \in \text{Spec } R \mid X \otimes \Gamma_{\mathfrak{p}}\mathbf{1} \neq 0\}$, with the set $\{\mathfrak{p} \in \text{Spec } R \mid X \otimes K(\mathfrak{p}) \neq 0\}$ is an easy consequence of the equality $\text{Loc}(K(\mathfrak{p})) = \text{Loc}(\Gamma_{\mathfrak{p}}\mathbf{1})$ mentioned above.

It remains to verify the moreover part of [Theorem 1.3](#). Let us begin by noting that, if $X \in \mathcal{K}$ is a compact object, we have

$$(4.6) \quad \text{supp}_R X = \text{supp}_R H^*X = \text{Supp}_R H^*X$$

by [\[Benson et al. 2008, Theorem 5.5 \(1\)\]](#) and [Lemma 2.1](#). Now let $\mathcal{L} \subseteq \mathcal{T}$ be such that $\mathcal{L} = \text{Loc}(\mathcal{L} \cap \mathcal{K})$. Then

$$\bigcup_{X \in \mathcal{L}} \text{supp}_R X = \bigcup_{X \in \mathcal{L} \cap \mathcal{K}} \text{supp}_R X = \bigcup_{X \in \mathcal{L} \cap \mathcal{K}} \text{Supp}_R H^*X$$

by (4.6), and the latter is a specialization closed subset of the spectrum. Conversely, if $S \subseteq \text{Spec } R$ is specialization closed the corresponding localizing subcategory $\{X \in \mathcal{T} \mid \text{supp}_R X \subseteq S\}$ is generated by compact objects by [\[Benson et al. 2008, Theorem 6.4\]](#), hence $\mathcal{L} = \text{Loc}(\mathcal{L} \cap \mathcal{K})$. This concludes the proof of the theorem.

It is well known that the assignments $\mathcal{C} \mapsto \text{Loc}(\mathcal{C})$ and $\mathcal{L} \mapsto \mathcal{L} \cap \mathcal{K}$ are mutually inverse bijections between thick subcategories $\mathcal{C} \subseteq \mathcal{K}$ and localizing subcategories $\mathcal{L} \subseteq \mathcal{T}$ which are generated by compact objects of \mathcal{T} (see [\[Neeman 1992b\]](#)). Together with (4.6), this shows how to deduce the classification of thick subcategories of [Corollary 1.2](#) from [Theorem 1.3](#).

Finally, there are several ways to derive the telescope conjecture of [Corollary 1.4](#) from the previous results. For instance, we may proceed as in [\[Benson et al. 2011, §6.2\]](#).

Remark 4.7. Using the theory of coherent functors, Benson, Iyengar and Krause have recently developed in [\[Benson et al. 2015\]](#) an analogue of their stratification theory of compactly generated categories that can be applied to general essentially small triangulated categories. Their theory, and more specifically [\[Benson et al. 2015, Theorem 7.4\]](#), provides an alternative way to derive [Theorem 1.1](#) from [Proposition 3.7](#).

The case of commutative dg algebras. We still owe readers a proof of [Theorem 1.6](#). Let A be a commutative dg algebra and let $D(A)$ be the derived category of (left,

say dg- A -modules. The following elementary fact was pointed out to us by the referee.

Lemma 4.8. *Every $f \in H^*A$ acts as zero on its own mapping cone $C(f)$.*

Proof. A (homogeneous) element $f \in H^*A$ of degree $|f| = -n$ is (represented by) a morphism $f : \Sigma^n A \rightarrow A$ of left dg- A -modules. Let us write sa ($a \in A$) for a generic element of degree $|a| - 1$ in the suspension $sA := \Sigma A$; here we use the Koszul sign convention and treat s as a symbol of degree -1 . The cone $C(f)$ has elements $(a, s^{n+1}b)$ (for $a, b \in A$). Then f acts on $C(f)$ by a morphism $s^n C(f) \rightarrow C(f)$ which, under the isomorphism $s^n C(f) \cong C(s^n f)$, is written as follows:

$$g : C(s^n f) \rightarrow C(f), \quad g(s^n a, s^{2n+1}b) = (f(s^n a), s^{n+1}f(s^n b))$$

(recall that the suspension $sh : sB \rightarrow sC$ of a morphism $h : B \rightarrow C$ is given by $(sh)(sb) = s(h(b))$). With these notations, the map $H : C(s^n f) \rightarrow C(f)$ defined by $H(s^n a, s^{2n+1}b) := (0, s^{n+1}a)$ is easily seen to satisfy $H(tx) = (-1)^{|t|}tH(x)$ (for $t \in A, x \in C(s^n f)$) and $dH + Hd = -g$; in other words, H is a homotopy $g \sim 0$ defined over A . □

As noted in [Remark 3.5](#), [Lemma 3.1](#) was the only place in all of our arguments where we made use of the hypothesis that R is concentrated in even degrees and that in the regular sequences we may choose the elements to be non-zero-divisors. But if we consider the example $\mathcal{K} := D(A)^c$, $\mathcal{T} := D(A)$ and $R := H^*A$, we see immediately that the conclusion of the lemma also follows from the above result ([Lemma 4.8](#)). Hence in this case we can get rid of the extra hypotheses, while the rest of our arguments go through unchanged. This proves [Theorem 1.6](#).

Indeed, in general in condition (b) of [Theorem 1.1](#) we could similarly renounce the evenness of R if we substitute the requirement that all elements f_i of the regular sequences be non-zero-divisors with the requirement that $f_i \cdot \mathbf{1} // f_i = 0$.

Acknowledgements

We are very grateful to Paul Balmer and an anonymous referee for their useful comments, and to the referee for suggesting [Theorem 1.6](#). We also thank Joseph Chuang for the reference [[Mathew 2016](#)].

References

[Balmer 2005] P. Balmer, “The spectrum of prime ideals in tensor triangulated categories”, *J. Reine Angew. Math.* **588** (2005), 149–168. MR 2196732 Zbl 1080.18007

[Balmer 2007] P. Balmer, “Supports and filtrations in algebraic geometry and modular representation theory”, *Amer. J. Math.* **129**:5 (2007), 1227–1250. MR 2354319 Zbl 1130.18005

- [Balmer 2010a] P. Balmer, “Spectra, spectra, spectra—tensor triangular spectra versus Zariski spectra of endomorphism rings”, *Algebr. Geom. Topol.* **10**:3 (2010), 1521–1563. MR 2661535 Zbl 1204.18005
- [Balmer 2010b] P. Balmer, “Tensor triangular geometry”, pp. 85–112 in *Proceedings of the International Congress of Mathematicians*, vol. II, edited by R. Bhatia et al., Hindustan Book Agency, New Delhi, 2010. MR 2827786 Zbl 1204.18004
- [Benson et al. 2008] D. Benson, S. B. Iyengar, and H. Krause, “Local cohomology and support for triangulated categories”, *Ann. Sci. Éc. Norm. Supér.* (4) **41**:4 (2008), 573–619. MR 2489634 Zbl 1171.18007
- [Benson et al. 2011] D. Benson, S. B. Iyengar, and H. Krause, “Stratifying triangulated categories”, *J. Topol.* **4**:3 (2011), 641–666. MR 2832572 Zbl 1239.18013
- [Benson et al. 2015] D. Benson, S. B. Iyengar, and H. Krause, “A local-global principle for small triangulated categories”, *Math. Proc. Cambridge Philos. Soc.* **158**:3 (2015), 451–476. MR 3335421
- [Bruns and Herzog 1993] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, Cambridge, 1993. MR 1251956 Zbl 0788.13005
- [Dell’Ambrogio 2010] I. Dell’Ambrogio, “Tensor triangular geometry and KK -theory”, *J. Homotopy Relat. Struct.* **5**:1 (2010), 319–358. MR 2812924 Zbl 1278.19010
- [Dell’Ambrogio and Tabuada 2012] I. Dell’Ambrogio and G. Tabuada, “Tensor triangular geometry of non-commutative motives”, *Adv. Math.* **229**:2 (2012), 1329–1357. MR 2855096 Zbl 1250.18012
- [Goto and Yamagishi 1983] S. Goto and K. Yamagishi, “Finite generation of Noetherian graded rings”, *Proc. Amer. Math. Soc.* **89**:1 (1983), 41–44. MR 706507 Zbl 0528.13015
- [Hochster 1969] M. Hochster, “Prime ideal structure in commutative rings”, *Trans. Amer. Math. Soc.* **142** (1969), 43–60. MR 0251026 Zbl 0184.29401
- [Keller 1994] B. Keller, “A remark on the generalized smashing conjecture”, *Manuscripta Math.* **84**:2 (1994), 193–198. MR 1285956 Zbl 0826.18004
- [Mandell 2012] M. A. Mandell, “The smash product for derived categories in stable homotopy theory”, *Adv. Math.* **230**:4–6 (2012), 1531–1556. MR 2927347 Zbl 1246.55010
- [Mathew 2015] A. Mathew, “A thick subcategory theorem for modules over certain ring spectra”, *Geom. Topol.* **19**:4 (2015), 2359–2392. MR 3375530 Zbl 06472920
- [Mathew 2016] A. Mathew, “Residue fields for a class of rational E_∞ -rings and spectra”, *J. Pure Appl. Algebra* (online publication July 2016).
- [Neeman 1992a] A. Neeman, “The chromatic tower for $D(R)$ ”, *Topology* **31**:3 (1992), 519–532. MR 1174255 Zbl 0793.18008
- [Neeman 1992b] A. Neeman, “The connection between the K -theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel”, *Ann. Sci. École Norm. Sup.* (4) **25**:5 (1992), 547–566. MR 1191736 Zbl 0868.19001
- [Park and Park 2011] C. H. Park and M. H. Park, “Integral closure of a graded Noetherian domain”, *J. Korean Math. Soc.* **48**:3 (2011), 449–464. MR 2815884 Zbl 1218.13001
- [Shamir 2012] S. Shamir, “Stratifying derived categories of cochains on certain spaces”, *Math. Z.* **272**:3–4 (2012), 839–868. MR 2995142 Zbl 1271.55010
- [Stevenson 2013] G. Stevenson, “Support theory via actions of tensor triangulated categories”, *J. Reine Angew. Math.* **681** (2013), 219–254. MR 3181496 Zbl 1280.18010
- [Thomason 1997] R. W. Thomason, “The classification of triangulated subcategories”, *Compositio Math.* **105**:1 (1997), 1–27. MR 1436741 Zbl 0873.18003

Received November 16, 2015. Revised May 9, 2016.

IVO DELL'AMBROGIO
LABORATOIRE DE MATHÉMATIQUES PAUL PAINLEVÉ
UNIVERSITÉ DE LILLE 1
BÂT. M2
CITÉ SCIENTIFIQUE
59665 VILLENEUVE D'ASCQ CÉDEX
FRANCE
ivo.dellambrogio@math.univ-lille1.fr

DONALD STANLEY
DEPARTMENT OF MATHEMATICS AND STATISTICS
UNIVERSITY OF REGINA
COLLEGE WEST 307.14
3737 WASCANA PARKWAY
REGINA, SK S4S 0A2
CANADA
donald.stanley@uregina.ca

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2016 is US \$440/year for the electronic version, and \$600/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 285 No. 1 November 2016

Iwahori–Hecke algebras for Kac–Moody groups over local fields	1
NICOLE BARDY-PANSE, STÉPHANE GAUSSENT and GUY ROUSSEAU	
A classification of spherical conjugacy classes	63
MAURO COSTANTINI	
Affine weakly regular tensor triangulated categories	93
IVO DELL’AMBROGIO and DONALD STANLEY	
Involutive automorphisms of N_{\circ}° -groups of finite Morley rank	111
ADRIEN DELORO and ÉRIC JALIGOT	
Schur–Weyl duality for Deligne categories, II: The limit case	185
INNA ENTOVA AIZENBUD	
A generalization of the Greene–Krantz theorem for the semicontinuity property of automorphism groups	225
JAE-CHEON JOO	
Gradient estimates for a nonlinear Lichnerowicz equation under general geometric flow on complete noncompact manifolds	243
LIANG ZHAO and SHOUWEN FANG	