

Pacific Journal of Mathematics

**SCALAR INVARIANTS OF SURFACES IN THE CONFORMAL
3-SPHERE VIA MINKOWSKI SPACETIME**

JIE QING, CHANGPING WANG AND JINGYANG ZHONG

SCALAR INVARIANTS OF SURFACES IN THE CONFORMAL 3-SPHERE VIA MINKOWSKI SPACETIME

JIE QING, CHANGPING WANG AND JINGYANG ZHONG

For a surface in the 3-sphere, by identifying the conformal round 3-sphere as the projectivized positive light cone in Minkowski 5-spacetime, we use the conformal Gauss map and the conformal transform to construct the associate homogeneous 4-surface in Minkowski 5-spacetime. We then derive the local fundamental theorem for a surface in the conformal round 3-sphere from that of the associate 4-surface in Minkowski 5-spacetime. More importantly, following an idea of Fefferman and Graham, we construct local scalar invariants for a surface in the conformal round 3-sphere. One distinct feature of our construction is to link the classic work of Blaschke to the work of Bryant and Fefferman and Graham.

1. Introduction

It is well-known that all local scalar invariants of a (pseudo-)Riemannian metric are Weyl invariants, based on Weyl's classical invariant theory for the orthogonal groups. A conformal structure on a manifold is described by an equivalent class of conformal Riemannian metrics. Two metrics g_1 and g_2 on a manifold M are conformal to each other if $g_1 = \lambda^2 g_2$ for some positive smooth function λ on M . There are several ways to set the theory of local conformal invariants, but it is no longer straightforward to account for local scalar conformal invariants because of the lack of Weyl Theorem for the group of conformal transformations. To tackle this problem, Fefferman and Graham, in a seminal paper [1985], described an ingenious construction for a Ricci-flat homogeneous Lorentzian ambient spacetime for a given conformal manifold, where the conformal manifold is represented by the homogeneous null hypersurface in the ambient spacetime. Their construction was motivated by the model case in which the conformal round sphere \mathbb{S}^n is the projectivized positive light cone \mathbb{N}_+^{n+1} in Minkowski spacetime $\mathbb{R}^{1,n+1}$. Thus they initiated the program of using local scalar (pseudo-)Riemannian invariants of the ambient metrics at the homogeneous null hypersurface to fully account for local scalar conformal invariants. Readers are referred to their recent expository paper

MSC2010: 53A30, 53B25.

Keywords: scalar invariant, conformal geometry, conformal Gauss map, Willmore surface.

[Fefferman and Graham 2012] to learn all the developments of this program, and also to [Bailey et al. 1994; Gover 2001]. This program has led to many significant advances in the global theory of conformal geometry, particularly via conformally invariant PDEs.

In this paper we build a model case for the study of local scalar invariants of submanifolds in a conformal manifold, following Fefferman and Graham's approach. The model case for us is to study 2-surfaces \hat{x} in the conformal round 3-sphere $(\mathbb{S}^3, [g_0])$. As in [Fefferman and Graham 1985], the conformal round 3-sphere is represented by the positive light cone \mathbb{N}_+^4 in Minkowski 5-spacetime $\mathbb{R}^{1,4}$. Given an immersed surface

$$\hat{x} : M^2 \rightarrow \mathbb{S}^3$$

or equivalently

$$y = (1, \hat{x}) : M^2 \rightarrow \mathbb{N}_+^4,$$

to incorporate all metrics in $[g_0]$ on the 3-sphere we consider the homogeneous extension

$$x^{\mathbb{N}} = \alpha(1, \hat{x}) : \mathbb{R}^+ \times M^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}.$$

Then we will use the conformal Gauss map ξ of \hat{x} to choose a canonical null vector y^* at each given point $y \in x^{\mathbb{N}} \subset \mathbb{N}_+^4$ to extend $x^{\mathbb{N}}$ further into a homogeneous timelike 4-surface

$$\tilde{x} = \alpha y + \alpha \rho y^* : \mathbb{R}^+ \times \mathbb{R}^+ \times M^2 \rightarrow \mathbb{R}^{1,4}.$$

We will also consider the associate ruled 3-surface

$$x^+ = \frac{1}{\sqrt{2}}(e^t y + e^{-t} y^*) : \mathbb{R} \times M^2 \rightarrow \mathbb{H}^4 \subset \mathbb{R}^{1,4}$$

where \mathbb{H}^4 is the hyperboloid in Minkowski 5-spacetime. The main idea, inspired by Fefferman and Graham's work, is to use the geometry of the associate 4-surface \tilde{x} in Minkowski spacetime $\mathbb{R}^{1,4}$ (the associate ruled 3-surface x^+ in the hyperboloid \mathbb{H}^4 and the spacelike surface as the image of the conformal Gauss map ξ in the de Sitter spacetime $S^{3,1}$ in Minkowski spacetime $\mathbb{R}^{1,4}$) to study the geometry of the surface \hat{x} in the conformal round 3-sphere \mathbb{S}^3 .

Our approach facilitates proofs of the local fundamental theorems (see [Theorem 3.3.1](#) and [Wang 1992; 1998]) and produces local scalar invariants of surfaces in the conformal round 3-sphere. It is more interesting to find scalar invariants and the PDE problems similar to the study of Willmore surfaces [Blaschke 1929; Bryant 1984; Li and Yau 1982; Marques and Neves 2014].

We remark that the key to our construction of associate surfaces is the conformal Gauss map ξ to a given surface \hat{x} in the conformal round 3-sphere. Conformal

Gauss maps have been introduced in several contexts [Blaschke 1929; Bryant 1984; Rigoli 1987]. We are searching for a definition that fits into the context of ambient spaces of Fefferman and Graham (Lemmas 2.3.1 and 2.3.2). It is fascinating to see how Blaschke [1929] introduced the conformal Gauss map as the map representing the family of mean curvature 2-spheres of the surface \hat{x} and the conformal transform \hat{x}^* (Definition 2.4.1) as the other envelope surface of the conformal Gauss map. One technical assumption for the null vector y^* to be well defined at each point $y \in x^{\mathbb{N}}$ is to require that the conformal Gauss map of the surface \hat{x} induces a spacelike surface in the de Sitter spacetime $\mathbb{S}^{1,3}$, which is equivalent to require the surface \hat{x} is free of umbilical point in the conformal 3-sphere \mathbb{S}^3 .

It is nice to know that in our construction the associate 4-surface \tilde{x} in Minkowski spacetime $\mathbb{R}^{1,4}$ is a minimal 4-surface (of vanishing mean curvature) if and only if the 2-surface \hat{x} is a Willmore surface with no umbilical point in \mathbb{S}^3 (Theorem 3.2.1). The same statement also holds for the associate ruled 3-surface x^+ in the hyperboloid \mathbb{H}^4 (Theorem 3.4.1) and the conformal Gauss map surface ξ in de Sitter spacetime $\mathbb{S}^{1,3}$ (Theorem 2.5.2).

Upon realizing that a different representative $\lambda^2 g_0$ in the conformal class $[g_0]$ on \mathbb{S}^3 is equivalent to a different parametrization for the associate surface

$$(1.0.1) \quad \tilde{x} = \alpha y_\lambda + \alpha \rho y_\lambda^* : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{M}^2 \rightarrow \mathbb{R}^{1,4},$$

where $y_\lambda = \hat{\lambda}(1, \hat{x})$ and $\hat{\lambda} = \lambda \circ \hat{x}$ for a conformal factor λ , the real issue is how we use the geometry of the surface \hat{x} in the 3-sphere $(\mathbb{S}^3, \lambda^2 g_0)$ to calculate the geometry of the associate surface \tilde{x} . The solution is to use as the realization of $(\mathbb{S}^3, \lambda^2 g_0)$ the following 3-sphere \mathbb{S}_λ^3 in the positive light cone \mathbb{N}_+^4 :

$$(1.0.2) \quad \lambda(1, x) : \mathbb{S}^3 \rightarrow \mathbb{N}_+^4.$$

For the convenience of readers we present the calculations of the geometry of \mathbb{S}_λ^3 as a spacelike 3-surface in Minkowski spacetime in Appendix B. But it starts with the observation that the conformal Gauss map

$$(1.0.3) \quad \xi = H_\lambda y_\lambda + \vec{n}_\lambda : \mathbb{M}^2 \rightarrow \mathbb{R}^{1,4},$$

where H_λ is the mean curvature of \hat{x} in $(\mathbb{S}^3, \lambda^2 g_0)$ and \vec{n}_λ is the unit normal to y_λ in $\mathbb{S}_\lambda^3 \subset \mathbb{N}_+^4$, is independent of the conformal metric $\lambda^2 g_0$ (cf. Lemma 2.3.4).

Using the calculations in Appendix B, we are able to show in the proof of Theorem 4.3.2 that the data $\{m, \omega^\lambda, \Omega_\lambda, \Omega_\lambda^*\}$ that determine the first and second fundamental forms of the associate surface \tilde{x} in Minkowski spacetime $\mathbb{R}^{1,4}$ can all be expressed in terms of covariant derivatives of the curvature of the surface \hat{x} in $(\mathbb{S}^3, \lambda^2 g_0)$ and the covariant derivatives of curvature of $(\mathbb{S}^3, \lambda^2 g_0)$ (including 0th order). In the exact same spirit as in [Fefferman and Graham 1985; 2012],

our construction of associate surfaces \tilde{x} provides a way to capture local scalar conformal invariants of a surface \hat{x} . Namely, one can obtain local scalar conformal invariants of the surface \hat{x} in the conformal round 3-sphere by computing the local scalar (pseudo-)Riemannian invariants of the associate surface \tilde{x} at the homogeneous surface $x^{\mathbb{N}}$ in the light cone in Minkowski 5-spacetime. The first nontrivial one is

$$(1.0.4) \quad \tilde{\Delta} \tilde{H}|_{\rho=0} = 2\alpha^{-3}(\Delta_\lambda H_\lambda + |\mathring{H}_\lambda|^2 H_\lambda + (\mathring{H}_\lambda)^{ij}(R^\lambda)_{i3j3} - (R^\lambda)_{3i}{}^i)$$

in a general parametrization (1.0.1), where $(R^\lambda)_{i3j3}$ and $(R^\lambda)_{3i}{}^i$ are the Riemann curvature and Ricci curvature of the metric $\lambda^2 g_0$ on \mathbb{S}^3 . Due to the homogeneity of \tilde{x} we automatically have

$$(1.0.5) \quad \mathcal{H}_\lambda = \Delta_\lambda H_\lambda + |\mathring{H}_\lambda|^2 H_\lambda + (\mathring{H}_\lambda)^{ij}(R^\lambda)_{i3j3} - (R^\lambda)_{3i}{}^i = \hat{\lambda}^{-3}(\Delta H + |\mathring{H}|^2 H)$$

which is the curvature that vanishes if and only if the surface \hat{x} is Willmore. Notice that extra curvature terms do not show up when we work with either the round metric g_0 or the Euclidean metric. Similar formulas have appeared in the literature [Hu and Li 2004; Gover and Waldron 2015] and are also used by R. Graham and N. Reichert (work in progress).

In (4.2.7) we obtain this conformal scalar invariant of higher order:

$$(1.0.6) \quad |\nabla \tilde{h}|^2|_{\rho=0} = \alpha^{-4}(|\nabla \Omega_\lambda|^2 + 8|dH_\lambda|^2 + 2\text{Ric}^\lambda(\vec{n}_\lambda, \nabla H_\lambda) + 3H_\lambda^2|\Omega_\lambda|^2 + 3K_\lambda^T|\Omega_\lambda|^2 + 6\Omega_\lambda \cdot \text{Hess}(H_\lambda)),$$

where K_λ^T is the sectional curvature of $(\mathbb{S}^3, \lambda^2 g_0)$ at the tangent plane to the surface \hat{x} . Another higher-order invariant is

$$(1.0.7) \quad \tilde{\Delta} \tilde{\Delta} \tilde{H}|_{\rho=0} = 8\alpha^{-5}(\Delta_\lambda \mathcal{H}_\lambda + 9|\omega^\lambda|^2 \mathcal{H}_\lambda - 3\text{Div}(\omega^\lambda) \mathcal{H}_\lambda - 6\omega^\lambda(\nabla \mathcal{H}_\lambda) - 6\mathcal{H}_\lambda|\mathring{H}_\lambda|^{-2} \mathring{H}_\lambda \cdot \Omega_\lambda^*),$$

where $\omega^\lambda = \langle dy_\lambda, y_\lambda^* \rangle$ and $\Omega_\lambda^* = -\langle dy_\lambda^*, d\xi \rangle$ are parts of the data that determine the geometry of the associate surface \tilde{x} ; they are given in (3.1.3) and (4.3.6) as invariants of the surface \hat{x} in $(\mathbb{S}^3, \lambda^2 g_0)$.

To end the introduction we remark that, for the sake of the production of local scalar invariants, the assumption of having no umbilical point in our construction is not an issue.

2. The associate surfaces in $\mathbb{R}^{1,4}$

In this section we introduce the associate surfaces in Minkowski space $\mathbb{R}^{1,4}$ for a given surface $\hat{x} : M^2 \rightarrow \mathbb{S}^3$. We then show that such associate surface is canonical

in doing conformal geometry for the surface \hat{x} . The construction relies on the conformal Gauss map and the conformal transform of \hat{x} . It is also very interesting to see how Blaschke and Bryant came to the conformal Gauss map and the conformal transform in very different perspectives [Blaschke 1929; Bryant 1984].

2.1. Surfaces in the 3-sphere. Suppose that

$$\hat{x} : M^2 \rightarrow S^3 \subset \mathbb{R}^4$$

is an immersed surface with isothermal coordinate (u^1, u^2) . Let

$$\mathbf{n} : M^2 \rightarrow \mathbb{R}^4$$

be the unit normal vector at each point on the surface. Then we obtain the first fundamental form

$$(2.1.1) \quad I = \langle d\hat{x}, d\hat{x} \rangle = E|du|^2$$

and the second fundamental form

$$(2.1.2) \quad II = -\langle d\hat{x}, d\mathbf{n} \rangle = e(du^1)^2 + 2fdu^1du^2 + g(du^2)^2.$$

Hence the mean curvature of the surface in the 3-sphere is

$$(2.1.3) \quad H = \frac{1}{2E}(e + g)$$

and the Gaussian curvature of the surface is

$$(2.1.4) \quad K = \frac{eg - f^2}{E^2} + 1.$$

Notice that

$$(2.1.5) \quad \mathbf{n}_{u^1} = -\frac{e}{E}\hat{x}_{u^1} - \frac{f}{E}\hat{x}_{u^2}, \quad \mathbf{n}_{u^2} = -\frac{f}{E}\hat{x}_{u^1} - \frac{g}{E}\hat{x}_{u^2}.$$

If one takes another conformal metric $\lambda^2 g_0$ on the 3-sphere S^3 , where λ is a positive function on S^3 , then the first fundamental form for the surface \hat{x} is

$$(2.1.6) \quad I_\lambda = \hat{\lambda}^2 I,$$

where $\hat{\lambda} = \lambda \circ \hat{x}$ and the second fundamental form is

$$(2.1.7) \quad II_\lambda = \hat{\lambda} II - \lambda_n I,$$

where $\lambda_n = \mathbf{n}(\lambda)$. Hence

$$(2.1.8) \quad H_\lambda = \hat{\lambda}^{-1} \left(H - \frac{\lambda_n}{\hat{\lambda}} \right) \quad \text{and} \quad \mathring{H}_\lambda = \hat{\lambda} \mathring{H},$$

where $\overset{\circ}{I}$ is the traceless part of the second fundamental form II . Here we see the easy scalar conformal invariant $|\overset{\circ}{I}|^2$, which can be considered to be the counter part of the square of the length of Weyl curvature on a conformal manifold.

2.2. Minkowski 5-spacetime. Let $\mathbb{R}^{1,4}$ be the Minkowski 5-spacetime, where we use the notation

$$\mathbb{R}^{1,4} = \{(t, x) : t \in \mathbb{R} \text{ and } x \in \mathbb{R}^4\}$$

with the Lorentz inner product

$$\langle (t, x), (s, y) \rangle = -st + x \cdot y.$$

Recall the positive light cone is given by

$$\mathbb{N}_+^4 = \{(t, x) \in \mathbb{R}^{1,4} : -t^2 + |x|^2 = 0 \text{ and } t > 0\};$$

the hyperboloid is given as

$$\mathbb{H}^4 = \{(t, x) \in \mathbb{R}^{1,4} : -t^2 + |x|^2 = -1 \text{ and } t > 0\};$$

and the de Sitter 4-spacetime is given as

$$\mathbb{S}^{1,3} = \{(t, x) \in \mathbb{R}^{1,4} : -t^2 + |x|^2 = 1\}.$$

Given a surface $\hat{x} : \mathbb{M}^2 \rightarrow \mathbb{S}^3 \subset \mathbb{R}^4$, we may consider the 2-surface

$$y = (1, \hat{x}) : \mathbb{M}^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$$

and the homogeneous extension

$$x^{\mathbb{N}} = \alpha y : \mathbb{R}^+ \times \mathbb{M}^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$$

for $\alpha \in \mathbb{R}^+$. There does not seem to be a way of doing “geometry” of the homogeneous 3-surface $x^{\mathbb{N}}$ in the positive light cone \mathbb{N}_+^4 .

To motivate our choice of the associate surface in $\mathbb{R}^{1,4}$ of \hat{x} we first introduce the so-called homogeneous coordinate for $\mathbb{R}^{1,4}$ used in the ambient space construction of [Fefferman and Graham 1985; 2012], that is,

$$(2.2.1) \quad (t, x) = x^0(1, \hat{x}) + x^0 x^\infty \frac{1}{2}(1, -\hat{x})$$

where

$$x^0 = \frac{1}{2}(r + t), \quad x^0 x^\infty = (-r + t)$$

and $r = |x|$ and $x = r\hat{x}$. In this coordinate the Minkowski metric is

$$\tilde{g}_0 = -2x^\infty(dx^0)^2 - 2x^0 dx^0 dx^\infty + (x^0)^2(1 - \frac{1}{2}x^\infty)^2 g_0(\hat{x}).$$

Hence, given a surface $\hat{x} : \mathbf{M}^2 \rightarrow \mathbb{S}^3$, we are looking to construct an associate homogeneous timelike 4-surface

$$(2.2.2) \quad \tilde{x} = \alpha y + \alpha \rho y^* : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbf{M}^2 \rightarrow \mathbb{R}^{1,4}$$

if we can have canonically the null vector y^* at a given null position y on $x^{\mathbb{N}}$. It is clear that the associate surface \tilde{x} is ruled by the positive quadrants of timelike 2-planes in Minkowski spacetime. One may consider the intersection of \tilde{x} with the hyperboloid \mathbb{H}^4 :

$$(2.2.3) \quad x^+ = \frac{1}{\sqrt{2}}(e^t y + e^{-t} y^*) : \mathbb{R} \times \mathbf{M}^2 \rightarrow \mathbb{H}^4,$$

which is called the associate ruled 3-surface since it is a 3-surface in hyperbolic 4-space ruled by geodesics lines. Recall that a geodesic line in the hyperboloid \mathbb{H}^4 is the intersection of the hyperboloid with a timelike 2-subspaces in Minkowski spacetime. In the following we will introduce the canonical choice of such y^* .

2.3. Conformal Gauss maps. Let us consider any unit spacelike normal vector to the homogeneous null 3-surface $x^{\mathbb{N}} = \alpha y$ in $\mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$. That is to ask a unit spacelike 5-vector ξ to satisfy

$$(2.3.1) \quad \langle \xi, x^{\mathbb{N}} \rangle = 0, \quad \langle \xi, x_{u^1}^{\mathbb{N}} \rangle = 0, \quad \langle \xi, x_{u^2}^{\mathbb{N}} \rangle = 0,$$

which implies that

$$\xi = \alpha y + \vec{n},$$

where $\vec{n} = (0, \mathbf{n})$ is the unit normal to the surface \hat{x} in the standard unit round 3-sphere in $\{1\} \times \mathbb{R}^4 \subset \mathbb{R}^{1,4}$. It turns out that there is a unique choice if we insist that the map

$$\xi : \mathbf{M}^2 \rightarrow \mathbb{S}^{1,3} \subset \mathbb{R}^{1,4}$$

is (weakly) conformal. Namely we have

Lemma 2.3.1. *Suppose that $\hat{x} : \mathbf{M}^2 \rightarrow \mathbb{S}^3$ is an immersed surface. For a unit normal vector ξ to the homogeneous null 3-surface $x^{\mathbb{N}} = \alpha y : \mathbb{R}^+ \times \mathbf{M}^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$,*

$$\langle \xi_{u^1}, \xi_{u^2} \rangle = 0$$

if and only if

$$\xi = H y + \vec{n}$$

and

$$(2.3.2) \quad \langle d\xi, d\xi \rangle = \frac{1}{2} E |\hat{H}|^2 |du|^2.$$

Proof. It is simply a straightforward calculation. We know

$$\xi_{u^i} = a_{u^i}(1, \hat{x}) + a(0, \hat{x}_{u^i}) + (0, \mathbf{n}_{u^i}).$$

Hence we have

$$\langle \xi_{u^1}, \xi_{u^2} \rangle = -2af + \frac{1}{E}(fe + fg) = 0,$$

which is equivalent to $a = H$. For the rest we calculate

$$\langle \xi_{u^1}, \xi_{u^1} \rangle = \langle \xi_{u^2}, \xi_{u^2} \rangle = \frac{1}{E^2} \left(f^2 + \left(\frac{e-g}{2} \right)^2 \right) E. \quad \square$$

Another way to identify a unique unit spacelike normal vector to the homogeneous null 3-surface $x^{\mathbb{N}} = \alpha y : \mathbb{R}^+ \times \mathbb{M}^2 \rightarrow \mathbb{N}_+^4$ is the following:

Lemma 2.3.2. *Suppose that $\hat{x} : \mathbb{M}^2 \rightarrow \mathbb{S}^3$ is an immersed surface. Then, for a unit spacelike normal vector ξ to $x^{\mathbb{N}} = \alpha y : \mathbb{R}^+ \times \mathbb{M}^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$,*

$$(2.3.3) \quad \xi = Hy + \vec{\mathbf{n}}$$

if and only if

$$(2.3.4) \quad \langle \Delta \xi, y \rangle = 0.$$

Proof. We simply calculate, for $\xi = a(1, \hat{x}) + (0, \mathbf{n})$,

$$\Delta_0 \xi = \xi_{u^1 u^1} + \xi_{u^2 u^2} = (\Delta_0 a)(1, \hat{x}) + 2\nabla a(0, \nabla \hat{x}) + a(0, \Delta_0 \hat{x}) + (0, \Delta_0 \mathbf{n})$$

and

$$\langle \Delta_0 \xi, (1, \hat{x}) \rangle = -2aE + 2HE.$$

Notice that $\Delta = E^{-1} \Delta_0$. □

Before we give a formal definition of the conformal Gauss map we remark that (2.3.4) is the integrability condition for the unit vector field ξ to be the conformal Gauss map (up to a sign) for the surface \hat{x} . This turns out to be the easiest way to see that \hat{x} is Willmore if and only if the conformal Gauss map ξ of \hat{x} is also the conformal Gauss map (up to a sign) of the conformal transform \hat{x}^* (see Definition 2.4.1).

Definition 2.3.3. Suppose that $\hat{x} : \mathbb{M}^2 \rightarrow \mathbb{S}^3$ is a surface. Then we will call

$$(2.3.5) \quad \xi = Hy + \vec{\mathbf{n}} : \mathbb{M}^2 \rightarrow \mathbb{S}^{1,3} \subset \mathbb{R}^{1,4}$$

the conformal Gauss map according to Blaschke (cf. [Bryant 1984; Rigoli 1987]).

For a positive function λ on the sphere \mathbb{S}^3 we consider the conformal metric $\lambda^2 g_0$ on the sphere \mathbb{S}^3 , which can be realized as the 3-sphere $\mathbb{S}_\lambda^3 : \lambda(1, x) : \mathbb{S}^3 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$ in Minkowski spacetime. It is essential here that the surface \hat{x} in the 3-sphere \mathbb{S}^3 with the conformal metric $\lambda^2 g_0$ is realized as the 2-surface $\hat{\lambda}(1, \hat{x}) :$

$M^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$ inside the 3-sphere \mathbb{S}_λ^3 . It is helpful to see the calculations in [Appendix B](#) about the geometry of the 3-sphere \mathbb{S}_λ^3 in Minkowski spacetime $\mathbb{R}^{1,4}$.

Lemma 2.3.4. *If one works with a conformal metric $\lambda^2 g_0$ in general, then*

$$(2.3.6) \quad \xi = \xi_\lambda = H_\lambda y_\lambda + \vec{n}_\lambda,$$

where $\vec{n}_\lambda = \vec{n} + (\log \lambda)_n y$ is the unit normal to the surface

$$y_\lambda = \hat{\lambda}(1, \hat{x}) : M^2 \rightarrow \mathbb{S}_\lambda^3 \subset \mathbb{N}_+^4.$$

Proof. It is easily seen that the normal direction to the surface y_λ inside \mathbb{S}_λ^3 is $\lambda_n(1, \hat{x}) + \lambda(0, \mathbf{n})$ and that $\langle \lambda_n(1, \hat{x}) + \lambda(0, \mathbf{n}), \lambda_n(1, \hat{x}) + \lambda(0, \mathbf{n}) \rangle = \lambda^2$. Therefore the unit normal for the surface y_λ in \mathbb{S}_λ^3 is $\vec{n}_\lambda = \vec{n} + (\log \lambda)_n y$. Hence [\(2.1.8\)](#) yields

$$H_\lambda y_\lambda + \vec{n}_\lambda = H y + \vec{n}. \quad \square$$

In light of [\(2.3.2\)](#), the conformal Gauss map gives rise a spacelike 2-surface

$$\xi : M^2 \rightarrow \mathbb{S}^{1,3} \subset \mathbb{R}^{1,4}$$

when the original surface $\hat{x} : M^2 \rightarrow \mathbb{S}^3$ is free of umbilical points. We will have more detailed discussions for the reasons to call ξ the conformal Gauss map in [Section 2.7](#).

It is interesting that Blaschke came across the conformal Gauss map from a different perspective. He considered the family of mean curvature 2-spheres to the surface \hat{x} in \mathbb{S}^3 . A round 2-sphere in 3-sphere can be thought of as the intersection of a timelike hyperplane and the 3-sphere at time $t = 1$ in Minkowski spacetime $\mathbb{R}^{1,4}$ and a timelike hyperplane in $\mathbb{R}^{1,4}$ is described by a unit normal vector lying in de Sitter 4-spacetime $\mathbb{S}^{1,3}$. Given a direction $(H, H\hat{x} + \mathbf{n}) \in \mathbb{S}^{1,3}$, the hyperplane perpendicular to that in $\mathbb{R}^{1,4}$ is given by the first equation in [\(2.3.1\)](#):

$$(2.3.7) \quad \langle (s, z), (H, H\hat{x} + \mathbf{n}) \rangle = 0,$$

which is

$$-sH + Hz \cdot \left(\hat{x} + \frac{1}{H} \mathbf{n} \right) = 0.$$

At the level $s = 1$ in the 3-sphere $|z| = 1$, we arrive at

$$1 - \hat{z} \cdot \left(\hat{x} + \frac{1}{H} \mathbf{n} \right) = 0.$$

We may rewrite this as

$$(2.3.8) \quad \left| \hat{z} - \left(\hat{x} + \frac{1}{H} \mathbf{n} \right) \right|^2 = \frac{1}{H^2}$$

which clearly is a round 2-sphere of mean curvature H when intersects with the 3-sphere $\mathbb{S}^3 \subset \mathbb{R}^4$ at $t = 1$ in $\mathbb{R}^{1,4}$. Hence the equations [\(2.3.1\)](#) exactly ask the

surface $y = (1, \hat{x}) : M^2 \rightarrow \mathbb{S}^3 \subset \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$ to be an envelope surface of the family of mean curvature 2-spheres described by the conformal Gauss map ξ .

It is known that a mean curvature sphere of a surface goes to the mean curvature sphere of the image surface under conformal transformations.

2.4. Conformal transforms. Assume that the surface $\hat{x} : M^2 \rightarrow \mathbb{S}^3$ is free of umbilical points. Then the conformal Gauss map induces a spacelike 2-surface in the de Sitter 4-space $\mathbb{S}^{1,3}$

$$\xi : M^2 \rightarrow \mathbb{S}^{1,3} \subset \mathbb{R}^{1,4}.$$

One notices that the equations (2.3.1) imply that $y = (1, \hat{x})$ is naturally a null normal vector the surface ξ in the de Sitter 4-spacetime $\mathbb{S}^{1,3}$. Because

$$\langle y, \xi_{u^i} \rangle = -\langle \xi, y_{u^i} \rangle = 0.$$

Hence it is natural to take the other null normal vector y^* such that

$$(2.4.1) \quad \begin{aligned} \langle y^*, y \rangle &= -1, & \langle y^*, y^* \rangle &= 0, & \langle y^*, \xi \rangle &= 0, \\ \langle y^*, \xi_{u^1} \rangle &= 0 & \text{and} & & \langle y^*, \xi_{u^2} \rangle &= 0. \end{aligned}$$

We may write

$$y^* = \hat{\mu}^*(1, \hat{x}^*).$$

Definition 2.4.1. Suppose that $\hat{x} : M^2 \rightarrow \mathbb{S}^3$ is a surface with no umbilical point. And suppose that

$$y^* = \hat{\mu}^*(1, \hat{x}^*) : M^2 \rightarrow \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$$

satisfies the equations (2.4.1) for $y = (1, \hat{x})$. Then the surface

$$\hat{x}^* : M^2 \rightarrow \mathbb{S}^3$$

is said to be the conformal transform of the surface \hat{x} according to [Bryant 1984] (cf. [Blaschke 1929]).

It is important that the conformal transform \hat{x}^* of a surface \hat{x} is independent of the conformal factor λ . Notice that the equations in (2.4.1) remain the same except the first one when replacing y by y_λ . It is again very interesting to recall how Blaschke discovered the surface \hat{x}^* . From the above discussions it is now easy to see that the surface \hat{x}^* is nothing but the other envelope surface of the family of round 2-spheres described by the conformal Gauss map ξ , i.e., the family of the mean curvature spheres of the surface \hat{x} , since y^* satisfies the last three equations in (2.4.1).

2.5. The geometry of the surface ξ in $\mathbb{S}^{1,3}$. Recall that the first fundamental form for the surface ξ in the de Sitter spacetime $\mathbb{S}^{1,3} \subset \mathbb{R}^{1,4}$ is

$$(2.5.1) \quad I^\xi = \langle d\xi, d\xi \rangle = m|du|^2,$$

where

$$(2.5.2) \quad m = \frac{1}{2}E|\mathring{H}|^2.$$

The first fundamental form I^ξ is usually called the Möbius metric on the surface \hat{x} . If one works with a conformal metric $\lambda^2 g_0$ instead, then the Möbius metric remains the same:

$$(2.5.3) \quad m = m_\lambda = \frac{1}{2}E_\lambda|\mathring{H}_\lambda|^2.$$

The second fundamental form for the surface ξ in $\mathbb{S}^{1,3}$ is given by

$$II^\xi = -\langle d\xi, dy \rangle y - \langle d\xi, dy^* \rangle y^* = \Omega y + \Omega^* y^* = \Omega_\lambda \hat{\lambda}^{-2} y_\lambda + \Omega_\lambda^* \hat{\lambda}^2 y_\lambda^*$$

and

$$(2.5.4) \quad \begin{aligned} \Omega_{ij} &= -\langle \xi_{u^i}, y_{u^j} \rangle, & \Omega_{ij}^* &= -\langle \xi_{u^i}, y_{u^j}^* \rangle, \\ (\Omega_\lambda)_{ij} &= -\langle \xi_{u^i}, (y_\lambda)_{u^j} \rangle = \hat{\lambda} \Omega_{ij}, & (\Omega_\lambda^*)_{ij} &= -\langle \xi_{u^i}, (y_\lambda^*)_{u^j} \rangle = \hat{\lambda}^{-1} \Omega_{ij}^*. \end{aligned}$$

In fact it is easy to calculate that

$$(2.5.5) \quad \Omega = \begin{bmatrix} \frac{e-g}{2} & f \\ f & \frac{g-e}{2} \end{bmatrix} = \mathring{H}.$$

Let us first calculate the mean curvature in the y^* direction. We notice that

$$\langle \Delta_0 \xi, y_\lambda^* \rangle = ((\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22})$$

while

$$\langle \Delta_0 \xi, y_\lambda \rangle = ((\Omega_\lambda)_{11} + (\Omega_\lambda)_{22}) = 0.$$

Based on the calculations

$$\begin{aligned} \langle \Delta_0 \xi, \xi \rangle &= -2m, \\ \langle \Delta_0 \xi, \xi_{u^1} \rangle &= \frac{1}{2}m_{u^1} - \frac{1}{2}m_{u^1} = 0, \\ \langle \Delta_0 \xi, \xi_{u^2} \rangle &= -\frac{1}{2}m_{u^2} + \frac{1}{2}m_{u^2} = 0, \end{aligned}$$

we obtain

$$(2.5.6) \quad \begin{aligned} \Delta_0 \xi &= -((\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22})y_\lambda - 2m\xi \\ &= (-((\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22}) - 2mH_\lambda)y_\lambda - 2m\vec{n}_\lambda. \end{aligned}$$

On the other hand, we directly calculate

$$(2.5.7) \quad \begin{aligned} \Delta_0 \xi &= \Delta_0(H_\lambda y_\lambda + \vec{n}_\lambda) \\ &= (\Delta_0 H_\lambda) y_\lambda + H_\lambda \Delta_0 y_\lambda + 2(H_\lambda)_{u^1} (y_\lambda)_{u^1} + 2(H_\lambda)_{u^2} (y_\lambda)_{u^2} + \Delta_0 \vec{n}_\lambda. \end{aligned}$$

It seems that the best way to calculate geometrically is to use the Lorentz orthogonal frame

$$\{y_\lambda, y_\lambda^\dagger, (y_\lambda)_{u^1}, (y_\lambda)_{u^2}, \vec{n}_\lambda\},$$

where

$$(2.5.8) \quad \langle y_\lambda^\dagger, y_\lambda \rangle = -1, \quad \langle y_\lambda^\dagger, y_\lambda^\dagger \rangle = \langle y_\lambda^\dagger, (y_\lambda)_{u^1} \rangle = \langle y_\lambda^\dagger, (y_\lambda)_{u^2} \rangle = \langle y_\lambda^\dagger, \vec{n}_\lambda \rangle = 0.$$

We find that

$$(2.5.9) \quad y_\lambda^\dagger = \frac{1}{\lambda} \left(\frac{1}{2} |\nabla \log \lambda|^2 y + y^\dagger - \nabla \log \lambda \right),$$

where $y^\dagger = \frac{1}{2}(1, -\hat{x})$ and ∇ is the gradient on the standard round 3-sphere. We will apply the inner product with the null vector y_λ^\dagger to both (2.5.6) and (2.5.7). To calculate $H_\lambda \langle \Delta_0 y_\lambda, y_\lambda^\dagger \rangle + \langle \Delta_0 \vec{n}_\lambda, y_\lambda^\dagger \rangle$ we rewrite

$$H_\lambda \langle \Delta_0 y_\lambda, y_\lambda^\dagger \rangle = -H_\lambda (\langle (y_\lambda)_{u^1}, (y_\lambda^\dagger)_{u^1} \rangle + \langle (y_\lambda)_{u^2}, (y_\lambda^\dagger)_{u^2} \rangle)$$

and

$$\langle \Delta_0 \vec{n}_\lambda, y_\lambda^\dagger \rangle = -\langle (\vec{n}_\lambda)_{u^1}, (y_\lambda^\dagger)_{u^1} \rangle - \langle (\vec{n}_\lambda)_{u^2}, (y_\lambda^\dagger)_{u^2} \rangle - \langle \vec{n}_\lambda, (y_\lambda^\dagger)_{u^i} \rangle_{u^i}.$$

Meanwhile one may calculate

$$(2.5.10) \quad \begin{cases} (\vec{n}_\lambda)_{u^1} = -\frac{e_\lambda}{E_\lambda} (y_\lambda)_{u^1} - \frac{f_\lambda}{E_\lambda} (y_\lambda)_{u^2} - \langle (\vec{n}_\lambda)_{u^1}, y_\lambda^\dagger \rangle y_\lambda \\ (\vec{n}_\lambda)_{u^2} = -\frac{f_\lambda}{E_\lambda} (y_\lambda)_{u^1} - \frac{g_\lambda}{E_\lambda} (y_\lambda)_{u^2} - \langle (\vec{n}_\lambda)_{u^2}, y_\lambda^\dagger \rangle y_\lambda. \end{cases}$$

Hence we have

$$(2.5.11) \quad \begin{aligned} H_\lambda \langle \Delta_0 y_\lambda, y_\lambda^\dagger \rangle + \langle \Delta_0 \vec{n}_\lambda, y_\lambda^\dagger \rangle \\ = E_\lambda^{-1} (\mathring{H}_\lambda)_{ij} \langle (y_\lambda)_{u^i}, (y_\lambda^\dagger)_{u^j} \rangle - \langle \vec{n}_\lambda, (y_\lambda^\dagger)_{u^i} \rangle_{u^i} \\ = -E_\lambda^{-1} (\mathring{H}_\lambda)_{ij} R_{i3j3}^\lambda + E_\lambda (R^\lambda)_{3i}{}^i \end{aligned}$$

due to (B.6), (B.7), and (B.8). Now we obtain the mean curvature of the surface ξ in the de Sitter spacetime $\mathbb{S}^{1,3}$.

Lemma 2.5.1. *Suppose that $\hat{x}: M^2 \rightarrow \mathbb{S}^3$ is an immersed surface with no umbilical point and that $\xi: M^2 \rightarrow \mathbb{S}^{1,3}$ is the conformal Gauss map. Then the surface ξ is*

spacelike and its mean curvature is a null vector

$$(2.5.12) \quad H^\xi = 2\hat{\lambda}^2 \frac{\mathcal{H}_\lambda}{|\mathring{H}_\lambda|^2} y_\lambda^*$$

for any positive function λ on the 3-sphere \mathbb{S}^3 , where

$$(2.5.13) \quad \mathcal{H}_\lambda = \Delta_\lambda H_\lambda + |\mathring{H}_\lambda|^2 H_\lambda + (\mathring{H}_\lambda)^{ij} (R^\lambda)_{i3j3} - (R^\lambda)_{3i}{}^i,$$

and $(R^\lambda)_{ijkl}$, $(R^\lambda)_{ij}$ are the Riemann curvature and Ricci curvature for the conformal metric $\lambda^2 g_0$ on the 3-sphere \mathbb{S}^3 .

Proof. We perform inner product to (2.5.6) and (2.5.7) by the null vector y_λ^\dagger and obtain that

$$(2.5.14) \quad (\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22} = E_\lambda (-\Delta_\lambda H_\lambda - |\mathring{H}_\lambda|^2 H_\lambda - (\mathring{H}_\lambda)^{ij} (R^\lambda)_{i3j3} + (R^\lambda)_{3i}{}^i)$$

in the light of (2.5.11). Then one can easily calculate the mean curvature for ξ in $\mathbb{S}^{1,3}$. □

We remark that (2.5.12) actually shows that

$$(2.5.15) \quad \mathcal{H}_\lambda = \hat{\lambda}^{-3} (-\Delta H - |\mathring{H}|^2 H)$$

for a surface \hat{x} in the conformal 3-sphere.

Most of the next theorem was known to Blaschke [1929] and Bryant [1984].

Theorem 2.5.2. *Suppose that $\hat{x} : M^2 \rightarrow \mathbb{S}^3$ is an immersed surface with no umbilical point. Then \hat{x} is a Willmore surface in \mathbb{S}^3 if and only if the conformal Gauss map induces a minimal spacelike surface in the de Sitter spacetime $\mathbb{S}^{1,3}$. Moreover its conformal transform \hat{x}^* is a dual Willmore surface in \mathbb{S}^3 .*

Proof. Because Lemma 2.3.2 implies that ξ is also the conformal Gauss map (up to the sign) for \hat{x}^* when H^ξ vanishes. The two dual Willmore surfaces are the two envelope surfaces of the family of round 2-spheres described by the conformal Gauss map ξ . □

Remark 2.5.3. It is also known to [Blaschke 1929] and [Bryant 1984] that if \hat{x} is a minimal surface in \mathbb{S}^3 , then $\hat{x}^* = -\hat{x}$; and that \hat{x} is a Willmore surface if and only if $\hat{x}^{**} = \hat{x}$. An interesting question then arises: is it possible to have $\hat{x}^{***} = \hat{x}$, and if so what would that equality imply for the surface?

2.6. Finding y_λ^* . Let us now solve y_λ^* for $y_\lambda = \hat{\lambda}(1, \hat{x}) = \hat{\lambda}y$, where $\hat{\lambda} = \lambda \circ \hat{x}$ and λ is a positive function on the sphere \mathbb{S}^3 . At each point on the surface we set

$$y_\lambda^* = \kappa y_\lambda + \kappa_\dagger y_\lambda^\dagger + b\vec{n}_\lambda + \frac{\omega_1^\lambda}{E_\lambda} (y_\lambda)_{u^1} + \frac{\omega_2^\lambda}{E_\lambda} (y_\lambda)_{u^2}.$$

We get from (2.4.1)

$$(2.6.1) \quad \left\{ \begin{array}{l} \kappa_{\dagger} = 1, \\ -2\kappa_{+}\kappa_{-} + b^2 + \frac{(\omega_1^\lambda)^2 + (\omega_2^\lambda)^2}{E_\lambda} = 0, \\ b = H_\lambda, \\ -(\Omega_\lambda)_{11}\omega_1^\lambda - (\Omega_\lambda)_{12}\omega_2^\lambda = (H_\lambda)_{u^1} E_\lambda, \\ -(\Omega_\lambda)_{21}\omega_1^\lambda - (\Omega_\lambda)_{22}\omega_2^\lambda = (H_\lambda)_{u^2} E_\lambda. \end{array} \right.$$

Lemma 2.6.1. *Suppose that $\hat{x} : M^2 \rightarrow S^3$ is an immersed surface with no umbilical point. Then*

$$(2.6.2) \quad y_\lambda^* = \frac{1}{2}(|\omega^\lambda|^2 + H_\lambda^2)y_\lambda + y_\lambda^\dagger + H_\lambda \vec{n}_\lambda - (\mathring{H})_\lambda^{-1} dH_\lambda$$

for any positive function λ on the 3-sphere, where

$$|\omega^\lambda|^2 = \frac{(\omega_1^\lambda)^2 + (\omega_2^\lambda)^2}{E_\lambda} = \frac{1}{m}((H_\lambda)_{u^1}^2 + (H_\lambda)_{u^2}^2).$$

In particular,

$$(2.6.3) \quad y^* = \frac{1}{2}(|\omega|^2 + H^2)y + \frac{1}{2}(1, -\hat{x}) + H(0, \mathbf{n}) - (0, (\mathring{H})^{-1}dH),$$

and

$$(2.6.4) \quad x^* = a\hat{x} + \frac{H}{1-a}\mathbf{n} - \frac{1}{1-a}(\mathring{H})^{-1}dH,$$

where

$$(2.6.5) \quad a = \frac{|\omega|^2 + H^2 - 1}{|\omega|^2 + H^2 + 1}.$$

Proof. One simply solves (2.6.1) if $\det \Omega_\lambda \neq 0$, which is equivalent to the fact that the surface has no umbilical point. □

2.7. Canonicity of y^* . We next show that the choice of y^* is canonical in terms of doing conformal geometry for the surface \hat{x} in S^3 . Two issues are involved; the first concerns the symmetry of the conformal 3-sphere. To be precise, for a conformal transformation

$$\phi : S^3 \rightarrow S^3$$

and the transformed surface

$$\phi(\hat{x}) : M^2 \rightarrow S^3,$$

is it true that

$$\tilde{\phi}(\tilde{x}) = \alpha\tilde{\phi}(y) + \alpha\rho\tilde{\phi}(y^*) : \mathbb{R}^+ \times \mathbb{R}^+ \times M^2 \rightarrow \mathbb{R}^{1,4}$$

is the associate 4-surface of $\phi(\hat{x})$ in $\mathbb{R}^{1,4}$, where $\tilde{\phi}$ is the corresponding Lorentz transformation on $\mathbb{R}^{1,4}$ to ϕ ? The other issue is whether or not the associate surface \tilde{x} is independent of metrics in the conformal class of the round 3-sphere. The first easy and important fact is that the conformal Gauss map is independent of the metrics in the conformal class.

Lemma 2.7.1. *Suppose that $\hat{x} : \mathbf{M}^2 \rightarrow \mathbb{S}^3$ is an immersed surface. Then the conformal Gauss map ξ is independent of the metrics in the conformal class of the round 3-sphere \mathbb{S}^3 . Meanwhile, the conformal Gauss map for the transformed surface $\phi(\hat{x})$ is exactly $\tilde{\phi}(\xi)$, where $\tilde{\phi}$ is the Lorentz transformation on the Minkowski spacetime $\mathbb{R}^{1,4}$ corresponding to a conformal transformation ϕ on \mathbb{S}^3 .*

Proof. First of all, one needs to realize that, for any given metric in the conformal class of the round 3-sphere, it simply amounts to consider the surface

$$y_\lambda = \hat{\lambda}(1, \hat{x}) : \mathbf{M}^2 \rightarrow \mathbb{N}_+^4$$

for some positive function $\lambda : \mathbb{S}^3 \rightarrow \mathbb{R}^+$ and $\hat{\lambda} = \lambda \circ \hat{x}$. But this might only alter the parametrization of the homogeneous null 3-surface $x^{\mathbb{N}} = \alpha \hat{\lambda}(1, \hat{x}) : \mathbb{R}^+ \times \mathbf{M}^2 \rightarrow \mathbb{N}_+^4$. Hence it will not alter the conformal Gauss map. Of course we have already seen this from [Lemma 2.3.4](#).

Next we consider the transformed surface $\phi(\hat{x})$. Recall that, given a conformal transformation ϕ of 3-sphere, we have a unique Lorentz transformation $\tilde{\phi}$ in the time and orientation preserving component of the Lorentz group on the Minkowski spacetime such that, for $\lambda(1, \hat{x}) \in \mathbb{R}^{1,4}$,

$$(2.7.1) \quad \tilde{\phi}(\lambda(1, \hat{x})) = \lambda \mu(1, \phi(\hat{x}))$$

for some positive number μ . By the definition, which requires that $\tilde{\phi}$ be a linear map and that

$$\langle \tilde{\phi}((t, \hat{x})), \tilde{\phi}((s, \hat{y})) \rangle = \langle (t, \hat{x}), (s, \hat{y}) \rangle,$$

we now easily see that $\tilde{\phi}(\xi)$ is the conformal Gauss map for the transformed surface $\phi(\hat{x})$. Since $\tilde{\phi}(\xi)$ is the unit normal vector field to the homogeneous null 3-surface $\tilde{\phi}(x)$ in \mathbb{N}_+^4 that is conformal map from \mathbf{M}^2 to $\mathbb{S}^{1,3}$. \square

Consequently:

Proposition 2.7.2. *Suppose that $\hat{x} : \mathbf{M}^2 \rightarrow \mathbb{S}^3$ is an immersed surface with no umbilical point. Then the associate surface*

$$\tilde{x} = \alpha y_\lambda + \alpha \rho y_\lambda^* : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbf{M}^2 \rightarrow \mathbb{R}^{1,4},$$

for any $y_\lambda = \hat{\lambda}(1, \hat{x})$ and $y^* = \hat{\lambda}^{-1} \lambda^*(1, \hat{x}^*)$ defined by the equations [\(2.4.1\)](#), is independent of the metrics in conformal class of the round 3-sphere \mathbb{S}^3 .

Proof. It suffices to verify that

$$(2.7.2) \quad (\hat{\lambda}y)^* = \hat{\lambda}^{-1}y^*.$$

Since it implies that the change of metrics in the conformal class will at most cause a possible change of parametrization of the associate surface \tilde{x} . \square

Lemma 2.7.3. *Suppose that $\hat{x} : M^2 \rightarrow S^3$ is an immersed surface with no umbilical point. Let $y_\lambda = \hat{\lambda}(1, \hat{x}) \in \mathbb{N}_+^4$ and let ϕ be a conformal transformation of 3-sphere. Then*

$$(2.7.3) \quad \tilde{\phi}(y_\lambda)^* = \tilde{\phi}(y_\lambda^*).$$

Hence

$$(2.7.4) \quad \phi(\hat{x}^*) = (\phi(\hat{x}))^*.$$

Proof. From Lemma 2.7.1 we know that the conformal Gauss map for the transformed surface $\phi(\hat{x})$ is $\tilde{\phi}(\xi)$. Then it is easy to verify (2.4.1) for $\tilde{\phi}(y^*)$ to be $\tilde{\phi}(y)^*$. Then the Equation (2.7.4) follows from (2.7.1) and (2.7.3):

$$\hat{\gamma}^*(1, (\phi(\hat{x}))^*) = \tilde{\phi}(y)^* = \tilde{\phi}(y^*) = \hat{\mu}^*\hat{\lambda}^*(1, \phi(\hat{x}^*)). \quad \square$$

Therefore:

Proposition 2.7.4. *Suppose that $\hat{x} : M^2 \rightarrow S^3$ is an immersed surface with no umbilical point. Let ϕ be a conformal transformation of 3-sphere. Then the associate 4-surface in $\mathbb{R}^{1,4}$ of the transformed surface $\phi(\hat{x})$ is exactly the 4-surface $\tilde{\phi}(\tilde{x})$ transformed from the associate 4-surface \tilde{x} of the original surface \hat{x} under the corresponding Lorentz transformation $\tilde{\phi}$ of ϕ .*

3. The geometry of the associate surfaces

In this section we calculate the first and second fundamental forms for the associate homogeneous timelike 4-surfaces \tilde{x} in $\mathbb{R}^{1,4}$ as well as for the associate ruled surface x^+ in the hyperboloid \mathbb{H}^4 , for a given immersed 2-surface \hat{x} in S^3 .

3.1. The first fundamental form for \tilde{x} in $\mathbb{R}^{1,4}$. To calculate the first fundamental form for the surface in the parametrization

$$(3.1.1) \quad \tilde{x} = \alpha y_\lambda + \alpha \rho y_\lambda^*$$

associated with a conformal metric $\lambda^2 g_0$ on the 3-sphere S^3 , we first calculate

$$d\tilde{x} = (y_\lambda + \rho y_\lambda^*)d\alpha + \alpha y_\lambda^*d\rho + (\alpha(y_\lambda)_{u^1} + \alpha\rho(y_\lambda^*)_{u^1})du^1 + (\alpha(y_\lambda)_{u^2} + \alpha\rho(y_\lambda^*)_{u^2})du^2.$$

Hence the first fundamental form for the associate 4-surface \tilde{x} in the coordinates (α, ρ, u^1, u^2) is

$$\begin{aligned} I^{\tilde{x}} = \langle d\tilde{x}, d\tilde{x} \rangle &= -2\rho d\alpha d\alpha - 2\alpha d\alpha d\rho \\ &+ 2\alpha^2 \langle (y_\lambda^*, (y_\lambda)_{u^1}) \rangle d\rho du^1 + 2\alpha^2 \langle y_\lambda^*, (y_\lambda)_{u^2} \rangle d\rho du^2 \\ &+ \langle \alpha (y_\lambda)_{u^1} + \alpha\rho (y_\lambda^*)_{u^1}, \alpha (y_\lambda)_{u^1} + \alpha\rho (y_\lambda^*)_{u^1} \rangle (du^1)^2 \\ &+ \langle \alpha (y_\lambda)_{u^2} + \alpha\rho (y_\lambda^*)_{u^2}, \alpha (y_\lambda)_{u^2} + \alpha\rho (y_\lambda^*)_{u^2} \rangle (du^2)^2 \\ &+ 2 \langle \alpha (y_\lambda)_{u^1} + \alpha\rho (y_\lambda^*)_{u^1}, \alpha (y_\lambda)_{u^2} + \alpha\rho (y_\lambda^*)_{u^2} \rangle du^1 du^2. \end{aligned}$$

In fact one may calculate

$$\begin{aligned} (3.1.2) \quad (y_\lambda)_{u^1} &= -\omega_1^\lambda y_\lambda - \frac{(\Omega_\lambda)_{11}}{m} \xi_{u^1} - \frac{(\Omega_\lambda)_{12}}{m} \xi_{u^2}, \\ (y_\lambda)_{u^2} &= -\omega_2^\lambda y_\lambda - \frac{(\Omega_\lambda)_{21}}{m} \xi_{u^1} - \frac{(\Omega_\lambda)_{22}}{m} \xi_{u^2}, \\ (y_\lambda^*)_{u^1} &= \omega_1^\lambda y_\lambda^* - \frac{(\Omega_\lambda^*)_{11}}{m} \xi_{u^1} - \frac{(\Omega_\lambda^*)_{12}}{m} \xi_{u^2}, \\ (y_\lambda^*)_{u^2} &= \omega_2^\lambda y_\lambda^* - \frac{(\Omega_\lambda^*)_{21}}{m} \xi_{u^1} - \frac{(\Omega_\lambda^*)_{22}}{m} \xi_{u^2}, \end{aligned}$$

where

$$(3.1.3) \quad \omega^\lambda = \langle dy_\lambda, y_\lambda^* \rangle = -I_\lambda (\Omega_\lambda^{-1} dH_\lambda)$$

based on (2.6.1). Now let us write $I^{\tilde{x}}$ in matrix form:

$$(3.1.4) \quad I_{\tilde{x}} = \begin{bmatrix} -2\rho & -\alpha & 0 & 0 \\ -\alpha & 0 & \alpha^2 \omega_1^\lambda & \alpha^2 \omega_2^\lambda \\ 0 & \alpha^2 \omega_1^\lambda & & \\ 0 & \alpha^2 \omega_2^\lambda & & \alpha^2 F \end{bmatrix}$$

where

$$(3.1.5) \quad \begin{cases} F_{11} = \frac{1}{m} (p^2 + q^2) + 2\rho (\omega_1^\lambda)^2, \\ F_{12} = F_{21} = \frac{1}{m} q (p + r) + 2\rho \omega_1^\lambda \omega_2^\lambda, \\ F_{22} = \frac{1}{m} (q^2 + r^2) + 2\rho (\omega_2^\lambda)^2, \end{cases}$$

and

$$\begin{bmatrix} p & q \\ q & r \end{bmatrix} = \Omega_\lambda + \rho \Omega_\lambda^*.$$

It can be calculated that

$$(3.1.6) \quad \begin{aligned} \det I_{\tilde{x}} &= -\frac{\alpha^6}{m^2}(pr - q^2)^2 \\ &= -\frac{\alpha^6}{4m^2}(E_\lambda^2|\Omega_\lambda + \rho\Omega_\lambda^*|^2 - \rho^2((\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22})^2) \end{aligned}$$

which can tell us where the associate surface \tilde{x} is degenerate. It is maybe a little surprising that it is actually not difficult to calculate the inverse of $I_{\tilde{x}}$. We present the calculations in [Appendix A](#) since they are straightforward calculations.

3.2. The second fundamental form for \tilde{x} in $\mathbb{R}^{1,4}$. It is clear from the definition that the conformal Gauss map ξ is the unit normal vector for the associate 4-surface \tilde{x} in $\mathbb{R}^{1,4}$. Hence the second fundamental form for \tilde{x} in $\mathbb{R}^{1,4}$ is

$$(3.2.1) \quad II_{\tilde{x}} = -\langle d\tilde{x}, d\xi \rangle = (\alpha(\Omega_\lambda)_{ij} + \alpha\rho(\Omega_\lambda^*)_{ij})du^i du^j,$$

or in matrix form

$$II_{\tilde{x}} = \begin{bmatrix} 0 & 0 \\ 0 & \alpha\Omega_\lambda + \alpha\rho\Omega_\lambda^* \end{bmatrix}.$$

Therefore the mean curvature for the associate 4-surface in $\mathbb{R}^{1,4}$ is

$$H^{\tilde{x}} = \text{Tr}(I_{\tilde{x}})^{-1}II_{\tilde{x}}.$$

To calculate the mean curvature $H^{\tilde{x}}$ one only needs to know the low-right 2×2 block in the inverse of the matrix $I_{\tilde{x}}$. According to the calculations in [Appendix A](#), particularly [\(A.3\)](#), [\(A.9\)](#) and [\(A.10\)](#), we therefore have

$$(3.2.2) \quad \begin{aligned} H^{\tilde{x}} &= \frac{m}{\alpha(pr - q^2)^2}((q^2 + r^2)p - 2q^2(p + r) + (p^2 + q^2)r) \\ &= \frac{m(p + r)}{\alpha(pr - q^2)}, \end{aligned}$$

where

$$pr - q^2 = \det \Omega_\lambda - \rho \text{Tr} \Omega_\lambda \Omega_\lambda^* + \rho^2 \det \Omega_\lambda^*$$

and

$$(3.2.3) \quad p + r = \rho((\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22}) = -\rho E_\lambda \mathcal{H}_\lambda$$

in the light of [\(2.5.14\)](#).

Theorem 3.2.1. *Suppose that $\hat{x} : \tilde{\mathbb{M}}^2 \rightarrow \mathbb{S}^3$ is an immersed surface with no umbilical point. Then \hat{x} is a Willmore surface in \mathbb{S}^3 if and only if the associate 4-surface \tilde{x} in $\mathbb{R}^{1,4}$ is minimal.*

Proof. Based on the above equations (3.2.3) and (3.2.2) we obtain

$$H^{\tilde{x}} = \frac{\rho \det \Omega_\lambda \mathcal{H}_\lambda}{\alpha(\det \Omega_\lambda - \rho \operatorname{Tr} \Omega_\lambda \Omega_\lambda^* + \rho^2 \det \Omega_\lambda^*)}. \quad \square$$

3.3. Local fundamental theorem for surfaces in conformal 3-sphere. In this subsection we state and prove a local fundamental theorem for surfaces in conformal 3-sphere. In the previous section we have introduced the associate surface \tilde{x} in Minkowski spacetime $\mathbb{R}^{1,4}$ from a given surface \hat{x} in \mathbb{S}^3 . From the geometric structure of the associate surface \tilde{x} one can tell that its intersection with the positive light cone \mathbb{N}_+^4 is a homogeneous null 3-surface whose projectivization will recover the original surface \hat{x} in \mathbb{S}^3 .

Given a surface \hat{x} in \mathbb{S}^3 with a isothermal coordinates (u^1, u^2) on the parameter space M^2 , we have the first fundamental form I in matrix form

$$I = \begin{bmatrix} E & 0 \\ 0 & E \end{bmatrix}$$

and the second fundamental form II in matrix form

$$II = \begin{bmatrix} e & f \\ f & g \end{bmatrix}$$

The local fundamental theorem for surfaces in Riemannian geometry states that, up to isometries of the standard round sphere \mathbb{S}^3 , locally the surface is uniquely determined by the first fundamental form I and the second fundamental form II in the standard round sphere \mathbb{S}^3 . Conversely, given a positive definite symmetric 2-form I and a symmetric 2-form II in the parameter domain, which satisfy some integrability conditions (Gauss-Codazzi equations), up to isometries, there is locally a unique surface \hat{x} in the standard round sphere \mathbb{S}^3 whose first and second fundamental forms are I and II . We are looking for the analogous local fundamental theorem for surfaces in conformal round 3-sphere \mathbb{S}^3 . The core idea of the local fundamental theorem in Riemannian geometry is to solve the structure equations, which are the equations of motion of Frenet frames on the surface and are determined from I and II .

Our strategy here is to use the local fundamental theorem for the associate surface \tilde{x} in the Minkowski spacetime $\mathbb{R}^{1,4}$ to establish the local fundamental theorem for a surface \hat{x} in the conformal sphere \mathbb{S}^3 . Since the association introduced in previous subsections requires that the surface \hat{x} has no umbilical point, we will always assume here that surfaces \hat{x} have no umbilical point.

To summarize the previous discussions, given a surface \hat{x} in \mathbb{S}^3 , we have $I = E|du|^2$ and $II = e(du^1)^2 + 2fdu^1du^2 + g(du^2)^2$. We also have the so-called Möbius metric $I^\xi = m|du|^2 = \frac{1}{2}E|\hat{I}|^2|du|^2$ induced from the Conformal Gauss

map ξ of the surface \hat{x} , where

$$\mathring{H} = \begin{bmatrix} \frac{e-g}{2} & f \\ f & \frac{g-e}{2} \end{bmatrix}$$

is the traceless part of the second fundamental form II . We then construct the associate surface

$$\tilde{x} = \alpha y_\lambda + \alpha \rho y_\lambda^* : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{M}^2 : \mathbb{R}^{1,4}.$$

The first fundamental form $I^{\tilde{x}}$ for \tilde{x} in matrix form is, from (3.1.4),

$$\begin{bmatrix} -2\rho & -\alpha & 0 & 0 \\ -\alpha & 0 & \alpha^2 \omega_1^\lambda & \alpha^2 \omega_2^\lambda \\ 0 & \alpha^2 \omega_1^\lambda & \frac{\alpha^2}{m}(p^2+q^2)+2\alpha^2 \rho(\omega_1^\lambda)^2 & \frac{\alpha^2}{m}q(p+r)+2\alpha^2 \rho \omega_1^\lambda \omega_2^\lambda \\ 0 & \alpha^2 \omega_2^\lambda & \frac{\alpha^2}{m}q(p+r)+2\alpha^2 \rho \omega_1^\lambda \omega_2^\lambda & \frac{\alpha^2}{m}(q^2+r^2)+2\alpha^2 \rho(\omega_2^\lambda)^2 \end{bmatrix},$$

where

$$\omega^\lambda = \omega_1^\lambda du^1 + \omega_2^\lambda du^2 = -d \log \hat{\lambda} - I(\Omega^{-1}(dH)) = d \log \hat{\lambda} + \omega.$$

The second fundamental form $II^{\tilde{x}}$ for \tilde{x} in $\mathbb{R}^{1,4}$ in matrix form is, from (3.2.1),

$$\begin{bmatrix} 0 & 0 \\ 0 & \alpha \Omega_\lambda + \alpha \rho \Omega_\lambda^* \end{bmatrix},$$

where $\Omega_\lambda = \hat{\lambda} \Omega$ and $\Omega_\lambda^* = \hat{\lambda}^{-1} \Omega^*$. Notice that $I^{\tilde{x}}$ and $II^{\tilde{x}}$ are exactly determined by the Möbius metric $I^\xi = m|du|^2$, the 1-form ω , the traceless symmetric 2-tensor Ω and the symmetric 2-tensor Ω^* , plus the conformal factor $\hat{\lambda}$.

Next we write the equations for the motion of the Frenet frames on the associate surface \tilde{x} according to $I^{\tilde{x}}$ and $II^{\tilde{x}}$. We consider the Frenet frame

$$\left\{ y_\lambda, y_\lambda^*, \frac{1}{\sqrt{m}} \xi_{u^1}, \frac{1}{\sqrt{m}} \xi_{u^2}, \xi \right\}$$

on the associate surface \tilde{x} , which are the orthonormal frames on \tilde{x} with respect to the Minkowski metric \tilde{G}_0 on $\mathbb{R}^{1,4}$. We now write

(3.3.1)

$$\frac{\partial}{\partial u^1} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix} = \begin{bmatrix} -\omega_1^\lambda & 0 & -\frac{1}{\sqrt{m}}(\Omega_\lambda)_{11} & -\frac{1}{\sqrt{m}}(\Omega_\lambda)_{12} & 0 \\ 0 & \omega_1^\lambda & -\frac{1}{m}(\Omega_\lambda^*)_{11} & -\frac{1}{m}(\Omega_\lambda^*)_{12} & 0 \\ \frac{1}{\sqrt{m}}(\Omega_\lambda)_{11} & \frac{1}{\sqrt{m}}(\Omega_\lambda^*)_{11} & 0 & -\frac{1}{2m}m_{u^2} & -\sqrt{m} \\ \frac{1}{\sqrt{m}}(\Omega_\lambda)_{21} & \frac{1}{\sqrt{m}}(\Omega_\lambda^*)_{21} & \frac{1}{2m}m_{u^2} & 0 & 0 \\ 0 & 0 & \sqrt{m} & 0 & 0 \end{bmatrix} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix}$$

and

(3.3.2)

$$\frac{\partial}{\partial u^2} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix} = \begin{bmatrix} -\omega_2^\lambda & 0 & -\frac{1}{\sqrt{m}} (\Omega_\lambda)_{21} & -\frac{1}{\sqrt{m}} (\Omega_\lambda)_{22} & 0 \\ 0 & \omega_2^\lambda & -\frac{1}{m} (\Omega_\lambda^*)_{21} & -\frac{1}{m} (\Omega_\lambda^*)_{22} & 0 \\ \frac{1}{\sqrt{m}} (\Omega_\lambda)_{21} & \frac{1}{\sqrt{m}} (\Omega_\lambda^*)_{21} & 0 & -\frac{1}{2m} m_{u^1} & 0 \\ \frac{1}{\sqrt{m}} (\Omega_\lambda)_{22} & \frac{1}{\sqrt{m}} (\Omega_\lambda^*)_{22} & \frac{1}{2m} m_{u^1} & 0 & -\sqrt{m} \\ 0 & 0 & 0 & \sqrt{m} & 0 \end{bmatrix} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix}.$$

Remember we also have the two trivial equations

$$\frac{\partial}{\partial \alpha} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix} = 0 \quad \text{and} \quad \frac{\partial}{\partial \rho} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix} = 0.$$

To solve the ODE systems (3.3.1) and (3.3.2), the necessary integrable condition is

$$(3.3.3) \quad \frac{\partial}{\partial u^1} \frac{\partial}{\partial u^2} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix} = \frac{\partial}{\partial u^2} \frac{\partial}{\partial u^1} \begin{bmatrix} y_\lambda \\ y_\lambda^* \\ \frac{1}{\sqrt{m}} \xi_{u^1} \\ \frac{1}{\sqrt{m}} \xi_{u^2} \\ \xi \end{bmatrix}.$$

It turns out that (3.3.3) is equivalent to the following six equations on the variables: the positive function m , the 1-form ω^λ , the traceless symmetric matrix Ω_λ and the symmetric matrix Ω_λ^* ,

$$(3.3.4) \quad \begin{cases} (\Omega_\lambda)_{11,2} - (\Omega_\lambda)_{12,1} = \omega_1^\lambda (\Omega_\lambda)_{12} - \omega_2^\lambda (\Omega_\lambda)_{11}, \\ (\Omega_\lambda)_{12,2} - (\Omega_\lambda)_{22,1} = \omega_1^\lambda (\Omega_\lambda)_{22} - \omega_2^\lambda (\Omega_\lambda)_{12}, \end{cases}$$

(3.3.5)

$$\begin{cases} (\Omega_\lambda^*)_{11,2} - (\Omega_\lambda^*)_{12,1} = -\omega_1^\lambda (\Omega_\lambda^*)_{12} + \omega_2^\lambda (\Omega_\lambda^*)_{11} + \frac{1}{2} \frac{(\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22}}{|\Omega_\lambda|^2} (|\Omega_\lambda|^2)_{u^2}, \\ (\Omega_\lambda^*)_{12,2} - (\Omega_\lambda^*)_{22,1} = -\omega_1^\lambda (\Omega_\lambda^*)_{22} + \omega_2^\lambda (\Omega_\lambda^*)_{12} + \frac{1}{2} \frac{(\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22}}{|\Omega_\lambda|^2} (|\Omega_\lambda|^2)_{u^2}, \end{cases}$$

$$(3.3.6) \quad \omega_{1,2}^\lambda - \omega_{2,1}^\lambda = \frac{1}{m} ((\Omega_\lambda)_{11} - (\Omega_\lambda)_{22}) (\Omega_\lambda^*)_{12} - ((\Omega_\lambda^*)_{11} - (\Omega_\lambda^*)_{22}) (\Omega_\lambda)_{12},$$

and

$$(3.3.7) \quad (\mathcal{K} - 1) = \frac{1}{m^2} \text{Tr } \Omega_\lambda \Omega_\lambda^*,$$

where \mathcal{K} is the Gaussian curvature of the Möbius metric $I^\xi = m|du|^2$. Of course, as one may verify, (3.3.4), (3.3.5), (3.3.6) and (3.3.7) are exactly the Gauss-Codazzi equations for the surface ξ in the de Sitter spacetime $\mathbb{S}^{1,3}$ induced by the conformal Gauss map ξ of the surface \hat{x} in conformal 3-sphere \mathbb{S}^3 .

Now we are ready to state and prove the local fundamental theorem for surfaces in conformal round 3-sphere \mathbb{S}^3 .

Theorem 3.3.1. *Suppose that, on a domain in $D \subset \mathbb{R}^2$, we are given*

- a traceless symmetric 2-form Ω ,
- a positive function m or equivalently E such that $m = \frac{-\det \Omega}{E}$,
- a 1-form ω ,
- a symmetric 2-form Ω^* ,

and that they satisfy the integrability conditions (3.3.4)–(3.3.7). Then, for a given point p_0 in D , there exists an open neighborhood D_0 of p_0 in D , a parametrized surface $\hat{x} : D_0 \rightarrow \mathbb{S}^3$ with no umbilical point, and a positive function $\hat{\lambda} : D_0 \rightarrow \mathbb{R}^+$ with $\hat{\lambda}(p_0) = 1$, such that

- $\Omega = \hat{\lambda} \hat{I}$, where \hat{I} is the traceless part of the second fundamental form of \hat{x} in the standard round \mathbb{S}^3 ,
- $m|du|^2 = \langle d\xi, d\xi \rangle$ is the Möbius metric induced by the conformal Gauss map ξ of \hat{x} ,
- $\omega = -I((\hat{I})^{-1}(dH)) - d \log \hat{\lambda}$, where I is the first fundamental form and H is the mean curvature of \hat{x} in the standard round \mathbb{S}^3 ,
- $\Omega^* = -\hat{\lambda}^{-1} \langle d\xi, dy^* \rangle$, where $y^* = \frac{1}{1-\hat{x} \cdot \hat{x}^*}(1, \hat{x}^*)$ and \hat{x}^* is the conformal transform of \hat{x} .

The surface \hat{x} is unique up to a conformal transformation of \mathbb{S}^3 .

Proof. We start by choosing starting values for $y, y^*, \xi_{u^1}, \xi_{u^2}, \xi$ at $p_0 = (u_0^1, u_0^2)$. First we take a null vector

$$y(u_0^1, u_0^2) = y_0 = (1, \hat{x}_0)$$

for some $\hat{x}_0 \in \mathbb{S}^3 \subset \mathbb{R}^4$. Then we choose $\xi(u_0^1, u_0^2) = \xi_0 \in \mathbb{R}^{1,4}$ such that

$$(3.3.8) \quad \langle y_0, \xi_0 \rangle = 0 \text{ and } \langle \xi_0, \xi_0 \rangle = 1.$$

Next we choose $\xi_{u^1}(u_0^1, u_0^2) = \xi_0^1 \in \mathbb{R}^{1,4}$ and $\xi_{u^2}(u_0^1, u_0^2) = \xi_0^2 \in \mathbb{R}^{1,4}$ such that

$$(3.3.9) \quad \begin{aligned} \langle \xi_0^1, \xi_0^1 \rangle &= \langle \xi_0^2, \xi_0^2 \rangle = m(u_0^1, u_0^2), \\ \langle \xi_0^1, \xi_0^2 \rangle &= \langle \xi_0, \xi_0^1 \rangle = \langle \xi_0, \xi_0^2 \rangle = \langle y_0, \xi_0^1 \rangle = \langle y_0, \xi_0^2 \rangle = 0. \end{aligned}$$

Finally choose the unique null vector $y^*(u_0^1, u_0^2) = y_0^*$ such that

$$(3.3.10) \quad \begin{aligned} \langle y_0^*, y_0 \rangle &= -1, \\ \langle y_0^*, y_0^* \rangle &= \langle y_0^*, \xi_0 \rangle = \langle y_0^*, \xi_0^1 \rangle = \langle y_0^*, \xi_0^2 \rangle = 0. \end{aligned}$$

Notice that for any other choice of $\{y_1, y_1^*, \xi_1^1, \xi_1^2, \xi_1\}$ satisfying the same orthonormal properties in (3.3.8)–(3.3.10), there is a Lorentz transformation that takes one to the other. With the integrability conditions assumed we may solve the systems (3.3.1) and (3.3.2) at least in an open neighborhood D_0 of p_0 in D . Using the uniqueness of solutions to systems of linear ODE one sees that the solution $\{y, y^*, \frac{1}{\sqrt{m}}\xi_{u^1}, \frac{1}{\sqrt{m}}\xi_{u^2}, \xi\}$ remains orthonormal in the Minkowski metric in D_0 .

Now one should realize that the $y = \hat{\lambda}(1, \hat{x})$ here is with some positive $\hat{\lambda}$ (not necessarily identically 1 in D_0). It is then clear from all previous calculations that the rest of the statements in the theorem can be easily verified. \square

3.4. The geometry of the associate ruled surface x^+ in hyperbolic space \mathbb{H}^4 . In this section we want to discuss the geometry of the associate ruled 3-surface x^+ in \mathbb{H}^4 , which is associated with a given surface \hat{x} in the conformal 3-sphere. It's relation to the associate surface \tilde{x} is very much analogous to the one between the ambient spacetime and the Poincaré-Einstein manifold of a given conformal manifold in the work of Fefferman and Graham. It is evidently useful to understand the geometry of the associate ruled 3-surface x^+ in \mathbb{H}^4 .

It is rather easy now to do calculations for x^+ after we have calculated the first fundamental form for the associate 4-surface \tilde{x} in Minkowski spacetime $\mathbb{R}^{1,4}$ in Section 3.1. We first have

$$dx^+ = \frac{1}{\sqrt{2}}(e^t y_\lambda - e^{-t} y_\lambda^*) dt + (e^t (y_\lambda)_{u^1} + e^{-t} (y_\lambda^*)_{u^1}) du^1 + (e^t (y_\lambda)_{u^2} + e^{-t} (y_\lambda^*)_{u^2}) du^2$$

and, using (3.1.2),

$$I^{x^+} = (dt)^2 - 2\omega_i^\lambda dt du^i + \left(\frac{e^{2t}}{2m} ((\Omega_\lambda)_{i1} (\Omega_\lambda)_{j1} + (\Omega_\lambda)_{i2} (\Omega_\lambda)_{j2}) \right. \\ \left. + (\omega_i \omega_j + \frac{1}{m} ((\Omega_\lambda)_{i1} (\Omega_\lambda^*)_{j1} + (\Omega_\lambda)_{i2} (\Omega_\lambda^*)_{j2})) \right. \\ \left. + \frac{e^{-2t}}{2m} ((\Omega_\lambda^*)_{i1} (\Omega_\lambda^*)_{j1} + (\Omega_\lambda^*)_{i2} (\Omega_\lambda^*)_{j2}) \right) du^i du^j.$$

One can calculate the determinant

$$(3.4.1) \quad \det I^{x^+} = \frac{1}{8m^2} (E_\lambda^2 |e^t \Omega_\lambda + e^{-t} \Omega_\lambda^*|^2 - e^{-2t} ((\Omega_\lambda^*)_{11} + (\Omega_\lambda^*)_{22})^2),$$

which can tell us where the associate ruled surface x^+ is degenerate.

To obtain the second fundamental form of the surface x^+ it suffices to see that the conformal Gauss map ξ is still the unit normal vector to the surface x^+ in the

hyperboloid \mathbb{H}^4 . Hence

$$(3.4.2) \quad II^{x^+} = -\langle dx^+, d\xi \rangle = \frac{1}{\sqrt{2}}(e^t \Omega_\lambda + e^{-t} \Omega_\lambda^*).$$

By the similar calculations as that in the previous section we have the mean curvature of the associate ruled surface x^+ as follows:

$$(3.4.3) \quad H^{x^+} = e^{-3t} \frac{\sqrt{2} \det \Omega_\lambda \mathcal{H}_\lambda}{(\det \Omega_\lambda - e^{-2t} \operatorname{Tr} \Omega_\lambda \Omega_\lambda^* + e^{-4t} \det \Omega_\lambda^*)}.$$

Theorem 3.4.1. *Suppose that \hat{x} is an immersed surface in the conformal sphere \mathbb{S}^3 with no umbilical point and that x^+ is the associate ruled surface in the hyperboloid \mathbb{H}^4 . Then \hat{x} is a Willmore surface in the conformal sphere if and only if the associate ruled 3-surface x^+ in the hyperboloid is a minimal surface.*

4. Scalar invariants of surfaces in conformal round 3-sphere

In this section we want to introduce scalar local invariants for surfaces in conformal round 3-sphere \mathbb{S}^3 . We will first recall what are scalar invariants for hypersurfaces in (pseudo-)Riemannian geometry. Inspired by the work of Fefferman and Graham on scalar local invariants in conformal geometry we are going to use the associate surface \tilde{x} in the Minkowski $\mathbb{R}^{1,4}$ of a given surface \hat{x} in the 3-sphere \mathbb{S}^3 , where one considers the standard conformal 3-sphere as the projectivized positive light cone of the Minkowski spacetime to construct scalar local invariant.

4.1. Scalar invariants of 4-surfaces in $\mathbb{R}^{1,4}$. For our purpose we will focus on the discussion of scalar (pseudo-)Riemannian invariants of 4-surfaces \tilde{x} in the Minkowski spacetime $\mathbb{R}^{1,4}$. Suppose that

$$\phi = \phi(v^2, v^3, v^4, v^5) : A \subset \mathbb{R}^4 \rightarrow \mathbb{R}^{1,4}$$

is a local parametrization of a surface \tilde{x} , where A is a domain in \mathbb{R}^4 . Hence it induces a local coordinate

$$\tilde{\phi} = \tilde{\phi}(v^1, v^2, v^3, v^4, v^5) : B \subset (-\epsilon, \epsilon) \times A \rightarrow \mathbb{R}^{1,4}$$

for $\mathbb{R}^{1,4}$ such that

$$\phi(v^2, v^3, v^4, v^5) = \tilde{\phi}(0, v^2, v^3, v^4, v^5).$$

We will use the Capital Latin letters to stand for indices from 1 to 5 and Latin letters to stand for the indices from 2 to 5. We will use $v = (v^1, v^2, \dots, v^5)$ and $\hat{v} = (v^2, \dots, v^5)$. Hence the Minkowski metric in this coordinate is given as

$$\tilde{G}_0 = \langle d\tilde{\phi}, d\tilde{\phi} \rangle = (\tilde{G}_0)_{IJ} dv^I dv^J$$

and the first fundamental form for \tilde{x} in $\mathbb{R}^{1,4}$ is given as

$$I^{\tilde{x}} = \langle d\phi, d\phi \rangle = \tilde{g}_{ij} dv^i dv^j = (\tilde{\mathcal{G}}_0)_{ij}|_{v^1=0} dv^i dv^j.$$

To be more restrictive we will assume that the surface \tilde{x} is timelike and let

$$\xi : B \rightarrow \mathbb{S}^{1,3}$$

be a unit normal vector field on \tilde{x} in $\mathbb{R}^{1,4}$. Then the second fundamental form for \tilde{x} is given as

$$II^{\tilde{x}} = -\langle d\phi, d\xi \rangle = \tilde{h}_{ij} dv^i dv^j,$$

and we have

$$\xi_{v^i} = -\tilde{h}_{ik} \tilde{g}^{kj} \phi_{v^j}.$$

Definition 4.1.1. Let $i : M^{n-1} \rightarrow N^n$ be an immersed hypersurface and let g be a (pseudo)-Riemannian metric on the ambient manifold N^n . A scalar (pseudo)-Riemannian invariant $I(i, N^n, g)$ for the hypersurface i in N^n at a point p_0 on the surface i is a polynomial in the variables that are the coordinate partial derivatives of g_{IJ} of any order and the reciprocal of the determinant of g_{IJ} at the point p_0 such that the value of $I(i, N^n, g)$ at p_0 is independent of choices of local coordinates $\tilde{\phi}$ of N^n which are induced from a parametrization ϕ of the surface i nearby the given point p_0 .

The well-known examples of scalar Riemannian invariants for \tilde{x} in $\mathbb{R}^{1,4}$ are

- $\tilde{H} = \tilde{g}^{ij} \tilde{h}_{ij}$,
- $|\tilde{h}|^2 = \tilde{g}^{ik} \tilde{g}^{jl} \tilde{h}_{ij} \tilde{h}_{kl}$ and $\tilde{H}^2 = \tilde{g}^{ij} \tilde{g}^{kl} \tilde{h}_{ij} \tilde{h}_{kl}$,
- $\tilde{\Delta} \tilde{H} = \tilde{g}^{kl} \tilde{g}^{ij} \tilde{h}_{ij,kl}$, $\text{Div Div } \tilde{h} = \tilde{g}^{ik} \tilde{g}^{jl} \tilde{h}_{ij,kl}$, $\tilde{H} |\tilde{h}|^2 = \tilde{g}^{ik} \tilde{g}^{jl} \tilde{g}^{mn} \tilde{h}_{ij} \tilde{h}_{kl} \tilde{h}_{mn}$, $\text{Tr}_{\tilde{g}} \tilde{h}^3 = \tilde{g}^{in} \tilde{g}^{jk} \tilde{g}^{km} \tilde{h}_{ij} \tilde{h}_{kl} \tilde{h}_{mn}$, and $\tilde{H}^3 = \tilde{g}^{ij} \tilde{g}^{kl} \tilde{g}^{mn} \tilde{h}_{ij} \tilde{h}_{kl} \tilde{h}_{mn}$,
- $|\tilde{\nabla} \tilde{h}|^2 = \tilde{g}^{ip} \tilde{g}^{jq} \tilde{g}^{kr} \tilde{h}_{ij,k} \tilde{h}_{pq,r}$, $\tilde{g}^{ip} \tilde{g}^{jr} \tilde{g}^{kq} \tilde{h}_{ij,k} \tilde{h}_{pq,r}$, $\tilde{g}^{ip} \tilde{g}^{jr} \tilde{g}^{kq} \tilde{h}_{ij,k} \tilde{h}_{pq,r}$, $|\tilde{\nabla} \tilde{H}|^2 = \tilde{g}^{ij} \tilde{g}^{pq} \tilde{g}^{kr} \tilde{h}_{ij,k} \tilde{h}_{pq,r}$, $|\tilde{\text{Div}} \tilde{h}|^2 = \tilde{g}^{ip} \tilde{g}^{jk} \tilde{g}^{qr} \tilde{h}_{ij,k} \tilde{h}_{pq,r}$, $\tilde{\text{Div}} \tilde{h} \cdot d\tilde{H}$,
- $\tilde{\Delta} \tilde{\Delta} \tilde{H}$

Each scalar invariant has an order. To find the order of each scalar invariant one simply scales the metric by a constant κ and see what is the dimension of the scalar invariant. For example, we can easily find that

$$\begin{aligned} \tilde{H}[\kappa^2 \tilde{\mathcal{G}}_0] &= \kappa^{-1} \tilde{H}[\tilde{\mathcal{G}}_0], & |\tilde{h}|^2[\kappa^2 \tilde{\mathcal{G}}_0] &= \kappa^{-2} |\tilde{h}|^2[\tilde{\mathcal{G}}_0], \\ \tilde{\Delta} \tilde{H}[\kappa^2 \tilde{\mathcal{G}}_0] &= \kappa^{-3} \tilde{\Delta} \tilde{H}[\tilde{\mathcal{G}}_0], & |\tilde{\nabla} \tilde{h}|^2[\kappa^2 \tilde{\mathcal{G}}_0] &= \kappa^{-4} |\tilde{\nabla} \tilde{h}|^2[\tilde{\mathcal{G}}_0], \\ \tilde{\Delta} \tilde{\Delta} \tilde{H}[\kappa^2 \tilde{\mathcal{G}}_0] &= \kappa^{-5} \tilde{\Delta} \tilde{\Delta} \tilde{H}[\tilde{\mathcal{G}}_0]. \end{aligned}$$

To understand what are scalar Riemannian invariants $I(\tilde{x}, \mathbb{R}^{1,4}, \tilde{\mathcal{G}}_0)$ we want to use the so-called Fermi coordinates. A Fermi coordinate is one such that 1) on the surface ϕ is a normal coordinate at a given point \tilde{x}_0 ; 2) the coordinate curves $\tilde{\phi}(t, v^2, v^3, v^4, v^5)$ is a geodesic perpendicular to the surface at $\phi(v^2, v^3, v^4, v^5)$ with unit speed (a line segment perpendicular to the surface in $\mathbb{R}^{1,4}$). Hence, for a Fermi coordinate,

$$(4.1.1) \quad \tilde{\phi}(v^1, \dots, v^5) = \phi(v^2, \dots, v^5) + v^1 \xi.$$

The following facts are well known.

Lemma 4.1.2. *Suppose that \tilde{x} is a timelike hypersurface in $\mathbb{R}^{1,4}$. Suppose that $\tilde{\phi}$ is a Fermi coordinate at a given point \tilde{x}_0 . Then*

$$\tilde{\mathcal{G}}_0 = \begin{bmatrix} 1 & 0 \\ 0 & [\mathcal{G}_{ij}] \end{bmatrix}$$

and

$$\mathcal{G}_{ij}(v^1, \hat{v}) = \tilde{g}_{ij}(\hat{v}) - 2\tilde{h}_{ij}(\hat{v})v^1 + \tilde{h}_{ik}(\hat{v})\tilde{h}_{jl}(\hat{v})\tilde{g}^{kl}(\hat{v})(v^1)^2,$$

where

$$\tilde{g}_{ij}(\hat{v}) = \eta_{ij} - \frac{2}{3}\tilde{R}_{ikjl}v^k v^l + \dots,$$

$$\tilde{h}_{ij}(\hat{v}) = \tilde{h}_{ij}(0) + \tilde{h}_{ij,k}(0)v^k + \dots;$$

here $\tilde{R}_{ijkl} = \tilde{h}_{ik}\tilde{h}_{jl} - \tilde{h}_{ij}\tilde{h}_{kl}$ is the Riemann curvature tensor for \tilde{x} and η is the standard matrix of signature $\{-1, 1, 1, 1\}$. All the coefficients in the Taylor expansions for G_{ij} are polynomials of \tilde{h}_{ij} and the covariant derivatives of \tilde{h}_{ij} at \tilde{x}_0 .

Therefore, in the light of Weyl's theorem on the invariants of orthogonal groups, we may conclude:

Proposition 4.1.3. *All scalar invariants $I(\tilde{x}, \mathbb{R}^{1,4}, \tilde{\mathcal{G}}_0)$ of a surface \tilde{x} in $\mathbb{R}^{1,4}$ are linear combinations of terms that are complete contractions of tensor product of the second fundamental form \tilde{h} and the covariant derivatives of \tilde{h} .*

Proof. From the above lemma it is easily that all scalar invariants of a surface \tilde{x} in $\mathbb{R}^{1,4}$ are polynomials of the first fundamental form \tilde{g} , the second fundamental form \tilde{h} and covariant derivatives of the second fundamental form \tilde{h} , if we evaluate them in a Fermi coordinate for the surface. Then, by the Weyl theorem on the invariants of orthogonal groups, we know they are linear combinations of full contractions of \tilde{h} and covariant derivatives of \tilde{h} . \square

4.2. Scalar invariants of the homogeneous associate surface \tilde{x} in $\mathbb{R}^{1,4}$. Let us work with the parametrization

$$\tilde{x} = \alpha\hat{\lambda}(1, \hat{x}) + \alpha\rho\hat{\lambda}^{-1} \frac{1}{1-a}(1, \hat{x}^*) = \alpha y_\lambda + \alpha\rho y_\lambda^*$$

and use the calculations given in [Section 3.1](#) and [Section 3.2](#). Now let us compute some scalar invariants for our associate surface \tilde{x} on the light cone where $\rho = 0$. Then the first fundamental form is

$$I_{\tilde{x}}|_{\rho=0} = \begin{bmatrix} 0 & -\alpha & 0 & 0 \\ -\alpha & 0 & \alpha^2\omega_1^\lambda & \alpha^2\omega_2^\lambda \\ 0 & \alpha^2\omega_1^\lambda & \alpha^2E_\lambda & 0 \\ 0 & \alpha^2\omega_2^\lambda & 0 & \alpha^2E_\lambda \end{bmatrix}$$

from (3.1.4), whose inverse is

$$I_{\tilde{x}}^{-1}|_{\rho=0} = \begin{bmatrix} |\omega^\lambda|^2 & -\frac{1}{\alpha} & \frac{\omega_1^\lambda}{\alpha E_\lambda} & \frac{\omega_2^\lambda}{\alpha E_\lambda} \\ -\frac{1}{\alpha} & 0 & 0 & 0 \\ \frac{\omega_1^\lambda}{\alpha E_\lambda} & 0 & \frac{1}{\alpha^2 E_\lambda} & 0 \\ \frac{\omega_2^\lambda}{\alpha E_\lambda} & 0 & 0 & \frac{1}{\alpha^2 E_\lambda} \end{bmatrix}.$$

And the second fundamental form at $\rho = 0$ is

$$II_{\tilde{x}}|_{\rho=0} = \begin{bmatrix} 0 & 0 \\ 0 & \alpha\Omega_\lambda \end{bmatrix}.$$

So the simplest (pseudo-)Riemannian invariants is the mean curvature \tilde{H} , but it is clear that

$$\tilde{H}|_{\rho=0} = \frac{1}{\alpha E_\lambda} ((\Omega_\lambda)_{11} + (\Omega_\lambda)_{22}) = 0.$$

The first nontrivial one is

$$(4.2.1) \quad |\tilde{h}|^2|_{\rho=0} = \tilde{g}^{ik} \tilde{g}^{jl} \tilde{h}_{ij} \tilde{h}_{kl}|_{\rho=0} = \alpha^{-2} |\Omega_\lambda|^2,$$

which produces the first nontrivial invariant $|\tilde{H}|^2$ for the surface \hat{x} in the conformal 3-sphere(cf. see the definition for scalar invariant of surfaces in the conformal 3-sphere in the next subsection). In fact the following nontrivial invariants without taking any derivative are all easy to calculate

$$\text{Tr}_{I_{\tilde{x}}} \tilde{h}^k|_{\rho=0} = \alpha^{-k} \text{Tr}_{I_{\hat{x}}} \Omega_\lambda^k$$

for any $k = 2, 3, \dots$. Obviously those are the ones that can be easily seen with no difficulty at all.

Next we want to calculate $|\nabla\tilde{H}|^2$ and $\tilde{\Delta}\tilde{H}$ at $\rho = 0$. To do so, let us first recall from [Section 3.2](#) the mean curvature

$$\tilde{H} = \frac{\rho \det \Omega_\lambda \mathcal{H}_\lambda}{\alpha (\det \Omega_\lambda - \rho \text{Tr} \Omega_\lambda \Omega_\lambda^* + \rho^2 \det \Omega_\lambda^*)}.$$

Hence $\tilde{H}_\alpha = \tilde{H}_{u^1} = \tilde{H}_{u^2} = 0$ and $|\nabla\tilde{H}|^2 = 0$ at $\rho = 0$, that is, $|\nabla\tilde{H}|^2$ gives no invariant for the surface \hat{x} . Let us set the convention to have a, b, c stand for α, ρ ;

i, j, k stand for u^1, u^2 , and A, B, C stand for all four variables. We then calculate, at $\rho = 0$,

$$\begin{aligned}
 (4.2.2) \quad \tilde{\Delta} \tilde{H} &= \frac{1}{\sqrt{|\tilde{g}|}} \partial_A (\sqrt{|\tilde{g}|} \tilde{g}^{AB} \partial_B \tilde{H}) \\
 &= \frac{1}{\alpha^3 E} (\partial_\alpha (\sqrt{|\tilde{g}|} \tilde{g}^{\alpha\rho} \partial_\rho \tilde{H}) + \partial_\rho (\sqrt{|\tilde{g}|} \tilde{g}^{\rho B} \partial_B \tilde{H}) + \partial_i (\sqrt{|\tilde{g}|} \tilde{g}^{i\rho} \partial_\rho \tilde{H})) \\
 &= \frac{1}{\alpha^3 E} (\partial_\alpha (\sqrt{|\tilde{g}|} \tilde{g}^{\alpha\rho} \partial_\rho \tilde{H}) + \partial_\rho (\sqrt{|\tilde{g}|} \tilde{g}^{\rho\alpha} \partial_\alpha \tilde{H}) + \sqrt{|\tilde{g}|} (\partial_\rho \tilde{g}^{\rho\rho}) \partial_\rho \tilde{H}) \\
 &= 2\alpha^{-3} \mathcal{H}_\lambda,
 \end{aligned}$$

where one needs to use the fact that $\tilde{g}^{\rho\rho}|_{\rho=0} = 0$ and $\partial_\rho \tilde{g}^{\rho\rho}|_{\rho=0} = \frac{2}{\alpha^2}$ based on calculations (A.8) in Appendix A. This confirms that \mathcal{H}_λ is indeed a conformal invariant of order 3 for a surface \hat{x} in the 3-sphere in general conformal metric $\lambda^2 g_0$.

The next invariant we want to calculate is $\tilde{\Delta} \tilde{\Delta} \tilde{H}$. To do so we observe, again from (A.8), that

$$\begin{aligned}
 (4.2.3) \quad \partial_\rho|_{\rho=0} \tilde{g}^{\rho\alpha} &= -\frac{2}{\alpha} |\omega^\lambda|^2, & \partial_\rho|_{\rho=0} \tilde{g}^{\rho\rho} &= \frac{2}{\alpha^2}, \\
 \partial_\rho|_{\rho=0} \tilde{g}^{\rho i} &= -\frac{2}{\alpha^2} \frac{\omega_i^\lambda}{E_\lambda}, & \partial_\rho \partial_\rho|_{\rho=0} \tilde{g}^{\rho\rho} &= \frac{8}{\alpha^2} |\omega^\lambda|^2.
 \end{aligned}$$

After a lengthy calculation we get

$$\begin{aligned}
 (4.2.4) \quad \tilde{\Delta} \tilde{\Delta} \tilde{H}|_{\rho=0} &= 8\alpha^{-5} \left(\Delta_\lambda \mathcal{H}_\lambda + 9|\omega^\lambda|^2 \mathcal{H}_\lambda - 3\text{Div}(\omega^\lambda) \mathcal{H}_\lambda \right. \\
 &\quad \left. - 6\omega^\lambda (\nabla \mathcal{H}_\lambda) - \frac{3 \text{Tr}(\Omega_\lambda \Omega_\lambda^*)}{2m^2} |\Omega_\lambda|^2 \mathcal{H}_\lambda \right).
 \end{aligned}$$

This tells us that the quantity in parentheses is a conformal invariant of order 5 for the surface \hat{x} in the 3-sphere.

We can also calculate the covariant derivatives of the second fundamental forms for the associate surface. We first list the relevant Christoffel symbols for the calculation

$$\begin{aligned}
 (4.2.5) \quad \tilde{\Gamma}_{\alpha\alpha}^k &= \tilde{\Gamma}_{\rho\rho}^k = \tilde{\Gamma}_{\alpha\rho}^k = 0, \\
 \tilde{\Gamma}_{\alpha j}^k &= \alpha^{-1} \delta_{jk}, \\
 \tilde{\Gamma}_{\rho j}^k &= \frac{1}{2E_\lambda} \left((\omega_k^\lambda)_{u^j} - (\omega_j^\lambda)_{u^k} + \frac{1}{m} ((\Omega_\lambda)_{jl} (\Omega_\lambda^*)_{kl} + (\Omega_\lambda)_{kl} (\Omega_\lambda^*)_{jl}) \right), \\
 \tilde{\Gamma}_{ij}^k &= (\Gamma_\lambda)_{ij}^k - \omega_k^\lambda \delta_{ij}.
 \end{aligned}$$

Then we calculate

$$\begin{aligned}
\tilde{h}_{ab,C} &= 0, \\
\tilde{h}_{ai,b} &= 0, \\
\tilde{h}_{\alpha j,k} &= -(\Omega_\lambda)_{jk}, \\
\tilde{h}_{\rho j,k} &= -\frac{\alpha}{2E_\lambda} \left((\Omega_\lambda)_{ij} ((\omega_i^\lambda)_{u^k} - (\omega_k^\lambda)_{u^i}) + \frac{1}{m} ((\Omega_\lambda)_{kl} (\Omega_\lambda^*)_{il} + (\Omega_\lambda)_{il} (\Omega_\lambda^*)_{kl}) \right), \\
(4.2.6) \quad \tilde{h}_{ij,\alpha} &= -(\Omega_\lambda)_{ij}, \\
\tilde{h}_{ij,\rho} &= \alpha (\Omega_\lambda^*)_{ij} \\
&\quad - \frac{\alpha}{2E} \left((\Omega_\lambda)_{lj} ((\omega_l^\lambda)_{u^i} - (\omega_i^\lambda)_{u^l}) + \frac{1}{m} ((\Omega_\lambda)_{kl} (\Omega_\lambda^*)_{ki} + (\Omega_\lambda)_{ki} (\Omega_\lambda^*)_{kl}) \right) \\
&\quad - \frac{\alpha}{2E} \left((\Omega_\lambda)_{il} ((\omega_l^\lambda)_{u^j} - (\omega_j^\lambda)_{u^l}) + \frac{1}{m} ((\Omega_\lambda)_{kl} (\Omega_\lambda^*)_{kj} + (\Omega_\lambda)_{kj} (\Omega_\lambda^*)_{kl}) \right), \\
\tilde{h}_{ij,k} &= \alpha (\Omega_\lambda)_{ij,k} + \alpha (\Omega_\lambda)_{lj} \omega_i^\lambda \delta_{ik} + \alpha (\Omega_\lambda)_{il} \omega_j^\lambda \delta_{jk}.
\end{aligned}$$

The easy ones are

$$\phi_\alpha = \tilde{h}_{\alpha j,k} \tilde{g}^{jk} = 0, \quad \phi_\rho = \tilde{h}_{\rho j,k} \tilde{g}^{jk} = \frac{1}{\alpha} \mathcal{H}_\lambda$$

in the light of (2.5.14). At the same time

$$\begin{aligned}
\phi_i &= \tilde{h}_{iB,C} \tilde{g}^{BC} = \tilde{h}_{ij,C} \tilde{g}^{jC} + \tilde{h}_{ib,k} \tilde{g}^{bk} = \tilde{h}_{ij,k} \tilde{g}^{jk} + \tilde{h}_{ij,\alpha} \tilde{g}^{j\alpha} + \tilde{h}_{i\alpha,k} \tilde{g}^{\alpha k} \\
&= \frac{1}{\alpha E_\lambda} (\Omega_\lambda)_{ij,j} + \frac{3}{\alpha E_\lambda} (\Omega_\lambda)_{ij} \omega_j^\lambda - \frac{1}{\alpha E_\lambda} (\Omega_\lambda)_{ij} \omega_j^\lambda - \frac{1}{\alpha E_\lambda} (\Omega_\lambda)_{ij} \omega_j^\lambda \\
&= \frac{1}{\alpha E_\lambda} (\Omega_\lambda)_{ij,j} + \frac{1}{\alpha E_\lambda} (\Omega_\lambda)_{ij} \omega_j^\lambda = 0
\end{aligned}$$

due to the integrability condition (3.3.4). Thus $|\widetilde{\text{Div}} \tilde{h}|^2 (= 0)$ does not give any invariant on the surface \hat{x} , nor does $\widetilde{\text{Div}} \tilde{h} \cdot d\tilde{H} (= 0)$, because $\tilde{g}^{\rho\rho}|_{\rho=0} = 0$.

We want to calculate $|\tilde{\nabla} \tilde{h}|^2$ since we have all the covariant derivatives $\tilde{h}_{AB,C}$ in (4.2.6). The calculation is direct yet very long. We omit details here.

$$\begin{aligned}
|\tilde{\nabla} \tilde{h}|^2|_{\rho=0} &= \alpha^{-4} \left(|\nabla \Omega|^2 + 8|dH|^2 - 6\Omega \cdot \Omega^* \right. \\
&\quad \left. - \frac{2}{E_\lambda^3} (\Omega_\lambda)_{ij} \omega_k^\lambda (R^\lambda)_{3ijk} - \frac{6}{E_\lambda^3} (\Omega_\lambda)_{ij} (\Omega_\lambda)_{ki,j} \omega_k^\lambda \right),
\end{aligned}$$

where the Codazzi equation for the surface \hat{x} in $(\mathbb{S}^3, \lambda^2 g_0)$

$$(\Omega_\lambda)_{ij,k} = (\Omega_\lambda)_{ik,j} + (R^\lambda)_{3ijk} + (H_\lambda)_{uj} E_\lambda \delta_{ik} - (H_\lambda)_{uk} E_\lambda \delta_{ij}$$

has been used. At this point we like to write each term as local scalar invariant of the surface \hat{x} in $(\mathbb{S}^3, \lambda^2 g_0)$. We first calculate

$$\begin{aligned}
& (\Omega_\lambda)_{ij} \omega_k^\lambda (R^\lambda)_{3ijk} \\
&= (\Omega_\lambda)_{ij} \omega_1^\lambda (R^\lambda)_{3ij1} + (\Omega_\lambda)_{ij} \omega_2^\lambda (R^\lambda)_{3ij2} \\
&= E_\lambda ((\Omega_\lambda)_{11} \omega_1^\lambda (R^\lambda)_{31} + (\Omega_\lambda)_{21} \omega_1^\lambda (R^\lambda)_{32} + (\Omega_\lambda)_{22} \omega_2^\lambda (R^\lambda)_{32} + (\Omega_\lambda)_{12} \omega_2^\lambda (R^\lambda)_{31}) \\
&= E_\lambda (\Omega_\lambda)_{ij} \omega_j^\lambda (R^\lambda)_{3i} = -E_\lambda^2 (H_\lambda)_{ui} (R^\lambda)_{3i} = -\text{Ric}^\lambda(\vec{n}_\lambda, \lambda H_\lambda).
\end{aligned}$$

Then we deal with the last term:

$$\begin{aligned}
(\Omega_\lambda)_{ij} (\Omega_\lambda)_{ki,j} \omega_k^\lambda &= (\Omega_\lambda)_{ij} ((\Omega_\lambda)_{ki} \omega_k^\lambda)_{,j} - (\Omega_\lambda)_{ij} (\Omega_\lambda)_{ki} \omega_{k,j}^\lambda \\
&= -E_\lambda (\Omega_\lambda)_{ij} (H_\lambda)_{i,j} - \frac{1}{2} E_\lambda^3 |\Omega_\lambda|^2 \text{Div}(\omega^\lambda),
\end{aligned}$$

where

$$\begin{aligned}
\text{Div}(\omega^\lambda) &= E_\lambda^{-1} \omega_{i,i}^\lambda = E_\lambda^{-1} (\omega_i^\lambda)_{,i} \\
&= E_\lambda^{-1} (\langle \Delta_0 y_\lambda, y_\lambda^* \rangle + \langle (y_\lambda)_{,ui}, (y_\lambda^*)_{,ui} \rangle) \\
&= H_\lambda^2 - |\omega^\lambda|^2 + (R^\lambda)_{1212} + E^{-1} \langle (y_\lambda)_{,ui}, (y_\lambda^*)_{,ui} \rangle \\
&= H_\lambda^2 + 2 \frac{\Omega_\lambda \cdot \Omega_\lambda^*}{|\Omega_\lambda|^2} + E_\lambda^{-1} (R^\lambda)_{1212}, \\
\Delta_0 y_\lambda &= 2E_\lambda H_\lambda \vec{n}_\lambda + 2E_\lambda y_\lambda^\dagger - (R^\lambda)_{1212} y_\lambda,
\end{aligned}$$

and

$$\sum_{i=1}^2 E^{-1} \langle (y)_{,ui}, (y^*)_{,ui} \rangle = |\omega^\lambda|^2 + 2 \frac{\Omega_\lambda \cdot \Omega_\lambda^*}{|\Omega_\lambda|^2}.$$

So we have obtained

$$\begin{aligned}
(4.2.7) \quad |\nabla \tilde{h}|^2|_{\rho=0} &= \alpha^{-4} (|\nabla \Omega_\lambda|^2 + 8|dH_\lambda|^2 + 2 \text{Ric}^\lambda(\vec{n}_\lambda, \nabla H_\lambda) + 3H_\lambda^2 |\Omega|^2 \\
&\quad + 3K_\lambda^T |\Omega_\lambda|^2 + 6\Omega_\lambda \cdot \text{Hess}(H_\lambda)),
\end{aligned}$$

where

$$K_\lambda^T = E_\lambda^{-1} (R^\lambda)_{1212}$$

is the sectional curvature of $(\mathbb{S}^3, \lambda^2 g_0)$ of the tangent plane to the surface \hat{x} .

4.3. Scalar invariants for surfaces in the conformal round 3-sphere. Let us start with the definition of scalar invariants for surfaces in conformal sphere.

Definition 4.3.1. Let $i : M^{n-1} \rightarrow N^n$ be an immersed hypersurface and let $[g]$ be a class of conformal metrics on the ambient manifold N^n . $I_c(i, N^n, g)$ is said to be a scalar conformal invariant of the hypersurface i in the conformal manifold $(N^n, [g])$ if it is a scalar Riemannian invariant and

$$(4.3.1) \quad I_c(i, N^n, \lambda^2 g) = \lambda^{-k} I_c(i, N^n, g).$$

for any positive function λ on N^n , where k is the order of the invariant $I_c(\hat{i}, N^n, g)$.

Recall that, for an immersed surface \hat{x} in S^3 , we have

$$\hat{I}(\hat{x}, S^3, \lambda^2 g_0) = \lambda \hat{I}(\hat{x}, S^3, g_0).$$

Hence it is easy to observe that

$$|\hat{I}|^2(\hat{x}, S^3, \lambda^2 g_0) = \lambda^{-2} g_0^{ik} \lambda^{-2} g_0^{jl} \lambda \hat{I}_{ij} \lambda \hat{I}_{kl} = \lambda^{-2} \|\hat{I}\|^2(\hat{x}, S^3, g_0)$$

and

$$\text{Tr}_{\lambda^2 g_0}(\hat{I})^k(\hat{x}, S^3, \lambda^2 g_0) = \lambda^{-k} \text{Tr}_{g_0}(\hat{I})^k(\hat{x}, S^3, g_0) \text{ for all } k = 2, 3, \dots$$

On the other hand, it does not seem easy to directly verify that \mathcal{H}_λ is a conformal invariant for a surface in the conformal 3-sphere, though this is a well-known one. We have verified this in computing the mean curvature (cf. (2.5.12)) of the surface ξ in the de Sitter spacetime $S^{1,3}$ as well as in the above calculation of $\tilde{\Delta}\tilde{H}$ (cf. (4.2.2)) of the homogeneous associate surface \tilde{x} . In general it takes tremendous, if not impossible, to verify whether an invariant $I(\hat{x}, S^3, \lambda^2 g_0)$ is conformally invariant, complicated by the six integrability conditions. The most important application of the construction of associate homogeneous surfaces is the following:

Theorem 4.3.2. *Suppose that $\hat{x} : M^2 \rightarrow S^3$ is an immersed surface with no umbilical point, and let*

$$\tilde{x} = \alpha y + \alpha \rho y^* : \mathbb{R}^+ \times \mathbb{R}^+ \times M^2 \rightarrow \mathbb{R}^{1,4}$$

be the associate surface for \hat{x} , where \hat{x}^ is the conformal transform of \hat{x} . Then any scalar (pseudo)-Riemannian invariant $I(\tilde{x}, \mathbb{R}^{1,4}, \tilde{\mathcal{G}}_0)$ evaluated at $\rho = 0$, if it is nontrivial, is a scalar conformal invariant $I_c(\hat{x}, S^3, \lambda^2 g_0)$ multiplied with $|\hat{I}_\lambda|^{2n}$ for some integer n .*

Proof. For any invariant $I(\tilde{x}, \mathbb{R}^{1,4}, \tilde{\mathcal{G}}_0)$, we know that it is a full contraction of tensor product of the second fundamental form and the covariant derivatives. For a choice of representative $\lambda^2 g_0$ on S^3 , in the corresponding parametrization (3.1.1), we claim that

$$(4.3.2) \quad I(\tilde{x}, \mathbb{R}^{1,4}, \tilde{\mathcal{G}}_0)|_{\rho=0} = \alpha^{-k} I(\hat{x}, S^3, \lambda^2 g_0) |\hat{I}_\lambda|^{2n}$$

for a positive integer k and a nonnegative integer n , due to the homogeneity of the associate surface. To see the right side of (4.3.2) is indeed a scalar Riemannian invariant multiplied with factor $|\hat{I}_\lambda|^{2n}$ for some integer n , we consider the tensors that determines the first and second fundamental forms of the associate surface in that parametrization. We recall from (2.5.5) that

$$\Omega_\lambda = \hat{I}_\lambda$$

is the traceless part of the second fundamental form for the surface \hat{x} in the 3-sphere with the conformal metric $\lambda^2 g_0$. We also know from (3.1.3) that

$$\omega^\lambda = -I_\lambda((\mathring{H}_\lambda)^{-1}(dH_\lambda)) = -\frac{2}{|\mathring{H}_\lambda|^2} \mathring{H}_\lambda(\nabla H_\lambda),$$

which causes us to include the possibly negative n in the right side of (4.3.2). We may also recall from (2.5.3) that

$$m = \frac{1}{2} E_\lambda |\mathring{H}_\lambda|^2.$$

Next we want to show that Ω_λ^* is also a tensor product of covariant derivatives of the 1-form ω^λ , covariant derivatives of the second fundamental form H_λ and covariant derivatives of Riemann curvature tensor of the conformal metric $\lambda^2 g_0$ on the 3-sphere(including 0th order). Recall the definition

$$(\Omega_\lambda^*)_{ij} = \langle y_\lambda^*, \xi_{u^i u^j} \rangle.$$

We use the same idea in the calculation of the trace of Ω^* in Section 2.5. Hence we write

$$(4.3.3) \quad \xi_{u^i u^j} = -(\Omega_\lambda^*)_{ij} y_\lambda - (\Omega_\lambda)_{ij} y_\lambda^* + (\Gamma_m)_{ij}^k \xi_{u^k} - m \delta_{ij} \xi.$$

From (2.6.2) we know that

$$\langle y_\lambda^*, y_\lambda^\dagger \rangle = -\frac{1}{2} (|\omega^\lambda|^2 + H_\lambda^2).$$

Using $\xi = H_\lambda y_\lambda + \vec{n}_\lambda$ from Lemma 2.3.4 and (B.6), we have

$$\langle \xi_{u^k}, y_\lambda^\dagger \rangle = -(H_\lambda)_{u^k} + (R^\lambda)_{3k}$$

and

$$\langle \xi, y_\lambda^\dagger \rangle = -H_\lambda.$$

Therefore we derive from (4.3.3) that

$$(4.3.4) \quad \begin{aligned} & \langle \xi_{u^i u^j}, y_\lambda^\dagger \rangle \\ &= (\Omega_\lambda^*)_{ij} + \frac{1}{2} (|\omega^\lambda|^2 + H_\lambda^2) (\Omega_\lambda)_{ij} + (\Gamma_m)_{ij}^k (-H_{u^k} + (R^\lambda)_{3k}) + H m \delta_{ij}, \end{aligned}$$

where

$$(\Gamma_m)_{ij}^k = \Gamma_{ij}^k + \frac{1}{2} |\Omega_\lambda|^{-2} (|\Omega_\lambda|_{u^i}^2 \delta_{jk} + |\Omega_\lambda|_{u^j}^2 \delta_{ik} - |\Omega_\lambda|_{u^k}^2 \delta_{ij})$$

represents the Christoffel symbols for the Möbius metric $m|du|^2$. On the other hand we have

$$\xi_{u^i u^j} = (H_\lambda)_{u^i u^j} y_\lambda + (H_\lambda)_{u^i} (y_\lambda)_{u^j} + (H_\lambda)_{u^j} (y_\lambda)_{u^i} + H_\lambda (y_\lambda)_{u^i u^j} + (\vec{n}_\lambda)_{u^i u^j}$$

which implies

$$\begin{aligned}
 (4.3.5) \quad \langle \xi_{u^i u^j}, y_\lambda^\dagger \rangle &= -(H_\lambda)_{u^i u^j} + H_\lambda \langle (y_\lambda)_{u^i u^j}, y_\lambda^\dagger \rangle + \langle (\vec{n}_\lambda)_{u^i u^j}, y_\lambda^\dagger \rangle \\
 &= -(H_\lambda)_{u^i u^j} - H_\lambda \langle (y_\lambda)_{u^i}, (y_\lambda^\dagger)_{u^j} \rangle - \langle (\vec{n}_\lambda)_{u^i}, (y_\lambda^\dagger)_{u^j} \rangle - \langle \vec{n}_\lambda, (y_\lambda^\dagger)_{u^i} \rangle_{u^j} \\
 &= -(H_\lambda)_{u^i u^j} + \frac{1}{E_\lambda} (\Omega_\lambda)_{ik} \langle (y_\lambda)_{u^k}, (y_\lambda^\dagger)_{u^i} \rangle - \langle \vec{n}_\lambda, (y_\lambda^\dagger)_{u^i} \rangle_{u^j} \\
 &= -(H_\lambda)_{u^i u^j} - \frac{1}{E_\lambda} (\Omega_\lambda)_{ik} (R^\lambda)_{i3k3} + ((R^\lambda)_{3i})_{u^j},
 \end{aligned}$$

by (B.7) and (B.6). Thus, comparing (4.3.4) and (4.3.5), we have

$$\begin{aligned}
 (4.3.6) \quad (\Omega_\lambda^*)_{ij} &= -(H_\lambda)_{u^i, u^j} - H_\lambda m \delta_{ij} - \frac{1}{E_\lambda} (\Omega_\lambda)_{ik} (R^\lambda)_{j3k3} + ((R^\lambda)_{3i})_{,u^j} \\
 &\quad - \frac{1}{2} (|\omega^\lambda|^2 + H_\lambda^2) (\Omega_\lambda)_{ij} \\
 &\quad + \frac{1}{2} |\Omega_\lambda|^{-2} (|\Omega_\lambda|_{u^i}^2 \delta_{jk} + |\Omega_\lambda|_{u^j}^2 \delta_{ik} - |\Omega_\lambda|_{u^k}^2 \delta_{ij}) ((H_\lambda)_{u^k} - (R^\lambda)_{3k}).
 \end{aligned}$$

The last factor that goes into the left side of Equation (4.3.2) is the reciprocal of the determinant:

$$\det \tilde{g}|_{\rho=0} = -\frac{\alpha^6}{m^2} (pr - q^2)^2|_{\rho=0} = \frac{\alpha^6}{m^2} (\det \Omega_\lambda)^2 = \alpha^6 E_\lambda^2 = \alpha^6 \det I_\lambda^{\hat{x}}.$$

due to (3.1.6), where $I_\lambda^{\hat{x}} = (\hat{x})^*(\lambda^2 g_0) = E_\lambda |du|^2$.

To verify that the right side of (4.3.2) is actually a conformal invariant, for a positive functions λ on 3-sphere, we simply compare the right side of (4.3.2) evaluated at $\alpha = 1$ with that evaluated at $\alpha = \hat{\lambda}$ and $\lambda = 1$. We then observe that

$$\mathbf{I}(\hat{x}, \mathbb{S}^3, \lambda^2 g_0) = \hat{\lambda}^{-k} \mathbf{I}(\hat{x}, \mathbb{S}^3, g_0).$$

Therefore it is a conformal scalar invariant for the surface \hat{x} in the 3-sphere. \square

Appendix A: The inverse of $I^{\tilde{x}}$ in general parametrizations

We consider the general parametrization

$$\tilde{x} = \alpha y_\lambda + \alpha \rho y_\lambda^* : \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbf{M}^2 \rightarrow \mathbb{R}^{1,4}.$$

Then the first fundamental form in matrix form is

$$(A.1) \quad I_{\tilde{x}} = \begin{bmatrix} -2\rho & -\alpha & 0 & 0 \\ -\alpha & 0 & \alpha^2 \omega_1^\lambda & \alpha^2 \omega_2^\lambda \\ 0 & \alpha^2 \omega_1^\lambda & & \alpha^2 F \\ 0 & \alpha^2 \omega_2^\lambda & & \end{bmatrix}$$

where

$$(A.2) \quad \begin{cases} F_{11} = \frac{1}{m}(p^2 + q^2) + 2\rho(\omega_1^\lambda)^2 \\ F_{12} = F_{21} = \frac{1}{m}q(p+r) + 2\rho\omega_1^\lambda\omega_2^\lambda \\ F_{22} = \frac{1}{m}(q^2 + r^2) + 2\rho(\omega_2^\lambda)^2 \end{cases} \text{ and } \begin{cases} F_{11}^* = \frac{1}{m}(p^2 + q^2) \\ F_{12}^* = F_{21}^* = \frac{1}{m}q(p+r) \\ F_{22}^* = \frac{1}{m}(q^2 + r^2) \end{cases}$$

and

$$\begin{bmatrix} p & q \\ q & r \end{bmatrix} = \Omega_\lambda + \rho\Omega_\lambda^*.$$

It is easily seen that

$$(A.3) \quad (F^*)^{-1} = \frac{m}{(pr - q^2)^2} \begin{bmatrix} r^2 + q^2 & -q(p+r) \\ -q(p+r) & p^2 + q^2 \end{bmatrix}$$

and

$$F|_{\rho=0} = F^*|_{\rho=0} = E \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Let

$$(I^{\tilde{x}})^{-1} = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}.$$

Therefore, for example,

$$(A.4) \quad \begin{aligned} -2\rho a_{11} - \alpha a_{12} &= 1, \\ -\alpha a_{11} + \alpha^2\omega_1 a_{13} + \alpha^2\omega_2 a_{14} &= 0, \\ \alpha^2\omega_1 a_{12} + \alpha^2 F_{11} a_{13} + \alpha^2 F_{21} a_{14} &= 0, \\ \alpha^2\omega_2 a_{12} + \alpha^2 F_{12} a_{13} + \alpha^2 F_{22} a_{14} &= 0. \end{aligned}$$

Subtracting the first of these equations multiplied by α from the second equation multiplied by 2, we get

$$(A.5) \quad \alpha^2 a_{12} + 2\alpha^2 \rho \omega_1 a_{13} + 2\alpha^2 \rho \omega_2 a_{14} = -\alpha.$$

Subtracting (A.5) multiplied by ω_1 from the third equation in (A.4) as well as subtracting (A.5) multiplied by ω_2 from the fourth equation in (A.4), we get

$$(A.6) \quad \alpha^2 F^* \begin{bmatrix} a_{13} \\ a_{14} \end{bmatrix} = \begin{bmatrix} \alpha\omega_1 \\ \alpha\omega_2 \end{bmatrix}$$

Plugging back the values of a_{13} and a_{14} into (A.5) we have

$$(A.7) \quad \begin{cases} a_{12} = \alpha^{-1} \left(-1 - 2\rho[\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix} \right), \\ a_{11} = -\frac{\alpha a_{12} + 1}{2\rho} = [\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix}. \end{cases}$$

Similarly one gets

$$\alpha^2 F^* \begin{bmatrix} a_{23} \\ a_{24} \end{bmatrix} = \begin{bmatrix} -2\rho\omega_1 \\ -2\rho\omega_2 \end{bmatrix}, \quad \alpha^2 F^* \begin{bmatrix} a_{33} \\ a_{34} \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \alpha^2 F^* \begin{bmatrix} a_{43} \\ a_{44} \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

which yield respectively

$$(A.8) \quad \begin{cases} a_{21} = \alpha^{-1} \left(-1 - 2\rho[\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix} \right), \\ a_{22} = \frac{2\rho}{\alpha^2} \left(1 + 2\rho[\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix} \right), \end{cases}$$

$$(A.9) \quad \begin{cases} a_{31} = \alpha^{-1} [\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \\ a_{32} = -\frac{2\rho}{\alpha^2} ([\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} 1 \\ 0 \end{bmatrix}), \end{cases}$$

$$(A.10) \quad \begin{cases} a_{41} = \alpha^{-1} (F^*)^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \\ a_{42} = \frac{2\rho}{\alpha^2} \left(1 + 2\rho[\omega_1, \omega_2](F^*)^{-1} \begin{bmatrix} \omega_1 \\ \omega_2 \end{bmatrix} \right). \end{cases}$$

Appendix B: The geometry of the 3-sphere \mathbb{S}_λ^3 in $\mathbb{R}^{1,4}$

Let us calculate the Gauss Theorem for the 3-sphere \mathbb{S}_λ^3 in Minkowski spacetime $\mathbb{R}^{1,4}$. There is nothing new or difficult about the calculation, but this helps to understand better about the geometry of the 3-sphere $\mathbb{S}_\lambda^3 \subset \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$. Crucial to our approach is that the induced metric on \mathbb{S}_λ^3 is exactly the conformal metric $\lambda^2 g_0$. We consider the Fermi parametrization induced from a parametrization of the surface $\hat{x} : M^2 \rightarrow \mathbb{S}^3$ such that

$$(B.1) \quad y_\lambda = \lambda(\hat{x}(u^1, u^2, u^3))(1, \hat{x}(u^1, u^2, u^3)) : M^3 \rightarrow \mathbb{S}_\lambda^3 \subset \mathbb{N}_+^4 \subset \mathbb{R}^{1,4}$$

with

$$(B.2) \quad \hat{x}(u^1, u^2, 0) = \hat{x}(u^1, u^2) \text{ and } (y_\lambda)_{u^3}|_{u^3=0} = \vec{n}_\lambda.$$

Notice that y_λ here is the extension of $\hat{\lambda}(1, \hat{x})$ before. We use the two null normal vectors $\{y_\lambda, y_\lambda^\dagger\}$ where

$$(B.3) \quad \langle y_\lambda^\dagger, y_\lambda \rangle = 1, \quad \langle y_\lambda^\dagger, (y_\lambda)_{u^1} \rangle = \langle y_\lambda^\dagger, (y_\lambda)_{u^2} \rangle = \langle y_\lambda^\dagger, (y_\lambda)_{u^3} \rangle = 0.$$

The first fundamental form is $I^{\mathbb{S}^3} = \lambda^2 g_0 = \langle dy_\lambda, dy_\lambda \rangle$, and the second is $II^{\mathbb{S}^3} = -\langle dy_\lambda, dy_\lambda^\dagger \rangle y_\lambda^\dagger - \langle dy_\lambda, dy_\lambda \rangle y_\lambda$. To find the curvature of the metric $g_\lambda = \lambda^2 g_0$ we calculate

$$(B.4) \quad \nabla_{\partial_{u^j}}^\lambda \nabla_{\partial_{u^i}}^\lambda \partial_{u^k} - \nabla_{\partial_{u^i}}^\lambda \nabla_{\partial_{u^j}}^\lambda \partial_{u^k} = R^\lambda(\partial_{u^i}, \partial_{u^j})\partial_{u^k} = (R^\lambda)_{ijk}{}^l \partial_{u^l}.$$

First

$$(B.5) \quad \nabla_{\partial_{u^j}}^\lambda \partial_{u^k} = (y_\lambda)_{u^k u^j} - \langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle y_\lambda - \langle (y_\lambda)_{u^j}, (y_\lambda)_{u^k} \rangle y_\lambda^\dagger;$$

then

$$\begin{aligned} \partial_{u^i} \nabla_{\partial_{u^j}}^\lambda \partial_{u^k} &= (y_\lambda)_{u^k u^j u^i} - \langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle_{u^i} y_\lambda - \langle (y_\lambda)_{u^j}, (y_\lambda)_{u^k} \rangle_{u^i} y_\lambda^\dagger \\ &\quad - \langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle (y_\lambda)_{u^i} - \langle (y_\lambda)_{u^j}, (y_\lambda)_{u^k} \rangle (y_\lambda^\dagger)_{u^i} \end{aligned}$$

and

$$\begin{aligned} \nabla_{\partial_{u^i}}^\lambda \nabla_{\partial_{u^j}}^\lambda \partial_{u^k} &= (\partial_{u^i} \nabla_{\partial_{u^j}}^\lambda \partial_{u^k})^{T\mathbb{S}^3} \\ &= (y_\lambda)_{u^k u^j u^i}^{T\mathbb{S}^3} - \langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle (y_\lambda)_{u^i} - \langle (y_\lambda)_{u^j}, (y_\lambda)_{u^k} \rangle (y_\lambda^\dagger)_{u^i} \end{aligned}$$

Hence

$$\begin{aligned} (R^\lambda)_{ijk}{}^l \partial_{u^l} &= \langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle (y_\lambda)_{u^i} + \langle (y_\lambda)_{u^j}, (y_\lambda)_{u^k} \rangle (y_\lambda^\dagger)_{u^i} \\ &\quad - \langle (y_\lambda)_{u^i}, (y_\lambda^\dagger)_{u^k} \rangle (y_\lambda)_{u^j} - \langle (y_\lambda)_{u^i}, (y_\lambda)_{u^k} \rangle (y_\lambda^\dagger)_{u^j}. \end{aligned}$$

One notices that $\langle (y_\lambda^\dagger)_{u^i}, y_\lambda^\dagger \rangle = 0$ and $\langle (y_\lambda^\dagger)_{u^i}, y_\lambda \rangle = 0$ and concludes that

$$(y_\lambda^\dagger)_{u^i} = (g_\lambda)^{ml} \langle (y_\lambda^\dagger)_{u^i}, (y_\lambda)_{u^m} \rangle (y_\lambda)_{u^l}.$$

Therefore

$$\begin{aligned} (R^\lambda)_{ijk}{}^l \partial_{u^l} &= (\langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle \delta_i^l + (g_\lambda)_{jk} (g_\lambda)^{ml} \langle (y_\lambda^\dagger)_{u^i}, (y_\lambda)_{u^m} \rangle \\ &\quad - \langle (y_\lambda)_{u^i}, (y_\lambda^\dagger)_{u^k} \rangle \delta_j^l - (g_\lambda)_{ik} (g_\lambda)^{ml} \langle (y_\lambda^\dagger)_{u^j}, (y_\lambda)_{u^m} \rangle) \partial_{u^l} \end{aligned}$$

and

$$\begin{aligned} (R^\lambda)_{ijkl} &= (R^\lambda)_{ijk}{}^n (g_\lambda)_{nl} \\ &= \langle (y_\lambda)_{u^j}, (y_\lambda^\dagger)_{u^k} \rangle (g_\lambda)_{il} + \langle (y_\lambda^\dagger)_{u^i}, (y_\lambda)_{u^l} \rangle (g_\lambda)_{jk} \\ &\quad - \langle (y_\lambda)_{u^i}, (y_\lambda^\dagger)_{u^k} \rangle (g_\lambda)_{jl} - \langle (y_\lambda^\dagger)_{u^j}, (y_\lambda)_{u^l} \rangle (g_\lambda)_{ik}. \end{aligned}$$

On the surface \hat{x} , where $u^3 = 0$, we have

$$[(g_\lambda)_{ij}] = \begin{bmatrix} E_\lambda & 0 & 0 \\ 0 & E_\lambda & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Therefore we have, for $i, j \in \{1, 2\}$,

$$\begin{aligned} & -\langle (y_\lambda)_{ui}, (y_\lambda^\dagger)_{uj} \rangle - \langle (y_\lambda)_{u^3}, (y_\lambda^\dagger)_{u^3} \rangle E_\lambda \delta_{ij} = (R^\lambda)_{i3j3}, \\ & -\langle (y_\lambda)_{uj}, (y_\lambda^\dagger)_{u^3} \rangle E_\lambda \delta_{jl} + \langle (y_\lambda)_{ul}, (y_\lambda^\dagger)_{u^3} \rangle E_\lambda = (R^\lambda)_{3jil}, \\ & -\langle (y_\lambda)_{ui}, (y_\lambda^\dagger)_{ui} \rangle E_\lambda - \langle (y_\lambda)_{uj}, (y_\lambda^\dagger)_{uj} \rangle E_\lambda = (R^\lambda)_{ijij}. \end{aligned}$$

We obtain, for $i, j \in \{1, 2\}$,

$$(B.6) \quad \langle \vec{n}_\lambda, (y_\lambda^\dagger)_{ui} \rangle = \frac{1}{E_\lambda} (R^\lambda)_{ijj3} = -(R^\lambda)_{i3},$$

and for $i \neq j$,

$$(B.7) \quad \begin{aligned} \langle (y_\lambda)_{ui}, (y_\lambda^\dagger)_{uj} \rangle &= -(R^\lambda)_{i3j3} \\ \langle (y_\lambda)_{ui}, (y_\lambda^\dagger)_{ui} \rangle &= -(R^\lambda)_{i3i3} + \frac{1}{2}((R^\lambda)_{33} - (R^\lambda)_{1212}) \\ \langle (y_\lambda)_{u^3}, (y_\lambda^\dagger)_{u^3} \rangle &= -\frac{1}{2}((R^\lambda)_{33} - (R^\lambda)_{1212}) \end{aligned}$$

Finally, for the induced Fermi coordinate from an isothermal coordinate, we can easily see that

$$(B.8) \quad \begin{aligned} (R^\lambda)_{3i},{}^i &= \frac{1}{E_\lambda} \left(\sum_{i=1}^2 R^\lambda \right)_{3i,i} = \frac{1}{E_\lambda} \sum_{i=1}^2 (((R^\lambda)_{3i})_{ui} - (R^\lambda)_{3k}(\Gamma_\lambda)_{ii}^k) \\ &= \frac{1}{E_\lambda} \sum_{i=1}^2 (((R^\lambda)_{3i})_{ui}). \end{aligned}$$

Indeed, we have $\sum_{i=1}^2 (\Gamma_\lambda)_{ii}^k = 0$ for $k = 1, 2$, where $(\Gamma_\lambda)_{ij}^k$ are the Christoffel symbols for the conformal metric $I_\lambda = E_\lambda |du|^2$ in isothermal coordinates.

References

- [Bailey et al. 1994] T. N. Bailey, M. G. Eastwood, and C. R. Graham, “Invariant theory for conformal and CR geometry”, *Ann. of Math. (2)* **139**:3 (1994), 491–552. [MR](#)
- [Blaschke 1929] W. Blaschke, *Vorlesungen über Differentialgeometrie und geometrische Grundlagen von Einsteins Relativitätstheorie, III: Differentialgeometrie der Kreise und Kugeln*, edited by G. Thomsen, Grundlehren der Math. Wissenschaften **1**, Springer, Berlin, 1929. [Zbl](#) [JFM](#)
- [Bryant 1984] R. L. Bryant, “A duality theorem for Willmore surfaces”, *J. Differential Geom.* **20**:1 (1984), 23–53. [MR](#) [Zbl](#)
- [Fefferman and Graham 1985] C. Fefferman and C. R. Graham, “Conformal invariants”, pp. 95–116 in *The mathematical heritage of Élie Cartan* (Lyon, 1984), Astérisque (numéro hors série), Société Mathématique de France, Paris, 1985. [MR](#) [Zbl](#)
- [Fefferman and Graham 2012] C. Fefferman and C. R. Graham, *The ambient metric*, Annals of Mathematics Studies **178**, Princeton University Press, Princeton, NJ, 2012. [MR](#) [Zbl](#)
- [Gover 2001] A. R. Gover, “Invariant theory and calculus for conformal geometries”, *Adv. Math.* **163**:2 (2001), 206–257. [MR](#) [Zbl](#)

- [Gover and Waldron 2015] A. R. Gover and A. Waldron, “Conformal hypersurface geometry via a boundary Loewner–Nirenberg–Yamabe problem”, 2015. [arXiv](#)
- [Hu and Li 2004] Z. Hu and H. Li, “Willmore submanifolds in a Riemannian manifold”, pp. 251–275 in *Contemporary geometry and related topics*, edited by N. Bokan et al., World Sci. Publ., 2004. [MR](#) [Zbl](#)
- [Li and Yau 1982] P. Li and S. T. Yau, “A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces”, *Invent. Math.* **69**:2 (1982), 269–291. [MR](#) [Zbl](#)
- [Marques and Neves 2014] F. C. Marques and A. Neves, “Min-max theory and the Willmore conjecture”, *Ann. of Math. (2)* **179**:2 (2014), 683–782. [MR](#) [Zbl](#)
- [Rigoli 1987] M. Rigoli, “The conformal Gauss map of submanifolds of the Möbius space”, *Ann. Global Anal. Geom.* **5**:2 (1987), 97–116. [MR](#) [Zbl](#)
- [Wang 1992] C. P. Wang, “Surfaces in Möbius geometry”, *Nagoya Math. J.* **125** (1992), 53–72. [MR](#) [Zbl](#)
- [Wang 1998] C. Wang, “Moebius geometry of submanifolds in S^n ”, *Manuscripta Math.* **96**:4 (1998), 517–534. [MR](#)

Received February 19, 2016.

JIE QING
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, SANTA CRUZ
SANTA CRUZ, CA 95064
UNITED STATES
qing@ucsc.edu

CHANGPING WANG
SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE
FUJIAN NORMAL UNIVERSITY
FUZHOU, 350108
CHINA
cpwang@fjnu.edu.cn

JINGYANG ZHONG
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, SANTA CRUZ
SANTA CRUZ, CA 95064
UNITED STATES

and

DEPARTMENT OF MATHEMATICAL SCIENCES
TSINGHUA UNIVERSITY
BEIJING, 100084
CHINA
jzhong2@ucsc.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 286 No. 1 January 2017

Elliptic curves, random matrices and orbital integrals	1
JEFFREY D. ACHTER and JULIA GORDON	
On the absolute continuity of p -harmonic measure and surface measure in Reifenberg flat domains	25
MURAT AKMAN	
On the geometry of gradient Einstein-type manifolds	39
GIOVANNI CATINO, PAOLO MASTROLIA, DARIO D. MONTICELLI and MARCO RIGOLI	
On the Fourier–Jacobi model for some endoscopic Arthur packet of $U(3) \times U(3)$: the nongeneric case	69
JAEHO HAAN	
A Kirchberg-type tensor theorem for operator systems	91
KYUNG HOON HAN	
Remarks on quantum unipotent subgroups and the dual canonical basis	125
YOSHIYUKI KIMURA	
Scalar invariants of surfaces in the conformal 3-sphere via Minkowski spacetime	153
JIE QING, CHANGPING WANG and JINGYANG ZHONG	
Action of intertwining operators on pseudospherical K -types	191
SHIANG TANG	
Local symmetric square L -factors of representations of general linear groups	215
SHUNSUKE YAMANA	