

*Pacific
Journal of
Mathematics*

**ALMOST EVERYWHERE CONVERGENCE
FOR MODIFIED BOCHNER–RIESZ MEANS
AT THE CRITICAL INDEX FOR $p \geq 2$**

MARCO ANNONI

ALMOST EVERYWHERE CONVERGENCE FOR MODIFIED BOCHNER–RIESZ MEANS AT THE CRITICAL INDEX FOR $p \geq 2$

MARCO ANNONI

Boundedness for a maximal modified Bochner–Riesz operator between weighted L^2 spaces is proved. As a consequence, we have sufficient conditions for a.e. convergence of the modified Bochner–Riesz means at the critical exponent $p_\lambda = 2n/(n - 2\lambda - 1)$.

1. Introduction	257
2. Reduction of Theorem 1.1 to Lemma 2.1	260
3. An upper bound for $ \widehat{w}_{\lambda, \mu} $	261
4. A useful weight comparable to $1/w_{\lambda, \mu}$	263
5. Reduction of Lemma 2.1 to Lemma 5.2	266
6. Proof of Lemma 5.2	270
Acknowledgement	274
References	274

1. Introduction

This paper contains the results proved in the author’s doctoral dissertation [Annoni 2010] and referenced by S. Lee and A. Seeger [2015], but yet unpublished in a mathematical journal. For $\lambda, R > 0$, let B_R^λ denote the Bochner–Riesz operators and m_λ the Fourier multipliers introduced in [Bochner 1936]:

$$B_R^\lambda(f)(x) = \int_{\mathbb{R}^n} \widehat{f}(\xi) m_\lambda\left(\frac{|\xi|}{R}\right) e^{2\pi i \xi x} d\xi, \quad m_\lambda(t) = (1 - t^2)_+^\lambda.$$

For $p < 2$, results related to almost everywhere convergence and maximal operators have been proved by Tao [1998; 2002], Ashurov [1983], and Ahmedov, Ashurov, and Mahmud [Ashurov et al. 2010]. For $p \geq 2$, partial results on almost everywhere convergence of $B_R^\lambda(f)$ to f as $R \rightarrow \infty$ have been achieved in [Carbery 1983; Christ 1985]. Carbery, Rubio de Francia, and Vega [Carbery et al. 1988] obtained a.e. convergence in the range $2 \leq p < p_\lambda$ and $\lambda > 0$.

MSC2010: primary 42B15; secondary 42B10, 42B25.

Keywords: Bochner–Riesz means, maximal Bochner–Riesz means, almost everywhere convergence, weighted inequalities, radial multipliers.

In this paper, the situation at the critical exponent $p_\lambda = 2n/(n - 2\lambda - 1)$ is studied by considering the modified Bochner–Riesz multipliers $m_{\lambda,\gamma}$

$$m_{\lambda,\gamma}(t) = \frac{(1 - t^2)_+^\lambda}{(1 - \log(1 - t^2))^\gamma},$$

which were introduced by Seeger [1987]. Seeger [1996] showed that $m_{\lambda,\gamma}$ is an $L^{p_\lambda}(\mathbb{R}^2)$ multiplier for $\gamma > 1/p'_\lambda$ (where $1/p_\lambda + 1/p'_\lambda = 1$). His results easily extend to dimensions $n \geq 3$ when $\lambda \geq (n - 1)/(2(n + 1))$ and had already been proven to be sharp in [Seeger 1987] when $n = 2$.

In order to investigate for which values of γ the means $B_R^{\lambda,\gamma}$ defined via $m_{\lambda,\gamma}$ converge a.e. for functions in L^{p_λ} , we study the maximal operator $B_*^{\lambda,\gamma}$. The following theorem is my main result.

Theorem 1.1. *Let $1 < 1 + 2\lambda < n$ and $0 \leq \mu < 2\gamma - 2$. Then there is a constant $C = C(n, \lambda, \gamma, \mu)$ such that*

$$(1) \quad \int_{\mathbb{R}^n} |B_*^{\lambda,\gamma}(f)(x)|^2 dx \leq C \int_{\mathbb{R}^n} |f(x)|^2 dx$$

for all $f \in L^2(\mathbb{R}^n, dx)$ and

$$(2) \quad \int_{\mathbb{R}^n} |B_*^{\lambda,\gamma}(f)(x)|^2 w_{\lambda,\mu}(x) dx \leq C \int_{\mathbb{R}^n} |f(x)|^2 w_{\lambda,\mu}(x) dx$$

for all $f \in L^2(\mathbb{R}^n, w_{\lambda,\mu}(x) dx)$, where $w_{\lambda,\mu} = \omega_{\lambda,\mu}(|x|)$ and

$$(3) \quad \omega_{\lambda,\mu}(t) = \begin{cases} \frac{1}{t^{2\lambda+1}} & \text{if } 0 < t \leq 1, \\ \frac{1}{t^{2\lambda+1}(\log(et))^\mu} & \text{if } t > 1. \end{cases}$$

For $(2\lambda + 1)/n < \mu$, we also have $L^{p_\lambda} \subseteq L^2 + L^2(w_{\lambda,\mu})$. Hence:

Corollary 1.2. *If $1 < 1 + 2\lambda < n$, $f \in L^{p_\lambda}(\mathbb{R}^n)$, and $\gamma > 1/p'_\lambda + 1/2$, we have*

$$(4) \quad \lim_{R \rightarrow \infty} B_R^{\lambda,\gamma}(f)(x) = f(x)$$

for almost every $x \in \mathbb{R}^n$. If $f \in L^p(\mathbb{R}^n)$ for $2 \leq p < p_\lambda$, then the condition $\gamma \geq 0$ suffices for (4) to hold.

When I first proved this result, it was natural to wonder whether the condition $\gamma > 1/p'_\lambda + 1/2$ was sharp. Lee, Rogers, and Seeger [Lee et al. 2014] have since proved among other things that, if

$$\frac{2(n+1)}{n-1} < p < \infty, \quad n \geq 2,$$

and $m \in B_{\alpha,q}^2$, then the maximal operator

$$M_m(f) := \sup_{t>0} |(\widehat{f} m(t|\cdot|))^\vee|$$

is bounded from $L^{p,q'}$ to L^p . This can be applied to $m = m_{\lambda,\gamma}$ to conclude that the condition $\gamma > 1/p'_\lambda + 1/2$ in [Corollary 1.2](#) can be replaced by $\gamma > 1/p'_\lambda$, if we further assume $(n-1)/(2(n+1)) < \lambda$.

Lee and Seeger [\[2015\]](#) have gone much further, proving that a.e. convergence of

$$S_t(f) := (\widehat{f} m_{\lambda,\gamma} \circ \rho(t(\cdot)))^\vee$$

to f (where ρ is an arbitrary homogeneous “distance” function, that is a homogeneous function that satisfies $\rho(\xi) > 0$ if $\xi \in \mathbb{R}^n \setminus \{0\}$ and $\rho(0) = 0$) holds for every $f \in L^{p,\lambda,q}$ when $q \geq 1$ if and only if $\gamma > 1/q'$, for all $0 < \lambda < (n-1)/2$. For $q = p_\lambda$ and $\rho(\xi) = |\xi|$, this implies [Corollary 1.2](#). In particular, they proved that the condition $\gamma > 1/q'$ is sharp.

The sufficiency of the condition $\gamma > 1/q'$ in [\[Lee and Seeger 2015\]](#) is presented as a consequence of a boundedness estimate between appropriate homogeneous Herz spaces — see [\[Baernstein and Sawyer 1985; Gilbert 1972\]](#) — of a maximal operator defined via an arbitrary quasiradial multiplier $h \circ \rho$, provided that h lies in an appropriate Besov space. A particular case of the same theorem also implies a characterization of boundedness for certain convolution operators on L^2 spaces that are weighted with power weights. In order to prove the sufficiency of the condition on γ , both of our papers use the approach of [\[Carbery et al. 1988\]](#), to some extent. However, much of my work is necessary to deal with the weight $w_{\lambda,\mu}$, that isn’t homogenous. The first choice of Lee and Seeger was to keep working with a homogeneous weight, but to use the observation that, for $p > 2$, the space $L^{p,2}$ is embedded in $L^2(|x|^{-n(1-2/p)} dx)$. By sharpening the analysis in [\[loc. cit.\]](#), this idea would only have yielded their result for $q = 2$. They solved the problem for all q by using Herz spaces, embedding theorems, and innovations that were needed to work with a more general “distance” function ρ and multiplier h .

The necessity of the condition $\gamma > 1/q'$ starts with the reminder that the operators S_t ($t > 0$) are naturally defined on the Schwartz class \mathcal{S} and extended on bigger spaces by using density. So, they proved that each operator S_t is continuous from \mathcal{S} — equipped with the $L^{p,\lambda,q}$ norm and topology — to \mathcal{S}' only if $\gamma > 1/q'$.

This paper. The proof of [Theorem 1.1](#) follows closely the idea developed in [\[Carbery et al. 1988\]](#), but accounts for the necessity to work with nonhomogeneous weights.

In [Section 2](#), [Theorem 1.1](#) is reduced to [Lemma 2.1](#), which is in turn reduced to [Lemma 5.2](#) in [Section 5](#). [Lemma 5.2](#) is proved in [Section 6](#).

In [Section 3](#), an upper bound is given for the Fourier transform of $w_{\lambda,\mu}^{(1)}$, which is $w_{\lambda,\mu}$ smoothened in a neighborhood of the spherical surface $\|x\| = 1$. An

analytic continuation argument is needed to prove that the upper bound holds for all $0 < \lambda < (n-1)/2$. This upper bound will be used to prove [Lemma 5.2](#).

In [Section 4](#), a new weight $\tilde{w}_{N,\lambda,\mu}$ is exhibited that is comparable to $1/w_{\lambda,\mu}$ and that has an algebraic form needed in the computations of [Section 5](#).

In [Section 5](#), [Lemma 2.1](#) is reduced to [Lemma 5.2](#). [Lemma 5.2](#) contains weighted Fourier inequalities for the special weight used in this paper. It is crucial that the “constants” appearing in both such inequalities have a certain functional form with respect to the parameter t . So, general results such as those in [[Benedetto and Heinig 2003](#)] were not sufficient.

[Section 6](#) contains the proofs of [Lemma 5.2](#) and [Corollary 1.2](#).

We shall refer to [[Carbery et al. 1988](#)] for every piece of the proof that doesn't differ significantly. Yet, the reader can find more details of the proof contained in that reference in [[Grafakos 2014](#), Subsection 10.5.2].

2. Reduction of [Theorem 1.1](#) to [Lemma 2.1](#)

We will only need to show (2), as the proof of (1) is contained in [[Grafakos 2014](#)] for the case $\gamma = 0$ (which implies it for all $\gamma \geq 0$). Let φ, ψ be smooth functions, supported in $[-\frac{1}{2}, \frac{1}{2}]$ and $[\frac{1}{8}, \frac{5}{8}]$ respectively, with values in $[0, 1]$, that satisfy

$$\varphi(t) + \sum_{k=0}^{\infty} \psi\left(\frac{1-t}{2^{-k}}\right) = 1$$

for all $t \in [0, 1)$. Let $e_1 = (1, 0, \dots, 0) \in \mathbb{R}^n$. We define $m_{\lambda,\gamma,00}(t) = m_{\lambda,\gamma}(te_1) \varphi(t)$ and

$$(5) \quad m_{\lambda,\gamma,k}(t) = 2^{k\lambda} m_{\lambda,\gamma}(te_1) \psi\left(\frac{1-t}{2^{-k}}\right), \quad k = 0, 1, 2, \dots$$

We define $\tilde{m}_{\lambda,\gamma,k}, (S_{\lambda,\gamma,k})_t, (S_{\lambda,\gamma,k})_*$, and $G_{\lambda,\gamma,k}$ from $m_{\lambda,\gamma,k}$, analogous to how $\tilde{m}^\delta, S_t^\delta, S_*^\delta$, and G^δ were defined from m^δ in [[Carbery et al. 1988](#)]. Similarly, we also define $(\tilde{S}_{\lambda,\gamma,k})_t, (\tilde{S}_{\lambda,\gamma,k})_*$, and $\tilde{G}_{\lambda,\gamma,k}$ by using $\tilde{m}_{\lambda,\gamma,k}$ instead of $m_{\lambda,\gamma,k}$. For $m_{\lambda,\gamma,k}$ we have the estimate

$$(6) \quad \sup_{0 \leq t \leq 1} \left| \frac{d^\ell}{dt^\ell} m_{\lambda,\gamma,k}(t) \right| \leq C_{\lambda,\gamma,\ell} \frac{2^{k\ell}}{k^\gamma}$$

for all $\ell \in \mathbb{Z}^+ \cup \{0\}$. As in [[loc. cit.](#)], these inequalities follow:

$$(7) \quad \|B_*^{\lambda,\gamma}\| \leq \|(S_{\lambda,\gamma,00})_*\| + \sum_{k=0}^{\infty} 2^{-k\lambda} \|(S_{\lambda,\gamma,k})_*\|,$$

$$(8) \quad \|(S_{\lambda,\gamma,k})_*(f)\|_{L^2(w_{\lambda,\mu})} \leq 2^{k+1} \|G_{\lambda,\gamma,k}(f)\|_{L^2(w_{\lambda,\mu})} \|\tilde{G}_{\lambda,\gamma,k}(f)\|_{L^2(w_{\lambda,\mu})}.$$

By reasoning as in [[loc. cit.](#)], one then shows without difficulty that the right-hand side in (8) can be controlled by the left-hand side of the inequality in the result we

are about to state:

Lemma 2.1. *For $k > 4$ we have*

$$\int_{\mathbb{R}^n} \int_1^2 |(S_{\lambda,\gamma,k})_{at}(f)(x)|^2 \frac{dt}{t} w_{\lambda,\mu}(x) dx \leq C_{n,\lambda,\mu,\gamma,k} \int_{\mathbb{R}^n} |f(x)|^2 w_{\lambda,\mu}(x) dx$$

for all $a > 0$ and for all functions f in $L^2(w_{\lambda,\mu})$, with

$$C_{n,\lambda,\mu,\gamma,k} = C_{n,\lambda,\mu,\gamma} \frac{2^{k(2\lambda-1)}}{k^{2\gamma-\mu}}.$$

We need not to worry about $k \leq 4$ because it is easily verified that $w_{\lambda,\mu}$ is an A_2 weight under the conditions of [Theorem 1.1](#) and therefore

$$\|(S_{\lambda,\gamma,k})_*(f)\|_{L^2(w_{\lambda,\mu})}^2 < \infty$$

for every k . Inequality (8) and [Lemma 2.1](#) then imply:

$$(9) \quad \|(S_{\lambda,\gamma,k})_*\|_{L^2(w_{\lambda,\mu}) \rightarrow L^2(w_{\lambda,\mu})} \leq C'(n, \lambda, \gamma) \left(\frac{2^{2k\lambda}}{k^{2\gamma-\mu}} \right)^{1/2}.$$

So, the right-hand side of (7) is finite if $\mu < 2\gamma - 2$. [Theorem 1.1](#) is now proved modulo [Lemma 2.1](#).

3. An upper bound for $|\widehat{w}_{\lambda,\mu}|$

The main result of this section will be used in [Section 6](#). Let $\theta \in C^\infty(\mathbb{R})$ satisfy $0 \leq \theta \leq 1$, $\text{supp}(\theta) \subset [\frac{9}{10}, \frac{11}{10}]$, $\theta \equiv 1$ on $[\frac{19}{20}, \frac{21}{20}]$. Now define

$$(10) \quad \omega_{\lambda,\mu}^{(1)}(t) = \omega_{\lambda,\mu}(t)(1 - \theta(t)) + \theta(t).$$

and $w_{\lambda,\mu}^{(1)}(x) = \omega_{\lambda,\mu}^{(1)}(|x|)$ for all $x \in \mathbb{R}^n \setminus \{0\}$. Then $w_{\lambda,\mu}^{(1)}$ is smooth on $\mathbb{R}^n \setminus \{0\}$ and $w_{\lambda,\mu}^{(1)} \approx_{\lambda,\mu} w_{\lambda,\mu}$; that is, $w_{\lambda,\mu}^{(1)}(x)$ and $w_{\lambda,\mu}(x)$ are comparable with comparability constant depending on λ and μ only. The goal of this section is to prove this result:

Theorem 3.1. *Let $w_{\lambda,\mu}$ and $w_{\lambda,\mu}^{(1)}$ be defined as above. Then for every λ satisfying $\frac{n-1}{4} < \lambda < \frac{n-1}{2}$ and every $\mu \geq 0$ there exists a constant $C_{n,\lambda,\mu}$ such that*

$$(11) \quad |\widehat{w}_{\lambda,\mu}(\xi)| \leq \Omega_{\lambda,\mu}(\xi) := \begin{cases} C_{n,\lambda,\mu} \frac{1}{|\xi|^{n-2\lambda-1} (\log \frac{e}{|\xi|})^\mu} & \text{if } |\xi| \leq 1, \\ C_{n,\lambda,\mu} \frac{1}{|\xi|^{n-2\lambda-1}} & \text{if } |\xi| \geq 1, \end{cases}$$

and, for all λ satisfying $0 < \lambda < \frac{n-1}{2}$ and μ as above, there exists a constant $C'_{n,\lambda,\mu}$ such that

$$(12) \quad |\widehat{w}_{\lambda,\mu}^{(1)}(\xi)| \leq C'_{n,\lambda,\mu} \Omega_{\lambda,\mu}(\xi)$$

for all $\xi \in \mathbb{R}^n \setminus \{0\}$.

Proof. We begin with the proof of (11) As $w_{\lambda,\mu}$ is radial, its Fourier transform is given by

$$\widehat{w}_{\lambda,\mu}(\xi) = \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} \int_0^\infty \omega_{\lambda,\mu}(r) J_{\frac{n-2}{2}}(2\pi|\xi|r) r^{\frac{n}{2}} dr,$$

where J_k denotes the k -th Bessel function. It is well known — see [Watson 1944] — that $|J_k(r)| \leq C_k r^k$ when $r \leq 2\pi$ and $|J_k(r)| \leq C_k r^{-\frac{1}{2}}$ when $r \geq 2\pi$. We control $|\widehat{w}_{\lambda,\mu}(\xi)|$ in two cases:

Case 1: $\frac{1}{|\xi|} \leq 1$. Then

$$\begin{aligned} |\widehat{w}_{\lambda,\mu}(\xi)| \leq C_n \left(\int_0^{\frac{1}{|\xi|}} r^{-2\lambda-1+\frac{n-2}{2}+\frac{n}{2}} dr \right) &+ \frac{C_n}{|\xi|^{\frac{n-1}{2}}} \left(\int_{\frac{1}{|\xi|}}^1 r^{-2\lambda-1-\frac{1}{2}+\frac{n}{2}} dr \right) \\ &+ \frac{C_n}{|\xi|^{\frac{n-1}{2}}} \left(\int_1^\infty \frac{r^{-2\lambda-1-\frac{1}{2}+\frac{n}{2}}}{(\log(er))^\mu} dr \right). \end{aligned}$$

Case 2: $\frac{1}{|\xi|} \geq 1$. Then

$$\begin{aligned} |\widehat{w}_{\lambda,\mu}(\xi)| \leq C_n \left(\int_0^1 r^{\frac{n-2}{2}+\frac{n}{2}-2\lambda-1} dr \right) \\ + C_n \left(\int_1^{\frac{1}{|\xi|}} \frac{1}{(\log(er))^\mu} r^{\frac{n-2}{2}+\frac{n}{2}-2\lambda-1} dr \right) \\ + \frac{C_n}{|\xi|^{\frac{n-1}{2}}} \left(\int_{\frac{1}{|\xi|}}^\infty \frac{1}{(\log(er))^\mu} r^{-\frac{1}{2}+\frac{n}{2}-2\lambda-1} dr \right). \end{aligned}$$

If $\lambda > \frac{n-1}{4}$ and $\lambda < \frac{n-1}{2}$, all integrals converge and (11) easily follows by using calculus.

The same holds with $w_{\lambda,\mu}$ replaced by $w_{\lambda,\mu}^{(1)}$ and the proof is almost identical. Then, an analytic continuation argument and the smoothness of $w_{\lambda,\mu}^{(1)}$ can be used to prove that (12) holds in the bigger range $0 < \lambda < \frac{n-1}{2}$. The argument involves many details that we omit but that may be split in two pieces.

In the first one, given any $\lambda' \in (0, \frac{n-1}{4}]$, we use more asymptotic estimates of the Bessel functions — see [Watson 1944] — and iterated integration by parts to rewrite the right-hand side of

$$(13) \quad \widehat{w}_{\lambda,\mu}^{(1)}(\xi) = \frac{2\pi}{|\xi|^{\frac{n-2}{2}}} \int_0^\infty \omega_{\lambda,\mu}^{(1)}(r) J_{\frac{n-2}{2}}(2\pi|\xi|r) r^{\frac{n}{2}} dr$$

in a way that also is well defined when λ ranges in a complex neighborhood $\mathcal{O}_{\lambda'}$ of the real interval $(\lambda', \frac{n-1}{2})$. We can call such extension $\tilde{w}_{\lambda,\mu}(\xi)$, and show that

$$|\tilde{w}_{\lambda,\mu}(\xi)| \leq C'_{n,\lambda,\mu} \Omega_{\lambda,\mu}(\xi),$$

as in (12).

In the second one, for the same value of λ' and the same neighborhood $\mathcal{O}_{\lambda'}$, we use the dominated convergence theorem to prove that, for a given test function φ defined on \mathbb{R}^n , the right-hand side of

$$(14) \quad \int_{\mathbb{R}^n} \widehat{\varphi}(\xi) w_{\lambda,\mu}^{(1)}(\xi) d\xi = \int_{\mathbb{R}^n} \varphi(\xi) \tilde{u}_{\lambda,\mu}(\xi) d\xi,$$

rewritten after the first piece of the argument, is holomorphic, hence analytic, on $\mathcal{O}_{\lambda'}$. It can be proved easily that the left-hand side of (14) is also analytic on $\mathcal{O}_{\lambda'}$. Since (14) holds when $\lambda \in (\frac{n-1}{4}, \frac{n-1}{2})$, we conclude from the analytic continuation theorem that (14) also holds when $\lambda \in (\lambda', \frac{n-1}{2})$. Then $\widehat{w}_{\lambda,\mu}^{(1)} = \tilde{u}_{\lambda,\mu}$, since φ is arbitrary. The arbitrariness of λ' concludes the proof. \square

4. A useful weight comparable to $1/w_{\lambda,\mu}$

In this section we show that $1/w_{\lambda,\mu}$ is comparable to another weight which can be written in a more useful way for our purposes, a fact that will be used in the next section. More precisely, let $u_{\lambda,\mu}$ and $\tilde{w}_{N,\lambda,\mu}$ be defined by:

$$(15) \quad u_{\lambda,\mu}(y) = \begin{cases} |y|^{-n-2\lambda-1} (\log \frac{e}{|y|})^\mu & \text{if } |y| < 1, \\ |y|^{-n-2\lambda-1} & \text{if } |y| \geq 1. \end{cases}$$

$$(16) \quad \tilde{w}_{N,\lambda,\mu}(x) = \int_{\mathbb{R}^n} |e^{i\langle x,y \rangle} - 1|^N u_{\lambda,\mu}(y) dy,$$

where N is a large enough integer independent of x .

The goal of this section is to prove that there exist constants $C_{1,n,\lambda,\mu,N}$ and $C_{2,n,\lambda,\mu,N}$ such that

$$(17) \quad \frac{C_{1,n,\lambda,\mu,N}}{w_{\lambda,\mu}(x)} \leq \tilde{w}_{N,\lambda,\mu}(x) \leq \frac{C_{2,n,\lambda,\mu,N}}{w_{\lambda,\mu}(x)}$$

for all $x \in \mathbb{R}^n \setminus \{0\}$. Let us write $\tilde{w}_{N,\lambda,\mu} = \tilde{w}_{N,\lambda,\mu,1} + \tilde{w}_{N,\lambda,\mu,2}$, where

$$(18) \quad \tilde{w}_{N,\lambda,\mu,1}(x) = \int_{|y| \leq \frac{1}{|x|}} |e^{i\langle x,y \rangle} - 1|^N u_{\lambda,\mu}(y) dy,$$

$$(19) \quad \tilde{w}_{N,\lambda,\mu,2}(x) = \int_{|y| > \frac{1}{|x|}} |e^{i\langle x,y \rangle} - 1|^N u_{\lambda,\mu}(y) dy.$$

Observe that in (18),

$$(20) \quad C_1 | \langle x, y \rangle | \leq |e^{i\langle x,y \rangle} - 1| \leq |x| |y|$$

for an absolute constant $0 < C_1 < 1$. Now, we estimate $\tilde{w}_{N,\lambda,\mu,1}$.

Case 1: $\frac{1}{|x|} \leq 1$. Given a positive constant $C > 0$, in view of (20),

$$\begin{aligned} \tilde{w}_{N,\lambda,\mu,1}(x) &= \int_{|y| \leq \frac{1}{|x|}} |e^{i\langle x,y \rangle} - 1|^N |y|^{-n-2\lambda-1} \left(\log \frac{e}{|y|} \right)^\mu dy \\ &\geq C_{C,N} \int_{\Omega(x)} |x|^N |y|^N |y|^{-n-2\lambda-1} \left(\log \frac{e}{|y|} \right)^\mu dy \\ &= C_{n,C,N} |x|^N \frac{|S^{n-1}|}{e^{2\lambda+1-N}} \int_{e^{|x|}}^\infty s^{2\lambda-N} (\log s)^\mu ds, \end{aligned}$$

where $\Omega(x) = \{y : |y| \leq \frac{1}{|x|} \text{ and } C \leq |\langle \frac{x}{|x|}, \frac{y}{|y|} \rangle|\}$. In order for this integral to converge, we need $N > 2\lambda + 1$. Later we will also need N to be even. So, we set $N = N_\lambda := 2\lceil 2\lambda + 1 \rceil$. It easily follows that there exist constants $C_{n,\lambda,N}$ and $C_{\lambda,\mu,N}$ such that $\tilde{w}_{N,\lambda,\mu,1}(x) \geq C_{n,\lambda,N}/w_{\lambda,\mu}(x)$ for all $x \in \mathbb{R}^n$ satisfying $|x| \geq C_{\lambda,\mu,N}$. An easier computation and (20) yield $\tilde{w}_{N,\lambda,\mu,1}(x) \leq C'_{n,\lambda,N}/w_{\lambda,\mu}(x)$ in **Case 1** for all $x \in \mathbb{R}^n$ satisfying $|x| \geq C_{\lambda,\mu,N}$. So, on $\{x \in \mathbb{R}^n : |x| \geq \max\{1, C_{\lambda,\mu,N}\}\}$ we have

$$(21) \quad \tilde{w}_{N,\lambda,\mu,1} \approx_{n,\lambda,N} 1/w_{\lambda,\mu}$$

Case 2: $\frac{1}{|x|} > 1$. Let us use the decomposition $\tilde{w}_{N,\lambda,\mu,1}(x) = I + II$, where

$$\begin{aligned} I &= \int_{|y| \leq 1} |e^{i\langle x,y \rangle} - 1|^N |y|^{-n-2\lambda-1} \left(\log \frac{e}{|y|} \right)^\mu dy \approx_{n,\lambda,\mu,N} |x|^N, \\ II &= \int_{1 < |y| \leq \frac{1}{|x|}} |e^{i\langle x,y \rangle} - 1|^N |y|^{-n-2\lambda-1} dy \approx_{n,\lambda,N} |x|^{2\lambda+1}, \end{aligned}$$

This proves that

$$(22) \quad \tilde{w}_{N,\lambda,\mu,1}(x) \approx_{n,\lambda,\mu,N} |x|^N + |x|^{2\lambda+1} \approx_{n,\lambda,N} |x|^{2\lambda+1} = \frac{1}{w_{\lambda,\mu}(x)}$$

on $\{x \in \mathbb{R}^n : |x| \leq 1\}$. If $C_{\lambda,\mu,N} \leq 1$, then relations (21) and (22) immediately imply that $\tilde{w}_{N,\lambda,\mu,1} \approx_{n,\lambda,\mu,N} 1/w_{\lambda,\mu}$ on \mathbb{R}^n . Otherwise, just observe that both functions $\tilde{w}_{N,\lambda,\mu,1}$ and $1/w_{\lambda,\mu}$ are positive and continuous on the compact annulus $1 \leq |x| \leq C_{\lambda,\mu,N}$. We still have to show that $\tilde{w}_{N,\lambda,\mu,2} \approx_{n,\lambda,\mu,N} 1/w_{\lambda,\mu}$. Let us define

$$(23) \quad \tilde{\tilde{w}}_{\lambda,\mu,2}(x) = \int_{|y| > \frac{1}{|x|}} u_{\lambda,\mu}(y) dy.$$

Then

$$(24) \quad \tilde{w}_{N,\lambda,\mu,2}(x) \leq 2^N \tilde{\tilde{w}}_{\lambda,\mu,2}(x).$$

We will prove that the inverse inequality also holds (with a constant different from 2^N), so that we have $\tilde{w}_{\lambda,\mu,2} \approx_{N,\lambda,\mu,n} \tilde{w}_{N,\lambda,\mu,2}$. Now, let us prove that

$$\tilde{w}_{\lambda,\mu,2} \approx_{N,\lambda,\mu,n} 1/w_{\lambda,\mu}.$$

Case 1: $\frac{1}{|x|} > 1$. Then:

$$\tilde{w}_{\lambda,\mu,2}(x) = \frac{|S^{n-1}|}{2\lambda + 1} |x|^{2\lambda+1} \approx_{\lambda,n} |x|^{2\lambda+1} = \frac{1}{w_{\lambda,\mu}(x)}.$$

Case 2: $\frac{1}{|x|} \leq 1$. Then:

$$\begin{aligned} \tilde{w}_{\lambda,\mu,2}(x) &= C_{n,\lambda} + \frac{|S^{n-1}|}{e^{2\lambda+1}} \int_e^{e^{|x|}} t^{2\lambda} (\log t)^\mu dt \\ &\approx_{\lambda,\mu,n} |x|^{2\lambda+1} (\log(e|x|))^\mu = \frac{1}{w_{\lambda,\mu}(x)}. \end{aligned}$$

This concludes the proof that $\tilde{w}_{\lambda,\mu,2} \approx_{\lambda,\mu,n} 1/w_{\lambda,\mu}$ on $\mathbb{R}^n \setminus \{0\}$. Now we need to prove that there exists a constant $C_{N,\lambda,\mu,n}$ such that the inequality

$$\tilde{w}_{\lambda,\mu,2} \leq C_{N,\lambda,\mu,n} \tilde{w}_{N,\lambda,\mu,2}$$

holds on $\mathbb{R}^n \setminus \{0\}$. Since both $\tilde{w}_{\lambda,\mu,2}$ and $\tilde{w}_{N,\lambda,\mu,2}$ are radial, it will be enough to prove that the functions $t \mapsto \tilde{w}_{N,\lambda,\mu,2}(te_1)$ and $t \mapsto \tilde{w}_{\lambda,\mu,2}(te_1)$ are comparable on \mathbb{R}^+ , where $e_1 = (1, 0, \dots, 0)$. Observe that $|e^{i\langle te_1, y \rangle} - 1| > \sqrt{2}$ on $G^t := \bigcup_{k \in \mathbb{Z}} G_k^t$, where

$$G_k^t := \left\{ y \in \mathbb{R}^n : \langle e_1, y \rangle \in \left[\frac{(4k+1)\pi}{2t}, \frac{(4k+3)\pi}{2t} \right) \right\}$$

for all $t > 0$ and $k \in \mathbb{Z}$. Therefore $u_{\lambda,\mu}(y) \approx_N |e^{i\langle te_1, y \rangle} - 1|^N u_{\lambda,\mu}(y)$ on G^t . In particular, there exists a constant C_N such that

$$\int_{G^t} u_{\lambda,\mu}(y) dy \leq C_N \int_{G^t} |e^{i\langle te_1, y \rangle} - 1|^N u_{\lambda,\mu}(y) dy.$$

If $t > 0$ and $k \in \mathbb{Z} \setminus \{0\}$ we define

$$R_k^t := \left\{ y \in \mathbb{R}^n : \langle e_1, y \rangle \in \left[\frac{(4k-1)\pi}{2t}, \frac{(4k+1)\pi}{2t} \right) \right\}$$

and

$$R_0^t := \left\{ y \in \mathbb{R}^n : \langle e_1, y \rangle \in \left[\frac{-\pi}{2t}, \frac{\pi}{2t} \right) \text{ and } |y| > \frac{1}{t} \right\}.$$

As

$$\int_{R_k^t} u_{\lambda,\mu}(y) dy \leq \int_{G_{k-1}^t} u_{\lambda,\mu}(y) dy$$

for all $k \in \mathbb{Z}^+$, and

$$\int_{R_k^t} u_{\lambda,\mu}(y) dy \leq \int_{G_k^t} u_{\lambda,\mu}(y) dy$$

for all $k \in \mathbb{Z}^-$, we also have

$$\int_{\bigcup_{k \in \mathbb{Z} \setminus \{0\}} R_k^t} u_{\lambda, \mu}(y) \, dy \leq \int_{G^t} u_{\lambda, \mu}(y) \, dy \leq C_N \int_{G^t} |e^{i\langle te_1, y \rangle} - 1|^{2N} u_{\lambda, \mu}(y) \, dy.$$

Since

$$\left\{ y : |\langle e_1, y \rangle| > \frac{\pi}{2t} \right\} = G^t \cup \bigcup_{k \in \mathbb{Z} \setminus \{0\}} R_k^t,$$

we have

$$\int_{|\langle e_1, y \rangle| > \frac{\pi}{2t}} u_{\lambda, \mu}(y) \, dy \leq 2C_N \int_{|y| > \frac{1}{t}} |e^{i\langle te_1, y \rangle} - 1|^{2N} u_{\lambda, \mu}(y) \, dy.$$

Since $u_{\lambda, \mu}$ is radial, we can replace e_1 by e_j in the inequality above for $j = 2, \dots, n$. Let $|y|_\infty := \sup_{1 \leq j \leq n} |\langle e_j, y \rangle|$. Then

$$(25) \quad \int_{\{y \in \mathbb{R}^n : |y|_\infty > \frac{\pi}{2t}\}} u_{\lambda, \mu}(y) \, dy \leq 2n C_N \int_{|y| > \frac{1}{t}} |e^{i\langle te_1, y \rangle} - 1|^{2N} u_{\lambda, \mu}(y) \, dy.$$

Inequality (25) and the Lemma 4.1 easily imply — see (19) and (23) for details — that $\tilde{w}_{\lambda, \mu, 2}(te_1) \leq C_{n, \lambda, \mu, N} \cdot \tilde{w}_{N, \lambda, \mu, 2}(te_1)$.

Lemma 4.1. *Let $u_{\lambda, \mu}$ be as in (15). Then, for all $n \in \mathbb{Z}^+$, $\lambda \in \mathbb{R}$, and $C > 1$ there exists a constant $D = D(n, \lambda, C) \in \mathbb{R}$ such that $u_{\lambda, \mu}(\frac{y}{C}) \leq D u_{\lambda, \mu}(y)$ for all $y \in \mathbb{R}^n \setminus \{0\}$. We can choose $D = C^{n+2\lambda+1} (\log(eC))^\mu$.*

The proof of Lemma 4.1 is left to the reader. This completes the proof that $\tilde{w}_{N, \lambda, \mu, 2} \approx_{n, \lambda, \mu, N} \frac{1}{w_{\lambda, \mu}}$ on $\mathbb{R}^n \setminus \{0\}$ and therefore the proof of (17), that is the claim of this section.

5. Reduction of Lemma 2.1 to Lemma 5.2

By duality, the inequality in Lemma 2.1 can be expressed as

$$(26) \quad \left\| \int_1^2 (S_{\lambda, \gamma, k})_{at}(h(t, \cdot))(x) \frac{dt}{t} \right\|_{L^2(\frac{dx}{w_{\lambda, \mu}(x)})} \leq C \|h(t, x)\|_{L^2(\frac{dt}{t} \frac{dx}{w_{\lambda, \mu}(x)})}$$

for all functions $h(t, x)$ in the appropriate space, where

$$C = C_{n, \lambda, \mu, \gamma, k} = \sqrt{C_{n, \lambda, \mu, \gamma} \frac{2^{k(2\lambda-1)}}{k^{2\gamma-\mu}}}.$$

In view of the result of Section 4, for every $f \in L^2(\mathbb{R}^n, \frac{1}{w_{\lambda, \mu}})$,

$$(27) \quad \|f\|_{L^2(\frac{dx}{w_{\lambda, \mu}(x)})}^2 \approx \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} u_{\lambda, \mu}(y) \left| \left(\sum_{j=0}^{N_\lambda/2} \hat{f}(g_{j, y}(\xi)) b_j \right) \right|^2 dy d\xi,$$

where

$$g_{j,y}(\xi) = \left(\xi - \left(\frac{N_\lambda/2 - j}{2\pi} \right) y \right), \quad b_j = (-1)^j \binom{N_\lambda/2}{j},$$

Plancherel’s identity was used, and the implicit comparability constants depend on λ, μ, n only. We can substitute the left-hand side of (26) by using (27) on the function

$$f(x) = \int_1^2 (S_{\lambda,\gamma,k})_{at}(h(t, \cdot))(x) \frac{dt}{t}.$$

For such a function we have

$$(28) \quad \left| \left(\sum_{j=0}^{N_\lambda/2} \widehat{f}(g_{j,y}(\xi)) b_j \right) \right|^2 = \left| \int_1^2 \left(\sum_{j=0}^{N_\lambda/2} \widehat{h}(t, g_{j,y}(\xi)) m_{\lambda,\gamma,k}(at|g_{j,y}(\xi)|) b_j \right) \frac{dt}{t} \right|^2.$$

Since $m_{\lambda,\gamma,k}$ is supported in $\left[1 - \frac{5}{8 \cdot 2^k}, 1 + \frac{1}{8 \cdot 2^k} \right]$, the Cauchy–Schwarz inequality in the t variable allows us to control the right-hand side of (28) by

$$(29) \quad \frac{C_\lambda}{2^k} \int_1^2 \left| \sum_{j=0}^{N_\lambda/2} \widehat{h}(t, g_{j,y}(\xi)) \cdot m_{\lambda,\gamma,k}(at|g_{j,y}(\xi)|) b_j \right|^2 \frac{dt}{t} =: H_{k,\lambda,\gamma}(y, \xi).$$

So, if we can show

$$(30) \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} u_{\lambda,\mu}(y) H_{k,\lambda,\gamma}(y, \xi) dy d\xi \leq C \frac{2^{k(2\lambda-1)}}{k^{2\gamma-\mu}} \|h(t, x)\|_{L^2(\frac{dx}{t} \frac{dx}{\omega_{\lambda,\mu}(|x|)})}^2$$

for a constant $C := C_{n,\lambda,\mu,\gamma}$, then (26) is proved. But (30) follows from the following pointwise (with respect to t) estimate:

$$(31) \quad \|(S_{\lambda,\gamma,k})_t(h)(x)\|_{L^2(\frac{dx}{\omega_{\lambda,\mu}(|x|)})}^2 \leq C_{n,\lambda,\mu,\gamma} \frac{2^{2k\lambda}}{k^{2\gamma-\mu}} \|h\|_{L^2(\frac{dx}{\omega_{\lambda,\mu}(|x|)})}^2$$

if (31) holds for all $t > 0$ rather than just $t \in [1, 2]$ (which allowed us to drop the parameter a), and for all $h \in L^2(\mathbb{R}^n, dx/w_{\lambda,\mu}(x))$. In order to see that (31) implies (30), just use (27) with $f(x) = (\widehat{h}(\cdot) m_{\lambda,\gamma,k}(t|\cdot|))^\vee(x) = (S_{\lambda,\gamma,k})_t(h)(x)$, to rewrite the left-hand side of (31).

By duality, (31) is equivalent to

$$(32) \quad \|(S_{\lambda,\gamma,k})_t(h)\|_{L^2(w_{\lambda,\mu})}^2 \leq C_{n,\lambda,\mu,\gamma} \frac{2^{2k\lambda}}{k^{2\gamma-\mu}} \|h\|_{L^2(w_{\lambda,\mu})}^2$$

for all $h \in L^2(w_{\lambda,\mu}(x) dx)$, $t > 0$. So, the latter also yields the inequality in Lemma 2.1 for every f in the appropriate space and every $a > 0$. We now need to prove (32).

We denote by $(K_{\lambda,\gamma,k})_t(x)$ the kernel of the operator $(S_{\lambda,\gamma,k})_t$, i.e., the inverse Fourier transform of the multiplier $m_{\lambda,\gamma,k}(t|\cdot|)$. $(K_{\lambda,\gamma,k})_t$ is radial on \mathbb{R}^n , and it is convenient to decompose it radially as

$$(K_{\lambda,\gamma,k})_t = (K_{\lambda,\gamma,k})_1^{(0)} + \sum_{j=1}^{\infty} (K_{\lambda,\gamma,k})_t^{(j)},$$

where

$$\begin{aligned} (K_{\lambda,\gamma,k})_1^{(0)}(x) &= (K_{\lambda,\gamma,k})_t(x) \theta(2^{-(k+3)}x/t), \\ (K_{\lambda,\gamma,k})_t^{(j)}(x) &= (K_{\lambda,\gamma,k})_t(x) (\theta(2^{-(j+k+3)}x/t) - \theta(2^{-(k+2+j)}x/t)), \end{aligned}$$

for some radial smooth function θ supported in the ball $B(0, 2)$ and equal to one on $B(0, 1)$.

To prove estimate (32) we make use of the subsequent lemmas.

Lemma 5.1. *For all $M \geq 2n$ there is a constant $C_{\lambda,\gamma,k,M} = C_{\lambda,\gamma,k,M}(n, \theta)$ such that for all $j = 0, 1, 2, \dots$,*

$$(33) \quad \sup_{\xi \in \mathbb{R}^n} |\widehat{(K_{\lambda,\gamma,k})_t^{(j)}}(\xi)| \leq C_{\lambda,\gamma,M} \frac{2^{-jM}}{k^\gamma}$$

and also

$$(34) \quad |\widehat{(K_{\lambda,\gamma,k})_t^{(j+l)}}(\xi)| \leq C_{\lambda,\gamma,M} \frac{2^{-(j+l)M}}{k^\gamma}$$

whenever $|t|\xi| - 1| \geq 2^{l-k-3}$ and $l \geq 4$. Also,

$$(35) \quad |\widehat{(K_{\lambda,\gamma,k})_t^{(j)}}(\xi)| \leq C_{\lambda,\gamma,M} \frac{2^{-(j+k+3)M}}{k^\gamma} (1 + t|\xi|)^{-M}$$

whenever $|t\xi| \leq \frac{1}{8}$ or $|t\xi| \geq \frac{15}{8}$.

Proof. The proof for $t = 1$ follows the lines of the proof of Lemma 10.5.5 in [Grafakos 2014, p. 413]. Just observe that estimate (10.5.9) in p. 409 of that reference is now replaced by (6), which explains why the factor $1/k^\gamma$ appears. The general case (any $t > 0$) is straightforward in view of the fact that

$$\widehat{(K_{\lambda,\gamma,k})_t^{(j)}}(\xi) = \widehat{(K_{\lambda,\gamma,k})_1^{(j)}}(t\xi). \quad \square$$

Lemma 5.2. *The inequalities*

$$(36) \quad \int_{|t\xi|-1| < \varepsilon} |\widehat{f}(\xi)|^2 d\xi \leq C_{n,\lambda,\mu} \omega_{\lambda,\mu}(t) \varepsilon \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)}$$

and

$$(37) \quad \int_{\mathbb{R}^n} |\widehat{f}(\xi)|^2 \frac{d\xi}{(1 + |t\xi|)^M} \leq C_{n,\lambda,\mu,M} \omega_{\lambda,\mu}(t) \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)},$$

hold for all Schwartz functions f , $t > 0$, $M \geq 2n$, all $0 < \varepsilon < 2$, λ , and μ as in Theorem 1.1.

The proof of Lemma 5.2 is postponed to Section 6.

By reasoning as in [Grafakos 2014, p. 414] and using the estimates in Lemmas 5.1 and 5.2 instead of those in Lemma 10.5.5 in [op. cit., p. 413] and Lemma 10.5.6 in [op. cit., p. 414], we can prove

$$(38) \quad \int_{\mathbb{R}^n} |((K_{\lambda,\gamma,k})_t^{(j)} * f)(x)|^2 dx \leq C \frac{2^{-2jM}}{2^k k^{2\gamma}} \omega_{\lambda,\mu}(t) \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)}$$

for another constant $C = C_{n,\lambda,\mu,\gamma,M}$. By duality, this is equivalent to

$$(39) \quad \int_{\mathbb{R}^n} |((K_{\lambda,\gamma,k})_t^{(j)} * f)(x)|^2 w_{\lambda,\mu}(x) dx \leq C \frac{2^{-2jM}}{2^k k^{2\gamma}} \omega_{\lambda,\mu}(t) \int_{\mathbb{R}^n} |f(x)|^2 dx.$$

Given a Schwartz function f , we write

$$f_0 = f \chi_{Q_0^{(n,k,j,t)}},$$

where $Q_0^{(n,k,j,t)}$ is a cube centered at the origin of side length $C_n 2^{j+k+4}t$ (note that $\text{supp} (K_{\lambda,\gamma,k})_t^{(j)} \subseteq B(0, 2^{j+k+4}t)$). Then for $x \in Q_0^{(n,k,j,t)}$ we have the inequality

$$|x| \leq \sqrt{n} C_n 2^{j+k+4}t;$$

hence, (39) implies

$$(40) \quad \int_{\mathbb{R}^n} |((K_{\lambda,\gamma,k})_t^{(j)} * f_0)(x)|^2 w_{\lambda,\mu}(x) dx \leq C_{n,\lambda,\mu,\gamma,M} \frac{2^{-2jM}}{2^k k^{2\gamma}} \frac{\omega_{\lambda,\mu}(t)}{\omega_{\lambda,\mu}(\sqrt{n} C_n 2^{j+k+4}t)} \int_{Q_0^{(n,k,j,t)}} |f_0(x)|^2 w_{\lambda,\mu}(x) dx$$

because the function $1/\omega_{\lambda,\mu}$ is increasing. A simple computation shows that

$$(41) \quad \sup_{t>0} \frac{\omega_{\lambda,\mu}(at)}{\omega_{\lambda,\mu}(t)} = \frac{1}{a^{2\lambda+1}} \quad \text{and} \quad \sup_{t>0} \frac{\omega_{\lambda,\mu}(at)}{\omega_{\lambda,\mu}(t)} = \frac{(\log(e/a))^\mu}{a^{2\lambda+1}}$$

if $a > 1$ and if $a \leq 1$, respectively. Therefore, for all j and k such that $j + k \geq C'_n$ for a suitable purely dimensional constant C'_n ,

$$(42) \quad \sup_{t>0} \frac{\omega_{\lambda,\mu}(t)}{\omega_{\lambda,\mu}(\sqrt{n} C_n 2^{j+k+4}t)} \leq C''_{n,\lambda,\mu} 2^{(j+k)(2\lambda+1)} (j^\mu + k^\mu),$$

where we used the hypothesis on j and k and the fact that

$$(j + k)^\mu \leq C_\mu (j^\mu + k^\mu).$$

It follows from (42) and (40) that $\int_{\mathbb{R}^n} |((K_{\lambda,\gamma,k})_t^{(j)} * f_0)(x)|^2 w_{\lambda,\mu}(x) dx$ is bounded by

$$C 2^{j(2\lambda+1-2M)} \frac{2^{2k\lambda}}{k^{2\gamma}} (j^\mu + k^\mu) \int_{Q_0^{(n,k,j,t)}} |f_0(x)|^2 w_{\lambda,\mu}(x) dx,$$

for $C = C_{n,\lambda,\mu,\gamma,M}$, provided that

$$(43) \quad j + k \geq C'_n.$$

Now write $\mathbb{R}^n \setminus Q_0^{(n,k,j,t)}$ as a mesh of cubes $Q_i^{(n,k,j,t)}$, indexed by $i \in \mathbb{Z} \setminus \{0\}$, of side lengths $C_n 2^{j+k+4} t$ (the same side length of $Q_0^{(n,k,j,t)}$) and centers c_{Q_i} . By using (33), reasoning as in [Grafakos 2014, p. 415] as well as simply noting that $2^{2k\lambda}(j^\mu + k^\mu) \geq 1$, we can find that the pieces

$$\int_{\mathbb{R}^n} |((K_{\lambda,\gamma,k})_t^{(j)} * f_i)(x')|^2 w_{\lambda,\mu}(x') dx'$$

are bounded by

$$C_{\lambda,\mu,\gamma,M} 2^{-2jM} \frac{2^{2k\lambda}}{k^{2\gamma}} (j^\mu + k^\mu) \int_{Q_i^{(n,k,j,t)}} |f_i(x)|^2 w_{\lambda,\mu}(x) dx$$

whenever f_i is supported in $Q_i^{(n,k,j,t)}$ and, in turn, that

$$(44) \quad \begin{aligned} \|((K_{\lambda,\gamma,k})_t^{(j)} * f)\|_{L^2(w_{\lambda,\mu})} \\ \leq C''_{n,\lambda,\mu,\gamma,M} 2^{j(\lambda+\frac{1}{2}-M)} \frac{2^{k\lambda}}{k^\gamma} (j^{\frac{\mu}{2}} + k^{\frac{\mu}{2}}) \|f\|_{L^2(w_{\lambda,\mu})} \end{aligned}$$

(in view of the argument in [Grafakos 2014]). Observe that condition (43) is satisfied if we assume $k \geq C'_n$, which we can as the convergence of (7) only depends on the estimates we have for k big enough. So, for $k \geq C'_n$, by using (44) and summing over $j = 0, 1, 2, \dots$, we deduce (32) if we just choose $M > n/2$ (remember that $n > 2\lambda + 1$). In turn, (32) is equivalent to (31), which is equivalent to (26), which is equivalent to the inequality in Lemma 2.1. Therefore, this completes the proof of the lemma, modulo Lemma 5.2

6. Proof of Lemma 5.2

6.1. Proof of inequality (36). We reduce estimate (36) by duality to

$$(45) \quad \int_{\mathbb{R}^n} |\widehat{g}(\xi)|^2 w_{\lambda,\mu}(\xi) d\xi \leq C_{n,\lambda,\mu} \omega_{\lambda,\mu}(t) \varepsilon \int_{||t x| - 1| \leq \varepsilon} |g(x)|^2 dx$$

for functions g supported in the annulus $||t x| - 1| \leq \varepsilon$. In Section 3 we observed that

$$w_{\lambda,\mu} \approx_{\lambda,\mu} w_{\lambda,\mu}^{(1)}$$

and proved in Theorem 3.1 that the function $|\widehat{w}_{\lambda,\mu}^{(1)}|$ is bounded by a scalar multiple of $\Omega_{\lambda,\mu}$ (see (12)) in the whole range $\lambda \in (0, (n-1)/2)$. Therefore, we can start

to prove (45) as follows:

$$\begin{aligned}
 (46) \quad \int_{\mathbb{R}^n} |\widehat{g}(\xi)|^2 w_{\lambda,\mu}(\xi) d\xi &\approx_{\lambda,\mu} \int_{\mathbb{R}^n} (\widehat{g} \widetilde{g})^\vee(x) \widehat{w}_{\lambda,\mu}^{(1)}(x) dx \\
 &\leq C_{n,\lambda,\mu} \int_{\mathbb{R}^n} (|g| * |\widetilde{g}|)(x) \Omega_{\lambda,\mu}(x) dx \\
 &= C_{n,\lambda,\mu} \iint_{\substack{||ty|-1| \leq \varepsilon \\ ||tx|-1| \leq \varepsilon}} |g(x)| |\widetilde{g}(y)| \Omega_{\lambda,\mu}(x-y) dx dy \\
 &\leq C_{n,\lambda,\mu} B(n, \lambda, \mu, \varepsilon, t) \|g\|_{L^2}^2
 \end{aligned}$$

where $\widetilde{g}(x) = g(-x)$ and

$$(47) \quad B(n, \lambda, \mu, \varepsilon, t) = \frac{1}{t^n} \sup_{\{x: ||x|-1| \leq \varepsilon\}} \int_{||y|-1| \leq \varepsilon} \Omega_{\lambda,\mu}^t(y-x) dy,$$

where $\Omega_{\lambda,\mu}^t(x) := \Omega_{\lambda,\mu}(x/t)$. The last inequality of (46) is proved by interpolation between the $L^1(S) \rightarrow L^1(S)$ and $L^\infty(S) \rightarrow L^\infty(S)$ estimates for the linear operator

$$L_{\lambda,\mu,t,\varepsilon}(g)(x) = \int_S g(y) \Omega_{\lambda,\mu}(y-x) dy,$$

where

$$S = \{y \in \mathbb{R}^n : ||ty|-1| \leq \varepsilon\},$$

using the Cauchy–Schwarz inequality. It remains to establish that

$$(48) \quad B(n, \lambda, \mu, \varepsilon, t) \leq C_{n,\lambda,\mu} \omega_{\lambda,\mu}(t) \varepsilon.$$

Then we reason as in [Grafakos 2014, pp. 417, 418]: we apply a rotation and a change of variable to the integrals in (47) to push the dependence on x to the domain of integration, then control the supremum in (47) by integrating $\Omega_{\lambda,\mu}$ over the bigger set

$$\{y : ||y-e_1|-1| \leq 2\varepsilon\},$$

finally we split this latter integral over the sets S_0, S_ℓ, S_* defined in [op. cit.] to be

$$\begin{aligned}
 S_0 &= \{y \in \mathbb{R}^n : ||y-e_1|-1| \leq 2\varepsilon, |y| \leq \varepsilon\}, \\
 S_\ell &= \{y \in \mathbb{R}^n : ||y-e_1|-1| \leq 2\varepsilon, \ell\varepsilon \leq |y| \leq (\ell+1)\varepsilon\}, \\
 S_* &= \{y \in \mathbb{R}^n : ||y-e_1|-1| \leq 2\varepsilon, |y| \geq 1\}.
 \end{aligned}$$

In the end, matters reduce to proving the estimates

$$(49) \quad \int_{S_0} \Omega_{\lambda,\mu}^t(y) dy \leq C'_{n,\lambda,\mu} t^n \omega_{\lambda,\mu}(t) \varepsilon^{2\lambda+1},$$

$$(50) \quad \sum_{\ell=1}^{\lceil \frac{1}{\varepsilon} \rceil + 1} \int_{S_\ell} \Omega_{\lambda,\mu}^t(y) dy \leq C_{\lambda,\mu} t^n \varepsilon \omega_{\lambda,\mu}(t),$$

$$(51) \quad \int_{S_*} \Omega_{\lambda,\mu}^t(y) dy \leq C_n \varepsilon \omega_{\lambda,\mu}(t) t^n.$$

In proving the inequalities above, we can assume without loss of generality that $t \geq 2$, because when $t < 2$ the proof of [Lemma 5.2](#) is an immediate consequence of [Lemma 10.5.6](#) in [\[op. cit., p. 414\]](#). We can also assume that $t \geq C_{n,\lambda,\mu}$, due to the compactness of $[2, C_{n,\lambda,\mu}]$ and the continuity and positivity of the functions involved. For a suitable constant $C_{n,\lambda,\mu}$ and $t \geq \max\{2, C_{n,\lambda,\mu}\}$, [\(49\)](#) is proved by using calculus (note that the integrand in [\(49\)](#) is radial and the domain of integration is a sphere); [\(50\)](#) is proved by using the maximum of the integrand over each set S_ℓ , then by comparing the sum with an integral, finally by using calculus to estimate the integral; [\(51\)](#) is proved by using the maximum of the integrand over S_* . The condition that $t \geq 2 > \varepsilon$ was used in both [\(49\)](#) and [\(50\)](#) and [\(41\)](#) was used in [\(51\)](#).

By combining estimates [\(49\)](#), [\(50\)](#), and [\(51\)](#), we obtain [\(48\)](#). This concludes the proof of [\(45\)](#) and, therefore, of [\(36\)](#). \square

6.2. Proof of inequality [\(37\)](#). Inequality [\(37\)](#) is already known for $t \leq 1$; see equation (10.5.22) in [\[Grafakos 2014, p. 414\]](#). Indeed, if $0 < t \leq 1$ then $\omega_{\lambda,\mu}(t) = 1/t^{2\lambda+1}$, and [\(37\)](#) follows by dilation from the case $t = 1$, the one shown in [\[op. cit.\]](#). For $t > 1$ define:

$$A_1^t = \left\{ \xi \in \mathbb{R}^n : |\xi| \leq \frac{1}{t} \right\}, \quad A_2^t = \left\{ \xi \in \mathbb{R}^n : \frac{1}{t} < |\xi| \leq \frac{2+\sqrt{t}}{t} \right\},$$

$$A_3^t = \left\{ \xi \in \mathbb{R}^n : \frac{2+\sqrt{t}}{t} < |\xi| \leq \frac{2+t}{t} \right\}, \quad A_4^t = \left\{ \xi \in \mathbb{R}^n : \frac{2+t}{t} < |\xi| \right\}.$$

We will prove [\(37\)](#) by proving that

$$(52) \quad I_j := \int_{A_j^t} |\widehat{f}(\xi)|^2 \frac{1}{(1+|t\xi|)^M} d\xi \leq C_{n,\lambda,\mu,M} \omega_{\lambda,\mu}(t) \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)}$$

for each $j = 1, 2, 3, 4$. For $j = 1$, first observe that $1/(1+|t\xi|)^M \approx_M 1$ on A_1^t and then argue as in the proof of [\(36\)](#), at the beginning of this section. By duality, we reduce [\(52\)](#) with $j = 1$ to

$$(53) \quad \int_{\mathbb{R}^n} |\widehat{f}(\xi)|^2 w_{\lambda,\mu}(\xi) d\xi \leq C_{n,\lambda,\mu} \omega_{\lambda,\mu}(t) \int_{A_1^t} |f(x)|^2 dx$$

for all functions f supported in the ball A_1^t . By proceeding as in (46), we can prove that

$$\int_{\mathbb{R}^n} |\widehat{f}(\xi)|^2 w_{\lambda,\mu}(\xi) d\xi \leq B'(n, \lambda, \mu, t) \|f\|_{L^2}^2$$

for every f supported in A_1^t , where $B'(n, \lambda, \mu, t)$, now, is defined by

$$\begin{aligned} (54) \quad B'(n, \lambda, \mu, t) &= \sup_{\{x:|x|\leq\frac{1}{t}\}} \int_{|y|\leq\frac{1}{t}} \Omega_{\lambda,\mu}(y-x) dy \\ &= \frac{1}{t^n} \sup_{\{x:|x|\leq 1\}} \int_{|y+x|\leq 1} \Omega_{\lambda,\mu}\left(\frac{y}{t}\right) dy \end{aligned}$$

and all we still need to show is that

$$(55) \quad B'(n, \lambda, \mu, t) \leq C_{n,\lambda,\mu} \omega_{\lambda,\mu}(t).$$

Since $|x| \leq 1$ and $|x + y| \leq 1$ we have $|y| \leq 2$. So, (55) is a consequence of

$$(56) \quad \frac{1}{t^n} \int_{|y|\leq 2} \Omega_{\lambda,\mu}\left(\frac{y}{t}\right) dy \leq C_{n,\lambda,\mu} \omega_{\lambda,\mu}(t),$$

which can be proved similarly to (49).

When $j = 2$, we use

$$(57) \quad I_2 \leq \sum_{\ell=0}^{\lceil\sqrt{t}\rceil} \frac{1}{(2+\ell)^M} \int_{\frac{1+\ell}{t} < |\xi| \leq \frac{2+\ell}{t}} |\widehat{f}(\xi)|^2 d\xi.$$

Next, we apply estimate (36) on each of the latter integrals. We are already assuming that $t > 1$. Since $\omega_{\lambda,\mu}(t) \approx_{\lambda,\mu,J} 1$ on any compact subinterval J of $(0, \infty)$, we can in fact assume $t \geq 3$. Now we control the right-hand side of (57) with

$$\begin{aligned} (58) \quad C_{n,\lambda,\mu} \sum_{\ell=0}^{\lceil\sqrt{t}\rceil} \frac{1}{(2+\ell)^M} \omega_{\lambda,\mu}\left(\frac{2t}{3+2\ell}\right) \frac{1}{3+2\ell} \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)} \\ \leq C_{n,\lambda,\mu,M} \omega_{\lambda,\mu}(t) \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)}, \end{aligned}$$

provided $M > 2\lambda + 1$. This proves that (52) holds for $j = 2$.

If $j = 3$ then $(2 + \sqrt{t})/t < |\xi|$, which implies that

$$\frac{1}{(1 + |t\xi|)^M} \leq \frac{1}{(3 + \sqrt{t})^M}.$$

Then apply (36). Observe that, as long as $t > 1$, we have that the quantity \tilde{t} that now plays the role of t in (36) is bounded above and below by absolute constants, so $\omega_{\lambda,\mu}(\tilde{t}) \approx_{\lambda,\mu} 1$. In addition, for t in the same range, we have $\tilde{\varepsilon} \leq 1$ ($\tilde{\varepsilon}$ being the quantity that now plays the role of ε in (36)). These considerations imply that

$$\begin{aligned}
 (59) \quad I_3 &\leq \frac{1}{(3 + \sqrt{t})^M} \int_{\frac{2+\sqrt{t}}{t} < |\xi| \leq \frac{2+t}{t}} |\widehat{f}(\xi)|^2 d\xi \\
 &\leq C'_{n,\lambda,\mu} \frac{1}{(3 + \sqrt{t})^M} \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)} \\
 &\leq C''_{n,\lambda,\mu,M} \omega_{\lambda,\mu}(t) \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)},
 \end{aligned}$$

last inequality holding for a suitable constant $C''_{n,\lambda,\mu,M}$, provided that $M > 4\lambda + 2$.

It only remains to prove (52) with $j = 4$. We have

$$(60) \quad I_4 \leq \sum_{\ell=\lfloor t \rfloor + 1}^{\infty} \int_{\frac{1+\ell}{t} < |\xi| \leq \frac{2+\ell}{t}} |\widehat{f}(\xi)|^2 \frac{1}{(1 + |t\xi|)^M} d\xi.$$

Again, we apply (36) to the integral in the last term of (60), which is therefore controlled by

$$\begin{aligned}
 C_{n,\lambda,\mu} \sum_{\ell=\lfloor t \rfloor + 1}^{\infty} \frac{1}{(2 + \ell)^M} \frac{1}{\left(\frac{2t}{3+2\ell}\right)^{2\lambda+1}} \frac{1}{(3 + 2\ell)} \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)} \\
 \leq C_{n,\lambda,\mu,M} \frac{1}{t^{2\lambda+1}} \int_{\mathbb{R}^n} |f(x)|^2 \frac{dx}{w_{\lambda,\mu}(x)} \frac{1}{(1 + t)^{M-2\lambda-1}},
 \end{aligned}$$

which yields the desired inequality, provided that $M > 2\lambda + 1$. By choosing any $M > 4\lambda + 2$ (as required after (59)), we conclude the proof of (37) and of the claimed statement.

Proof of Corollary 1.2. The proof in [Carbery et al. 1988] can be used with $m_{\lambda,\gamma}$ instead of m_1^λ to account for the case where $\gamma \geq 0$ and $2 \leq p < p_\lambda$. When $p = p_\lambda$ and $\gamma > 1/p'_\lambda + 1/2$, values of μ satisfying $(2\lambda + 1)/n < \mu < 2\gamma - 2$ exist. For such μ , since $1 < 1 + 2\lambda < n$, we can use Theorem 1.1. Since (4) trivially holds for all $f \in \mathcal{S}$, the boundedness of $B_*^{\lambda,\gamma}$ implies that it also holds for every $f \in L^2(\mathbb{R}^n, dx)$ and every $f \in L^2(\mathbb{R}^n, w_{\lambda,\mu})$. But then it must hold for every $f \in L^2 + L^2(w_{\lambda,\mu})$. Since $(2\lambda + 1)/n < \mu$ we have $L^{p_\lambda} \subseteq L^2 + L^2(w_{\lambda,\mu})$, concluding the proof. \square

Acknowledgement

I would like to thank professor Loukas Grafakos for suggesting to work on this topic, for the conversations we have had on it, and for contributing to the economic support of this work.

References

[Annoni 2010] M. Annoni, *Almost everywhere convergence for modified Bochner Riesz means at the critical index for p greater than or equal to 2*, Ph.D. thesis, University of Missouri, 2010, available at <http://search.proquest.com/docview/9111786885>. MR

- [Ashurov 1983] R. R. Ashurov, “Summability almost everywhere of Fourier series from L_p in eigenfunctions”, *Mat. Zametki* **34**:6 (1983), 837–843. In Russian; translated in *Math. Notes Acad. Sci. USSR* **34**:6 (1983), 913–916. [MR](#) [Zbl](#)
- [Ashurov et al. 2010] R. Ashurov, A. Ahmedov, and A. Rodzi b. Mahmud, “The generalized localization for multiple Fourier integrals”, *J. Math. Anal. Appl.* **371**:2 (2010), 832–841. [MR](#) [Zbl](#)
- [Baernstein and Sawyer 1985] A. Baernstein, II and E. T. Sawyer, *Embedding and multiplier theorems for $H^p(\mathbb{R}^n)$* , Mem. Amer. Math. Soc. **318**, American Mathematical Society, Providence, RI, 1985. [MR](#) [Zbl](#)
- [Benedetto and Heinig 2003] J. J. Benedetto and H. P. Heinig, “Weighted Fourier inequalities: new proofs and generalizations”, *J. Fourier Anal. Appl.* **9**:1 (2003), 1–37. [MR](#) [Zbl](#)
- [Bochner 1936] S. Bochner, “Summation of multiple Fourier series by spherical means”, *Trans. Amer. Math. Soc.* **40**:2 (1936), 175–207. [MR](#) [Zbl](#)
- [Carbery 1983] A. Carbery, “The boundedness of the maximal Bochner–Riesz operator on $L^4(\mathbb{R}^2)$ ”, *Duke Math. J.* **50**:2 (1983), 409–416. [MR](#) [Zbl](#)
- [Carbery et al. 1988] A. Carbery, J. L. Rubio de Francia, and L. Vega, “Almost everywhere summability of Fourier integrals”, *J. London Math. Soc. (2)* **38**:3 (1988), 513–524. [MR](#) [Zbl](#)
- [Christ 1985] M. Christ, “On almost everywhere convergence of Bochner–Riesz means in higher dimensions”, *Proc. Amer. Math. Soc.* **95**:1 (1985), 16–20. [MR](#) [Zbl](#)
- [Gilbert 1972] J. E. Gilbert, “Interpolation between weighted L^p -spaces”, *Ark. Mat.* **10** (1972), 235–249. [MR](#) [Zbl](#)
- [Grafakos 2014] L. Grafakos, *Modern Fourier analysis*, 3rd ed., Graduate Texts in Mathematics **250**, Springer, 2014. [MR](#) [Zbl](#)
- [Lee and Seeger 2015] S. Lee and A. Seeger, “On radial Fourier multipliers and almost everywhere convergence”, *J. Lond. Math. Soc. (2)* **91**:1 (2015), 105–126. [MR](#) [Zbl](#)
- [Lee et al. 2014] S. Lee, K. M. Rogers, and A. Seeger, “Square functions and maximal operators associated with radial Fourier multipliers”, pp. 273–302 in *Advances in analysis: the legacy of Elias M. Stein*, Princeton Mathematical Series **50**, Princeton University Press, 2014. [MR](#) [Zbl](#)
- [Seeger 1987] A. Seeger, “Necessary conditions for quasiradial Fourier multipliers”, *Tohoku Math. J. (2)* **39**:2 (1987), 249–257. [MR](#) [Zbl](#)
- [Seeger 1996] A. Seeger, “Endpoint inequalities for Bochner–Riesz multipliers in the plane”, *Pacific J. Math.* **174**:2 (1996), 543–553. [MR](#) [Zbl](#)
- [Tao 1998] T. Tao, “The weak-type endpoint Bochner–Riesz conjecture and related topics”, *Indiana Univ. Math. J.* **47**:3 (1998), 1097–1124. [MR](#) [Zbl](#)
- [Tao 2002] T. Tao, “On the maximal Bochner–Riesz conjecture in the plane for $p < 2$ ”, *Trans. Amer. Math. Soc.* **354**:5 (2002), 1947–1959. [MR](#) [Zbl](#)
- [Watson 1944] G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, 1944. [MR](#) [Zbl](#)

Received November 4, 2014.

MARCO ANNONI
 1513 INDIANA ST.
 APT. A
 ST CHARLES, IL 60174
 UNITED STATES
marcoannoni.81@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jlhu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 286 No. 2 February 2017

Almost everywhere convergence for modified Bochner–Riesz means at the critical index for $p \geq 2$	257
MARCO ANNONI	
Uniqueness of conformal Ricci flow using energy methods	277
THOMAS BELL	
A functional calculus and restriction theorem on H-type groups	291
HEPING LIU and MANLI SONG	
Identities involving cyclic and symmetric sums of regularized multiple zeta values	307
TOMOYA MACHIDE	
Conformally Kähler Ricci solitons and base metrics for warped product Ricci solitons	361
GIDEON MASCHLER	
Calculating Greene’s function via root polytopes and subdivision algebras	385
KAROLA MÉSZÁROS	
Classifying resolving subcategories	401
WILLIAM SANDERS	
The symplectic plactic monoid, crystals, and MV cycles	439
JACINTA TORRES	
A note on torus actions and the Witten genus	499
MICHAEL WIEMELER	