

*Pacific  
Journal of  
Mathematics*

**CALCULATING GREENE'S FUNCTION  
VIA ROOT POLYTOPES AND SUBDIVISION ALGEBRAS**

KAROLA MÉSZÁROS

Volume 286    No. 2

February 2017



## CALCULATING GREENE’S FUNCTION VIA ROOT POLYTOPES AND SUBDIVISION ALGEBRAS

KAROLA MÉSZÁROS

Greene’s rational function  $\Psi_P(x)$  is a sum of certain rational functions in  $x = (x_1, \dots, x_n)$  over the linear extensions of the poset  $P$  (which has  $n$  elements), which he introduced in his study of the Murnaghan–Nakayama formula for the characters of the symmetric group. In recent work Boussicault, Féray, Lascoux and Reiner showed that  $\Psi_P(x)$  equals a valuation on a cone and calculated  $\Psi_P(x)$  for several posets this way. In this paper we give an expression for  $\Psi_P(x)$  for any poset  $P$ . We obtain such a formula using dissections of root polytopes. Moreover, we use the subdivision algebra of root polytopes to show that in certain instances  $\Psi_P(x)$  can be expressed as a product formula, thus giving a compact alternative proof of Greene’s original result and its generalizations.

### 1. Introduction

Given a poset  $P$  on the set  $[n] = \{1, \dots, n\}$ , Greene’s rational function is defined by

$$(1-1) \quad \Psi_P(x) = \sum_{w \in \mathcal{L}(P)} w \left( \frac{1}{(x_1 - x_2)(x_2 - x_3) \cdots (x_{n-1} - x_n)} \right),$$

where  $\mathcal{L}(P)$  denotes the set of linear extensions of  $P$  and for  $w \in \mathcal{L}(P)$  and a function  $f(x_1, \dots, x_n)$  we have that  $w(f(x_1, \dots, x_n)) = f(x_{w(1)}, \dots, x_{w(n)})$ . It was introduced by Greene [1992] in his work on the Murnaghan–Nakayama formula. Boussicault, Féray, Lascoux and Reiner [Boussicault et al. 2012] showed that

$$(1-2) \quad \Psi_P(x) = s(K_P^{\text{root}}; \mathbf{x}),$$

where

$$(1-3) \quad K_P^{\text{root}} = \mathbb{R}_+\{e_i - e_j \mid i <_P j\} = \mathbb{R}_+\{e_i - e_j \mid i \prec_P j\}$$

---

The author was partially supported by a National Science Foundation grant (DMS 1501059).

MSC2010: 05E10.

Keywords: Greene’s function, root polytope, subdivision algebra.

and

$$(1-4) \quad s(K; \mathbf{x}) := \int_K e^{-\text{span}_{\mathbb{R}_+}(\mathbf{x}, v)} dv,$$

for  $K$  a polyhedral cone in a Euclidean space  $V$  with inner product  $\text{span}_{\mathbb{R}_+}(\cdot, \cdot)$ .

Next we explain two important results about calculating  $\Psi_P(\mathbf{x})$ . Further work on  $\Psi_P(\mathbf{x})$  appeared in [Boussicault 2007; 2009; Boussicault and Féray 2009; Ilyuta 2009].

**Greene’s theorem.** Let  $P$  be a *strongly planar* poset, meaning that the Hasse diagram of  $P \sqcup \{\hat{0}, \hat{1}\}$  has a planar embedding with all edges directed upward in the plane. For a strongly planar poset  $P$  the edges of the Hasse diagram of  $P$  dissect the plane into bounded regions  $\rho$  such that the set of vertices of  $P$  in the boundary of  $\rho$  are two chains starting and ending at the same two elements,  $\min(\rho)$  and  $\max(\rho)$ , respectively. Denote by  $b(P)$  the set of bounded regions into which the Hasse diagram of  $P$  dissects the plane.

**Greene’s theorem** [Greene 1992]. *For any strongly planar poset  $P$ ,*

$$(1-5) \quad \Psi_P(\mathbf{x}) = \frac{\prod_{\rho \in b(P)} (x_{\min(\rho)} - x_{\max(\rho)})}{\prod_{i <_P j} (x_i - x_j)}.$$

**Boussicault’s, Féray’s, Lascoux’s and Reiner’s theorem.** A beautiful theorem appearing in [Boussicault et al. 2012] gives an expression for  $\Psi_P(\mathbf{x})$  for some posets  $P$  whose Hasse diagrams are bipartite graphs in terms of certain lattice paths. The setup is as follows. Let  $D$  be a skew Ferrers diagram in English notation, and let us label its rows from top to bottom by  $1, 2, \dots, r$  and its columns from right to left by  $1, 2, \dots, c$ . See the left of Figure 1. With this labeling the northeasternmost point of  $D$  is  $(1, 1)$  and the southwesternmost is  $(r, c)$ . The *bipartite poset*  $P_D$  is a poset on the set  $\{x_1, \dots, x_r, y_1, \dots, y_c\}$  with order relations  $x_i <_{P_D} y_j$  if and only if  $(i, j) \in D$ .

**BFLR theorem** [Boussicault et al. 2012]. *For any skew diagram  $D$ ,*

$$(1-6) \quad \Psi_{P_D}(\mathbf{x}) = \sum_{\pi} \frac{1}{\prod_{(i,j) \in \pi} (x_i - y_j)},$$

where the sum runs over all lattice paths  $\pi$  from  $(1, 1)$  to  $(r, c)$  inside  $D$  that take steps either one unit south or one unit west.

**Roadmap of the paper.** The objective of this paper is to (1) give a combinatorial expression of  $\Psi_P(\mathbf{x})$  for any poset  $P$ , (2) give an alternative proof of the BFLR theorem and (3) generalize Greene’s theorem. We accomplish (1) and (2) in Section 2, while we do (3) in Sections 3 and 4. In Sections 3 and 4 we also study the integer point transform of the root cone, which can be seen as a more refined

invariant of the cone than Greene's function. The integer point transform of the root cone and generalizations of Greene's theorem were also investigated in [Boussicault et al. 2012]. Our tools will be root polytopes and their subdivision algebras, the latter of which were introduced in [Mészáros 2011] and put to use in [Escobar and Mészáros 2015a; 2015b; Mészáros 2015a; 2015b; 2016a; 2016b; Mészáros and Morales 2015].

## 2. Greene's function for an arbitrary poset

The purpose of this section is twofold. First we show how to express  $\Psi_P(x)$  for any poset  $P$  in terms of  $\Psi_P(x)$  for posets  $P$  whose Hasse diagrams are alternating graphs. Then we give an expression for  $\Psi_P(x)$  for posets whose Hasse diagrams are alternating graphs, thereby also obtaining an expression for  $\Psi_P(x)$  for any poset  $P$ . Finally, we show that for certain posets  $P$  whose Hasse diagrams are bipartite graphs we can write  $\Psi_P(x)$  as a nice summation formula. The latter result originally appeared in the work of Boussicault, Féray, Lascoux and Reiner [Boussicault et al. 2012], who used triangulations of order polytopes in their proof. We phrase our proof in terms of root polytopes. The point of view of this paper is that (dissections of) root polytopes (and the root cone) are the unifying approach to the calculation of  $\Psi_P(x)$ .

A *root polytope* (of type  $A_{n-1}$ ) is the convex hull of the origin and some of the points  $e_i - e_j$  for  $1 \leq i < j \leq n$ . Given a graph  $G$  on the vertex set  $[n]$  we associate to it the root polytope

$$(2-1) \quad \tilde{Q}_G = \text{ConvHull}(0, e_i - e_j \mid (i, j) \in E(G), i < j).$$

It can be seen that  $\tilde{Q}_G$  is a simplex if and only if  $G$  is acyclic and to emphasize this we sometimes denote  $\tilde{Q}_G$  for acyclic graphs  $G$  by  $\tilde{\Delta}_G$ . In the proof of Lemma 4.2 we will also use the notation

$$(2-2) \quad \Delta_F = \text{ConvHull}(e_i - e_j \mid (i, j) \in E(F), i < j)$$

for a forest  $F$ .

The posets  $P$  we work with in this section are on the set  $[n]$  and they are labeled naturally; that is to say that if  $i <_P j$  then  $i < j$  in the order of natural numbers. Note that this does not pose a restriction on the results, it only makes them easier to state. Denote by  $\mathcal{H}(P)$  the graph of the Hasse diagram of  $P$ . The directed transitive closure of a graph  $H$  is denoted by  $\bar{H}$ , and it is the graph on vertex set  $V(G)$  with edges  $(i, j) \in \bar{H}$  if there is an increasing path from  $i$  to  $j$  in  $H$ .

**$\Psi_P(x)$  in terms of alternating posets.** This subsection explains how to reduce the computation of  $\Psi_P(x)$  to the computation of  $\Psi_P(x)$  for posets  $P$  whose Hasse diagram is an alternating graph. A graph  $G$  on the vertex set  $[n]$  is called *alternating*

if there are no edges  $(i, j)$  and  $(j, k)$  in it with  $i < j < k$ . We call a poset on  $[n]$  an *alternating poset* if its Hasse diagram is an alternating graph.

**Proposition 2.1.** *For any naturally labeled poset  $P$  on  $[n]$  we can write*

$$(2-3) \quad \Psi_P(\mathbf{x}) = \sum_{L,R} \Psi_{P_{L,R}}(\mathbf{x}),$$

where the summation runs over all  $L, R$  such that  $L \sqcup R = [n]$ , and

$$G_{L,R} = ([n], \{(i, j) \in E(G) \mid i \in L, j \in R, i < j\})$$

is a connected graph, where  $G = \overline{\mathcal{H}(P)}$ . Furthermore,  $\mathcal{H}(P_{L,R}) = G_{L,R}$  for a naturally labeled alternating poset  $P_{L,R}$ .

*Proof.* Recall that  $\Psi_P(\mathbf{x}) = s(K_P^{\text{root}}; \mathbf{x})$ . If  $K_P^{\text{root}} = \bigcup_{i=1}^l K_i$  for interior disjoint cones  $K_i$  with  $i \in [l]$  then  $s(K_P^{\text{root}}; \mathbf{x}) = \sum_{i=1}^l s(K_i; \mathbf{x})$ . If  $K_i = K_{P_i}^{\text{root}}$  for some posets  $P_i$  with  $i \in [l]$  then  $\Psi_P(\mathbf{x}) = \sum_{i=1}^l \Psi_{P_i}(\mathbf{x})$ . Therefore, to prove (2-3), it suffices to show that  $K_P^{\text{root}} = \bigcup_{L,R} K_{P_{L,R}}^{\text{root}}$ , where the union runs over all  $L, R$  such that  $L \sqcup R = [n]$ ,  $G_{L,R}$  is a connected graph ( $G = \overline{\mathcal{H}(P)}$ ) and  $\mathcal{H}(P_{L,R}) = G_{L,R}$  for a naturally labeled poset  $P_{L,R}$ .

Since  $K_P^{\text{root}} = \mathbb{R}_+\{e_i - e_j \mid i <_P j\}$ , if  $\tilde{Q}_G = \bigcup \tilde{Q}_{G_{L,R}}$  (the  $\tilde{Q}_{G_{L,R}}$  are interior disjoint), where the union runs over all  $L, R$  such that  $L \sqcup R = [n]$ , and  $G_{L,R}$  is a connected graph, then we also obtain that  $K_P^{\text{root}} = \bigcup_{L,R} K_{P_{L,R}}^{\text{root}}$  for interior disjoint cones  $K_{P_{L,R}}^{\text{root}}$ . The equation  $\tilde{Q}_G = \bigcup \tilde{Q}_{G_{L,R}}$  follows from [Postnikov 2009, Proposition 13.3] together with the observation that  $G = \overline{G}$  for our choice of  $G$ .  $\square$

We note that the cones  $K_{P_{L,R}}^{\text{root}}$  are generally not simplicial. One way to compute  $\Psi_{P_{L,R}}(\mathbf{x})$  would be to triangulate  $K_{P_{L,R}}^{\text{root}}$  into simplicial cones with rays of the form  $e_i - e_j$ , since for such a cone the following simple lemma gives the value of Greene’s function.

**Lemma 2.2** [Boussicault et al. 2012]. *The cone  $K_P^{\text{root}}$  is simplicial if and only if the Hasse diagram of  $P$  contains no cycles. In this case it is also unimodular and*

$$\Psi_P(\mathbf{x}) = \frac{1}{\prod_{i <_P j} (x_i - x_j)}.$$

We remark that a proof of Lemma 2.2 different from that given in [Boussicault et al. 2012] follows immediately using the subdivision algebra of root polytopes defined in [Mészáros 2011].

**Calculating  $\Psi_P(\mathbf{x})$  for an alternating poset  $P$ .** In light of Proposition 2.1, if we can calculate  $\Psi_P(\mathbf{x})$  for an alternating poset  $P$ , then we can in turn calculate  $\Psi_P(\mathbf{x})$  for any poset  $P$ . In this section we accomplish the former, building on the results of Li and Postnikov [2015]. The next paragraph follows the exposition of that paper.

Given an alternating graph  $G$  on the vertex set  $[n]$ , pick a linear order  $\mathcal{O}$  on the edges of  $G$ . Let  $T$  be a spanning tree of  $G$ , and let  $e$  be an edge that does not belong to  $T$ . Let  $C$  be the unique cycle contained in the graph  $([n], E(T) \cup \{e\})$ . Let  $e^*$  be the maximal edge in the cycle  $C$  in the linear ordering  $\mathcal{O}$  of the edges. We say that an edge  $e$  is *externally semiactive* if either  $e = e^*$  or there is an odd number of edges in  $C$  between  $e$  and  $e^*$ . (Since  $G$  is alternating, all cycles in  $G$  have an even length.) Let  $\text{ext}_{\mathcal{O}}^G(T)$  be the number of externally semiactive edges of  $G$  with respect to a spanning tree  $T$ .

**Theorem 2.3** [Li and Postnikov 2015]. *Given an alternating graph  $G$  and a linear ordering  $\mathcal{O}$  of its edges, let  $\mathcal{T}_{\mathcal{O}}^G$  be the set of spanning trees  $T$  with  $\text{ext}_{\mathcal{O}}^G(T) = 0$ . Then*

$$(2-4) \quad \tilde{Q}_G = \bigcup_{T \in \mathcal{T}_{\mathcal{O}}^G} \tilde{\Delta}_T,$$

where the simplices  $\tilde{\Delta}_T$  are interior disjoint.

**Corollary 2.4.** *For any naturally labeled poset  $P$  on  $[n]$  we can write*

$$(2-5) \quad \Psi_P(\mathbf{x}) = \sum_{L,R} \sum_{T \in \mathcal{T}_{G_{L,R}}^{\mathcal{O}_{L,R}}} \frac{1}{\prod_{(i,j) \in E(T), i < j} (x_i - x_j)},$$

where the summation runs over all  $L, R$  such that  $L \sqcup R = [n]$ , and

$$G_{L,R} = ([n], \{(i, j) \in E(G) \mid i \in L, j \in R, i < j\})$$

is a connected graph, where  $G = \overline{\mathcal{H}(P)}$ . Furthermore,  $\mathcal{O}_{L,R}$  is an arbitrary linear order of the edges of  $G_{L,R}$ .

*Proof.* The proof follows from Proposition 2.1, Lemma 2.2 and Theorem 2.3.  $\square$

We remark that we obtained Corollary 2.4 from a particular dissection of the root polytope  $\text{ConvHull}(0, e_i - e_j \mid e_i <_P e_j)$  into simplices. Such a dissection then induced a dissection of  $K_P^{\text{root}} = \mathbb{R}_+\{e_i - e_j \mid e_i <_P e_j\}$  into simplicial cones. Since we know that  $K_P^{\text{root}} = \mathbb{R}_+\{e_i - e_j \mid e_i \leq_P e_j\}$ , instead of  $\text{ConvHull}(0, e_i - e_j \mid e_i <_P e_j)$  one could also dissect  $\text{ConvHull}(0, e_i - e_j \mid e_i \leq_P e_j)$  into simplices and obtain an expression with fewer terms for  $\Psi_P(\mathbf{x})$ . However, since such a dissection also would not in general yield significantly fewer terms, we find the expression presented in Corollary 2.4 a fine representative of what a general formula for  $\Psi_P(\mathbf{x})$  for an arbitrary poset  $P$  can look like. We devote the next section to particularly nice formulas for  $\Psi_P(\mathbf{x})$  for special posets  $P$ , also demonstrating that in certain instances we can expect the formula presented in Corollary 2.4 to be far better than the formula given in (1-1), although this is not always the case.

**An alternative proof of the BFLR theorem.** Let  $P_D$  be the poset of a connected skew diagram  $D$  as in the BFLR theorem. Let  $G_D$  be the graph  $\mathcal{H}(P_D)$  drawn on a line with vertices from left to right,  $x_r, \dots, x_1, y_1, \dots, y_c$ , and with edges as arcs above this line. Note that the condition that  $G_D$  comes from  $P_D$  can be translated into the conditions that  $G_D$  is bipartite on parts  $\{x_1, \dots, x_r\}$  and  $\{y_1, \dots, y_c\}$  and for each  $i \in [r]$ ,  $x_i$  is connected to  $y_j$  for  $j \in [a_i, b_i]$ ,  $i \in [r]$ , where  $a_1 \leq \dots \leq a_r$  and  $b_1 \leq \dots \leq b_r$  and  $[1, c] = \bigcup_{i=1}^r [a_i, b_i]$ .

Given a drawing of a graph  $G$  such that its vertices  $v_1, \dots, v_n$  are arranged in this order on a horizontal line and its edges are drawn above this line, we say that  $G$  is *noncrossing* if it has no edges  $(v_i, v_k)$  and  $(v_j, v_l)$  with  $i < j < k < l$ . A vertex  $v_i$  of  $G$  is said to be *nonalternating* if it has both an incoming and an outgoing edge; it is called *alternating* otherwise. The graph  $G$  is alternating if all its vertices are alternating.


**Lemma 2.5.** *The root polytope  $\tilde{Q}_{G_D}$  decomposes into  $\tilde{Q}_{G_D} = \bigcup_T \tilde{\Delta}_T$ , where the union runs over all noncrossing alternating trees of  $G_D$  and the simplices  $\tilde{\Delta}_T$  are interior disjoint.*

Since noncrossing depends on the drawing of the graph it is essential that we remember that we drew  $G_D$  with vertices from left to right:  $x_r, \dots, x_1, y_1, \dots, y_c$ .

*Proof of Lemma 2.5.* Consider the following ordering  $\mathcal{O}$  on the edges of  $G_D$ . The edges incident to  $y_i$  precede the edges incident to  $y_j$  in the ordering  $\mathcal{O}$  if  $1 \leq i < j \leq c$ . Moreover, if edges  $(x_a, y_k)$  and  $(x_b, y_k)$  are incident to  $y_k$  for some  $k \in [c]$  with  $1 \leq a < b \leq r$ , then  $(x_a, y_k)$  precedes  $(x_b, y_k)$  in the ordering  $\mathcal{O}$ . We claim that then the spanning trees  $T$  of  $G_D$  with  $\text{ext}_{G_D}^{\mathcal{O}} = 0$  are exactly the noncrossing alternating trees of  $G_D$  and then the lemma follows from Theorem 2.3. Indeed, note that given any noncrossing alternating tree  $T$  of  $G_D$  and an edge  $e \in E(G_D) - E(T)$ , in the unique cycle  $C$  of the graph  $T$  with the edge  $e$  adjoined, the edge  $e$  is always 0 edges away from the largest edge of  $C$  in the ordering  $\mathcal{O}$ . Thus, for any noncrossing alternating tree  $T$  of  $G_D$  we have  $\text{ext}_{G_D}^{\mathcal{O}} = 0$ . On the other hand, given a crossing alternating spanning tree  $T'$  of  $G_D$  (note that all spanning trees of  $G_D$  are alternating) let the edges  $(x_i, y_j)$  and  $(x_k, y_l)$  cross with  $k > i$  and  $l < j$ . Since  $D$  is a connected skew diagram, both of the edges  $(x_k, y_j)$  or  $(x_i, y_l)$  are contained in  $G_D$ . Since  $T'$  is a spanning tree of  $G_D$ , it follows that exactly one of the edges from  $\{(x_k, y_j), (x_i, y_l)\}$  is in it. Adjoining the other edge as edge  $e$  we see that it is an externally semiactive edge for  $T$ , concluding the proof.  $\square$

**Lemma 2.6.** *The noncrossing alternating spanning trees of  $G_D$  are in bijection with the lattice paths  $\pi$  from  $(1, 1)$  to  $(r, c)$  inside  $D$  that take steps either one unit south or one unit west.*





**Figure 1.** The correspondence between noncrossing alternating spanning trees of  $G_D$  and lattice paths from  $(1, 1)$  to  $(r, c)$  inside  $D$  that take steps either one unit south or one unit west.

*Proof.* The bijection is given by the map that takes a noncrossing alternating spanning tree  $T = (\{x_r, \dots, x_1, y_1, \dots, y_c\}, \{(x_i, y_j) \mid (i, j) \in S(T)\})$  of  $G_D$  to the path  $\pi = S(T)$ . See Figure 1. □

Given a graph  $G$  on the vertex set  $[n]$  such that if  $(i, j) \in E(G)$  then the only increasing path from  $i$  to  $j$  in  $G$  is the edge  $(i, j)$  itself, we can define the naturally labeled poset  $P_G$  to be one on the set  $[n]$  with Hasse diagram given by (the edges of)  $G$ .

**Corollary 2.7** (BFLR theorem). *For any skew diagram  $D$ ,*

$$(2-6) \quad \Psi_{P_D}(\mathbf{x}) = \sum_{\pi} \frac{1}{\prod_{(i,j) \in \pi} (x_i - y_j)},$$

where the sum runs over all lattice paths  $\pi$  from  $(1, 1)$  to  $(r, c)$  inside  $D$  that take steps either one unit south or one unit west.

*Proof.* By Lemma 2.5 we have that the cone  $K_{P_D}^{\text{root}}$  is triangulated into simplicial cones  $K_{P_T}^{\text{root}}$ , where the  $T$ 's run over all noncrossing alternating spanning trees of  $G_D$ . By Lemma 2.6 the latter trees are in bijection with lattice paths  $\pi$  from  $(1, 1)$  to  $(r, c)$  inside  $D$  that take steps either one unit south or one unit west, and thus by Lemma 2.2 we obtain the corollary. □

Our proof for Corollary 2.7 is a special case of the proof of Corollary 2.4. We note that the formula for  $\Psi_{P_D}(\mathbf{x})$  given in Corollary 2.7 is substantially different from the expression given in (1-1). We can see this for example by looking at the number of terms that can appear in each. When  $D$  is a diagram in the shape of an  $r \times c$  rectangle, then in (1-1) we are summing over all linear extensions of the poset  $P_D$  yielding  $r!c!$  terms. In comparison, in Corollary 2.7 we have  $\binom{r+c-2}{r-1}$  terms corresponding to the lattice paths from  $(1, 1)$  to  $(r, c)$  inside  $D$ . The latter in general can be larger than the former. However, if instead we take  $D$  to be the skew shape  $D = (n, n - 1, \dots, 1) \setminus (n - 2, n - 3, \dots, 1)$ , then in Corollary 2.7 we have a single term and in (1-1) we are summing over all linear extensions of the

zigzag poset  $P_D$ . In this case the number of terms in (1-1) is larger than  $n!(n-1)!$ , which is many more than the one term in Corollary 2.7.

### 3. Lifting Greene's theorem to the subdivision algebra

The objective of this section is to generalize Greene's theorem to a relation in the subdivision algebra of root polytopes. Subdivision algebras of root polytopes were introduced and studied in [Mészáros 2011], where they were used for triangulating root polytopes. Subdivision algebras were also utilized for subword complexes and flow polytopes in [Escobar and Mészáros 2015a; Mészáros 2015a; 2015b; 2016a; 2016b; Mészáros and Morales 2015]. We will see in this section that both Greene's theorem and an analogous one for the integer point transform of the root cone are special cases of a relation in the subdivision algebra.

We begin by explaining how to use subdivision algebras to subdivide root cones  $K_P^{\text{root}}$ . Since Greene's function of a poset  $P$  is a valuation on a root cone  $K_P^{\text{root}}$  and we know its expression for unimodular root cones, if we triangulate  $K_P^{\text{root}}$  into unimodular root cones, then we obtain a way to calculate Greene's function of  $P$ .

**Root cones  $\mathcal{C}(G)$  and their subdivisions.** We establish a simpler notation for root cones here. For an arbitrary loopless graph  $G$ , define the *root cone*

$$(3-1) \quad \mathcal{C}(G) := \text{span}_{\mathbb{R}_+}(e_i - e_j \mid (i, j) \in E(G), i < j).$$

In order for  $\mathcal{C}(G)$  and  $\mathcal{C}(H)$  to be distinct for distinct graphs  $G$  and  $H$ , we will mostly consider *good graphs*  $G$ , which are loopless graphs such that if there is an increasing path from vertex  $i$  to vertex  $j$  in  $G$ , which is not the edge  $(i, j)$ , then the edge  $(i, j)$  is not present in  $G$ . (In particular,  $G$  contains no multiple edges.) Given a graph  $H$  let  $g(H)$  be the unique good graph on the vertex set  $V(H)$  such that  $\mathcal{C}(H) = \mathcal{C}(g(H))$ . The graph  $g(H)$  can be obtained from  $H$  by repeated removal of edges  $(i, j)$  for which there is an increasing path between  $i$  and  $j$  other than the edge  $(i, j)$ . In particular, all multiple edges are removed in order to obtain  $g(H)$ . An important property of root cones is given in the cone reduction lemma below, which can be expressed through reduction rules on graphs, as we now explain.

The *reduction rule for graphs*: given a graph  $G_0$  on the vertex set  $[n]$  and  $(i, j), (j, k) \in E(G_0)$  for some  $i < j < k$ , let  $G_1, G_2, G_3$  be graphs on the vertex set  $[n]$  with edge sets

$$(3-2) \quad \begin{aligned} E(G_1) &= E(G_0) \setminus \{(j, k)\} \cup \{(i, k)\}, \\ E(G_2) &= E(G_0) \setminus \{(i, j)\} \cup \{(i, k)\}, \\ E(G_3) &= E(G_0) \setminus \{(i, j), (j, k)\} \cup \{(i, k)\}. \end{aligned}$$

We say that  $G_0$  *reduces* to  $G_1, G_2$  and  $G_3$  under the reduction rules defined by equations (3-2).

For a good graph  $G$  we define two edges  $(i, j), (j, k) \in E(G), i < j < k$ , to be a *good pair of edges* of  $G$  if they belong to a common cycle in  $G$ , or if neither of them belongs to any cycle in  $G$ .

**Lemma 3.1** (cone reduction lemma; cf. [Mészáros 2011]). *Given a good graph  $G_0$  let  $(i, j), (j, k) \in E(G_0)$  be a good pair of edges of  $G_0$  for some  $i < j < k$  and  $G_1, G_2$  as described by equations (3-2). Then*

$$(3-3) \quad \mathcal{C}(G_0) = \mathcal{C}(G_1) \cup \mathcal{C}(G_2)$$

and

$$(3-4) \quad \mathcal{C}(G_3) = \mathcal{C}(G_1) \cap \mathcal{C}(G_2),$$

where the cones  $\mathcal{C}(G_0), \mathcal{C}(G_1), \mathcal{C}(G_2)$  are of the same dimension and  $\mathcal{C}(G_3)$  is a facet of both  $\mathcal{C}(G_1)$  and  $\mathcal{C}(G_2)$ .

For convenience we include a proof of Lemma 3.1 here. It is an adaptation of the proof from [Mészáros 2011], where it was written for acyclic graphs.

*Proof.* Let the edges of  $G_0$  be  $f_1 = (i, j), f_2 = (j, k), f_3, \dots, f_k$ . Let  $v(f_1), v(f_2), v(f_3), \dots, v(f_k)$  denote the vectors that the edges of  $G_0$  correspond to under the correspondence  $v : (i, j) \mapsto e_i - e_j$ , where  $i < j$ . By equations (3-2),

$$\begin{aligned} \mathcal{C}(G_0) &= \text{span}_{\mathbb{R}_+}(v(f_1), v(f_2), v(f_3), \dots, v(f_k)), \\ \mathcal{C}(G_1) &= \text{span}_{\mathbb{R}_+}(v(f_1), v(f_1) + v(f_2), v(f_3), \dots, v(f_k)), \\ \mathcal{C}(G_2) &= \text{span}_{\mathbb{R}_+}(v(f_1) + v(f_2), v(f_2), v(f_3), \dots, v(f_k)), \\ \mathcal{C}(G_3) &= \text{span}_{\mathbb{R}_+}(v(f_1) + v(f_2), v(f_3), \dots, v(f_k)). \end{aligned}$$

Thus, if  $\mathcal{C}(G_0)$  is  $d$ -dimensional, so are the cones  $\mathcal{C}(G_1)$  and  $\mathcal{C}(G_2)$ , while cone  $\mathcal{C}(G_3)$  is at least  $(d-1)$ -dimensional (and at most  $d$ -dimensional). We note that  $\dim(\mathcal{C}(G_3)) \neq d$  because  $G_0$  is a good graph and  $f_1$  and  $f_2$  are a good pair of edges.

Clearly,  $\mathcal{C}(G_1) \cup \mathcal{C}(G_2) \subset \mathcal{C}(G_0)$ . Given an expression of a vector  $v \in \mathcal{C}(G_0)$  as a nonnegative linear combination of the vectors  $v(f_1), v(f_2), v(f_3), \dots, v(f_k)$  it satisfies either that the coefficient of  $v(f_1)$  in such an expression is greater than or equal to the coefficient of  $v(f_2)$  in the expression, or it is not. In the former case we see that  $v \in \mathcal{C}(G_1)$  and in the latter case  $v \in \mathcal{C}(G_2)$ . Therefore,  $\mathcal{C}(G_0) = \mathcal{C}(G_1) \cup \mathcal{C}(G_2)$ .

Clearly,  $\mathcal{C}(G_3) \subset \mathcal{C}(G_1) \cap \mathcal{C}(G_2)$ . Given an expression of a vector  $v \in \mathcal{C}(G_1)$  as a nonnegative linear combination of the vectors  $v(f_1), v(f_2), v(f_3), \dots, v(f_k)$ , the coefficient of  $v(f_1)$  is greater than or equal to the coefficient of  $v(f_2)$ . Similarly, given an expression of a vector  $v \in \mathcal{C}(G_2)$  as a nonnegative linear combination of the vectors  $v(f_1), v(f_2), v(f_3), \dots, v(f_k)$ , the coefficient of  $v(f_1)$  is less than or equal to the coefficient of  $v(f_2)$ . Thus, there is an expression of  $v \in \mathcal{C}(G_1) \cap \mathcal{C}(G_2)$  as a nonnegative linear combination of the vectors  $v(f_1), v(f_2), v(f_3), \dots, v(f_k)$

such that the coefficient of  $v(f_1)$  is equal to the coefficient of  $v(f_2)$ . Therefore,  $\mathcal{C}(G_1) \cap \mathcal{C}(G_2) \subset \mathcal{C}(G_3)$ , leading to  $\mathcal{C}(G_1) \cap \mathcal{C}(G_2) = \mathcal{C}(G_3)$ .  $\square$

**The subdivision algebra, Greene’s theorem and the integer point transform of a root cone.** In this subsection we explain the subdivision algebra and show how it yields a slick proof for Greene’s theorem and its generalization.

A graph  $G$  can be encoded by the monomial  $m[G] = \prod_{(i,j) \in E(G), i < j} x_{ij}$  and the reduction rule going from  $G_0$  to  $G_1, G_2$  and  $G_3$  can be encoded by the equation  $x_{ij}x_{jk} = x_{ik}(x_{ij} + x_{jk} + \beta)$ . We define the *subdivision algebra*  $\mathcal{S}_n$  of root polytopes as the commutative algebra generated by the variables  $x_{ij}, 1 \leq i < j \leq n$ , subject to the relations  $x_{ij}x_{jk} = x_{ik}(x_{ij} + x_{jk} + \beta)$  for  $1 \leq i < j < k \leq n$ .

Let us explain the connection of the subdivision algebra to Greene’s function. If we set  $\beta = 0$ , then the relation  $x_{ij}x_{jk} = x_{ik}(x_{ij} + x_{jk})$  of  $\mathcal{S}_n$  is satisfied by  $x_{ij} := 1/(x_i - x_j)$ , which are the kind of terms appearing in Greene’s function. If instead, we set  $\beta = -1$ , then the relation  $x_{ij}x_{jk} = x_{ik}(x_{ij} + x_{jk} - 1)$  of  $\mathcal{S}_n$  is satisfied by  $x_{ij} := 1/(1 - x_i/x_j)$ . The latter will play a part in calculating the *integer point transform*  $\sigma_{K_P^{\text{root}}}(\mathbf{x})$  of the root cone  $K_P^{\text{root}} \subset \mathbb{Z}^d$  defined as

$$(3-5) \quad \sigma_{K_P^{\text{root}}}(\mathbf{x}) := \sum_{m \in K_P^{\text{root}} \cap \mathbb{Z}^d} \mathbf{x}^m.$$

The function  $\sigma_{K_P^{\text{root}}}(\mathbf{x})$  can be seen as a finer invariant of the cone than  $\Psi_P(\mathbf{x})$ , as explained in [Boussicault et al. 2012, Section 2.4]. We note that in that paper the integer point transform  $\sigma_{K_P^{\text{root}}}(\mathbf{x})$  is denoted as  $H(K_P^{\text{root}}; \mathbf{X})$  and is referred to as the Hilbert series of the affine semigroup ring of the root cone. We chose to follow the more geometric name and notation of [Beck and Robins 2007, Section 3.2].

We are now ready to prove the following generalization of Greene’s theorem via the subdivision algebra, which first appeared in [Boussicault et al. 2012]:

**Theorem 3.2** [Boussicault et al. 2012, Corollary 8.10]. *For any (connected) strongly planar poset  $P$  on  $[n]$  we have*


$$(3-6) \quad \sigma_{K_P^{\text{root}}}(\mathbf{x}) = \frac{\prod_{\rho \in b(P)} (1 - x_{\min(\rho)}/x_{\max(\rho)})}{\prod_{i <_P j} (1 - x_i/x_j)}$$

and

$$(3-7) \quad \Psi_P(\mathbf{x}) = \frac{\prod_{\rho \in b(P)} (x_{\min(\rho)} - x_{\max(\rho)})}{\prod_{i <_P j} (x_i - x_j)},$$

where  $\rho$  runs through all bounded regions of the Hasse diagram.

*Proof.* Since  $P$  is a connected strongly planar poset, it follows that its Hasse diagram is a good graph on the vertex set  $[n]$  such that for every cycle  $C$  of  $G$  the only alternating vertices of  $C$  (considered within  $C$ ), that is vertices that have




**Figure 2.** In reducing an increasing path we always pick the top-most leftmost edges in the path and its offsprings to do reductions on. For a graph  $G_0$  the arrow to the left points to  $G_1$ , the middle arrow to  $G_3$ , and the right arrow to  $G_2$ , as in equations (3-2).

only incoming or only outgoing edges, are its minimal and maximal vertices. Thus we have that  $K_p^{\text{root}} = \mathcal{C}(G)$  for a good graph  $G$ . Note that a root cone  $\mathcal{C}(H)$  is unimodular if and only if  $g(H)$  is acyclic. We will use the cone reduction lemma to write  $\mathcal{C}(G)$  as a union of unimodular cones. Note that the cone reduction lemma applies to good graphs, and thus if we want to repeatedly apply it to the outcome cones  $\mathcal{C}(G_i)$ ,  $i \in [3]$ , we need to apply it to  $g(G_i)$ ,  $i \in [3]$ .

We claim that we can apply the cone reduction lemma repeatedly in such a fashion that at the end we have trees  $T_1, \dots, T_k$  (with  $n - 1$  edges), and forests  $F_{n-i}^j$ ,  $2 \leq i \leq n - 1$ ,  $j \in I_{n-i}$  (for some index sets  $I_{n-i}$ ), with  $n - i$  edges, where  $\mathcal{C}(T_1), \dots, \mathcal{C}(T_k)$  are unimodular cones triangulating  $\mathcal{C}(G)$  and the  $\mathcal{C}(F_{n-i}^j)$  are their intersections.

We now prove the above claim. When  $G$  has no cycles, the claim is obvious. Suppose that  $G$  has  $m > 0$  linearly independent cycles. Fix a strongly planar drawing of  $P$ . In it there are  $m$  bounded regions, and the boundaries of these regions are  $m$  linearly independent cycles in  $G$ . Let  $C$  be one of these cycles, such that it bounds a region in the drawing of  $P$  which is adjacent to the infinite region. The cycle  $C$  consists of two increasing paths  $p$  and  $p'$  from  $i$  to  $j$  for some  $i < j$ . Let  $p = (i = i_0 \rightarrow i_1 \rightarrow i_2 \rightarrow \dots \rightarrow i_l = j)$  be the path bordering the infinite region in the drawing of  $P$ . We can perform consecutive reductions on the edges of the path  $p$  and its offsprings, ultimately obtaining all noncrossing alternating forests on the vertices  $\{i_0, i_1, \dots, i_l\}$  containing the edge  $(i_0, i_l)$ . We do this by picking the topmost leftmost edges that we can do a reduction on in  $p$  and its offsprings in the reduction process. See Figure 2 for an illustration. (A proof of the previous claim can be obtained by induction on the length of the path and is given in detail in [Mészáros 2011].) Until we arrive at the aforementioned noncrossing alternating forests on the vertices  $\{i_0, i_1, \dots, i_l\}$  containing the edge  $(i_0, i_l)$  all



**Figure 3.** Top left shows a strongly planar drawing of our poset, with the cycle  $C$  in bold. The path  $p$  is  $(1 \rightarrow 3 \rightarrow 4)$  and  $p'$  is  $(1 \rightarrow 2 \rightarrow 4)$ . Top right shows the graph  $G$ . Below are the graphs  $G_1, G_3, G_2$  obtained by applying the reduction on the topmost leftmost edges of  $p$ , which are  $(1, 3), (3, 4)$ . The last row shows  $g(G_1), g(G_3), g(G_2)$  (which are  $G_1, G_3, G_2$  with the edge  $(1, 4)$  removed since there is an increasing path  $1 \rightarrow 2 \rightarrow 4$ ), on which we can keep applying the cone reduction lemma as in the proof of Theorem 3.2.

graphs obtained in this fashion from  $G$  are good graphs. We can see that once we obtain the noncrossing alternating forests on the vertices  $\{i_0, i_1, \dots, i_l\}$  containing the edge  $(i_0, i_l)$  the offspring of  $G$  is not good anymore, as there is still  $p'$  in it, which is an increasing path between the vertices  $i_0$  and  $i_l$ . We need to now remove the edge  $(i_0, i_l) = (i, j)$  from all the aforementioned offsprings in order to obtain good graphs and be able to apply the cone reduction lemma further. However, once we remove the edge  $(i, j)$  from all these offsprings we will have good graphs with the number of bounded regions one less than it was for  $G$ . We can now repeat the same process we just described for each of these graphs and their offsprings until they are all acyclic. We demonstrate the basic idea of this argument in Figure 3.

If we inspect what edges we had to drop in the process to make sure we always apply the cone reduction lemma to good graphs and obtain the acyclic graphs described in the previous paragraph, we find the following relation in the subdivision

algebra:

$$(3-8) \quad m[G] = \prod_{\rho \in b(P)} x_{\min(\rho), \max(\rho)} \left( \sum_{T_i} m[T_i] + \sum_{F_{n-i}^j} \beta^{i-1} m[F_{n-i}^j] \right).$$

Note that

$$(3-9) \quad \sigma_{K_P^{\text{root}}}(\mathbf{x}) = \left( \sum_{T_i} m[T_i] + \sum_{F_{n-i}^j} (-1)^{i-1} m[F_{n-i}^j] \right) \Big|_{x_{ij}=1/(1-x_i x_j^{-1})}$$

and

$$(3-10) \quad \Psi_P(\mathbf{x}) = \sum_{T_i} m[T_i] \Big|_{x_{ij}=1/(x_i - x_j)}$$

Equations (3-8), (3-9) and (3-10) together with the observations that  $x_{ij} = 1/(1 - x_i x_j^{-1})$  satisfies  $x_{ij} x_{jk} = x_{ik}(x_{ij} + x_{jk} - 1)$  and that  $x_{ij} = 1/(x_i - x_j)$  satisfies  $x_{ij} x_{jk} = x_{ik}(x_{ij} + x_{jk})$  immediately yield equations (3-6) and (3-7).  $\square$

We can see (3-8) is the main theorem of this section, so we bestow it with that title:

**Theorem 3.3.** *Let  $G = \mathcal{H}(P)$  for a naturally labeled connected strongly planar poset  $P$ . Then, using the notation of the proof of Theorem 3.2, we have that*

$$m[G] = \prod_{\rho \in b(P)} x_{\min(\rho), \max(\rho)} \left( \sum_{T_i} m[T_i] + \sum_{F_{n-i}^j} \beta^{i-1} m[F_{n-i}^j] \right)$$

*holds in the subdivision algebra.*

Both statements of Theorem 3.2 are special cases of Theorem 3.3 as shown in the proof of Theorem 3.2.

#### 4. Generalizing Greene's theorem beyond strongly planar posets

In this section we will examine a special family of posets for which Greene's function factors linearly. These posets were first identified by Boussicault, Féray, Lascoux and Reiner [Boussicault et al. 2012], who proved the aforementioned result by studying the affine semigroup ring of the root cone. We will give a short alternative proof via root polytopes.

We give some definitions following the exposition of [Boussicault et al. 2012]. In a finite poset  $P$ , say that a triple of elements  $(a, b, c)$  forms a notch of  $\vee$  shape (dually, a notch of  $\wedge$  shape) if  $a <_P b, c$  (dually,  $b, c <_P a$ ), and in addition,  $b, c$  lie in different connected components of the poset  $P \setminus P_{\leq a}$  (dually,  $P \setminus P_{\geq a}$ ). When  $(a, b, c)$  forms a notch of either shape in a poset  $P$ , say that the quotient poset  $\bar{P} := P/\{b=c\}$ , having one fewer element and one fewer Hasse diagram edge, is obtained from  $P$  by closing the notch, and that  $P$  is obtained from  $\bar{P}$  by opening a notch.

**Theorem 4.1.** *Let  $P$  be a connected poset in which  $(a, b, c)$  forms a notch, and let  $\bar{P} := P/\{b = c\}$ . We assume without loss of generality that  $P$  and  $\bar{P}$  are naturally labeled. Then the root polytope  $\tilde{Q}_{\mathcal{H}(P)}$  has a triangulation with top-dimensional simplices  $\tilde{\Delta}_{T_1}, \dots, \tilde{\Delta}_{T_k}$ , and  $\tilde{Q}_{\mathcal{H}(\bar{P})}$  has a triangulation with top-dimensional simplices  $\tilde{\Delta}_{T'_1}, \dots, \tilde{\Delta}_{T'_k}$ , where  $(a, b) \in T'_i$ ,  $(a, b), (a, c) \in T_i$ ,  $i \in [k]$ , and moreover  $T_i|_{b=c} = T'_i$  (we ignore multiple edges).*

To prove Theorem 4.1 we use the following criterion.

**Lemma 4.2** (cf. [Postnikov 2009, Lemma 12.6]). *For two trees  $T$  and  $T'$  on the vertex set  $[n]$ , the intersection  $\tilde{\Delta}_T \cap \tilde{\Delta}_{T'}$  is a common face of the simplices  $\tilde{\Delta}_T$  and  $\tilde{\Delta}_{T'}$  if and only if the directed graph*

$$U(T, T') = ([n], \{(i, j) \mid (i, j) \in E(T), i < j\} \cup \{(j, i) \mid (i, j) \in E(T'), i < j\})$$

*has no directed cycles of length at least 3.*

The following proof of Lemma 4.2 is a straightforward adaptation of the proof of [Postnikov 2009, Lemma 12.6] to our more general setting. We include the proof here for convenience.

*Proof of Lemma 4.2.* Suppose that  $U(T, T')$  has a directed cycle  $C$  of length at least 3. Let  $E$  be the set of edges of  $T$  in  $C$  and  $E'$  be the set of edges of  $T'$  in  $C$ . Then  $\sum_{(i,j) \in E} (e_i - e_j) = \sum_{(i,j) \in E'} (e_i - e_j)$ . Let  $k = \max(|E|, |E'|)$ . Then

$$x := \frac{1}{k} \sum_{(i,j) \in E} (e_i - e_j) = \frac{1}{k} \sum_{(i,j) \in E'} (e_i - e_j) \in \tilde{\Delta}_T \cap \tilde{\Delta}_{T'}.$$

However, the minimal face of the simplex  $\tilde{\Delta}_T$  containing  $x$  is  $\Delta_{([n], E)}$  if  $k = |E|$  and  $\tilde{\Delta}_{([n], E)}$  if  $k > |E|$ . Similarly, the minimal face of the simplex  $\tilde{\Delta}_{T'}$  containing  $x$  is  $\Delta_{([n], E')}$  if  $k = |E'|$  and  $\tilde{\Delta}_{([n], E')}$  if  $k > |E'|$ . In any case, the minimal faces of  $\tilde{\Delta}_T$  and  $\tilde{\Delta}_{T'}$  containing  $x$  are not equal. Thus,  $\tilde{\Delta}_T \cap \tilde{\Delta}_{T'}$  is not their common face.

Next, assume that  $U(T, T')$  has no directed cycles of length at least 3. Let  $F = ([n], E(T) \cap E(T'))$ . Since  $U(T, T')$  has no directed cycles of length at least 3 we can find a function  $h : [n] \rightarrow \mathbb{R}$  such that (1)  $h$  is constant on connected components of  $F$ ; and (2) for any directed edge  $(a, b) \in U(T, T')$  that joins two different components of  $F$  we have  $h(a) < h(b)$ . Thus, if  $(a, b)$  is the edge  $(i < j)$  of  $T$  then  $h(i) < h(j)$ , and if  $(a, b)$  is the edge  $(i < j)$  of  $T'$  then  $h(i) > h(j)$ . The function  $h$  defines a linear form  $f_h$  on the space  $\mathbb{R}^n$  with the coordinates  $h(1), \dots, h(n)$  in the standard basis. The above conditions imply (1) for any vertex in the common face  $\tilde{\Delta}_F$  of  $\tilde{\Delta}_T$  and  $\tilde{\Delta}_{T'}$  we have  $f_h(x) = 0$ ; (2) for any vertex  $x \in \tilde{\Delta}_T \setminus \tilde{\Delta}_F$  we have  $f_h(x) < 0$ ; and (3) for any vertex  $x \in \tilde{\Delta}_{T'} \setminus \tilde{\Delta}_F$  we have  $f_h(x) > 0$ . Thus, the hyperplane  $f_h(x) = 0$  intersects  $\tilde{\Delta}_T$  and  $\tilde{\Delta}_{T'}$  at their common face  $\tilde{\Delta}_F$  as desired. □



*Proof of Theorem 4.1.* The criterion of Lemma 4.2 is sufficient to establish the above theorem, since we also have that  $\tilde{Q}_{\mathcal{H}(\bar{P})}$  has a triangulation with top-dimensional simplices  $\tilde{\Delta}_{T'_1}, \dots, \tilde{\Delta}_{T'_k}$ , where  $(a, b) \in T'_i$ , as  $e_a - e_b$  is a vertex of  $\tilde{Q}_{\mathcal{H}(\bar{P})}$ .  $\square$

When we calculate  $\sigma_{K_{\bar{P}\text{root}}}(\mathbf{x})$  and  $\Psi_{\bar{P}}(\mathbf{x})$  using triangulations of the root cones as implied by Theorem 4.1, we immediately get:

**Corollary 4.3** [Boussicault et al. 2012, Theorem 8.6]. *When  $\bar{P}$  is obtained from  $P$  by closing a  $\vee$ -shaped notch  $(a, b, c)$ , then*

$$\sigma_{K_{\bar{P}\text{root}}}(\mathbf{x}) = (1 - x_a x_b^{-1}) \sigma_{K_{P\text{root}}}(\mathbf{x})|_{x_b=x_c} \quad \text{and} \quad \Psi_{\bar{P}}(\mathbf{x}) = (x_a - x_b) \Psi_P(\mathbf{x})|_{x_b=x_c}.$$

A consequence of Theorem 4.1 is the following generalization of Greene's theorem pertaining to posets  $P$  to which we can repeatedly apply the opening notch operation and obtain a poset whose Hasse diagram has only cycles as biconnected components. Such posets  $P$  we call *admissible*. We now recall the definition of biconnected components following [Boussicault et al. 2012]. Given a graph  $G = (V, E)$  we say that two edges of it are cycle-equivalent if there is a cycle which contains both edges. Let  $E_i$  be the equivalence classes of this relation. Let  $V_i$  be the set of vertices which are the endpoint of at least one edge in  $E_i$ . Then the biconnected components of  $G$  are the graphs  $G_i = (V_i, E_i)$ .

**Theorem 4.4.** *Let  $P$  be an admissible planar poset. Then, we have*

$$\sigma_{K_P^{\text{root}}}(\mathbf{x}) = \frac{\prod_{\rho \in b(P)} (1 - \prod_{i \in \min(\rho)} x_i \prod_{j \in \max(\rho)} x_j^{-1})}{\prod_{i <_P j} (1 - x_i x_j^{-1})}$$

and

$$\Psi_P(\mathbf{x}) = \frac{\prod_{\rho \in b(P)} (\sum_{i \in \min(\rho)} x_{\min(i)} - \sum_{j \in \max(\rho)} x_j)}{\prod_{i <_P j} (x_i - x_j)},$$

where  $\rho$  runs through all bounded regions of the Hasse diagram of  $P$ .

*Proof.* This theorem can be deduced from Corollary 4.3 together with Corollaries 8.2 and 8.3 in [Boussicault et al. 2012]. We note that the latter corollaries also have simple proofs using the root polytope considerations of this paper, and we leave such alternative proofs as an exercise for the interested reader.  $\square$

### Acknowledgements

I am grateful to Vic Reiner for bringing Greene's function to my attention as well as for many informative and valuable exchanges about this work. I am also grateful to Alex Postnikov for sharing his knowledge generously. I thank the anonymous referee for many thoughtful suggestions which significantly improved the exposition.

## References

- [Beck and Robins 2007] M. Beck and S. Robins, *Computing the continuous discretely: integer-point enumeration in polyhedra*, Springer, New York, 2007. MR Zbl
- [Boussicault 2007] A. Boussicault, “Operations on posets and rational identities of type A”, 2007, available at <http://www.tinyurl.com/operations-on-posets>. Presented at *Formal Power Series and Algebraic Combinatorics* (Tianjin, China, 2007).
- [Boussicault 2009] A. Boussicault, *Action du groupe symétrique sur certaines fractions rationnelles suivi de Puissances paires du Vandermonde*, Ph.D. thesis, Université Paris-Est, 2009, available at <https://tel.archives-ouvertes.fr/tel-00502471>.
- [Boussicault and Féray 2009] A. Boussicault and V. Féray, “Application of graph combinatorics to rational identities of type A”, *Electron. J. Combin.* **16**:1 (2009), Research Paper 145, 39. MR
- [Boussicault et al. 2012] A. Boussicault, V. Féray, A. Lascoux, and V. Reiner, “Linear extension sums as valuations on cones”, *J. Algebraic Combin.* **35**:4 (2012), 573–610. MR Zbl
- [Escobar and Mészáros 2015a] L. Escobar and K. Mészáros, “Subword complexes via triangulations of root polytopes”, preprint, 2015. arXiv
- [Escobar and Mészáros 2015b] L. Escobar and K. Mészáros, “Toric matrix Schubert varieties and their polytopes”, 2015. To appear in *Proc. Amer. Math. Soc.* arXiv
- [Greene 1992] C. Greene, “A rational-function identity related to the Murnaghan–Nakayama formula for the characters of  $S_n$ ”, *J. Algebraic Combin.* **1**:3 (1992), 235–255. MR Zbl
- [Ilyuta 2009] G. Ilyuta, “Calculus of linear extensions and Newton interpolation”, preprint, 2009. arXiv
- [Li and Postnikov 2015] N. Li and A. Postnikov, “Slicing zonotopes”, preprint, 2015.
- [Mészáros 2011] K. Mészáros, “Root polytopes, triangulations, and the subdivision algebra, I”, *Trans. Amer. Math. Soc.* **363**:8 (2011), 4359–4382. MR Zbl
- [Mészáros 2015a] K. Mészáros, “ $h$ -polynomials via reduced forms”, *Electron. J. Combin.* **22**:4 (2015), Paper 4.18, 17. MR
- [Mészáros 2015b] K. Mészáros, “Product formulas for volumes of flow polytopes”, *Proc. Amer. Math. Soc.* **143**:3 (2015), 937–954. MR Zbl
- [Mészáros 2016a] K. Mészáros, “ $h$ -polynomials of reduction trees”, *SIAM J. Discrete Math.* **30**:2 (2016), 736–762. MR Zbl
- [Mészáros 2016b] K. Mészáros, “Pipe dream complexes and triangulations of root polytopes belong together”, *SIAM J. Discrete Math.* **30**:1 (2016), 100–111. MR Zbl
- [Mészáros and Morales 2015] K. Mészáros and A. H. Morales, “Flow polytopes of signed graphs and the Kostant partition function”, *Int. Math. Res. Not.* **2015**:3 (2015), 830–871. MR Zbl
- [Postnikov 2009] A. Postnikov, “Permutohedra, associahedra, and beyond”, *Int. Math. Res. Not.* **2009**:6 (2009), 1026–1106. MR Zbl

Received August 13, 2015. Revised March 9, 2016.

KAROLA MÉSZÁROS  
 DEPARTMENT OF MATHEMATICS  
 CORNELL UNIVERSITY  
 212 GARDEN AVE.  
 ITHACA, NY 14853  
 UNITED STATES  
 karola@math.cornell.edu

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Igor Pak  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pak.pjm@gmail.com](mailto:pak.pjm@gmail.com)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 286 No. 2 February 2017

---

Almost everywhere convergence for modified Bochner–Riesz means at the critical index for $p \geq 2$	257
MARCO ANNONI	
Uniqueness of conformal Ricci flow using energy methods	277
THOMAS BELL	
A functional calculus and restriction theorem on H-type groups	291
HEPING LIU and MANLI SONG	
Identities involving cyclic and symmetric sums of regularized multiple zeta values	307
TOMOYA MACHIDE	
Conformally Kähler Ricci solitons and base metrics for warped product Ricci solitons	361
GIDEON MASCHLER	
Calculating Greene’s function via root polytopes and subdivision algebras	385
KAROLA MÉSZÁROS	
Classifying resolving subcategories	401
WILLIAM SANDERS	
The symplectic plactic monoid, crystals, and MV cycles	439
JACINTA TORRES	
A note on torus actions and the Witten genus	499
MICHAEL WIEMELER	



0030-8730(201702)286:2;1-X