# Pacific Journal of Mathematics

# APPROXIMATION TO AN EXTREMAL NUMBER, ITS SQUARE AND ITS CUBE

JOHANNES SCHLEISCHITZ

Volume 287 No. 2

April 2017

## APPROXIMATION TO AN EXTREMAL NUMBER, ITS SQUARE AND ITS CUBE

JOHANNES SCHLEISCHITZ

We study rational approximation properties for successive powers of extremal numbers defined by Roy. For  $n \in \{1, 2\}$ , the classic approximation constants  $\lambda_n(\zeta)$ ,  $\hat{\lambda}_n(\zeta)$ ,  $w_n(\zeta)$ ,  $\hat{w}_n(\zeta)$  connected to an extremal number  $\zeta$ have been established and in fact much more is known. However, so far almost nothing had been known for  $n \ge 3$ . In this paper we determine all classic approximation constants as above for n = 3. Our methods will more generally provide detailed information on the combined graph defined by Schmidt and Summerer assigned to an extremal number, its square and its cube. We provide some results for n = 4 as well. In the course of the proofs of the main results we establish a very general connection between Khintchine's transference inequalities and uniform approximation.

#### 1. Approximation constants and extremal numbers

Let  $\zeta$  be a real transcendental number and  $n \ge 1$  be an integer. For  $1 \le j \le n+1$  we define the approximation constants  $\lambda_{n,j}(\zeta)$  as the supremum of  $\eta \in \mathbb{R}$  such that the system

(1) 
$$|x| \le X, \qquad \max_{1 \le i \le n} |\zeta^i x - y_i| \le X^{-\eta}$$

has (at least) *j* linearly independent solutions  $(x, y_1, y_2, ..., y_n) \in \mathbb{Z}^{n+1}$  for arbitrarily large values of *X*. Moreover, let  $\hat{\lambda}_{n,j}(\zeta)$  be the supremum of  $\eta$  such that (1) has (at least) *j* linearly independent solutions for all sufficiently large *X*. In the case of j = 1 we also only write  $\lambda_n(\zeta)$  and  $\hat{\lambda}_n(\zeta)$  respectively, which are just the classical approximation constants defined by Bugeaud and Laurent [2005]. By Dirichlet's theorem for all transcendental real  $\zeta$  and  $n \ge 1$  these exponents satisfy the estimate

(2) 
$$\lambda_n(\zeta) \ge \hat{\lambda}_n(\zeta) \ge \frac{1}{n}$$

Supported by the Austrian Science Fund FWF grant P24828.

MSC2010: primary 11J13; secondary 11H06.

Keywords: extremal numbers, Diophantine approximation constants, geometry of numbers, lattices.

Moreover from the definition we see that

$$\lambda_1(\zeta) \ge \lambda_2(\zeta) \ge \cdots$$
 and  $\hat{\lambda}_1(\zeta) \ge \hat{\lambda}_2(\zeta) \ge \cdots$ .

Similarly, let  $w_{n,j}(\zeta)$  and  $\widehat{w}_{n,j}(\zeta)$  be the supremum of  $\eta \in \mathbb{R}$  such that the system

(3) 
$$H(P) \le X, \qquad 0 < |P(\zeta)| \le X^{-\eta}$$

has (at least) *j* linearly independent polynomial solutions  $\sum_{i=0}^{n} a_i T^i$  of degree at most *n* with integers  $a_j$  for arbitrarily large *X* and all large *X* respectively, where  $H(P) = \max_{0 \le j \le n} |a_j|$ . Again for j = 1 we also write  $w_n(\zeta)$  and  $\widehat{w}_n(\zeta)$  which coincide with classical exponents. Again by Dirichlet's theorem we have

(4) 
$$w_n(\zeta) \ge \widehat{w}_n(\zeta) \ge n$$

Moreover it is obvious that

$$w_1(\zeta) \le w_2(\zeta) \le \cdots$$
 and  $\widehat{w}_1(\zeta) \le \widehat{w}_2(\zeta) \le \cdots$ 

The exponents defined above are connected via Khintchine's [1926] transference inequalities.

(5) 
$$\frac{w_n(\zeta)}{(n-1)w_n(\zeta)+n} \le \lambda_n(\zeta) \le \frac{w_n(\zeta)-n+1}{n}.$$

Similarly thanks to German [2012] we know that the uniform exponents are connected via

(6) 
$$\frac{\widehat{w}_n(\zeta) - 1}{(n-1)\widehat{w}_n(\zeta)} \le \widehat{\lambda}_n(\zeta) \le \frac{\widehat{w}_n(\zeta) - n + 1}{\widehat{w}_n(\zeta)}.$$

We point out that the estimates (5) and (6) hold more generally for the analogue exponents concerning vectors  $\underline{\zeta} \in \mathbb{R}^n$  whose coordinates are Q-linearly independent together with {1}; see for example [Schmidt and Summerer 2009]. This will be of some importance in Remark 3.2. Moreover in this case all estimates in (5) and (6) are known to be optimal.

It is known due to Davenport and Schmidt [1969] that  $\widehat{w}_2(\zeta) \leq \frac{3+\sqrt{5}}{2}$  for all real transcendental  $\zeta$ . Roy [2004a] proved that there exist countably many real transcendental numbers for which equality holds, and called such numbers extremal numbers. Their approximation properties have been intensely studied in dimensions  $n \in \{1, 2\}$ . We gather below some of the known facts which will be of importance for this paper. Throughout the paper let

$$\rho = 2 + \sqrt{5}, \qquad \tau = \frac{3 + \sqrt{5}}{2}, \qquad \nu = \frac{1 + \sqrt{5}}{2} \quad \text{and} \quad \gamma = \frac{\sqrt{5} - 1}{2}.$$

These values are linked via  $\tau = \nu^2$ ,  $\rho = \nu^3$  and  $\gamma = \nu^{-1}$ . Moreover  $\tau = \nu + 1$  and  $\nu^2 - \nu - 1 = 0$ . It is known that for  $\zeta$  an extremal number, the identities

(7) 
$$w_1(\zeta) = \lambda_1(\zeta) = \lambda_2(\zeta) = 1$$
,  $\hat{\lambda}_2(\zeta) = \gamma$ ,  $w_2(\zeta) = \rho$ ,  $\hat{w}_2(\zeta) = \tau$ 

hold. Concerning the higher successive minima functions it is immediate by Roy's results that any extremal number satisfies

- (8)  $w_{2,2}(\zeta) = \tau$ ,  $w_{2,3}(\zeta) = \nu$ ,  $\lambda_{2,2}(\zeta) = \gamma$ ,  $\lambda_{2,3}(\zeta) = \gamma^2$ ,
- (9)  $\widehat{w}_{2,2}(\zeta) = \nu, \qquad \widehat{w}_{2,3}(\zeta) = 1, \qquad \widehat{\lambda}_{2,2}(\zeta) = \gamma^2, \qquad \widehat{\lambda}_{2,3}(\zeta) = \gamma^3.$

In fact even more detailed approximation properties are known for n = 2. There is concise information on the integral approximation vectors inducing very good approximations in (1) such as for the polynomials inducing very good approximations in (3). We will concretely utilize the following consequence of Roy's results, which is part of the claim of [Roy 2004b, Theorem 7.2]. See also [Roy 2004a, Proposition 8.1, Theorem 8.2]. As usual  $a \simeq b$  means both  $a \ll b$  and  $b \ll a$  are satisfied everywhere it occurs in the sequel.

**Theorem 1.1** (Roy). For any extremal number  $\zeta$  there exists a sequence of irreducible polynomials  $(P_k)_{k\geq 1} \in \mathbb{Z}[T]$  of degree precisely two such that

 $H(P_{k+1}) \simeq H(P_k)^{\nu}$  and  $|P_k(\zeta)| \simeq H(P_k)^{-\rho}$ .

Moreover we have

(10)  $|P'_k(\zeta)| \asymp H(P_k).$ 

All the implied constants depend on  $\zeta$  only.

For the irreducibility and (10), see [Roy 2004a, Proposition 8.1, Theorem 8.2]. The other claims are part of the claims of [Roy 2004b, Theorem 7.2]. In fact the irreducibility is easily deduced from  $\lambda_1(\zeta) = 1$  in (7) and (42) below. Indeed these relations imply that  $P_k$  in the theorem cannot have a rational root at least for large k and are thus indeed irreducible. In context of (8), (9) we finally mention that for n = 2, extremal numbers induce the regular graph defined by Schmidt and Summere [2013b].

This paper aims to provide a better understanding of the classic approximation constants for extremal numbers in higher dimension n > 2. More generally we will provide a description of the behavior of the approximation functions  $L_j(q)$  and  $L_j^*(q)$  defined by Schmidt and Summerer [2009] in the course of their study of parametric geometry of numbers, for n = 3 and partially for n = 4. We recall basic facts on parametric geometry of numbers at the start of Section 3. Our results will arise as a combination of the known results on extremal numbers for  $n \in \{1, 2\}$  recalled above with estimates from parametric geometry of numbers. So far only a

few nontrivial quantitative results on classical approximation constants for extremal numbers in dimension n > 2 exist. The estimates

$$w_n(\zeta) \le \exp\{c(\zeta) \cdot (\log(3n))^2 (\log\log(3n))^2\}$$

for all  $n \ge 1$  and some constant  $c(\zeta) > 0$  are due to Adamczewski and Bugeaud [2010]. It was recently proved [Bugeaud and Schleischitz 2016] that  $\widehat{w}_3(\zeta) \le 4$  for extremal numbers  $\zeta$ , which improves the upper bound  $3 + \sqrt{2}$  valid for all transcendental real  $\zeta$  from the same paper (which in turn improved the bound 2n - 1 = 5 of Davenport and Schmidt [1969, Theorem 2b]). However, we will determine the precise value of  $\widehat{w}_3(\zeta)$  in Theorem 2.1. Besides approximation to extremal numbers by cubic algebraic integers has been investigated. Roy [2004a] showed that for extremal number  $\zeta$  and any algebraic integer  $\alpha$  of degree three we have

$$|\zeta - \alpha| \gg H(\alpha)^{-\tau - 1}.$$

Moreover in [Roy 2003, Theorem 1.1] he showed that for some extremal numbers the exponent  $-1 - \tau$  can be replaced by  $-\tau$ . The exponent  $-\tau$  is optimal since

$$|\zeta - \alpha| \ll H(\alpha)^{-\tau}$$

has solutions in algebraic integers  $\alpha$  of degree at most three and arbitrarily large height  $H(\alpha)$  for any given real number  $\zeta$ , as shown by Davenport and Schmidt [1969]. It follows that for any real  $\zeta$  there are monic polynomials of degree at most three and arbitrarily large height H(P) such that

$$|P(\zeta)| \ll H(P)^{-\nu}$$

It follows from [Roy 2004a] that the exponent  $\nu$  is optimal as well, since again the reverse inequality holds at least for some class of extremal numbers and arbitrarily large H(P).

#### 2. New results

*The case* n = 3. The first major result of the paper is the following.

**Theorem 2.1.** Let  $\zeta$  be an extremal number. Then we have

(11) 
$$w_3(\zeta) = w_2(\zeta) = \rho, \qquad \lambda_3(\zeta) = \frac{1}{\sqrt{5}},$$

and

(12) 
$$\widehat{w}_3(\zeta) = 3, \qquad \widehat{\lambda}_3(\zeta) = \frac{1}{3},$$

See the comments subsequent to Lemma 3.3 below for additional information on the dynamic behavior of the successive minima as parametric functions. This

dynamical point of view will also enable us to derive the following Theorem 2.2 from Theorem 2.1. As usual for an algebraic number  $\alpha$  we write  $H(\alpha) = H(P)$  where  $P \in \mathbb{Z}[T]$  is the irreducible minimal polynomial of  $\alpha$  over  $\mathbb{Z}[T]$  with coprime coefficients.

**Theorem 2.2.** Let  $\zeta$  be an extremal number and  $\epsilon > 0$ . Then the estimate

$$|Q(\zeta)| \le H(Q)^{-3-}$$

has only finitely many irreducible solutions  $Q \in \mathbb{Z}[T]$  of degree precisely three. In particular

 $\epsilon$ 

(14) 
$$|\zeta - \alpha| \le H(\alpha)^{-4-\epsilon}$$

has only finitely many algebraic solutions  $\alpha$  of degree precisely three. On the other hand the estimates

(15) 
$$|Q(\zeta)| \le H(Q)^{-3+\epsilon} \quad and \quad |\zeta - \alpha| \le H(\alpha)^{-4+\epsilon}$$

have solutions in irreducible polynomials Q of degree precisely three and algebraic  $\alpha$  of degree precisely three of arbitrarily large heights H(Q) and  $H(\alpha)$ . Moreover there are arbitrarily large X such that

(16) 
$$H(Q) \le X, \qquad |Q(\zeta)| \le X^{-\sqrt{5}-\epsilon}$$

has no irreducible solution  $Q \in \mathbb{Z}[T]$  of degree precisely three. In particular for arbitrarily large X the system

(17) 
$$H(\alpha) \le X, \qquad |\zeta - \alpha| \le H(\alpha)^{-1} X^{-\sqrt{5}-\epsilon}$$

has no algebraic solution  $\alpha$  of degree precisely three.

We strongly expect that the exponents in (16) and (17) are optimal as well. See the comments below the proof of Theorem 2.2 for a heuristic argument that supports this belief. Compare Theorem 2.2 with the estimates concerning approximation by algebraic integers  $\alpha$  at the end of Section 1.

*The case n* = 4. We want to establish a lower bound for the exponent  $\lambda_4(\zeta)$ . Our result, based on parametric geometry of numbers, is the following.

**Theorem 2.3.** Let  $\zeta$  be an extremal number. Then

(18) 
$$\lambda_4(\zeta) \ge \frac{\gamma}{2} = \frac{\sqrt{5-1}}{4}.$$

If  $w_4(\zeta) = w_2(\zeta) = \rho$ , then there is equality in (18) and moreover

(19) 
$$\widehat{w}_4(\zeta) = 4, \qquad \widehat{\lambda}_4(\zeta) = \frac{1}{4}.$$

Observe that  $\rho \approx 4.2361 > 4$ , so the assumption of the conditioned results are natural and thus we believe that there is actually equality in (18) and that (19) holds. Theorems 2.1 and 2.2 also support this belief. On the other hand (4) prohibits  $w_n(\zeta) = \rho$  for  $n \ge 5$ , which in general prohibits the methods of the paper from working for  $n \ge 5$ .

The constant in (18) is approximately  $\gamma/2 \approx 0.3090$ . Observe that this improves the lower bound derived from  $w_4(\zeta) \ge w_2(\zeta) = \rho$  in combination with Khintchine's transference inequalities (5), which turns out to be  $\frac{2+\sqrt{5}}{10+3\sqrt{5}} \approx 0.2535$ , only slightly larger than the trivial bound  $\frac{1}{4}$  from (2).

#### 3. Preparatory results

*Parametric geometry of numbers.* For the proofs of the new results we introduce some concepts of the parametric geometry of numbers following Schmidt and Summerer [2009, 2013a], where we develop the theory only as far as it is needed for our purposes and slightly deviate from their notation. In particular we restrict to the case of successive powers. Some more specific properties will be carried out in Section 4 for immediate application to preliminary results. Let  $\zeta \in \mathbb{R}$  be given and Q > 1 a parameter. For  $n \ge 1$  and  $1 \le j \le n+1$ , define  $\psi_{n,j}(Q)$  as the minimum of  $\eta \in \mathbb{R}$  such that

$$|x| \le Q^{1+\eta}, \quad \max_{1 \le j \le n} |\zeta^j x - y_j| \le Q^{-(1/n)+\eta}$$

has (at least) *j* linearly independent solutions  $(x, y_1, ..., y_n) \in \mathbb{Z}^{n+1}$ . The functions  $\psi_{n,j}(Q)$  can be equivalently defined via a lattice point problem, see [Schmidt and Summerer 2009]. They have the properties

$$-1 \le \psi_{n,j}(Q) \le \frac{1}{n}$$
 and  $Q > 1$ ,  $1 \le j \le n+1$ .

Let

$$\underline{\psi}_{n,j} = \liminf_{Q \to \infty} \psi_{n,j}(Q) \text{ and } \overline{\psi}_{n,j} = \limsup_{Q \to \infty} \psi_{n,j}(Q).$$

These values clearly all lie in the interval [-1, 1/n]. From Dirichlet's theorem it follows that  $\psi_{n,1}(Q) \leq 0$  for all Q > 1 and hence  $\overline{\psi}_{n,1} \leq 0$ . For our purposes, even more important will be the functions  $\psi_{n,j}^*(Q)$  from [Schmidt and Summerer 2009]. For  $1 \leq j \leq n+1$  and a parameter Q > 1, define the value  $\psi_{n,j}^*(Q)$  as the minimum of  $\eta \in \mathbb{R}$  such that

$$|H(P)| \le Q^{(1/n)+\eta}, \quad |P(\zeta)| \le Q^{-1+\eta}$$

has (at least) *j* linearly independent solutions in polynomials  $P \in \mathbb{Z}[T]$  of degree at most *n*. See the same work for the connection of the functions  $\psi_{n,j}^*$  to a related

lattice point problem, similarly as for simultaneous approximation. Again put

$$\underline{\psi}_{n,j}^* = \liminf_{Q \to \infty} \psi_{n,j}^*(Q) \quad \text{and} \quad \overline{\psi}_{n,j}^* = \limsup_{Q \to \infty} \psi_{n,j}^*(Q).$$

For transcendental  $\zeta$  Schmidt and Summerer [2013a, (1.11)] established the inequalities

$$j\underline{\psi}_{n,j} + (n+1-j)\overline{\psi}_{n,n+1} \ge 0$$
 and  $j\overline{\psi}_{n,j} + (n+1-j)\underline{\psi}_{n,n+1} \ge 0$ ,

for  $1 \le j \le n+1$ . The dual inequalities

(20) 
$$j\underline{\psi}_{n,j}^* + (n+1-j)\overline{\psi}_{n,n+1}^* \ge 0$$
 and  $j\overline{\psi}_{n,j}^* + (n+1-j)\underline{\psi}_{n,n+1}^* \ge 0$ ,

hold as well for the same reason. As pointed out in [Schmidt and Summerer 2009] Mahler's inequality implies

(21) 
$$|\psi_{n,j}(Q) + \psi^*_{n,n+2-j}(Q)| \ll \frac{1}{\log Q} \quad \text{for} \quad 1 \le j \le n+1.$$

In particular we have

(22) 
$$\underline{\psi}_{n,j} = -\overline{\psi}_{n,n+2-j}^*$$
 and  $\overline{\psi}_{n,j} = -\underline{\psi}_{n,n+2-j}^*$  for  $1 \le j \le n+1$ .

In particular all values  $\underline{\psi}_{n,j}^*$ ,  $\overline{\psi}_{n,j}^*$  lie in the interval [-1/n, 1], and  $\overline{\psi}_{n,1}^* \leq 0$  follows again from Dirichlet's theorem. The constants  $\underline{\psi}_{n,j}$ ,  $\overline{\psi}_{n,j}$ ,  $\underline{\psi}_{n,j}^*$ ,  $\overline{\psi}_{n,j}^*$  relate to the classical approximation constants  $\lambda_{n,j} = \lambda_{n,j}(\zeta)$ ,  $w_{n,j} = w_{n,j}(\zeta)$  assigned to real  $\zeta$  via

(23) 
$$(1 + \lambda_{n,j})(1 + \underline{\psi}_{n,j}) = (1 + \hat{\lambda}_{n,j})(1 + \overline{\psi}_{n,j}) = \frac{n+1}{n}$$
 for  $1 \le j \le n+1$ ,

and

(24) 
$$(1+w_{n,j})\left(\frac{1}{n}+\psi_{n,j}^*\right) = (1+\widehat{w}_{n,j})\left(\frac{1}{n}+\overline{\psi}_{n,j}^*\right) = \frac{n+1}{n} \quad \text{for} \quad 1 \le j \le n+1.$$

See [Schmidt and Summerer 2009, Theorem 1.4] for a proof of j = 1 which can be readily extended to the case of arbitrary  $1 \le j \le n + 1$  as noticed in [Schleischitz 2013]. From repeated application of (22), (23) and (24) one can deduce

(25) 
$$\lambda_{n,j}(\zeta) = \frac{1}{\widehat{w}_{n,n+2-j}(\zeta)} \quad \text{and} \quad \hat{\lambda}_{n,j}(\zeta) = \frac{1}{w_{n,n+2-j}(\zeta)},$$

for  $1 \le j \le n + 1$ , already noticed in [Schleischitz 2014]. For q > 0 we also define the functions

(26) 
$$L_{n,j}(q) = q\psi_{n,j}(Q)$$
 and  $L_{n,j}^*(q) = q\psi_{n,j}^*(Q)$ ,

where  $Q = e^q$ . They are piecewise linear with slopes among  $\{-1, 1/n\}$  and  $\{-1/n, 1\}$  respectively. More precisely locally any  $L_{n,j}$  coincides with some

(27) 
$$L_{\underline{x}}(q) = \max\left\{ \log |x| - q, \max_{1 \le j \le n} \log |\zeta^j x - y_j| + \frac{q}{n} \right\}$$

where  $\underline{x} = (x, y_1, ..., y_n) \in \mathbb{Z}^{n+1}$  for  $y_j$  the closest integer to  $\zeta^j x$ , see [Schmidt and Summerer 2009, page 75]. Similarly any  $L_{n,j}^*$  coincides locally with

(28) 
$$L_P^*(q) = \max\left\{\log H(P) - \frac{q}{n}, \log |P(\zeta)| + q\right\}$$

for some  $P \in \mathbb{Z}[T]$  of degree at most *n*. Observe that for fixed *P* the left expression in (28) decays with slope -1/n whereas the right expression rises with slope 1 in the parameter *q*. Consequently, at a local maximum of some  $L_{n,j}^*$ , the rising right expression of some  $L_P^*(q)$  meets the falling left expression of some  $L_Q^*(q)$  with H(Q) > H(P), and similarly for local maxima of  $L_{n,j}$ . On the other hand, at any local minimum *q* of some  $L_{n,j}^*$  there is either equality in the expressions in (28) for some *P*, or the rising phase of some  $L_P^*$  meets the falling phase of some  $L_Q^*$  for some *Q* with H(Q) > H(P). In the first case, which always applies for j = 1, the function  $L_{n,j}^*$  coincides with  $L_P^*$  in a neighborhood of *q*. The situation is again very similar for  $L_{n,j}$ . The identity (24) has a parametric version in the sense that for any  $(Q, \psi_{n,j}^*(Q))$  in the graph of some function  $\psi_{n,j}^*$  there exist *j* linearly independent polynomials  $P_1, \ldots, P_j \in \mathbb{Z}[T]$  of degree at most *n* such that

(29) 
$$(1+w_n^{(j)})\left(\frac{1}{n}+\psi_{n,j}^*(Q)\right)=\frac{n+1}{n}+o(1), \qquad Q\to\infty.$$

holds where

$$w_n^{(j)} := \frac{\min_{1 \le i \le j} (-\log |P_i(\zeta)|)}{\max_{1 \le i \le j} \log H(P_i)},$$

and vice versa. Very similarly a dual parametric version of (23) for the functions  $\psi_{n,j}(Q)$  can be obtained. Both versions are basically inherited from the proof of [Schmidt and Summerer 2009, Theorem 1.4]. A crucial observation for the parametric geometry of numbers developed in [Schmidt and Summerer 2009, 2013a] is that Minkowski's second lattice point theorem translates into

(30) 
$$\left|\sum_{j=1}^{n+1} L_{n,j}(q)\right| \ll 1 \text{ and } \left|\sum_{j=1}^{n+1} L_{n,j}^*(q)\right| \ll 1.$$

This implies that in any interval  $I = (q_1, q_2)$ , the sum of the differences  $L_{n,j}(q_2) - L_{n,j}(q_1)$  and  $L_{n,j}^*(q_2) - L_{n,j}^*(q_1)$  over  $1 \le j \le n+1$  are bounded in absolute value as well by a fixed constant independent of *I*. We will implicitly use this fact in the proof of Theorem 2.1. This argument is widely used in [Schmidt and Summerer 2013a].

*Two technical lemmas.* For the conditioned result (19) we need parts of Lemma 3.1 below, which is of some interest on its own. For its proof we will use that every local maximum of  $L_{n,1}$  is a local minimum of  $L_{n,2}$  (note: the analogue is in general false for  $L_{n,j}$ ,  $L_{n,j+1}$  when j > 1). This follows from the elementary fact that for any vector  $\underline{x} = (x, y_1, \dots, y_n) \in \mathbb{Z}^{n+1}$  clearly any integral multiple  $N\underline{x}$  cannot lead to a smaller value in (1). Hence if two functions  $L_{\underline{x}_1}$ ,  $L_{\underline{x}_2}$  as in (27) induce two (successive) falling slopes -1 of  $L_{n,1}$ , with some rising phase of  $L_{n,1}$  of slope 1/n in between, then the corresponding vectors  $\underline{x}_1$ ,  $\underline{x}_2$  are linearly independent, and the claim follows. Moreover we use  $L_{n,1}(q) < 0$  for all q > 0, which is equivalent to Dirichlet's theorem.

**Lemma 3.1.** Let  $n \ge 1$  be an integer and  $\zeta$  be a real transcendental number. Assume there is equality in either inequality of (5), that is, either

(31) 
$$n\lambda_n(\zeta) + n - 1 = w_n(\zeta)$$

or

(32) 
$$\lambda_n(\zeta) = \frac{w_n(\zeta)}{(n-1)w_n(\zeta) + n}$$

holds. Then  $\hat{\lambda}_n(\zeta) = 1/n$  and  $\widehat{w}_n(\zeta) = n$ .

*Proof.* Assume there is equality in the right inequality, that is  $n\lambda_n(\zeta) + n - 1 = w_n(\zeta)$ . In case of  $\lambda_n(\zeta) = \infty$  we have  $\hat{\lambda}_n(\zeta) = 1/n$  and  $\widehat{w}_n(\zeta) = n$  anyway by [Schleischitz 2016, Theorem 1.12 and Theorem 5.1]. Hence we can assume  $\lambda_n(\zeta) < \infty$ , which will simplify the estimates. It suffices to show  $\hat{\lambda}_n(\zeta) = 1/n$  since the two claims are well known to be equivalent, which follows for example from (6). It was shown by Schmidt and Summerer [2009, remark on page 80, after the proof of Theorem 1.4] that the right inequality in (5) is equivalent to  $\underline{\psi}_{n,1} + n\overline{\psi}_{n,n+1} \ge 0$ . It follows directly from their deduction of the mentioned remark that more generally the identity (31) implies that for any  $\varepsilon > 0$  there exist arbitrarily large parameters Q such that

$$|\psi_{n,1}(Q) + n\psi_{n,j}(Q)| < \varepsilon \quad \text{for} \quad 2 \le j \le n+1,$$

where Q can be chosen so that simultaneously  $\psi_{n,1}(Q)$  is arbitrarily close to  $\underline{\psi}_{n,1}$ and  $\psi_{n,j}(Q)$  is arbitrarily close to  $\overline{\psi}_{n,j}$  for  $2 \le j \le n+1$ . In particular, the identity (31) implies

(33) 
$$\underline{\psi}_{n,1} = -n\overline{\psi}_{n,2} = -n\overline{\psi}_{n,3} = \dots = -n\overline{\psi}_{n,n+1}$$

and that for any  $\epsilon > 0$  and the (arbitrarily large) parameters Q as above the estimate

(34) 
$$0 < \psi_{n,n+1}(Q) - \psi_{n,2}(Q) < \epsilon$$

is satisfied. Moreover, since  $\psi_{n,1}(Q)$  is close to  $\underline{\psi}_{n,1}$ , we may assume that at such Q the function  $\psi_{n,1}$  has a local minimum, or equivalently  $L_{n,1}$  has a local minimum

at log Q (otherwise we get a contradiction to the definition of  $\psi_1$  either for some  $\tilde{Q} < Q$  or some  $\tilde{Q} > Q$  dependent on whether  $\psi_{n,1}$  rises in some interval  $(Q - \delta, Q)$  or decays in some interval  $(Q, Q + \delta)$ ). Let  $\epsilon > 0$  and  $Q_1$  be any fixed large value as above that in particular satisfies (34). Further let  $q_1 = \log Q_1$ . The estimate (34) can be written in terms of the functions  $L_{n,.}$  as

(35) 
$$0 < L_{n,n+1}(q_1) - L_{n,2}(q_1) < \epsilon \cdot q_1.$$

From (30) we know that  $L_{n,1}(q_1)$  approximately equals  $-\sum_{j=2}^{n+1} L_{n,j}(q_1)$  up to addition of some constant, that is

$$\left|L_{n,1}(q_1) + \sum_{j=2}^{n+1} L_{n,j}(q_1)\right| \le C.$$

Since all  $L_{n,2}(q_1), \ldots, L_{n,n+1}(q_1)$  are roughly equal by (35), we further deduce

$$|L_{n,1}(q_1) + nL_{n,2}(q_1)| = \left| \left( L_{n,1}(q_1) + \sum_{j=2}^{n+1} L_{n,j}(q_1) \right) + \sum_{j=2}^{n+1} (L_{n,2}(q_1) - L_{n,j}(q_1)) \right|$$
  
$$\leq C + n\epsilon q_1,$$

and hence in particular

(36) 
$$L_{n,2}(q_1) \ge -\frac{L_{n,1}(q_1)}{n} - \epsilon q_1 - \tilde{C},$$

where  $\tilde{C} = C/n$  is another constant. Now let  $q_0$  be the largest value smaller than  $q_1$  at which the function  $L_{n,1}(q)$  has a local maximum. Then by the assumption that  $q_1$  is a local minimum of  $L_{n,1}$  justified above, the function  $L_{n,1}$  decays in the interval  $[q_0, q_1]$  with slope -1 so that

(37) 
$$L_{n,1}(q_1) - L_{n,1}(q_0) = q_0 - q_1.$$

On the other hand

(38) 
$$L_{n,2}(q_1) - L_{n,2}(q_0) \le \frac{q_1 - q_0}{n},$$

since the function  $L_{n,2}(q)$  has slope at most 1/n. Moreover, since any local maximum of  $L_{n,1}(q)$  is a local minimum of  $L_{n,2}(q)$ , we have

$$L_{n,1}(q_0) = L_{n,2}(q_0).$$

Combining this with (37) and (38) yields

$$L_{n,2}(q_1) - L_{n,1}(q_1) \le \left(1 + \frac{1}{n}\right)(q_1 - q_0).$$

Together with (36) we obtain

$$L_{n,1}(q_1) \ge L_{n,2}(q_1) - \left(1 + \frac{1}{n}\right)(q_1 - q_0) \ge -\frac{L_{n,1}(q_1)}{n} - \epsilon q_1 - \tilde{C} - \left(1 + \frac{1}{n}\right)(q_1 - q_0),$$

which yields

$$L_{n,1}(q_1) \ge -\frac{n\epsilon}{n+1}q_1 - \tilde{C} - (q_1 - q_0).$$

Together with (37) we infer

$$L_{n,1}(q_0) \ge -\frac{n\epsilon}{n+1}q_1 - \tilde{C}$$

Now the assumption  $\lambda_n(\zeta) < \infty$  implies with (23) that  $\psi_{n,1} > -1$  and from this it is not hard to see that  $q_1 \ll q_0$  for all  $q_0, q_1$  as above with a constant depending only on  $\lambda_n(\zeta)$  or equivalently  $\psi_{n,1}$ . Hence, for  $q_0 > 1$ , we have

$$0 > L_{n,1}(q_0) \gg -\epsilon q_0.$$

Since by the transcendence of  $\zeta$  the values  $q_0$  induced from  $q_1$  as above clearly tend to infinity as  $q_1$  does, we infer  $\overline{\psi}_{n,1} = 0$  as we may choose  $\epsilon$  arbitrarily small. By (23) this is again equivalent to  $\hat{\lambda}_n(\zeta) = 1/n$ . The proof in case of equality in the right inequality is finished.

We only sketch the deduction of the dual result. Assume the identity (32) holds. The dual characterization  $\psi_{n,1}^* + n\overline{\psi}_{n,n+1}^* \ge 0$  from [Schmidt and Summerer 2009] for the related left inequality in (5) yields the dual characterization for the equality (32) for the same reasons. Proceeding as above yields very similarly as above  $0 < \psi_{n,n+1}^*(Q) - \psi_{n,2}^*(Q) < \epsilon$  for large Q for which log Q are local minima of  $L_{n,1}^*$  and such that  $\psi_{n,1}^*(Q)$  is close to  $\psi_{n,1}^*$ , dual to (34). For such Q we now look at the smallest local maximum of  $L_{n,1}^*$  greater than log Q. Since all  $L_{n,j}^*$  have slope within  $\{-1/n, 1\}$ , the claim  $\widehat{w}_n(\zeta) = n$  follows very similarly incorporating that any local maximum of  $L_{n,1}^*$  is a local minimum of  $L_{n,2}^*$  again.

**Remark 3.2.** We point out that the proof of Lemma 3.1 does not require that the point lies on the Veronese curve defined as  $\{(t, t^2, ..., t^k) : t \in \mathbb{R}\}$ . The only point where we used the special form of successive powers was for  $\lambda_n(\zeta) = \infty$ , and in this case more concise estimates show the claim as well. Hence the claim extends naturally to the analogue exponents assigned to  $\zeta \in \mathbb{R}^k$  whose coordinates are linearly independent together with  $\{1\}$ .

It will be convenient to utilize the following Lemma 3.3 for the proof of Theorem 2.1. Roughly speaking, it shows that multiplication of a polynomial P with a polynomial Q for which  $|Q(\zeta)| \approx H(Q)^{-1}$  holds induces an increase of the corresponding function  $L_{3,.}^*$  by  $\frac{1}{3}$  in some interval. For fixed real  $\zeta$  we will say a polynomial  $P \in \mathbb{Z}[T]$  of degree at most 3 *induces a point*  $(q, L_P^*(q))$  *in the* 

3-dimensional Schmidt–Summerer diagram if  $(q, L_P^*(q))$  is the local minimum of  $L_P^*$  implicitly defined via H(P),  $P(\zeta)$  by

(39) 
$$L_P^*(q) = \log H(P) - \frac{q}{3} = \log |P(\zeta)| + q,$$

consistent with (28). Recall that any local minimum of some successive minimum function  $L_3^*$  is obtained as in (39) for some  $P \in \mathbb{Z}[T]$ .

**Lemma 3.3.** Let  $P, Q, R \in \mathbb{Z}[T]$  be of large heights and such that R = PQ and R has degree at most three. Assume P induces the point  $(q_1, L_P^*(q_1))$  and R induces the point  $(q_2, L_R^*(q_2))$  in the 3-dimensional Schmidt–Summerer diagram. Further assume

$$(40) \qquad \qquad |Q(\zeta)| = H(Q)^{-1+\delta}$$

for  $\delta$  of small absolute value, and that  $(\log H(Q))^{-1} = O(\delta)$ . Then

(41) 
$$\frac{L_R^*(q_2) - L_P^*(q_1)}{q_2 - q_1} = \frac{1}{3} + O(\delta).$$

*Proof.* From (39) we calculate

 $q_1 = \frac{3}{4} \cdot (\log H(P) - \log |P(\zeta)|)$  and  $L_P^*(q_1) = \frac{3}{4} \cdot \log H(P) + \frac{1}{4} \cdot \log |P(\zeta)|$ . Similarly, we infer

$$q_{2} = \frac{3}{4} \cdot (\log H(R) - \log |R(\zeta)|)$$
  
=  $\frac{3}{4} \cdot (\log H(P) + \log H(Q) + \Delta - (\log |P(\zeta)| + \log |Q(\zeta)|)),$ 

and

$$L_R^*(q_2) = \frac{3}{4} \cdot (\log H(P) + \log H(Q) + \Delta) + \frac{1}{4} \cdot (\log |P(\zeta)| + \log |Q(\zeta)|),$$

where  $\Delta$  is bounded by virtue of (42) below. Inserting yields

$$\frac{L_R^*(q_2) - L_P^*(q_1)}{q_2 - q_1} = \frac{\frac{3}{4}\log H(Q) + \frac{1}{4}\log |Q(\zeta)| + \frac{3}{4}\Delta}{\frac{3}{4}\log H(Q) - \frac{3}{4}\log |Q(\zeta)| + \frac{3}{4}\Delta},$$

and with the assumption (40) further

$$\frac{L_R^*(q_2) - L_P^*(q_1)}{q_2 - q_1} = \frac{\left(\frac{1}{2} + \frac{1}{4}\delta\right)\log H(Q) + \frac{3}{4}\Delta}{\left(\frac{3}{2} - \frac{3}{4}\delta\right)\log H(Q) + \frac{3}{4}\Delta}.$$

The claim follows by simply rearranging and assuming  $(\log H(Q))^{-1} = O(\delta)$ .  $\Box$ 

Conversely (41) implies that  $\log |Q(\zeta)| / \log H(Q) + 1$  is small by a very similar argument, but we will not use this. Again the proposition did not use the fact that we deal with successive powers of a number, and can be generalized to any dimension.

#### 4. Proofs of Theorems 2.1, 2.2 and 2.3

Apart from Theorem 1.1 and the concepts of the parametric geometry of numbers discussed on pages 490–492, we will use that for any polynomials  $Q_1$ ,  $Q_2$  with integral coefficients of degree bounded by n we have

(42) 
$$H(Q_1Q_2) \asymp_n H(Q_1)H(Q_2).$$

See [Wirsing 1961, Hilfssatz 3]. As in our applications the dimensions *n* are fixed we can assume absolute constants in (42). We will sometimes implicitly use the consequence that if  $Q = Q_1Q_2$  then  $|Q(\zeta)| \le H(Q)^{-z}$  implies that either  $|Q_1(\zeta)| \ll H(Q_1)^{-z}$  or  $|Q_2(\zeta)| \ll H(Q_2)^{-z}$  must be satisfied, which was essentially used by Wirsing [1961]. We start with the proof of Theorem 2.3 since it is the least technical one.

*Proof of Theorem 2.3.* We will prove that any extremal number  $\zeta$  satisfies

(43) 
$$w_{4,4}(\zeta) \ge \rho.$$

Assume we have already shown (43). Then the unconditional claim (18) follows from iterated use of results from parametric geometry of numbers. Indeed, from (43) applying (24) with n = j = 4 we first obtain

(44) 
$$\underline{\psi}_{4,4}^* \le \frac{2 - \sqrt{5}}{4(3 + \sqrt{5})}$$

In view of (22) and (20) applied with n = j = 4, we obtain

(45) 
$$\underline{\psi}_{4,1} = -\overline{\psi}_{4,5}^* \le 4 \cdot \underline{\psi}_{4,4}^* \le \frac{2-\sqrt{5}}{3+\sqrt{5}}.$$

Eventually computing the corresponding value of  $\lambda_4$  by applying (23) with n = 4, j = 1 leads precisely to the lower bound  $\gamma/2$  in the theorem.

We are left to prove (43). For this we use the characterization of the polynomials  $P_k \in \mathbb{Z}[T]$  of degree 2 for n = 2 from Theorem 1.1. Consider for fixed large k three successive polynomials  $P_{k-2}$ ,  $P_{k-1}$ ,  $P_k$ . Then we know from Theorem 1.1 that

(46) 
$$|P_j(\zeta)| \asymp H(P_j)^{-\rho}$$
 for  $j \in \{k-2, k-1, k\}.$ 

Applied with j = k it is obvious that the polynomials  $R_k(T) = TP_k(T)$  and  $S_k(T) = T^2P_k(T)$  have degrees 3 and 4, heights  $H(P_k) = H(R_k) = H(S_k)$ , and satisfy

$$|P_k(\zeta)| \asymp_{\zeta} |R_k(\zeta)| \asymp_{\zeta} |S_k(\zeta)| \asymp_{\zeta} H(P_k)^{-\rho}$$

as well. The polynomials  $P_k$ ,  $R_k$ ,  $S_k$  are obviously linearly independent and hence  $w_{4,3}(\zeta) \ge \rho$ . As the fourth polynomial  $T_k$  we take the product of  $P_{k-1}$  and  $P_{k-2}$ . First we show that  $\{P_k, R_k, S_k, T_k\}$  are linearly independent. Otherwise

 $T_k = P_{k-1}P_{k-2}$  would lie in the 3-dimensional space spanned by  $P_k$ ,  $R_k$ ,  $S_k$ , which by the special form of  $R_k$ ,  $S_k$  means  $T_k = P_k Z$  for some polynomial  $Z(T) \in \mathbb{Q}[T]$ of degree 2. However we know from Theorem 1.1 that the best approximating polynomials  $P_j$  are irreducible over  $\mathbb{Z}[T]$  for all large j. Hence by the unique factorization in  $\mathbb{Z}[T]$  the polynomial  $P_k$  must equal (up to sign) either  $P_{k-1}$  or  $P_{k-2}$ , which is clearly false, so we have a contradiction.

Moreover from (46) and the characterization in Theorem 1.1 it is known that  $H(P_{k-2})^{\nu^2} \simeq H(P_{k-1})^{\nu} \simeq H(P_k)$ . Since  $\nu^{-1} + \nu^{-2} = 1$  and  $H(T_k) \simeq H(P_{k-1})H(P_{k-2})$  by (42), we deduce  $H(T_k) \simeq H(P_k)$ . Together with property (46) for j = k - 1 and j = k - 2 we infer

$$|T_k(\zeta)| = |P_{k-1}(\zeta)P_{k-2}(\zeta)| \approx_{\zeta} H(P_{k-1})^{-\rho} H(P_{k-2})^{-\rho} \approx H(P_{k-1}P_{k-2})^{-\rho} \approx H(P_k)^{-\rho}$$

Summing up, we have found four linearly independent polynomials  $P_k$ ,  $R_k$ ,  $S_k$ ,  $T_k$  with the properties

$$H(P_k) \simeq H(R_k) \simeq H(S_k) \simeq H(T_k)$$

and

$$|P_k(\zeta)| \asymp_{\zeta} |R_k(\zeta)| \asymp_{\zeta} |S_k(\zeta)| \asymp_{\zeta} |T_k(\zeta)| \asymp_{\zeta} H(P_k)^{-\rho}$$

Since this holds for any large k we have established (43).

Finally we show the conditioned results. The equality  $\lambda_4(\zeta) = \gamma/2$  follows immediately from Khintchine's inequalities (5) since the upper bound for  $\lambda_4(\zeta)$ that arises from n = 4,  $w_4(\zeta) = \rho$ , coincides with the lower bound  $\gamma/2$  established above (the argument essentially used the characterization (33), (34) for equality (31) from [Schmidt and Summerer 2009] used in the proof of Lemma 3.1). Finally (19) follows from Lemma 3.1 since we have just shown that  $w_4(\zeta) = \rho$  implies the identity (31) for any extremal number  $\zeta$  and n = 4.

**Remark 4.1.** It was essentially shown in the proof of [Bugeaud 2010, Theorem 2] that the condition

(47) 
$$w_1(\zeta) = w_2(\zeta) = \dots = w_n(\zeta)$$

implies (31). If the hypothesis  $w_4(\zeta) = \rho$  of Theorem 2.3 holds then its assertion and (7) show that extremal numbers provide counterexamples for the reverse implication for n = 4. In this context note that if  $\lambda_n(\zeta) > 1$  the claims (47) and (31) are indeed equivalent by [Schleischitz 2016, Theorem 5.4]. Note also that from Lemma 3.1 and the above implication we could deduce that (47) implies  $\hat{\lambda}_n(\zeta) = 1/n$  and  $\hat{w}_n(\zeta) = n$ . However, the weaker condition  $w_1(\zeta) \ge n$  already implies  $\hat{\lambda}_n(\zeta) = 1/n$  and  $\hat{w}_n(\zeta) = n$  as established in [Schleischitz 2016, Theorem 5.1].

The proof of Theorem 2.3 in fact provides upper bounds for the frequency of good simultaneous rational approximations to  $(\zeta, \zeta^2, \zeta^3, \zeta^4)$ . More precisely the

proof shows that there exists a sequence  $(x_k)_{k\geq 1}$  of positive integers that satisfy

$$x_{k+1} \ll x_k^{\nu}$$
 and  $\max_{1 \le j \le 4} \|x_k \zeta^j\| \ll x_k^{-\gamma/2}$ .

In case of the conjectured equality in (18) we even have

(48) 
$$x_{k+1} \asymp x_k^{\nu} \quad \text{and} \quad \max_{1 \le j \le 4} \|x_k \zeta^j\| \asymp x_k^{-\gamma/2}.$$

Here as usual  $\|\cdot\|$  denotes the distance to the nearest integer. We briefly sketch how to deduce these facts from the proof above. The polynomials  $P_k$ ,  $R_k$ ,  $S_k$ ,  $T_k$ in the proof which induce the bound for the value  $\psi_{4,4}^*$  in (44) appear with frequency  $H(P_{k+1}) \simeq H(P_k)^{\nu}$  (and very similarly for  $R_k, S_k, T_k$ ). The last minimum  $\psi_{4,5}^{*}(Q)$  at the corresponding positions Q in the Schmidt–Summerer diagram is asymptotically bounded below as in (45) and the corresponding polynomials appear with the same logarithmic asymptotic height frequency  $\nu$ . We now flip the diagram along the horizontal axis according to (21) to obtain (roughly) the dual problem of simultaneous approximation. Thereby with simple geometric considerations involving (27) and reinterpreting to classical exponents  $\lambda_{4,..}$  we see that the first coordinates of best approximations related to the bound for  $\psi_{4,1}$  in (45) appear with frequency  $x_{k+1} \ll x_k^{\nu}$  as well (with a technical proof it possible to show that a single  $x_k$  cannot induce the good approximations for two consecutive values of Q obtained this way). In case of equality in (18) the functions  $\psi_{4,1}(Q)$  must have a local minimum at such places Q and (48) follows. It is tempting to further conjecture that for the corresponding approximation vectors  $(x_k, y_{k,1}, \ldots, y_{k,4})_{k\geq 1}$ , where  $x_k$ is as in (48) and  $y_{k,j}$  is the closest integer to  $\zeta^j x_k$ , similar general recursive patterns as for n = 2 noticed in [Roy 2004a] exist. However, we do not further investigate this topic here.

We turn to the case n = 3. For a real number  $\zeta$  we define the sequence of 1-dimensional *best approximation polynomials*  $(E_l)_{l\geq 1}$  attached to  $\zeta$ . They are given by linear polynomials  $E_l(T) = a_l T + b_l$  with  $a_l, b_l \in \mathbb{Z}$  defined by  $E_1(T) = T - \lfloor \zeta \rfloor$  and  $E_{l+1}$  is recursively defined via  $E_l$  as the linear polynomial of least height for which  $0 < |E_{l+1}(\zeta)| < |E_l(\zeta)|$ . These polynomials obviously satisfy  $H(E_1) < H(E_2) < \cdots$  and

$$E_l(\zeta) = \min\{|Q(\zeta)| : Q \in \mathbb{Z}[T], \quad \deg(Q) = 1, \quad 1 \le H(Q) \le H(E_l)\}.$$

It follows from the theory of continued fractions that the rational numbers  $b_l/a_l$  are precisely the convergents to  $\zeta$ . Moreover by Dirichlet's theorem the best approximating polynomials satisfy

(49) 
$$|E_l(\zeta)| \ll_{\zeta} H(E_l)^{-1} \text{ for } l \ge 1.$$

Furthermore it is well known and follows from elementary results on the theory of continued fractions that  $|E_l(\zeta)| \simeq_{\zeta} H(E_{l+1})^{-1}$  for all irrational  $\zeta$ , which readily implies

(50) 
$$1 \leq \liminf_{l \to \infty} \frac{\log H(E_{l+1})}{\log H(E_l)} \leq \limsup_{l \to \infty} \frac{\log H(E_{l+1})}{\log H(E_l)} = \lambda_1(\zeta).$$

In view of the rather technical proof of (11), for the convenience of the reader we give a brief outline of some facts we will show in the course of the proof. We will establish a rather precise description of the functions  $L_{3,1}^*(q), \ldots, L_{3,4}^*(q)$  on  $q \in (0, \infty)$  induced by an extremal number, its square and its cube. Denote by |I|the length of an interval *I*. We will show there exists a partition of the positive real numbers in successive intervals  $I_1, J_1, I_2, J_2, \ldots$  with the following properties:

- $\lim_{k\to\infty}|I_k|/|J_k|=1.$
- $\lim_{k \to \infty} |I_{k+1}| / |I_k| = \lim_{k \to \infty} |J_{k+1}| / |J_k| = \nu.$
- At the beginning of every *I<sub>k</sub>* all *L*<sup>\*</sup><sub>3,i</sub>(*q*) are all small (more precisely *o*(*q*) as *q* → ∞) by absolute value. Then in *I<sub>k</sub>* the functions *L*<sup>\*</sup><sub>3,1</sub>(*q*), *L*<sup>\*</sup><sub>3,2</sub>(*q*) basically decay with slope -1/3, whereas *L*<sup>\*</sup><sub>3,3</sub>(*q*), *L*<sup>\*</sup><sub>3,4</sub>(*q*) basically rise with slope <sup>1</sup>/<sub>3</sub> in any not too short subinterval of *I<sub>k</sub>* (clearly not in too short intervals, since the *L*<sup>\*</sup><sub>3</sub>, have slope within {-<sup>1</sup>/<sub>3</sub>, 1}).
- At the end of  $I_k$  and beginning of  $J_k$  the opposite behavior appears; that is,  $L_{3,1}^*(q), L_{3,2}^*(q)$  basically rise with slope  $\frac{1}{3}$  on any not too short subinterval of  $J_k$ , whereas  $L_{3,3}^*(q), L_{3,4}^*(q)$  basically decay with slope  $-\frac{1}{3}$  until the functions  $L_{3,1}^*, \ldots, L_{3,4}^*$  asymptotically meet again at the end of  $J_k$  which is the beginning of  $I_{k+1}$ .
- The functions  $|L_{3,1}^*(q) L_{3,2}^*(q)|$  such as  $|L_{3,3}^*(q) L_{3,4}^*(q)|$  are bounded uniformly in *q*.

All above is basically true for the simultaneous approximation functions  $L_{3,j}(q)$  as well by (22). Observe that by the last point above in particular

(51)	$w_{3,1}(\zeta) = w_{3,2}(\zeta),$	$w_{3,3}(\zeta) = w_{3,4}(\zeta),$	$\widehat{w}_{3,1}(\zeta) = \widehat{w}_{3,2}(\zeta),$	$\widehat{w}_{3,3}(\zeta) = \widehat{w}_{3,4}(\zeta),$
(52)	$\lambda_{3,1}(\zeta) = \lambda_{3,2}(\zeta),$	$\lambda_{3,3}(\zeta) = \lambda_{3,4}(\zeta),$	$\hat{\lambda}_{3,1}(\zeta) = \hat{\lambda}_{3,2}(\zeta),$	$\hat{\lambda}_{3,3}(\zeta) = \hat{\lambda}_{3,4}(\zeta),$

which extends the claim of Theorem 2.1. See also Remark 4.2 below. We point out that roughly speaking the decay phases of  $L_{3,..}^*$  are induced by the polynomials  $P_k$  from Theorem 1.1. The rising phases are induced by products  $P_k E_l$  for fixed  $P_k$  and suitable successive best approximating polynomials  $E_l$  defined above, which indeed lead to asymptotic increase by  $\frac{1}{3}$  as stated in the description above, basically in view of Lemma 3.3.

*Proof of Theorem 2.1.* First we prove (12). We show that

(53) 
$$w_{3,4}(\zeta) \ge 3.$$

Provided this is true it follows immediately that  $w_{4,3}(\zeta) = \hat{w}_3(\zeta) = 3$ , since  $w_{3,4}(\zeta) = \hat{\lambda}_3(\zeta)^{-1} \leq 3$  by (25) and (2). This argument in fact utilizes parametric geometry of numbers. Actually it is well known and follows, for example, from (6) that both claims in (12) are equivalent.

For (12) it remains to be shown that (53) holds. Let *k* be fixed large and consider the polynomials  $P_k$ ,  $P_{k+1}$ , ... from Theorem 1.1, and let  $R_j(T) = TP_j(T)$  for  $j \ge k$ . Further let  $X = H(P_{k+1})$ . Then obviously  $P_{k+1}(T)$  and  $R_{k+1}(T) = TP_{k+1}$  satisfy

(54) 
$$H(P_{k+1}) = H(R_{k+1}) = X$$
 and  $|P_{k+1}(\zeta)| \asymp_{\zeta} |R_{k+1}(\zeta)| \asymp_{\zeta} X^{-\rho} < X^{-3}$ .

Let  $\epsilon > 0$ . We shall construct polynomial multiples

(55) 
$$Q_{k,1} = R_{k,1} \cdot P_k$$
 and  $Q_{k,2} = R_{k,2} \cdot P_k$ 

of  $P_k$  with  $R_{k,i} \in \mathbb{Z}[T]$  polynomials of degree one such that  $\{R_{k,1}, R_{k,2}\}$  and hence also  $\{Q_{k,1}, Q_{k,2}\}$  are linearly independent and satisfy

(56) 
$$H(Q_{k,i}) \ll X$$
 and  $|Q_{k,i}(\zeta)| \ll X^{-3+\epsilon}$  for  $i \in \{1, 2\}$ 

One readily verifies that  $\{Q_{k,1}, Q_{k,2}\}$  span the same space as  $\{P_k, TP_k\}$  regardless of which linear polynomials  $R_{k,i}$  we choose. Observe that the space spanned by  $\{P_{k+1}, R_{k+1}, Q_{k,1}, Q_{k,2}\}$  consequently has dimension 4. Indeed, otherwise the polynomial identity  $P_k(T)Y_1(T) = P_{k+1}(T)Y_2(T)$  would have linear integer polynomial solutions  $Y_1, Y_2$ , which is a contradiction since  $P_k, P_{k+1}$  have degree two and are irreducible and not proportional and  $\mathbb{Z}[T]$  has unique factorization. Hence from (54) and (56) indeed the claim (53) follows by considering  $\{P_{k+1}, R_{k+1}, Q_{k,1}, Q_{k,2}\}$  as  $\epsilon$  can be chosen arbitrarily small. To finally prove (56), for the given  $X = H(P_{k+1})$ we let  $R_{k,1} = E_l$  and  $R_{k,2} = E_{l+1}$  be two successive best approximating polynomials in dimension n = 1 as introduced before the proof with l chosen largest possible such that still  $H(R_{k,i})H(P_k) \leq X$  for  $i \in \{1, 2\}$ . It follows from (42) and (55) that

(57) 
$$H(Q_{k,i}) \ll X \text{ for } i \in \{1, 2\}$$

On the other hand, since extremal numbers satisfy  $\lambda_1(\zeta) = 1$  as mentioned in (7), by (50) the sequence  $(E_l(T))_{l\geq 1}$  of best approximating polynomials in dimension 1 satisfies

(58) 
$$\lim_{l \to \infty} \frac{\log H(E_{l+1})}{\log H(E_l)} = 1 \quad \text{and} \quad \lim_{l \to \infty} -\frac{\log |E_l(\zeta)|}{H(E_l)} = 1.$$

Since  $R_{k,1} = E_l$ ,  $R_{k,2} = E_{l+1}$  and by our maximal choice of *l*, it is easy to see that

$$H(Q_{k,i}) \ge X^{1-\epsilon} \quad \text{for} \quad i \in \{1, 2\}.$$

It further follows from (42) and the fact that  $H(P_{k+1}) \simeq H(P_k)^{\nu}$ , or equivalently  $H(P_k) \simeq H(P_{k+1})^{\gamma}$  in view of Theorem 1.1, that we have

$$H(R_{k,i}) \gg H(Q_{k,i})H(P_k)^{-1} \gg X^{1-\epsilon}H(P_k)^{-1} \gg X^{1-\gamma-\epsilon}$$
 for  $i \in \{1, 2\}$ .

Together with (49) this leads to

$$R_{k,i}(\zeta) \ll_{\zeta} X^{-1+\gamma+\epsilon}$$
 for  $i \in \{1, 2\}$ .

Hence

$$|Q_{k,i}(\zeta)| = |P_k(\zeta)| \cdot |R_{k,i}(\zeta)| \ll_{\zeta} X^{-\rho\gamma} \cdot X^{-1+\gamma+\epsilon} = X^{-3+\epsilon} \quad \text{for} \quad i \in \{1, 2\},$$

where we used  $\rho\gamma + 1 - \gamma = 3$ , which can be readily checked. Thus recalling (57) we have proved (56) and hence together with (54) finally (12).

Now we prove the more technical identities (11). In the proof of (12) above we have shown that for any large k, with  $X = H(P_{k+1})$  we have four linearly independent polynomials  $\{T_1, \ldots, T_4\} = \{P_{k+1}, R_{k+1}, Q_{k,1}, Q_{k,2}\}$  with  $H(T_i) \ll X$ and  $|T_i(\zeta)| \le X^{-3+\epsilon}$ . Following the proof of (24), this means that for arbitrarily small  $\varepsilon > 0$ , any large k induces  $q_k > 0$  such that all

(59) 
$$|L_{3,i}^*(q_k)| \le \varepsilon q_k \quad \text{for} \quad 1 \le i \le 4,$$

where  $\lim_{k\to\infty} q_k / \log H(P_{k+1}) = 3$  in view of (28). Since by Theorem 1.1 any polynomial  $P_{k+1}$  induces an approximation of quality

$$-\frac{\log|P_{k+1}(\zeta)|}{\log H(P_{k+1})} = \rho + o(1) > 3, \qquad k \to \infty,$$

and so does  $R_{k+1}(T) = TP_{k+1}(T)$ , it follows that  $L_{3,1}^*$  and  $L_{3,2}^*$  decay with asymptotic slope -1/3 in some interval  $(q_k, b_k)$  and  $(q_k, c_k)$  respectively, for  $b_k$  and  $c_k$  local minima of  $L_{3,1}^*$  and  $L_{3,2}^*$  respectively. More precisely, the local minima  $(d_k, L_{P_{k+1}}^*(d_k))$  and  $(e_k, L_{R_{k+1}}^*(e_k))$  of the functions  $L_{P_{k+1}}^*$  and  $L_{R_{k+1}}^*$  as in (28), almost coincide with local minima  $(b_k, L_{3,1}^*(b_k))$  and  $(c_k, L_{3,2}^*(c_k))$ , respectively. By this more precisely we mean that all differences

$$|b_k - d_k|, |b_k - e_k|, |c_k - d_k|, |c_k - e_k|$$

as well as the corresponding differences of the  $L^*$  evaluations

$$\begin{split} |L_{3,1}^*(b_k) - L_{P_{k+1}}^*(d_k)|, & |L_{3,1}^*(b_k) - L_{R_{k+1}}^*(e_k)|, \\ |L_{3,2}^*(c_k) - L_{P_{k+1}}^*(d_k)|, & |L_{3,2}^*(c_k) - L_{R_{k+1}}^*(e_k)|, \end{split}$$

at these points are bounded by a fixed constant for all k. Very similarly it is obvious from the fact that  $P_{k+1}(\zeta)$  and  $R_{k+1}(\zeta)$  differ only by the factor  $\zeta$  that  $b_k$  and  $c_k$ are asymptotically equal, by which we mean their ratio  $b_k/c_k$  tends to one (in fact their difference  $|b_k - c_k|$  is again bounded) as  $k \to \infty$ . Hence with the parametric formula (29) for the parameter  $w_3^{(1)} = w_3^{(2)} = \rho$ , with

$$Q_k := e^{b_k} \quad \text{for} \quad k \ge 1,$$

(not to confuse with the polynomials  $Q_{k,i}$ ) we calculate

(60) 
$$\lim_{k \to \infty} \psi_{3,1}^*(Q_k) = \lim_{k \to \infty} \psi_{3,2}^*(Q_k) = \frac{1 - \sqrt{5}}{3(3 + \sqrt{5})}.$$

Since  $L_{3,1}^*$  and  $L_{3,2}^*$  both decay with asymptotic slope  $-\frac{1}{3}$  in intervals  $I_k := (q_k, b_k)$ , that is,

$$L_{3,1}^*(b_k) - L_{3,1}^*(q_k) = (b_k - q_k) \left( -\frac{1}{3} + \varepsilon \right)$$
 and  $L_{3,2}^*(b_k) - L_{3,2}^*(q_k) = (b_k - q_k) \left( -\frac{1}{3} + \varepsilon \right)$ ,

it follows from (30) that the sum  $L_{3,3}^* + L_{3,4}^*$  asymptotically increases with constant slope  $\frac{2}{3}$  in  $I_k$ , that is,

$$L_{3,3}^*(b_k) + L_{3,4}^*(b_k) - L_{3,3}^*(q_k) - L_{3,4}^*(q_k) = (b_k - q_k) \left(\frac{2}{3} + \varepsilon\right)$$

Consequently, if we can show that both  $L_{3,3}^*$  and  $L_{3,4}^*$  increase at most by  $\frac{1}{3}$  in any large subinterval of  $I_k$ , that is, for any  $q_k \le a < b \le b_k$ , we have

(61) 
$$L_{3,3}^*(b) - L_{3,3}^*(a) \le (b-a) \left(\frac{1}{3} + \varepsilon\right)$$
 and  $L_{3,4}^*(b) - L_{3,4}^*(a) \le (b-a) \left(\frac{1}{3} + \varepsilon\right)$ ,

then both must have asymptotically constant increase by precisely  $\frac{1}{3}$  in the entire interval  $I_k$ , i.e., equality in (61). We more precisely show the following claims:

**Claim A:** For any parameter  $\tilde{X} \in (H(P_k), \infty)$ , let

$$U_{k,\tilde{X}} = P_k \cdot E_t$$
 and  $V_{k,\tilde{X}} = P_k \cdot E_{t+1}$ 

with  $t = t(k, \tilde{X})$  chosen as the largest integer such that  $\max\{H(U_{k,\tilde{X}}), H(V_{k,\tilde{X}})\} \leq \tilde{X}$ . Then the functions  $L_{3,..}^{*}(q)$  arising from the succession (equals the pointwise minimum) of the  $L_{U_{k,\tilde{X}}}^{*}, L_{V_{k,\tilde{X}}}^{*}$  as  $\tilde{X}$  runs through  $(H(P_{k}), \infty)$  via (28) have asymptotically constant slope  $\frac{1}{3}$  in  $(b_{k-1}, \infty)$ . By this more precisely we mean that for any  $b_{k-1} \leq \tilde{X} < \tilde{Y}$  if  $(a, L_{U_{k,\tilde{X}}}(a))$  or  $(a, L_{V_{k,\tilde{X}}}(a))$  lies in the graph of  $L_{U_{k,\tilde{X}}}$  or  $L_{V_{k,\tilde{X}}}$ , respectively, and similarly for  $(b, L_{U_{k,\tilde{X}}}(b))$  or  $(b, L_{V_{k,\tilde{X}}}(b))$ , then we have

$$L^*_{U_{k,\tilde{Y}}}(b) - L^*_{U_{k,\tilde{X}}}(a) = (b-a) \left(\frac{1}{3} + \varepsilon\right) \text{ and } L^*_{V_{k,\tilde{Y}}}(b) - L^*_{V_{k,\tilde{X}}}(a) = (b-a) \left(\frac{1}{3} + \varepsilon\right).$$

**Claim B:** Moreover if we restrict to  $\tilde{X} \in (H(P_{k+1}), H(P_{k+2}))$ , then the functions  $L^*_{U_{k,\tilde{X}}}$  and  $L^*_{V_{k,\tilde{X}}}$  induce  $L^*_{3,3}$  and  $L^*_{3,4}$  on  $I_k$ , respectively (remark: as we will see later on they induce  $L^*_{3,1}$  and  $L^*_{3,2}$  in intervals  $(b_{k-1}, q_k)$  if we let  $\tilde{X} \in (H(P_k), H(P_{k+1}))$ ).

First recall that at the beginning  $q_k$  of the interval  $I_k$  the successive minima are induced basically by  $\{P_k, TP_k, P_{k+1}, TP_{k+1}\}$ . Claim A follows basically directly

from Lemma 3.3, where  $E_t$  and  $E_{t+1}$  respectively play the role of Q and  $P_k$  the role of P. Note also that  $\delta$  from Lemma 3.3 tends to 0 in our context in view of (58), which also implies that the minima (in fact the entire functions) of consecutive functions of the form  $L_{U_{k,\bar{X}}}^*$  or  $L_{V_{k,\bar{X}}}^*$  do not differ much. Finally, it should be pointed out that the condition that  $1/\log H(Q) = O(\delta)$  does not cause problems since for any fixed  $\delta > 0$  and smaller heights H(Q) only minor changes of the function  $L_{3,.}^*(q)$  can appear in intervals  $(b_{k-1}, b_{k-1} + O(1))$ , so that the global behavior of the function is not affected. For Claim B further observe that  $\{U_{k,\bar{X}}, V_{k,\bar{X}}\}$  span the same space as  $\{P_k, TP_k\}$  for all  $\tilde{X} \in (H(P_k), \infty)$ , and we have already noticed that polynomials in the space  $\{P_{k+1}, TP_{k+1}\}$  induce the first two successive minima in  $I_k$  and  $\{P_k, TP_k, P_{k+1}, TP_{k+1}\}$  are linearly independent. Hence  $L_{3,3}^*$  and  $L_{3,4}^*$  are bounded above by  $L_{U_{k,\bar{X}}}^*$  in  $I_k$  respectively, and thus each increase at most by  $\frac{1}{3}$ . As noticed above, we may conclude  $L_{3,3}^*$  and  $L_{3,4}^*$  must actually coincide with the functions induced by  $L_{U_k,\bar{X}}^*$  and  $L_{V_k,\bar{X}}^*$ , respectively.

Thus together with (60) we have proved

(62) 
$$\lim_{k \to \infty} \psi_{3,1}^*(Q_k) = \lim_{k \to \infty} \psi_{3,2}^*(Q_k)$$
$$= \lim_{k \to \infty} -\psi_{3,3}^*(Q_k) = \lim_{k \to \infty} -\psi_{3,4}^*(Q_k) = \frac{1 - \sqrt{5}}{3(3 + \sqrt{5})}$$

We show next that in the interval  $J_k := (b_k, q_{k+1})$  the functions  $L_{3,1}^*, L_{3,2}^*$  have slope  $-\frac{1}{3}$  whereas the functions  $L_{3,3}^*, L_{3,4}^*$  have (asymptotic) slope  $\frac{1}{3}$  until they all meet (asymptotically) at  $q_{k+1}$ . More precisely

$$L_{3,1}^*(q_{k+1}) - L_{3,1}^*(b_k) = (q_{k+1} - b_k) \left( -\frac{1}{3} + \varepsilon \right),$$
  
$$L_{3,2}^*(q_{k+1}) - L_{3,2}^*(b_k) = (q_{k+1} - b_k) \left( -\frac{1}{3} + \varepsilon \right)$$

whereas

$$L_{3,3}^{*}(q_{k+1}) - L_{3,3}^{*}(b_{k}) = (q_{k+1} - b_{k}) \left(\frac{1}{3} + \varepsilon\right),$$
  
$$L_{3,4}^{*}(q_{k+1}) - L_{3,4}^{*}(b_{k}) = (q_{k+1} - b_{k}) \left(\frac{1}{3} + \varepsilon\right)$$

and

$$L_{3,4}^*(q_{k+1}) - L_{3,1}^*(q_{k+1}) \le \varepsilon q_{k+1}.$$

Again by (59) with index shift k to k + 1 we know that for arbitrarily small  $\varepsilon$  and all large  $k \ge k_0(\varepsilon)$  we indeed have

(63) 
$$|L_{3,i}^*(q_{k+1})| \le \varepsilon q_{k+1} \quad \text{for} \quad 1 \le i \le 4.$$

Since we have shown that  $L_{3,1}^*$  and  $L_{3,2}^*$  decay in  $I_k$  with slope  $-\frac{1}{3}$  and (62) holds it suffices to show that  $J_k$  has asymptotically equal length to  $I_k$ , that is,  $\lim_{k\to\infty} |J_k|/|I_k| = 1$ , to conclude that  $L_{3,3}^*$  and  $L_{3,4}^*$  must decay with the minimum

possible slope  $-\frac{1}{3}$  in the entire interval  $J_k$  and more precisely (64)

$$\lim_{k \to \infty} -\frac{L_{3,1}^*(b_k)}{q_{k+1} - b_k} = \lim_{k \to \infty} -\frac{L_{3,2}^*(b_k)}{q_{k+1} - b_k} = \lim_{k \to \infty} \frac{L_{3,3}^*(b_k)}{q_{k+1} - b_k} = \lim_{k \to \infty} \frac{L_{3,4}^*(b_k)}{q_{k+1} - b_k} = \frac{1}{3}$$

We show the claim that  $I_k$  and  $J_k$  have asymptotically equal length, that is,  $|I_k|/|J_k| = 1 + o(1)$  as  $k \to \infty$ . By construction this is equivalent to  $b_k$  being asymptotically equal to  $(q_k + q_{k+1})/2$ , that is,  $b_k = (q_k + q_{k+1})/2 + o(q_k)$ . Since  $\lim_{k\to\infty} \log H(P_{k+1})/\log H(P_k) = v$  and  $L_{3,..}^*(q_k) = o(q_k)$  and  $L_{3,..}^*(q_{k+1}) = o(q_{k+1})$  as  $k \to \infty$ . Further notice that  $L_{3,.1}^*, L_{3,2}^*$  decay in  $(q_k, b_k)$  induced by  $P_{k+1}, TP_{k+1}$  and thus by (28) we have  $L_{3,.j}^*(q_i) = \log H(P_{i+1}) - q_i/3 + O(1)$  for  $1 \le j \le 4$  and all  $i \ge 1$ . Putting all this together leads to

(65) 
$$\lim_{k \to \infty} \frac{q_{k+1}}{q_k} = \nu$$

Thus the claimed asymptotic relation  $b_k = (q_k + q_{k+1})/2 + o(q_k)$  is equivalent to  $b_k = q_k \cdot (1 + \nu)/2 + o(q_k)$ . We know that at  $Q_k = e^{b_k}$ . We have asymptotically

(66) 
$$\psi_{3,3}^*(Q_k) = \frac{b_k - q_k}{3} + o(q_k), \qquad k \to \infty,$$

since  $L_{3,3}^*$  and  $L_{3,3}^*$  are small at  $q_k$  by (59) and rise with slope  $\frac{1}{3}$  in  $I_k$ . We remark that the asymptotic (66) holds for  $\psi_{3,4}^*(Q_k)$  as well. On the other hand (62) provides an asymptotic formula for  $\psi_{3,3}^*(Q_k)$  and  $\psi_{3,4}^*(Q_k)$ . It follows directly from the definition of  $L_{3,j}^*$  via  $\psi_{3,j}^*$  in (26) that  $\psi_{3,3}^*(Q_k)$  is the slope from the origin to  $(b_k, L_{3,3}^*(b_k))$  of  $L_{3,3}^*$  in the Schmidt–Summerer diagram (and similarly for  $L_{3,4}^*$ ). Hence asymptotically

(67) 
$$\psi_{3,3}^*(Q_k) = \psi_{3,4}^*(Q_k) = \frac{\sqrt{5-1}}{3(3+\sqrt{5})}b_k + o(b_k), \qquad k \to \infty$$

Again the asymptotic (67) holds for  $\psi_{3,4}^*(Q_k)$  as well. Comparing the two expressions for  $\psi_{3,3}^*(Q_k)$  in (66) and (67), with a short computation, indeed we verify  $b_k = q_k \cdot (1+\nu)/2 + o(q_k)$ , so we have proved that  $I_k$  and  $J_k$  have asymptotically equal length.

Since consequently  $L_{3,3}^*$  and  $L_{3,4}^*$  both asymptotically decay with slope  $-\frac{1}{3}$  in  $J_k$ , from (30) again we deduce that the sum  $L_{3,1}^* + L_{3,2}^*$  must asymptotically increase by  $\frac{2}{3}$  in  $J_k$ . Now recall in Claim A we showed that  $L_{U_{k,\tilde{X}}}^*$ ,  $L_{V_{k,\tilde{X}}}^*$  asymptotically induce an increase with slope at most  $\frac{1}{3}$  in the entire interval  $(b_{k-1}, \infty)$  if we let  $\tilde{X}$ run through  $(H(P_k), \infty)$ . Hence if we restrict to  $\tilde{X} \in (H(P_k), H(P_{k+1}))$ , by a very similar argument as in Claim B, in the interval  $(b_{k-1}, q_k)$  they induce  $L_{3,1}^*$  and  $L_{3,2}^*$ such that they both asymptotically increase precisely with slope  $\frac{1}{3}$ . By index shift the analogous claim is clearly also true for  $(b_k, q_{k+1}) = J_k$ . Hence indeed both  $L_{3,1}^*$  and  $L_{3,2}^*$  must asymptotically increase with slope precisely  $\frac{1}{3}$  in the entire interval  $J_k$ . Observe that the end of  $J_k$  is the beginning of  $I_{k+1}$ , so that we have basically established a complete description of all functions  $L_{3,1}^*, \ldots, L_{3,4}^*$  on  $(0, \infty)$ . The characterizations of the graphs of  $L_{3,i}^*(q)$  established above show that asymptotically at the values  $q = b_k$  both the smallest local minima of  $L_{3,1}^*(q), L_{3,2}^*(q)$  (in the sense of minimal values of  $\psi_{3,1}^*(Q), \psi_{3,2}^*(Q)$ ) and the largest local maxima of  $L_{3,3}^*(q), L_{3,4}^*(q)$  (in the sense of maximal values of  $\psi_{3,3}^*(Q), \psi_{3,4}^*(Q)$ ) are attained. Moreover both  $|L_{3,1}^*(b_k) - L_{3,2}^*(b_k)|$  and  $|L_{3,3}^*(b_k) - L_{3,4}^*(b_k)|$  are bounded uniformly in k, in fact more generally  $|L_{3,1}^*(q) - L_{3,2}^*(q)|$  and  $|L_{3,3}^*(q) - L_{3,4}^*(q)|$  are uniformly bounded for  $q \in (0, \infty)$ . Thus with (62) we have

$$\underline{\psi}_{3,1}^* = \underline{\psi}_{3,2}^* = \frac{1 - \sqrt{5}}{3(3 + \sqrt{5})}$$
 and  $\overline{\psi}_{3,3}^* = \overline{\psi}_{3,4}^* = \frac{\sqrt{5} - 1}{3(3 + \sqrt{5})}$ 

With (22), (23) and (24) we derive

(68) 
$$w_3(\zeta) = w_{3,2}(\zeta) = \rho$$
 and  $\lambda_3(\zeta) = \lambda_{3,2}(\zeta) = \frac{1}{\sqrt{5}}$ .

This contains in particular the claims in (11).

**Remark 4.2.** We can also determine the remaining constants  $w_{3,i}$ ,  $\lambda_{3,i}$ ,  $\hat{w}_{3,i}$ ,  $\hat{\lambda}_{3,i}$  for extremal numbers. From (25) and (68) we deduce

 $\square$ 

(69) 
$$\widehat{w}_{3,3}(\zeta) = \widehat{w}_{3,4}(\zeta) = \sqrt{5}, \qquad \widehat{\lambda}_{3,3}(\zeta) = \widehat{\lambda}_{3,4}(\zeta) = \frac{1}{\rho}.$$

Moreover the above characterizations of the functions  $L_{3,i}^*$  imply

$$\overline{\psi}_{3,1}^* = \overline{\psi}_{3,2}^* = \underline{\psi}_{3,1}^* = \underline{\psi}_{3,2}^* = 0.$$

With (24) and (25) this is equivalent to

(70)  
$$w_{3,3}(\zeta) = w_{3,4}(\zeta) = \widehat{w}_3(\zeta) = \widehat{w}_{3,2}(\zeta) = 3, \\\lambda_{3,3}(\zeta) = \lambda_{3,4}(\zeta) = \widehat{\lambda}_3(\zeta) = \widehat{\lambda}_{3,2}(\zeta) = \frac{1}{3}.$$

The description of the combined graph of the functions  $L_{3,j}^*(q)$  and the information on the structure of the polynomials inducing them from the proof of Theorem 2.1 allows one to estimate the approximation to an extremal number by algebraic numbers of degree precisely three.

*Proof of Theorem 2.2.* It follows from the proof of Theorem 2.1 and the description above that the first two successive minima functions of the linear form problem related to  $\psi_{3,1}^*$ ,  $\psi_{3,2}^*$  are induced by polynomial multiples of  $P_k$  from Theorem 1.1, and for each *k* these multiples span the same space as  $\{P_k, TP_k\}$ . Since  $P_k$  have degree two, there is no irreducible polynomial of degree three which lies in the space spanned by  $\{P_k, TP_k\}$  for some *k*. Thus the optimal exponent in (13) is not larger than  $w_{3,3}(\zeta)$ . On the other hand it was shown in the proof of Theorem 2.1

that  $w_{3,3}(\zeta) = 3$ , see (70). Thus, combining these facts, we see that indeed (13) has only finitely many solutions in  $Q \in \mathbb{Z}[T]$  an irreducible polynomial of degree precisely three. From (13) we infer (14) by a standard argument. Indeed if *R* is the minimal polynomial of some  $\alpha$  then  $|R(\zeta)| = |R(\zeta) - R(\alpha)| = |\zeta - \alpha| \cdot R'(z)$  for some *z* between  $\alpha$  and  $\zeta$  by the intermediate theorem of differentiation. On the other hand  $|R'(z)| \ll H(R)$  for bounded *z* is easy to see, and the claim (14) follows from (13).

Next we show (16) and (17). By essentially the argument from the proof of (13) again  $\widehat{w}_{3,3}(\zeta)$  is an upper bound for the exponent in (16) for some large X. On the other hand we have noticed in (69) that  $\widehat{w}_{3,3}(\zeta) = \widehat{w}_{3,4}(\zeta) = \sqrt{5}$ . Combining these yields (16) and we deduce (17) from it very similarly as (14) from (13).

For (15) recall that in the proof of Theorem 2.1 we showed that for any large k there exists a linear polynomial  $E_l$  such that, with  $X := H(P_{k+1})$  and  $Q_{k,1} := P_k E_l$ , we have

(71) 
$$H(Q_{k,1}) \asymp H(P_{k+1}) = X,$$
$$|Q_{k,1}(\zeta)| \le X^{-3+\epsilon}, \qquad |P_{k+1}(\zeta)| \le X^{-3+\epsilon}$$

Since  $Q_{k,1}$  is not irreducible by construction and  $P_{k+1}$  has degree only 2, we consider the polynomials  $S_{k,j}(T) := Q_{k,1}(T) + jT \cdot P_{k+1}(T)$  for  $j \in \{1, 2\}$ . We show that at least one of these two polynomials has the desired properties (in fact we need the distinction only for the right hand side of (15); the left follows for both j = 1 and j = 2). The polynomials  $S_{k,j}(T)$  obviously have degree three and height  $H(S_{k,j}) \ll X$ . Moreover with (71) we infer

(72) 
$$|S_{k,j}(\zeta)| = |Q_{k,1}(\zeta) + j\zeta P_{k+1}(\zeta)| \le |Q_{k,1}(\zeta)| + j|\zeta| \cdot |P_{k+1}(\zeta)| \ll_{\zeta} X^{-3+\epsilon},$$

for  $1 \le j \le 2$ . Next we check that  $S_{k,j}$  are irreducible for large k and  $1 \le j \le 2$ . Consider j fixed and suppose  $S_{k,j}$  is reducible. Then we may write  $S_{k,j}(T) = M(T)N(T)$  for  $M, N \in \mathbb{Z}[T]$  each of degree one or two. Then  $|S_{k,j}(\zeta)| = |M(\zeta)| \cdot |N(\zeta)|$  and it follows from (42) and (72) that at least one of the inequalities

$$|M(\zeta)| \le H(M)^{-3+2\epsilon}$$
 or  $|N(\zeta)| \le H(N)^{-3+2\epsilon}$ 

must be satisfied; see also the remark subsequent to (42). Without loss of generality say this holds for *M*. However, since  $w_{2,2}(\zeta) \le \tau < 3$ , see (8), and *M* has degree at most two, it follows from Theorem 1.1 that the inequality can only be satisfied if *M* is some  $P_l$  from Theorem 1.1. However, by construction of  $S_{k,j}$  we clearly cannot have  $P_k|S_{k,j}$  or  $P_{k+1}|S_{k,j}$ . Thus  $M = P_l$  for some  $l \le k - 1$ . Theorem 1.1 further implies

$$H(M) \ll H(P_{k-1}) \ll H(P_{k+1}) \cdot \frac{H(P_{k-1})}{H(P_{k+1})} = X \cdot \frac{H(P_{k-1})}{H(P_{k+1})} \ll X^{1/\nu^2} = X^{1/\tau}$$

and it follows further that

(73) 
$$|M(\zeta)| \asymp H(M)^{-\rho} \gg X^{-\rho/\tau} = X^{-\nu}.$$

Since  $M = P_l$  has degree two and  $S_{k,j}$  degree three, the polynomial N must have degree one such that by  $\lambda_1(\zeta) = 1$  from (7) we have

(74) 
$$|N(\zeta)| \gg H(N)^{-1-\epsilon} \gg X^{-1-\epsilon}$$

Combining (73) and (74) yields

$$|S_{k,j}(\zeta)| = |M(\zeta)| \cdot |N(\zeta)| \gg X^{-\nu - 1 - \epsilon} = X^{-\tau - \epsilon}.$$

Again from  $\tau < 3$  we obtain a contradiction to (72) for small  $\epsilon$ . Hence the assumption was wrong and indeed  $S_{k,j}$  must be irreducible for  $j \in \{1, 2\}$ , and in view of (72) we have finished the proof of the left-hand side of (15).

For the right-hand side of (15) suppose we have already shown that for all large k and some  $j = j(k) \in \{1, 2\}$  we have

(75) 
$$|S'_{k,j}(\zeta)| \gg X^{1-\epsilon}.$$

Then the claim follows together with (72) from the left-hand side for  $\alpha$  some root of the corresponding  $S_{k,j}$  by a similar standard argument as in the deduction of (14) from (13). Indeed it is well known that any polynomial  $U \in \mathbb{Z}[T]$  has a root  $\beta$  that satisfies  $|\beta - \zeta| \ll |U(\zeta)|/H(U)$ , see for example [Roy 2004a]. The claim follows with  $U = S_{k,j}$ . It remains to be checked that (75) holds, for which we use (10). First note that the derivative of  $S_{k,j}$  can be written

(76) 
$$|S'_{k,j}(\zeta)| = |Q'_{k,1}(\zeta) + jP_{k+1}(\zeta) + j\zeta P'_{k+1}(\zeta)|, \qquad 1 \le j \le 2.$$

Obviously the term  $jP_{k+1}(\zeta)$  in the sum is negligible since it is very small. Hence (76) can be small only if  $Q'_{k,1}(\zeta)$  is of the same order (and reverse sign) as  $j\zeta P'_{k+1}(\zeta)$ . On the other hand (10) implies for all large k the estimate

$$|j\zeta P'_{k+1}(\zeta)| \ge j|\zeta|H(P_{k+1})^{1-\epsilon} \gg_{\zeta} X^{1-\epsilon}$$
 for  $j \in \{1, 2\}$ .

and very similarly the difference between the right-hand sides in (76) for j = 2and j = 1 is at least of order  $X^{1-\epsilon}$  as well. It follows that (75) can be violated for at most one index  $j \in \{1, 2\}$ , and for the other index (75) must be satisfied. This finishes the proof of (15).

We finish by giving a heuristic argument why the exponents in (16) and (17) should be optimal as well. For any  $\tilde{X}$  we can again consider linear combinations  $S_{k,j}(T) = jTP_{k+1}(T) + P_k(T)E_t(T)$  for  $k = k(\tilde{X})$  largest possible such that  $H(P_{k+1}) \leq \tilde{X}$  and some  $E_t$  of degree one from the proof of Theorem 2.1 such that (16) is satisfied for  $Q(T) = Q_{k,1}(T) = P_k(T)E_t(T)$ . Given the irreducibility of

 $S_{k,j}$  for all large k and j rather small, we can again basically proceed as in the proof of (15). However, the method from the proof of (15) to guarantee the irreducibility of some of the arising  $S_{k,j}(T)$  does not work here.

#### Acknowledgement

The author warmly thanks the anonymous referee for the careful reading and for pointing out inaccuracies.

#### References

- [Adamczewski and Bugeaud 2010] B. Adamczewski and Y. Bugeaud, "Mesures de transcendance et aspects quantitatifs de la méthode de Thue–Siegel–Roth–Schmidt", *Proc. Lond. Math. Soc.* (3) **101**:1 (2010), 1–26. MR Zbl
- [Bugeaud 2010] Y. Bugeaud, "On simultaneous rational approximation to a real number and its integral powers", *Ann. Inst. Fourier (Grenoble)* **60**:6 (2010), 2165–2182. MR Zbl
- [Bugeaud and Laurent 2005] Y. Bugeaud and M. Laurent, "Exponents of Diophantine approximation and Sturmian continued fractions", *Ann. Inst. Fourier (Grenoble)* **55**:3 (2005), 773–804. MR Zbl
- [Bugeaud and Schleischitz 2016] Y. Bugeaud and J. Schleischitz, "On uniform approximation to real numbers", *Acta Arith.* **175** (2016), 255–268.
- [Davenport and Schmidt 1969] H. Davenport and W. M. Schmidt, "Approximation to real numbers by algebraic integers", *Acta Arith.* **15** (1969), 393–416. MR Zbl
- [German 2012] O. N. German, "On Diophantine exponents and Khintchine's transference principle", *Mosc. J. Comb. Number Theory* **2**:2 (2012), 22–51. MR Zbl
- [Khintchine 1926] A. Khintchine, "Über eine Klasse linearer diophantischer Approximationen", *Rend. Circ. Mat. Palermo* **50**:2 (1926), 170–195. Zbl
- [Roy 2003] D. Roy, "Approximation to real numbers by cubic algebraic integers, II", *Ann. of Math.*(2) 158:3 (2003), 1081–1087. MR Zbl
- [Roy 2004a] D. Roy, "Approximation to real numbers by cubic algebraic integers, I", *Proc. London Math. Soc.* (3) **88**:1 (2004), 42–62. MR Zbl
- [Roy 2004b] D. Roy, "Diophantine approximation in small degree", pp. 269–285 in *Number theory*, edited by H. Kisilevsky and E. Z. Goren, CRM Proc. Lecture Notes 36, Amer. Math. Soc., Providence, RI, 2004. MR Zbl
- [Schleischitz 2013] J. Schleischitz, "Diophantine approximation and special Liouville numbers", *Commun. Math.* **21**:1 (2013), 39–76. MR Zbl
- [Schleischitz 2014] J. Schleischitz, "Two estimates concerning classical diophantine approximation constants", *Publ. Math. Debrecen* 84:3-4 (2014), 415–437. MR Zbl
- [Schleischitz 2016] J. Schleischitz, "On the spectrum of Diophantine approximation constants", *Mathematika* **62**:1 (2016), 79–100. MR Zbl
- [Schmidt and Summerer 2009] W. M. Schmidt and L. Summerer, "Parametric geometry of numbers and applications", *Acta Arith.* **140**:1 (2009), 67–91. MR Zbl
- [Schmidt and Summerer 2013a] W. M. Schmidt and L. Summerer, "Diophantine approximation and parametric geometry of numbers", *Monatsh. Math.* **169**:1 (2013), 51–104. MR Zbl
- [Schmidt and Summerer 2013b] W. M. Schmidt and L. Summerer, "Simultaneous approximation to three numbers", *Mosc. J. Comb. Number Theory* **3**:1 (2013), 84–107. MR Zbl

[Wirsing 1961] E. Wirsing, "Approximation mit algebraischen Zahlen beschränkten Grades", J. Reine Angew. Math. 206 (1961), 67–77. MR Zbl

Received March 9, 2016. Revised August 22, 2016.

Johannes Schleischitz Institute of Mathematics Universität für Bodenkultur Wien Gregor-Mendel-Strasse 33 1180 Vienna Austria

johannes.schleischitz@boku.ac.at

### PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

#### msp.org/pjm

#### EDITORS

Don Blasius (Managing Editor) Department of Mathematics University of California Los Angeles, CA 90095-1555 blasius@math.ucla.edu

Vyjayanthi Chari Department of Mathematics University of California Riverside, CA 92521-0135 chari@math.ucr.edu

Kefeng Liu Department of Mathematics University of California Los Angeles, CA 90095-1555 liu@math.ucla.edu

Igor Pak Department of Mathematics University of California Los Angeles, CA 90095-1555 pak.pjm@gmail.com

Paul Yang Department of Mathematics Princeton University Princeton NJ 08544-1000 yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

#### SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI CALIFORNIA INST. OF TECHNOLOGY INST. DE MATEMÁTICA PURA E APLICADA KEIO UNIVERSITY MATH. SCIENCES RESEARCH INSTITUTE NEW MEXICO STATE UNIV. OREGON STATE UNIV.

Paul Balmer

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

balmer@math.ucla.edu

Robert Finn

Department of Mathematics

Stanford University

Stanford, CA 94305-2125

finn@math.stanford.edu

Sorin Popa

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

popa@math.ucla.edu

STANFORD UNIVERSITY UNIV. OF BRITISH COLUMBIA UNIV. OF CALIFORNIA, BERKELEY UNIV. OF CALIFORNIA, DAVIS UNIV. OF CALIFORNIA, LOS ANGELES UNIV. OF CALIFORNIA, RIVERSIDE UNIV. OF CALIFORNIA, SAN DIEGO UNIV. OF CALIF., SANTA BARBARA Daryl Cooper Department of Mathematics University of California Santa Barbara, CA 93106-3080 cooper@math.ucsb.edu

Jiang-Hua Lu Department of Mathematics The University of Hong Kong Pokfulam Rd., Hong Kong jhlu@maths.hku.hk

Jie Qing Department of Mathematics University of California Santa Cruz, CA 95064 qing@cats.ucsc.edu

UNIV. OF CALIF., SANTA CRUZ UNIV. OF MONTANA UNIV. OF OREGON UNIV. OF SOUTHERN CALIFORNIA UNIV. OF UTAH UNIV. OF WASHINGTON WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.



nonprofit scientific publishing http://msp.org/

© 2017 Mathematical Sciences Publishers

# **PACIFIC JOURNAL OF MATHEMATICS**

Volume 287 No. 2 April 2017

Maximal operators for the <i>p</i> -Laplacian family	257
PABLO BLANC, JUAN P. PINASCO and JULIO D. ROSSI	
Van Est isomorphism for homogeneous cochains	297
ALEJANDRO CABRERA and THIAGO DRUMMOND	
The Ricci–Bourguignon flow	337
GIOVANNI CATINO, LAURA CREMASCHI, ZINDINE DJADLI, CARLO MANTEGAZZA and LORENZO MAZZIERI	
The normal form theorem around Poisson transversals	371
PEDRO FREJLICH and IOAN MĂRCUŢ	
Some closure results for <i>C</i> -approximable groups	393
DEREK F. HOLT and SARAH REES	
Coman conjecture for the bidisc	411
ŁUKASZ KOSIŃSKI, PASCAL J. THOMAS and WŁODZIMIERZ ZWONEK	
Endotrivial modules: a reduction to $p'$ -central extensions	423
CAROLINE LASSUEUR and JACQUES THÉVENAZ	
Infinitely many positive solutions for the fractional	439
Schrödinger–Poisson system	
WEIMING LIU	
A Gaussian upper bound of the conjugate heat equation along	465
Ricci-harmonic flow	
XIAN-GAO LIU and KUI WANG	
Approximation to an extremal number, its square and its cube	485
JOHANNES SCHLEISCHITZ	